

Gravitational Shielding Research

The following list of possible keywords relevant to the research of E.E. Podkletnov in the area of gravitational shielding can be implemented with a Web search algorithm to locate related documents on the Web and sort them by relevance.

Gravitational shielding theory and research

1. Podkletnov
 - a. Variations:
 - i. Yevgeni
 - ii. Yevgeniy
 - iii. Yevgeny
 - iv. Yevgenij
 - v. Eugene
 - vi. Ye.
 - vii. Ye.Ye.
 - viii. Evgenny
 - ix. Evgeniy
 - x. Evgenij
 - xi. E.
 - xii. E.E.
2. Podkletnov Effect
3. Gravitational Meissner effect
 - a. Variations
 - i. Meissner effect
 - ii. Gravitoelectromagnetic effects
4. Gravitational shielding
 - a. Variations
 - i. Gravity Shielding
 - ii. Anti-Gravity Shielding
 - iii. Gravitational screening
 - iv. Gravitational force shielding
 - v. Superconductor shield gravity
5. Anti-Gravity
 - a. Variations
 - i. Antigravity
6. Electrogravitics
7. London equation
 - a. Variations
 - i. Maxwell-London equations
8. Mass density
9. Weight reduction
10. Project Greenglow
11. Project Grasp
12. Tampere University of Technology

Related terms:

1. Electromagnetism
 - a. Gravitational effects
 - b. Maxwell equations
 - c. Current density
 - d. Electric charge
 - e. Light velocity
 - f. Permittivity
 - g. Magnetic permeability
 - h. Superconducting materials
 - i. Electromagnetic dispersion
 - j. Electromagnetic shielding
 - k. Magnetic levitation

1. Buoyancy correction
- m. RF Magnetron Sputtering
2. Mechanics
 - a. Rotating machinery
 - b. Flywheels
 - c. Torque
 - d. Angular momentum
 - e. Rotational kinetic energy
 - f. Levitating superconducting disks
 - i. Disk rotation speed
 - ii. Weight loss
 - iii. Electrooptical balance system
3. Materials Physics
 - a. Refractory materials
 - b. Cerium compounds
 - c. Indium compounds
 - d. High temperature properties
 - e. Electric conductivity of solids
 - f. Porosity
 - g. Thermodynamic stability
 - h. Strength of materials
 - i. Combustion
 - j. Reduction
 - k. Electroconducting ceramics
 - l. Propane combustion
 - m. N-type conductivity
 - n. Mixed ceramics
 - o. Cryostats
 - p. Yttrium compounds
 - i. Yttrium barium copper oxides
 1. $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$
 - ii. Oxide Superconductors
- q. Copper oxides
- r. Ceramic materials
- s. Crystal structure
4. Boeing
 - a. Phantom Works
 - b. George Mueller
 - c. Project Grasp
5. British Aerospace
 - a. BAe
 - b. BAe Systems PLC (U.BA)
 - c. Project Greenglow
6. NASA
7. Tampere University of Technology
 - a. Finland

Podkletnov's collaborators & other researchers

1. Buzea, C.Gh
 - a. Institute of Technical Physics, Iasi, Romania
 - b. Technical University 'Gh. Asachi', Iasi, Romania
 - c. Agop, M
 - d. Nica, P.
2. Bass, Robert
 - a. Journal of New Energy
3. de Podesta, Michael
 - a. Bull, Martyn
 - b. Birkbeck College, London , England
4. Akopov, F.A.
 - a. Barykin, B.M.

- b. Novov, Yu.D.
- c. Pakhomov, E.P.
- d. Institut Vysokikh Temperatur RAN, Moscow, Russia (High Temperature Institute, Russian Academy of Science, Moscow)
- 5. Nieminen, R.
 - a. Tampere University of Technology, Finland
- 6. Jaervinen, R.J.O.
 - a. Vuorinen, P.T.
 - b. Maentylae, T.A.
 - c. Tampere University of Technology, Finland
- 7. Romanov, A.I.
 - a. Valjano, G.E.
- 8. Modanese, G.
- 9. Unnikrishnan, C.S.
 - a. Tata Institute of Fundamental Research, Bombay, India
- 10. Holwerda, Martin
- 11. Selin, Alexey

[4. Yahoo Physics](#)

Publications


- 1. Journal of New Energy
- 2. Physica C: Superconductivity and its Applications
- 3. Journal of Crystal Growth
- 4. Proceedings of the Unified International Technical Conference on Refractories - UNITECR '89
- 5. Europhysics Letters
- 6. Physics Letters
- 7. American Institute of Aeronautics & Astronautics
- 8. Jane's Information Group
 - a. Jane's Defence Weekly

Research Organizations

- 1. [Tampere University of Technology, Finland](#)
- 2. [Institut Vysokikh Temperatur RAN, Moscow, Russia](#)
(High Temperature Institute, [Russian Academy of Science](#), Moscow)
- 3. [Tata Institute of Fundamental Research, Bombay, India](#)
- 4. [Institute of Technical Physics, Iasi, Romania](#)
[Technical University 'Gh. Asachi', Iasi, Romania](#)
- 5. [Birkbeck College, London , England](#)
- 6. [Boeing](#)
 - a. [A&M Systems](#)
 - i. [Phantom Works](#)
 - 1. Project Grasp
 - ii. [George Muellner](#)
- 7. [British Aerospace](#)
 - a. [BAe Systems PLC](#)
 - i. Project Greenglow
- 8. [NASA](#)

Databases

- 1. [Engineering Information](#)
 - a. [Engineering Village 2](#)
 - i. Compendex
- 2. [American Institute of Aeronautics & Astronautics](#)
- 3. [Jane's Information Group](#)
 - a. [Jane's Defence Weekly](#)
 - b. [Online Research](#)



[[Latest Headlines](#) | [Market Overview](#) | [News Alerts](#) | [More Dow Jones Business News](#)]

Monday July 29, 6:12 am Eastern Time

Dow Jones Business News

Boeing Attempting To Create Gravity-Defying Device - BBC

Related Quote

[BA](#) 40.90 -0.75
delayed 20 mins - [disclaimer](#)
Quote Data provided by [Reuters](#)

LONDON -(Dow Jones)- Researchers at the world's largest aircraft maker, Boeing Co . , are using the work of a controversial Russian scientist to try to create a device that will defy gravity, the BBC reported Monday on its Web site.

ADVERTISEMENT

The screenshot shows the Yahoo! Finance Real-Time Package interface. At the top, it says "YAHOO! Finance Real-Time Package". Below that, there's a "Quote" section with a dropdown menu set to "My First Portfolio". The portfolio data table includes:

Symbol	Last	Change	Time
Dow Jones	10911.90	+39.30	4:03 PM
S&P 500	1255.82	+7.74	4:23 PM
MCD	30.28	-0.41	4:00 PM
^GDAIG	6123.26	+82.04	2:15 PM
Nasdaq	2110.49	+25.99	5:16 PM
GE	49.00	-0.02	4:01 PM
BASF.F	47.75	+0.85	1:30 PM
YHOO	18.11	+0.44	4:00 PM

Below the portfolio data, there's a "Recent News and Analysis" section with a table:

Symbol	Time	Source
T	May 31 7:54 PM	AP
AAC	May 31 7:31 PM	Reuters Securities
CSCO	May 31 7:20 PM	Industry Standard

The BBC said Boeing is examining an experiment by Yevgeny Podkletnov, who claims to have developed a device which can shield objects from the Earth's pull.

Podkletnov is viewed with suspicion by many conventional scientists, but the involvement of Boeing has lent his work a new credibility, the BBC said.

The project is being run by the top-secret Phantom Works in Seattle, the part of the company which handles Boeing's most sensitive programs. The head of the Phantom Works, George Muellner, told the security analysis journal Jane's Defence Weekly that the science appeared to be valid and plausible.

Podkletnov says he countered the effects of gravity in an experiment at the Tampere University of Technology in Finland in 1992.

The scientist says he found that objects above a superconducting ceramic disc rotating over powerful electromagnets lost weight.

The reduction in gravity was small, about 2%, but the implications - for example, in terms of cutting the energy needed for a plane to fly - were immense, the BBC said.

Scientists who investigated Podkletnov's work, however, said the experiment was fundamentally flawed and that negating gravity was impossible.

But documents obtained by Jane's Defence Weekly and seen by the BBC show Boeing is taking Podkletnov's research seriously.

The hypothesis is being tested in a program codenamed Project Grasp. Boeing is the latest in a series of high-profile institutions trying to replicate Podkletnov's experiment.

The military wing of the U.K. hi-tech group BAe Systems PLC (U.BA) is working on an antigravity program, dubbed Project Greenglow.

The US space agency, NASA, is also attempting to reproduce Podkletnov's findings, but a preliminary report indicates the effect does not exist. -0-

[Email this story](#) - [Most-emailed articles](#) - [Most-viewed articles](#)

More Quotes and News: The Boeing Co (NYSE:[BA](#) - [News](#))

Related News Categories: [aerospace/defense](#)

ADVERTISEMENT

Special Offers

- [Access Your PC from Anywhere - Free Download](#)
- [\\$8.95 Domain Name Registrations & Transfers at GoDaddy.com!](#)
- [Paying too much for car insurance? Find out](#)
- [Interest rates won't stay this low forever. Yahoo! Mortgage Center](#)

[Help](#)

Copyright © 2002 Yahoo! Inc. All rights reserved. [Privacy Policy](#) - [Terms of Service](#)

Copyright © 2002 Dow Jones. All rights reserved.

[Questions or Comments?](#)

You are in: **Science/Nature**

Monday, 29 July, 2002, 03:23 GMT 04:23 UK

News Front Page**Africa****Americas****Asia-Pacific****Europe****Middle East****South Asia****UK****Business****Entertainment****Science/Nature****Technology****Health****Talking Point****Country Profiles In Depth****Programmes****COMMONWEALTH GAMES****BBC SPORT****BBC WEATHER****SERVICES****Daily E-mail****News Ticker****Mobile/PDAs****Text Only****Feedback****Help****EDITIONS****Change to UK****Boeing tries to defy gravity**

An anti-gravity device would revolutionise air travel

Researchers at the world's largest aircraft maker, Boeing, are using the work of a controversial Russian scientist to try to create a device that will defy gravity.

The company is examining an experiment by Yevgeny Podkletnov, who claims to have developed a device which can shield objects from the Earth's pull.

Dr Podkletnov is viewed with suspicion by many conventional scientists. They have not been able to reproduce his results.

The project is being run by the top-secret Phantom Works in Seattle, the part of the company which handles Boeing's most sensitive programmes.

The head of the Phantom Works, George Mueller, told the security analysis journal Jane's Defence Weekly that the science appeared to be valid and plausible.

Dr Podkletnov claims to have countered the effects of gravity in an experiment at the Tampere University of Technology in Finland in 1992.

The scientist says he found that objects above a superconducting ceramic disc rotating over powerful electromagnets lost weight.

WATCH/LISTEN REAL MEDIA**ON THIS STORY**

[] The BBC's Andrew Gilligan

"The idea is still highly experimental"

[] Professor Robin Tucker and writer Nick Cook

"Boeing wants to build its own impulse gravity generator"

See also:

29 Jul 02 | Science/Nature

[Q&A: Boeing and anti-gravity](#)

27 Mar 00 | Science/Nature

[Gravity research gets off the ground](#)


29 Jul 02 | Media reports

[Russia's 'gravity-beating' scientist](#)**Internet links:****[Nasa Breakthrough Propulsion Physics Program](#)****[Quantum Cavorite](#)****[Project Greenglow](#)****[Boeing](#)****[Jane's](#)**

The BBC is not responsible for the content of external internet sites

Top Science/Nature stories now:**[Galaxy 'may cause ice ages'](#)****[Fresh doubt over America map](#)****[Into the eye of the hurricane](#)****[Hypersonic jet launch raises hopes](#)****[UN's 'risky' Earth Summit gambit](#)****[UK defends smallpox vaccine decision](#)****[Antimatter mystery deepens](#)****[Asteroid to miss - this time around](#)**

Links to more Science/Nature stories are at the foot of the page.



1. Solenoids create magnetic field
2. Spinning, super-conducting ceramic ring
3. Liquid Nitrogen acts as coolant
4. Dr Podkletnov claims weight can be reduced by 2% (1kg = 980g)

[Q&A: What chance floating jumbos?](#)

The reduction in gravity was small, about 2%, but the implications - for example, in terms of cutting the energy needed for a plane to fly - were immense.

Scientists who investigated Dr Podkletnov's work, however, said the experiment was fundamentally flawed and that negating gravity was impossible.

Research explored

But documents obtained by Jane's Defence Weekly and seen by the BBC show that Boeing is taking Dr Podkletnov's research seriously.

The hypothesis is being tested in a programme codenamed Project Grasp.

Boeing is the latest in a series of high-profile institutions trying to replicate Dr Podkletnov's experiment.

The military wing of the UK hi-tech group BAE Systems is working on an anti-gravity programme, dubbed Project Greenglow.

The US space agency, Nasa, is also attempting to reproduce Dr Podkletnov's findings, but a preliminary report indicates the effect does not exist.

 [E-mail this story to a friend](#)

Links to more Science/Nature stories

© BBC

[^^ Back to top](#)

[News Front Page](#) | [Africa](#) | [Americas](#) | [Asia-Pacific](#) | [Europe](#) | [Middle East](#) |
[South Asia](#) | [UK](#) | [Business](#) | [Entertainment](#) | [Science/Nature](#) |
[Technology](#) | [Health](#) | [Talking Point](#) | [Country Profiles](#) | [In Depth](#) |
[Programmes](#)

[To BBC Sport>>](#) | [To BBC Weather>>](#) | [To BBC World Service>>](#)

[© MMII](#) | [News Sources](#) | [Privacy](#)

You are in: [Science/Nature](#)

Monday, 29 July, 2002, 14:52 GMT 15:52 UK

News Front[Page](#)[World](#)[UK](#)[England](#)[N Ireland](#)[Scotland](#)[Wales](#)[UK Politics](#)[Business](#)[Entertainment](#)[Science/Nature](#)[Technology](#)[Health](#)[Education](#)[Talking Point](#)[Country](#)[Profiles](#)[In Depth](#)[Programmes](#)**COMMONWEALTH GAMES**[BBC SPORT](#)[BBC WEATHER](#)[CBBC news](#)**SERVICES**[Daily E-mail](#)[News Ticker](#)[Mobile/PDAs](#)[Text Only](#)[Feedback](#)[Help](#)**EDITIONS**[Change to](#)[World](#)**Q&A: Boeing and anti-gravity**

The aircraft manufacturer Boeing is reported to be investigating anti-gravity research. But our science correspondent Dr David Whitehouse says jumbo jets are unlikely to start floating off the ground any time soon.

What has fired the interest of Boeing?

According to company documents seen by the respected journal Jane's Defence Weekly, Boeing is intrigued by the Russian of Russian Dr Eugene Podkletnov. It wants to see if it really is possible to build an "impulse gravity generator".

**[Boeing tries to defy gravity](#)****What does Dr Podkletnov claim to have achieved?**

He carried out an experiment involving a supercold, spinning ceramic ring.

[Click here to see how an anti-gravity device could work](#)

An object held above the ring lost about 2% of its weight. The pull of gravity on the object was reduced, he claims.

When did Dr Podkletnov do this?

It seems that the first time was at a Finnish university in 1992. However, the research paper that described the results was submitted to a scientific journal but then withdrawn. The same happened in 1996 when news of the paper's imminent publication leaked. However, Dr Podkletnov has continued to work on his ideas and is reported to have seen similar effects in his studies.

Has the experiment been repeated by others?

Several teams have tried but none has seen the same thing. The American space agency Nasa tried but said it couldn't find any effect.

Is anti-gravity possible?

The vast majority of the world's qualified scientists would fervently say that it is not. If it were possible to shield something from the force of gravity it would mean a rewrite of the most fundamental, cherished, and rigorously tested laws of physics. Most would put it alongside a perpetual-motion machine. That is, impossible.

But George Muellner of Boeing has been quoted as saying that the science of the antigravity device appeared to be valid

[WATCH/LISTEN](#) [REAL MEDIA](#)**ON THIS STORY**

[The BBC's Andrew Gilligan](#)
"The idea is still highly experimental"

[Professor Robin Tucker and writer Nick Cook](#)
"Boeing wants to build its own impulse gravity generator"

See also:

29 Jul 02 | Science/Nature

[Boeing tries to defy gravity](#)

27 Mar 00 | Science/Nature

[Gravity research gets off the ground](#)**Internet links:**[Nasa Breakthrough Propulsion Physics Program](#)[Quantum Cavorite](#)[Project Greenglow](#)[Boeing](#)[Jane's](#)

The BBC is not responsible for the content of external internet sites

Top Science/Nature stories now:[Galaxy 'may cause ice ages'](#)[Fresh doubt over America map](#)[Into the eye of the hurricane](#)[Hypersonic jet launch raises hopes](#)[UN's 'risky' Earth Summit gambit](#)[UK defends smallpox vaccine decision](#)[Antimatter mystery deepens](#)[Asteroid to miss - this time around](#)

Links to more Science/Nature stories are at the foot of the page.

and possible. Why did he say that?

There are three possibilities. Perhaps he knows something the rest of the world's physicists don't, or he does not understand the laws of physics, which would be highly surprising for a man in his position, or he has been misquoted.

If it were possible what would be the outcome?

We could build spacecraft that would not need any rockets to power them. Aircraft would not need engines and could be any size and stay up indefinitely using no fuel. Supporters have claimed military and medical applications as well.


Have I heard this before?

If you read HG Wells, yes. In his First Men On The Moon, he writes of an eccentric inventor called James Cavor who creates a substance called Cavorite that repels gravity. Coating a capsule with it makes the vehicle fly up towards the Moon.

But is not history full of geniuses who changed the world after their colleagues had dismissed them as crazy?

There are not as many of these as you might think. Besides, just because a few geniuses have been laughed at in the past does not mean that every one who is laughed at today will turn out to be a genius in the future.

HOW AN ANTI-GRAVITY DEVICE COULD WORK



[**Click here to return**](#)

[**E-mail this story to a friend**](#)

[**Links to more Science/Nature stories**](#)

You are in: [Sci/Tech](#)[Front Page](#)

Monday, 27 March, 2000, 23:34 GMT 00:34 UK

[World](#)[UK](#)[UK Politics](#)[Business](#)[Sci/Tech](#)[Health](#)[Education](#)[Entertainment](#)[Talking Point](#)[In Depth](#)[AudioVideo](#)

Such devices would shield planes from the Earth's pull

A leading UK company is challenging what we understand to be the fundamental laws of physics.

The military wing of the hi-tech group BAe Systems, formerly British Aerospace, has confirmed it has launched an anti-gravity research programme.

It hopes that Project Greenglow will draw scientists from different backgrounds to work on future technologies that will have echoes of the propellantless propulsion systems being investigated by Nasa's Breakthrough Propulsion Physics Program.

Gravitation shielding

If any of the work is successful, it could lead to dramatic developments in the way we travel - anti-gravity devices could make it much easier for aeroplanes, spacecraft and even the next generation of cars to get off the ground.

In 1996, the experiments of a Russian scientist were jeered at by the physics world. Writing in the journal *Physica C*, Dr Yevgeny Podkletnov claimed that a spinning, superconducting disc lost some of its weight. And, in an unpublished paper on the weak gravitation shielding properties of a superconductor, he argued that such a disc lost as much as 2% of its weight.

However, most scientists believe that such anti-gravity research is fundamentally flawed. It goes against what we know about the physical Universe and is therefore impossible, they say.

Pascal's Wager

"I find it rather peculiar that they've done this," said Bob Park

[Search BBC News Online](#)[Advanced search options](#)**Launch console
for latest audio/video**[BBC RADIO NEWS](#)[BBC ONE TV NEWS](#)[WORLD NEWS SUMMARY](#)[BBC NEWS 24 BULLETIN](#)[PROGRAMMES GUIDE](#)

Internet links:

[Nasa Breakthrough Propulsion Physics Program](#)[Electrogravity](#)[Project Greenglow](#)

The BBC is not responsible for the content of external internet sites

Links to other Sci/Tech stories are at the foot of the page.

from the American Physical Society, in reaction to the BAe Systems admission. "One can only conclude that at the higher levels of these organisations there are people who don't have a very sound grounding in fundamental physics.

"You can invest a little money in far-out projects if they have some chance of success - it's called Pascal's Wager. In this case, most scientists would say there is zero chance of success."

Nonetheless, this view will not stop anti-gravity devices from continuing to be a popular feature of science fiction and the inspiration for countless websites.

 [E-mail this story to a friend](#)

Links to more Sci/Tech stories



^^ [Back to top](#)

[News Front Page](#) | [World](#) | [UK](#) | [UK Politics](#) | [Business](#) | [Sci/Tech](#) | [Health](#) | [Education](#) |
[Entertainment](#) | [Talking Point](#) | [In Depth](#) | [AudioVideo](#)

[To BBC Sport>>](#) | [To BBC Weather>>](#)

[© MMII](#) | [News Sources](#) | [Privacy](#)

BBC NEWS

You are in: [World: Europe](#)[Front Page](#)

Monday, 27 March, 2000, 13:58 GMT 14:58 UK

[World](#)

British probe anti-gravity device

A British defence contractor, BAe Systems, has confirmed that it is funding research into a device which is said to modify the effects of gravity.

The concept was pioneered by a Russian physicist, Yevgeny Podkletnov, working in Finland, and employs electromagnets and a rotating disc.

Dr Podkletnov has said experiments showed that it eased gravitational force by two per cent or more, which could have a dramatic effect on means of travelling.

BAe Systems says it will call its anti-gravity research programme Project Greenglow.

However, the BBC Science Correspondent says the possibility that a basic law of physics can be altered has attracted ridicule from many scientists, who believe that the concept belongs in the realm of science fiction.

[From Our Own Correspondent](#)

[Letter From](#)

[America](#)

[UK](#)

[UK Politics](#)

[Business](#)

[Sci/Tech](#)

[Health](#)

[Education](#)

[Entertainment](#)

[Talking Point](#)

[In Depth](#)

[AudioVideo](#)

From the newsroom of the BBC World Service

 [E-mail this story to a friend](#)

[Links to more Europe stories](#)

[Search BBC News Online](#)

[Advanced search options](#)

**Launch console
for latest audio/video**



[BBC RADIO NEWS](#)

[BBC ONE TV NEWS](#)

[WORLD NEWS SUMMARY](#)

[BBC NEWS 24 BULLETIN](#)

[PROGRAMMES GUIDE](#)

[Europe Contents](#)

[Country profiles](#)

Links to other Europe stories are at the foot of the page.

[Front Page](#)

Friday, February 27, 1998 Published at 04:58 GMT

Despatches

Nigel Margerison
London

An international team of astronomers say they have found evidence that the universe is expanding at an accelerating rate which can only be explained by the existence of anti-gravity. Their findings are published today in the journal, Science. As the BBC's Nigel Margerison reports, if true, the data would challenge the current thinking on the history of space and time.

The accepted view up to now is that the birth of the Universe in the Big Bang catapulted matter outwards and the Universe has been expanding ever since. But physics decrees that the force of gravity will cause the acceleration to slow.

This new research shows that instead of slowing, the expansion is speeding up and that, say the scientists, can only be explained by anti-gravity, a force opposing the effects of gravity. The findings are based on data gathered from observatories around the world and the Hubble Space telescope.

Some of the most distant objects supernovae or exploding stars, up to 10bn light years from earth, were observed over a period of time and their speed of acceleration measured. Albert Einstein first proposed the concept of anti-gravity but later discarded the idea.

Even the scientists themselves can't believe their results saying they have looked for an error but so far can't find one. If no error is found scientists say the findings will mean the Big Bang happened longer ago than we thought and would mean that in billions of years time many of the stars we now see would have accelerated out of range, making the universe look a very lonely place.

In this section

[Historic day for East Timor](#)[Despatches Contents](#)

Relevant Stories

27 Feb 98 | Sci/Tech
[Universe's expansion speeds up](#)

Internet Links

[Einstein revealed](#)[Einstein: Image and Impact](#)[Overview of the Hubble Space Telescope](#)[Journal of Science](#)


The BBC is not responsible for the content of external internet sites.

[Front Page](#)[UK](#)[World](#)[Business](#)[Sci/Tech](#)[Sport](#)[Despatches](#)[World Summary](#)[On Air](#)[粵語廣播](#)[Talking Point](#)[Feedback](#)[Low Graphics](#)[Help](#)[Site Map](#)

Friday, February 27, 1998 Published at 11:13 GMT

Sci/Tech


Universe's expansion speeds up



Hubble pinpoints distant supernovae

An international team of scientists has found evidence of a mysterious anti-gravity force that is causing the universe to expand at an accelerating rate.

The finding supports a concept first proposed by Albert Einstein, who later discarded the idea and called it his biggest blunder.



Einstein discarded his theory of anti-gravity

"It is such a strange result we are still wondering if there is some other sneaky little effect climbing in there," said Adam Riess, an astronomer at the University of California, Berkeley.

The 15-member team that made the discovery "have looked hard for errors," but found none, he said.

If true, the discovery could challenge the current thinking on the history of space and time.

Galaxies far, far away

In this section

[World's smallest transistor](#)[Scientists join forces to study Arctic ozone](#)[Mathematicians crack big puzzle](#)

From Business
[The growing threat of internet fraud](#)

Who watches the pilots?

From Health
[Cold 'cure' comes one step closer](#)

Sci/Tech Contents


Internet Links

[Einstein revealed](#)[Einstein: Image and Impact](#)[Overview of the Hubble Telescope](#)[Journal of Science](#)

The BBC is not responsible for the content of external internet sites.

Scientists say they discovered the force using the Hubble Space Telescope and ground-based telescopes in Hawaii, Australia and Chile while analysing the light arriving from 14 supernovae, or exploding stars, that are seven billion to 10 billion light-years from Earth.

A light-year is the distance that light travels in one year - about six trillion miles.



1997 photo of a supernova

1997 photo of a supernova said. "It will be very lonely."

The scientists expected to find that the expansion of the universe was slowing slightly from the effect of gravity. Instead, they say, it is actually speeding up.

One of the scientists involved, Robert Kirshner said the acceleration would continue and within billions of years many of the stars now seen would disappear from view.

"The universe will be a very different place to look at," he

Questions answered

If the speed of expansion of the universe is accelerating, it could solve one problem for astronomers.

Some measurements have put the age of the universe at about 10 billion years. This is younger than the measured ages of some stars, a dilemma that has confounded astronomers.

With the acceleration of the universe factored in, the universe would have to be about 14 billion years old, some two billion years older than the oldest star, according to UC Berkeley's Mr Riess.

The team's conclusion will go through an intensive review by many astronomers before the results are accepted, according to Mr Kirshner. He noted however that preliminary results from a parallel study by another astronomy group agreed with the findings.

You are in: [Monitoring: Media reports](#)

Monday, 29 July, 2002, 20:53 GMT 21:53 UK

News Front[Page](#)[World](#)[UK](#)[England](#)[N Ireland](#)[Scotland](#)[Wales](#)[UK Politics](#)[Business](#)[Entertainment](#)[Science/Nature](#)[Technology](#)[Health](#)[Education](#)[Talking Point](#)[Country Profiles](#)[In Depth](#)[Programmes](#)**COMMONWEALTH GAMES**[BBC SPORT](#)[BBC WEATHER](#)[CBBC news](#)**SERVICES**[Daily E-mail](#)[News Ticker](#)[Mobile/PDAs](#)[Text Only](#)[Feedback](#)[Help](#)**EDITIONS**[Change to](#)[World](#)

Are anti-gravity devices science fiction or fact?

Aircraft manufacturer Boeing's work on an experimental anti-gravity device is based on a concept pioneered by Russian physicist Yevgeny Podkletnov.

BBC Monitoring looks back at his long and often controversial career.

[Country Profiles](#)[In Depth](#)[Programmes](#)

Yevgeny Podkletnov was born into a highly educated family in the Soviet Union in the mid-1950s.

His father was a scientist and a professor in St Petersburg, while his mother had previously conducted research in the field of medicine.

Dr Podkletnov himself followed his father into a career in science, first earning a degree from the Mendeleev Institute in Moscow, and then moving on to work at an institute within the Russian Academy of Sciences.

In the late 1980s, Dr Podkletnov left for Finland to pursue his research work at the Tampere University of Technology in Finland.

He first came to prominence in 1992, when he claimed to have conducted a successful experiment into what he called "gravity shielding".

Scepticism

Other scientists however, were initially unable to reproduce the results of Dr Podkletnov's experiment in their own laboratories, and the Russian scientist's research was viewed with considerable scepticism.

Undeterred, Dr Podkletnov submitted a paper on the research to the Institute for Physics in London. It was accepted for print, and was to be published in October 1996.

But, a month before that, news of his research entered into the wider public domain, when the Sunday Telegraph published an article highlighting the great potential for "anti-gravitation" technology.

The material for the article, which had been leaked, provoked a

See also:[29 Jul 02 | Science/Nature](#)[Q&A: Boeing and anti-gravity](#)[27 Mar 00 | Science/Nature](#)[Gravity research gets off the ground](#)[27 Mar 00 | Europe](#)[British probe anti-gravity device](#)**Internet links:**[NASA](#)[Quantum Cavorite: Podkletnov](#)[BAe Systems](#)[Boeing Corporation](#)

The BBC is not responsible for the content of external internet sites

Top Media reports stories now:[Ukrainian, Russian papers assess plane crashes](#)[Romanian Nobel winner returns home](#)[Russian air crash survivor tells story](#)[Spain press scorns Gibraltar vote](#)[Russia's 'gravity-beating' scientist](#)[Abdul Kalam urges drive against poverty](#)[Kursk mothers reject disaster report](#)[Israeli press uneasy over Gaza attack](#)

Links to more Media reports stories are at the foot of the page.

broadly negative reaction among the scientific community at large, and among Dr Podkletnov's research colleagues.

A week after the article appeared, Dr Podkletnov asked the Institute of Physics to withdraw his paper from publication.

Despite these problems however, major players have continued to maintain an interest in Dr Podkletnov's work.



Only in space are the effects of gravity ameliorated

In 2000, the British defence contractor BAe Systems confirmed that it would be funding research into a device said to modify the effects of gravity. And now Boeing, the world's largest aircraft manufacturer, is also using Dr Podkletnov's research in its own programmes.

BBC Monitoring, based in Caversham in southern England, selects and translates information from radio, television, press, news agencies and the Internet from 150 countries in more than 70 languages.

 [E-mail this story to a friend](#)

[Links to more Media reports stories](#)

© BBC

[^ Back to top](#)

[News Front Page](#) | [World](#) | [UK](#) | [England](#) | [N Ireland](#) | [Scotland](#) | [Wales](#) |
[UK Politics](#) | [Business](#) | [Entertainment](#) | [Science/Nature](#) | [Technology](#) |
[Health](#) | [Education](#) | [Talking Point](#) | [Country Profiles](#) | [In Depth](#) |
[Programmes](#)

[To BBC Sport>>](#) | [To BBC Weather>>](#) | [To BBC World Service>>](#)


[© MMII](#) | [News Sources](#) | [Privacy](#)

[News in Audio](#)[News in Video](#)[Newyddion](#)[Новости](#)[Noticias](#)[أخبار](#)[国际新闻](#)[粵語廣播](#)[Front Page](#)[World](#)[UK](#)[UK Politics](#)[Business](#)[Sci/Tech](#)[Health](#)[Education](#)[Sport](#)[Entertainment](#)[Talking Point](#)[On Air](#)[Feedback](#)[Low Graphics](#)[Help](#)

Tuesday, October 27, 1998 Published at 14:08 GMT

Sci/Tech

Hubble v Big Bang



Through a lens clearly - multiple images from across the universe

By our Science Editor David Whitehouse

New observations made by the Hubble space telescope suggest that some of our ideas about the Big Bang, the explosion that started the universe, may need revising.

A team from the University of Arizona has obtained a Hubble image of a distant galaxy, designated PG 115+080, seen through a much nearer galaxy.

PG 115+080 is eight billion light years away. The closer galaxy, three billion light years distant, acts as a 'gravitational lens' bending light so that the picture we see of PG 115+080 consists of several distorted images.

This is useful to astronomers because it gives them the chance to do some simple geometry to measure distances across the universe based on the path that the light in each image has travelled

The good news is that using the new Hubble image, it is possible to measure how fast the universe is expanding. The new figure agrees with previous estimates.

The bad news is that the big bang cannot easily explain this observed rate of expansion with the matter the cosmos seems to contain.

So it seems that the Big Bang theory needs patching up. There are three possible ways to do this but each adjustment is unpalatable to astronomers.

- They may have to change what they thought were the conditions in the big bang

[Sci/Tech Contents](#)

Relevant Stories

28 Oct 98 | Sci/Tech
[Looking back 12 billion years](#)

Internet Links

[Hubble observations](#)
[University of Arizona](#)

The BBC is not responsible for the content of external internet sites.

In this section

[World's smallest transistor](#)
[Scientists join forces to study Arctic ozone](#)
[Mathematicians crack big puzzle](#)
 From Business
[The growing threat of internet fraud](#)
[Who watches the pilots?](#)
 From Health
[Cold 'cure' comes one step closer](#)

- They may have to postulate new, so far undiscovered, types of matter
- Or they may have to bring back an idea that Albert Einstein discarded.

Earlier this century, before it was discovered that the universe was expanding, Einstein tried to explain why it did not collapse. He came up with an anti-gravity force that holds it up.

When it was discovered in the 1920's that the cosmos expands, Einstein's new force was dropped. However some astronomers, while still not liking the idea, point out that it could come in useful today to explain a few things about the universe.

Despite these problems it is clear that the fundamental basis of the Big Bang theory, that the universe began in an explosion of time and space about 13 billion years ago, is not in dispute.

It is just that we do not understand what came next as well as we thought we did.

[Advanced options](#) | [Search tips](#)

[2 Images](#)

HEADLINES

Date Posted: July 29, 2002

JANE'S DEFENCE WEEKLY - JULY 31, 2002

Anti-gravity propulsion comes 'out of the closet'

NICK COOK *JDW Aerospace Consultant*
London

Boeing, the world's largest aircraft manufacturer, has admitted that it is working on experimental **anti-gravity** projects that could overturn a century of conventional aerospace propulsion technology if the science that underpins them - science that senior Boeing officials describe as "valid" - can be engineered into hardware.

As part of the effort, which is being run out of Boeing's **Phantom** Works advanced research and development facility in Seattle, the company is trying to solicit the services of a Russian scientist who claims he has developed 'high-' and 'low-power' **anti-gravity** devices in **Russia** and **Finland**. The approach, however, has been thwarted by Russian officialdom.

The Boeing drive to develop a collaborative relationship with the scientist in question, Dr Evgeny Podkletnov, has its own internal project name: 'GRASP' - **Gravity** Research for Advanced Space Propulsion.

A briefing document on GRASP obtained by *Jane's Defence Weekly* sets out what Boeing believes to be at stake. "If gravity modification is real," it says, "it will alter the entire aerospace business." The report was written by Jamie Childress, principal investigator for Boeing's propellantless propulsion work at the **Phantom** Works in Seattle.

GRASP's objective is to explore propellantless propulsion (the aerospace world's more formal term for **anti-gravity**), determine the validity of Podkletnov's work and "examine possible uses for such a technology". Applications, the company says, could include space launch systems, artificial gravity on spacecraft, aircraft propulsion and 'fuelless' electricity generation - so-called 'free energy'.

But it is also apparent that Podkletnov's work could be engineered into a radical form of weapon system. The GRASP paper focuses on Podkletnov's claims that his high-power experiments, using a device called an 'impulse gravity generator', are capable of producing a beam of 'gravity-like' energy that can exert an instantaneous force of 1,000g on any object - enough, in principle, to vaporise it, especially if the object is moving at high speed.

Podkletnov maintains that a laboratory installation in [Russia](#) has already demonstrated the 4in (10.16cm) wide beam's ability to repel objects a kilometre away and that it exhibits negligible power loss at distances of up to 200km (*JDW* 24 July). Such a device, observers say, could be adapted for use as an anti-satellite weapon or a ballistic missile shield.

The GRASP paper details the beam's reported characteristics: that it is immune to electromagnetic shielding, that it can penetrate any intermediate barriers (objects placed between the generator and the target), that it propagates at very high speed ("possibly light speed or greater") and that the total force is proportional to target mass - that its effect, in other words, is exactly the same as gravity's.

Podkletnov's claims first surfaced in 1992 when he published a paper detailing his low-power experiments into gravity-shielding using superconductors, materials that lose their electrical resistance at low temperatures. The original experiments were conducted at the University of Technology in Tampere, [Finland](#), before moving to [Russia](#).

Podkletnov, who has a PhD in materials science from Tampere and the University of Chemical Technology in Moscow, declared that any object placed above his rapidly spinning superconducting apparatus lost up to 2% of its weight.

Although he was vilified by traditionalists who claimed that gravity-shielding was impossible under the known laws of physics, the US National Aeronautics and Space Administration (NASA) went on to attempt a replication of his work in the mid-1990s.

Because NASA lacked Podkletnov's unique formula for the 30cm yttrium-barium copper oxide (YBCO) superconducting ceramic discs - a formula the Russian maintains is critical to the experiment's success - the attempt failed. NASA's Marshall Space Flight Center in Alabama will shortly conduct a second set of experiments, this time using apparatus built to Podkletnov's specifications.

In August 2001, Podkletnov published a paper revealing his experimental high-power work and it is this that forms the focus of the GRASP report. Boeing wants to build its own impulse gravity generator at Seattle but admits that it lacks vital knowledge in the area of the YBCO emitter - Podkletnov's special superconducting apparatus - which forms the heart of the generator.

As a result, Boeing recently approached Podkletnov directly, but promptly fell foul of Russian technology transfer controls. George Muellner, the outgoing head of the Boeing [Phantom](#) Works, confirmed that attempts by Boeing to work with Podkletnov had been blocked by Moscow, which is seeking to stem the exodus of Russian high-technology to the West. Muellner is convinced, however, that the science underpinning Podkletnov's work is real.

"The physical principles - and Podkletnov's device is not the only one - appear to be valid," he said. He confirmed that Boeing had conducted tests on a number of other **anti-gravity** devices, some of which


were detailed in *JDW* 24 July.

"There is basic science there. They're not breaking the laws of physics. The issue is whether the science can be engineered into something workable," Muellner said.

The GRASP briefing document reveals that [BAE Systems](#) and Lockheed Martin have also contacted Podkletnov "and have some activity in this area".

It is also possible, Boeing admits, that "classified activities in gravity modification may exist". The paper points out that Podkletnov is strongly anti-military and will only provide assistance if the research is carried out in the 'white world' of open development.

GRASP concludes that a "positive result from experiments would give Boeing a substantial advantage in the aerospace industry".



Reported operation of high-power gravity device: the YBCO emitter is cooled to superconducting conditions and the inner electric coil produces a magnetic field inside it. A very high voltage pulse is input to the emitter which then discharges to the collector. The discharge is guided and contained by the magnetic field in the outer coil

(Source: Jane's/Boeing)





Diagram showing configuration of low-power gravity modification experiment
(Source: Jane's/Boeing)

© 2002 Jane's Information Group


© Jane's Information Group 2002

[Terms of Use](#)

[Powered by Verity](#)



Gravity experiment apparatus



[7 Images](#)

FEATURE

Date Posted: July 18, 2002

JANE'S DEFENCE WEEKLY - JULY 24, 2002

Exotic propulsion: Air power ELECTRIC

NICK COOK *JDW Aerospace Consultant
London*

Intriguing, perplexing, confusing - projects in the field of '**anti-gravity**' propulsion may seem the stuff of science-fiction but are very much the science-fact of today. Nick Cook looks at the research - and finds some high profile companies involved.

In July 2001, officials from a small Alabama-based research company called Transdimensional Technologies (TDT) - a concern with strong links to the National Aeronautics and Space Administration (NASA) - lifted the lid on a set of experiments they boldly claimed would "change virtually every aspect of our daily lives." Under laboratory conditions, TDT had just tested a tethered subscale model of a device it called the 'Lifter' that had demonstrated its ability to 'fly' using a propulsion methodology that science is grappling to explain.

The Lifter is the brainchild of TDT's chief scientist, Jeff Cameron, an electro-optics engineer by training, who had come by the idea after noticing some anomalous effects during his experiments with lasers. Sensing that the twisting motion of a foil and wire component within a part of the laser called a pre-ioniser was attributable to some unknown force, Cameron adapted the pre-ionising apparatus into a 3-D triangular shape with foil around its edges and a thin wire supported by balsa wood poles running above its upper surface. When positive and negative charges were run through the wire and the foil, the model Lifter levitated off the bench, making a crackling noise and spraying a colourful corona discharge.

By treating the triangular shape as a 'cell', TDT began scaling its Lifter, using increasing numbers of cells to make bigger and heavier 'craft'. Lifters have no moving parts and do not need to be aerodynamic in shape to get off the ground.

Soon after TDT published its findings, a French physicist, Jean-Louis Naudin, replicated the US company's experiments and quantified the lift produced. The biggest craft, the Lifter 4, used 36 cells joined together to form a basic hexagonal shape 820mm by 700mm in dimension. Constructed using the same materials as the TDT Lifters - aluminium foil, balsa wood and wire - Naudin's Lifter Version 4 weighed 32g and needed approximately 81W of power (40 kV at 2.01 mA) to levitate. To maintain stability in flight, it required 132.9W (44 kV at 3.02 mA). "The Lifter v4.0 is able to accelerate upwards very quickly and silently," Naudin reported, "and it is very stable during hovering." When word spread on the Internet of the TDT and Naudin tests, dozens of amateur engineers got in on the act. Using materials that can be found in hardware stores, they have consistently demonstrated that their lifters can and do 'fly' in and out of the laboratory.

It would be difficult for anyone to dispute the reality of lifters - films and still photos depict a plethora of lifter designs 'flying' on multiple websites. The \$64,000 dollar question is: why are they flying at all? Some believe that their lift derives from an effect known as 'ion wind' - a 'breeze' caused by the movement of electrons crossing the gap between the wire and the foil. In 1964, Alexander De Seversky, founder of the Republic aviation company, patented a lifter-like design called the 'ionocraft' that relied on just such an effect for its postulated lift - but no such craft was ever built.

Although lifters can be steered by varying the power to the cells, the models are so light that if they are flown in the open they get blown around, even in a light breeze. What is needed, according to Graham Ennis, who organised the First International Field Propulsion Conference in Brighton, UK, last year, is for the effect to be tested on a 10m craft with a much higher voltage throughput - 10 or 20 times higher than the 20-45 kV typically applied to date. "The actual thrust is not linear. It increases by the square of the voltage. If you double the voltage you'll get four times as much thrust. To be effective, lifters not only need more voltages, but something hitherto not done: a very lightweight, very high voltage power supply," Ennis says.

TDT's Cameron admits that ion wind is generated by his company's Lifters, but it is far too slight, he says, to explain the levitational effect. If an obstruction is placed between the wire and the foil, thereby blocking the flow of electrons, the Lifter continues to levitate, he maintains. "We know enough about ion wind to determine how much is created and what it can push. There is insufficient ion wind to lift this test article." Tim Ventura, a Seattle-based lifter-builder, who has studied the phenomenon and the technology at length, is also sceptical that ion wind explains a lifter's ability to fly. "I think that ion wind is a component, but it's not the whole answer," he says. The question, however, remains. Is the lifter effect a laboratory curiosity or does it have some practical utility to aerospace development?

The TDT work has sparked further controversy via claims that it is not a new effect at all but a reworking of a phenomenon discovered by a US inventor called Thomas Townsend Brown in the late 1920s. Brown postulated a novel form of lift based on the assertion that an 'asymmetric capacitor' - a disc-shaped plate, slightly domed on one side, capable of retaining a large electrical charge - would experience thrust in the direction of its positive pole when charged negatively on one side and positively on the other. Intentionally or not, Cameron's Lifters are actually three dimensional representations of Brown's asymmetric plates, only on the lifters it is the air between the wire and the foil that acts as the capacitor.

In 1952, Brown went on to submit a proposal for a 'joint services' technology demonstration

programme, 'Project Winterhaven', that was designed to lead to a M3 disc-shaped interceptor powered via his electro-kinetic propulsion principles. A series of tests demonstrated to the US Air Force in the mid-1950s, in which 3ft diameter disc-shaped capacitors were supposedly charged at 150 kV, were said by some contemporary sources to have exhibited results "so impressive as to be highly classified." Brown also fell victim to the claim that the 'Biefeld-Brown effect' was attributable to ion wind - a claim he sought to refute by demonstrating the ability of his capacitors to levitate in a vacuum. According to Ventura, lifters have also performed successfully under vacuum conditions. But ion wind or not - and, in a sense, proponents say, what does it matter, if it leads to an entirely new form of aero-propulsion - the lifter debate and Brown's work are attracting the attention of some high-profile organisations. Indeed, there are signs that after years on the shelf as a taboo science, electrogravitics - antigravity by another name - is under the microscope of government agencies and aerospace companies as they seek to find breakthrough propulsion technologies that could catapult aviation into a new era. The US Congress recently voted to give \$4.75 million of US taxpayers money to the West Virginia-based Institute of Software Research (ISR) to see if its scientists could independently verify the bold claims being made for the lifters.

Earlier this year, NASA claimed it had discovered a "method for generating thrust from two dimensional asymmetrical capacitor modules" and filed US Patent No. 6,317,310 to register its 'breakthrough'. The space agency's observation that implementation of the patent would result in "potentially greater efficiencies and improved reliability over currently available electric thrusters" points to its manifest interest in the 'modules' as potential orbital transfer engines for satellites or other kinds of space vehicle. But in lodging the patent, NASA opened itself to a barrage of criticism from electrogravitic aficionados that it is patenting a discovery already made - and patented - by Brown a half century earlier. What is significant, however, is NASA's open acknowledgment of a form of thrust generation that neither it nor anyone else can adequately explain by known scientific principles.

NASA's quest for leap-ahead propulsion technologies was launched in 1996 under the Breakthrough Propulsion Physics (BPP) programme, run out of the NASA Glenn Research Center in Cleveland, Ohio. BPP is a subset of NASA's Advanced Space Transportation Plan (ASTP) managed by the NASA Marshall Space Flight Center in Huntsville, Alabama. ASTP's wide-ranging remit was to scout out bold-strokes technologies designed to lead to visionary forms of space propulsion that in the near term could reduce the cost of space access and within 100 years or so, as the project evolved, lead to true quantum leap propulsion advances capable perhaps of taking man to the stars. Under BPP, NASA Glenn was handed a remit to fund research into emerging principles that might lead to forms of propulsion hitherto regarded as impossible.

For example, it has been looking at ways of overcoming the supposedly unbreachable light-speed barrier as well as the manipulation of poorly understood physical forces such as gravity and inertia.

Despite its public appeal and relative low-cost - a total budget over seven years of \$1.7 million - BPP was notified at the end of June that its funds were being cut and diverted into nearer-term NASA space-launch and propulsion research. Although BPP may still be resurrected - attempts are being made to 'park' it into a non-profit institute that could be configured to receive funds from other government agencies or industry sponsors - the outlook, according to officials close to the programme, is not good. BPP, they say, has ended up a "collateral casualty" of fallout from an investigation by the Office of Management and Budget (OMB) into what OMB has perceived as overlapping research by NASA Glenn, NASA Marshall and the ISR in 'breakthrough' areas of propulsion physics.

Among other criticisms, according to Marc Millis, who founded the BPP programme, OMB objected to the amount of funding that had been devoted to the testing of claims that gravity may be 'shielded' to

produce a levitation effect. This task, though related to the BPP effort, was actually pursued as an in-house effort by NASA Marshall, which from 1996 has been seeking to verify claims by a Russian materials scientist that objects can lose their weight by up to 5% when placed above superconducting ceramic discs rotated at high speed. The scientist, Dr Evgeny Podkletnov, discovered the effect while working at the University of Tampere, [Finland](#), in the early 1990s. When he published his claims, however, he was vilified by traditionalists who said that 'gravity shielding' was impossible.

But Podkletnov has demonstrated to the satisfaction of many that some kind of weight reduction effect is taking place and with its obvious potential utility in reducing launch weights - and therefore launch costs - NASA Marshall quickly registered its interest. It had already been approached by a researcher at the University of Alabama in Huntsville, Ning Li, who independently claimed that she could produce a 'gravity-like field' capable of repelling or attracting matter using rapidly spinning superconductors, materials that lose their electrical resistance at low temperatures. In the mid-1990s, NASA Marshall unsuccessfully tried to replicate Podkletnov's experiments, but without the Russian scientist's unique formula for his 30cm yttrium-barium copper oxide superconducting discs, the agency admitted it was largely operating in the dark. In 1999, it paid \$600,000 to a Columbus, Ohio, company called Superconductive Components (SCI) to construct discs like the ones Podkletnov had been using in [Russia](#) and [Finland](#) and Podkletnov was hired as a consultant. After some delay, it is expected shortly to take delivery of SCI's superconducting apparatus. Ron Koczor, who heads up the programme at NASA, is confident that the experiment will go ahead, despite the OMB's investigation into what it sees as wasteful and unnecessary duplication of US government-funded research.

Li and Podkletnov have continued with their work. While Li has gone to ground, apparently to produce a workable derivative of her 'AC gravity machine' - which, she says, will be able to produce a beamed, 'gravity-like' force in any direction - Podkletnov, now based at the Moscow Chemical Scientific Research Centre, has taken his ideas further, publishing claims - backed by an Italian physicist, Giovanni Modanese of the California Institute for Physics and Astrophysics and the University of Bolzano, northern Italy - that he has developed what he refers to as an 'impulse gravity generator' capable of exerting a repulsive force on all matter.

Using a two- to 10-million Volt power supply developed recently in [Russia](#) - it is, significantly, Podkletnov says, no bigger than a fridge - a strong electrical discharge source and a rapidly rotating superconductor, Podkletnov's equipment has produced a 'gravity impulse', he claims, "that is very short in time and propagates with great speed (practically instantaneously) along the line of discharge passing through different objects without any observable loss of energy".

The result, he maintains, is a repulsive action on any object the beam hits that is proportional to its mass. Using a laser pointer, Podkletnov says that his 'laboratory installation' has demonstrated its ability to knock over objects more than a kilometre away.

"Our theoretical studies give grounds to claims that the gravity impulse will have negligible power loss at a distance of up to 200km when propagating through air and space," he adds. He is already proposing practical applications of the device. "Even our laboratory-scale installation, when combined with a precise modern radar system gives us a possibility to change the trajectories of satellites". While there are no independent verifications of Podkletnov's claims, the fact that his work is being taken seriously by NASA, at least, should lend some credence to them, proponents say.


Podkletnov says that his impulse gravity generator could be engineered into a radical new form of aero-engine, but it is also apparent that it could be developed into a new form of weapon - one that could knock satellites from their orbits or ballistic missiles from their impact trajectories. *Jane's*

Defence Weekly understands that one US aerospace company - Boeing - could already be working on a derivative of Podkletnov's impulse generator. Boeing admits that its [Phantom](#) Works advanced projects division "is aware of Podkletnov's work on an impulse gravity generator and would be interested in seeing further development work being done."


The company, however, may have already taken steps towards assembling a device - possibly under classified auspices. Informed sources maintain that it has proposed a 'gravity modification experimental setup' using a superconducting disc housed in a stainless steel vacuum chamber and linked to a 3 mV Van de Graaff generator. The sources told *JDW*: "Some preliminary work has been done by Boeing in assembling the electrical part of the 'gravity gun', but the company still has a problem with the superconducting emitter." The claims and the company's response to them show, if nothing else, that Boeing, like NASA, is engaging in new and little-understood areas of physics in attempts to develop a fundamental understanding of phenomena that could lead to breakthroughs.

Boeing denies that it is funding "any activities" in this area, but it declined to be interviewed on any aspect of its interest in Podkletnov's work - or its other '**anti-gravity**' type research projects. One of these was well documented in a publication called 'Infinite Energy', which reported that the [Phantom](#) Works ran tests in late 1999 of a 'two-rotor reactionless drive unit' built by a California-based inventor called Robert Cook. The tests were led by Jamie Childress, 'principal investigator' in this field for the [Phantom](#) Works in Seattle. On 29 November, 1999, the magazine reported, the reactionless drive unit "proved itself by accelerating a series of 5lb weights from zero to over 1,000ft per minute in a fraction of a second and then decelerating them without producing the 90 to 100lb negative reaction force required by Newton's Third Law of action-reaction." The Cook Inertial Propulsion (CIP) engine uses a pair of counter-rotating arms spinning about a common axle to convert centrifugal force into linear force. By adding additional rotors, Cook says that he can markedly increase the thrust.

The reason for Boeing's interest in the CIP and Podkletnov's impulse gravity work remains unclear, but it is apparent, just as it was in the mid-1950s - when Convair, Martin and other US aerospace companies revealed their deep involvement in gravity-led research programmes, then fell silent on the subject - that a quest for radically alternative propulsion methodologies is firmly rooted in its agenda.



TDT's basic, single triangular cell Lifter photographed using light-enhanced photography shows the device's electrical corona - a side-effect of high-voltage charge transfer
(Source: American Antigravity)



TDT's third Lifter measures 1m on each side and generates lift through four triangular cells
(Source: American Antigravity)



Close-up of Lifter 3 in flight at TDT's laboratory in Alabama
(Source: American Antigravity)



*The Cook Inertial Propulsion unit being prepared for a strain-gauge test at Boeing's facility in Seattle
(Source: Boeing)*



*Californian inventor Robert Cook, right, pictured at the tests of his inertial propulsion system by Boeing
(Source: Boeing)*



*Boeing's stainless steel vacuum chamber proposed for the Podkletnov experiments
(Source: Boeing)*





*Boeing's Van de Graaff generator proposed for the Podkletnov experiments
(Source: Boeing)*


© 2002 Jane's Information Group


© Jane's Information Group 2002
[Terms of Use](#)


[Powered by Verity](#)
















[1 Image](#)

FEATURES

Date Posted: July 24, 2000

JANE'S DEFENCE WEEKLY - JULY 26, 2000

Electrogravitics: the positive approach

One area of research not included in the NASA Breakthrough Propulsion Physics study, but which is touted by some as having the potential to leap the current state of the art in aerospace technology, is 'electrogravitics'.

This is a postulated form of lift based on the controversial assertion that a capacitor charged positively on one side and negatively on the other will experience weight loss, exhibiting thrust in the direction of its positive pole.

In 1952, the man generally described as the 'father of electrogravitics', US inventor Thomas Townsend Brown, submitted a proposal for a 'joint services research and development contract' outlining an electrogravitics aerospace programme called 'Project Winterhaven'. This set out how the US military could go about constructing a Mach 3 manned interceptor utilising electrogravitics technology. Brown based his data on a series of demonstrations he gave to the US Navy immediately after the Second World War.


In these and subsequent tests, 2ft (61cm) diameter disc-shaped capacitors charged with 50 kilovolts and a continuous energy input of 50 watts are said to have achieved a speed of 17ft/s in a circular air course 20ft in diameter. A second series of tests, in which slightly larger discs were supposedly charged at 150 kilovolts, were rumoured in the mid-1950s to have exhibited results "so impressive as to be highly classified". This has led to claims by proponents of the technology that the US government has conducted a clandestine electrogravitics effort ever since under the auspices of a 'deep black' test programme.

Lending possible credence to that assertion is this photograph, published here for the first time. It is purportedly of a sub-scale electrogravitics experimental air vehicle mounted on a wind-tunnel test-stand somewhere in the USA. Although *Jane's Defence Weekly* cannot verify its origins, it appears to have been taken in the late 1950s or early 1960s. Intriguingly, Brown recommended the construction of a 10ft diameter disc charged at 500 kilovolts as part of 'Project Winterhaven'.

Officially, the US Air Force and Navy lost interest in electrogravitics when their analysis showed that the movement of Brown's capacitors was caused by 'ionic wind' - the displacement of ions around the surface of the charged plates. Brown later claimed to refute this by conducting the tests in a vacuum.

With one or two exceptions, aerospace companies have studiously ignored breakthrough propulsion concepts such as those proposed by Brown and Podkletnov. [BAE Systems](#) describes its 'Project Greenglow' initiative as a "speculative research programme" into esoteric propulsion concepts (such as **anti-gravity**) and the Boeing [Phantom](#) Works recently conducted tests on a 'reactionless drive' device that apparently contravenes Newton's Third Law of action-reaction. Boeing concluded that the device, which uses two spinning arms transferred through three-dimensions, did work, though not yet at the required impulses for endo-atmospheric propulsion or practical space flight.

Echoing Frank Whittle's early pioneering work on turbojets in the 1920s and 1930s, however, there are those who say it might just be the start of something with far-reaching consequences for the entire future of transportation.




*This late 1950s or early 1960s picture is said to be a sub-scale electrogravitics experimental air vehicle mounted on a wind-tunnel test stand
(Source: Jane's)*

© 2000 Jane's Information Group

© Jane's Information Group 2002

[Terms of Use](#)

[Powered by Verity](#)



A-27772

(c) Rear view at minimum test height.

Figure 1.- Concluded.

[Click for more...](#)[My Account](#)[Jane's Services](#)[Online Research](#)[Online Channels](#)[Data Search](#) | [Data Browse](#) | [Image Search](#) | [Web Search](#) | [News/Analysis](#)[5 Images](#)

FEATURES

Date Posted: July 24, 2000

JANE'S DEFENCE WEEKLY - JULY 26, 2000

WARP DRIVE: WHEN?

INTRODUCTION:

NASA recently awarded a set of contracts that aim to prove if propulsion concepts behind the warp-drive system of the Starship *Enterprise* may be closer to reality than we think. Nick Cook reports on the breakthrough science that may one day make current aerospace technology as relevant as the Steam Age

Nick Cook *JDW's Aviation Editor*
London

In three years' time it will be exactly a century since the Wright Brothers achieved sustained powered flight at Kitty Hawk, North Carolina. Compared to ships and trains, aviation is still a young mode of transport, but already some people are asking whether it has hit the buffers of scientific limits.

On both sides of the Atlantic, aerospace companies and government research centres spend millions of dollars annually to reduce airframe drag factors by increments of a percentage point or two.

In the propulsion field, an investment of hundreds of millions of dollars is required from one generation of engine to the next to reduce fuel consumption figures by fractions. In spaceflight, the rockets that loft satellites into orbit at a cost of \$5,000 per kilogram comprise the same basic technology as the [V2](#) ballistic missiles developed by Germany during the Second World War. To go beyond the [Space Shuttle Orbiter](#), the US government will spend billions more dollars developing a totally reusable launch vehicle, the VentureStar.

Yet, even VentureStar, which - if it ever goes ahead - will represent an enormous leap forward, will fall victim to the same laws of physics as any other rocket. At the moment of launch, fuel will account for almost 90% of its weight, and this simply to deliver a relatively small payload into low earth orbit. Brute force solutions to air and space flight require high energy levels to sustain them; and this - for aircraft as well as rockets - means heavy, expensive and potentially explosive fuel loads.

At the US National Aeronautics and Space Administration (NASA), a tiny research effort costing approximately half a million dollars has been set up to pose, rather than answer, a simple series of questions: does aerospace propulsion have to remain locked into this incremental improvement cycle? Or are there quantum leaps in science that can be tapped into to circumvent the evolutionary process - as was achieved with the development of the jet engine more than half a century ago?

The NASA effort is called the Breakthrough Propulsion Physics (BPP) programme and it is run out of the administration's Glenn Research Center at Cleveland, Ohio.

Although only peripherally seeking endo-atmospheric payoffs, as the big picture is mainly concerned with space travel, BPP does raise fundamental questions about the future of aerospace as a scientific and engineering discipline. That it is doing so now hinges on the fact that NASA officials believe there may be "emerging clues" to new propulsion methods, based on some recent "provocative developments", that could point the way to some startling advances.

BPP is part of the Advanced Space Transportation Plan (ASTP), managed by the NASA Marshall Space Flight Center in Alabama. ASTP is a long-range effort that seeks to develop new low-cost space launch and transfer technologies.

Its remit is also to come up with visionary ideas for future space transport systems - ideas that if ever translated into hardware might make future trips to other star systems possible.

BPP stemmed from some fundamental questions asked by NASA administrator Dan Goldin in 1996 about perceived physical limits to the advancement of aerospace and the exploration of space - areas such as light speed and gravity control. The former was until recently deemed to be unsurpassable, the latter little more than a notional curiosity.

One of the people within the organisation also asking these questions was Marc Millis at NASA Glenn (formerly the NASA Lewis Research Center). Millis and a number of colleagues had been scanning physics literature for years for theoretical ideas; many of them more closely allied to science-fiction than physical reality; that might one day be exploited for deep space travel. The results were posted on a website called 'Warp-Drive: When?'

The website managed to catch the attention of the ASTP team at Marshall, which had staff of its own examining similar kinds of issues. 'Warp-Drive: When?' examines theories ranging from the potential existence of 'worm-holes' - portals connecting distant parts of the universe - and the feasibility of bending space-time, the principle at the heart of the Starship *Enterprise*'s warp-drive system in the 'Star Trek' television series.

BPP became official when NASA Marshall asked Millis to propose a formal programme based on what, until then, had largely been a hobby.

"In the NASA charter we're supposed to sustain pre-eminence in the science and technology of air- and space-flight," Millis says. "To do that we have to look at the edge." Determining the location of the 'edge' was not easy.

Millis illustrates the point with a chart that shows technology on a scale spanning 'routine' - that which is in use today - to 'conjecture' - ideas that are based on "visions without knowledge", the stuff of science-fiction.

"We're also supposed to be prudent investors of the tax-payer's money, which means you have to strike a balance," he says. Consequently, BPP, which is categorised as 'emerging science', is positioned on Millis' chart somewhere between 'science' ("nature understood: when you know what is theoretically possible") and 'speculation' ("knowledgeable conjecture: when you know what you know, and know what you don't"). It is so cutting-edge that much of it is based on science that traditionalists would write off as impossible.

BPP's aim is to deliver progress towards achieving breakthroughs, rather than delivering the breakthroughs themselves. In seeking submissions that would challenge thinking in many areas of accepted physics, it has received 13 proposals that Millis and his panel of expert reviewers considered acceptable. But with its \$450,000 budget, it could only afford to let five contracts. These were awarded between December 1999 and February this year.

The successful bidders placed considerable importance on experimental observation via what Millis describes as "affordable, short-term, relevant research". They must submit their results within the next 12 months to three years.

The first experiment, by Dr John Cramer of Washington University, Seattle, will test a transient inertia effect supposedly observed a decade ago by a scientist called James Woodward in which it was asserted via experiment and theory that a change in inertia is possible with a sudden energy density change.

Manipulate the inertia of an object like a spacecraft and you have the ability to dramatically reduce, if not negate altogether, its need for propellant. But this implies a fundamentally new way of creating motion and a fundamental shift in our understanding of physics. Cramer's experiment is designed to prove one way or another whether Woodward's original experimentation was sound or flawed. The second experiment by Dr [Jordan](#) Maclay and a small industry team from Huntsville, Alabama, will examine the existence of the so-called Zero Point Energy (ZPE) field and whether it exists at the magnitudes that have been postulated. If so, it could be of immense practical value for space travel as well as leading to entirely new sources of non-polluting energy here on earth.

For some years there has been a developing understanding that space is not the empty vacuum of traditional theory, but a seething mass of energy, with particles arising and disappearing and fields continuously fluctuating about their 'zero-point' baselines. These fluctuations of the vacuum remain even at absolute zero or the zero point of temperature - hence the name. If the ZPE field can be 'mined' in space for practical use it would yield an infinite energy source for propulsion.

The third experiment is a test of 'electrodynamic torsion tensor theory'. This will examine the as yet unproven link between electromagnetism and gravity and its impact on space-time. "The immediate utility (for propulsion purposes) is not obvious," Millis says, "but it's like a foot in the door." Perturbations of space-time, however, open up esoteric possibilities such as time travel, as well as the potential for crossing vast oceans of the universe.

The fourth experiment, by Dr Kevin Malloy of the University of New [Mexico](#) in Albuquerque, looks at 'superluminal quantum tunnelling': faster-than-light speed. Recent tests have shown light pulses apparently accelerating beyond the speed of light, thereby shattering Einstein's theory of relativity, which says the light speed barrier cannot be breached.

The BPP experiment aims to prove definitively whether the superluminal effect is indeed real. "It's

really only touching on the make or break issue of whether you can have a faster than light phenomenon," Millis says. "If the answer is 'yes', how that translates into space travel is way down the pike."

The fifth experiment is no less controversial. Conducted under the auspices of the Marshall Space Flight Center itself, it will set out to prove or disprove a highly contentious claim by Russian scientist Evgeny Podkletnov, that he has negated the forces of gravity using super-conducting ceramic discs rotated above powerful electromagnets. The claims were first made public in 1996 when Podkletnov, a materials scientist, was working at the University of Tampere in [Finland](#).

Since then, NASA Marshall and the University of Alabama at Huntsville (UAH) have been putting his claim to the test. They have found cause to believe that something unusual is occurring, although the precise nature of the effect is a matter of conjecture. In his initial tests, Podkletnov claimed that objects he placed above superconductors spinning at 5,000rpm were losing 2% of their weight and, crucially, that the tests were repeatable.

More recently, at a lecture organised by [BAE Systems](#) at the University of Sheffield in the UK in February this year, Podkletnov openly claimed that for a short period of time objects placed in the area of influence above the disc were losing up to 5% of their weight. Higher rotation speeds, Podkletnov told *Jane's Defence Weekly*, had shown that even more dramatic reductions were possible.

Podkletnov, who is now working with the Japanese industry to fine-tune his findings and produce hardware, maintains that the key to success lies in the way in which the superconducting discs are fabricated - a secret he is keeping close to his chest. Part of the problem faced by the NASA/UAH team has been its inability to reproduce superconductors of the size and fidelity Podkletnov says are needed to achieve the desired results.

Under a small business industry research grant outside the BPP programme but running in parallel, a company in Columbus, Ohio, will seek to demonstrate the feasibility of building discs suitable for the BPP gravity shielding experiments - a term with which even Millis is uncomfortable. "It's more accurate to call it an anomalous gravitational effect, but even whether or not it's a gravitational effect is jumping the gun," he says. It could, he adds, simply be an experimental error.


If Podkletnov has discovered a genuine effect, however, its impact on aerospace could be enormous, which is why NASA feels it cannot afford to ignore his work. "If using controllable hardware, you can be creating forces at a distance or manipulating gravity and inertia, it would be a breakthrough and have substantial utility to a NASA mission," Millis says.

Although physicists do not like to use the term '**anti-gravity**', because of its pseudo-scientific connotations, such hardware would usher in an entire new era of propulsion. Podkletnov himself is convinced that his work will be applied to endo-atmospheric aerospace vehicles as well as spacecraft - and sooner, perhaps, than the world of conventional aerospace engineering would like to think.

In experiments using rotation speeds far in excess of those used in his early work, Podkletnov said that his superconducting discs had actually taken off.

The US Air Force maintains close links with BPP and is updated at significant milestones. It conducted investigations of its own into breakthrough propulsion physics - one such was the 1990 Electric Propulsion Study commissioned by the Space Systems Division of Air Force Systems Command at Edwards Air Force Base, California. "If the US Air Force or Navy are doing restricted work in this area, it's not a part of this programme," Millis says, as the BPP programme aims to keep work in the public domain and unrestricted.


And if NASA does hit upon a real breakthrough? "We have to keep an eye out for how things are advancing," Millis says. "For example, if we were to have a breakthrough where you could immediately engineer a device, then we have to reassess what is the impact of this. By having the air force and other military people as part of the network means we're just poised to do that should it happen."




NASA's Breakthrough Propulsion Programme want to determine if a space flight system like the starship Enterprise's warp drive system could be a reality
(Source: Paramount Pictures)



Hypothetical spacecraft with a "negative energy" induction ring, inspired by recent theories describing how space could be warped with negative energy to produce hyperfast transport
(Source: NASA/Les Bossinas)



Recognising a pattern from historical breakthroughs
(Source: Jane's)



Status of potential breakthrough propulsion
(Source: Jane's)



<http://www.grc.nasa.gov/WWW/PAO/warp.htm>
(Source: Jane's)


© 2000 Jane's Information Group

© Jane's Information Group 2002

[Terms of Use](#)

[Powered by Verity](#)






Recognising a pattern from historical breakthroughs

(adapted from Posner, 1986)


To exceed the limits of existing methods, seek out entirely different methods



When is it time to seek the next breakthrough?


When...


- (1) existing methods are reaching their theoretical limits
and
(2) clues are emerging for new methods



Status of potential breakthrough propulsion

(compared to a scale of technological advancement)





[Click for more.](#)

[Log In](#)
[Log Out](#)
[Help](#)
[Feedback](#)

[My Account](#)[Jane's Services](#)[Online Research](#)[Online Channels](#)[Data Search](#) | [Data Browse](#) | [Image Search](#) | [Web Search](#) | [News/Analysis](#)

 Print friendly page

6 Images

VISION 2020, TURNING SCIENCE FICTION INTO FACT

p 30

JANE'S DEFENCE WEEKLY

DATE: JUNE 10, 1995

EDITION: 1995

VOLUME/ISSUE: 023/023

BY LINE:

Nick Cook

INTRODUCTION:

Not only the US Air Force is investing time and money in future air vehicle technology, as Nick Cook reports

TEXT:

Technologists who dabble in the interpretation of visions do not always get it right. Take this example from a specialist US aviation magazine in 1956: "We're already working with equipment to cancel out gravity," Lawrence D Bell, founder of the company that bears his name, was quoted as saying.

Bell, apparently, was not the only one working in this field. Others said to be seeking to master this arcane 'science' included the Glenn L Martin Company, Convair, Lear and Sperry Gyroscope. Within a few years, we were assured, aircraft, cars, submarines and power stations would all be driven by this radical new propulsion technology. Sadly, it was not to be.

However, despite some notable setbacks, the US Air Force has never given up on its 'vision'. The word recurs again and again in service literature and is always on the lips of those who run its science and technology (S&T) programmes. In no other air force - in no other nation - is such prominence given to the techno-visionary.

USAF is constantly looking 25 to 30 years ahead. It encourages those

who work within the service - and the companies that build systems for it - to gaze into the future, to set ambitious goals and to strive to realize them. It goes so far as to map out this vision in unclassified literature - the periodically released 'Air Force S&T Program' - and has a formal mechanism for bringing revolutionary technologies to the boil. The latest example of this, Project Forecast II - the third in a series of studies going back to 1944 - was published in 1985.

Between the S&T Program and Forecast II, we are afforded a glimpse of what might be in 2020. By comparing these documents with the Pentagon's last Defense Technology Plan, published last September, which gazes a maximum of 10 years ahead, it is also possible to zero in on what revolutionary advances USAF may be looking for. The picture is only partly representative because of the enormous resources allocated to 'black' projects, programmes so sensitive that, technically, they do not exist. By their very nature, many future projects fall into this category. The picture is also blurred by USAF's failure to publish an 'S&T Program' since 1990 and the enormous political and military upheavals that have taken place in the interim.

Yet, revolutionary technology, according to USAF, is only partly dependent on these external forces. As well as what USAF calls 'capability needs' - requirements specified by its user commands - the other big driver is 'technology innovation': "the process," says the S&T Program, "whereby an air force laboratory or industry team conceives of a new technology and interests users in exploiting its capability." It goes on: "Most of the really significant improvements in military technology - radar, jet engines and the atomic bomb - occur this way." Even so, quantum leaps require a great deal of funding, and the USAF's S&T community, like the rest of the US military, finds itself with a static budget and short on personnel. USAF R&D and S&T funding for FY95, however, still hovers around the \$11.5 billion mark - a considerable amount of money. Enough, certainly, to transition some key technologies to the user commands and so maintain technological superiority, considered by USAF to be the 'backbone of deterrence', well into the 21st century.

HYPERSONICS

Despite a rash of observer 'sightings' of unidentified high speed, high flying air vehicles in the late 1980s and early 1990s, the US Government has consistently denied it has developed or is developing an aircraft to replace the Mach 3-plus Lockheed SR-71 strategic reconnaissance platform. Its blanket denials of an 'Aurora-type' capability included hypersonic aircraft for missions other than reconnaissance.

That USAF entertained hypersonic ambitions for a range of roles in the mid-to-late 1980s, however, is not in dispute. Four out of eight 'mission area summaries' for air vehicles in the '1990 S&T Plan' - strategic offence, strategic defence, space and recce/intel - called for a NASP-derived vehicle to replace current bombers, orbital launch vehicles and reconnaissance aircraft.

NASP was the X-30

National Aero-Space Plane, a NASA-led effort, since cancelled on funding and technical grounds.

Forecast II said: "The air force will be able to go into space and de-orbit on demand as a result of the improved understanding of

hypersonic aerodynamics, particularly in the Mach 8 and Mach 25 regime." With [NASP](#) cancelled and, apparently, no Aurora capability in the offing either, does this mean USAF has abandoned all hope of fielding a hypersonic platform in the future? Apparently not.

In last year's Defense Technology Plan, a 'roadmap of technology objectives' lists a "flight demonstration of (a) Mach 8 hydrocarbon-fuelled scramjet (supersonic combustion ramjet)" as one of its key goals by 2005 in the aerospace propulsion field. Elsewhere, the document says, "the demonstration of a scramjet to M>12 may lay the foundation for single-stage to orbit space launch." That such a capability is not undergoing significant development in the classified R&D arena would, many qualified observers believe, be highly unusual. Mastering hypersonics is certainly consistent with one of two current overriding drives within USAF labs - operating at ever increasing range from the Continental USA (the other is making current technologies more affordable) as US overseas forces contract.

GAS TURBINE INCREMENTS

Research into more conventional aero-engine propulsion continues via the Integrated High-Performance Turbine Engine Technology (IHPTET) programme and other comparable efforts outside the USA. By 2005, the IHPTET will be delivering 100 per cent thrust-to-weight improvements over current fighter-type jet engines.

In the UK, Rolls-Royce is locked into a technology demonstration schedule that, all things being equal, will see it running a combat engine in the same time period with a 60 per cent improvement in thrust-to-weight ratio over the [Eurofighter](#) 2000's EJ 200 - an airframe-powerplant combination that, as Jane's Defence Weekly went to press, is yet to fly. Though less dramatic improvements will always be sought in other aspects of engine development - fuel burn, life-cycle and acquisition cost, for example - radical enhancements of the jet engine itself are finite. Doubling the power-to-weight ratio again by 2020 may simply not be worth the effort or cost. Alternatively, pushing for all-out power may have an adverse effect in other areas, such as 'signature management', ensuring engines remain as stealthy as their airframes.

THE SUPER COCKPIT

"Machines," according to Forecast II, "must respond reliably to voice commands and eye motion signals, and man and machine must interact to share the sense of touch." An area of research that will be critical in achieving this is virtual reality. It is predicted that the virtual cockpit - the pilot deriving his or her entire view of the outside world via cockpit and helmet displays - will begin equipping operational aircraft in 2020. Research at Aeronautical Systems Center laboratories at Wright-Patterson Air Force Base, Ohio, embodies the essence of what USAF calls the Super Cockpit. According to Forecast II, a two-way flow of visual and aural information between pilot and aircraft will enable him or her to aim and fire weapons or to activate other cockpit functions simply by looking or talking. Add 'thinking' to that list and the picture is even more complete.

Revolutionary advances in the cockpit will be helped by a shift towards commercial off-the-shelf avionics, especially in the computer processor arena.

Modular avionics, based on slide-in, slide-out processor cards, will allow users to improve sensors and cockpit systems without implementing costly mid-life upgrades. Situational awareness will also be vastly improved thanks to real-time access to satellite communications, imagery and other intelligence information distributed between every echelon of the command and fighting forces. Also growing in importance are mission planning and rehearsal tools - most of them PC based - that allow pilots to fly sorties 'virtually' before they have even taken off to strike the target.

WEAPONS

In the 'conventional armament' field, USAF's goal is to develop 'affordable, autonomous, all-weather, day and night stand-off weapon technologies that provide near-perfect accuracy'. By 2020, it will probably be dependent on an advanced ramjet-powered version of, or successor to, the AMRAAM medium-range air-to-air missile and the short-range [AIM-9X](#) dogfight missile or its pre-planned product improvement. Available P3I options are likely to include multi-spectral, stealth-defeating seekers. Beyond improvements to the next-generation air-to-surface weaponry typified by the Joint Direct-Attack Munition and the Joint StandOff Weapon, other more exotic technologies are being sought for soft and hard kill options. "We will transition high and medium power microwave technologies to developers for use as weapons, broadband countermeasure devices, radars and enhanced jammers," the S&T Program reports. "We anticipate breakthroughs in long-range, high-altitude, very high-velocity impact weaponry for use against a variety of hardened (including very deep) targets," Forecast II adds. This has been given added impetus by the development of Third World weapons of mass destruction, which might typically be stored in such places. The current development of an airborne laser for boost phase interception of theatre ballistic missiles is one of those rare leaps in capability that will change the way wars are fought. By 2020, miniaturization techniques may result in deploying lasers in smaller, tactical aircraft. Increased application of photonic as opposed to electronic-based equipment will make user aircraft less susceptible to damage from the EW threat. 'Smart skins' - outer panels containing embedded sensors - will allow aircraft to sense and communicate across multiple frequency bands in any direction from any attitude.

STEALTH

In 1985 and 1990, Stealth was so sensitive that Forecast II and the S&T Program barely gave it a mention. Stealth has changed irrevocably the way in which war in the air is being and will be fought. The USA - and other Western technical powers like France, the UK and Germany - are pouring considerable funds into the mastery of both Stealth and Counter-Stealth technologies.

Take the UK, for example. Since the mid-late 1980s, [British Aerospace](#) Warton has pumped over Pds100 million (\$160 million) - about half of it coming from government sources - into infrastructure improvements geared towards Stealth. Much of this activity is aimed at meeting the RAF's need for a Future Offensive Aircraft (FOA) to replace the [Tornado IDS](#). Traditionally, US firms such as Lockheed and Northrop have enjoyed a US-only monopoly in black programming - the development of aircraft that are invisible to public scrutiny.

Recent revelations that BAe is establishing a 'Skunk Works' on its own remote site at Warton and that the Germans tested manned wind-tunnel models of their Stealth fighter, the Lampyridae, in the mid-1980s - when the first-generation Lockheed [F-117](#) was still so classified its existence was officially denied - show an emerging trend towards 'black' programme secrecy in Europe, mirroring that of the USA.

Together with advanced, lean manufacturing techniques that will allow companies to rapid-prototype and build small batches of aircraft at low cost, expect more from these countries in the way of classified technology demonstration and even low-volume production programmes, including full-sized air vehicles. 'More', because according to numerous sources, such activity is well advanced already. In the USA, it is widely accepted that a third generation of true Stealth aircraft (beyond the first generation [F-117](#) and the second generation [B-2](#)) is already under test and perhaps even in service for missions spanning strike and defence suppression as well as tactical and strategic reconnaissance.

BEYOND 2001

Groom Lake, Nevada, is the epicentre of classified USAF research into Stealth and other exotic aerospace technologies. Several years after the collapse of the Soviet threat, activity and investment at this remote, highly secret air base (so secret its presence is, as yet, unacknowledged by the US Government) is still on the increase. While research into less sensitive technologies such as two-dimensional thrust-vectoring and advanced short take-off and vertical landing (ASTOVL) are pursued in the open at nearby Edwards AFB in California, Groom Lake is set to hang onto its secrets. The USAF's recent confiscation of 1600 ha of public land bordering the facility is consistent with the Pentagon's desire to maintain its lead in quantum leap technologies - some of which, according to well qualified observers in and around the Nevada area, defy current thinking into the predicted direction of aerospace engineering.

That aerospace companies continue to look at highly radical alternative air vehicle concepts is evidence of the ongoing quest for breakthrough designs. Glimpses into this world are rare, but provide some insight into likely 21st century research activity. The 1990 unclassified 'Electric Propulsion Study' (a quest for an antigravity propulsion system by another name) conducted by the USA's Science Applications International Corp on behalf of USAF's then-Astronautics Laboratory at Edwards AFB, shows that USAF's visionaries are still being given free rein. Until recently, BAe also provided internal resources for its own **anti-gravity** studies and even went so far as to outline this thinking with artists' concepts - a case of Lawrence Bell's vision perhaps being not so wide of the mark after all.

Before he died, Ben Rich, who headed Lockheed's Skunk Works from 1975-1991, was quoted as saying: "We have some new things. We are not stagnating. What we are doing is updating ourselves, without advertising. There are some new programmes, and there are certain things - some of them 20 or 30 years old - that are still breakthroughs and appropriate to keep quiet about. Other people don't have them yet."

Thirty years from now, we may still not know the half of what is currently being tested in and around Groom Lake.



Lampyridae missile fighter

A manned wind-tunnel model of Germany's medium-range missile fighter, *Lampyridae*

Lampyridae missile fighter

CAPTION:

A manned wind-tunnel model of Germany's medium-range missile fighter, *Lampyridae*



Future Offensive Aircraft

CAPTION:

Artist's concept of British Aerospace's Future Offensive Aircraft for the UK



*Air vehicle using **anti-gravity***

CAPTION:

BAe Military Aircraft Division's concept for an air vehicle employing **anti-gravity** propulsion



*Air vehicle using **anti-gravity***

CAPTION:

Another BAe concept showing an **anti-gravity** propulsion system in action



Heavy lift vehicle concept

CAPTION:


BAe concept for a heavy lift vehicle, also using an **anti-gravity** propulsion system
















THE JANE'S INTERVIEW SCIENCE FICTION AUTHOR ARTHUR C [CLARKE](#)

p 32

JANE'S DEFENCE WEEKLY

DATE: JANUARY 08, 1994

EDITION: 1994

VOLUME/ISSUE: 021/001

BY LINE:

Nick Cook
[Sri Lanka](#)

TEXT:

World famous author Arthur C [Clarke](#) has recently found himself in demand by the military to talk on how science fiction is becoming science fact.

These days, Arthur C [Clarke](#), the author of 2001: A Space Odyssey, rarely ventures beyond his home in Colombo, [Sri Lanka](#). He admits he is ``fed up'' with talking to the media. ``However,'' he adds, ``I'm always ready to comment on any major development in my areas of interest - for example, an ET landing on the White House lawn, or the first genuine message from space.''

One recent exception to this self-imposed rule was a talk that he gave to the Pacific Area Senior Officer Logistics Seminar (PASOLS) in Colombo, a gathering of high ranking military officers from countries across the Pacific Rim, including the USA.

Diplomatic observers on the ground were surprised by [Clarke's](#)

appearance at PASOLS. At the time, it seemed to gel neither with his own leanings - he confesses to being something of a pacifist - nor with the overall theme, a forum for improving the provision of logistics among allies in the region.

The substance of [Clarke's](#) address was a short dissertation on the process known as 'cold fusion'. Almost five years after two US scientists, Pons and Fleischmann, claimed they had hit upon the reaction that fuels the sun under laboratory, or room temperature conditions - hence the misleading description, 'cold fusion' - it is a subject that consumes him greatly.

It is also a highly controversial one. Soon after the two reported their discovery, that more energy was coming out of their experiments than was being put in - the all-important breakthrough - many other laboratories tried to repeat their tests and failed. Pons and Fleischmann were ``laughed out of court'' in [Clarke's](#) words, for a number of years. Lately, however, there has been a resurgence of interest in their work, ``an underground movement of scientists,'' according to [Clarke](#), ``who believed there might be something in all this business.'' Fresh experiments, many of them backed by the Japanese, others by the US military, confirmed positive results.

``It is now beyond serious dispute that anomalous amounts of energy are being produced from hydrogen by some unknown reaction,'' [Clarke](#) says. However, the theoretical basis of cold fusion is still a mystery, he adds. Though it could be a laboratory curiosity, ``frankly, I doubt this. Anything so novel indicates a breakthrough of some kind.''

For a man who told JDW that he ``won't get contaminated by security'', adding, ``I've never been involved in anything I'm not supposed to talk about,'' [Clarke](#) has been doing a fair amount of talking to the military lately. He recently addressed the US Air Force's Air War College at Maxwell Air Force Base, Alabama, via video-phone link on the future of technology in general.

The USAF, unlike most other air forces, is not afraid to solicit the views of science-fiction writers once in a while in an effort to predict the future direction of air warfare. A few years ago, the USAF instituted Project Forecast II, an initiative aimed to produce a handful of 'quantum leap' technologies for the 21st century. In 1990, the air force's then Astronautics Laboratory (now Phillips Propulsion Labs) published a report under the Electric Propulsion Study that examined theories and experimental approaches for an '**anti-gravity**' propulsion system. The chances are, therefore, that when he talks to the US military, [Clarke's](#) views, including those on cold fusion, have not fallen on deaf ears.

In 2010: *Odyssey II*, published in 1982, [Clarke](#) pretended that the Russians had invented a spaceship engine based on cold fusion principles worked on by Andrei Sakharov during his exile in Gorky. ``He didn't, of course, so that's a piece of fictitious history. However, three Russian scientists who have indeed been working on nuclear propulsion for rockets have now got into the cold fusion act, and they have just published some startling results.

``They are obtaining about five times their energy input in gas mixtures, not solids, and at temperatures of up to 1800 deg C. Now this is not exactly 'cold' fusion - but it's certainly ice-cold compared with the tens of millions of degrees the hot fusioneers are talking about.'' If a plasma fusion rocket could be developed, it would open up the solar system, ``just as the aeroplane opened up this planet,'' he says.

Back on earth, cold fusion would essentially mean the end of the Fossil Fuel Age, [Clarke](#) says; an era of cheap, clean power. ``The environmental benefits would be overwhelming.'' Eschewing the ways of the military as he does, perhaps this is the real message Arthur C [Clarke](#) wants the military men to take away with them.

CAPTION:

Photograph:

© 1994 Jane's Information Group

[© Jane's Information Group 2002](#)

[Terms of Use](#)

[Powered by Verity](#)



COLLABORATE OR PROTECT?

p 56

JANE'S DEFENCE WEEKLY

DATE: JUNE 12, 1993

EDITION: 1993

VOLUME/ISSUE: 019/024

TEXT:

In an atmosphere of increasing tension over its trade links with Europe and

Japan, the USA is having to define how much of this impressive science and technology base it protects (see box 2) and how much it shares with its traditional allies in future joint collaborative efforts.

Since the evaporation of the Soviet threat, defence R&D between the USA and its NATO allies has drifted through lack of focus and funding. The Pentagon has promised Congress renewed emphasis on the programme, including more speedy negotiations and implementation of bilateral agreements, but its commitment to many of these initiatives is unclear.

Twenty-six areas of potential co-operative research between USAF's Electronic Systems Center (ESC) and Japan were identified by some 70 programme managers and engineers in an ESC/Japan initiative conference in February. The Japan initiative was designed to target Japan's defence industry and technology market for possible international co-operative R&D efforts with ESC.

Potential areas of co-operation include: the miniaturization of radar

technology, fibre optics, tactical communications, sensors, lasers, smart skin radars, computer language translation, robotics, advanced wafer technology, theatre missile defence, improved over-the-horizon radar, modelling and simulation, use of sonar and acoustics technology to test materials and Japanese AWACS technology. Various USAF offices will now pursue co-operative efforts in these areas.

Within Europe, the UK is the chief beneficiary of shared S&T research with the United States. Through the Master Information Exchange Agreement (MIEA) -- the technology transfer provision of the US/UK 'special relationship' -- the UK's government research establishments enjoy a two-way exchange of technical data that is the envy of other US allies.

The depth of the relationship is illustrated by the fact that, according to reliable UK sources, the UK's Royal Aeronautical Establishment (now the Defence Research Agency) was passed technical data on the Lockheed [F-117](#) Stealth Fighter in the mid-1980s, long before it ceased to be a heavily compartmentalized programme in the USA.

According to UK officials, the USA only conducts extensive (and often highly sensitive) co-operative research with an ally when that nation comes along with a technological advantage in a given area, or areas. It is significant, for example, that USAF's ESC initiative with [Japan](#) focuses on many of the five technologies (mentioned at the beginning of this article) in which [Japan](#) was deemed to be "significantly ahead" of the USA by USAF.

Of course, when the USA, or any other country, feels it has an overwhelming advantage in a particular field of military science, it classifies it.

The idea of 'facetting' an aircraft to make it stealthy (the approach adopted for the [F-117](#)), only transitioned from the white, to the black world -- in the form of the DARPA Have Blue technology demonstrator programme -- when DoD officials believed what US scientists had been saying openly for some years: that near absolute radar invisibility was achievable.

In the same way, if any Project Forecast II-type technologies were to result in breakthrough -- a quantum leap such as the key to the secret of gravity, say -- the USAF would undoubtedly move to bury it beneath so many layers of classification that it would not see the light of day for years.

As a post-script, anyone who believes the 'science' of **anti-gravity** to be too arcane, too esoteric, even for the US Air Force should

consider the Electric Propulsion Study undertaken by SAIC for the Astronautics Laboratory, now part of Phillips Lab, in the late 1980s.

The study's primary objective was to "outline physical methods to test theories of inductive coupling between electromagnetic and gravitational forces to determine the feasibility of such methods as they apply to space propulsion" -- in simplified terms, an **anti-gravity** propulsion system.

It remains, for some, the ultimate quantum leap.

CAPTION:

Drawing: AFMC: guardian of USAF's wealth in science and technology

CAPTION:

Photograph: The X-30 -- outgrowing of a secret USAF reconnaissance aircraft or pioneering hypervelocity demonstrator?

CAPTION:

Drawing: Myth or reality? Aurora has come to embody secret 'black' aircraft projects (Julian Cook)

CAPTION:

Drawing: The 'More Electric' Aircraft Programme will minimize on-board hydraulics

CAPTION:

Photograph: Sandia National Laboratories, Kirtland AFB: a key component of the New Mexico 'superlab'

CAPTION:

Photograph: Cuts in US nuclear weapons spending is threatening the competitiveness of laboratories like Los Alamos

CAPTION:

Drawing: SAIC proposal for a rail gun satellite launch system reflects US lead in hypervelocity weapons

CAPTION:

Drawing: Los Alamos free electron laser, packaged in a 747, would knock out ballistic missiles soon after launch (Julian Cook)

© 1993 Jane's Information Group

© Jane's Information Group 2002

[Terms of Use](#)

[Powered by Verity](#)

[1 Image](#)

FEATURES

Date Posted: July 24, 2000

JANE'S DEFENCE WEEKLY - JULY 26, 2000

Electrogravitics: the positive approach

One area of research not included in the NASA Breakthrough Propulsion Physics study, but which is touted by some as having the potential to leap the current state of the art in aerospace technology, is '**electrogravitics**'.

This is a postulated form of lift based on the controversial assertion that a capacitor charged positively on one side and negatively on the other will experience weight loss, exhibiting thrust in the direction of its positive pole.

In 1952, the man generally described as the 'father of **electrogravitics**', US inventor Thomas Townsend Brown, submitted a proposal for a 'joint services research and development contract' outlining an **electrogravitics** aerospace programme called 'Project Winterhaven'. This set out how the US military could go about constructing a Mach 3 manned interceptor utilising **electrogravitics** technology. Brown based his data on a series of demonstrations he gave to the US Navy immediately after the Second World War.


In these and subsequent tests, 2ft (61cm) diameter disc-shaped capacitors charged with 50 kilovolts and a continuous energy input of 50 watts are said to have achieved a speed of 17ft/s in a circular air course 20ft in diameter. A second series of tests, in which slightly larger discs were supposedly charged at 150 kilovolts, were rumoured in the mid-1950s to have exhibited results "so impressive as to be highly classified". This has led to claims by proponents of the technology that the US government has conducted a clandestine **electrogravitics** effort ever since under the auspices of a 'deep black' test programme.

Lending possible credence to that assertion is this photograph, published here for the first time. It is purportedly of a sub-scale **electrogravitics** experimental air vehicle mounted on a wind-tunnel test-stand somewhere in the USA. Although *Jane's Defence Weekly* cannot verify its origins, it appears to have been taken in the late 1950s or early 1960s. Intriguingly, Brown recommended the construction of a 10ft diameter disc charged at 500 kilovolts as part of 'Project Winterhaven'.

Officially, the US Air Force and Navy lost interest in **electrogravitics** when their analysis showed that the movement of Brown's capacitors was caused by 'ionic wind' - the displacement of ions around the surface of the charged plates. Brown later claimed to refute this by conducting the tests in a vacuum.

With one or two exceptions, aerospace companies have studiously ignored breakthrough propulsion concepts such as those proposed by Brown and Podkletnov. [BAE Systems](#) describes its 'Project Greenglow' initiative as a "speculative research programme" into esoteric propulsion concepts (such as anti-gravity) and the Boeing [Phantom](#) Works recently conducted tests on a 'reactionless drive' device that apparently contravenes Newton's Third Law of action-reaction. Boeing concluded that the device, which uses two spinning arms transferred through three-dimensions, did work, though not yet at the required impulses for endo-atmospheric propulsion or practical space flight.

Echoing Frank Whittle's early pioneering work on turbojets in the 1920s and 1930s, however, there are those who say it might just be the start of something with far-reaching consequences for the entire future of transportation.



*This late 1950s or early 1960s picture is said to be a sub-scale **electrogravitics** experimental air vehicle mounted on a wind-tunnel test stand
(Source: Jane's)*

© 2000 Jane's Information Group

© Jane's Information Group 2002

[Terms of Use](#)

[Powered by Verity](#)

Technology

SEARCH:

STAY UP-TO-DATE WITH ALL OF THE LATEST ADVANCES

EXPERIMENTS

Weighty Implications: NASA Funds Controversial Gravity Shield

By Jack Lucentini
Special to SPACE.com
posted: 11:00 am ET
28 September 2000

Brushing aside controversy and a few glitches, NASA officials are forging ahead with plans to build a device that they say could work as an antigravity machine.

Most scientists say the idea of such a gadget is ludicrous. But given the stakes, NASA officials say, it's worth a try.



French astronaut Philippe Perrin floating in France's "Vomit Comet" -- the Airbus 300. These training planes are still the only way to change what you weigh -- short of a trip to space.

A machine that even slightly reduces gravity at spacecraft launch sites, agency officials believe, could save significant amounts of money.

"The fact that it had appeared in a credible scientific journal is what really caught our eye."

The opportunity to try out such a machine is expected to come this May, when an Ohio company is scheduled to finish a prototype of the device for NASA.

Not that the space agency's officials themselves have high hopes.

"To say this is highly speculative is probably putting it mildly," acknowledged Ron Koczor, assistant director for science and technology at the Space Science Laboratory in NASA's [Marshall Space Flight Center](#), Huntsville, Alabama.

Nonetheless, NASA awarded a \$600,000 contract last year to Superconductive Components Inc. in Columbus, Ohio to build the device.

Critics say the notion of a "gravity shield" violates Einstein's

The Effects of Gravity

Videos of how gravity governs the universe:

- [Fate of The Milky Way and Andromeda Galaxies](#)
- [Mars' Surface Gravity](#)
- [Gravitational Microlensing: How massive stars can distort light](#)

More Stories

- [Book Review: Is 'The Accelerating Universe' Beautiful?](#)
- [Low Gravity Risky for Ships As Well As Crews, Says Report](#)
- [Physicists' Work Forces Earth to Drop Weight](#)
- [Einstein Experiment Faces Crucial Test This Summer](#)

Multimedia

▼ advertisement

CLASSIC
NASA
VIDEOSOWN ONE OR
COLLECT THEM
ALL TODAY!

SpaceStore

SPACE
COM

BUY NOW!

fundamental [laws of physics](#).

"The theory of gravity is fairly well established, and I don't see it reversing itself," said Francis Slakey, a professor of physics at Georgetown University. The NASA project is "wasted money that could have been used to do legitimate space science," he added.

Koczor portrayed that view as closed-minded.

Scientists such as Slakey "don't seem to be amenable to observing that maybe the laws [of physics] are incomplete," Koczor said.

Throughout history, new discoveries have rocked old assumptions, he pointed out. "People used to talk about laws of conservation of mass, conservation of energy. Then all of a sudden, Einstein comes along and says those are really parts of the same thing."

Einstein wrote that gravity can be considered a bending of space-time that inevitably occurs around massive objects such as planets and stars. That, the conventional view holds, means no mere machine or invention can make it go away; it is not a "force" that can be counteracted.

The conventional scientists aren't the only critics of the NASA project. The agency is also drawing fire from some of its former collaborators in the effort. To see why, it helps to start from the beginning.

In 1992, a Finnish scientist, Eugene Podkletnov, claimed to have built a device that produced a gravity-shielding effect.

It consisted of a hot, fast-spinning, 12-inch (30-centimeter) disk of a superconducting ceramic, levitating within a magnetic field. Objects above the disk, Podkletnov reported, showed a loss of weight of between about 0.5 percent and 2 percent.

In 1996, researchers at Marshall Space Flight Center decided to investigate the claims. "The fact that it had appeared in a credible scientific journal is what really caught our eye," Koczor said.

Actually, Podkletnov had withdrawn his most recent article from publication under unclear circumstances. But he and others had published research on antigravity phenomena in several peer-reviewed journals.

Koczor assembled a team that worked together with scientists at the nearby University of Alabama at Huntsville, to build a device partially simulating the one Podkletnov had used. But the researchers were unable to replicate Podkletnov's results, and the partnership fell apart last year with bad blood between the two sides.

The university's Larry Smalley, a physics professor, says NASA simply failed to assemble a competent team of scientists who could give the project a serious chance.

The events "amused me, stunned me and upset me," said Smalley, who said he was involved as an observer of the project at the time. "It made me feel like they wasted time, a lot of money and a really golden opportunity to do something."

Smalley said he remains skeptical that Koczor and NASA have the know-how to do anything meaningful with the project.

The main university professor involved with the project, Ning Li, has since left the school. She said she has founded a company in Huntsville that also will market a gravity-shield device.

Li said she dropped the NASA collaboration and decided to work independently after the agency "wasted" the project's money and resources.

Koczor said the project fell apart not because of incompetence, but because Li was primarily interested in proving her theories of why the "gravity shield" would work. That differed from NASA's goal of simply building a working device, he said.

"She wanted the research to focus on her particular theory. Our intent was simply to show there was a gravity effect, without saying 'theory A is right' or 'theory B is right,'" he explained.

Last year, NASA decided to try again, this time by contracting out the construction of the device. Superconductive Components is in communication with Podkletnov as they attempt to build it, Koczor said.

The project is on or ahead of schedule, said J.R. Gaines, vice president of Superconductive Components.

"The superconductor is built. The rest has been designed and fabrication is proceeding," Gaines said. However, he said, he can't offer an opinion on whether the device will actually work. The company's job is simply to build it to the assigned specifications.

"We don't necessarily have a technical opinion," he said, though "we would certainly love to see this work."

■ [Fate of The Milky Way and Andromeda Galaxies](#)

■ [Mars' Surface Gravity](#)

■ [Gravitational Microlensing: How massive stars can distort light](#)



Dark Energy: Astronomers Still 'Clueless' About Mystery Force Pushing Galaxies Apart



**SCIENCE
TUESDAY**

By [Andrew Chaikin](#)
Editor, Space & Science
posted: 07:00 am ET
15 January 2002

It sounds like something out of a *Star Trek* episode: Dark energy, a mysterious force that no one understands, is causing the universe to fly apart faster and faster. Only a few years ago, if you'd suggested something like that to astronomers, they would have told you to spend less time in front of the TV and more time in the "real" world.

But dark energy is real — or at least, a growing number of astronomers think it is. No one, however, can truly explain it.

"Frankly, we just don't understand it," says Craig Hogan, an astronomer at the University of Washington at Seattle. "We know what its effects are," Hogan says, but as to the details of dark energy, "We're completely clueless about that. And everybody's clueless about it."

Dark energy entered the astronomical scene in 1998, after two groups of astronomers made a survey of exploding stars, or supernovas, in a number of distant galaxies. These researchers found that the supernovas were dimmer than they should have been, and that meant they were farther away than they should have been. The only way for that to happen, the astronomers realized, was if the expansion of the universe had sped up at some time in the past.

Until then, astronomers had generally believed that the cosmic expansion was gradually slowing down, due to the gravitational tugs that individual galaxies exert on one another. But the [supernova results](#) implied that some mysterious force was acting against the pull of gravity, causing galaxies to fly away from each other at ever greater speeds.

It was a stunning realization.

At first, other researchers questioned the result; perhaps the supernovas were dimmer because their light was being blocked by clouds of interstellar dust. Or maybe the supernovas themselves were intrinsically dimmer than scientists thought. But with careful checking, and more data, those explanations have largely been put aside, and the dark energy hypothesis has held up.

In one sense, the idea is not completely new. Einstein had included such an "anti-gravity" effect in his theory of general relativity, in his so-called cosmological constant. But Einstein himself, and later many other astronomers, came to regard this as a kind of mathematical contrivance

▼ advertisement

BUY NOW!

Starry night pro.

**Advanced
Software
for the
Astronomy
Enthusiast**




SCIENCE TUESDAY


Visit SPACE.com to explore a new science feature each Tuesday.

[>>Go to Science Tuesday archive page](#)

Images



This chart shows how much of the universe is made up of dark energy, dark matter, and ordinary matter.



This diagram reveals changes in the rate of expansion since the universe's birth 15 billion years ago. The more shallow the curve, the faster the rate of expansion. The curve changes noticeably about 7.5 billion years ago, when objects in the universe began to move away from us at an increasing rate.

CLICK HERE



Related SPACE.com STORIES

that had little relationship to the real universe. By the 1990s no one expected that the effect would turn out to be real.

Still, anti-gravity isn't the right way to describe dark energy, says Virginia Trimble of the University of Southern California at Irvine.

"It doesn't act opposite to gravity," Trimble says. "It does exactly what general relativity says it should do, if it has negative pressure."

Trimble has a fairly simple way of imagining the phenomenon.

"If you think in terms of the universe as a very large balloon," she says, "when the balloon expands, that makes the local density of the [dark energy] smaller, and so the balloon expands some more because it exerts negative pressure. While it's inside the balloon it's trying to pull the balloon back together again, and the lower the density of it there is, the less it can pull back, and the more it expands. This is what happens in the expanding universe."

The supernova evidence suggests that the acceleration kicked in about 5 billion years ago. At that time, galaxies were far enough apart that their gravity (which weakens with distance) was overwhelmed by the relatively gentle but constant repulsive force of dark energy. Since then, dark energy's continuing push has been causing the cosmic expansion to speed up, and it seems likely now that this expansion will continue indefinitely.

"It means that if you look out at the universe today, and if we wait many billions of years," says Hogan, "everything will be flying away faster and faster, and eventually we'll be left quite alone."

Aside from such grim forecasts, dark energy is causing quite a bit of upset for astronomers who have to adjust to an unexpected and outlandish new view of the universe. Already, they have had to accept the notion of [dark matter](#), which is now thought to far outnumber ordinary matter in the universe, but which has never been detected in any laboratory. Now, the arrival of an unknown force that rules cosmic expansion has added insult to injury.

"I'm as big a fan of dark matter and dark energy as anybody else," says astronomer Richard Ellis of Caltech. But, he adds, "I find it very worrying that you have a universe where there are three constituents, of which only one [i.e., ordinary matter] is really physically understood."

"When you teach undergraduates, and they say, 'Well, what is dark matter?' Well, nobody's really sure. 'What is dark energy?' We're even less sure. So you have to explain to a student, that ... 90 percent of the universe, 95 percent, is in two ingredients that nobody really understands," says Ellis. "This isn't really progress."

No one argues that dark energy is difficult to comprehend. And as Trimble points out, it is hardly the first strange idea scientists have had to accept.

"It took two generations for people to be comfortable with quantum mechanics," she says. "The fact that you do not have good intuition about [dark energy] is true for quantum mechanics, general relativity, and lots of other things, because we can't easily mock them up in the laboratory."

And for cosmologists, dark energy has solved at least one cosmological conundrum raised by studies of the Cosmic Microwave Background, or CMB.

Next page: An infinitely expanding universe?

■ [Astrophysics Challenged By Dark Energy Finding](#)

■ [Farthest Supernova Detected, 'Dark Energy' Suspected](#)

■ ['Groundbreaking' Discovery: First Direct Observation of Dark Matter](#)

■ [Understanding Dark Matter and Light Energy](#)

TODAY'S DISCUSSION

What do you think of this story?

[>>Uplink your views](#)





A02-14241

AIAA-2002-1131

**"Outside the Box" Space and Terrestrial
Transportation and Energy Technologies for the
21st Century**

Theodore C. Loder, III

Institute for the Study of Earth, Oceans and Space

University of New Hampshire

Morse Hall

Durham, NH 03824

40th AIAA Aerospace Sciences Meeting and Exhibit

14-17 January 2002

Reno, Nevada

For permission to copy or to republish, contact the copyright owner named on the first page.

**For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.**

"OUTSIDE THE BOX" SPACE AND TERRESTRIAL TRANSPORTATION AND ENERGY TECHNOLOGIES FOR THE 21ST CENTURY

Theodore C. Loder, III¹

Institute for the Study of Earth, Oceans and Space
University of New Hampshire
Durham, NH 03824

Abstract

This paper reviews the development of antigravity research in the US and notes how research activity seemed to disappear by the mid 1950s. It then addresses recently reported scientific findings and witness testimonies - that show us that this research and technology is alive and well and very advanced. The revelations of findings in this area will alter dramatically our 20th century view of physics and technology and must be considered in planning for both energy and transportation needs in the 21st century.

Historical Background

Townsend Brown's Technology of Electrogravitics¹

In the mid 1920's Townsend Brown² discovered that electric charge and gravitational mass are coupled. He found that when he charged a capacitor to a high voltage, it had a tendency to move toward its positive pole. This became known as the Biefeld-Brown effect. His findings were opposed by conventional minded physicists of his time.

The Pearl Harbor Demonstration. Around 1953, Brown conducted a demonstration for military top brass. He flew a pair of 3-foot diameter discs around a 50-foot course tethered to a central pole. Energized with 150,000 volts and emitting ions from their leading edge, they attained speeds of several hundred miles per hour. The subject was thereafter classified.

Project Winterhaven. Brown submitted a proposal to the Pentagon for the development of a Mach 3 disc shaped electrogravitic fighter craft. Drawings of its basic design are shown in one of his patents. They are essentially large-scale versions of his tethered test discs.

Review of Issues From the 1950s

Once Brown's findings became well known, some scientists began to openly speak about the flying technology of UFOs, which had been observed extensively since the 1940s. None other than Professor

Hermann Oberth, considered by some to be one of the fathers of the space age, who later worked in the US with Wernher von Braun, the Army Ballistic Missile Agency and NASA, stated the following in 1954.³

"It is my thesis that flying saucers are real and that they are space ships from another solar system." Perhaps of more interest to our present discussion on propulsion, he then stated that: "They are flying by the means of artificial fields of gravity... They produce high-tension electric charges in order to push the air out of their paths, so it does not start glowing, and strong magnetic fields to influence the ionized air at higher altitudes. First, this would explain their luminosity... Secondly, it would explain the noiselessness of UFO flight..." We now know that he was fundamentally correct in his assessment.

In 1956, a British research company, Aviation Studies (International) Ltd. published a classified report on Electrogravitics Systems examining various aspects of gravity control. They summarized the pioneering work of Townsend Brown and then described the use of electrogravitic thrust as follows:

"The essence of electrogravitics thrust is the use of a very strong positive charge on one side of the vehicle and a negative on the other. The core of the motor is a condenser and the ability of the condenser to hold its charge (the K-number) is the yardstick of performance. With air as 1, current dielectrical materials can yield 6

¹ Copyright © 2002 by Theodore C. Loder. Published by the American Institute of Aeronautics and Astronautics, Inc. with permission. Presented at the 40th AIAA Aerospace Sciences Meeting and Exhibit, Reno NV. Paper number AIAA-2002-1131.

and use of barium aluminate can raise this considerably, barium titanium oxide (a baked ceramic) can offer 6,000 and there is a promise of 30,000, which would be sufficient for supersonic speed.”⁴

In one of their conclusions, based on Brown’s work, they suggested that: “Electrostatic energy sufficient to produce a Mach 3 fighter is possible with megavolt energies and a k of over 10,000.”⁵

In spite of Brown’s solid research, they later stated that, “One of the difficulties in 1954 and 1955 was to get aviation to take electrogravitics seriously. The name alone was enough to put people off.”⁶ It seems that is as true today as it was in the 1950s.

A report by another British company, Gravity Rand, Ltd. in 1956, agrees with this assessment and states: “To assert electrogravitics is nonsense is as unreal as to say it is practically extant. Management should be careful of men in their employ with a closed mind or even partially closed mind on the subject.”⁷

However, a trade press magazine, The Aviation Report, made a numerous references to antigravity projects and listed many of the companies pursuing research in this area. Quotes from The Aviation Report listed in the Aviation Studies (International) Ltd. report⁸ are suggestive of what was going on behind the scenes.

In 1954 they predicted that: “... progress has been slow. But indications are now that the Pentagon is ready to sponsor a range of devices to help further knowledge.”... “Tentative targets now being set anticipate that the first disk should be complete before 1960 and it would take the whole of the ‘sixties to develop it properly, even though some combat things might be available ten years from now.” (Aviation Report 12 October 1954)⁹

During this time period many of the major defense and technology companies were cited as either having research projects or activities in this new field. For example: “Companies studying the implications of gravitics are said, in a new statement, to include Glenn Martin, Convair, Sperry-Rand, and Sikorsky, Bell, Lear Inc. and Clark Electronics. Other companies who have previously evinced interest include Lockheed, Douglas and Hiller.” (Aviation Report 9 December 1955)¹⁰.

Others of these reports mention: AT&T, General Electric, as well as Curtiss-Wright, Boeing and North American as having groups studying electrogravitics.

During the same time period, the Gravity Rand report notes that: “Already companies are specializing in evolution of particular components of an electogravitics disk.”¹¹

However, in the area of predictions, the Aviation Report stated the following based on an extrapolation of technology development: “Thus this century will be divided into two parts – almost to the day. The first half belonged to the Wright Brothers who foresaw nearly all the basic issues in which gravity was the

bitter foe. In part of the second half, gravity will be the great provider. Electrical energy, rather irrelevant for propulsion in the first half becomes a kind of catalyst to motion in the second half of the century.” (Aviation Report 7 September 1954).¹²

Looking back it is easy to say that they missed the mark. Did they really miss it by a half a century? Reading through these reports it is quite obvious that there was much interest in antigravity among a number of very high profile companies, as well as in the Department of Defense. What happened to this interest and why was it all downplayed during the following four plus decades? After all, T. Brown had shown that there is a demonstrable connection between high voltage fields and gravity. Why has it taken until the 1990s for more than just a few scientists to look at these results and publish on them in the open literature? A review of recent statements by former military personnel and civilians connected to covert projects begins to shed light on research activity in these areas over the last half century. And it appears that there had been significant breakthroughs during this time period, well shielded from both the scientific and public eye.

Recent Scientific Developments

In this section we consider developments in the antigravity field since the late 1980s and why the confluence of scientific findings and the testimony of witnesses associated with the military and covert groups indicates that a gravity solution with technological implications has been found.

Although general relativity has not been able to explain Brown’s electrogravitic observations, or any other antigravity phenomenon, the recent physics methodology of quantum electrodynamics (QED), appears to offer the theoretical framework to explain electrogravitic coupling. Recent papers by members of the Institute for Advanced Study Alpha Foundation are putting a solid theoretical foundation onto the antigravity effects within the theory of electrodynamics and include papers by Evans¹³ and Anastasozki et al.¹⁴

Earlier in a breakthrough paper in 1994, Alcubierre showed that superluminal space travel is, in principle, physically possible and will not violate the tenants of the theory of relativity¹⁵. Puthoff¹⁶ later analyzed these findings in light of the present SETI (Search for Extraterrestrial Intelligence) paradigms that insist that we could not be visited by extraterrestrial civilizations because of the speed-of-light limitations dictated by the general relativity theory. He suggests that superluminal travel is indeed possible. This leads to reduced-time interstellar travel and the possibility of extraterrestrial visitation, which our limited understanding of physics and scientific arrogance has “forbidden” in some sectors for most of the 20th century.

The second aspect of these physics findings deals with the zero point or vacuum state energy shown by the Casimir effect, which predicts that two metal plates close together attract each other due to imbalance in the quantum fluctuations. The implications of this zero point or vacuum state energy are tremendous and are described in several papers by Puthoff¹⁷ starting during the late 1980s. A detailed bibliography on this and similar topics is available on the WWWeb.¹⁸ Bearden¹⁹ and colleagues have also written extensively on the theoretical physics of zero point energy and additionally have described various technological means of extracting this energy (for example see the recent paper by Anastasozki et al.²⁰). A theoretical book on zero point energy by Bearden will soon be available.²¹ There is significant evidence that scientists since Tesla have known about this energy, but that its existence and potential use has been discouraged and indeed suppressed over the past half century or more.²²

The coupling of the electrogravitic phenomena observations and the zero point energy findings are leading to a new understanding of both the nature of matter and of gravity. This is just now being discussed in scientific journals (though some evidence suggests that it has been understood for decades within the black project covert community). The question that is being addressed is: what keeps the universe running? Or more specifically, where do electrons get their energy to keep spinning around atoms? As electrons change state they absorb or release energy, and where does it come from? The simplistic answer is that it is coming from the vacuum state. Puthoff²³ describes the process as follows: "I discovered that you can consider the electron as continually radiating away its energy as predicted by classical theory, but SIMULTANEOUSLY ABSORBING a COMPENSATING AMOUNT of energy from the ever-present sea of zero-point energy in which the atom is immersed. An equilibrium between these two processes leads to the correct values for the parameters that define the lowest energy, or ground-state orbit (see "Why atoms don't collapse," NEW SCIENTIST, July 1987). Thus there is a DYNAMIC EQUILIBRIUM in which the zero-point energy stabilizes the electron in a set ground-state orbit. It seems that the very stability of matter itself appears to depend on an underlying sea of electromagnetic zero-point energy."

Furthermore, it appears that it is the spinning of electrons that provides inertia and mass to atoms. These theories, linking electron spin, zero point energy, mass, and inertia have been presented in a number of recent papers, such as those by Haisch²⁴ and colleagues and provide us with a possible explanation of the Biefeld-Brown effect. It appears that an intense voltage field creates an electromagnetic barrier that blocks the atomic structure of an atom from interacting

with the zero point field. This slows down the electrons, reducing their gyroscopic effect, and thus reducing atomic mass and inertia, making them easier to move around.

Evidence of Extensive Antigravity Technology

The B-2 Advanced Technology Bomber

In 1993, LaViolette wrote a paper²⁵ discussing the B-2 bomber and speculating on its probable antigravity propulsion system, based on a solid understanding of electrogravitics,²⁶ the aircraft's design and the materials used in its manufacture. It appears that the craft is using a sophisticated form of the antigravity principles first described by T. Brown. Support for this thesis came from the Aviation Week and Space Technology (March 9, 1992), which reported that the B-2 bomber electrostatically charges its leading edge and its exhaust stream. Their information had come from a small group of former black project research scientists and engineers suggesting the B-2 utilizes antigravity technology. This information was supported by Bob Oechsler, an ex-NASA mission specialist who had publicly made a similar claim in 1990. These findings support the contention that there have been major developments in the area of antigravity propulsion which are presently being applied in advanced aircraft.

LaViolette later states the obvious that "the commercial airline industry could dramatically benefit with this technology which would not only substantially increase the miles per gallon fuel efficiency of jet airliners, but would also permit high-speed flight that would dramatically cut flight time."²⁷

The Disclosure Project Witnesses

On May 9, 2001 a private organization, "The Disclosure Project"²⁸ held a press conference at the National Press Club in Washington DC. They presented nearly two dozen witnesses including retired Army, Navy and Air Force personnel, a top FAA official, members of various intelligence organizations including the CIA and NRO, and industry personnel, all of whom who had witnessed UFO events or had inside knowledge of government or industrial activities in this area. They also produced a briefing document²⁹ for members of the press and Congress and a book³⁰ which includes the testimony of nearly 70 such witnesses from a pool of hundreds. Although they all spoke of the reality of the UFO phenomena, many also spoke of covert projects dealing with antigravity, zero point energy technologies and development of alien reproduction vehicles (ARVs) by US black project and covert interests. The following excerpted quotes from these witnesses support the above contentions.

Dan Morris³¹ is a retired Air Force career Master Sergeant who was involved in the extraterrestrial

projects for many years. After leaving the Air Force, he was recruited into the super-secret National Reconnaissance Organization (NRO), during which time he worked specifically on extraterrestrial-connected operations. He had a cosmic top-secret clearance (38 levels above top secret) which, he states, no U.S. president, to his knowledge, has ever held.

"UFOs are both extraterrestrial and man made. Well, the guys that were doing the UFOs, they weren't sleeping, and Townsend Brown was one of our guys who was almost up with the Germans. So we had a problem. We had to keep Townsend Brown - what he was doing on anti-gravity electromagnetic propulsion secret." He then describes a type of zero point energy device.

"Well, if you have one of these units that's about sixteen inches long and about eight inches high and about ten inches wide, then you don't need to plug into the local electric company. These devices burn nothing. No pollution. It never wears out, because there are no moving parts. What moves are electrons, in the gravity field, in the electronic field, and they turn in opposite directions, okay?"

Dr. B.³² (name withheld since he still works in this area) is a scientist and engineer who has worked on top-secret projects almost all his life. Over the years he has directly worked on or had involvement with such projects involving anti-gravity, chemical warfare, secure telemetry and communications, extremely high-energy space based laser systems, and electromagnetic pulse technology.

"Anti-gravity. As a matter of fact, I used to go out to the Hughes in Malibu. They had a big think tank up there. Big anti-gravity projects; I used to talk to them out there. I'd give them ideas, because they bought all my equipment. But the American public will never, never hear about that." . . . "This flying disc has a little plutonium reactor in it, which creates electricity, which drives these anti-gravity plates. We also have the next level of propulsion, it is called virtual field, which are called hydrodynamic waves..."

Captain Bill Uhouse³³ served 10 years in the Marine Corps as a fighter pilot, and four years with the Air Force at Wright-Patterson AFB as a civilian doing flight-testing of exotic experimental aircraft. Later, for the next 30 years, he worked for defense contractors as an engineer of antigravity propulsion systems: on flight simulators for exotic aircraft - and on actual flying discs.

"I don't think any flying disc simulators went into operation until the early 1960s - around 1962 or 1963. The reason why I am saying this is because the simulator wasn't actually functional until around 1958. The simulator that they used was for the extraterrestrial craft they had, which is a 30-meter one that crashed in Kingman, Arizona, back in 1953 or 1952."

"We operated it with six large capacitors that were charged with a million volts each, so there were six million volts in those capacitors." . . . "There weren't any windows. The only way we had any visibility at all was done with cameras or video-type devices." . . . "Over the last 40 years or so, not counting the simulators - I'm talking about actual craft - there are probably two or three-dozen, and various sizes that we built."

A.H.³⁴ formally of Boeing Aerospace, is a person who has gained significant information from inside the UFO extraterrestrial groups within our government, military, and civilian companies. He has friends at the NSA, CIA, NASA, JPL, ONI, NRO, Area 51, the Air Force, Northrup, Boeing, and others.

"Most of the craft operate on antigravity and electrogravitic propulsion. We are just about at the conclusion state right now regarding antigravity. I would give it maybe about 15 years and we will have cars that will levitate using this type of technology. We're doing it up at Area 51 right now. That's some of the stuff that my buddy worked on up at Area 51 with Northrup, who lives now in Pahrump, Nevada. We're flying anti-gravity vehicles up there and in Utah right now..."

Lieutenant Colonel Williams³⁵ entered the Air Force in 1964 and became a rescue helicopter pilot in Vietnam. He has an electrical engineering degree and was in charge of all the construction projects for the Military Air Command. During his time in the military he knew that there was a facility inside of Norton Air Force Base in California that no one was to know about.

"There was one facility at Norton Air Force Base that was close hold - not even the wing commander there could know what was going on. During that time period it was always rumored by the pilots that that was a cover for in fact the location of one UFO craft."

Note that all he knew was of the rumor, however, it is confirmed by the next testimony, which also confirms some of the comments made by Captain Uhouse.

Mark McCandlish³⁶ is an accomplished aerospace illustrator and has worked for many of the top aerospace corporations in the United States. A colleague, with whom he studied, has been inside a facility at Norton Air Force Base, where he witnessed alien reproduction vehicles, or ARVs, that were fully operational and hovering. He states that the US not only has operational antigravity propulsion devices, but we have had them for many, many years, and they have been developed through the study, in part, of extraterrestrial vehicles over the past fifty years.

A close friend, Brad Sorensen, told him of visiting The Big Hangar, during an air show at Norton Air Force Base on November 12, 1988 and how he had

seen flying saucers in this hangar. "There were three flying saucers floating off the floor—no cables suspended from the ceiling holding them up, no landing gear underneath—just floating, hovering above the floor. He said that the smallest was somewhat bell-shaped. They were all identical in shape and proportion, except that there were three different sizes. They had little exhibits with a videotape running, showing the smallest of the three vehicles sitting out in the desert, presumably over a dry lakebed, some place like Area 51. It showed this vehicle making three little quick, hopping motions; then [it] accelerated straight up and out of sight, completely disappearing from view in just a couple of seconds—no sound, no sonic boom—nothing."

"Well, this craft was what they called the Alien Reproduction Vehicle; it was also nicknamed the Flux Liner. This antigravity propulsion system—this flying saucer—was one of three that were in this hangar at Norton Air Force Base. [Its] synthetic vision system [used] the same kind of technology as the gun slaving system they have in the Apache helicopter: if [the pilot] wants to look behind him, he can pick a view in that direction, and the cameras slew in pairs. [The pilot] has a little screen in front of his helmet, and it gives him an alternating view. He [also] has a little set of glasses that he wears—in fact, you can actually buy a 3-D viewing system for your video camera now that does this same thing—so when he looks around, he has a perfect 3-D view of the outside, but no windows. So, why do they have no windows? Well, it's probably because the voltages that we're talking about [being] used in this system were probably something between, say, half a million and a million volts of electricity." Brad Sorensen stated that at the ARV display, "a three star general said that these vehicles were capable of doing light speed or better."

All of these witness testimonies do not prove the existence of a successful US covert program in antigravity technologies. Only the demonstration of such craft coupled with the accompanying government and technical specification documents would 'prove' this. However, these testimonies coupled with information from other substantial sources such as Nick Cook's new book mentioned below, strongly supports this contention.

The Hunt for Zero Point³⁷

Although this very recent book (August, 2001) is not yet available in the US, it contains some of the strongest evidence yet for major efforts and success in the field of antigravity technology. The author, Nick Cook, who for the past 15 years has been the Aviation Editor and Aerospace Consultant for Jane's Defense Weekly, spent the last 10 years collecting information for the book. This included archival research on Nazi

Germany's antigravity technology and interviews with top officials at NASA, the Pentagon and secret defense installations. He shows that America has cracked the gravity code and classified the information at the highest security levels. Because antigravity and its allied zero point energy technologies potentially offer the world a future of unlimited, non-polluting energy it has been suppressed because of the "huge economic threat". His findings support those reported by many of the Disclosure Project witnesses cited above.

Antigravity Technology Demonstrations

Although T. Brown reported many of his findings nearly a half century ago, other experimenters have just recently begun to reproduce his work and report it in the open literature and on the WWW. For example, Davenport³⁸ published the results of his work in 1995 supporting the findings of T. Brown. More recently, Transdimensional Technologies³⁹ in the USA and J. Naudin⁴⁰ labs in France have posted on the WWW: diagrams, web videos, and data on their versions of antigravity "Lifters" based on an extension of Brown's work. It is a sad commentary on this whole area of research to see that public science is requiring us to demonstrate principles that were demonstrated nearly fifty years ago.

There have also been a number of other demonstrations of "antigravity" phenomena by researchers throughout the world. This includes the work of Brazilian physics professor, Fran De Aquino, and such devices as: the Searl Electrogravity Disc, the Podkletnov Gravity Shield and Project Greenglow, the Zinsser Kineto-baric Field Propulsion and the Woodward Field Thrust Experiments on Piezoelectrics. All of these are described in more detail by Greer and Loder.⁴¹

Implications of This Research

- Antigravity and zero point energy research and their applications are finally being addressed by some of the open scientific community. This means there will have to be a rewriting of textbooks in this area so our new generation of students can apply this "new knowledge." Its application will lead to major breakthroughs in transportation technologies both earthside and in outer space. The implications are that we have the potential for human exploration of our solar system and beyond, if we have the will, within our lifetimes. It also means that the majority of 20th century space technology will be obsolete and in fact may already be so.

- The zero point or vacuum state energy source is seen as a totally non-polluting energy source, which has the potential to replace all the fossil fuels on this planet. It also will provide the energy needed for long range space flights. This means that fuel cells and solar cells

in common use today for space flight energy applications will only be needed until we transition to these new energy technologies.

- Based on an analysis of trends in antigravity research over the last half-century and the information provided by numerous witnesses, it appears that there is both good and bad news. The good news is that it appears that we (at least covert projects) have already developed the theories of antigravity, and additionally have developed working spacecraft based on these principles. The bad news is that these technologies have been developed for at least several decades, at the public's expense and that human kind has been deprived of these technologies, while continuing to waste energy using less efficient and pollution enhancing technologies.

Supporting this contention is the following quote from Ben Rich, former head of the Lockheed Skunkworks. Just prior to his death, he stated to a small group after a lecture that: "We already have the means to travel among the stars, but these technologies are locked up in black projects and it would take an act of God to ever get them out to benefit humanity..." He further went on to say, 'anything you can imagine we already know how to do.' Strong words from a knowledgeable deep insider and words that support what a number of the witnesses stated as well.

- As the reality of this knowledge begins to be understood, there will be an outcry among space scientists not on the inside for release of these technologies to allow all of us to explore space. There will be major changes in the way that NASA does its business, though predicting these changes is difficult.

- Not only has space exploration in the public sector suffered, but our planet's environment has suffered as well. Thus as this knowledge begins to sink in there will be an outcry among all concerned citizens on this planet for release of these technologies to allow all of us to reduce and ultimately eliminate global warming and environmental pollution that so threatens our way of life. These technologies will not only affect space travel technologies, but will also have a profound effect on transportation and energy production on the earth's surface.

- In conclusion, we might consider the observation made by Halton Arp⁴²:

"We are certainly not at the end of science.

Most probably we are just at the beginning!"

Acknowledgements

I thank the following people for many discussions while preparing and writing this paper including: S. Greer, A. Craddock, T. Bearden, P. LaViolette, M. McCandlish, D. Hamilton, T. Valone, E. Mallove, T. Loder, C. Loder, S. Patel and many of the courageous Disclosure Project witnesses.

References Cited

-
- ¹ Quoted from: LaViolette, P. A. 2000. Moving Beyond the First Law and Advanced Field Propulsion Technologies. In T. Loder (ed.), "Outside-the-Box" Technologies, Their Critical Role Concerning Environmental Trends, and the Unnecessary Energy Crisis. Report prepared for The U.S. Senate Environment and Public Works Comm. www.senate.gov/~epw/loder.htm.
- ² Brown, T. T. 1929. How I control gravity. Science and Information Magazine, Aug. 1929. Reprinted in Psychic Observer 37 (1): 66-67.
- ³ Oberth, Hermann: "Flying Saucers Come from a Distant World," *The American Weekly*, October 24, 1954.
- ⁴ Aviation Studies (International) Ltd. 1956. Electrogravitics Systems: An examination of electrostatic motion, dynamic counterbary and barycentric control. p. 14. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005.
- ⁵ Ibid. p. 27.
- ⁶ Ibid. p. 19.
- ⁷ Gravity Rand Ltd. 1956. The Gravitics Situation. p. 54. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005.
- ⁸ Aviation Studies (International) Ltd. 1956. Electrogravitics Systems: An examination of electrostatic motion, dynamic counterbary and barycentric control. p. 11. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005.
- ⁹ Ibid. p. 34.
- ¹⁰ Ibid p. 41.
- ¹¹ Gravity Rand Ltd. 1956. The Gravitics Situation. p. 47. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005.
- ¹² Aviation Studies (International) Ltd. 1956. Electrogravitics Systems: An examination of electrostatic motion, dynamic counterbary and barycentric control. p. 32. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005.
- ¹³ Evans, M. W. 2002. The link between the Sachs and O(3) theories of electrodynamics. In Evans, M.W. (ed.), Modern Nonlinear Physics, Part 2. 2nd ed., Advances in Chemical Physics 19:469-494.
- ¹⁴ Anastasovski, P.K., T.E. Bearden, C. Ciubotariu, W.T. Coffey, L.B. Crowell, G.J. Evans, M.W. Evans, R. Flower, A. Labounsky, B. Lehnert, M. Mészáros, P.R. Molnár, S. Roy, and J.-P. Vigier. (In Press). Anti

- gravity effects in the Sachs theory of electrodynamics Foundations of Physics Letters.
- ¹⁵ Alcubierre, M. 1994. The Warp Drive: Hyper-fast travel within general relativity. Classical and Quantum Gravity, 11, L73.
- ¹⁶ Puthoff, H. E. 1996. SETI, The Velocity-of-Light Limitation, and the Alcubierre Warp Drive: An Integrating Overview, Physics Essays 9:156.
- ¹⁷ Puthoff, H. 1989. Gravity as a Zero-Point Fluctuation Force." Phys. Rev A., 39(5):2333-2342.
- Puthoff, H. 1989. Source of Electromagnetic Zero-Point Energy." Phys. Rev A, 40(9):4597-4862.
- ¹⁸ www.motionsciences.com
- ¹⁹ See the Tom Bearden web site for an extensive listing and copies of his papers at: www.cheniere.org.
- ²⁰ Anastasovski, P.K., T.E. Bearden, C. Ciubotariu, W.T. Coffey, L.B. Crowell, G.J. Evans, M.W. Evans, R. Flower, A. Labounsky, B. Lehnert, M. Mészáros, P.R. Molnár, J.K. Moscicki, S. Roy, and J.P. Vigier. 2001. Explanation of the motionless electromagnetic generator with O(3) Electrodynamics. Foundations of Physics Letters, 14(1):87-93.
- ²¹ Bearden, T. 2002. Energy from the Vacuum: Concepts and Principles. World Scientific (In Press).
- ²² Valone, T. 2000. The Right Time to Develop Future Energy Technologies. in T. Loder (ed.), "Outside-the-Box" Technologies, Their Critical Role Concerning Environmental Trends, and the Unnecessary Energy Crisis. Report prepared for The U.S. Senate Environment and Public Works Comm. www.senate.gov/~epw/loder.htm.
- ²³ Puthoff, H. 1990. Everything for Nothing. New Scientist, 28 July 1990, pp. 52-55.
- ²⁴ Haisch, B., Rueda, A. and Puthoff, H. 1994. Beyond E=mc²; A First Glimpse of a Postmodern Physics, in which Mass, Inertia and Gravity Arise from Underlying Electromagnetic Processes. The Sciences, 34:26.
- Haisch, B., Rueda, A., and Puthoff, H. 1997. Physics of the Zero-Point Field: Implications for Inertia, Gravitation and Mass. Speculations in Science and Technology, 20:99.
- Haisch, B. and Rueda, A. 1998. An Electromagnetic Basis for Inertia and Gravitation: What Are the Implications for 21st Century Physics and Technology? in El-Genk, M. S. (ed.), Space Technology and Applications International Forum-1998, DOE CNF-980103, CP420, p. 1443.
- Haisch, B. and Rueda, A. 1999. The Zero-Point Field and the NASA Challenge to Create the Space Drive. Proc. NASA Breakthrough Propulsion Physics Workshop, NASA/CP-1999-208694, p. 55.
- ²⁵ LaViolette, P. 1993. The U.S. Antigravity Squadron. p. 82-101. In Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology.

- Integrity Research Institute, Washington, DC 20005.p.82-101.
- ²⁶ LaViolette, P. A. 1992. Electrogravitics: Back to the future. Electric Spacecraft, Issue 4, pp. 23-28.
- LaViolette, P. A. 1993. A theory of electrogravitics. Electric Spacecraft, Issue 8, pp. 33-36
- ²⁷ LaViolette, P. A. 2000. Moving Beyond the First Law and Advanced Field Propulsion Technologies. in T. Loder (ed.). "Outside-the-Box" Technologies, Their Critical Role Concerning Environmental Trends, and the Unnecessary Energy Crisis. Report prepared for The U.S. Senate Environment and Public Works Comm. www.senate.gov/~epw/loder.htm.
- ²⁸ Information available at: www.disclosureproject.org.
- ²⁹ Greer, S.M. and T.C. Loder III. 2001. Disclosure Project Briefing Document, 492 pp. Available on CD from: The Disclosure Project, P.O. Box 2365, Charlottesville, VA 22902. Also available from: www.disclosureproject.org.
- ³⁰ Greer, S. M. 2001. Disclosure: Military and government witnesses reveal the greatest secrets in modern history. Crossing Point, Inc. Crozet, VA, 573 pp.
- ³¹ Ibid. pp. 357-366.
- ³² Ibid. pp. 262-270.
- ³³ Ibid. pp. 384-387.
- ³⁴ Ibid. pp. 391-403.
- ³⁵ Ibid. pp. 388-389.
- ³⁶ Ibid. pp. 497-510.
- ³⁷ Cook, N. 2001. The Hunt for Zero Point. Available from: www.amazon.co.uk.
- ³⁸ Deavenport, L. 1995. "T.T. Brown Experiment replicated. Electric Spacecraft Journal. Issue 16. Oct. 1995. (Reprinted in: Valone, T. (ed.), 1994. Electrogravitics Systems: Reports on a new propulsion methodology. Integrity Research Institute, Washington, DC 20005)
- ³⁹ Transdimensional Technologies, 906-E Bob Wallace, Ave., Huntsville, AL 35801.
- ⁴⁰ http://jnaudin.free.fr
- ⁴¹ Greer, S.M. and T.C. Loder III. 2001. Disclosure Project Briefing Document, 492 pp. Available on CD from: The Disclosure Project, P.O. Box 2365, Charlottesville, VA 22902. Also available from: www.disclosureproject.org.
- ⁴² Arp, H. 1998. Seeing Red: Redshifts, Cosmology and Academic Science. Montreal: Aperion. (p. 249).



A97-36388

AIAA 97-3209

Some Observations on Avoiding Pitfalls in Developing Future Flight Systems

Gary L. Bennett
Metaspace Enterprises
Emmett, Idaho; U.S.A.

**33rd AIAA/ASME/SAE/ASEE Joint Propulsion
Conference & Exhibit**
July 6 - 9, 1997 / Seattle, WA

SOME OBSERVATIONS ON AVOIDING PITFALLS IN DEVELOPING FUTURE FLIGHT SYSTEMS

Gary L. Bennett*
5000 Butte Road
Emmett, Idaho 83617-9500

Abstract

A number of programs and concepts have been proposed to achieve breakthrough propulsion. As an cautionary aid to researchers in breakthrough propulsion or other fields of advanced endeavor, case histories of potential pitfalls in scientific research are described. From these case histories some general characteristics of erroneous science are presented. Guidelines for assessing exotic propulsion systems are suggested. The scientific method is discussed and some tools for skeptical thinking are presented. Lessons learned from a recent case of erroneous science are listed.

Introduction

Over the past few years a number of speculative propulsion/transportation ideas have been advanced which, if they can be verified, promise to revolutionize space transportation.¹⁻⁴ In addition, a question has been asked and some proposed answers given on the possible propulsion system(s) for unidentified flying objects (UFOs) or "flying saucers", if the existence of UFOs is accepted.^{5,6} Researchers at the U.S. Air Force (USAF) Phillips Laboratory have proposed research into "... more advanced ideas ... [such as]... fundamental physics concepts which require basic research and/or substantial development to create some sort of breakthrough. These include vacuum zero point energy, energetic species (i.e. nuclear or electronic metastables), ball lightning, and various 'breakthrough physics' concepts. In addition, a number of new emerging technologies and popular ideas will be followed such as nanotechnology, above unity devices, and cold fusion".⁷

Separately, NASA has recently initiated a "breakthrough propulsion physics" program which "... implies discovering fundamentally new ways to create motion, presumably by manipulating inertia and gravity or by harnessing other interactions of matter, fields and space-time".^{8,9}

Copyright © 1997 by Gary L. Bennett. Published by the American Institute of Aeronautics and Astronautics, Inc., with permission.

*Fellow, AIAA

Given the speculative proposals and the interest in developing breakthrough propulsion systems it seems prudent and appropriate to review some of the pitfalls that have befallen other programs in "speculative science" so that similar pitfalls can be avoided in the future. And, given the interest in UFO propulsion, some guidelines to use in assessing the reality of UFOs will also be presented.

This paper will summarize some of the principal areas of "speculative science" in which researchers were led astray and it will then provide an overview of guidelines which, if implemented, can greatly reduce the occurrence of errors in research.

Some Case Histories of Errors in Science

This section will briefly describe several case histories involving errors in science: (1) N rays; (2) Davis and Barnes experiment; (3) polywater; (4) infinite dilution; and (5) cold fusion.

N Rays

In 1903, French physicist René-Prosper Blondlot claimed to have discovered that by heating a filament inside an iron tube containing an aluminum-covered opening one could see faintly illuminated objects better in a darkened room. Some of his early experiments were performed with pieces of paper which were barely illuminated by a glowing calcium sulfide screen or a lamp shining through pinholes. Since the heated filament inside the iron tube improved the viewing, Blondlot concluded that a new type of ray was being produced, a ray which was stopped by the iron but not by the aluminum. He called this new ray an "N ray" to honor the University of Nancy, where he worked.¹⁰

Blondlot discovered that certain objects could be made to store these N rays. For example, if a brick was wrapped in black paper and exposed to sunlight it would give off N rays when taken back into the darkened laboratory. Blondlot noted that if the brick was held close to one's head it made it easier to see the paper in the darkened room. Holding the brick near the paper had a similar effect.¹⁰

Blondlot reported that the N rays had some odd effects. For example there was no improvement in the effect of improved illumination if more than one brick was used. Loud noises (such as someone entering the laboratory) could spoil the effect. Heat increased the effect but Blondlot said N rays were not the same as heat. Blondlot also claimed to have found negative N rays, which he called N' rays that could nullify the effect of the N rays. It was necessary to spend some time in the darkened room to see the effect.¹⁰

Blondlot published a number of papers on N rays. Other scientists also published papers with about half confirming Blondlot's work.¹⁰

The American physicist R. W. Wood visited Blondlot to witness some of the N ray experiments. Wood watched as Blondlot used a large aluminum prism with a 60° angle to measure the refractive index of the N rays to two or three significant figures.¹⁰

When Wood questioned Blondlot as to how he could measure the position of the beam to within 0.1 mm when the slits in the Nernst filament were 2-mm wide, Blondlot replied to the effect "That's one of the fascinating things about the N rays. They don't follow the ordinary laws of science that you ordinarily think of".¹⁰

At Wood's request, Blondlot repeated the experiments; however, unbeknownst to Blondlot in the darkened room, Wood had removed the aluminum prism. Blondlot got the same measurements! Wood subsequently published this result, effectively ending the era of the N rays.¹⁰

Later some German researchers investigated how Blondlot had been deceived. They observed that in a darkened room it is difficult to see the paper so one is tempted to pass one's hand in front of the paper to see if the paper is being illuminated. The interesting result the German researchers found was that the hand could be "observed" whether it was in front of the paper or behind it. Langmuir has remarked that this "... is the natural thing, because this is a threshold phenomenon. And a threshold phenomenon means that you don't know, *you really don't know*, whether you are seeing it or not. But if you have your hand there, well, of course, you see your hand because you *know* your hand's there and that's just enough to win you over to where you know that you see it. But you know it just as well if the paper happens to be in front of your hand instead of in back of your hand, because you don't know where the paper is but you *do* know where your hand is".¹⁰

The phenomenon of accepting a false belief and then continuing to defend it in the face of information showing the falsity of the belief is not new. In a landmark study published in 1956 it was stated that "A man with a conviction is a hard man to change. Tell him you disagree and he turns away. Show him facts or figures and he questions your sources. Appeal to logic and he fails to see your point".¹¹ As this study demonstrated, showing someone unequivocal and undeniable evidence that a person's belief is wrong means "The individual will frequently emerge, not only unshaken, but even more convinced of the truth of his beliefs than ever before. Indeed, he may even show a new fervor about convincing and converting other people to his view".¹¹

Davis and Barnes Experiment

Around 1930, two American physicists, Bergen Davis and Arthur Barnes conducted a series of experiments in which electrons from a hot filament were accelerated to match the speed of alpha particles being emitted from a polonium source. By carefully adjusting the voltage used to accelerate the electrons the experimenters claimed that they could cause the electrons to be captured by the alpha particles. The measurements were made in a darkened room by counting the number of scintillations on a zinc sulfide screen with a microscope. To save time, the experiments were focused on the voltages where it was calculated that the peaks should occur according to the Bohr theory. Davis and Barnes claimed to be able to achieve electron capture at each of the energy levels of the Bohr theory of the helium atom with measured accuracies of 0.01 volts.¹⁰

Irving Langmuir and Willis R. Whitney (then both at the General Electric Research Laboratory in Schenectady, New York) visited the Columbia University laboratory of Davis and Barnes and witnessed some of the experiments. They found that Barnes was not counting for a fixed two minutes as claimed and that he was ruling out some counts arbitrarily. When Langmuir secretly had the technician try a wider range of voltages Barnes got measurements that no longer correlated with the Bohr energy levels. Barnes had an *ad hoc* excuse for every criticism Langmuir raised about the way the experiments were being conducted.¹⁰

Langmuir, who was to earn the 1932 Nobel Prize in chemistry, subsequently wrote Barnes a 22-page letter giving Barnes all the data obtained by Whitney and himself "... and showing really that the whole approach to the thing was wrong; [Barnes] was counting

hallucinations, which I find is common among people who work with scintillations if they count for too long. Barnes counted for six hours a day and it never fatigued him. Of course it didn't fatigue him, because it was all made up out of his head".¹⁰ The next year Davis and Barnes admitted in a short article in *Physical Review* that they hadn't been able to reproduce the effect. They noted that the scintillations they were measuring were a threshold phenomenon.¹⁰

Polywater

In the 1960s and early 1970s reports from N. N. Fedyakin of the Kostrama Polytechnical Institute of the former Soviet Union and Boris V. Derjaguin of the Institute of Physical Chemistry of the Academy of Sciences of the former Soviet Union claimed there was a new form of water, later called polywater, that was more dense and viscous than normal water. Polywater, which was claimed to be formed in capillary tubes in an atmosphere nearly saturated with water, reportedly froze at -50 °C and boiled near 300 °C. Infrared spectroscopy showed that polywater produced a spectrum entirely different from normal water. Many papers were published on polywater and several theories were developed.¹²

As part of a study of polywater, Denis L. Rousseau and Sergio Porto, then at the University of Southern California, irradiated a sample of polywater with a laser to obtain a Raman spectrum. The laser turned the polywater into a black char, indicating that it "... was no polymer of water but more likely a carbonaceous material".¹² Whenever negative results like these were obtained, the proponents of polywater would always claim that the samples giving the negative results were contaminated but that their own samples, which showed polywater, were clean. Rousseau later showed that one could obtain the infrared spectrum for polywater from ordinary human sweat. This showed "... that polywater is not water at all but a product of organic contamination in the capillary tubes".¹²

Rousseau subsequently observed that "The polywater episode illustrates the loss of objectivity that can accompany the quest for great new discoveries. The quantities of polywater available were so small that many useful experiments could not be done. Many theories were put forward to describe the structure of polywater without even considering the thermodynamic difficulty of accounting for its very existence. Finally, definitive experiments showing high levels of contamination were done but not accepted, until overwhelming evidence showed that a new polymer of water had not been discovered".¹²

Infinite Dilution

In 1988 Jacques Benveniste of the University of Paris and his collaborators reported that water retained a memory or template of a molecule even when there should be no molecules present. This phenomenon is referred to as "infinite dilution" meaning that a biological effect is produced even when the "... biologically active solution is diluted so many times that no active molecules can be present ...".¹²

After Benveniste's paper was published in *Nature*, John Maddox, the editor of *Nature*, James Randi (a professional magician), and Walter Stewart (an experienced fraud investigator) spent three weeks in Benveniste's laboratory witnessing the experiments. When an elaborate series of double-blind experiments were conducted the biological effect was not measured. This committee of three reported "... that the original experiments were poorly controlled and that no effort had been made to exclude systematic error or observer bias".¹²

Many of the traits reported in the earlier episodes of erroneous science were present with infinite dilution. The effect was weak and independent of the causative agent. The counting was done visually and it was extremely difficult to do. When the control produced results at variance with the experimenters's beliefs the counting was redone. Negative results were overlooked. Sometimes experiments would not work for several months and one investigator seemed to be the best at making the experiments work.¹²

Rousseau has written that "Benveniste and his colleagues were not doing fraudulent work. They observed the effects that they reported. But they so believed in the phenomenon that they could ignore or reinterpret any questionable findings. In replying to the Maddox committee, Benveniste wrote, 'It may be that all of us are wrong in good faith. This is no crime but science as usual and only the future knows.' But, self-delusion is not science as usual".¹²

Cold Fusion

In 1989 two research groups in Utah announced that they had discovered cold fusion by means of simple electrochemical cells. Both research groups used heavy water although their salts were different. The heavy water would split into deuterioxyl ions (OD^-) and deuterons when a current was passed through the cells. What they hoped to achieve was to collect enough deuterons at the palladium cathode to cause them to pack so tightly that they would fuse at room

temperature and release energy, a process normally requiring temperatures on the order of 10^8 °C.¹²

B. Stanley Pons of the University of Utah and Martin Fleischmann of the University of Southampton reported achieving enough cold fusion that the thermal output exceeded the energy input by 4.5 times. To explain away the absence of the neutrons which should have been present at a level sufficient to kill the experimenters they reported that "... the bulk of the energy release is due to a hitherto unknown nuclear process".¹² It was later learned that their electrochemical cell was not a closed system so they did not have a good heat balance. Moreover, their method of calculating the energy release was flawed.^{13,14}

David E. Williams, the Thomas Graham Professor of Chemistry at University College London and previously a research group leader at the Harwell Laboratory of the U.K. Atomic Energy Authority responsible for that laboratory's attempts to verify cold fusion, has written about one staunch advocate of cold fusion repeating "... the common assertion that the excess energies associated with the electrolysis of D₂O with a Pd cathode are on the order of tens to hundreds of MJ/cm³ and hence can only find an explanation in some phenomenon outside the chemistry of the system. In this he displays an ignorance of the experimental measurement and a naive belief in the significance of impressively large numbers. The great majority of experiments measure power; the large numbers are obtained by multiplying an (often small) power by a large time (the duration of the experiment) and dividing by a small volume (that of the Pd cathode). In comparison with the total energy applied to the electrolytic cell, the excesses are much less impressive, on the order of a few percent. More properly, in comparison with the power applied to the cell, the claimed excess power is often also small ...".¹⁵

Steven Jones of Brigham Young University and his collaborators found evidence of increased neutron production but nothing like the heat output reported by Pons and Fleischmann.¹²

Other researchers were unable to duplicate the experiments of the two Utah groups. The neutron emissions measured by Pons and Fleischmann were inferred from a gamma-ray emission spectrum that was more narrow than the resolution of their detector. The claim that tritium, one product of a deuteron-deuteron fusion reaction, had been found was later discredited when it was learned that tritium was a contaminant in the palladium electrode.¹²

As Rousseau has summarized it, "Cold fusion was doomed from the start when a race to be first took precedence over the desire to be right. Most measurements reporting nuclear effects from cold fusion were barely above the background noise, and extended periods of failed experiments afflicted even Pons's laboratory. The proponents of cold fusion attributed the failure to several causes: differences in the materials, the size of the electrodes, impurities in the electrodes, and low current density. The list goes on.

"Nuclear reactions, however, are very well understood. Any theory offered to account for the reported observations must postulate new nuclear processes that only occur in the palladium electrodes ... The investigators of cold fusion also ignored definitive experiments".¹²

The whole sorry episode of cold fusion, which the co-chair of the Department of Energy Energy Research Advisory Board panel on cold fusion, Professor John R. Huizenga, has termed the "scientific fiasco of the century", has been well documented in Refs. 13,14, 16-18. Suffice it to say that it is not a pretty picture but as Prof. Huizenga stated: "The cold fusion fiasco illustrates once again that the scientific process works by exposing and correcting its own errors".¹³

Despite all these problems the U.S. Air Force has considered studying (as noted earlier) "... a number of new emerging technologies and popular ideas ... such as nanotechnology, above unity devices, and cold fusion".⁷ Some USAF personnel apparently foresee using cold fusion for satellite power and for fast acting thrusters for survivability. The payoffs are listed as (1) lower weight; (2) near limitless source of heat; (3) high reliability and operability; and (4) simplicity and robustness. The only technology needs identified were the scaling up of cold fusion and the development of specialized pumps, valves, and nozzles.⁷ The fact that no reputable, objective scientist has been able to prove the existence of cold fusion seems to be beside the point! (Even NASA got in the cold fusion act briefly in response to the NASA Administrator's order that no nuclear power sources be flown again on NASA spacecraft. After researchers at Lewis Research Center were unable to make cold fusion work and mission planners at the Jet Propulsion Laboratory showed to the Administrator's satisfaction that there wasn't much sunlight in the outer Solar System he recanted his order against nuclear power although he does want a more sporty -- and possibly unachievable within current funding and schedule limits -- conversion system for the next generation of radioisotope power sources.)

As shown by the original hullabaloo over cold fusion and the more recent NASA activities followed by the proposed USAF program, an unfortunate aspect to bad science such as cold fusion is that it diverts time and funding from real science. Science writer Gary Taubes has written that "What cold fusion had proven, nonetheless, was that the nonexistence of a phenomenon is by no means a fatal impediment to continued research. As long as financial support could be found, the research would continue. And that support might always be found so long as the researchers could obtain positive results. In fact, the few researchers still working in the field would have little incentive to acknowledge negative results as valid, because such recognition would only cut off their funds. It promised to be an endless loop".¹⁴

Something like this has certainly occurred with thermionic reactor research in the U.S. where hundreds of millions of dollars have been spent yet no long-lived (>5 years), high efficiency ($\geq 10\%$), high specific power ($>5.5 \text{ We/kg}$) in-reactor-core thermionic system has been built in the U.S.¹⁹⁻²¹ Still the studies continue, now with a focus on bimodal power/propulsion systems.²² And despite the investment of over \$500 million in today's dollars with no U.S. in-core thermionic reactor being built and meeting the aforementioned goals, General Atomics of San Diego, California was recently awarded a \$5.4 million cost-plus-fixed-fee contract by the Defense Special Weapons Agency "... for the research, development, test and evaluation of advanced in-core thermionic technologies to support long duration space missions for national security purposes. Focus is on the advancement of thermionic performance rather than an overall system design improvement since no specific mission has yet been identified on which to base detailed system requirements" [emphasis added].²³ Given the end of the Cold War and the lack of military reactor missions coupled with the lack of any real progress in U.S. in-core thermionic research it is doubtful if there will ever be any real military system requirements but contracts will continue to be awarded in an "endless loop". Is this pathological science or pork barrel politics -- or both?

Symptoms of Pathological Science

On 18 December 1953, Nobel-laureate Irving Langmuir gave a colloquium on the subject of what he termed "pathological science", the science of things that aren't so. As Langmuir stated it, "These are cases where there is no dishonesty involved but where people are tricked into false results by a lack of understanding about what human beings can do to themselves in the way of being led astray by subjective effects, wishful

thinking or threshold interactions".¹⁰

Based on his study of the Davis and Barnes experiment, the incident of the N rays and other early mistakes, Langmuir listed six symptoms of pathological science:¹⁰

- *The maximum effect that is observed is produced by a causative agent of barely detectable intensity, and the magnitude of the effect is substantially independent of the intensity of the cause.*
- *The effect is of a magnitude that remains close to the limit of detectability or, many measurements are necessary because of the very low statistical significance of the results.*
- *There are claims of great accuracy.*
- *Fantastic theories contrary to experience are suggested.*
- *Criticisms are met by ad hoc excuses thought up on the spur of the moment.*
- *The ratio of supporters to critics rises up to somewhere near 50% and then falls gradually to oblivion.*

Working with more recent erroneous science such as polywater, infinite dilution and cold fusion, Denis L. Rousseau, at the time a Distinguished Member of the Technical Staff at AT&T Bell Laboratories, condensed Langmuir's six symptoms into two characteristics and added a third, which he believes to be the most important:¹²

1. *The effect being studied is often at the limits of detectability or has a very low statistical significance.*
2. *There is a readiness to disregard prevailing ideas and theories.*
3. *The investigator finds it nearly impossible to do the critical experiments that would determine whether or not the effect is real.*

In looking at the cases of pathological science presented in this paper it is clear that most of them are at the limits of detectability and they often were based on subjective visual observations. The fact that there may be no direct connection between the causative agent (e.g., number of hot bricks supposedly emitting N rays) and the effect (better viewing) is not seen as an impediment to the "new" scientific "discovery". News accounts of the advertised anti-gravity effect also place

the effect at the limits of detectability.^{1,2}

Cold fusion represents perhaps the most extreme example of concocting new theories to explain a poorly understood and poorly conducted set of experiments.^{13,14} In contrast to the ready belief of the cold fusion proponents in the fantastic new physics invoked to explain their results, the author found it refreshing to hear from several of the original nuclear fission researchers (who were coincidentally meeting in a special 50th anniversary conference at the same time as the cold fusion fiasco erupted) that they were unwilling to accept their experimental results until all other avenues had been explored.²⁴ Would that other researchers had been as careful!

In none of the examples considered in the previous section did the researchers conduct experiments to determine if their results could be wrong. And in those cases where other researchers came to different conclusions they were criticized for having made mistakes or using bad samples or the wrong procedures.

Carl Sagan has written about people who make extraordinary claims, "The burden of proof is on them, not on those who might be dubious".²⁵ In short, if someone believes he/she has discovered cold fusion or anti-gravity it is up to him/her to prove it beyond a shadow of doubt. It is not up to the skeptics to disprove it. Unfortunately, as Sagan has observed pseudoscience operates just the opposite from normal science: "Hypotheses are often framed precisely so they are invulnerable to any experiment that offers a prospect of disproof, so even in principle they cannot be invalidated. Practitioners are defensive and wary. Skeptical scrutiny is opposed. When the pseudoscientific hypothesis fails to catch fire with scientists, conspiracies to suppress it are deduced".²⁶

As Sagan has so eloquently put it, "I believe that the extraordinary should certainly be pursued. But extraordinary claims require extraordinary evidence".²⁵ To date the extraordinary evidence has not been reported for cold fusion or anti-gravity.

Guidelines for Assessing Other Exotic Propulsion Systems

One writer has pondered the question of what sort of propulsion system a UFO would have to have to match the reported observations of the motions of so-called flying saucers.^{5,6} Even though people have seen strange things in the sky for centuries, the "Age of Confusion" about flying saucers began with the sighting near Mount Rainier by private pilot Kenneth

Arnold on 24 June 1947.²⁷ Following Arnold's claimed sighting there have been thousands of reports of strange craft in the skies.^{27,28}

Unfortunately, the sightings remain for the most part anecdotal and the photographs are generally suspect. Supposed artifacts have been shown to be of terrestrial manufacture. In short, while there remain some unexplained episodes, the vast majority of "sightings" are of natural phenomena or they are delusions or hoaxes.²⁵⁻²⁸

Astronomer Alan Hale, who is co-discoverer of Comet Hale-Bopp and director of the Southwest Institute for Space Research, has listed three basic principles as guides he uses in confronting beliefs about UFOs or other paranormal phenomena:²⁹

1. Extraordinary claims require extraordinary evidence.

As an example of the kind of extraordinary evidence he is seeking, Hale lists "... the actual physical aliens themselves, where I and other trustworthy and competent scientists and individuals can study and communicate with them. I'd like to examine their spacecraft and learn the physical principles under which it operates. I'd like a ride on that spacecraft."²⁹ And he lists other things he would like to see and investigate directly.

Personally, I would like to see the UFO land on the Mall in Washington, D.C. and be subjected to a complete inspection using all the resources (scientists, engineers, technicians, equipment, etc.) at humanity's disposal. Like Sagan^{25,26}, I am highly skeptical about these reported aliens in UFOs since they never seem to contact any officials, they never demonstrate or leave any technology we don't already have and they never leave any messages that we don't already know.

2. The burden of proof is on the positive.

Hale demands that the one making the claim produce the extraordinary evidence to prove the correctness of the claim, noting that the burden is not on the skeptic to prove that the advocate is wrong. He goes on to state that "... you must prove your case by providing the direct and compelling evidence for it; you can't prove it by eliminating a few token explanations and then crying, 'Well, what else can it be?'"²⁹

3. Occam's Razor: If one is confronted with a series of phenomena for which there exists more than one viable explanation, one should choose the simplest

explanation which fits all the observed facts.

Hale notes that people can make mistakes in their observations and that most people are not aware of the natural phenomena one can observe in the sky. Many people are not trained to observe what they claim to observe. People can be deceived by their preconceived notions and expectations and some people will create hoaxes. Hale asserts that "Taking all these undeniable facts together, the simplest explanation--to me, anyway--for the UFO phenomenon is that every report is either a hoax or is a mistake of some sort. If this explanation is incorrect, then you have to increase the sphere of undeniable facts; and for this, see points 1) and 2) above".²⁹

Another "breakthrough physics" concept being studied is based on the belief that energy can be extracted from the zero-point fluctuations of the vacuum.⁷ In searching for the origin of this idea, R. L Park of the American Physical Society wrote that "'The New Energy News' ... credits the idea to physicist Harold E. Puthoff and proclaimed him 'The New Energy News Theorist of the Year.' ... Puthoff's ideas are controversial; but he's accustomed to controversy. In 1972, at the Stanford Research Institute, Puthoff and Russell Targ were promoting psychic spoon-bender Uri Geller; five years later, they published 'Mind Reach,' a book about remote-viewing that inspired the CIA to invest in psychic espionage. Reportedly, Puthoff himself once sent his mind to explore the surface of planet Mercury".³⁰ Clearly, Hale's principles should be applied to this type of research!

Author's Note: For those not familiar with the *New Energy News*, it has been published by something called the Fusion Information Center which also disseminated information on cold fusion. This publication states that it is interested in "... papers ... covering both theory and practice of energy producing devices and systems such as cold nuclear fusion, rotating N-Machines, Solid-State energy systems, Magnetic over-unity machines, Tapping Space Energy (Zero-Point Energy), gravity control techniques, energetic transmutations (nuclear reactions), and other new energy research".³¹ Those interested in some of the activities of Harold Puthoff, Russell Targ and the spoon-bender Uri Geller are encouraged to read *Science, Good, Bad and Bogus* by Martin Gardner.³²

The Scientific Method

Physicist Alan Cromer has described science as "... the search for a consensus of rational opinion among all competent researchers ... The products of science give

empirical proofs of its theories. Mathematics and experimentation provide powerful arguments with which to convince and persuade".³³ Cromer states that "Science is the heretical belief that the truth about the real nature of things is to be found by studying the things themselves".³³ Cromer also notes that the scientific way of thinking is not natural to humans; it is something which has to be learned.³³ Sagan has observed that "Science is a way of thinking much more than it is a body of knowledge".²⁵

To study things requires the use of the scientific method. While there is no single absolute detailed process to be followed for all science, the American philosopher Morris R. Cohen has written that "The problem of how to get rid of illusion and see what truly goes on in nature requires that persistent and arduous use of reason which we call scientific method ... Scientific method is a systematic effort to eliminate the poison of error from our common knowledge".³⁴ In short, science is as much a process as it is an assembly of facts, but it is a process that requires assembling of the relevant facts (quantitative measurements) and the testing of hypotheses. It is a process that any scientist in the field should be able to duplicate in order to check claims of new discoveries.^{25,26,29,32-35}

To aid in avoiding pitfalls in developing advanced propulsion (or exploring any new field), Sagan's tools of skeptical thinking are highly recommended.²⁶

- *Obtain independent confirmation of the "facts".*
- *"Encourage substantive debate on the evidence by knowledgeable proponents of all points of view."*
- *Give little weight to arguments from authority.*
- *Develop and test more than one hypothesis in a fair and objective manner.*
- *Do not become overly attached to your own hypothesis.*
- *Quantify your observations wherever possible.*
- *Test the entire argument -- "If there's a chain of argument, every link in the chain must work (including the premise)--not just most of them".*
- *Use Occam's Razor -- when two or more hypotheses explain the data equally well, choose the simplest hypothesis.*
- *"Always ask whether the hypothesis can be, at*

least in principle, falsified. Propositions that are untestable, unfalsifiable are not worth much ... You must be able to check assertions out. Inveterate skeptics must be given the chance to follow your reasoning, to duplicate your experiments and see if they get the same result."

A useful guide in conducting scientific research is the National Academy of Sciences publication *On Being a Scientist*.³⁵

Lessons Learned

In the concluding chapter of his book, *Cold Fusion: The Scientific Fiasco of the Century*, Professor John R. Huizenga lists 15 lessons from which we all may profit in conducting research. These lessons are¹³

- **Handling far-out ideas and claims**

Far-out ideas and claims should be presented first informally at a meeting of colleagues to address the hard questions. Where the work involves science or technology outside the advocate's main area of expertise the briefings should include experts from those areas. When all questions have been answered and all the experimental checks made one should present the results to other experts outside the laboratory or organization and then submit to the peer-review process.

- **Judging hypotheses**

One should avoid the extreme position of rejecting all previous work in the field just because one has obtained some anomalous result. If a hypothesis requires the belief in several miraculous occurrences then the hypothesis is probably pathological.

- **Premature publication**

Researchers should submit to the full peer review process to avoid being caught later with an unsound paper. Editors and reviewers also have a responsibility to ensure the technical soundness of a paper. Where there is some justifiable need for urgent publication the editor can always put a disclaimer on the paper noting that it has not been peer reviewed. (The author proposed something like this for speculative papers submitted to the annual Intersociety Energy Conversion Conference not only as a caution for other researchers but as a warning to potential investors.)

- **Publication by press conference**

The announcement of a new scientific discovery at a

press conference should only be done where it is justified and where there is enough supporting information to back up the claimed discovery. Generally, it is far better and safer to "announce" the discovery through the normal peer review process.

- **Publication of primary data**

Researchers should publish enough of their primary data and experimental procedures to enable other researchers to check their results. If mistakes are later found these should be formally noted in the same publication that carried the original data.

- **Reproducibility in science**

Researchers should publish sufficient information about how they conducted their experiments so that other researchers can check their work. As Huizenga has written, "The foundation of science requires that experimental results must be reproducible. Validation is an integral part of the scientific process".¹³

- **Scientific isolation in research**

Breakthroughs in science generally are made by researchers who are fully knowledgeable in their fields and in contact with other researchers. In contrast, University of Utah officials claimed that the isolation of Pons and Fleischmann from both the traditional centers of fusion research and from nuclear scientists was what led them to make their claimed breakthrough discovery of cold fusion. Unfortunately, when their cold fusion results could not be duplicated by reputable, objective (and skeptical) scientists, cold fusion was dubbed another example of the "Utah Effect" (a term first used in connection with University of Utah Professor Edward Eyring's 1972 mistaken claim of discovering the x-ray laser).^{13,14}

- **Control of information**

Researchers must be open with and willing to share their research data and experimental techniques. Free and open communication is central to the scientific process.

- **Secrecy in basic research**

Basic research is best conducted openly with the peer review process. If the research is done in secret the kinds of errors that afflicted so-called cold fusion can occur. Similar problems occurred when the former Strategic Defense Initiative Organization (SDIO) tried to develop a nuclear rocket code-named "Timber Wind" using a technology (particle bed reactor) that

was highly questioned by outside experts. In that case people who had technical concerns about the viability of the concept were either ignored or denied further access to information on the program.³⁶

- **Discovery by outsiders**

There seems to be a growing belief that only people who are not experts can make breakthrough discoveries because their minds are not clouded by "official" science. The facts are quite different—it is very rare that a non-expert makes a breakthrough discovery outside his or her field. As Prof. Huizenga has observed, "Most fundamental discoveries are made by persons intimately familiar with their research discipline because they not only know the subject matter of the field but also know the pitfalls and traps and have made many of the obvious mistakes".¹³

- **Lobbying before Congressional committees**

Lobbying before Congressional committees is contrary to the scientific process of peer review and it is particularly dangerous when the advocates are championing something like cold fusion which has not been confirmed. Not only is lobbying for unproven concepts damaging to science it wastes the country's limited research funds. Prof. Huizenga has estimated that "It has taken upwards of some fifty to one hundred million dollars of research time and resources to show that there is no convincing evidence for room temperature fusion".¹³ Timber Wind cost the U.S. at least \$139 million between Government Fiscal Years 1987 and 1991 largely on the basis of Congressional lobbying.³⁶ Thermionic reactor research has cost the country on the order of \$500 million in today's dollars.¹⁹ These monies could have been better spent on peer-reviewed space technology of more benefit to the country.

- **Funding large initiatives**

Large initiatives should be funded only when there is clear, objective, peer-reviewed evidence in support of the concept.

- **Patents and revenues from basic science**

If the concern is over protecting patent and intellectual property rights, the normal process is to verify the scientific results first, then file the patents and then announce the work. The process of announcing the results first with very limited information and no scientific verification was a key contributor to the cold fusion fiasco.

- **The press and basic science**

The press should explore claims of breakthrough science thoroughly making sure that the public is being given a true picture of what is claimed. The preferred process for publicly reporting new discoveries is to wait until the research paper has been peer reviewed and accepted for publication by a respected journal in the field. Using press conferences before complete scientific papers were accepted contributed to the confusion about and eventually distrust of claims of cold fusion.

- **The scientific process**

Claims of breakthrough discoveries should be made through the usual scientific process of peer review and validation through independent reproduction of the results.

Concluding Remarks

Advanced propulsion research must be continued because of the tremendous payoff it offers to our ability to conduct scientific, exploration, and commercial space missions. However, in our studies of advanced propulsion concepts we should be guided by the scientific method so that we can avoid the types of pitfalls that have ensnared other scientists in what Irving Langmuir has called "pathological science". Physicist Richard P. Feynman said it best in his minority report on the Challenger accident: "For a successful technology, reality must take precedence over public relations, for nature cannot be fooled".³⁷ Unfortunately, what has often occurred with pseudoscience is that "... for the survival of an unsuccessful technology, public relations must take precedence over reality".¹⁴ If we ignore reality then everyone loses and our research field is damaged.

References

1. "NASA's Fling With Anti-Gravity", *Science*, Vol. 274, p. 183, 11 October 1996.
2. M. Burkey and L. David, "Changing Gravity: A Weighty Issue", *Space News*, p. 15, 11-17 November 1996.
3. J. H. Vance, "A Force Sink in Classically Defined Non-Inertial Space", AIAA paper 92-3781, prepared for the AIAA/SAE/ASME/ASEE 28th Joint Propulsion Conference and Exhibit, held in Nashville, Tennessee, 6-8 July 1992.

4. L. T. Cox, Jr., *Calculation of Resonant Values of Electromagnetic Energy Incident Upon Dielectric Spheres*, PL-TR-93-3002, Phillips Laboratory, Edwards Air Force Base, California, February 1994.
5. P. Murad, "An Electromagnetic Rocket Stellar Drive - Myth or Reality? Electromagnetic Evidence and Relativistic Phenomenon", AIAA paper 95-2602, prepared for the 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, held in San Diego, California, 10-12 July 1995.
6. P. Murad, "An Electromagnetic Rocket Stellar Drive - Myth or Reality? Fluid Dynamic Interactions and an Engine Concept", AIAA paper 95-2894, prepared for the 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, held in San Diego, California, 10-12 July 1995.
7. F. Mead and G. Olson, "Advanced Concepts for Consideration at Phillips Laboratory Propulsion Directorate", presentation given at the Sixth Advanced Space Propulsion Workshop held at the Jet Propulsion Laboratory, Pasadena, California, 21-23 March 1995.
8. L. David, "Mind Fields", *Final Frontier*, Vol. 10, No. 1, pp. 19-23, January/February 1997.
9. M. G. Millis letter to G. L. Bennett, dated 3 May 1997.
10. I. Langmuir, "Pathological Science", transcribed and edited by R. N. Hall, *Physics Today*, pp. 36-48, October 1989. See also the "Letters" column in the March 1990 and April 1990 issues of *Physics Today*. Langmuir gave this talk on 18 December 1953 at General Electric's Knolls Atomic Power Laboratory.
11. L. Festinger, H. W. Riecken and S. Schachter, *When Prophecy Fails*, University of Minnesota Press, Minneapolis, Minnesota, 1956.
12. D. L. Rousseau, "Case Studies in Pathological Science", *American Scientist*, Vol. 80, pp. 54-63, January-February 1992.
13. J. R. Huizenga, *Cold Fusion: The Scientific Fiasco of the Century*, University of Rochester Press, Rochester, New York, 1992.
14. G. Taubes, *Bad Science, The Short Life and Weird Times of Cold Fusion*, Random House, New York, New York, 1993.
15. D. E. Williams, letter to *Physics Today*, p. 94, March 1994.
16. F. Close, *Too Hot To Handle, The Race for Cold Fusion*, Princeton University Press, Princeton, New Jersey, 1991.
17. N. Hoffman, *A Dialogue on Chemically Induced Nuclear Effects, A Guide for the Perplexed About Cold Fusion*, American Nuclear Society, La Grange Park, Illinois, 1995.
18. Energy Research Advisory Board, *Cold Fusion Research*, DOE/S-0073, DE90 005611, A Report of the Energy Research Advisory Board to the United States Department of Energy, Washington, D.C., November 1989.
19. Anon., "Thermionic Space Nuclear Power, An Option for NASA Missions", General Atomics briefing package, July 1991.
20. Technology Group, "Space Nuclear Reactor Power Systems Report of Technology Group to Program Planning Group", U.S. Department of Energy, Germantown, Maryland, 18 February 1992.
21. R. J. Sovie, "Presentation on Thermionic Technology Status to SP-100 Program Review Group", Dallas, Texas, 19-20 December 1989.
22. E. L. James, W. D. Ramsey and G. J. Talbot, "Thermionic Converters for ISUS", *Proceedings of the Space Technology and Applications International Forum (STAIF-97)*, Part One, pp. 479-484, AIP Conference Proceedings 387 published by the American Institute of Physics, Woodbury, New York. Proceedings of a conference held in Albuquerque, New Mexico, 26-30 January 1997.
23. Department of Defense Contracts News Release No. 194-97 posted on the World Wide Web, 23 April 1997.
24. J. W. Behrens and A. D. Carlson, editors, *50 Years with Nuclear Fission*, 2 volumes, American Nuclear Society, Inc., La Grange Park, Illinois (proceedings of a conference commemorating 50 years with nuclear fission held at the National Academy of Sciences, Washington, D.C. and the National Institute of Standards and Technology, Gaithersburg, Maryland, 25-28 April 1989).
25. C. Sagan, *Broca's Brain, Reflections on the Romance of Science*, Ballantine Books, New York, New York, 1980. Originally published in 1978 by Random House, New York, New York.
26. C. Sagan, *The Demon-Haunted World, Science as*

a Candle in the Dark, Random House, New York, New York, 1995.

27. C. Peebles, *Watch the Skies! A Chronicle of the Flying Saucer Myth*, Smithsonian Institution Press, Washington, D.C. and London, 1994.
28. C. Sagan and T. Page, editors, *UFO's-- A Scientific Debate*, Barnes and Noble Books, New York, New York, 1996. Originally published in 1972 by Cornell University Press, Ithaca, New York.
29. A. Hale, "An Astronomer's Personal Statement on UFOs", *Skeptical Inquirer*, Vol. 21, No. 2, pp. 29-30, March/April 1997.
30. R. L. Park, "What's New for Mar 11, 1994", Internet message sent on 11 March 1994.
31. *New Energy News*, Monthly Newsletter of the Institute for New Energy, Salt Lake City, Utah. February 1994.
32. M. Gardner, *Science, Good, Bad and Bogus*, Prometheus Books, Buffalo, New York, 1989.
33. A. Cromer, *Uncommon Sense, The Heretical Nature of Science*, Oxford University Press, New York and Oxford, 1993.
34. M. R. Cohen, *Reason and Nature, An Essay on the Meaning of Scientific Method*, Dover Publications, Inc., New York, New York, 1978. The first edition of this work was originally published by Harcourt, Brace and Company, New York, in 1931.
35. Committee on the Conduct of Science, *On Being a Scientist*, National Academy of Sciences, National Academy Press, Washington, D.C., 1989.
36. Office of the Inspector General, *The Timber Wind Special Access Program*, Audit Report Number 93-033, Department of Defense, Arlington, Virginia, 16 December 1992.
37. R. P. Feynman, "Appendix F--Personal Observations on Reliability of Shuttle", in *Report of the Presidential Commission on the Space Shuttle Challenger Accident*, Volume II, Washington, D.C., 1986.

A01-34345



AIAA 2001-3660

An Experimental Investigation of the Physical Effects in a Dynamic Magnetic System

V. Roschin and S. Godin
Institute for High Temperatures
Russian Academy of Science
Moscow, Russia

**37th AIAA/ASME/SAE/ASEE
Joint Propulsion Conference and Exhibit
8-11 July 2001
Salt Lake City, Utah**

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

AN EXPERIMENTAL INVESTIGATION OF THE PHYSICAL EFFECTS IN A DYNAMIC MAGNETIC SYSTEM.

Vladimir Roschin, E-mail: rochtchin@mail.ru

Sergei Godin, E-mail: smgodin@online.ru

Institute for High Temperatures, Russian Academy of Science
Izhorskaya st. 13/19, Moscow 127412, Russia

Abstract

In the current paper the results of experimental research of magnetic-gravity effects are presented. Anomalous magnetic and thermal changes within a radius of 15 meters from the researched device were measured as well.

PACS: 41.20.-q; 44.60.+k; 76.50.+q

Introduction

We have experimentally studied the physical effects in a system based on rotating permanent magnets [1]. Below we describe the technology of manufacture, assembly, and the results of testing this experimental setup, which is referred to as the converter.

Received effects:

- Generation of mechanical energy in a self-governing mode of operations;
- Change of weight of the converter;
- Formation of a local magnetic and temperature fields as concentric cylinders around converter.

Description of the Experimental Setup

The basic difficulty arises in choosing the materials and maintaining the necessary magnetic pattern ("imprinting") on the plate and roller surfaces. To simplify the technology we decided to use a one-ring design with one-ring plate (stator) and one set of rollers (rotor). It is obvious, that it was necessary to strengthen the rollers on a rotor by the bearings and balance the rollers well. In the suggested design, air bearings were used which provided the minimum losses due to friction.

From the available description [1] it was not clear how it is possible to make and magnetize the stator with a diameter of about one meter. In order to make the stator from separate magnetized segments executed with rare earth magnets with the residual induction of 1T, the segments were magnetized in a usual way by discharging a capacitor-battery energizer through a coil. Afterwards the segments were assembled and glued together in a special iron armature, which reduced magnetic energy.

"Copyright © 2001 by Vladimir Roschin and Sergei Godin. Published by American Institute of Aeronautics and Astronautics, Inc. with permission."

To manufacture the stator 110 kg of rare earth magnets were used. To manufacture the rotor 115 kg of that material was used. High-frequency field magnetization was not applied. It was decided to replace imprinting technology described in [1] with cross-magnetic inserts having a flux vector directed at 90 degrees to the primary magnetization vector of the stator and rollers. For these cross-inserts the modified rare earth magnets with residual magnetization of 1.2 T and coercive force a little bit greater than in a base material were used. Fig.1 and Fig.2 show the joint arrangement of stator 1 and rotor, made up of rollers 2, and a way of their mutual gearing or sprocketing by means of cross magnetic inserts 19. Between the stator and roller surfaces air gap δ of 1 mm is maintained.




Fig. 1. Variant of one-ring converter.

No layered structure was used except a continuous copper foil of 0.8 mm thickness, in which the stator and rollers was wrapped up. This foil had a direct electrical contact to magnets of the stator and rollers. Distance between the inserts in rollers is equal to distance between the inserts on the stator. In other words $t_1 = t_2$ on a Fig.2.




Fig.2. Sprocket effect of magnetic stator and roller inserts.

The ratio of parameters of the stator 1 and the rotor 2 on Fig.2 is chosen so that the relation of stator diameter D to the roller diameter d is an integer equal to or greater than 12. Choosing such ratio allowed us to achieve a resonant mode between elements of a working body of the device.

The elements of magnetic system were assembled in a uniform design on an aluminum platform. Fig.3 displays the general view of the platform with the one-ring converter. This platform was supplied with springs, shock absorbers and had a possibility of moving vertically on three supports. The possible vertical displacement was 10mm and the induction meter 14 was used for measuring this displacement. Thus, the instantaneous change of platform weight was defined during the experiment in real time. Gross weight of the platform with magnetic system in an initial condition was 350 kg.

The stator 1 was mounted motionlessly and the rollers 2 were assembled on a mobile common rotor - separator 3, which is connected with the basic shaft 4 of the converter. Through this shaft the rotary moment was transferred. The basic shaft by the means of friction

muff 5 was connected to a starting engine 6, which accelerated rotor of the converter up to a mode of self-sustained rotation. An ordinary DC electrodynamics generator 7 also was connected to the basic shaft as a main loading of the converter. Adjacent to the rotor, electromagnetic inductors 8 with open cores 9 were located.

The magnetic rollers 2 crossed the open cores of inductors and closed the magnetic flux circuit through electromagnetic inductors 8, inducing an electromotive force in them, which acted directly on an auxiliary active load 10 - a set of incandescent lamps with total active power of 1 kW. The electromagnetic inductors 8 were equipped with an electrical drive 11 on supports 12. Driven coils were used for smooth stabilization of the rotor rpm. The speed of the rotor also could be adjusted by changing the main loading 10.

To study influence of high voltage on characteristics of the converter, a system for radial electrical polarization was mounted. On a periphery of rotor the ring electrodes 13 were set between the electromagnetic inductors 8 and had an air gap of 10 mm with the rollers 2. The electrodes are connected to a high-voltage source; the positive potential was connected to the stator, and the negative - to the polarization electrodes. The polarizing voltage was adjusted in a range of 0-20 kV. In the experiments, a constant value of 20 kV was used.

In case of emergency braking, a friction disk from the ordinary car braking system was mounted on a basic shaft of the rotor. The electrodynamics generator 7 was connected to an ordinary passive resistive load through a set of switches guaranteeing a step connection to the load from 1 kW to 10 kW - a set of ten ordinary electric water heaters.




Fig.3. The general view of the converter and its platform.

Converter under testing had in its inner core the oil friction generator of thermal energy 15 intended for directing a superfluous power (more than 10 kW) into the

thermo-exchange contour. But since the real output power of the converter in experiment has not exceeded 7 kW, the oil friction thermal generator was not used.




Fig.4. A diagram illustrating various operation regimes of the converter,
-G, +G changes in weight of the platform vs. rpm

Experimental results

The magnetic converter was mounted in a laboratory room on three concrete supports at a ground level. The ceiling height of lab room was 3 meter. A common working area of the laboratory was about 100 sq. meters. Besides a presence of an iron-concrete ceiling in the immediate proximity from the magnetic system there was a generator and electric motor, which contained tens of kilos of iron and could potentially deform the field structures.

The device was started by the electric motor, which accelerated rpm of the rotor. The revolutions were smoothly increased up to the moment the ammeter included in a circuit of the electric motor begin to show zero or a negative value of consumed current. The negative value indicated a presence of back current. This back current was detected at approximately 550 rpm under clockwise rotation. The displacement meter 14 starts to detect the change in weight of the whole installation at 200 rpm. Afterwards, the electric motor is completely disconnected by the electromagnetic muff and an ordinary electrodynamics generator is connected to the switchable 10 kW resistive load. The converter rotor continues self-accelerating and approaches the critical regime of 550 rpm when the weight of the whole installation quickly changes.

In addition to dependence on a speed of rotation, the common weight of device depends from output power on the load and from applied polarizing voltage

as well. As seen on Fig.4, under maximum output power is equal to 6-7 kW the change of weight ΔG of the whole platform (total weight is about 350 kg), reaches 35 % of the weight in an initial condition G_i . Applying a load of more than 7 kW results in a gradual decrease of rotation speed and an exit from self-sustained mode (right sides of the curves on Fig.4 for a 7 kW loading).

The net weight G_n of the platform can be controlled by applying high voltage to polarization ring electrodes located at a distance of 10 mm from external surfaces of the rollers. Under a high 20 kV voltage (electrodes have a negative polarity) the increase of tapped power of the basic generator more than 6 kW does not influence ΔG if rotation speed is kept above 400 rpm. "Tightening" of this effect is observed as well as the effect of hysteresis on ΔG at rotation of a rotor on a clockwise and counter-clockwise (a kind of "residual induction"). The experimental diagrams given on Fig.4 illustrate the +G and -G changes in weight of the converter vs. rotor rpm. The effect of a local change of the platform weight is reversible, relative to the direction of rotor revolution, and has the same hysteresis. A clockwise rotation causes the critical regime to occur in area of 550 rpm and a propulsion force against the direction of gravitation vector is created. Correspondingly, a counter-clockwise rotation causes the critical mode to occur in area of 600 rpm and a force in the direction of gravitation vector is created.

The difference in approach to a critical regime of 50 - 60 rpm was observed. It is necessary to mention that the most interesting region are situated above the critical area of 550 rpm, but due to of a number of circumstances the implementation of such research was not possible. Probably, there are also other resonant modes appropriate to higher rpm of a rotor and to the significant levels of useful loading and weight changing. Proceeding from the theoretical assumptions, the dependence of tapped mechanical energy from the pa-

rameters of magnetic system of the converter and rpm of a rotor has a non-linear character and the received effects are not optimum. From this point of view, the revealing of a maximal output power, of maximal change of weight and converter resource represents a large practical and scientific interest. In tested sample of the converter the using of higher rpm was inadmissible because of unsufficient mechanical durability of the magnetic system, which has been stuck together from separate pieces.




Fig.5 Diagrams of rotor accelerating and loading of the converter.

On Fig.5 the dependence of total weight of whole installation and its output power directed into active loading from rpm of a rotor of the converter is explained in detail. The diagrams are constructed for a case of HV polarization ON (top diagram) and HV polarization OFF (bottom diagram). The time from starting of the engine up to a mode of self-sustaining of the converter at the rotation of a rotor clockwise approximately is equal 1.5 minutes. Power of starting DC engine was about of 2 kW and reduction on a shaft of the converter was equal 1/10. At achievement of a critical mode (550 rpm.) the change of gross weight of a platform already achieves +/-30% from G_i . Under transition to resonant mode the revolutions with the large acceleration have increases up to 590 rpm and weight has changes up to +/-35% from G_i . This point on the diagram begins at once after a critical point (inclination of

a curve α_1). At achievement of 590 rpm the first stage of active loading 1 kW is connected to the electrodynamics generator. The revolutions have a sharply reducing and ΔG also is changing. As soon as the revolutions begin to grow again, the second switchable loading is connected and rotor rpm are stabilized at a level of 590-595 rpm. ΔG continues to change. The increasing of switchable loading occurs by steps on 1 kW up to total power of 6 kW. All intervals are equal approximately of 10-30 sec. Afterwards the short-term increasing of revolutions and then the full stabilization of the 6 kW output during of 12-15 min was observed.

More than 50 launches of the converter with an absolute repeatability within three months were carried out. It is necessary to note that revolutions will grow with acceleration shown on Fig.5 by angles $\alpha_1 \dots \alpha_5$, if do not switch on the next step of loading to the genera-

tor at rpm increasing. For returning to a previous rpm mode it is necessary to switch on a twice more loading.

The words above concern a mode with switched ON a high voltage polarization of 20 kV, "plus" is on a grounded stator. Without polarization voltage (lower curve on Fig.5) the diagram is approximately the same, but is well indicated the more soft character of a loading and faster changing of weight of a platform due to decreasing of rpm.

Other interesting effect is corona discharges, which was observed at the work of the converter in a dark room. At this, around the converter rotor a blue-pink glowing luminescence and a characteristic ozone smell were noticed. On Fig.6 the cloud of ionization covers area of the stator and rotor and is having accordingly a toroidal form.




Fig.6. Corona discharges around the converter.

On a background of luminescence glowing on the roller surfaces, we distinguished a separate «wave picture». A number of more vigorous strips of discharges around the rollers were observed. These discharges were of a white-yellow color but the characteristic sound for arc discharges was not audible. There were not noticed any erosive damages by arc discharges on surfaces of the stator and the rollers as well.

One more effect previously not mentioned anywhere was observed - the vertical concentric magnetic "walls" around the installation. We noticed and measured a permanent magnetic field around the converter within a radius of 15 meters. For this magnetic field measurement a Russian made magnetometer F4354/1 was used. Magnetometer had a Hall-effect sensor in a copper shielding. The zones of increased intensity of a magnetic flux 0.05 T, located concentrically from the center of the installation were detected. The direction of the magnetic field vector in these walls coincided with direction of magnetic field vector of rollers. The structure of these zones was like the Bessel function of zero order of two arguments. No any magnetic fields were registered between these zones by portable magnetometer. The layers of increased intensity were distributed practically without losses up to a distance of about 15 meters from a center of the converter and had a quick decreasing at a border of this zone. The thickness of each layer was approximately of 5 - 6 cm the border of each layer was very sharp. The distance between layers was about of 50 - 80 cm where the upper value is seen when moving from center of the converter. A stable picture of this field was observed as well as at a height of 6 m above the installation (on the second floor above the lab). Above the second floor, measurements were not carried out. The similar picture was observed and outside of a room of laboratory, directly in the street, on the ground. The concentric walls were strictly vertical and no had appreciable distortions. The Fig.7 illustrates the schematic placing of the converter in a room of laboratory and arrangement of concentric magnetic and thermal fields around the installation.




Fig.7. Schematic placing of the converter and field patterns in lab room.

An anomalous decrease of temperature in a vicinity of the converter was also found. While the common temperature background in laboratory was + 22°C ($\pm 2^\circ\text{C}$) the stable fall of temperature of 6-8°C was noticed. The same phenomenon was observed in concentric vertical magnetic walls as well as. The measurements of temperature inside the magnetic walls were carried out by an ordinary alcohol thermometer with an inertia of indication about 1.5 min. Inside the magnetic walls the temperature changes can even be distinctly observed by hand. When the hand is placed into this magnetic wall the cold is felt at once. A similar thermal picture was observed at height above the installation, i.e. on a second floor of the laboratory as well as despite the steel-reinforced concrete blocks of a ceiling and also on an open air outside of the laboratory.

Concentric magnetic walls and accompanied thermal effects begin to appear approximately from 200 rpm and have a linearly increasing with speeding up of revolutions up to a critical regime. The measurements above 600 rpm were not made because of fear of destruction of magnetic system. On Fig.8 the curve of intensity of magnetic field in mT and change of temperature in Celsius degrees due to rpm changing is represented.




Fig.8. The dependence of intensity of magnetic field and changes of temperature vs. rotor rpm of the converter.

Discussion

All the results we obtained are extremely unusual and require some theoretical explanation. Some theoretical approaches and phenomenological descriptions can be found in our book [2]. Unfortunately, interpretation of these results within the framework of conventional physical theory cannot explain all observed phenomena especially the change of weight. We can interpret change of weight as a local change of gravitation or

as some propulsion force like a propeller work in air. Only a role of a propeller the magnetic system is playing, and a role of air - the quantum environment - ether filled with a "dark matter and energy". Direct experiment, which would have confirming the presence of a propulsion force was not performed, but in any case both interpretations of the weight change do not correspond to the modern physics paradigm.

In conclusion we would like to emphasize that issues of biological influence and especially effects of unknown radiation around of the converter were not considered at all. Our own experience allows us to do only cautious assumption that the short-term stay (dozen minutes) in a working zone of the converter with fixed output power of 6 kW remains without observed consequences for exposed persons.

Bibliography

1. J. A. Thomas, Jr., ANTI-GRAVITY: the Dream Made Reality: the Story of John R. Searl (Direct International Science Consortium, London, 1994), Vol.VI, Iss.2.
2. V.F. Zolotarev, V.V. Roschin, S.M. Godin. About the structure of space-time and some interactions. (The theory and original experiments with extraction of an internal atomic energy). Moscow, 'Prest', 2000, 309p. with illustrations. (in Russian).

EXPLORATION OF ANOMALOUS GRAVITY EFFECTS BY MAGNETIZED HIGH-T_C SUPERCONDUCTING OXIDES

Glen A. Robertson* and Ron Litchford†

NASA Marshall Space Flight Center

Huntsville, AL

Bryan Thompson‡
TMET
Winchester, TN

Dr. Randall Peters§
Mercer University
Macon, GA

ABSTRACT

Driven by the knowledge that mass-ejection from a rocket engine is a major drawback in the exploration of space, investigations of fringe effects (or abnormalities) in known science and dealing with mass reduction was undertaken. This research, then examines the possible connection between gravity and electro-magnetic affects on the Type II, YBCO superconductor, as reported by the Russian scientist, Eugene Podkletnov. It is suggested that the quantum fluctuations of the electrons across the multitude of superconductor grain boundaries in a properly prepared Type II; superconductors may produce a measurable force on the vacuum that could counteract the effect of gravity, an acceleratory force. Within known physicists, the driving phenomena appears to relate to both the Maxwell Stress Tensor as derived by Oliver Heaviside and Woodward's transient mass theory. As a means of improving this understanding, a simplified laboratory experiment has been constructed using a modified-automated commercial Cavendish balance. The larger lead masses used in this balance was replaced by a system to EM modulate a superconductor. Tests results were inconclusive because at both room temperature and at liquid nitrogen temperatures the application of the electromagnetic (EM) or rf energy resulted in an upward climb in the data.

* Research Scientist; Propulsion Research Center

† Project Scientist; Advanced Space Transportation Program Office

‡ Research Engineer

§ Research Consultant; Department of Physics and Earth Science

"Copyright © 2001 by the American Institute of Aeronautics and Astronautics, Inc. No copyright is asserted in the United States under Title 17, U.S. Code. The U.S. Government has a royalty-free license to exercise all rights under the copyright claimed herein for Governmental Purposes. All other rights are reserved by the copyright owner."

INTRODUCTION

Engineers working in the aerospace fields dealing with rocket engine technology quickly learn that mass-ejection is a major drawback in the exploration of space. Using current rocket technology a trip to the next star would easily consume the mass-energy equivalent of a planet in order to arrive within a reasonable lifetime with reasonable hardware and expendables for the journey. Technologies like nuclear fission and fusion offer some hope, but still will not support the "Star Trek" vision of space exploration. Therefore, the NASA Propulsion Research Center at the Marshall Space Flight Center in response to the propulsion challenges specified by the NASA Breakthrough Propulsion Physics (BPP)** project, specially to investigations fringe effects or abnormalities in known science and dealing with mass reduction, proposed to explore the recently reported observation of anomalous gravitation behavior in experiments using high temperature superconductor.

The thesis of this research is to see if there is a possibility to circumvent the rocket equation without violating physical laws and to produce valid experimental data that can be used to show credible validation of these effects. The intent here is then to examining the possible connection between gravity and electro-magnetic affects on the Type II, YBCO superconductor, as reported by the Russian scientist, Eugene Podkletnov and provide a rigorous, independent, empirical conformation (or refutation) of the effect.

It has been suggested that the quantum fluctuations (or electromagnetic nature) of the electrons across the multitude of small

** Lead by the NASA Glen Research Center

superconductor grains, called Josephson junctions, in properly prepared sintered Type II, superconductors may produce a measurable force on the vacuum (i.e., space vacuum) that could counteract the acceleratory force of gravity.

The experiment that was proposed utilizes a commercially available torsion balance called a Cavendish balance, which is commonly used by physics students to measure the value of the gravitational constant. The extent of the experiment is not to measure the gravitational constant, but to measure the change in the dynamic angle induced on a torsion beam as a result of the attraction between the beam masses and external test masses (i.e., superconductor). In theory, the values of the calculated dynamic angles between two tests should be different if the masses are of different weight values. Assuming that the characteristics of the balance do not change between tests, the difference between the two dynamic angles can be used to correlate the mass change. Whereby, if a change is detected between a superconductor and an electromagnetically radiated superconductor, one can deduce that there is a possibility that an interaction with the vacuum had occurred.

The research conducted here is but a first step in the possible application of a theory into an applicable engineering space drive model, which can then be used to design a purely massless propulsion system for interplanetary applications.

A successful or null test would however, not indicate the full benefit of the phenomena nor that it is truly a physical effect. Whereby, further testing would need to be conducted to validate the results and to devise the true nature and applicability to a space propulsion system.

BACKGROUND

A number of anomalous gravitational effects (or acceleratory forces) have been reported in the scientific literature during recent years, but there has been no independent confirmation with regard to any of these claims. One such experiments was reported by the Russian scientist, Eugene Podkletnov, in which he reported anomalous weight loss (0.05–2.1%) for a variety of test masses suspended above a rotating YBCO[#], type-II superconductor.^[1,2] Further experiments using simplified apparatus without rotation have reported transients of up to 5% weight loss.^[3,4] Still, a great deal of skepticism continues to

be expressed, mainly due to uncertainties associated with experimental technique. Other researchers, for example, have yet to duplicate Podkletnov's rotating disk experiments and obtained null results in a set of simplified experiments using a stationary disk.^[5]

The technical goal was then to critically test this revolutionary physical claim and provide a rigorous, independent, empirical confirmation (or refutation) of anomalous effects related to the manipulation of gravity by rf-pumped magnetized type-II superconductors. Because the current empirical evidence for gravity modification is anecdotal, our objective was to design, construct, and meticulously implement a discriminating experiment, which would put these observations on a more firm footing within the scientific community. Our approach is unique in that we advocate the construction of an extremely sensitive torsion balance with which to measure gravity modification effects by rf-pumped type-II superconductor test masses.

Three competing theoretical explanations have been proposed to explain these gravitational anomalies: (1) gravity shielding,^[1,2] (2) absorption via coupling to a Bose condensate,^[3,4] and (3) a gravito-magnetic force.^[5-9, 18] To date, however, there has been no definitive corroboration between any of these theories and empirical observations. Therefore, it is clear that carefully designed and meticulously executed experiments are needed to explore these anomalies and to convincingly demonstrate the alleged effects. However, validation of a new theory is in itself a long and mischievous task. This is more so when you have to consider the nature of electrons at the atomic scale.

In light of the granular nature of a sintered YBCO superconductor disk, one can address the much larger grain interfaces in more macroscopic terms using electric-potentials, displacement currents, and magnetic fields. This is due to the Josephson junction effect at the interface, which is somewhat like an AC capacitor.

A search of the literature has produced several experiments using capacitors to interact with the vacuum to cause a force; 1) the Trouton and Noble (T-N) experiment^[21], 2) the Biefeld-Brown (B-B) experiment^[10], 3) the Graham and Lahoz (Heaviside) experiment^[20], and 4) the Woodward (Transient Mass) experiment^[17].

ENGINEERING APPLICATION OF QUANTUM VACUUM

[#]Yttrium, Barium, Copper, and Oxygen.

As aerospace engineers we deal more with the technology development of machines that have proven to work within the physical boundaries of known physical laws, which govern the rocket equations. Speculative theories can only lead to misunderstanding and lost paths. Therefore, space propulsive systems are designed to overcome gravitational forces by the application of time varying the mass of a vehicle, i.e., the exhausting of onboard mass at high velocities. What seems to be lost in this rational is that as the propellant becomes smaller with higher and higher exhaust velocities, as in the case of a laser or photon drive, the mass approaches a more quantum state. The logical next step would then be to connect propulsion to the quantum vacuum through acceleratory forces, i.e., gravity.

Long ago internationally renowned physicists hypothesized that gravity is an induced effect associated with zero point fluctuations (ZPF) of the quantum vacuum.^[13-14] Zeldovich first suggested that gravitational interactions could lead to a small disturbance in the non-zero quantum fluctuations of the vacuum and thus give rise to a finite value of Einstein's cosmological constant.^[13] Sakharov later derived a value for Newton's gravitational constant G using frequency w as the only free parameter.^[14]

$$G = c^5/h \int w dw \quad (1)$$

where c is the speed of light and h is the Plank constant. The integral is carried out over all frequencies using the Plank frequency on observable electromagnetic phenomena ($w_p \sim 10^{-33}$ cm) as a cutoff value.

Using this hypothesis as a basis, Puthoff has further extended Sakharov's condition in a relativistically consistent manner.^[15] As a result of this work, it is possible to envision the attractive force of gravity in terms of the radiative interaction between oscillating charges. That is, the zero point field applied to subatomic particles. From this standpoint, it is plausible that MHz frequency irradiation of superconductors rich in Josephson junction sites, as occurred in Podkletnov's experiments, could lead to a gravity modification effect through quantum ZPF interaction.

Scientific evidence continues to mount in favor of a frequency dependent interpretation of gravity as an induced effect associated with the zero point fluctuations of the vacuum. Accelerating theoretical progress combined with the anomalous gravity modification effects observed in experiments with

irradiated Type II superconductors leads one to strongly suspect a deep physical connection.

THE PODKLETNOV EXPERIMENTS

Podkletnov's gravity modification experiments were conducted in the early 1990's. Nevertheless, skepticism persists, especially since the experiments have not been adequately documented and repeated. Podkletnov reports the use of fairly large superconductor disks, 10 and 12 inches in diameter and approximately $\frac{1}{2}$ inches thick, which were magnetically levitated and magnetically rotated in the presence of an rf electromagnetic field. Samples placed over the rotating disk initially demonstrated a weight loss of 0.2–0.5%. When the rotation speed was slowly reduced, the shielding effect became considerably higher and reached a maximum reduction of 1.9–2.1%.

Of what is known of the YBCO superconductor disk used in these experiments, it seems certain that a large number of superconductor-oxide Josephson junctions exist within the disk. These types of Josephson junctions, when conversed by an ac current, emit electromagnetic waves in the rf frequency range, and when radiated at rf frequencies, generate an ac current. In a general sense, Josephson junctions are very small capacitors with the electrodes composed of superconductor material and the dielectric composed of an oxide layer.^[12] The junction is modeled as shown in figure 1, where the superconductors (SC) are small sintered grains (noting that one or both of the grains could be a normal conductor), rf is the rf energy applied, JJ is the Josephson junction site, and i is the induced or applied current.




Figure 1. SC Josephson Junction Model.

A superconductor with a structure of sintered grains would have many flux pinning sites around which Josephson junction sites exist. Flux pinning is a well-known phenomenon associated with type-II superconductors (like YBCO) where magnetic flux penetrates the superconductor and is held in place by

self-generated super-currents. Flux pinning results from any spatial inhomogeneity of the material, such as impurities, grain boundaries, voids, etc. where the magnetic flux can become trapped. To be most effective, these inhomogeneities must be on the scale of the order of the penetration depth or the coherence length, i.e. ~10.6 to 10.5 cm, rather than on the atomic scale where inhomogeneity causes electronic scattering which limits the mean free path.^[12]

The Josephson junction sites at the grain boundaries would generally not produce flux pinning due to the resistive nature of the boundary. The exceptions might be under very high static magnetic field conditions. Radiation of the sites with rf energy would allow current flows, but could cause the flux to vibrate and jump from one site to another depending on the frequency.

Experiments conducted on both sintered and melt textured YBCO superconductors for the purpose of magnetic flux compression have shown that the rapidly moving flux with a millisecond rise time to approximately 1 kilogauss penetrates the superconductor with little (<35 gauss) compression of the field.[¶] This would indicate that a magnetic field could easily move through the body of a sintered superconductor to produce currents within the grain structure. Contradictory to this, it has also been shown during tests to repeat the Podkletnov experiment that an AC magnetic field will levitate a sintered (12 inch) disk.⁺⁺

Podkletnov's experiment used both AC magnetic fields and rf energy. It would therefore seem that Podkletnov has produced a device to enhance the production of rf energy and rf energy to enhance the production of superconductor currents. These reinforcing phenomena should lead to high electron densities in the superconductor disk, focused at the Josephson junction sites and generated at the junction frequency.

It is then suggested here that the quantum fluctuations of the electrons across the multitude of Josephson junctions in properly prepared Type II, superconductors may produce a measurable force on the vacuum that could counteract the effect of gravity.

OTHER RELATED EXPERIMENTS

The capacitive like nature of the Josephson junction would make one wonder if other experiments have been conducted using capacitors to affect the vacuum. Research of the literature indicates that experiments using capacitors as a

coupling mechanism to the quantum vacuum is not a new idea. In 1904, Trouton and Noble (T-N) reported that a mechanical force could be detected from a charged capacitor, which was free to rotate^[21]. And in 1929, Townsend Brown reported translational motion using the now famous Biefeld-Brown (B-B) effect, which utilizes capacitors with extremely high electrical potentials (>70 kV)^[10]. To the author's knowledge,

[¶] Conducted by the first two authors.
⁺⁺This work is being conducted under a NASA SBIR Phase II. no one has report a successful duplication of the B-B experiment.^[11] However in 1998, Cornille, Naudin, and Szames reported a successful duplication of the T-N experiment also using voltages near 70 kV.^[22]

Trying to connect these two experiments to Podkletnov's experiment is somewhat deceptive as only statically charged capacitors with no magnetic fields were used. One could speculate that the leakage current across the dielectric medium could occur at some (low) frequency associate with the atomic electron energy states. Also, stray magnetic fields could have been present; at the least, the magnetic field of the earth was. From a more physical sense, the time varying magnetic fields in the Podkletnov experiment would have created time varying high electrical potentials in the superconductor.

In more recent times, Graham and Lahoz (in 1980) reported the use of a coaxial capacitor to produce rotational motion from the vacuum by setting up a non-vanishing Poynting vector, as Maxwell and Poynting foresaw and predicted by Heaviside's^{**} time variation of Maxwell's equations.^[20] Further, Woodward has recently done some very interesting work with capacitors, both theoretically and experimentally to validate the notion of a transient mass effect.^[17] These two experiments do have similarity to the Josephson junction and a further clarification follows.

Heaviside Force

Oliver Heaviside in 1886 obtained an express from the divergence of the Maxwell stress tensor, which is a vector with units of force density (N/m^3), and therefore implies momentum transfer. Corum names this the Heaviside force f_H and gives it in vector form as

^{**}Referred to as the Heaviside Force by Corum^[19].

$$f_H = \frac{\partial(DxB)}{\partial t}, \quad (2)$$

where D is the electric displacement and B is the magnetic induction^[19].

Corum goes on to present a space-drive that was first presented in an essay by Joseph Slepian in 1949. In the essay, Slepian models the space-drive by employing an rf source to drive two solenoids and a parallel-plate capacitor electrically wired in series. The rf energy was directed between the plates and perpendicular to the electric field of the plates. In this arrangement, the current passing through the coils must also cross the capacitor. This is shown in figure 2.

In the Slepian model, one may say that the current can only cross the capacitor in the presence of an rf field. This is the case for a superconductor Josephson junction. In fact, Slepian space-drive model is a crude approximation of a Josephson junction and is only a short stretch to the junction model of figure 1.




Figure 2. Slepian Space Drive Model.

Transient Mass Shifts

Woodward has come up with an equation for a transient mass shift (TMS) derived from Mach's Principle (Woodward, 1991). Woodward presented the transient mass shift δm_0 in general terms as:

$$\delta m_0 = \frac{bwP_0}{2pGr_0c^2}, \quad (3)$$

where δm_0 is the transient mass; b is the ratio F/c^2 (F is the gravitational potential due to all the matter of the universe) and is approximately 1 and unitless; w is the frequency of the driving voltage into the capacitors in radians per second; P_0 is the power applied to the capacitors in Watts; G is the gravitational constant = 6.673×10^{-11} N m²/kg²; r_0 is the density of the capacitors; and c is the velocity of light = 2.9979×10^8 m/s.

A connection between Woodward's transient mass and Podkletnov's gravity modification

experiment was presented in a previous paper.^[16] In the paper, a model of the Josephson junction, transient mass relationship was given similar to figure 3.

As with figure 1, figure 3 presents a two-grain Josephson junction where one grain is a normal conductor NC and the other is superconductive SC. The current has been represented as a function $f(i)$ due to the uncertainty of this mechanism (i.e., time varying fields and rotation). The prospective is that electron charges e formed in the normal conductor. The application of the rf energy at the appropriate frequency allows the flow of electron to cross the junction as pairs $2e$. Noting the reverse effect is also possible.




Figure 3. Josephson Junction TMS Model.

As cited in the previous paper, Woodward's TMS formula has commonality with Modanese's anomalous coupling theory (ACT)^[3,4] and Woodward's capacitor experiment has commonality with the layered superconductor disk of Podkletnov's second experiment where the top part was a superconductor and the bottom portion a normal conductor. The TMS formula derives a mass fluctuation from a time-varying energy density. The ACT suggests that the essential ingredient for the gravity phenomenon is the presence of strong variations or fluctuations of the Cooper pair density (a time-varying energy density). Woodward's experiment used a small array of capacitors whose energy density was varied by an applied 11 kHz signal. When these are vibrated up and down at the correct frequency so that they are going up when their mass is minimum and going down when their mass is maximum, then a small, constant, mass-force change is possible. Podkletnov's superconductor disk contained many Josephson junctions, which were radiated with a 3-4 MHz signal. At the layered interface, the Cooper pairs are moving upward, while the electron pair separations are moving downward.

These commonalities allow for ease in rewriting

superconductor mass shift ∂m as

$$\partial m_{sc} = \frac{\mathbf{b} f_{jj} P_{jj}}{G \mathbf{r}_{sc} c^2}, \quad (4)$$

where f_{jj} is the resonance frequency (in Hz) of the superconductor Josephson junctions, P_{jj} the effective combined power (in watts) of all the junctions, and \mathbf{r}_{sc} is the density of the superconductor. Equation 4 then represents the mass change of the superconductor.

EXPERIMENTAL APPROACH

Repeating the original Podkletnov experiment has been a major undertaking within the Marshall Space Flight Center for several years. Confusion over the original experimental design and the ability to produce the large superconductors have been the major problems. Given these problems a much simpler experimental approach was devised to investigate the possible gravity connection. This approach has not been without problems.

The approach involved the replacement of the large (~1 kg) lead masses in a commercially available computerized torsion or Cavendish balance with a system that magnetically modulates an YBCO superconductor. A sketch of the proposed experiment is given in figure 4.




Figure 4. Sketch of the Cavendish balance experiment.

The measurement of the gravitation constant G ($6.67 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$) using a Cavendish balance is a simple experiment, which is routinely performed by beginning physics students. Using Newton's universal law of gravitation, it is possible to express the angular displacement \mathbf{f} of the beam in terms of directly measurable quantities

$$\mathbf{f} = \frac{2GMml}{R^2 \mathbf{k}} \quad (5)$$

where M is the test mass, m is the mass attached to each end of the beam, l is the separation length of the beam mounted masses, R is the distance from the center of each test mass to the center of each beam mounted mass, and \mathbf{k} is the torsion constant of the fiber supporting the beam (not shown).

Corrections

The use of the Cavendish balance as supplied with the large lead masses requires several corrections. Such as a correction for the gravitational torque on the beam and the cross torque between the opposite masses. These corrections are easily approximated and are applied to the smaller masses. Other corrections such as the variations in R can be averaged out over many data points.

In this experiment, the spherical large lead masses are being replaced by a much larger mass that is more like a rectangular box. Such a shape makes the calculation of these corrections much more complicated without extensive testing. The simple solution is to look at the things that are measurable versus those that are not and see how these change from one condition to the other. Equation 5 is then rewritten as,

$$k_c \left(\frac{M}{R^2} \right) = \left(\frac{\mathbf{k}}{2Gml} \right) \mathbf{f} \quad (6)$$

where k_c is the correction to m the small mass.

Equation 6 can then be used in the formulation of a percent mass change $M\%$ even though the left side of the equation is unknown. For example, a percent mass change $M\%$ between a non-modulated superconductor mass M_1 and a modulated superconductor mass M_2 is just the ratio of the angular displacements \mathbf{f}_1 and \mathbf{f}_2 , given by

$$M\% = \frac{M_2}{M_1} = \frac{\mathbf{f}_2}{\mathbf{f}_1} \quad (7)$$

where the angular displacements are measurable.

Sensitivity

Equation 6 also is true for two different tests using two different mass weights. The change in the measured angular displacement of the torsion fiber will then be directly proportional to the change in the test mass. For example, the change in angular displacement $d\mathbf{f}$ associated with an effective change in the test masses dM is given by

$$\frac{d\mathbf{f}}{dM} = K_c G \quad (8)$$

The sensitivity of the device is therefore dependent on the magnitude of

$$K_c = \left(\frac{k_c}{R^2} \right) \frac{2ml}{k} \quad (9)$$

$$K_c = \left(\frac{1}{R^2} \right) \left(\frac{2T^2}{lp^2} \right) \quad (10)$$

which is independent of the smaller mass and any correction to it as long as the period T is measurable.

Based on the published characteristics of the as delivered Cavendish balance a numerical estimate for the sensitivity was determined to be

$$\frac{d\mathbf{f}}{dM} = 0.3 \text{ microradians / gram} \quad (11)$$

for $T = 120$ sec, $l = 30$ cm, and $R = 4.6$ cm.

The commercially available Cavendish balance was chosen because it contained a computerized electronic detector known as a symmetric differential capacitive (SDC) control unit** to electronically measure the position of the beam as it rotates. The SDC is easily capable of measuring a displacement angle of 1 micro-radian. Therefore for a 0.5% weight change of a 300-gram superconductor, a displacement angle of 1.5 micro-radians would be detectable.

Testing has shown that the calculated displacement angle varies not with M but with M/R^2 . This is due to the uneven mass density of the superconductor containment system or modulator that will be described later. That is, as the mass of the superconductor (or test mass) changes, the average density displacement of the modulator also changes, which in turn shifts the position of the modulators center of mass. This makes the determination of the percent mass change or equation 7 more difficult to determine. However, since we are only concerned at this point with seeing a change in the relationship between tests, the sensitivity to $1/R^2$ in equation 9 is a plus. In the general sense, the sensitivity as given in equation 11 is enhanced by the square of the difference in the shifted R values as a result of the test mass weight change.

Test Apparatus

** The SDC is an invention of Dr. Randall Peters; Research Consultant.

In order to operate the balance in the hand-off operation inside a liquid nitrogen cyro-tank, the balance was fitted into a structure and the motion of the large masses was automated using a National Instruments, nuDrive (model 4SX-411) stepper motor controller and a National Instrument's rack (PXI-1010) chassis that contains a PXI-8136 D/A converter (PXI-8136 D/A p/T)² where using Labview 5.1 software written specifically for this operation. The modified balance is shown in figure 5.

The large masses were replaced by a system, referred to as the modulator, which was required to modulate the superconductor sample with electromagnetic (EM) or rf energy. The modulator was composed of a superconductor, permanent magnets, an iron frame, a kHz coil, and an MHz antenna. Electrodes were placed adjacent to the superconductor to detect the hall current induced in the superconductor. A magnetic shield (1006 steel) was added due to an attraction problem with the



balance's aluminum beam. A picture of the modified balance with the modulators is shown in figure 6 and a sketch of the modulator is shown in figure 7.

Figure 5. Automated Cavendish Balance.




Figure 6. Modulator & Balance.




Figure 7. Sketch of Modulator.

The iron frame forms a magnetic circuit that directs the field through the superconductor and parallel to the balance's beam. The placement of the coil around the iron frame induces a time varying kHz field in the static field. The MHz antenna is placed near the coils also inducing a time varying field in the static field. The effect of these time varying fields were measured perpendicular to the static field.

The purpose of the static magnetic field produced by the permanent magnets was to induce currents in the superconductor about pinned flux sights and possibly the weaker holes about the grains, which would be much higher than that seen by the AC levitated superconductor in the Podkletnov experiment. The kHz modulation field would then act to break the pinning sites freeing the pinning currents to move according to the motion of the magnetic field in the superconductor material. In theory, the radiated MHz field reduces the resistance of the Josephson junction between the grain boundaries and to allow resistiveless passage of the currents throughout the superconductor.

The configuration of the superconductor then determines the mode of operation. That is, if the superconductor is composed entirely of sintered grains, the Heaviside force at the Josephson junction sights will dominate any gravity effect. On the other hand, if a non-superconducting, conductive layer is placed on the outward side of the sintered superconductor away from the balance; the electron motion across the boundary will produce a mass transit effect.

Instabilities

Several months were spent after the initial completion of the automated balance in determining

and eliminating instabilities caused by the automation mechanism. The major mechanical problem was caused by the support apparatus, which would bend downward (ah gravity) as the modulator moved through the zero position. This allowed the spur gear to hit the support structure. Repositioning of the gear only made it hit other structures. The problem was fixed by grinding the top and bottom of the spur gear at an angle.

Placement of the drive motor also presented a problem. Due to the lack of support perpendicular to the balance beam, lead to the introduction of vibration. Placement of the motor such that the shaft rotation was in the plane of the modulator's motion and adding support structures provided a major reduction in the induced vibrations.

The deduction of these problems and the resulting fix thereof was hindered by the enclosure of the spur gear and the fact that some of the vibrations could only be detected by the analysis of the data. A data run was typically done overnight to allow the balance to stabilize. Three to five data cycles were typically required before stabilization occurred. One data cycle of about 35 minutes was required to get one dynamic angle measurement.

The only problem to arise during cooling in liquid nitrogen was with the electrical connections. Cooling below 170C caused intermittent signal disturbances. This was fixed by insulating the connections exposed to the liquid nitrogen temperatures.

TESTS RESULTS

Only one type of superconductor sample has been tested. It was composed of two layers; one of YBCO and one of PrBCO (Pr – Praseodymium). The substitution of Pr for Y was done to cause the layer to be a conductor with similar crystal structure to the YBCO. This sample was fabricated by the same manufacture producing the samples for the repeat of the Pokletnov experiment under a NASA SBIR phase II. Tests were conducted at room temperature and at liquid nitrogen (-196 C) temperature.

The YBCO superconductor only required < -180 C to become superconductive. The type K thermocouples used in the experiments flat lined between -186 C and -188 C in liquid nitrogen. Therefore to insure that the superconductor was in a superconductive state, tests were not conducted until a temperature < -186 C was optioned. The following graphs (figure 8 – 12) represent a data set using the same torsion wire. The numbers across the bottom (x

– axis) of each chart refers to the number of cycles from which each dynamic angle was calculated. The sequence is in order of time from start to finish of a data run. Each dynamic angle (or cycle) depended on the period of the torsion beam, but typically was between 30 and 40 minutes. The voltage value on the left (y -axis) of each chart refers to the calculated dynamic angle. The values have been left in its voltage value because the conversion factor for the control unit was stable between tests.

The results of the superconductor tests are base lined against a copper (Cu) sample.

No-Modulation

The results of the room temperature and the liquid nitrogen (i.e., superconductive) tests for the non-modulated or static magnetic field cases are shown in figure 8 and figure 9, respectfully.

As seen in the room temperature tests of figure 8, the calculated dynamic angles between the varying weighted masses increased with decrease mass weight due to the non-uniform density of the modulator (i.e., $1/R^2$). Additional tests were conducted using other weights composed of fiberglass epoxy, aluminum, and lead, which showed the same result.




Figure 8: Dynamic Angle of copper sample (Cu) and superconductor sample (PR123) at room temperatures.

In the liquid nitrogen tests of figure 9, the tests were started when the superconductor was at superconductive temperatures and allow to warm-up over night. As shown, at the non-superconductive temperature of – 175 C there is a noticeable change in the values of the dynamic angles. (Approximately 0.06 v for the superconductor sample and 0.05 for the copper sample.) Earlier tests down to –170 C with the original lead masses in the balance, also show good results in the calculation of the gravitational constant, which changed by less than 2% from the room temperature value.

Instabilities noted at the beginning of the liquid nitrogen test of the superconductor sample warranted a repeat at the lower temperature. This was conducted immediately following the first test as not to disturb the balance. Figure 10 shows the data from the repeated run.

Figures 9 and 10 then show that the superconductor and the copper samples produced similar results at the lower temperatures.

EM Modulation

EM modulation tests were conducted at room temperatures and at liquid nitrogen temperatures for only the superconductor. These tests are shown in




Figure 9: Dynamic Angle of copper sample (Cu) and superconductor sample (PR123) from superconductive

temperatures to non-superconductive temperature.




Figure 10: Dynamic Angle of the superconductor sample (PR123) at superconductive temperatures.

figure 11 and figure 12. The room temperature tests of figure 11 shows the effect of the MHz and kHz frequencies separately. In the liquid nitrogen test shown in figure 12, both the MHz and KHz frequencies are used together.

In both cases, the calculated dynamic angle increased over time. In the liquid nitrogen tests, the first two data points are with no EM modulation. This was done to detect a change before the boil off of the liquid nitrogen due to the rf heating of the iron frame, which was partially submerged in the liquid nitrogen. Rapid boil off reduced the superconductive run time from approximately six hours with no EM modulation down to two hours with EM modulation.





Figure 11: Dynamic Angle of the superconductor sample (PR123) at room temperatures with EM energy applied.




Figure 12: Dynamic Angle of the superconductor sample (PR123) at superconductive temperatures with EM energy applied.

CONCLUSIONS AND RECOMMENDATIONS

To summarize, we note that these exploratory experiments have been carried out in an attempt to quantify the effects of EM energy on a superconductor. The general conclusion is that the results of these tests gave a null result. That is, no conclusion at this time can be made to the EM effects on the superconductor. This conclusion is reached based on the increasing dynamic angle over time in both the room temperature and liquid nitrogen temperature tests.

Further, it is concluded that the balance is sensitive to mass changes at room temperature and down to approximately -175 C but not when the temperature is < -186. This conclusion was reached based on the similarities in the data for both the copper and superconductor samples in figure 9. However because no time varying temperature data was taken on these tests, further testing is required to pin point the actual shift point between -175 C and -186 C.

If a temperature at which the superconductor becomes superconductive and within the sensitivity of the balance is determined, it is recommended that

non-EM modulated tests on the masses reported here and on a non-layered superconductor be conducted. Regardless of the results, a redesign of the balance is recommended to eliminate the EM modulation effects on the balance control unit and to reduce the heating effect of the modulator.

REFERENCES

- [1.] E. Podkletnov and R. Niemen, :"A Possibility of Gravitational Force Shielding by Bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ Superconductor," *Physica C*, Vol. 203, 1992, pp. 441 - 444.
- [2.] E. E. Podkletnov, "Weak gravitation shielding properties of composite bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ superconductor below 70K under e.m. field," cond-mat/9701074 v3, 16 Sept. 1997.
- [3.] G. Mandanese, *Europhys Lett.*, Vol. 35, 1996, p. 413; *Phys. Rev. D*, Vol. 54, 1996, p. 5002.
- [4.] Modanese, Giovanni, "On the theoretical interpretation of E. Podkletnov's experiment," LANL gr-qc/9612022, Presented for the World Congress of the International Astronautical Federation, 1997, nr. IAA-97-4.1.07.
- [5.] N. Li, D. Noever, T. Robertson, R. Koczor, and W. Brantley, "Static Test for a Gravitational Force Coupled to Type-II YBCO Superconductors," *Physica C*, Vol. 281, 1997, pp. 260-267.
- [6.] N. Li and D. G. Torr, *Phys. Rev. D*, Vol. 43, 1990, p. 457.
- [7.] N. Li and D. G. Torr, "Gravitational effects on the magnetic attenuation of superconductors," *Phys. Rev. B*, Vol. 46, 1992, p. 5489.
- [8.] D. G. Torr and N. Li, "Gravito-Electric Coupling Via Superconductivity," *Found. Phys. Lett.*, Vol. 6, 1993, p. 371.
- [9.] C. S. Unnikrishnan, "Does a superconductor shield gravity," *Physica C*, Vol. 266, 1996, pp. 371-383.
- [10.] T. T. Brown, "How I control gravitation," *Science & Invention Magazine*, 1929.
- [11.] R. L. Talley, "Twenty-First Century Propulsion Concept", Phillips Laboratory (Propulsion Directorate), Air Force Systems Command, Final Report No. PL-TR-91-3009, Project 3058, 1991.
- [12.] M. Tinkham and G. McKay, *Introduction to Superconductivity*, McGraw-Hill, Inc., 1996.
- [13.] Ya. B. Zeldovich, *JETP Letters*, Vol. 6, 1967. p. 345.
- [14.] A. Sakharov, "Vacuum quantum fluctuations in curved space and the theory of gravitation, *Sov. Phys. Doklady*, Vol. 12, 1968, pp. 1040-1041.
- [15.] H. E. Puthoff, "Gravity as a zero-point-fluctuation force," *Physical Review A*, Vol. 39, No. 5, pp. 2333-2342, 1989.
- [16.] Glen A. Robertson, "Search for a Correlation Between Josephson Junctions and Gravity," Space Technology and Applications International Forum - 2000, pp. 1026-1031.
- [17.] J. F. Woodward, "Mach's Principle of Weight Reduction = Propellantless Propulsion," *Foundation of Physics Letters*, Vol. 9, No. 3, 1996, pg. 247 – 293.
- [18.] Agop, A., C. Gh. Buzea, and P. Nica, "Local Gravitoelectromagnetic Effects on a Superconductor," *Physica C*, 339, pp. 120-128, 2000.
- [19.] Corum, James F., John P. Dering, Philip Pesavento, and Alexana Donne, ""EM Stress-Tensor Space Drive," "Space Technology and Applications International Forum 1999," American Institute of Physics, AIP Conference Proceedings 458. (The papers on BPP topics are in pages 875-937 and 954-1059.)
- [20.] G. M Graham and D. G. Lahoz, "Observation of static electromagnetic angular momentum in vacuo," *Nature* Vol. 285, pg 154-155, May 1980.
- [21.] F. T. Trouton and H. R. Noble, "The mechanical forces acting on a charged condenser moving through space," *Philosophical Transactions of the Royal Society of London*, 202A, 1904, pp. 165-181.
- [22.] Patrick Cornille, Jean-Louis Naudin, and Alexandre Szames, "Stimulated Forces Demonstrated: Why the Trouton-Noble Experiment Failed and How to Make it Succeed," Space Technology and Applications International Forum – 1999, pp. 1005-1013.



A01-34136

AIAA 2001-3361

Antimatter Production at a Potential Boundary

Michael LaPointe
Ohio Aerospace Institute
Cleveland, OH 44135

**37th AIAA/ASME/SAE/ASEE Joint Propulsion Conference
Salt Lake City, UT
July 8 – 11, 2001**

For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

ANTIMATTER PRODUCTION AT A POTENTIAL BOUNDARY

Michael LaPointe*
Ohio Aerospace Institute
Cleveland, OH

ABSTRACT

Current antiproton production techniques rely on high-energy collisions between beam particles and target nuclei to produce particle and antiparticle pairs, but inherently low production and capture efficiencies render these techniques impractical for the cost-effective production of antimatter for space propulsion and other commercial applications. Based on Dirac's theory of the vacuum field, a new antimatter production concept is proposed in which particle-antiparticle pairs are created at the boundary of a steep potential step formed by the suppression of the local vacuum fields. Current antimatter production techniques are reviewed, followed by a description of Dirac's relativistic quantum theory of the vacuum state and corresponding solutions for particle tunneling and reflection from a potential barrier. The use of the Casimir effect to suppress local vacuum fields is presented as a possible technique for generating the sharp potential gradients required for particle-antiparticle pair creation.

INTRODUCTION

Present chemical engines and electric propulsion thrusters are well suited for near-Earth applications and robotic space flight, but advanced propulsion technologies must be developed to enable fast piloted and robotic deep space missions. Of all the known energy sources, none provides more specific energy than the annihilation of matter and antimatter. The energy released per kilogram of combined matter and antimatter is nearly 250 times the specific energy released in nuclear fusion, and over 8 orders of magnitude greater than the specific energy released in chemical combustion.¹ The possibility of producing photon rockets using gamma rays from electron-positron annihilation was investigated over half a century ago,² but the efficiency of the engines were

curtailed by an inability to collimate the energetic photons. With the experimental discovery of the antiproton in 1955, attention turned to the use of proton-antiproton annihilation as an energy source for spacecraft propulsion. The higher rest mass energy of the proton-antiproton pair yields 1877 MeV per annihilation event, compared with 1.02 MeV released by electron-positron annihilation. Equally important, a significant fraction of the proton-antiproton annihilation energy appears in the kinetic energy of charged particles,³⁻⁵ which may be collimated for direct thrust or used to heat an expellant more effectively than electron-positron gamma radiation. Several antiproton-powered rocket designs have been proposed over the past few decades, ranging from low thrust, high specific impulse pion engines to higher thrust, lower specific impulse solid and gas core thermal rockets.⁶⁻²⁰ Recent modeling efforts have simulated the performance of magnetically confined hydrogen plasma engines heated by charged proton-antiproton annihilation byproducts²¹⁻²⁴ and have investigated antiproton-boosted fission reactions as a driver for an inertial confinement fusion rocket.^{25,26}

Although a number of potential antiproton propulsion concepts have been analyzed, their transition from theoretical design to experimental validation and practical use has been constrained by the prohibitive cost of creating and storing the antiprotons. The following section discusses current antiproton production methods, and outlines near-term prospects for efficient antiproton production and storage.

Current Antiproton Production Methods

The two leading facilities for antiproton production and storage are the European Laboratory for Particle Physics (formerly CERN, the Center for European Nuclear Research) in Geneva, and the Fermi National Accelerator Laboratory (FNAL) in the United States. At the

* Senior Research Associate; Member, AIAA

This material is declared a work of the U.S. Government and is not subject to copyright protection in the United States.

CERN facility, protons are accelerated by a linear accelerator to 50 MeV (8×10^{-12} J), injected into a booster ring and accelerated to 800 MeV, and then sent to a proton synchrotron, where they are further accelerated to 26 GeV. The high-energy protons are then focused into a 2-mm beam and directed into a 3-mm diameter, 11-cm long copper wire target. The relativistic protons collide with the target nuclei, producing a spray of gammas, pions, kaons, and baryons, including antiprotons. On leaving the target, the antiprotons have a peak momentum of 3.5 GeV/c, corresponding to a peak energy of roughly 3 GeV. A short focal length, pulsed magnetic horn is used to capture antiprotons that have momenta within 1.5% of their peak value, at angles up to 50 mrad from the target centerline. The captured antiprotons are sent to a storage ring in bursts of about 10^7 antiprotons every few seconds, and around 10^{11} antiprotons can be accumulated before space charge effects scatter the circulating beam. The antiprotons are sent back to the proton synchrotron, which decelerates them to an energy of 200 MeV, and then to the low energy antiproton ring, where the circulating beam is further decelerated, stochastically cooled, and stored. Similar techniques are used to create antiprotons at FNAL.

During the high-energy collisions, approximately one antiproton is created for every 10^5 - 10^6 high-energy protons incident on the target. The energy efficiency, defined as the energy released in a proton-antiproton annihilation event (1.88 GeV at rest) divided by the energy required to create an antiproton, is abysmally low. On average, CERN creates 1 antiproton for every 2.5×10^6 protons; at an average energy of 26 GeV per proton, the corresponding energy efficiency is approximately 3×10^{-8} . FNAL, which uses a 120 GeV proton beam to strike the target, creates 1 antiproton for every 3.3×10^4 protons, corresponding to an energy efficiency of around 4×10^{-7} . Assuming a "wall-plug" efficiency for each accelerator of around 5%, the total antiproton production efficiencies are roughly 1.5×10^{-9} for CERN and 2×10^{-8} for FNAL. The total annihilation energy contained in 1-mg of antiprotons (roughly 6×10^{20} antiprotons) is 1.8×10^{11} J; an efficiency of 1.5×10^{-9} means that it would take nearly 1.2×10^{20} J (3.3×10^{13} kW-hr) to create 1 mg of antiprotons. Assuming a conservative energy cost of \$0.05/kW-hr, the estimated production cost is a staggering $\$1.6 \times 10^{11}$ per milligram of antiprotons. Most antimatter propulsion concepts require milligrams to grams of antiprotons, indicating that current antiproton production techniques are inadequate for future spacecraft propulsion applications. However, as discussed by Forward¹ and Schmidt *et al.*,²⁷ neither CERN nor FNAL were designed as dedicated antiproton production facilities. As such, a

number of upgrades to the current facilities could be made to improve antiproton production and storage capabilities. Magnetic fields produced by electric currents flowing through the metal wire targets could be used to keep the spray of antiprotons closer to the target axis, reducing their angular spread. Multiple targets could be employed, with magnetic lenses used to refocus the antiprotons between each section. Angular capture efficiencies could be improved by going to higher beam energies, creating a forward-peaked distribution that allows more antiprotons to be captured. Material lenses could be replaced with current-carrying plasma lenses, which are less likely to absorb the antiprotons and would not need active cooling. Using linear rather than synchrotron accelerators to produce the initial high-energy proton beams could increase the accelerator energy efficiency by an order of magnitude over the current 5% wall plug efficiencies.

Taken together, the potential facility improvements could result in the yearly production and storage of microgram quantities of antiprotons at a potential cost²⁷ of around $\$6.4 \times 10^6/\mu\text{g}$ ($\$6.4 \times 10^9/\text{mg}$). While these production numbers and costs are approaching those required for ground testing antimatter propulsion concepts, they are not adequate for antimatter-based propulsion systems. Forward^{1,20} calculates that antiproton propulsion becomes cost competitive with chemical propellant systems at an antiproton production cost of approximately $\$10^7/\text{mg}$, and antiproton propulsion becomes the most cost effective propulsion source available if the production costs can be lowered to $\$2 \times 10^6/\text{mg}$. Because the near term facility modifications outlined above are unlikely to produce the necessary reduction in antiproton production costs, a number of alternative antiproton production techniques have been suggested. Chapline²⁸ has proposed colliding heavy ion beams, made up of singly charged uranium atoms, to produce up to 10^{18} antiprotons/sec. Unfortunately, the antiprotons will be emitted isotropically and will be very difficult to collect. Equally problematic, the colliding heavy ion beams will produce a significant amount of nuclear debris and radiation, which would have to be safely and efficiently removed from the spray of antiprotons. Cassenti²⁹ has suggested that the pions generated during the collision of high-energy protons with heavy target nuclei could be redirected toward the target to increase the number of antiprotons and improve the efficiency of current antiproton production techniques. Although promising, the collection and redirection of the pions and antiprotons remains a major challenge to this concept.

Hora³⁰ proposed the use of a high intensity laser that could generate sufficiently strong electric fields to pro-

duce proton-antiproton pairs from the vacuum, and Crowe³¹ separately proposed the use of high intensity lasers to produce electron-positron pairs. At present, however, there are no known lasers that can produce the high intensity electric fields needed for pair production. Forward^{1,20} and Haloulakos and Ayotte³² have investigated the possibility of building and operating an antiproton factory in space, where the proton accelerator could be powered by solar energy. However, the estimated cost to produce and store the antiprotons is still nearly \$10⁹/mg, which is a factor of 10² too high for cost-effective space propulsion applications.³²

Rather than rely on high-energy proton beam collisions with a stationary target, this paper outlines a new concept that may lead to the more efficient production of antimatter in quantities sufficient for propulsion and other commercial applications. The proposed technique is based upon particle-antiparticle pair production at the steep potential boundary created by the suppression of local vacuum field energies. The premise is based on Dirac's relativistic theory of the vacuum state, which is outlined in the following section. The theory underlying particle-antiparticle pair creation at a potential boundary is discussed, followed by an explanation of the technique proposed to create the required potential step. The paper concludes with an overview of an experimental approach designed to demonstrate the feasibility of this new antimatter production concept.

DIRAC'S THEORY OF THE VACUUM STATE

Dirac was the first to develop a relativistic wave equation that correctly describes the interaction of spin-1/2 particles, such as electrons and protons.³³ Dirac's equation contains both positive and negative energy solutions, the latter identified with the continuum energy of the vacuum state (Fig. 1). As defined by Dirac, the vacuum state is characterized by the absence of all real electrons in positive energy states, but has electrons filling all negative energy states (the "Dirac sea"). Because of the Pauli exclusion principle, real electrons cannot transition into negative energy states since all such states are already occupied; however, an electron in a negative energy state can absorb radiation and transition to a positive energy state, leaving behind a "hole" in the negative energy continuum. The hole behaves like a positive electron and represents the antiparticle of the electron. The creation of an electron and an anti-electron (positron) is identified as pair creation and requires a minimum energy of $2m_e c^2$. Pair annihilation occurs when an electron drops back into the (unoccupied) hole, with the resulting transition energy emitted as radiation.




Figure 1. Energy Levels of the Dirac Equation

More generally, in Dirac's theory the vacuum represents a continuum of negative energy states occupied by negative energy particles. Pair creation is the process in which sufficient energy is given to a particle in the negative energy state to raise it to a positive energy state (creating a real particle and leaving behind a hole, or antiparticle); annihilation occurs when the particle falls back into the hole, with the energy carried away as radiation. The vacuum itself should have zero energy, zero mass, and no charge, which is clearly not satisfied by the simple form of the theory. Instead, there are infinitely many negative energy states, which together have an infinitely large negative energy, and, in the case of electrons populating the negative energy continuum, an infinitely large negative charge. These difficulties are removed by renormalizing the zero point of charge and energy in such a way that the vacuum has no mass, energy, or charge. This renormalization process is not pleasing from an aesthetic viewpoint, but it does satisfy the constraint that only departures from the vacuum state are observable and hence relevant.

TUNNELING AND POTENTIAL BARRIERS

Related to Dirac's theory of the vacuum is the quantum mechanical process of particle tunneling in the presence of a steep potential step. An overview of this process is provided by Greiner,³⁴ the salient features of which are given here.

Consider a spin-1/2 particle (for example, an electron or proton) with energy, E, and momentum, p, traveling along the z-axis (Figure 2). The particle encounters a step potential of magnitude V_0 that rises to full value in a distance equal to the Compton wavelength of the particle, λ_c :

$$\lambda_c = \frac{h}{m_0 c} \quad (1)$$

where h is Planck's constant (6.626×10^{-34} J-s), m_0 is the

particle rest mass, and c is the speed of light.




Figure 2. Particle incident on a potential step.

The Dirac equation describing the propagation of the particle in Region I is:

$$i\hbar \frac{\partial \psi}{\partial t} = \left[\frac{\hbar c}{i} \left(\hat{\alpha}_1 \frac{\partial}{\partial x^1} + \hat{\alpha}_2 \frac{\partial}{\partial x^2} + \hat{\alpha}_3 \frac{\partial}{\partial x^3} \right) + \hat{\beta} m_0 c^2 \right] \psi = H_f \psi \quad (2)$$

where ψ is the particle wave function, \hbar is the reduced Planck constant ($h/2\pi$), H_f is the Hamiltonian, and $\hat{\alpha}$, $\hat{\beta}$ are the standard Dirac matrices. Noting that the momentum operator \vec{p} is given by:

$$\vec{p} = \frac{\hbar c}{i} \left(\hat{\alpha}_1 \frac{\partial}{\partial x^1} + \hat{\alpha}_2 \frac{\partial}{\partial x^2} + \hat{\alpha}_3 \frac{\partial}{\partial x^3} \right) = \frac{\hbar}{i} \nabla \quad (3)$$

the Dirac equation can be written in the more compact form:

$$i\hbar \frac{\partial \psi}{\partial t} = [c(\hat{\alpha} \cdot \vec{p}) + \hat{\beta} m_0 c^2] \psi = H_f \psi \quad (4)$$

The Hamiltonian for Region I (zero potential) is the total particle energy, E , while in Region II the Hamiltonian becomes $(E - V_0)$. The Dirac equation for a particle wave traveling along the $+z$ direction in Region I is then:

$$[c(\hat{\alpha}_3 \cdot \vec{p}_1) + \hat{\beta} m_0 c^2] \psi = (E - V_0) \psi \quad (5)$$

In Region II, the Dirac equation for the traveling particle wave becomes:

$$[c(\hat{\alpha}_3 \cdot \vec{p}_1) + \hat{\beta} m_0 c^2] \psi = E \psi \quad (6)$$

The solution for the particle wave function in Region I is:

$$\psi_I = A \begin{pmatrix} 1 \\ 0 \\ \frac{p_1 c}{E + m_0 c^2} \\ 0 \end{pmatrix} \exp \left[\frac{i p_1 z}{\hbar} \right] \quad (7)$$

where A is a constant and the particle momentum, $p_1 c$, is given by:

$$p_1 c = \sqrt{E^2 - m_0^2 c^4} \quad (8)$$

At the potential boundary, part of the particle wave will be reflected and part will be transmitted. The reflected wave solution in Region I is:

$$\psi'_I = B \begin{pmatrix} 1 \\ 0 \\ \frac{-p_1 c}{E + m_0 c^2} \\ 0 \end{pmatrix} \exp \left[\frac{-i p_1 z}{\hbar} \right] \quad (9)$$

and the transmitted solution in Region II is:

$$\psi_{II} = D \begin{pmatrix} 1 \\ 0 \\ \frac{-p_2 c}{V_0 - E - m_0 c^2} \\ 0 \end{pmatrix} \exp \left[\frac{i p_2 z}{\hbar} \right] \quad (10)$$

where again B and D are constants. The particle momentum in Region II is given by:

$$p_2 c = \sqrt{(E - V_0)^2 - m_0^2 c^4} \quad (11)$$

The incident and reflected wave functions must equal the transmitted wave function at the step boundary ($z=0$):

$$\psi_I|_{z=0} + \psi'_I|_{z=0} = \psi_{II}|_{z=0} \quad (12)$$

from which the following conditions are obtained for the coefficients A , B and D :

$$A + B = D \quad (13)$$

$$\begin{aligned} A - B &= -D \frac{p_2}{p_1} \left(\frac{E + m_0 c^2}{V_0 - E - m_0 c^2} \right) \\ &= -D \sqrt{\frac{(V_0 - E + m_0 c^2)(E + m_0 c^2)}{(V_0 - E - m_0 c^2)(E - m_0 c^2)}} = -D\gamma \end{aligned} \quad (14)$$

where γ is defined as:

$$\gamma = \sqrt{\frac{(V_0 - E + m_0 c^2)(E + m_0 c^2)}{(V_0 - E - m_0 c^2)(E - m_0 c^2)}} \quad (15)$$

Upon rearrangement, Equations 13 and 14 yield:

$$\frac{B}{A} = \left(\frac{1-\gamma}{1+\gamma} \right) \quad (16)$$

Dividing Equation 14 by the coefficient A and substituting Equation 16 for (B/A) yields:

$$\frac{D}{A} = \frac{2}{1-\gamma} \quad (17)$$

The particle current j is defined to be:

$$j(z) = c\psi^\dagger(z)\hat{\alpha}\psi(z) \quad (18)$$

where $\psi^\dagger(z)$ is the adjoint of $\psi(z)$. The values of $\psi^\dagger\hat{\alpha}$ in Region I are:

$$\begin{aligned} \psi_I^\dagger\hat{\alpha}_1 &= A^* \left(0, \frac{p_1 c}{E + m_0 c^2}, 0, 1 \right) \exp \left[\frac{-ip_1 z}{\hbar} \right] \\ \psi_I^\dagger\hat{\alpha}_2 &= A^* \left(0, -i \frac{p_2 c}{E + m_0 c^2}, 0, -i \right) \exp \left[\frac{-ip_1 z}{\hbar} \right] \\ \psi_I^\dagger\hat{\alpha}_3 &= A^* \left(\frac{p_1 c}{E + m_0 c^2}, 0, 1, 0 \right) \exp \left[\frac{-ip_1 z}{\hbar} \right] \end{aligned} \quad (19)$$

from which j_I , the incident particle current in Region I, is:

$$j_I = -AA^* \frac{2p_1 c^2}{E + m_0 c^2} \quad (20)$$

Similarly, the reflected (j_I^r) and transmitted (j_{II}) particle currents are:

$$j_I^r = -BB^* \frac{2p_1 c^2}{E + m_0 c^2} \quad (21)$$

$$j_{II} = DD^* \left(\frac{-2p_1 c^2}{V_0 - E - m_0 c^2} \right) \quad (22)$$

Equations 20-22 can now be used to calculate the reflection and transmission coefficients for the particle wave function impacting the potential boundary. Taking the ratio of the reflected current to the incident particle current yields:

$$\frac{|j_I^r|}{|j_I|} = \left| \frac{BB^* \frac{2p_1 c^2}{E + m_0 c^2}}{AA^* \frac{2p_1 c^2}{E + m_0 c^2}} \right| = \left| \frac{BB^*}{AA^*} \right| = \left| \frac{B^2}{A^2} \right| = \frac{(1+\gamma)^2}{(1-\gamma)^2} \quad (23)$$

The ratio of the transmitted current to the incident current is given by:

$$\begin{aligned} \frac{|j_{II}|}{|j_I|} &= \left| \frac{DD^* \frac{2p_2 c^2}{V_0 - E - m_0 c^2}}{AA^* \frac{2p_1 c^2}{E + m_0 c^2}} \right| = \left| \frac{DD^*}{AA^*} \right| \gamma \\ &= \left| \frac{D^2}{A^2} \right| \gamma = \frac{4\gamma}{(1-\gamma)^2} \end{aligned} \quad (24)$$

For a potential step $V_0 > (E + m_0 c^2)$, the value $\gamma > 1$. From Equation 23, this indicates that the reflected particle current exceeds the incident particle current in Region I ($|j_I^r| > |j_{II}|$). It appears that electrons are entering Region I from Region II, but there are no electrons initially present in Region II. This result, known as Klein's Paradox, is most often interpreted as particle-antiparticle pair creation at the potential boundary.

Discussion of Results.

Applying a potential $V_0 > E + m_0 c^2$ raises the energy in Region II sufficiently for there to be an overlap between the negative energy continuum ($z > 0$) and the positive energy continuum ($z < 0$), as shown in Figure 3:




Figure 3. Energy Continuum of the Dirac Equation at a Potential Barrier.³⁴ (x) = particle, (o) = antiparticle

When $V_0 > E + m_0 c^2$, the particles striking the barrier from the left are able to knock additional particles out of the vacuum on the right, leading to an antiparticle current flowing from the left to the right in Region II and a particle current flowing from right to left in Region I. This pair creation is depicted schematically in Figure 3, with the additional particles entering Region I from the right accounting for the increase in return cur-

rent. As noted by Greiner,³⁴ this process is most readily understood as particle-antiparticle pair creation at the potential barrier and is related to the decay of the vacuum in the presence of supercritical fields. When the potential function V_0 is less than $E + m_0c^2$, the particle momentum in Region II is imaginary (Eq. 11) and the wave solution will be exponentially damped (Eq. 10); all of the incident current is then reflected back into Region I, and no particle current is transmitted into Region II. It is only when $V_0 > E + m_0c^2$ that the momentum in Region II becomes real and the particle wave function in Region II again becomes a traveling wave.

The above derivations may be summarized as follows:

- The Dirac equation is used to represent the evolution of spin-1/2 particle wave functions.
- Dirac's equation permits both positive and negative energy solutions; the negative energy states are filled with virtual particles, which prevent particle transitions from positive energy states to negative energy states via the Pauli exclusion principle.
- Particles will be completely reflected from a potential barrier (V_0) when their energy $E < V_0$; the transmitted particle wavefunction is exponentially damped within the potential barrier.
- For potential steps with $V_0 > E + m_0c^2$, an incident particle will induce pair creation at the potential boundary, resulting in a (real) return particle current and a (real) transmitted antiparticle current.

The question now arises as to whether this effect can actually be applied to the production of antimatter. The potential step must be greater than $E + m_0c^2$, where E is the total particle energy (rest mass plus kinetic), and the potential must rise to its full value over a distance comparable to the Compton wavelength of the particle, $\hbar/(m_0c)$. For an electron, the minimum potential step height is 1.02 MeV (plus the kinetic energy of the electron), and the Compton wavelength is approximately 2.4×10^{-12} m. For a proton, the minimum potential step height is 1876 MeV and the Compton wavelength is approximately 1.3×10^{-16} m. These supercritical potentials are too large to be generated over such short distances using laboratory electric fields, but there may be another option: rather than use externally applied fields to raise the vacuum energy in Region II, it may be possible to use the Casimir effect to lower the vacuum energy in Region I. This concept, shown schematically in Figure 4, may be able to produce the same pair creation effects depicted in Figure 3 without requiring the application of supercritical external fields. A brief discussion of the Casimir effect is provided below, followed by an outline of a possible experimental test of the pro-

posed antimatter production mechanism.




Figure 4. Lowering the vacuum energy via the Casimir effect.

VACUUM FIELDS AND THE CASIMIR EFFECT

Dirac's vacuum is a negative energy continuum populated by particles that prevent positive energy particles from transitioning into the negative energy states. To avoid obvious problems associated with infinite vacuum charges and energies, the vacuum state is renormalized to zero; only deviations from the vacuum state are measurable. As such, the potential step shown in Figure 3 represents an applied field measured with respect to the background vacuum; the energy in Region II has been raised by the applied field such that the negative energy states now overlap the positive energy states in Region I. However, raising the background vacuum in Region II requires a tremendous amount of energy ($V_0 > E + m_0c^2$) over a very small distance, which is clearly beyond present capabilities. Rather than raising the vacuum energy in Region II, it is proposed that the same effect can be generated by suppressing the relative vacuum energy in Region I, as shown in Figure 4. This process can be viewed either as lowering the positive energy states such that they now overlap with the negative energy states in Region II, or the background vacuum in Region I can be renormalized to zero to yield an energy diagram similar to Figure 3. In either instance, a particle wave traveling from Region I into Region II will be described by the same solutions outlined in the section above, leading to particle-antiparticle pair creation at the potential boundary. Unlike Figure 3, pair production is achieved not by raising the potential but by lowering the relative vacuum energy, an effect that might be accomplished through the use of a Casimir cavity.

Before discussing the Casimir effect, it should be noted that Dirac's interpretation of the vacuum as a continuum of negative energy states occupied by negative energy particles, though somewhat dated, is not in disagreement with the current quantum electrodynamic (QED) interpretation of the vacuum as an infinite sea of electromagnetic radiation populated with virtual particle pairs. Because exchange interactions occur in Dirac's theory, virtual electron-positron pairs are continuously created and annihilated in the vacuum; an electron in a bound or free state can fill a virtual hole in the Dirac sea, with a virtual electron taking its place. Renormalization of the vacuum energy and charge is required both in Dirac's original theory and in QED, and as noted by Greiner,³⁴ the physical content of the Dirac theory forms the basis of current quantum electrodynamics. This is mentioned because the Casimir effect is generally discussed in terms of the QED interpretation of the vacuum state, and it is necessary to point out that both the Dirac and QED vacuum interpretations are complementary.

The Casimir Effect

As in the Dirac theory, the vacuum state in quantum electrodynamics is interpreted to be the state of lowest energy. This lowest energy state is not at rest, but fluctuates with a "zero-point" energy.³⁵⁻³⁹ The vacuum fluctuations have measurable effects, including the experimentally observed Lamb-Rutherford shift between the *s* and *p* energy levels of the hydrogen atom, and the attractive Casimir force that occurs between closely spaced uncharged conductors. As discussed by Mostepanenko *et al.*,³⁵ the Casimir effect can be accounted for by assuming the force is a consequence of the separation-dependent vacuum field energy trapped between the conductors. For example, assume that two square conducting plates with side dimensions *L*, separated by a distance *z*, are placed in a vacuum. In the QED interpretation, the vacuum is teeming with electromagnetic radiation (although mathematically the vacuum state is renormalized to zero), hence the plates may be considered to constitute a cavity that supports vacuum fluctuation modes with wave numbers down to about *z*⁻¹. The vacuum energy trapped between the plates is approximately given by:

$$U = \sum_{k=\frac{1}{2}}^K \frac{1}{2} \hbar \omega \approx (L^2 z) \int_{z^{-1}}^K \hbar c k^2 dk \cong \frac{1}{4} L^2 \hbar c [z K^4 - z^{-3}] = U_U - U_L \quad (25)$$

where *U_U* is the upper energy bound, *U_L* is the lower energy bound, and *K* represents a high frequency cut-off to make the total energy finite. The negative rate of change of the lower cut-off energy *U_L* with separation *z*

constitutes a force of attraction, *F*, per unit area, given by:

$$\frac{F}{A} = -\frac{1}{L^2} \frac{dU_L}{dz} \approx \frac{\hbar c}{z^4} \quad (26)$$

A more careful analysis leads to the exact relationship for the force per unit area between the parallel conducting plates:³⁹

$$\frac{F}{A} = \frac{\pi^2 \hbar c}{240 z^4} \quad (27)$$

In other words, the vacuum energy density between the plates is lower than the vacuum energy density outside the plates by an amount equal to the right hand side of Equation 27. For reasonable plate sizes and small separation distances, the change in the vacuum energy density can be appreciable. Assuming the plates have length *L*=0.1 m and are separated by a distance of 1-μm (10^{-6} m), the change in the vacuum energy density is approximately 1.3×10^{-4} J/m³, corresponding to an inward pressure of 1.3×10^{-4} N/m² on the plates. For a separation distance of 0.1-μm (10^{-7} m), the change in the vacuum energy density is equal to 1.3 J/m³, corresponding to an inward pressure of 1.3 N/m². The calculations can be carried down to separation distances that are approximately equal to the cut-off wavelength of the conducting material, at which point the plates can no longer be considered good electromagnetic reflectors.

Equations 26 and 27 demonstrate that the vacuum energy density between the conducting plates is lower than the external vacuum energy density. Multiplying Equation 27 by the plate surface area (*L*²) and the separation distance between the plates (*z*) yields an expression relating the decrease in the vacuum energy between the plates compared to the external vacuum field energy:

$$\Delta E_{VAC} = -\frac{\pi^2 \hbar c L^2}{720 z^3} \quad (28)$$

For square plate dimensions of *L* = 0.1 m and a separation distance of 0.1-μm, the change in vacuum energy is calculated to be 4.3×10^{-9} J, or roughly 2.7×10^{10} eV (27 GeV). The vacuum energy between the plates is thus substantially lower than the vacuum energy external to the plates, or conversely, if the vacuum energy between the plates is renormalized to zero, the vacuum energy external to the plates is substantially higher than the vacuum energy between the plates. By adjusting the plate dimensions and separation distance, it may thus be possible to significantly suppress the vacuum energy in a given region and generate a condition similar to that shown in Figure 4. A particle generated in the suppressed vacuum fields of Region I will see a higher vacuum field energy outside of the plates, and if the relative change in the vacuum energy exceeds $E + m_0 c^2$, it may be possible to generate particle-antiparticle pairs

at the vacuum energy step. Because the Dirac solutions hold equally well for electrons or protons, the possibility exists that low energy proton-antiproton pairs might be created at the steep potential boundary created by vacuum field suppression in a Casimir cavity.

Creating a Potential Gradient

The Casimir effect provides an avenue for creating sufficiently large potential steps, but the question arises as to whether these steps can be generated over a distance comparable to the Compton wavelength of the particle. From Equation 1, the Compton wavelength for electrons is 2.42×10^{-12} m, while for protons the Compton wavelength is 1.32×10^{-15} m. Either distance is several orders of magnitude smaller than any realistic plate thickness separating the interior and exterior vacuum fields, hence typical flat plates will not provide a sufficiently steep potential step for pair creation to occur.

A possible solution to this dilemma is to create a suitable potential gradient over a distance larger than the Compton wavelength, such that the required step change in potential occurs over a distance comparable to the Compton wavelength. Assuming the required potential step has magnitude V and the change in vacuum field energy due to the Casimir effect has magnitude ΔE_{vac} , the gradient relation can be expressed:

$$\frac{V}{\lambda_c} \approx \frac{\Delta E_{vac}}{d} \quad (29)$$

where λ_c is the Compton wavelength of the particle and d is the plate thickness, or more properly the thickness of the region over which the change in vacuum field energy occurs. For pair production to occur at the step boundary, the potential V must exceed $E + m_0 c^2$, where E is the total energy of the particle (rest mass plus kinetic energy). Assuming the particle kinetic energy at the boundary is small compared to its rest mass energy, the requirement for V becomes:

$$V \geq 2m_0 c^2 \quad (30)$$

Inserting Equations 28 and 30 into Equation 29 yields the following expression for the plate area, thickness, and separation as a function of particle mass and Compton wavelength:

$$\frac{2m_0 c^2}{\lambda_c} \approx \frac{\pi^2 \hbar c L^2}{720 z^3 d} \quad (31)$$

which reduces to:

$$\frac{L^2}{z^3 d} \approx 4.15 \times 10^{44} \frac{m_0}{\lambda_c} \quad (32)$$

Equation 32 expresses necessary conditions for a flat parallel plate geometry to form a sufficiently steep potential gradient for particle-antiparticle pair production to occur. For electrons, this condition can be written:

$$\frac{L^2}{z^3 d} \approx 1.56 \times 10^{26} (m^{-2}) \quad (33)$$

Assuming a plate separation distance $z = 10^{-7}$ m and a gradient length $d = 10^{-6}$ m, the required plate area, L^2 , is around 0.156 m^2 ; for square plates this corresponds to a side length of approximately 0.4 m. Plates of this size and flatness are well within current manufacturing capabilities, offering some encouragement for experimental verification of the proposed concept. For protons,

$$\frac{L^2}{z^3 d} \approx 5.26 \times 10^{32} (m^{-2}) \quad (34)$$

Using the same plate separation and gradient distances above would require a plate length of 725 m to provide a suitable potential gradient. However, a number of Casimir force experiments have been performed over the past several years with plate separations down to several nanometers, and a reasonable lower limit of 10^{-8} m can be assumed for the plate separation z . The distance over which the potential gradient is formed, d , can also be reduced by locally thinning the plate support structure; a more realistic lower bound on d is thus assumed to be 10^{-9} m. Given these values, the plate size for producing the required potential step for proton-antiproton pair production is around 0.73 m, which is difficult but not impossible to manufacture.

Ideally, the potential gradient would be formed in a region devoid of plate material; this can be accomplished by designing the Casimir plate with a small hole whose diameter is of the same order or smaller than the plate separation distance. Vacuum electromagnetic fields with wavelengths larger than the hole diameter will still be blocked by the cavity, but the potential gradient formed by the plates is now in a material-free region that more faithfully reproduces the assumptions behind Figures 3 and 4. Possible effects due to the fringing of vacuum electromagnetic fields at the hole boundaries remain to be evaluated, but this method appears promising to provide the required potential gradients in a material-free region.

Summary. In summary, it appears possible to produce a Casimir cavity geometry that will provide sufficiently steep potential gradients for particle-antiparticle pair creation. By introducing a small hole in the plate material with a diameter similar to the plate separation distance, a potential step may be created in a material-free region, as depicted in Figure 4. Electron-positron pairs and proton-antiproton pairs can conceivably be gener-

ated using this technique, with significantly less infrastructure and presumably lower cost than current antimatter production methods.

EXPERIMENTAL DESIGN

The following experiment is proposed to evaluate the possibility of producing particle-antiparticle pairs at a potential boundary created within a Casimir cavity. The proposed experiment is a modified version of a similar experiment performed by the author under a prior contract to the NASA Marshall Space Flight Center.⁴⁰ In that experiment, flat parallel plates were used to form a Casimir cavity to investigate pair creation, but the plate geometries were not properly designed to generate sharp potential gradients. Based on that effort and the additional analysis in this report, the following experiment is suggested for a proof-of-concept test.

Electron-Positron Production

To investigate the formation of electron-positron pairs, it is proposed that a Casimir cavity be constructed from two flat, square metallic plates, each with an area of $1.61 \times 10^{-2} \text{ m}^2$ ($L = 12.7 \text{ cm}$). From Equation 33, spacing the plates a distance $z = 10^{-7} \text{ m}$ apart should provide a suitably steep potential gradient for electron-positron pair creation to occur. To create a material free region for the steep potential step, one plate should include a central hole, drilled perpendicular to the plate boundary, of diameter $d \leq 10^{-7} \text{ m}$; vacuum electromagnetic fields with wavelengths larger than d will still be blocked by the cavity, although there may be minor effects on the potential step due to field fringing. On the opposite plate, a small amount of radioactive material can be deposited to act as a source of electrons within the cavity; Ni⁶³ is a readily available commercial source, and the kinetic energy of the emitted electrons (0.067 MeV) are sufficiently below the rest mass energy of the electron that Equation 30 remains a viable approximation. Alternatively, the metallic plate can be irradiated to produce subsequent electron emissions within the cavity.

The plate surfaces must be aligned and moved to within 10^{-7} m to form the required potential step, which can be accomplished using commercially available piezoelectric transducers. The close separation distance requires that the surface flatness of the plates not exceed 10^{-8} m , which is a stringent but commercially attainable constraint. The entire system should be mounted in a vacuum system capable of achieving a hard vacuum ($\leq 10^{-8} \text{ torr}$), and isolated from vibrations. To evaluate whether the proposed pair production method works, a small target placed outside the central plate hole can be used

to intercept any positrons emitted at the potential boundary. The resulting annihilation of the positrons with the target material will produce 0.511-MeV gamma rays that can be measured with a detector located behind the target.

Proton-Antiproton Production

If successful, the experimental arrangement outlined above will demonstrate the basic feasibility of the proposed pair production process. However, for propulsion applications it is desirable to produce antiprotons rather than positrons. As previously discussed, the constraints on plate size, flatness, and separation become significantly more demanding, but remain within the capability of current manufacturing techniques. For proton-antiproton pair production, square plates with areas of 0.526 m^2 ($L = 0.725 \text{ m}$) would have to be separated by a distance of 10^{-8} m , indicating that the surface flatness of the plates would have to be on the order of angstroms. Larger plate areas would relax this constraint by allowing larger separation distances, and the ability to machine flat surface areas must be traded against the fabrication of larger plate dimensions. The central hole diameter would again be on the order of the plate separation distance to provide a material-free potential step, and the electron source would be replaced with a proton source to provide the particles within the cavity. If this scheme is successful, the antiprotons can be captured upon exiting through the hole in the Casimir plate and stored in portable Penning traps for later use.

CONCLUDING REMARKS

A new concept has been described for creating matter-antimatter particle pairs at a steep potential boundary. The potential step is created using the Casimir effect to suppress the vacuum energy between parallel conducting plates. Preliminary calculations indicate that a sufficiently steep potential gradient can be formed for reasonable plate dimensions and separation distances. A preliminary experimental design is outlined as a proof-of-concept test for the proposed antimatter production scheme. Additional analysis remains to be performed to validate the concept, including an evaluation of material and temperature effects on the plate boundaries, the effect of fringing fields on the potential step at the plate central hole, and the consequences of imperfect parallel plate alignment on the required potential gradient. Nevertheless, based on the preliminary analysis presented in this paper, the proposed concept appears to be a potentially viable alternative to the high-energy antimatter production methods currently in use.

ACKNOWLEDGEMENTS

Funding for this research was provided through NASA SBIR Contract NAS8-98109 during the period April – November 1998.

REFERENCES

1. Forward, R.L., *Antiproton Annihilation Propulsion*, AFRL-TR-85-034, Forward Unlimited, Oxnard, CA, 1985.
2. Sanger, E., "Zur Theorie der Photoneraketen" ("The Theory of Photon Rockets"), *Ingenieur-Archiv.*, **21**, 1953, pp. 213-226.
3. Morgan, D.L. and Hughes, V.M., "Atomic Processes Involved in Matter-Antimatter Annihilation", *Phys. Rev.*, **D-2** (8), 1979, pp. 1389-1391.
4. Agnew, L.E., Jr., Elioff, T., Fowler, W.G., Lander, R.L., Powell, W.M., Segre, E., Steiner, H.M., White, H.S., Wiegand, C., and Ypsilantis, T., "Antiproton Interactions in Hydrogen and Carbon Below 200 MeV," *Phys. Rev.*, **118** (5), 1960, pp. 1371-1391.
5. Morgan, D.L., *Annihilation of Antiprotons in Heavy Nuclei*, AFRL TR-86-011, Lawrence Livermore National Laboratory, Livermore, CA, Apr 1986.
6. Massier, P., "The Need for Expanded Exploration of Matter-Antimatter Annihilation for Propulsion Applications," *J. Br. Interplanetary Soc.*, **35** (9), 1982, pp. 387-390.
7. Forward, R.L., "Antimatter Propulsion," *J. Br. Interplanetary Soc.*, **35** (9), 1982, pp. 391-395.
8. Cassenti, B.N., "Design Considerations for Relativistic Antimatter Rockets," *J. Br. Interplanetary Soc.*, **35** (9), 1982, pp. 396-404.
9. Morgan, D.L., "Concepts for the Design of an Antimatter Annihilation Rocket," *J. Br. Interplanetary Soc.*, **35** (9), 1982, pp. 396-404.
10. Cassenti, B.N., "Antimatter Propulsion for OTV Applications," *J. Propulsion and Power*, **1** (2), 1985, pp. 143-149.
11. Vulpetti, G., "Antimatter Propulsion for Space Exploration," *J. Br. Interplanetary Soc.*, **39** (9), 1986, pp. 391-409.
12. Forward, R.L., *Advanced Space Propulsion Study*: *Antiproton and Beamed Power Propulsion*, AFAL-TR-87-070, Hughes Research Laboratories, Malibu, CA, Oct 1987.
13. Cassenti, B.N., "Energy Transfer in Antiproton Annihilation Rockets," in *Antiproton Science and Technology*, B.W. Augenstein *et al.*, eds., World Scientific, Singapore, 1988, pp. 574-602.
14. Cassenti, B.N., "Conceptual Designs for Antiproton Propulsion Systems," AIAA Paper No. 89-2333, presented at the 25th Joint Propulsion Conference, Monterey, CA, Jul 10-12, 1989
15. Vulpetti, G., and Pecchioli, M., "Considerations About the Specific Impulse of an Antimatter-Based Thermal Engine", *J. Propulsion and Power*, **5** (5), 1989, pp. 591-595.
16. Howe, S.D., and Metzger, J.D., "Antiproton-Based Propulsion Concepts and Potential Impact on a Manned Mars Mission," *J. Propulsion and Power*, **5** (3), 1989, pp. 295-300.
17. Forward, R.L., *21st Century Space Propulsion Study*, AL-TR-90-030, Forward Unlimited, Malibu, CA, Oct 1990.
18. Tarpley, C., Lewis, M.J., and Kothari, A.P., "Safety Issues in SSTO Spacecraft Powered by Antimatter Rocket Engines," AIAA Paper No. 90-2365, 26th Joint Propulsion Conference, Orlando, FL, Jul 16-18, 1990.
19. Cassenti, B.N., "High Specific Impulse Antimatter Rockets," AIAA Paper No. 91-2548, presented at the 27th Joint Propulsion Conference, Sacramento, CA, June 24-26, 1991.
20. Forward, R.L., and Davis, J., *Mirror Matter: Pioneering Antimatter Physics*, J. Wiley & Sons, NY, NY, 1988.
21. LaPointe, M.R., *Antiproton Annihilation Propulsion Using Magnetically Confined Plasma Engines*, Dissertation, U. New Mexico, Dept. Chemical and Nuclear Engineering, Albuquerque, NM, 1989.
22. LaPointe, M.R., "Antiproton Powered Propulsion with Magnetically Confined Plasma Engines", *J. Propulsion and Power*, **7** (5), pp. 749-759.
23. Callas, J.L., *The Application of Monte Carlo Modeling to Matter-Antimatter Annihilation Propulsion Concepts*, JPL D-6830, Jet Propulsion Laboratory, Pasadena, CA, Oct 1989.

24. Huber, F., Antimatterie-Annihilationsantriebe fur Interplanetare Raumfahrtmissionen, Ph.D. Dissertation, Universitat Stuttgart, Institute fur Raumfahrtssysteme, 1994.
25. Lewis, R.A., Newton, R., Smith, G.A., Toothacker, W.S., and Kanzleiter, R.J., "An Antiproton Catalyst for Inertial Confinement Fusion Propulsion," AIAA Paper No. 90-2760, presented at the 26th Joint Propulsion Conference, Orlando, FL, Jul 16-18, 1990.
26. Lewis, R.A., Smith, G.A., Toothacker, W.S., Kanzleiter, R.J., Surratt, M.S., Higman, K.I., and Newton, R.J., "An Antiproton Driver for Inertial Confinement Fusion Propulsion", AIAA Paper No. 91-3618, presented at AIAA/NASA/OAI Conf. on Advanced SEI Technologies, Cleveland, OH, Sep 4-6, 1991.
27. Schmidt, G. R., Gerrish, H. P., Martin, J. J., Smith, G. A., and Meyer, K. J., "Antimatter Requirements and Energy Costs for Near-Term Propulsion Applications", *J. Propulsion and Power*, **16** (5), Sep-Oct 2000, pp. 923-928.
28. Chapline, G. "Antimatter Breeders," *J. Br. Interplanetary Soc.*, **35** (9), pp. 423-424, 1982.
29. Cassenti, B.N., "Concepts for the Efficient Production and Storage of Antimatter," AIAA-93-2031, presented at the 29th Joint Propulsion Conference, Monterey, CA, Jun 28-30, 1993.
30. Hora, H., "Estimates of the Efficient Production of Antihydrogen by Lasers of Very High Intensities," *OptoElectronics*, **5**, 1973, pp. 491-501.
31. Crowe, E.G., "Laser Induced Pair Production as a Matter-Antimatter Source," *J. Br. Interplanetary Soc.*, **36**, 1983, pp. 507-508.
32. Haloulakos, V., and Ayotte, A., "The Prospects for Space-Based Antimatter Production," AIAA Paper 91-1987, presented at the 27th Joint Propulsion Conference, Sacramento, CA, Jun 24-26, 1991.
33. Dirac, P.A.M., Directions in Physics, John Wiley & Sons, New York, NY, 1978, pp. 1-37, 55-70.
34. Greiner, W., Relativistic Quantum Mechanics: Wave Equations, Springer-Verlag, Inc., Berlin, 1994, pp. 261-267.
35. Mostepanenko, V. M. and N. N. Trunov, The Casimir Effect and its Applications, Oxford Science Publications, Clarendon Press, Oxford, 1997.
36. Weinberg, S., The Quantum Theory of Fields, Vol. 1, Cambridge University Press, 1995, pp. 578-593.
37. Forward, R.L., Mass Modification Experiment Definition Study, PL-TR-96-3004, Forward Unlimited, Malibu, CA, Feb 1996.
38. Boyer, T.H., "The Classical Vacuum," *Sci. Am.*, **253** (3), Aug 1985, pp. 70-78.
39. Milonni, P.W., The Quantum Vacuum: An Introduction to Quantum Electronics, Academic Press, New York, NY, 1994.
40. LaPointe, M. R., Antimatter Production at a Potential Boundary, NASA SBIR Phase I Final Report, Contract NAS8-98109, prepared for the NASA Marshall Space Flight Center, Huntsville, AL, Nov 1998.

A01-34137



AIAA 2001-3363

Gravity Modification by High-Temperature Superconductors

**C. Woods
University of Sheffield
Sheffield, UK**

**S. Cooke
Cooke Computers,
Jersey, British Isles**

**J. Helme
University of Sheffield
Sheffield, UK**

**C. Caldwell
Motorola Ltd.
Basingstoke, UK**

**37th AIAA/ASME/SAE/ASEE
Joint Propulsion Conference and Exhibit
8-11 July 2001
Salt Lake City, Utah**

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

GRAVITY MODIFICATION BY HIGH-TEMPERATURE SUPERCONDUCTORS

R.C. Woods, S.G. Cooke*, J. Helme, C.H. Caldwell†

*Department of Electronic and Electrical Engineering, University of Sheffield,
Mappin Street, Sheffield, S1 3JD, U.K.*

ABSTRACT

Claims that the weight of test masses can be changed in non-relativistic experiments have been investigated. Amongst the most prominent reports has been a paper by Podkletnov & Nieminen (*Physica C* 203 441, 1992); more recently, Rounds (*Proc. NASA Breakthrough Propulsion Phys. Workshop*, Cleveland, 297, 1998) used a simpler experimental arrangement. Both of these experiments measure the gravitational field above $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ (superconducting below $T_c \approx 93\text{K}$). Podkletnov & Nieminen specify that the superconductor must be cooled below 70K, magnetically rotated at ~ 5000 rpm, and simultaneously levitated magnetically using two separate frequency excitations; weight changes of $\sim 1\%$ were reported. Rounds specifies less stringent conditions, mainly that the YBCO is cooled to 77K whilst stationary. The Rounds experiment has been repeated virtually exactly, and the Podkletnov experiment has been investigated by reproducing some of the conditions specified. No measurable gravity modification (within $\pm 0.03\%$) was observed using the subset of the Podkletnov conditions, (but this does *not* preclude gravity modification arising from the full specified conditions). The Rounds configuration also has not shown any effects reliably ascribable to gravity modification, although there was unusual thermal behaviour.

1. INTRODUCTION

A number of reports recently claimed that the weight of test masses can be changed as a result of using various materials in various configurations and using various excitations. One of the most prominent of these has been that by Podkletnov and Nieminen¹; as far as the present authors are aware, not only is this the only such paper that has appeared in a peer-reviewed

journal, but it is also one of the few that cannot immediately be dismissed as spurious.

The experimental results reported by these authors appear to contradict conventional gravitational theory, because their claims amount to apparent modifications to the gravitational field in a laboratory-based system that – at face value – does not require analysis by General Relativity. This experiment is therefore potentially highly important scientifically because of the enormous technological implications for the design of current transportation vehicles and handling methods for bulk materials if gravity modification (and, in particular, gravitation reduction) were demonstrated to be feasible.

Objections to the Podkletnov and Nieminen experiment have been raised. Podesta and Bull² discuss the measurements¹ in terms of buoyancy effects that could be present because of the helium cooling the air surrounding the test weight. Although interesting, the air temperature required to achieve a buoyancy effect of the same magnitude as seen by Podkletnov is 150K which seems to be too low to make this a likely explanation. Unnikrishnan³ includes a description of static levitation of a large superconducting disk that finds no gravity effect, but is more interesting for its discussion of Podkletnov's claim that the gravitational effects have no dependence on the height above the spinning disk, which appears extremely strange at first sight. This, along with Podkletnov's claims to have seen increases as well as reductions in test mass weight, confirms that the effect claimed must be "Gravity modification" rather than passive "Gravitational shielding".

Pokeltnov's report¹ was therefore selected for particular attention and experimental testing in the present work, along with a subsequent and broadly similar (though smaller) effect reported by Rounds⁴

* Present address: Cooke Computers, Jersey, British Isles.

† Present address: Motorola Ltd., Basingstoke, U.K..

Copyright © 2001 by the American Institute of Aeronautics and Astronautics, Inc.. All rights reserved.

using a much simplified experimental arrangement. The experiment due to Rounds⁴ has been repeated virtually exactly, and that due to Podkletnov¹ (experimentally much more challenging) has been investigated by examining some (although not yet all) of the conditions reported as necessary for demonstrating gravity modification.

Both of these experiments have in common that they require the gravitational field to be measured above the high-temperature superconductor $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ (which is superconducting below the critical temperature $T_c \approx 93\text{K}$). In the Podkletnov arrangement, the superconductor (in the form of a multiphase circular disk having a minimum diameter 10cm) must be cooled below 70K, and magnetically rotated at high speed (e.g., 5000 rpm) and simultaneously levitated magnetically using two separate frequency excitations; weight changes (in a test mass) of the order of 1% were reported. In the Rounds arrangement, the conditions are much less stringent and mainly require the YBCO to be cooled to 77K in a stationary magnetic field.

1.1 Other claims of gravity modification

Reiss⁵ has reported work on the weight of a PTFE capsule containing pellets of various materials including 2212 BSCCO high- T_c superconductor when the capsule is immersed in liquid nitrogen. All of Reiss's measurements showed the apparent weight rising with immersion time to some final figure after about 5 minutes. This rise is a result of the thermal contraction of the holder (giving a steadily reducing buoyancy in the liquid nitrogen) and shows some scatter as well as differences for different materials contained within the holder. However, Reiss's figures indicate that the capsules containing the superconductor have the biggest apparent weight increase (although not markedly bigger). Reiss interprets this in terms of the superconductor gaining weight as the inside of the sample holder cools below his superconductor's critical temperature ($\sim 95\text{K}$). Reiss points out that the variation in the buoyancy drop found for his three non-superconducting materials (alumina, copper and PVC, in order of the lowest to highest drop) does not match variations in their specific heat. However, the variation in buoyancy drop *does* occur in the same order as the thermal expansion of these materials (i.e. alumina has a thermal expansion coefficient smaller than that of copper, which in turn is smaller than that of PVC). The room-temperature expansion coefficient of BSCCO (which shows the largest buoyancy drop) is roughly the same size as that of copper. Reiss's calculations show that the shrinkage of the PTFE capsule should lead to a

buoyancy drop three times larger than that seen in any of his experiments; this inconsistency alone throws doubt upon gravity modification having been demonstrated conclusively.

Gravity experiments by Schnurer⁶ use a 1" diameter superconducting disk attached to the top of three coils in a triangular arrangement and a metallic test mass is attached to the top of the superconductor. This assembly forms the weight on one end of a balance; the counter weight then sits on digital laboratory scales so that a weight decrease in the superconductor assembly produces an increase in the scales reading. The superconductor assembly is immersed in liquid nitrogen; once it has cooled the scales reading is noted, and power applied to the coils. The scales show an increase proportional to the sample mass. Nitrogen boiling due to the resistive heating by the coils, and possible flexing of the wires supplying the coils, may be possible explanations for these observations.

2. TESTS OF WEIGHT REDUCTION USING ROTATING SUPERCONDUCTORS

2.1 Background

Fully reproducing the Podkletnov conditions¹ is not straightforward. As far as the present authors are aware, no others have demonstrated gravity modification using similar experimental arrangements, though other partial tests⁷ have been reported.

The present approach to this experiment was to start by spinning a superconductor at high speed. The basic equipment used in initial tests is illustrated in Fig. 1. A nominal 2" diameter (the die size, not the precise final disk size; where appropriate this is given in mm) granular disk fabricated ten years previously was employed. Although this disk was able to produce significant levitation ($\sim 5\text{mm}$) of a 10g FeCo magnet, it could not levitate a 40g, 1" 0.45T NdFe magnet, nor could it sustainably levitate itself above an array of these magnets. A number of tests where the superconductor was cooled to 77K, spun at up to 5000 rpm, then slowed, and finally tested with a small magnet to ensure that the Meissner effect was still present, never produced any change in the test weight.

2.2 Present Project

In developing this investigation further, it was clear that reproducing the full Podkletnov conditions would take considerable time. Dr. Podkletnov has visited our laboratory to observe our experiments, and has made

some helpful comments based upon his own experience⁸. The strategy adopted was as follows:

- to be limited initially to 3" diameter disks,
- to develop a melt texturing process for disk fabrication (Podkletnov partially melt textured his disks to achieve much greater levitation for the same magnetic flux density),
- to develop a new weight measurement system,
- to use permanent magnets to provide levitation and/or rotating fields, and
- to produce low field, high frequency excitation using a 13.56MHz generator.

These points will be discussed in turn below.





Fig. 1: The basic gravity test equipment. The aim was to spin the nitrogen container at high speed, while keeping the superconductor below T_c .

2.2.1 Melt Texturing and Disk Fabrication

Melt texturing has been found to improve YBCO's levitation force significantly for a given field and so was developed to produce disks that could self-levitate above NdFe magnets.

When YBCO is raised above 1008°C it begins to decompose according to the reaction:



i.e., the high temperature form of the superconducting YBCO phase decomposes to a liquid that is mostly BaCuO₂ and solid green phase YBCO. This reaction is partially reversible, i.e. slow cooling back through 1008°C re-forms the 123 (superconducting) phase with inclusions of the 211 (green) phase and solidified copper/barium material. The resulting YBCO is referred to as "melt textured"; essentially the grain boundaries which limit the bulk superconductivity of a granular sample have been reduced as the grains have melted into each other. Most importantly the crystal axes in a melt textured sample tend to be aligned (usually so that the larger *c*-axis is vertical) so that the sample properties are no longer an average of the good in-plane and poor out-of-plane superconductivity exhibited by YBCO.

One step further than melt texturing is top seeded melt texturing. Seeding is carried out using small 123 phase crystals with the yttrium replaced with neodymium or samarium. This substitution produces a material with the same crystal structure (except that the larger rare earths have a tendency to be disordered because their size means that they can partially substitute for barium) but having a higher decomposition temperature. These crystals when placed on the surface of an YBCO pellet above the decomposition temperature can, on cooling, seed the growth of large single domain pseudo-crystals of 123 YBCO (they are not true single crystals because of the 211 and Cu/Ba inclusions).

Practical problems such as liquid flowing out from the pellet above 1008°C can be controlled by adding extra 211 phase to increase the solid content at high temperatures, but precise temperature control, without significant temperature gradients, is necessary. (Temperature gradients lead to different crystallisation rates across the disks, causing disk cracking.)

The aim of this work was to produce 3" disks with an even distribution of reasonably sized, randomly oriented domains across their surface. Although single YBCO domains up to 70mm across have been demonstrated⁹, owing to the level of furnace temperature control currently available to the present work the largest single domain produced was approximately 12mm × 12mm. In addition Podkletnov considered inter-grain junctions to be crucial to the gravity effect and we did not want to eliminate grain or domain junctions entirely.

It was found that attempting to distribute Nd123 crystals across the disk surface did not produce the even domain distribution necessary. Not all of the Nd seeds produced large domains and in any case

distributing small seed crystals in a furnace at 1020°C is not an easy process. A disk resulting from multiple top seeding is shown in Fig. 2; it clearly showed visibly uneven crystal distribution, giving a levitation force varying from place to place across the disk.



Fig. 2: Top seeded 3" disk

A better disk was produced not by seeding, but by dropping the post-decomposition temperature from the 982°C used for seeded growth to 975°C. This lower temperature produces self-seeding and smaller randomly shaped but evenly distributed domains resulted (Fig. 3). This 110g disk, following oxygenation between 500 and 200°C for a week, could self-levitate above an array of seven NdFe magnets at a height of approximately 15mm, and could levitate a single NdFe magnet plus a further 80g at 2mm above its surface.

As an alternative to homogeneous self-seeding a new process, heterogeneous self-seeding, was tried. This involves including some Nd211 phase in the Y123/211 mix so that the Nd/Y ratio was 2/98. When the disk was raised to 1100°C (above the decomposition temperature of Nd123) and then cooled to 982°C, Nd123 is able to solidify from the 211/liquid mix first. This is because of its higher decomposition temperature and because the neodymium's similar size to barium makes Nd much more soluble than Y in the melt – enhancing diffusion to a growing crystal

surface. The Nd123 crystals formed in this way then act as seeds distributed evenly throughout the disk for the Y123 growth. A disk grown by this method (Fig. 4) had levitation characteristics similar to the homogeneously seeded disk; it is expected that further development of this process would lead to still higher levitation forces.




Fig. 3: Homogeneously seeded 3" disk




Fig. 4: Heterogeneously self-seeded 3" disk. The crystallisation is more evenly distributed with a more random orientation than the top seeded disk

The chief problem found in this work is that of disk cracking. It was found that any indication of a crack in

the unfired disk that had been compacted in a die from the precursor powders using a fifty tonne hydraulic press would lead to catastrophic cracking during firing. Only one in four disks as-pressed was good enough to be fired (although unfired disks could usually be reground and pressed again). One in two that were fired held together, with or without some evidence of cracking, after melt texturing. The extra shrinkage due to the densification that occurs during melt texturing, plus the uneven crystallisation resulting from temperature gradients, puts a big strain on the mechanical stability of the disk.

One final innovation used during disk manufacture was the use of Saffil fabric (Alumina/Silica fibres pressed into a loose fabric/board) between disk and substrate on firing. This served two purposes; the main one was to ensure that disk could be separated from the substrate post partial melting. Secondly, since both aluminium and silicon chemically poison YBCO, incorporation of the fabric into the lower layer of the disk during melt texturing meant that there was a graduated poisoning of the superconductivity from top to bottom. In effect, the disks were bi-layered as specified by Podkletnov¹.

2.2.2 Experimental Apparatus

A beam balance was designed having the necessary resolution (0.004g or better for a 100g test mass, i.e. 0.004%) using a knife-edge pivot. Damping was essential, and was incorporated using a magnet that induced a Lorentz current in an aluminium plate opposing motion in proportion to velocity. The final design is shown in Fig. 5. The balance counter-mass was adjustable to give sensitivities that ranged from 1mm deflection for 0.02g difference, to 5mm deflection for 0.02g difference. The test mass used was 100g in all the tests undertaken as part of the present work, so that the best resolution obtainable was of the order of 0.004%, certainly sufficient for this application.

The cryostat body was made of aluminium and uses chemically-expanding foam as a thermal insulator. An aluminium heat-exchanger, for thermal coupling of the superconductor to the liquid nitrogen refrigerant, was also used.

Tests on this cryostat assembly showed that although it performed better than previous cryostats (fabricated from nylon) in maintaining superconductivity during static tests, after spinning at 5000 rpm and slowing to rest the superconductor had warmed above T_c (determined using the Meissner effect). Over 20 tests were performed using this cryostat (on the basis that

the disk would be superconducting for at least some of the time that it was being spun) with the old 2" YBCO disk and various magnet configurations. No balance deflections were seen that could be definitely ascribed to anything except vibration and so using this configuration a gravity effect larger than 0.02% can be formally ruled out.

Following this work a foil liner was added to the cryostat between the liquid nitrogen refrigerant and the foam insulator to reduce the friction on the nitrogen during acceleration. With this liner in place and using a new partially crystallised 2" disk (an early melt-textured sample that could only levitate the NdFe magnets a few millimetres) the superconductor could be consistently accelerated to a top speed of 6000 rpm, stopped, and then still levitate a magnet afterwards.

In this new configuration 32 full runs were performed using a video camera to record any balance deflections. The kinds of magnet configurations used are illustrated in Fig. 6. In some tests the superconductor was cooled through T_c in the presence of a magnetic field ("field cooling") and in others the superconductor was cooled without a field.

One variation (two NdFe magnets held over the edge of the disk, one with north up and one with north down, and the superconductor field-cooled) seemed to show balance deviations that indicated a very small weight loss during disk acceleration and deceleration. However, the vibration produced by the rotating cryostat was such that rotation alone produced balance deviations as large as any possible gravity effect. The movements were not consistent enough to be ascribed gravity effects, which in this configuration can be formally ruled out down to 0.03%.

2.3 Experimental Results from Rotating Field Tests

The aim of this work was to rotate a levitated three inch disk. However, whilst a levitated cylindrical magnet can be freely rotated about its axis in any orientation, levitated superconducting disks resist rotation in a stationary field. Therefore, the superconductor was fixed in place at the levitation height – 10 to 20mm – above a rotating array of NdFe magnets (Fig. 7). If the superconductor was not fixed in these experiments then it tended to rotate unstably with the rotating magnets. If a container was used that only just contained the superconducting disk, then sometimes the disk was seen to rise out of the position in which it had been lodged and rotate slowly when the magnets were rotating fast.




Fig. 5: The final assembly for the rotating disk tests




Fig. 6: Sample magnet configurations




Fig. 7: Rotating field apparatus

If the superconductor was fixed in place and the container was unsecured, then when the magnet rotation speed reached around 2000 rpm the superconductor lifted the whole container with liquid

nitrogen, and the container then vibrated violently. This agrees with Podkletnov's observation of his disks rising when rotated⁸, but it seems more likely that this is due to Lorentz force rather than a weight loss.

When the superconductor was firmly fixed in position the field linkage between superconductor and magnets meant that at least ~10% more power was required to start the motor turning. If the superconductor was then allowed to warm through T_c with the magnets still rotating below, the motor immediately accelerated, indicating that high rotation speeds do not diminish the torque between superconductor and magnets.

Several magnet arrangements were tried (see Fig. 8). The only noticeable result arising from the different arrangements was that greater variation in the field distribution produced greater rotational stiffness of the superconductor rotating in that field. Both the homogeneously and heterogeneously self-seeded disks discussed in Section 2.2.1 were used with no obvious differences between their behaviour. No balance deflections were seen for magnet rotation speeds of up to 10000 rpm (using three pairs of opposed magnets at this speed is equivalent to a 500Hz levitation field) and a gravitational effect can be ruled out down to 0.01% (these experiments did not require the "see-saw" balance to be moved for each run, giving better resolution).




Fig. 8: Two different rotating magnet configurations tried during the present work

A variation on the above was to use four vertical magnets placed around the superconductor (Fig. 9). These magnetically contained the superconductor which was not secured in place mechanically. When the magnets below were rotating at speed the superconductor rotated slowly from the face of one containment magnet to the next. No balance deflections were observed.

Finally, five turns of copper tube were placed around the superconductor and connected through a matching unit to a 13.56MHz r.f. supply. The matching unit was designed for capacitive loads and had very little effect in minimising the reflected power. In fact the generator

instrumentation suggested that for 200W forward power only 2W was absorbed by the load. Nevertheless, significant r.f. radiation was present in the vicinity of the superconductor, evidenced by the fact that the magnetic pick-up coil used for measuring rotation speed developed 3V peak to peak at r.f., swamping the output from the small magnets bonded to the rotating assembly. The r.f. had no other effect on the experiment and a gravity effect could still be ruled out to 0.01%.




Fig. 9: Positioning of containment magnets to prevent the superconductor rotating uncontrollably

2.4 Discussion

It can be argued that the fact that no effect has been observed in tests such as those described above is that these tests have not fulfilled the specified conditions for a gravity effect. These are:

- A disk with a diameter greater than 100mm
- A disk containing $\sim 30\%$ non-superconducting YBCO, preferably organised into two layers
- A disk capable of self-levitation, but still containing large numbers of intergrain junctions
- An a.c. levitation field with a frequency $\sim 10\text{kHz}$
- A second a.c. excitation field with a frequency of $\sim 1\text{MHz}$, for disk rotation
- Disk rotation speeds of a few thousand rpm for large ($>0.05\%$) gravity effects (the largest effects ($>2\%$) were reported¹ when the disk was being decelerated at around 3000 rpm).

The difficulties involved in meeting Podkletnov's experimental conditions¹ have already been discussed. In 1999 another team⁷ working on reproducing Podkletnov's work followed a broadly similar route to the present work, i.e. using melt textured disks

levitated with low frequencies. This group was not able to achieve stable static levitation of large disks at frequencies greater than 600Hz, and with disks that were not melt-textured stable levitation was achieved only with frequencies up to 45Hz. No gravity effects greater than 0.000001% were found.

The present work has investigated melt-texturing technology and produced high quality bi-layer disks with a final diameter of 70mm. There seems no *a priori* reason why this process may not produce larger diameter YBCO samples, particularly if annuli are formed so that larger diameters can be achieved at similar surface areas. An annular sample may also significantly reduce cracking due to shrinkage during the sintering process.

Rotating a melt-textured bi-layer 45mm disk at up to 7500 rpm through small static magnetic fields produced no effects that could be extracted from the system noise. Nor did various arrangements of high quality bi-layer 70mm superconducting disks held at levitation height above magnets rotating at up to 10000 rpm produce any gravity effect. The addition of a small 13.56MHz excitation field did not alter this negative result.

3. ROUNDS'S EXPERIMENT: STATIC SUPERCONDUCTOR

3.1 Introduction

Rounds⁴ discussed a series of experiments in which a superconductor in the presence of a permanent magnet and immersed in liquid nitrogen was slowly allowed to warm. Rounds measured the weight of the whole system, reasoning that as the superconductor passed through T_c any weight anomaly associated with the Podkletnov effect would show up as a deviation from the monotonic weight loss as the nitrogen boiled off.

Rounds found deviations from linearity in his weight against time curves for superconducting disks, and less so for copper disks. However, as with the Reiss experiment⁵, it is possible to postulate a number of thermal effects that could account for this observation.

Nevertheless, the experiment was interesting and has been repeated. Exactly the same type of superconducting disk and magnet (Edmund Scientific, part numbers CR37-446 and C52-867 respectively) were used, along with a somewhat larger nitrogen container than used by Rounds and also with a thermocouple bonded to the top of the YBCO and copper disks used.

3.2 Results

The results from a typical run with a thermocouple, an YBCO disk and a magnet separated from the YBCO by 10mm of plasticine are shown in Fig. 10, and for a copper sample in Fig. 11. Different ambient conditions along with the larger container meant that our experiments took longer (210s to the prominent inflection in the curve) than those of Rounds⁴ (48s) and the initial portion of the weight loss curve shows greater deviation from a straight line. However, the deviation of our weight loss curve from a straight line (Fig. 12) shows behaviour similar to that reported by Rounds, in particular an anomaly at around 20s elapsed time. Unfortunately our results contain too much scatter (probably a result of vibrations in the thermocouple wires) for us to ascribe this anomaly to anything more than noise.




Fig. 10: The changes in weight of the experimental apparatus as nitrogen boils off in the present work




Fig. 11: As for Fig. 10 except that the YBCO disk has been replaced with a copper disk of the same size

However, the inclusion of a thermocouple does allow identification of the point at which the superconductor sample's temperature begins to rise. Inspection of Fig.

13 shows this to be after 205s, i.e. at the same time as the inflection in the weight loss curve. This concurrence can be explained by hypothesising that the heat flux into the experimental apparatus prior to 205s produced only nitrogen refrigerant boil-off, but post-205s was also required to raise the superconductor temperature, thus resulting in a slower boil-off rate.




Fig. 12: The deviation from a straight line of the weight changes in Fig. 10

After 220s, as shown in Fig. 13, the rate of increase of temperature slowed dramatically, and the temperature effectively paused for a period of 10s at 93–94K. This is, of course, the critical temperature of the superconductor. Thereafter the rate of increase was slow until after 320s when no further liquid nitrogen refrigerant remained. This behaviour was reproducible using the YBCO superconductor sample but no such temperature pause was seen when the superconductor was replaced with a copper disc of similar dimensions.




Fig. 13: Graph of thermocouple temperature vs. time in the Rounds-configuration experiment

3.3 Discussion

With the superconductor sample (and to a lesser extent the copper sample), the inflection in the weight loss

curve corresponded to the sample starting to warm. A consequence of this (and presumably Rounds's original equipment⁴ behaved similarly) is that any anomalies in the weight loss curve before the inflection point occur while the superconductor is at a constant 77K and are therefore most unlikely to be related to modification of gravitation.

Of more interest in the results is the slowing of warming rate at T_c which is reminiscent of a first order phase transition (like, for example, the melting of ice). However, the only first order phase transition reported in high T_c superconductors is flux vortex lattice melting and this has only been observed in small, high quality, flux grown single crystals, not in granular samples like ours¹⁰. Whilst this is an interesting observation and in need of further investigation, it appears to be a purely thermal effect without any implications for gravity research.

4. CONCLUSIONS

The conclusions from the present work are that the subset of the Podkletnov conditions examined here does not produce gravity modification measurable with our equipment (a resolution of the order of $\pm 0.004\%$); it remains an open question whether gravity modification can be *repeatably* observed using the full set of Podkletnov conditions. This may, of course, only be a confirmation that not all of the correct and prescribed conditions have been achieved so far and that the present experiments could not be expected to show positive results.

The Rounds configuration also has not shown any effects unequivocally indicating gravity modification. Although some interesting effects have been produced, no effects reliably ascribable to gravity modification have been observed using either experimental arrangement.

From our repetition of the Rounds experiment, it appears likely that Rounds⁴ was seeking a gravitational anomaly at an incorrect temperature. There was, however, unusual thermal behaviour in the YBCO results for which as yet there is no adequate explanation. Unusual behaviour around T_c has been found by other authors and it is known that high T_c superconductors with very short coherence lengths can show short-lived superconducting fluctuations up to 15K either side of T_c (the width of the distribution describing these fluctuations is proportional to coherence length). It is not clear that such fluctuations or thermal effects have any relation to any gravity effect.

4.1 Simplified summary of conclusions

Two recent experiments, one in Finland and the other in the U.S.A., have claimed to demonstrate reduction of the earth's gravity in a laboratory. Such observations are potentially of great importance to the future of spaceflight, air travel, and other endeavours in which gravity plays an important part. These experiments are extremely challenging technically and so an exact repetition is difficult; nevertheless, a partial repetition of the specified experimental conditions has been undertaken. No reduction in the earth's gravity has been observed by the present researchers to date. However, this may simply be because the full specified conditions have not yet been met.

5. FURTHER WORK

The next step in this work is to attempt a more complete test of the Podkletnov conditions. Work is already in hand to increase the disk size to 10cm diameter, to construct a magnetic levitation system, to construct rotation coils and to construct a disk braking system. Investigation of the detailed theory of such effects, in our view, should properly follow a repeatable demonstration of the Podkletnov observations beyond reasonable doubt.

6. ACKNOWLEDGEMENTS

It is a pleasure to acknowledge the continued support of Dr. R.A. Evans, BAE Systems, Warton, U.K., under "Project Greenglow".

7. REFERENCES

- ¹ E. Podkletnov & R. Nieminen, Physica C **203** 441 (1992)
- ² M. de Podesta & M. Bull, Physica C **253** 199 (1995)
- ³ C.S. Unnikrishnan, Physica C **266** 133 (1996))
- ⁴ F.N. Rounds, Proc. NASA Breakthrough Propulsion Phys. Wkshp., Cleveland 297 (1998)
- ⁵ H. Reiss, Proc. 15th Europ. Conf. Thermophys. Prop., Würzburg (1999)
- ⁶ J. Schnurer, website url: <http://inetarena.com/~noetic/pls/schnurer.html> (1997)
- ⁷ R. Koczor & D. Noever, AIAA/ASME/SAE/ASEE Joint Propulsion Conference AIAA 99-2147 (1999)
- ⁸ E. Podkletnov, private communication (2000)
- ⁹ P. Gautier-Picard, X. Chaud, E. Beaugnon, A. Erraud & R. Tournier, Mat. Sci. Eng. **B53** 66 (1998)
- ¹⁰ A. Schilling, R.A. Fisher, N.E. Phillips, U. Welp, D. Dasgupta, W.K. Kwok & G.W. Crabtree, Nature **382** 791 (1996)



A01-34340

AIAA 2001-3653

**A PROPOSED EXPERIMENTAL ASSESSMENT
OF A POSSIBLE PROPELLANTLESS
PROPULSION SYSTEM**

D. Goodwin

U.S. Department of Energy
Germantown, Maryland

37th Joint Propulsion Conference
8-11 July 2001
Salt Lake City, Utah

For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

Abstract. A proof-of-principle experiment is described for a possible propellantless propulsion system (PPS) concept presented during the Space Technology and Applications International Forum of 2001. A newly available, high power, solid state switch and recent improvements in superconducting magnets, may have made a propellantless propulsion system (PPS) possible. Propulsion may occur during the non-steady state condition of the ramp-up of a very rapidly pulsed, high power magnet. Propulsion would not occur after the first 100 nanoseconds of each pulse, since the magnetic field will have reached steady state. The switch can provide 100 nanosecond ramp-ups at a rate of 0.4 megahertz, and at 9,000 volts and 30 amperes. To produce an asymmetry in the magnetic field, a plate would need to be attached to one end of the magnet. The primary components of such a PPS would be an electrical power supply, the switch, and a low temperature, superconducting magnet, capable of 2,000 amperes per square millimeter. Other applications may include propulsion with significantly reduced thermal and acoustical signatures; and a means to dampen inertia.

Introduction

A proof-of-principle experiment is described for a possible propellantless propulsion system (PPS) concept presented during the Space Technology and Applications International Forum of 2001 (Goodwin, 2001). The effect under investigation for a possible propellantless propulsion system (PPS) is the movement of high power magnets, under certain conditions, during initial power ramp-ups. A newly available, high power, solid state switch, coupled with recent improvements in superconducting magnets, may have made possible a PPS based on this effect. This new concept could be termed "a solid state PPS".

If successful, this type of PPS would meet one of the NASA's Breakthrough Propulsion Physics (BPP) technical challenge #1 for "propulsion methods that eliminate ... the need for propellant"; and may meet BPP technical challenge #2 to "dramatically reduce transit times" (Millis, 2000).

Background

High power magnets are designed to avoid movements since such movements have detrimental effects on the experiments. When movements of the magnets are encountered, the

magnets are redesigned to eliminate such movements (e.g., eliminate vibrations). By designing a system to exploit and enhance this Lenz's Law based movement, it may be possible to produce propulsion during the non-steady state conditions of the ramp-ups of very rapidly pulsed magnets.

The propulsion, if any, would only occur during the 100-nanosecond ramp-up of each pulse. After this ramp-up, the magnetic field will have reached steady state and no propulsion would occur. The non-steady, high power pulse might be producing an inertia drive effect, but without the need to mechanically move massive objects (which are limited by the strength of materials and produce only small amounts of propulsion, with supplied power; and, perhaps, no propulsion with on-board power). With sufficient power, it may be possible to accelerate the mass of the plate to 100s of g's.

The time for the magnetic field to reach steady state is related to the pulse duration (100 nanoseconds) and the size of the system (100 feet of wire). Given the size of the system (less than 1,000 feet), the magnetic field reaches steady state in less than 1 microsecond. This

This paper is declared a work of the U.S. Government and is not subject to copyright protection in the United States.

effect, if it exists, is different from magnetic "levitation" or Lenz's Law "jumping rings", both of which require the object to push off (or pull towards) other objects.

Until the recent improvements in technology, the motion produced by pulsed magnets has been an experimental interference. The motion is produced due to the Lenz's Law interaction of a pulsed magnet with an attached component producing an asymmetric magnet field. A large magnet for a particle physics moved about a centimeter every time the upgraded magnet was powered up, due to an attached component from a previous experiment. Since the component was no longer required, it was removed and the motion ceased. A small magnet for a fusion experiment moved several centimeters with the once per second pulses of 2,000 volts and 600 amperes. Since the attached asymmetric component could not be removed, the magnet was allowed to slide back into its original position. In this case, other interactions were discounted because the experiment was located in a laboratory with concrete block walls and with no rebar in the floor.

The form of Lenz's Law for a loop with fixed area is: $V = -A \frac{d\Phi}{dt}$, where V is the voltage, A is the area of the plate, and B is the magnetic field. In addition to mass penalties and costs, increasing the magnetic field (B) is limited by the physical stresses on the magnet. Superconducting magnets can produce about 10 Tesla magnetic fields for years, but a 100 Tesla field destroys the magnet after about 500 pulses and a 500 Tesla field destroys the magnet after 1 pulse. There are substantial mass penalties and costs associated with increasing the area (A). The newly available switch allows more rapidly varying the magnetic field (dt) without significant mass penalties or costs. Modifying the waveform, by adding more switches and/or modifying the switches, could increase the pulses per second and reduce the size and mass of the system. Further system optimization would be to increase the amperage (superconducting magnets have been run at kiloamps); to modify the waveform of the switch (to provide more of the pulse in the ramp-up and less in the flat-top); and to reduce the size, mass, and complexity of the interfaces between the power supply and the switch and between the switch and the magnet (e.g., reducing the need for transformers and/or using superconducting transformers).

Concept

The primary components of such a PPS would consist of the switches; low temperature, superconducting magnets; and an electrical power supply.

In FY 2000, an IR-100 award was earned by Diversified Technologies of Bedford, MA (<http://www.divtecs.com>), for the development of a high power solid-state switch. The \$6K, 7"X7"X3" switch can provide 100 nanosecond ramp-ups at a rate of 0.4 megahertz (MHz), and at 9,000 volts and 30 amperes. Allowing for the duty cycle of the ramp-ups and ramp-downs, the switch provides an integrated power of about 250 KWe. Compared to previous switches, the new switch is more compact, of lower mass, 10 times more energy efficient (only a 0.2% energy loss), and has 1000s of times higher reliability than vacuum tubes. Since the new switch has a pulse rate 10 times faster than previously available, the mass of the magnet can be reduced by a factor of 10.

Another recent improvement in technology which contributes to the possible enabling of a PPS, is a the low temperature superconductor that has recently become available which operates at 2,000 amperes per square millimeter; i.e., 10,000 times more than a non-superconductor (Barletta, 1999). High temperature superconductors carry 20 times less amperage and would probably not be able to tolerate the magnetic fields and/or the pulsed power.

To produce an asymmetry in the magnetic field for the Lenz's Law interaction, a plate would need to be attached to one end of the magnet. To prevent the energy from the pulse being carried away as induced current, the plate would need to be insulated during the ramp-ups. To prevent oscillations during the ramp-downs, the plate would be uninsulated until the next ramp-up (by using a voltage-regulated switch with a timer to connect the plate to a drain during the flattop of the pulse and then disconnect the plate from the drain before the next ramp-up).

Experiment

The proof-of-principle laboratory experiment should initially use 1 switch, 1 magnet, and externally supplied power. The experiment could use the commercial-off-the-shelf (COTS)

switch and a commercially available cable for the superconducting magnet. As with the designs for other types of experiments, the cryostats would need to be electromagnetically non-interacting.

Superconducting cable consists on up to 60 superconducting wires in a parallel conductor configuration. Allowing 500 nanoseconds during each power pulse to connect the plate to the drain, drain the plate, and reinsulate the plate, indicates about 2,000 feet of superconducting wire would be required for the propulsion magnet (to convert the electrical energy to propulsion in the remaining 2,000 nanoseconds of the power pulse). Reconfiguring 20 of the wires in the cable to be a series conductor would result in 100 feet of cable being required for the propulsion magnet. This configuration would convert each power pulse from the switch into 20 pulses of the magnet (8 MHz) and allow 20 passes through the magnet to convert each power pulse into propulsion, if the returns can be tuned to increase the ramp-up without producing a flat-top. Only about 5 KWe (2%) of the 250 KWe may be converted to propulsion since the ramp-up is only 4% of the duty cycle and the ramp-up starts at 0% power and increases to 100% rated power (i.e., averages 50% of rated power). As the ramp-up continues to build, the wires would be carrying 300 amperes (20 passes at 50% rated power). If a 2 g acceleration is produced, each pulse of the magnet would result, on average, in 0.6 milliwatt-seconds being converted into 1.2 microns of lateral motion on a supporting surface. With 32 windings and a 1-foot diameter, the magnet would have a height of about 3 feet. The cable would have a mass of about 25 kilograms. The magnet, with the cryostat, would have a mass of \geq 50 kilograms (excluding the plate).

A holdup magnet (without a plate) would be required to allow the unused energy to be returned to the switch with the correct timing for the next power pulse. Reconfiguring the 60 wires in the cable into a series conductor would result in 40 feet of cable being required to return this 2,400 nanoseconds of the power pulse. The holdup magnet would be similar in size and mass to the propulsion magnet. The wires in holdup magnet would only be carrying 30 amperes.

PPS assessment criteria (Millis, 1997) include: "must ... induce a unidirectional acceleration ... be controllable ... sustainable ... be effective

enough to propel the vehicle ... satisfy conservation of momentum (and) energy ... can be ... carried ... on the vehicle (and) consistent with empirical observations." (Millis, 1997).

Although the proposed PPS appears consistent with empirical observations and conservation of energy is not an issue, the following tests are required to confirm this and to determine if the other PPS criteria are met:

1. Confirm uni-directional motion (net external force) is produced (e.g., not limited to oscillations).
2. Confirm momentum is conserved without pushing off or pulling toward other objects (voltage regulated switch locked to drain should result in no motion; and locked to insulated mode should result in oscillations).
3. Confirm the system is effective enough to be carried on a vehicle and propel a vehicle (by measuring acceleration produced against power input). Given the potential for high energy efficiency, this type of PPS may be capable of producing near the theoretical maximum conversion of 20 kg/kWh for a 1 g thrust-per-power.
4. Confirm the system is sustainable (by long duration runs). The DOE Office of High Energy and Nuclear Physics is providing funds for the Alpha Magnetic Spectrometer (AMS) to be flown on the International Space Station. Most of the mass of this 7-ton detector is a superconducting magnet, which will operate for 3 years without maintenance. The magnet has 90 km of superconducting cable and an 800-watt cooling system.

Although the proposed PPS appears readily controllable, this could be confirmed by follow-on laboratory experiments with additional switches and magnets.

Demonstration

A sub-orbital demonstration vehicle would require an on-board electrical power supply, such as the 30 KWe "microturbine", commercially available from Capstone Turbine Corporation of Chatsworth, CA (<http://www.capstoneturbine.com>). Extensive

modifications would be required for this COTS power supply.

To match the microturbine, 6 switches (@ 5 KWe) would be required to convert the 30 KWe into propulsion. By configuring all 60 wires of the propulsion magnet into 3 sets of 20 conductors in series, the same propulsion magnet (and holdup magnet) could be used from the above experiment, if the switches can be tuned to have their ramp-ups build and not produce a flattop. With 2 switches for each set of 20 conductors in series, the propulsion magnet would be carrying 600 amperes, whereas the holdup magnet would be carrying 180 amperes (from the 6 switches). Since the propulsion magnet would still be operating at 8 MHz, a 2 g acceleration (a net of 1 g over gravity) would result in each pulse of the magnet, on average, converting 3.6 milliwatt-seconds into 0.6 microns of vertical motion (for this more massive object moving against gravity).

Given the maximum conversion of energy to propulsion noted above (10 kg per KWH for a 2 g acceleration), the mass of the demonstration vehicle would need to be kept below about 300 kilograms (including about 2 kg of fuel/oxidizer per KWH). Test stand runups could initially be performed with air, and then with an on-board oxidizer (e.g., LOX).

Applications

Possible NASA missions for this type of PPS would include the manned mission to Mars and the interstellar "precursor" probe. If the solid state PPS can provide a continuous acceleration of up to 0.2 g, the transit times for robotic missions would be reduced from years to months and for manned missions from months to weeks. As discussed below, the mass of the reactors/PPS may be only about 20% of the vehicle mass for a 0.2 g acceleration.

For launches within gravitational wells, specific power (KWe/kg) is the critical factor (which requires chemical based power supplies). For transits through the solar system and beyond, specific energy (KWH/kg) is the critical factor (which requires nuclear based power supplies). A space-based reactor with a specific mass of ≤ 18 kilograms per KWe and a conversion efficiency of $\leq 30\%$ can be available with existing technology in 5 years for units with ≤ 0.3 MWe and 10 years for units with ≥ 1 MWe

(Houts 2001, Lenard 2001, Lipinski 2001, Polk 2001, and Poston 2001).

The solid state PPS, with about a 0.1 MWe reactor, should be suitable for most robotic missions. For example, this sized system would provide more payload, power, and data return capability for such missions as the (precursor) "interstellar probe" to the heliopause at about 200 AU and on into the interstellar medium to about 400 AU (Liewer, 2000). Using the maximum conversion efficiency noted above (100 kg per 0.2 g acceleration), 90 KWe may result in a vehicle mass of up to 9 tons (what a Delta IV heavy can take to an escape orbit) and would use an optimized propulsion magnet. To match the 90 KWe, would require 18 switches (@ 5 KWe), with 6 switches connected to each of the 20 conductors in series, if the switches can be tuned to produce ramp-ups and not a flat-top. The wires in the propulsion magnet would be carrying 1,800 amperes and the wires in the holdup magnet would be carrying 540 amperes. The size and mass of the magnets would be the same as for the experiment and the demonstration. Since the propulsion magnet would still be operating at 8 MHz, a 0.2 g acceleration, would result in each pulse of the magnet, on average, converting 11 milliwatt-seconds into 0.1 microns of motion (for this much more massive object). Scaling from the AMS magnet, ≥ 10 watts of cooling would be required for each magnet.

At the maximum conversion efficiency, the solid state PPS, with a 1 MWe power supply, may be able to propel a 100-ton, manned vehicle at 0.2 g for manned missions within the solar system. Reactors can be placed in Earth orbit in a non-critical configuration. All subsequent departures and returns could be to the lunar polar orbit which faces the maximum distance away from the Earth. For launches to and returns from LEO and the polar lunar orbit, the same type of vehicle could be used, except with a chemical based electrical power supply. With the maximum conversion energy to propulsion and fuel/oxidizer consumption per KWH noted above, a vehicle to LEO would require at least 7% of the vehicle mass for fuel/oxidizer. The solid state PPS may also be able to provide substantial maneuver capability for manned and unmanned vehicles. Providing an acceleration of 0.4 g (increasing the mass of the reactors/PPS to 40% of the vehicle mass), would allow for

nuclear powered landers for Mars and most moons.

To convert 1 MWe into propulsion would require 200 switches (@ 5 KWe); i.e., 11 or 12 of the 90 KWe (optimized) propulsion magnets. For the propulsion magnets, the rep rate (8 MHz), size, mass, power conversion per pulse, motion, cooling, and amperes would be the same as described above for the robotic missions. Three holdup magnets would be required (each operating at 2,000 amperes), and with the same size, mass, and cooling as described above.

The NASA BPP interest in PPS is because the potential ISP is in the billions of seconds, as compared to \leq 480 seconds for rockets and \leq 5,000 seconds for ion drives.

The solid state PPS may be able to provide:

1. Substantially reduced thermal signatures for aerospace craft and substantially reduced acoustical signatures for military and commercial aircraft, surface ("electric") ships and submarines (for the protection of the astronauts and the ISS, the AMS magnetic is being designed to minimize stray electromagnetic fields).
2. Cost-effective boost and maneuver for trans-atmospheric vehicles and for orbital vehicles to transfer from one orbit to another (possibly including real time, varying adjustments over each ISS orbit).
3. Commercial supersonic flight.
4. Significantly reduced vibrations and acceleration/deceleration from launch/deorbit.
5. Elimination of the need for tiles.
6. Powered landings.
7. The additional safety of twin engines, either of which may be able to return the vehicle (with a lifting body fuselage for additional safety inside the atmosphere).

With this type of propulsion system, it might be possible to dampen inertia. Using < 20 femtosecond pulses, a 10 g acceleration could be divided into 0.01 angstrom increments, which may allow the biological cells time to recover

from each short burst of acceleration. This approach should lend itself to microfabrication. For example, using a 1 megahertz switch (under development by the vendor), an integrated circuit with 50 million micromagnets would be required to convert 5 KWe to propulsion without inertia. Each micromagnet would have about 5 microns of wire to convert 0.1 nanowatt-seconds per pulse into propulsion. Since the micromagnets would be in series, they would need to fail closed. Since 10 square microns of wire are required for each unit, 10 micromagnets in parallel in each unit would allow the micromagnets to fail closed. By more rapidly varying the magnetic field, such a system may also increase the conversion efficiency of electrical energy to propulsion. The cooling requirements may also be reduced.

Summary/Conclusions

By using very rapidly pulsed magnets, it may be possible to develop a propulsion system that does not rely on propellants. If successful, such a system would substantially reduce the cost of the spacecraft and would reduce the transit times for robotic missions from year to months and for manned missions from months to weeks.

Acknowledgments

The development of the high power, solid state switch and the high performance superconductors were funded, in part, by the U.S. Dept. of Energy, Office of High Energy and Nuclear Physics.

References

Barletta, W., et al. "1999 Accelerator and Fusion Research and Development Research Highlights," published by the Lawrence Berkeley National Laboratory, page 4-14.

Goodwin, D. (DOE), "A Possible Propellantless Propulsion System", in Proceedings of *Space Technology and Applications International Forum* (STAIF-2001), Feb. 13, 2001.

Houts, M. (NASA MSFC, formerly with Los Alamos National Laboratory), et al, "Options for the Development of Space Fission Propulsion Systems" in Proceedings of STAIF-2001, Feb 14, 2001.

Lenard, R. (Sandia National Laboratory), et al,
“Evolutionary Strategy for the Use of Nuclear
Electric Propulsion in Planetary Exploration”, in
the Proceedings of STAIF-2001, Feb. 13, 2001.

Liewer, P. (JPL) “Interstellar Probe: Exploring
the Interstellar Medium and the Boundaries of
the Heliosphere”,
(<http://interstellar.jpl.nasa.gov/>), November 17,
2000.

Lipinski, R. (Sandia National Laboratory), et al,
“Mass Estimates for Very Small Reactor Cores
Fueled by U-235, U-233, and Cm-245” in
Proceedings of STAIF-2001, Feb. 13, 2001.

Millis, M., (NASA GRC) “Project Plan for
Breakthrough Propulsion Physics” (NASA GRC
document TD15-PLN-015), December 4, 2000.

Millis, M., (NASA GRC) “The Challenge to
Create The Space Drive”, in AIAA Journal of
Propulsion and Power, V. 13, N. 5, pp. 577-582,
Sept-Oct 97.

Polk, J., (Jet Propulsion Laboratory) “The JPL
Advanced Propulsion Concepts Research
Program” in the Proceeding of the University of
Alabama “Advanced Space Propulsion
Workshop”, April 3, 2001.

Poston, D. (Los Alamos National Laboratory).
“The Heatpipe-Operated Mars Explorer
Reactor,” in Proceedings of STAIF-2001, Feb.
13, 2001.



A01-34341

AIAA 2001-3654
The Electromagnetic Stress-Tensor
As A Possible Space Drive Propulsion
Concept

J.F. Corum, T.D. Keech, S.A. Kapin, D.A. Gray,
P.V. Pesavento, M.S. Duncan, and J.F. Spadaro

37th AIAA/ASME/SAE/ASEE JPC
Conference and Exhibit
8-11 July 2001
Salt Palace – Salt Lake City, Utah

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

AIAA 2001-3654

THE ELECTROMAGNETIC STRESS-TENSOR AS A POSSIBLE SPACE DRIVE PROPULSION CONCEPT

by

J.F. Corum, T.D. Keech, S.A. Kapin, D.A. Gray,
P.V. Pesavento, M.S. Duncan, and J.F. Spadaro

Institute for Software Research, Inc.
1000 Technology Drive
Fairmont, WV 26554

ABSTRACT

The Heaviside force, by virtue of its nonzero value in vacuum, would appear to offer the germ of a real space-drive. Classical electrodynamic interactions may be viewed as transmitted continuously through deformations in the state of the electromagnetic field, which acts like a stressed elastic medium. Maxwell stresses not only provide a controllable net momentum flow, but also a physical mechanism acting on the fabric of space-time. In response to this unbalanced force it has been asserted that a spacecraft will move off with equal and opposite momentum. The electromagnetic interaction mechanism under consideration suggests the basis for a novel development in electrical machine technology and a possible space-drive for space transportation. The physical basis of this concept is examined in this paper, and experimental investigations are described. Supporting analyses and historical background are included.

INTRODUCTION

Perhaps the most intriguing challenge facing twenty-first century space-flight is the novel concept of a "Space-Drive".¹ John W. Campbell² and Sir Arthur C. Clarke³ are usually credited with conceiving this visionary hypothesis. Without proposing any physical mechanism (for which one might cultivate some emerging technology to exploit) they articulated what such an astonishing apparatus will do: *a space-drive is a propulsion mechanism that acts directly upon the fabric of free-space.* (Actually, the notion of a space-drive was discussed in the engineering literature almost a dozen years earlier by Joseph Slepian. See below.) Remarkably, spacecraft employing such space-drive devices would not have to convey any reaction mass to eject as propellant.

The grand challenge is, how might such a striking proposition be supported within the framework of conventional physics? And, even more thought-provoking, how might such a concept be realized in actual engineering hardware? There exist several enchanting candidates for making progress along these lines. We will address only one such concept,

and let us confess from the offset that we have no startling breakthrough to disclose. However, we do want to report on a phenomenological approach to the problem, which, to us, would seem both to encompass the concept of a space-drive, and also to hold out the promise of achievability with technology that is within reach.

ELECTROMAGNETIC FOUNDATION

In Maxwell's field theory electrodynamic interactions are transmitted continuously through deformations in the state of electromagnetic fields, which act like elastic stresses. (Note that Maxwell's equations were already relativistically covariant when he wrote them down in 1864, so what we are exploring has little, if any, relation to the old "luminiferous aether" theories.) Recall that in General Relativity (also a field theory) the interactions are transmitted as deformations in the underlying metrical structure of space-time. Ultimately, any true space drive mechanism must be expressible within such a framework. The *original* force-field derivation, actually constructed by Maxwell⁴ (and extended to vector differential form by Heaviside)^{5,6} provides the archetypical classical space-drive: an electromagnetic force acting directly upon the electrical fabric of space-time itself. The classical derivation of the Maxwell stress tensor is

Copyright © 2001 by the Institute for Software Research, Inc. Published by the American Institute of Aeronautics and Astronautics, Inc. with permission.

presented in many places and we have discussed it in several previous publications.^{7,8}

The most direct approach for developing the stress tensor is to start with the traditional coordinate independent Gibbsian, vector differential form of Maxwell's four celebrated equations in practical engineering (MKSA) units:

$$\nabla \bullet \bar{D} = \rho \quad (1)$$

$$\nabla \bullet \bar{B} = 0 \quad (2)$$

$$\nabla \times \bar{E} = - \frac{\partial \bar{B}}{\partial t} \quad (3)$$

$$\nabla \times \bar{H} = \bar{J} + \frac{\partial \bar{D}}{\partial t} \quad (4)$$

where ρ is the free electric charge density and \mathbf{J} is the sum of the electric conduction (σE) and convection (ρv) current densities. The stress tensor is quadratic in the fields and is erected by formally crossing Ampere's law into the magnetic induction vector, \mathbf{B} , and Faraday's law into the electric displacement vector, \mathbf{D} , and adding Gauss' laws (both electric and magnetic) multiplied by the field strengths \mathbf{E} and \mathbf{H} , respectively, to obtain the expression

$$\nabla \bullet \bar{T} = \rho \bar{E} + \bar{J} \times \bar{B} + \frac{\partial (\bar{D} \times \bar{B})}{\partial t} \quad (5)$$

where \bar{T} is the classical Maxwell stress tensor, and may be expressed by

$$\bar{T} = \bar{E} \bar{D} + \bar{H} \bar{B} - \frac{1}{2} (\bar{E} \bullet \bar{D} + \bar{H} \bullet \bar{B}) \bar{I} \quad (6)$$

in the common dyadic notation of Gibbs. (The product of two vectors is a dyadic quantity and \mathbf{I} is the unit dyadic.) For practical calculations one usually writes down the covariant or contravariant components of the tensor in some desired system of coordinates. For example, the components of the coordinate independent tensor quantity may be expressed on the tangent basis vectors, e_μ , or on the dual basis covectors, w^ν , of some particular system of coordinates through

$$\bar{T} = T^{\mu\nu} e_\mu \otimes e_\nu = T_{\mu\nu} w^\mu \otimes w^\nu \quad (7)$$

where $\langle e_\mu, w^\nu \rangle = \delta_\mu^\nu$ expresses the dual nature of the basis. The covariant components of the Maxwell stress tensor are given by

$$T_{\mu\nu} = h_\mu h_\nu (E_\mu D_\nu + H_\mu B_\nu - \delta_{\mu\nu} W) \quad (8)$$

and the contravariant components of the tensor are given by

$$T^{\mu\nu} = \frac{1}{h_\mu h_\nu} (E^\mu D^\nu + H^\mu B^\nu - \delta^{\mu\nu} W) \quad (9)$$

where $\mu, \nu = 1, 2, 3$ and $W = \frac{1}{2} (\mathbf{E} \bullet \mathbf{D} + \mathbf{H} \bullet \mathbf{B})$. In orthogonal curvilinear coordinates, the metrical line element is expressed as

$$ds^2 = (h_1 dx^1)^2 + (h_2 dx^2)^2 + (h_3 dx^3)^2 \quad (10)$$

with coordinate "scale factors" h_μ . [We note in passing that Maxwell's equations, and the stress tensor, \bar{T} itself, are independent of any metrical relation either Riemannian or conformal. Not only is a metric tensor not needed for writing down Maxwell's equations, but also no connection, neither Riemannian nor projective nor conformal, is required. Van Dantzig^{9,10} has shown that, in spite of this fact, the divergence of the four-dimensional stress energy-momentum tensor, which gives the Lorentz force density as well as the Poynting theorem, cannot be formulated without the aid of a metric.]

The divergence of the stress tensor, Equation (5), has the units of N/m^3 , a force density, and, therefore, implies momentum transfer. How is this to be understood? The equation actually indicates that the classical electromagnetic force expressions (the Coulomb force and the Lorentz force) are, in fact, inherent in Maxwell's equations. Since the last term on the right does not vanish in empty space it further suggests the idea of a volume force on vacuum or, at least, on the permeability, μ_0 , and permittivity, ϵ_0 , of free-space by means of vacuum displacement current interaction. Note that since \bar{T} contains only field terms, $\nabla \bullet \bar{T}$ is a force exerted by the fields.

FIELD MOMENTUM

If one volume integrates both sides of Equation (5), one obtains

$$\bar{F} = \iiint_V (\rho \bar{E} + \bar{J} \times \bar{B}) dV + \iiint_V \frac{\partial \bar{g}}{\partial t} dV \quad (11)$$

which has the units of newtons, where we have set

$$\bar{g} = \bar{D} \times \bar{B}. \quad (12)$$

In Equation (11), the first two terms on the right are the familiar forces on charges and currents.

However, a third force mechanism has emerged which is not zero in empty space:

$$\bar{F} = \iiint_V \frac{\partial \bar{g}}{\partial t} dV . \quad (13)$$

Concerning this curious term, Abraham and Becker have observed, "*The remarkable thing about this force is that its existence does not presuppose the presence of matter, the Maxwellian stresses provide a 'moving' force at points of empty space, even though there is no matter there for the force to move.*"¹¹

Recognizing that force is the time rate of change of momentum, Heaviside identified $\mathbf{g} = \mathbf{D} \times \mathbf{B}$ as the electromagnetic field *momentum density* more than one hundred years ago. With the advent of the Special Theory of Relativity, Hermann Minkowski incorporated the notion of field momentum into a four-dimensional formulation of the Maxwell stress tensor. (See Appendix I, below.) Since then, a variety of expressions have appeared for the electromagnetic field momentum, and until recently, with the measurements of Graham and Lahoz, there existed no experimental evidence to resolve the issue as to which expression is physically correct.

There is another interesting feature associated with the electromagnetic momentum. It has been observed that mechanical momentum is *not* conserved by itself alone, and that action is equal to reaction only when *both* mechanical and electromagnetic momenta are considered. It is the total (mechanical plus electromagnetic) momentum (both angular and linear) that is conserved. Action and reaction forces are always equal in magnitude and opposite in direction, and Newton's third law is satisfied - but only if electromagnetic momentum is included and correctly accounted for. O'Leary¹² gives an example of such a calculation.

Oddly, this third term appears to be *precisely what is desired* for a true "Space-Drive", and its manipulation is of engineering interest. In fact, the phenomenon seems to have first been suggested as a possible electromagnetic space-drive concept as early as 1949.

SLEPIAN AND THE SPACE DRIVE

In one of his many "Electrical Essays for Recreation", Dr. Joseph Slepian, of the Westinghouse Research Laboratories in Pittsburgh, hypothesized an electromagnetic space-ship propelled by RF forces on vacuum.¹³ (Though related, this is not radiation pressure.) His 1949

essay clearly illustrates the notion of what has since come to be known as a "Space-Drive": the craft was to be propelled, in Slepian's words, by "*a means of propulsion which does not require any material medium upon which the propelling thrust is exerted*", and, as such, the structure might be thought to be appropriate for use in space.

The mechanism that Slepian described was, essentially, a parallel plate, conformally mapped version of what later became the Graham-Lahoz cylindrical capacitor experiment.¹⁴ For his conceptual arrangement, Slepian concludes that the structure really does experience an unbalanced linear force - but for the arrangement given (\mathbf{E}, \mathbf{H} sinusoidal; stationary volume of interaction) the force is an alternating one, with zero time-average.

LINEAR FARADAY DISK MACHINE

In modern notation, what Heaviside actually wrote in 1886 was that the force to be accounted for by a state of stress in the fields is

$$\bar{f} = \rho \bar{E} + \left(\bar{J} + \frac{\partial \bar{D}}{\partial t} \right) \times \bar{B} + \bar{D} \times \frac{\partial \bar{B}}{\partial t} \quad (14)$$

The electric current (convection plus conduction) and displacement current are given by

$$\begin{aligned} \bar{J} &= \rho \bar{v} + \sigma \bar{E} \\ \bar{J}_D &= \frac{\partial \bar{D}}{\partial t} = \epsilon_0 \frac{\partial \bar{E}}{\partial t} + \frac{\partial \bar{P}}{\partial t} \end{aligned} \quad (15a,b)$$

where, in linear media

$$\bar{D} = \epsilon_0 \bar{E} + \bar{P} = \epsilon_r \epsilon_0 \bar{E} . \quad (16)$$

Equation (5) is obtained by combining the last two product terms of Equation (14) according to the product rule for differentiation.

The classic Faraday homopolar machine is a simple geometry for obtaining rotary motion by exploiting the conduction current $\mathbf{J} \times \mathbf{B}$ force. The structure may also be arranged in the form of a linear pump for conducting fluids (as patented by Einstein and Szilard in 1928), or as a linear motor by making the conductor stationary and permitting the electric current source and magnet to translate. Alternatively, Wood¹⁵ has proposed a novel neutral media accelerator by utilizing the polarization current $(\partial \bar{P} / \partial t)$ component of the displacement current in a crossed magnetic field: $\mathbf{J}_P \times \mathbf{B}$. Further, Slepian, in his space-drive system, proposed using the displacement current and the $\mathbf{J}_D \times \mathbf{B}$ force for propulsion. A machine based on these mechanisms might look something like that in Figure 1. In the limit, as $\epsilon_r \rightarrow 1$, we have Slepian's space-drive.




Figure 1. A dielectric linear motor. Crossed magnetic fields and displacement currents account for the propulsion.




Figure 2. Force density in an insulator as a function of relative permittivity for Heaviside, Abraham and Minkowski.

GRAHAM-LAHOZ

What is the correct form for the electromagnetic force density in material media? What expression is the true representation for the electromagnetic field momentum in matter? Is it possible for static electric and magnetic fields from different sources to combine to provide field momentum (angular or linear) in vacuum? In an attempt to resolve these thorny issues Walker, Graham and Lahoz have carried out a number of experimental investigations. Their results would seem to support the conclusion that the field force density does not approach zero as $\epsilon_r \rightarrow 1$. As part of the ISR BPP research project, we have subcontracted the Physics Department at West Virginia University to re-examine these basic experiments. (See Appendix III, below.)

Graham, Lahoz, and Walker have asserted that their experiments not only verify the reality of electromagnetic field angular momentum in vacuum for quasi-static fields,¹⁶ but surprisingly, they have determined that both the Minkowski and the Abraham stress tensor formulations predict results incommensurate with experimental observations.^{17, 18} (See Table I, which summarizes their results.)

HARTLEY/MANLEY-ROWE

Slepian concluded that the field force is, indeed substantial, but that it is an alternating one with zero time-average. The inherent presupposition in Slepian's work was that his field system possessed stationary boundary conditions. This assumption was first delineated by H.A. Lorentz. (See Appendix II.) The issue is, how does one spectrally translate an

oscillating force to have a DC component? We have discovered among the works of R.V.L. Hartley (of Hartley oscillator fame), published while he was at the Bell Laboratories, a remarkable phenomenon for accomplishing just this.

In the course of Hartley's work on nonlinear and time varying reactive modulators, he observed the peculiar artifact of a mechanical DC offset in the equilibrium position of the plates of an AC excited capacitor where no DC fields were applied.¹⁹ Also, there were no conventional means of rectification present to provide such DC offsets. This work, which started in 1917, led to several publications through the 1920's and 1930's and culminated in the Manley-Rowe^{20,21} papers in 1956. They conclude that time-varying and nonlinear elements both result in spectral translation. (Linear time-invariant systems never produce new frequency components.)

Setting up the Hartley criteria in Slepian's conceptual spacecraft results in nonstationary boundary conditions and a unidirectional force, as shown in Figure 3. (This should be of interest to others attempting to create a space drive out of alternating forces. Perhaps this can be adapted for use with coherent zero-point fluctuations.)

INSTANTANEOUS RADIATION PRESSURE

While the momentum of the radiation field and the phenomenon of radiation pressure emerge naturally from the Maxwell stress tensor, it should be noted that radiation pressure is somewhat different from




Figure 3. Volume integrated Heaviside force in a nonstationary medium as a function of time..

the field force exploited by Slepian in his uniform field coil-capacitor resonator. In Slepian's resonator, the orthogonal \mathbf{E} and \mathbf{H} fields are actually in time quadrature. (True radiation fields are in-phase, of course.) As a result, Slepian's fields produced a huge instantaneous force, but one with zero average value. However, as we have just discussed, the Hartley/Manley-Rowe boundary variations will provide Slepian with a spectrally translated longitudinal DC force. (See Appendix II.)

We also need to mention that most pedagogical treatments of radiation pressure present only the time-average value of the pressure of the incident radiation field at the interface. It should be noted from Appendix I that each distinct formulation of the Maxwell stress tensor predicts a different *instantaneous* radiation pressure. The instantaneous Minkowski, Abraham, and Maxwell-Heaviside radiation pressures are plotted in Figure 4 for a slightly lossy interface.




Figure 4. Instantaneous radiation pressure for the Heaviside, Minkowski, and Abraham tensors.

EXPERIMENTAL PROGRAM

Approach

In studying the linearization of the Faraday Disk machine, in both its $\mathbf{J} \times \mathbf{B}$ and $\mathbf{J}_D \times \mathbf{B}$ [as defined in Equation (15)] configurations, a way of implementing it in a laboratory environment was sought. It became obvious that one of the constraints was that there was always going to be some $\mathbf{J} \times \mathbf{B}$ component in the $\mathbf{J}_D \times \mathbf{B}$ apparatus, and that a way of separating out the two components would be required. This would be done by making force measurements with a \mathbf{J} the same length as the metallic conductor portions of the \mathbf{J}_D experiment and seeing if the force were greater when the \mathbf{J}_D length was added. After searching through the early literature on $\mathbf{J} \times \mathbf{B}$ experiments, we came across the work of Andre Marie Ampere and Hypolyte Pixii in what has become known as Ampere's Rail Motor. In modified form it is now known as the Rail Gun. The very first version of this apparatus consisted of troughs of mercury on which floated a bridge made from iron wire. When a DC current was passed through the mercury and iron wire, the bridge was found to float away from the source of DC current. Later versions replaced the mercury by metal rails and the iron bridge with two wheels connected by an axle. Either grooves in the rail prevented the wheels from rolling off, or classical railroad wheels with inner flanges were used to prevent the wheels from rolling off. What we found curious was that, invariably, in almost all reference books the experiment is never attributed to Ampere (or anyone else for that matter). Further, the magnetic field is always provided independent from the current flowing through the system, unlike in the original experiment where the \mathbf{J} current also generated the \mathbf{B} -field. And finally, the Ampere's Rail Motor is *actually a Universal Motor*: it provides a unidirectional force with both DC and AC excitation – a phenomenon which we demonstrated recently. (Strangely, we have not been able to find any references in the literature to this behavior.) Apparently, we are the first group to demonstrate this. Even the rail gun community always use pulsed DC. It was because of the capability of the Ampere-Pixii Rail Motor to operate as a universal motor that we chose to continue with that experimental configuration through the first phase of our experimental program.

Phase I

In order to use the Ampere-Pixii apparatus to demonstrate the possible unidirectional or oscillating force on a dielectric from the $\mathbf{J}_D \times \mathbf{B}$ component of the Heaviside formulation of the Maxwell stress tensor, we would have to make a break in the circuit (as was

TABLE 1. EXPERIMENTAL RESULTS

Year	Medium	Observation	Conclusion
1975	Matter	$f \neq 0$	Minkowski Wrong
1976	Matter	$f \neq \frac{(\epsilon_r \mu_r - 1)}{c^2} \frac{\partial (\mathbf{E} \times \mathbf{H})}{\partial t}$	Abraham Wrong
1980	Vacuum	$f \approx \frac{1}{c^2} \frac{\partial (\mathbf{E} \times \mathbf{H})}{\partial t}$	Correct

done in the Slepian configuration) and insert a capacitor. We decided to insert the capacitor in place of the axle between the wheels, which would make the displacement current generate the force that caused the wheels to roll away from the power generator. Discussions with several manufacturers of capacitors revealed that the largest two-plate capacitor we could obtain was only 20,000 pF, utilizing a dielectric constant of approximately 5000; we also obtained two-plate capacitor discs with dielectrics from about 1000 down to 10. In order to obtain a current of close to 50 amperes flowing through the capacitor (this was the current required with a metallic conducting axle to get the wheels to roll consistently), we could not use 60 Hz AC; based on commercially available components and our budget constraints, we chose 30 kHz as the frequency at which we would run this first experiment. The self-field at 50 amperes was only a couple of gauss for the B-field, so we also built a set of 30 kHz Helmholtz coils, which would provide a factor of between 20 and 40 improvement in the B-field (50-100 gauss, depending on the circuit Q). The phasing between the capacitor current and the Helmholtz coil current can be varied continuously over a wide range with a specially designed and constructed phasing network. Several series resonating inductors of various values to resonate the capacitor-axles at 30 kHz have been built. It is assumed that the force may be oscillating, notwithstanding the Universal motor characteristics, so a piezoelectric transducer will be used to measure the magnitude of the oscillating force. Should the Universal motor characteristic provide a unidirectional force at 30 kHz, we will be quite pleased, to say the least.

The DC power supplies are based on autotransformers followed by full wave rectifiers and capacitor filters. (The parts are refurbished

components that were once used in the coal mining industry here in West Virginia.) These power supplies will be used to drive the Class D amplifiers/power generators.

The two 30 kHz power generators are based on a full H-bridge, Class D amplifier topology. We are using Insulated Gate Bipolar Transistors (IGBT's) from IXYS Corp. for the H-bridge final drive devices, and are utilizing International Rectifier's driver chips to simplify the overall circuit. The IGBT's are rated for 1200V @ 150A, though our DC power supply is limited to 300V @ 150A, so we will not be subjecting them to their full voltage stress. The design can be easily expanded through the use of multiple devices to handle up to 900A at the same 1200V. We have also mocked up the amplifiers using IXYS BIMOS transistors, which, though they are only rated at 1200V @ 60A, are capable of operation up to several hundred kHz, as opposed to the IGBT's limit of 30-40 kHz.

The Helmholtz coils employed in our research are unique. They are a modified *spiral* coil design, instead of the classical solenoid or multi-turn loop. The reason we chose the spiral coil was that it required less wire for the same number of ampere-turns, and we got a higher field strength with the same length of wire as compared to a solenoid, as determined from our electromagnetic modeling software. The overall height of the spiral coil system is smaller than a solenoidal system. And, with some modifications to the inner turns, we were able to obtain the same field uniformity as with the classical Helmholtz coil. Laboratory measurements confirmed this. With uncompensated spirals, the magnetic field distribution was sharply peaked in the center.

The reason we want to make the tests with several different dielectrics is to establish whether, when the

force is plotted versus dielectric constant, it intersects at zero or if it intersects above zero on the force axis. (See Figure 2, above.) This will determine if the force is nonzero when the dielectric is vacuum (dielectric equal to 1). We have not been able to find any commercially available parallel plate vacuum capacitors with capacitances greater than 100 pF with plate spacing's greater than a few millimeters, which is too small at the frequency and current levels that we can obtain at the present time. Therefore, we are taking the indirect approach of measuring forces for dielectrics approaching 1 and finding out what the intercept is. As a point of note, the Graham-Lahoz experiments showed that the intercept is above zero on the force axis for angular momentum; we are expecting to be able to demonstrate the same intercept bias for linear momentum.

As a backup to this experiment, in case we run into unforeseen problems, we have a pulsed experiment, using a large permanent magnet for the B-field, and a stack of disc capacitors that will be configured as a Marx bank. Everything will be held stationary during the parallel charging of the bank; then the stops will be released and the discharge of the bank in series will generate a large current, which coupled with the 10 \times larger B-field (~1000 gauss) will result in a several orders of magnitude larger force. Because this is the equivalent of a quarter sine wave, the force will be unidirectional. The reason we are performing the rail experiment first is that the short time constant (microseconds) of the larger force will be difficult to observe because of the greatly increased mass of the second experiment (several hundred kilograms) to that of the first (a few kilograms) compared to the magnitude and duration of the force (newtons for microseconds versus millinewtons for seconds)

Phase II

The second phase of our experimental program will be to 'rectify' the force. This experiment will utilize the Hartley-Peterson-Manley-Rowe effect to perform the rectification as described earlier. Because of the requirement to operate the electronics to wobble the capacitor plate at twice the drive frequency, it will be a challenge to obtain commercially available components with these capabilities at the high power levels required to measure a detectable force (several 10's of kilowatts minimum). We also want to operate the apparatus first at the 30 kHz for which we already have several high power generators; then at between 10 and 30 MHz, which will require us to build new amplifiers. It has been our experience that the Class D topology is no longer feasible at

frequencies above about 5MHz, so we will be utilizing the Class E topology. As with Class D, this also offers very high efficiency (>90%), as well as some ruggedness to impedance mismatches, though we will be protecting the generator from high reflected power by using lumped element hybrid couplers in a back-to-back configuration, resulting in an economical circulator. The usual method of protecting RF equipment from high-reflected power is to use a circulator/isolator, but these Faraday rotation devices are not commercially available for frequencies much below 70 MHz. The dual hybrid coupler can be built or purchased for any frequency from 60 Hz to low GHz frequencies and for power levels from mW to MW. We are utilizing this lumped element coupler technique in connection with other ongoing research efforts.

CONCLUSIONS

The goal of finding an appropriate physical phenomenon to exploit and actually creating a true space-drive is a tall order. Assuming that such a device is possible, where will it be found? A variety of strategies may be employed to discover this elusive mechanism. It may be searched for along the frontiers of modern science, it may be sought in the risky hinterlands of unconventional physics, or looked for among the foundation stones of conventional knowledge. This paper reports on research along the latter path.

We have taken the position that electromagnetic forces follow directly from Maxwell's equations and, consequently, that the Heaviside force is worth experimentally studying. If the Heaviside hypothesis is correct, then, indeed, a phenomenology exists as the basis of a true space drive. While the forces might (initially) seem small, an emerging technology may then be cultivated to exploit this phenomenon. If the Heaviside hypothesis should prove to be in error, then the payoff for this research is the experimental delineation between the various classical formulations of the Maxwell stress tensor and electromagnetic momentum.

In conclusion, the appropriateness of our research effort may be measured against the five "Tests for Breakthrough Propulsion Physics." (1) Is the candidate research 'Grand'? (That is, does it possess the *potential* to respond to the 'Grand Challenge' of a true space-drive?) (2) Is the research aimed at a real 'Breakthrough'? (It must be something more than merely a refinement of conventional practice.) (3) Is the candidate research 'Credible'? (It must be scientifically sound and not what Irving Langmuir

once called ‘Pathological Science’.) (4) Is the research ‘Measurable’? (Can quantifiable advances be identified?) (5) ‘Near Term’? (Incremental progress must be inherently possible in the concept under investigation.) We believe that, when measured against each of these criteria, the answer is ‘yes’, and the present research is certainly justifiable.

SUMMARY

A space-drive advance is being sought within the domain of conventional theory. It is our hypothesis that there exists within classical electrodynamics the germ of a mechanism which may be engineered into a method of propulsion not requiring the expulsion of reaction mass. Inspired by the original electromagnetic momentum concept of Oliver Heaviside, the relativistic field formulations of Hermann Minkowski and Max Abraham, and the nonstationary networks of R.V.L. Hartley, the BPP group at ISR is attempting to exploit a modified space propulsion arrangement due to Joseph Slepian. Heaviside formulated the electromagnetic field momentum notion within the framework of Maxwell’s equations. Minkowski, Abraham, and others created four-dimensional versions of the stress tensor incorporating the field momentum concept. And, Slepian conceived the modern notion of an electrical space-drive: an electromagnetic propulsion mechanism acting directly upon space. However, Slepian only disclosed a propulsion configuration for which the space-drive delivered an alternating longitudinal force with zero average value. By incorporating the varying parameter systems of Hartley, Manley, and Rowe, one is quickly led to a Slepian-like space-drive that delivers a nonzero average force for rectilinear propulsion.

The theoretical basis has been explored, analytical models have been examined, and computer simulations have been carried out. However, the ultimate validation of the concept can only be determined experimentally in the laboratory. Such an investigation is currently being conducted.

ACKNOWLEDGEMENTS

One of the authors (JFC) would like to extend his appreciation to Professor W.R. Mellen (of the University of Lowell), and to Professors H.C. Ko, R.G. Kouyoumjian and J.D. Kraus (all of the Ohio State University) for introducing him to this enchanting topic almost forty years ago. The current research effort was supported in part by NASA Contract NAS3-000124 with the National Aeronautics and Space Administration.

APPENDIX I. OTHER FORMULATIONS

In the present paper, we have attempted to discuss the Maxwell stress tensor within the framework of conventional vector analysis. The whole formulation may be expressed more elegantly by employing either Cartan forms or tensor components in a four dimensional space-time manifold. We will follow the traditional path. There are also several ways to formulate the four dimensional stress-energy momentum tensor in material media, and each has significant differences and consequences.²² The most commonly discussed are the treatments by Minkowski,^{23,24} Abraham,²⁵ Einstein and Laub,^{26,27} and what is called the Amperian²⁸ (or Chu) formulation. While the features are different, all follow the prescription discovered by Minkowski in 1910.

Minkowski first assumed that Maxwell’s equations were valid in a frame of reference in which the material media may undergo deformation and translation. Second, he assumed that the divergence of the three-dimensional stress tensor is as obtained directly from Maxwell’s equations in Equation (5), above. Then, (and this is a critical step) by calling the sum of $\rho\mathbf{E}$ plus $\mathbf{J}\times\mathbf{B}$ the total force, i.e. by christening the Lorentz force density

$$\bar{\mathbf{f}}_L = \rho\bar{\mathbf{E}} + \bar{\mathbf{J}} \times \bar{\mathbf{B}}, \quad (\text{A.I.1})$$

as the total force, Minkowski was able to manipulate Equation (5) into the expression

$$\bar{\mathbf{f}}_L = \nabla \bullet \bar{\mathbf{T}} - \frac{\partial(\bar{\mathbf{D}} \times \bar{\mathbf{B}})}{\partial t} \quad (\text{A.I.2})$$

where the divergence is a three dimensional spatial operation. The form of Equation (A.I.2) inspires the introduction of a four-divergence operator. By combining Equation (A.I.2) with Poynting’s Theorem in the form

$$\bar{\mathbf{E}} \bullet \bar{\mathbf{J}} = -\nabla \bullet \bar{\mathbf{S}} - \frac{\partial \mathbf{W}}{\partial t} \quad (\text{A.I.3})$$

(where $\mathbf{S} = \mathbf{E} \times \mathbf{H}$), Minkowski was able to obtain a sixteen-component, four dimensional tensor, $T_{\mu\nu}$, that may be represented schematically as

$$T_{\mu\nu} = \begin{bmatrix} \bar{\mathbf{T}} & -\bar{\mathbf{g}} \\ -\bar{\mathbf{S}} & -\mathbf{W} \end{bmatrix}. \quad (\text{A.I.4})$$

The four-divergence of his tensor is a four-vector whose three spatial components present the Lorentz force density (the force on charges and currents), as it was designed to, and whose fourth component gives the power dissipated per unit volume:

$$f_\mu = \nabla^\nu T_{\mu\nu} = (\rho \bar{E} + \bar{J} \times \bar{B}; \bar{E} \cdot \bar{J}). \quad (\text{A.I.5})$$

Where the various prominent formulations differ is in their choice of an expression for the electromagnetic field momentum density, \mathbf{g} . Minkowski chose

$$\bar{g} = \bar{D} \times \bar{B} \quad (\text{A.I.6})$$

while Abraham chose

$$\bar{g} = \frac{\bar{E} \times \bar{H}}{c^2} \quad (\text{A.I.7})$$

and others make additional choices. Their selections differ only in material media, of course. None-the-less, there has been an (almost) century-long controversy over which formulation is physically correct. The various formulations predict different force expressions in matter. Minkowski has

$$\bar{f} = \nabla \bullet \bar{T} - \frac{\partial(\bar{D} \times \bar{B})}{\partial t} = \bar{f}_L. \quad (\text{A.I.8})$$

Abraham has

$$\begin{aligned} \bar{f} &= \nabla \bullet \bar{T} - \frac{1}{c^2} \frac{\partial(\bar{E} \times \bar{H})}{\partial t} \\ &= \bar{f}_L + \frac{\epsilon_r \mu_r - 1}{c^2} \frac{\partial(\bar{E} \times \bar{H})}{\partial t} \end{aligned} \quad (\text{A.I.9})$$

And, of course, Maxwell-Heaviside have

$$\bar{f} = \nabla \bullet \bar{T} = \bar{f}_L + \frac{\partial(\bar{D} \times \bar{B})}{\partial t}. \quad (\text{A.I.10})$$

Of these, only the Maxwell-Heaviside expression survives in vacuum to give a space-drive.

APPENDIX II. ZERO TIME-AVERAGE

Many years ago, an objection was raised, not to the reality of the Heaviside force, but to its accessibility. Since the force density is proportional to the time derivative of $\mathbf{E} \times \mathbf{H}$, and the time average of the derivative is zero, won't the force average to zero? This was an issue raised by Slepian in his 1949 space-drive proposal. The "zero time-average theorem"^{29,30} can be stated succinctly as follows:

The average of the derivative of any bounded quantity is zero:

$$\begin{aligned} \left\langle \frac{dg}{dt} \right\rangle &= \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \left(\frac{dg}{dt} \right) dt \\ &= \lim_{T \rightarrow \infty} \frac{g(T) - g(0)}{T} = 0 \end{aligned} \quad (\text{A.II.1})$$

Since $g(t)$ varies only within finite limits, then as the time interval T increases without

limit, the average value of dg/dt clearly goes to zero.

Panofsky and Phillips have asserted that, on a macroscopic level, the only classical quantities that have physical significance are those that have non-vanishing macroscopic space-time averages,³¹ and this would appear to dispose of any phenomenology

arising from the Heaviside term, $\left\langle \iiint_V \left(\frac{\partial \bar{g}}{\partial t} \right) dV \right\rangle$.

However, the veiled presuppositions of the theorem should be examined carefully. These are most clearly distinguished in the statement of the theorem as presented by H.A. Lorentz, himself. "*If the system is in a stationary state, the mean value of any quantity that is the derivative with respect to time of a quantity determined by the state of the system will vanish... The numerator is finite, but the denominator can be made to exceed any limit.*"³² Lorentz was, indeed, aware of the inherent assumptions nested within the theorem.

Nonstationary Systems

The question to be answered is, "What happens if the physical system is *nonstationary*?" That is, what happens if the boundary conditions or volume limits of integration (and not just the integrands) are time varying? If the boundaries can somehow be manipulated to effectively *rectify* the integrand by means of a temporally discontinuous (*nonstationary*) physical action, such as might occur with time-varying boundary conditions, then the time-average of the *volume integrated* derivative will not necessarily vanish.

When one considers vector quantities, the theorem has rather interesting implications. For a vector function \bar{F} that has been defined as the volume integral of the time-derivative of some vector density, \bar{g} , one has:

$$\left\langle \bar{F} \right\rangle = \left\langle \iiint_V \frac{\partial \bar{g}}{\partial t} dV \right\rangle \neq \iiint_V \left\langle \frac{\partial \bar{g}}{\partial t} \right\rangle dV. \quad (\text{A.II.2})$$

Consistent with Lorentz's remarks, the zero time-average theorem does *not* imply that $\langle \bar{F} \rangle = 0$ for nonstationary systems.

The Leibnitz Theorem

To illustrate this conclusion, consider the Leibnitz integral theorem from undergraduate calculus:

$$\frac{d}{dt} \left[\int_a^{b(t)} g(x, t) dx \right] = \int_a^b \frac{\partial g}{\partial t} dx + \frac{\partial b}{\partial t} g(b, t) - \frac{\partial a}{\partial t} g(a, t) \quad (\text{A.II.3})$$

The theorem shows how to differentiate an integral whose integrand, $g(x, t)$, as well as whose limits of integration, $a(t)$ and $b(t)$, are functions of the variable of differentiation.

Example

As an example of interest, suppose we have the following linear momentum density (notice that the corresponding cavity fields are reactive) and the simple *nonstationary* distributed system described by the parameters $a(t)$ and $b(t)$:

$$\begin{aligned} g(x, t) &= g_0 \cos(\omega t) \sin(\omega t) \\ a(t) &= 0 \\ b(t) &= b_0 [1 + \cos(2\omega t)] \end{aligned} \quad (\text{A.II.4})$$

where we have fixed one spatial boundary, a , and let the other boundary, b , oscillate sinusoidally about b_0 . For the given parameters, the time average Heaviside force is found from Equation (13) and Leibnitz's Theorem as

$$\begin{aligned} \langle \bar{F} \rangle &= \left\langle \iiint_V \frac{\partial \bar{g}}{\partial t} dV \right\rangle \\ &= \left\langle \frac{d}{dt} \left[\int_a^{b(t)} g(x, t) dx \right] - \frac{\partial b}{\partial t} g(b, t) \right\rangle \quad (\text{II.5}) \\ &= \frac{1}{2} \omega g_0 b_0 \end{aligned}$$

which is nonzero. The issue is that for nonstationary systems the order of volume and time integrations becomes important. As with nonergodic systems, *the expected value of the integral is not identical to the integral of the expected value*. (Obviously!) Therefore, the zero time-average theorem does *not* imply that the time-average value of the Heaviside force must be zero.

APPENDIX III. WVU EXPERIMENTS

In an attempt to independently verify the conclusions of Graham and Lahoz, a subcontract was let to the Physics Department at West Virginia University where Dr. Alan Barnes is repeating the oscillating cylinder experiments. Professor Barnes has written this Appendix III as a description of his work.

Replication Of Graham-Lahoz

The momentum density in an electromagnetic field is described by Poynting's vector, $\mathbf{E} \times \mathbf{B}/c^2$. Although easily demonstrated in the case of free radiation or

large power flows, the observation of momentum density in static electric and magnetic fields is technically difficult. Indeed only the 1980 experiment of Graham and Lahoz³³ has demonstrated a quantitative agreement between theory and experiment and their errors are about 20% of the size of the effect. To further study this fundamental aspect of momentum, we have begun a new series of measurements at West Virginia University.

These new measurements, like the Graham and Lahoz experiment, use a torsion pendulum to measure the change in angular momentum when a coaxial cylindrical capacitor is charged in a magnetic field to produce an $\mathbf{E} \times \mathbf{B}$ field between the capacitor plates. In their experiment Graham and Lahoz mounted the capacitor assembly on a torsion rod in the field of a superconducting magnet and charged the capacitor through leads running almost parallel to the magnetic fields. This gave rise to two unwanted effects: an unbalanced electromotive force on the capacitor leads during charging and a dependence of the oscillator Q-value on the alignment of the torsion pendulum with the magnetic field. The present experiment avoids these complications by putting all of the parts onboard the pendulum bob. The magnetic field is provided by permanent magnets and the battery powered unit is controlled optically from outside. Only forces arising from interactions with external fields need to be taken into account.

To give an idea of the challenge facing the experiment, the angular momentum content of the fully charged capacitor in the magnet is about 5×10^{-13} Kg m²/sec. That is enough to set the torsion bob spinning at a rate of about 1.25×10^9 radians/sec or one revolution per 160 years.

Although this sounds difficult, the experiment is not impossible. The optical system used to measure this angle is sensitive to far better than 10^{-7} radians and driving the pendulum and observing the response synchronously at resonance greatly enhances the signal to noise ratio. Our goal is to have a one percent statistical error arising from the thermal noise in the system. Early measurements of the vibrational environment of the lab where this is being undertaken indicate we should be able to achieve this thermally limited goal without going to a special low vibration lab.

The conceptual and mechanical design of the apparatus is complete and first pendulum bob is presently under construction. A second design powered by a coaxial feed down the pendulum fiber is being pursued in parallel.

REFERENCES

-
- ¹ Millis, M. "Challenge to Create the Space Drive," *Journal of Propulsion and Power (AIAA)*, **13**(5), pp. 577-682 (Sept.-Oct. 1997).
- ² Campbell, J.W., "The Space Drive Problem," *Astounding/Analog (US)*, June, 1960, pp. 83-106. Discussions: Nov., 1960, pp. 170-173; March, 1961, p. 172.
- ³ Clarke, A.C., "Beyond Gravity," published in *Profiles of the Future*, Bantum Books, New York, 1964, H2734, pp. 46-60, 235.
- ⁴ Maxwell, J.C., *Treatise on Electricity and Magnetism*, Clarendon Press, 3rd ed., 1891, Vol. I, pp. 157-166; Vol. II, pp. 276-283, 440-441.
- ⁵ Heaviside, O., *Electrical Papers*, MacMillan, 1892, Vol. I, pp. 545-548 [or The Electrician, Jan. 15, 1886, p. 187 (in Heaviside's notation, $C = J$ and $G = \partial B / \partial t$)]; Vol. II, pp. 509, 527, and 557-558.
- ⁶ Heaviside, O., "On the Electromagnetic Effects due to the Motion of Electrification Through a Dielectric," *Philosophical Magazine*, **27**, April, 1889, pp. 324-339.
- ⁷ Corum, J.F., *An Examination of the Maxwell Stress-Tensor in Minkowski Space-Time*, MSEEE Thesis, Dept. of Electrical Engineering, Ohio State Univ., Columbus, Ohio, 1967, pp. 3-6.
- ⁸ Corum, J.F., et. al., "EM Stress-Tensor Space Drive," Proc. Conf. on Breakthrough Propulsion Physics, Space Technology and Applications International Forum (STAIF-99), Albuquerque, NM, January, 1999.
- ⁹ van Dantzig, D., "The Fundamental Equations of Electromagnetism, Independent of Metrical Geometry," *Proc. Cambridge Phil. Soc.*, **30**, 421-427 (1934).
- ¹⁰ van Dantzig, D., "Electromagnetism, Independent of Metrical Geometry, Part 4 – Momentum and Energy; Waves," *Proc. Kon. Ak. Amsterdam*, 825-836 (1934).
- ¹¹ Abraham, M. and Becker, R., *The Classical Theory of Electricity and Magnetism*, Blackie and Sons, 2nd edition, 1950, p. 245.
- ¹² O'Leary, A.J., "Addition of a Third Term in the Lorentz Force Equation," *American Journal of Physics*, **14**, 63 (1946).
- ¹³ Slepian, J., "Electromagnetic Space-Ship," *Electrical Engineering*, March, 1949, pp. 145-146; "Discussion," *Electrical Engineering*, April, 1949, p. 245
- ¹⁴ Corum, J.F. et. al., "Resource Letter: The Classical Maxwell Stress Tensor," (2001). In review.
- ¹⁵ Wood, L.T., "Electromagnetic Acceleration of Neutral Molecules," *American Journal of Physics*, **42**, 1020 (1974).
- ¹⁶ Graham, G.M. and D. H. Lahoz, "Observation of Static Electromagnetic Angular Momentum in Vacuo," *Nature*, **285**, 154-155 (1980)
- ¹⁷ Walker, G.B. and D.G. Lahoz, "Experimental Observation of Abraham Force in a Dielectric," *Nature*, **253**, 339-340 (1975).
- ¹⁸ Walker, G.B. and G. Walker, "Mechanical Forces of Electromagnetic Origin," *Nature*, **263**, 401 (1976).
- ¹⁹ Hartley, R.V.L., "Oscillations in Systems with Nonlinear Reactance," *BSTJ*, **15**, 424-440 (1936).
- ²⁰ Manley, J.M. and H.E. Rowe, "Some General Properties of Nonlinear Elements – Part I: General Energy Relations," *Proc. IRE*, **44**, 904-913 (1956).
- ²¹ Rowe, H.E., "Some General Properties of Nonlinear Elements – Part II: Small Signal Theory," *Proc. IRE*, **46**, 850-860 (1958).
- ²² Pauli, W., *Theory of Relativity*, (Pergamon, 1958), pp. 109-111.
- ²³ Minkowski, H., "Die Grundgleichungen für die elektromag. Vorgänge in bewegten Körpern," *Math. Annalen*, **68**, 472-525 (1910).
- ²⁴ Minkowski, H., "Eine Ableitung der Grundgleich. für die elektromagnetischen Vorgänge in bewegten Körpern vom der Standpunkte der Elektronentheorie," *Math. Ann.*, **68**, 525-551 (1910).
- ²⁵ Abraham, M., *Rendiconti mat. Palermo*, **28**, 1 (1909); **30**, 33 (1910).
- ²⁶ Einstein, A. and J. Laub, "Über die elektromag. Grundgleichungen für bewegte Körper," *Ann. d. Physik*, **26**, 532-540 (1908).
- ²⁷ Einstein, A. and J. Laub, "Über die im elektromagnetischen Felde auf ruhende Körper ausgeübten ponderomotorischen Kräfte," *Annalen der Physik*, **26**, 541-550 (1908).
- ²⁸ Fano, R., L.J. Chu, and R. Adler, *Electromagnetic Fields, Energy, and Forces* (Wiley, 1960), Appendix A.
- ²⁹ Landau, L. and E.M. Lifshitz, *Classical Mechanics*, Pergamon Press, 3rd Edition, 1976, p. 23.
- ³⁰ Landau, L.D. and E.M. Lifshitz, *The Classical Theory of Fields*, Pergamon Press, Revised 2nd Edition, 1962, p. 96.
- ³¹ Panofsky, W.K.H. and M. Phillips, *Classical Electricity and Magnetism*, Addison-Wesley, 2nd Edition, 1962, p. 192.
- ³² Lorentz, H.A., *Problems of Modern Physics*, Ginn and Co., 1927, p. 128.
- ³³ Graham, G.M. and D. H. Lahoz, "Observation of Static Electromagnetic Angular Momentum in Vacuo," *Nature*, **285**, 154-155 (1980).



A01-34342

AIAA 2001-3656

Research on Achieving Thrust by EM Inertia Manipulation

H. Brito
Instituto Universitario Aeronautico
Cordoba, Argentina

37th AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit

8 -11 July 2001
Salt Lake City, Utah

For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

RESEARCH ON ACHIEVING THRUST BY EM INERTIA MANIPULATION

Hector H. Brito*

*Instituto Universitario Aeronautico
Córdoba, Argentina*

ABSTRACT

One of the challenges of interstellar travel being to discover a self-contained means of propulsion that requires no propellant, it has already been shown that a solution to the problem does exist, provided the system is endowed with tensor mass properties. The aim of this paper is to discuss a "propellantless" propulsion concept based on electromagnetic fields to modify the inertial properties of the system. Mass tensor properties were found to be replicated in Electromagnetism in connection with the still standing Abraham-Minkowski controversy. An experiment to settle the question has been implemented which consists of mounting the device as a seismic mass atop a mechanical suspension. Results gathered during the 1993 – 1997 period consistently point to a mechanical vibration induced by matter-EM field momentum exchange, as predicted by Minkowski's formulation, after all other sources of vibration were taken into account, or removed when possible. Tests conducted since 1999 on a redesigned experiment aiming at getting rid of most of the identified spurious effects, yield comparatively sharper and clearer evidence of sustained thrusting. However, it must yet be seen as an "anomalous" effect, since the system momentum is not conserved; thus further work remains to be done to confirm these results.

INTRODUCTION

Human interstellar exploration may not be possible without the discovery of a self-contained means of propulsion that requires no propellant.¹ This formally translates into the problem of achieving "jet-less" propulsion of spaceships that can then be seen as closed systems, i.e., without external assistance or mass/energy exchanges with the surrounding medium. It follows

from this picture that the 4-Momentum of the system should be conserved. As already shown, a formal solution to the problem does exist, provided the system is endowed with tensor mass properties. This is embodied in the Covariant Propulsion Principle (CPP) which derives from a mass tensor description of the closed system consisting of the rocket driven spaceship and its propellant mass, provided the "solidification" point is other than the system center of mass, within a relativistic covariant formulation.²

The mass tensor formulation shows that the propulsion effect is to be related to the deviatoric part of the tensor, which exhibits the particularity of producing a non-vanishing linear momentum in the spaceship comoving Lorentzian frame. This can be seen as the physical signature of a non-diagonal 4-mass tensor. The situation is reminiscent of the concept of static EM field momentum which can develop in the rest frame of a physical arrangement of electric and magnetic sources including polarizable media, as depicted in Fig. 1.

Different theoretical results are possible depending whether Planck's principle of inertia of energy is satisfied or not in the relationship between the Poynting vector (energy flow density) and the electromagnetic momentum density.³ These different results are basically Abraham's and Minkowski's forms of the EM momentum density, as the three dimensional expression of the so called "Abraham-Minkowski controversy" about the correct Energy-Momentum tensor of EM fields in polarizable media. The controversy, lasting since 1909, strikingly remains as a yet unsolved issue of Physics.^{4,5} Supporters on the theoretical aspect split about equally between the two forms, according to literature reviews, while existing experimental evidence does not allow to draw definite conclusions.

* Project Manager, Applied Research Center, Member AIAA.

Copyright © 2001 by Instituto Universitario Aeronautico. Published by the American Institute of Aeronautics and Astronautics, Inc. with permission.




Fig. 1 Different determinations of the EM momentum density in the “matter” rest frame with polarizable media.

In the present work, by using Minkowski’s formalism,⁶ a non-vanishing momentum of electromagnetic origin is shown to arise for the particular device depicted in Fig. 1. The whole system would then enjoy tensor mass properties of electromagnetic origin, when viewed as a single particle at rest in the “matter” subsystem frame. It follows that the EM field, given suitable charge and current distributions, can modify the inertial properties of the generating device, giving rise to the possibility of obtaining mechanical impulses on the device, not undergoing any exchange of mass-energy with the surrounding medium, as stated by the CPP. A propulsion concept based upon this kind of inertia manipulation mechanism is subsequently drawn; the experimental setup built to test that concept is discussed, as well as the obtained results and the applied signal processing techniques.

THRUST BY TENSOR MASS “WARPING”

The Relativistic Rocket Approach

A fully covariant reformulation of the relativistic rocket motion has been achieved and the two-particle and one-particle models of the related physical system were analyzed in Ref. 2, together with the effect of a generalized “solidification point” location, w.r.t. the center of mass (c.m.) location. When the whole system (Rocket + Ejecta) is analyzed as a single particle located at the “Rocket” system c.m. (or any “structural” point), a mass tensor is readily found as related to the whole system which reads in geometric notation

$$\mathbf{M} = \left(m_0 + m_E^* \right) \mathbf{I} + (\mathbf{p}_E \wedge \mathbf{v}) / c_0^2. \quad (1)$$

where m_0 and m_E^* represent the masses of the rocket and of the ejecta in the rocket rest frame, \mathbf{I} is the identity 4-tensor, \mathbf{p}_E the 4-momentum of the ejecta and \mathbf{v} the 4-velocity of the “solidification point”. For a closed system ($\mathbf{f} = \mathbf{0}$),

$$d(\mathbf{M} \cdot \mathbf{v}) = \mathbf{0} \quad \Rightarrow \quad \mathbf{M} \cdot d\mathbf{v} = - d\mathbf{M} \cdot \mathbf{v}, \quad (2)$$

thus, the 4-acceleration of the chosen “solidification point” is to be related to the rate of variation of the mass tensor deviator, this being written as

$$m_0 d\mathbf{v} = - (\mathbf{d}\mathbf{p}_E \wedge \mathbf{v}) \cdot \mathbf{v} / c_0^2. \quad (3)$$

Eq. (3) shows that to have zero “Rocket” 4-acceleration for any 4-velocity, $(\mathbf{d}\mathbf{p}_E \wedge \mathbf{v})$ must vanish, i.e., the deviatoric part of the mass tensor variation must vanish. It is also shown that this comes out as a necessary condition for vanishing acceleration.

The Covariant Propulsion Principle

Now, to achieve 4-acceleration, anisotropy of the mass tensor must follow at some instant and, as a result, a non-vanishing 3-momentum shows up in the spacecraft comoving Lorentz frame. An elementary algebraic analysis shows that a mass 4-ellipsoid arises as a “geometrical” representation of the mass tensor; a given variation of this tensor, related to a propulsion effect, can be viewed as a mass 4-ellipsoid warping, giving rise to the Covariant Propulsion Principle: *A spaceship undergoes a propulsion effect when the whole system mass 4-ellipsoid warps.*²

The C.P.P. furnishes an alternative wording for the Action and Reaction Principle (A.R.P.) which is known to fail in relativistic mechanics even in its weak form; it does not substitute the four-momentum conservation law, but extends its meaning to complex systems when they are represented as pointlike particles which are not collocated with the whole system center of mass. They are both derived concepts which allow, in their domains of validity, for an intuitive representation of the involved propulsion mechanisms. In the same way the A.R.P. allowed for the “mass-spring” analogy as an intuitive model, the C.P.P. allows for the essentially geometrical analogy with the *jétée* motion of the ballet dancer, during which she gets angular acceleration by retracting her extremities onto the gyration axis, i.e., by *warping its inertia tensor* in 3-D space. As for the propulsion case, a linear acceleration/deceleration is achieved by *warping the mass tensor*, which must be defined in

Minkowski's 4-D space. A pictorial representation of the propulsive effect is shown in Figs. 2 and 3, where a mass ellipsoid is assigned to a "Flatland" rocket system.




Fig. 2 "Flatland" rocket at rest in its 3-D spacetime and the corresponding mass ellipsoid.




Fig. 3 "Flatland" rocket after acceleration in its 3-D spacetime and the corresponding mass ellipsoid.

Studies about inertia as something that could be manipulated for propulsion purposes are not new, a tentative explanation has already been undertaken on the basis of the relativistic mechanics of extended bodies under electrostatic pressures.⁹ However, the C.P.P. allows for a general formulation of the problem, provided a "propulsion field" of 4-momentum \mathbf{p}_F can be ascribed to the whole system, analog to the rocket ejecta subsystem. The system can then be viewed as a single particle located at the "matter" system c.m. (or any "structural" point) so that a mass tensor is readily found as related to the whole system. The 4-acceleration of the chosen "solidification point" now becomes

$$m_0 d\mathbf{v} = -(\mathbf{d}\mathbf{p}_F \wedge \mathbf{v}) \cdot \mathbf{v} / c_0^2. \quad (4)$$

Thus, the 4-thrust on the single particle, in any arbitrary frame, is given by

$$\mathbf{F} = -\frac{d\mathbf{p}_F}{d\tau}. \quad (5)$$

Equation (5) expresses, as expected, the law of conservation of the total system energy-momentum, consistently with Equation (2). The change of the mechanical (matter) momentum exactly balances the change of the propulsion field momentum; momentum is then being exchanged within the whole closed system. The device works as a propulsion field momentum "accumulator" whereas the mechanical momentum that can be drawn from it is, by present Physics paradigms, limited to the propulsion field momentum amount.

ELECTROMAGNETIC MASS TENSOR

Since mass tensor anisotropy means non-vanishing 3-momentum in some convenient frame, question arises about the general existence conditions of momentum of electromagnetic origin in the "matter" comoving frame. A fully covariant formulation of the problem requires to consider the Energy-Momentum tensors for a closed physical system consisting of "matter" and EM fields. By applying the Law of 4-Momentum Conservation to the particles and fields contained in any four-dimensional region of space-time bounded by a closed, three-dimensional surface, the system 4-momentum is found, for closed systems, to be conserved in time. Now, if the observer's frame coincides with the frame where the "matter" is at rest when no EM field is present, the condition for anisotropic mass tensor when the EM field is ON, means that in no case the system 4-momentum aligns with the observer's 4-velocity.¹⁰ These are global consequences of mass tensor anisotropy.

When consideration is given to the locality of the energy-momentum conservation law, the following relationship can be found for the volume integral of the momentum densities¹¹

$$\int_V (\mathbf{g}^{(m)} + \mathbf{g}^{(f)}) dV = - \int_V \mathbf{x} \operatorname{div} (\mathbf{g}^{(m)} + \mathbf{g}^{(f)}) dV. \quad (6)$$

By introducing the relationship between energy flow and momentum density, and assuming that Plank's principle of inertia of energy does not necessarily hold for the EM energy flux, such as a field of group velocity c of electromagnetic waves propagation exists, Eq. (6) becomes

$$\int_V \left(g^{(m)} + g^{(f)} \right) dV = \int_V \frac{x}{c_0^2} \left[\frac{\partial w}{\partial t} - \text{grad} \left(\frac{c_0}{c} \right)^2 \cdot s^{(f)} + \left(\frac{c_0^2}{c^2} - 1 \right) \text{div } s^{(f)} \right] dV, \quad (7)$$

with w standing for the mass-energy density.

Transient Regimes

It can be seen that to obtain non-zero total momentum for specific matter-field configurations, a non-vanishing energy density variation rate is a sufficient condition. It is a sufficient condition for any matter-field configuration, provided Planck's principle of inertia holds within polarizable matter too.

Mass tensor anisotropy, as related to a special frame, can thus arise when net mass-energy fluxes take place within closed systems where Planck's principle of inertia holds everywhere, or, in other words, when the system is under anisotropic non-equilibrium conditions.

Stationary Regimes

Eq. (7) becomes, for stationary regimes and any matter-field configuration

$$\int_V g^{(f)} dV = - \int_V \left[\text{grad} \left(\frac{1}{c^2} \right) \cdot s^{(f)} + \left(\frac{1}{c^2} \right) \text{div } s^{(f)} \right] dV \quad (8)$$

The quantity between brackets being $\text{div } g^{(f)}$, a non-zero LHS is possible provided $g^{(f)}$ is not divergence-free everywhere. This can be achieved for arbitrary matter-field configurations if gradients of EM wave propagation group velocity occur in the integration region, i.e., as assumed for the derivation of Eq. (7), Planck's principle of inertia does not hold within polarizable media, in which case the energy-momentum tensor becomes unsymmetrical. This is the case for the setup shown in Fig. 4, where $\text{div } s^{(f)} = 0$ everywhere and a non-vanishing total EM momentum can only arise from the RHS first term of Eq. (8). The contributions for the volume integral come from the free surfaces of the dielectric, through which jumps of the velocity of light take place in the direction of the EM energy flux.

For this particular setup, transient regimes do not allow to produce an EM momentum contribution since energy density variation rates distribute symmetrically throughout the setup regions.




Fig. 4 Stationary regime in the 'matter rest frame with polarizable media.'

THE ABRAHAM-MINKOWSKI CONNECTION

A non-vanishing EM momentum in the stationary case implies that the whole system Energy-Momentum tensor is unsymmetrical; this is a rather uncomfortable property for a system assumed to be a closed one. Either the assumption that Planck's principle of inertia of energy does not hold everywhere is false, or the system is open and one must consider an extended closed system to which the present one belongs. The first possibility, as far as stationary regimes are considered, completely precludes a closed system to bear a tensor mass; the second possibility leads to the question about what that extended closed system is like. A hint comes from the fact that the "excess" EM momentum behaves as an "external" stress in 4-space. As a conjecture, if ZPF (Zero Point Field) were a physical reality for describing inertia,¹² that "excess" EM momentum could be explained as a form of "directed", anisotropic vacuum fluctuations of EM energy. The sought extended system would then happen to be space-time itself.

The existence conditions are consistent with the use of Minkowski's Energy-Momentum tensor for the EM field.³ By definition, the Relativistic Mechanics Laws of Conservation are satisfied; the same being true, nevertheless, for Abraham's Energy-Momentum tensor, together with other forms of the electromagnetic Energy-Momentum tensor. This is precisely the still-standing^{4,5} Abraham-Minkowski controversy about the form of that Energy-Momentum tensor, specially for low frequency or quasi-stationary fields.^{3,13-15} It reduces, basically, to the discrepancy around the mathematical expression of the EM momentum density:

$$\text{Abraham's claim: } \mathbf{g}^A = \frac{(\mathbf{E} \times \mathbf{H})}{c_0^2}, \quad (9)$$

$$\text{Minkowski's claim: } \mathbf{g}^M = (\mathbf{D} \times \mathbf{B}). \quad (10)$$

The issue is thus highly relevant to “propellantless” propulsion, since the resultant total EM 4-momentum acts exactly as the generic propulsion field \mathbf{p}_F in Eq. (5), so EM inertia manipulation becomes a theoretical possibility. Experiments to definitely settle the question were still missing besides some partialized attempts,¹⁶⁻¹⁹ leading to non conclusive enough results. A positive answer for Minkowski’s EM tensor would allow to obtain “jet-less” propulsive effects by EM fields manipulation; furthermore, it could mean an indirect demonstration of the physical reality of ZPF.

A propulsion concept based upon this kind of inertia manipulation mechanism can subsequently be drawn. It basically consists on suitably grouping the sources of electric and magnetic fields within a rigidly connecting device, as depicted in Fig. 4. By doing so, a stationary Minkowski’s EM field momentum can develop thanks to the dielectric filled region; by controlling the intensities of these fields, the inertia properties of the system as a whole, when represented by its “matter” part – the device –, are allowed to change so that a conversion of the EM field momentum into mechanical momentum of the device is expected to happen, and reciprocally; again if Minkowski is right. Nevertheless, it must be realized that this device works as an EM momentum “accumulator”. The mechanical momentum that can be drawn from, is, in accordance with present Physics paradigms, limited to the “accumulated” EM momentum amount.

ALTERNATE THRUST EXPERIMENTS

Experimental Setup Rationale

An electromagnetic momentum generator (EMMG), based on the schematics of Fig. 5, was engineered up to the “proof of concept” level and an experiment was designed aimed to verify that: a) Minkowski’s EM Energy-Momentum tensor does properly describe the electromagnetic field-matter interactions in polarizable media. b) Global EM momentum in the matter rest frame of a closed system is being generated, or, equivalently for such a system, a non scalar 4-mass tensor behaviour is being obtained. c) The experimental thruster is applying mechanical forces on the test stand without expenditure of mass, besides that equivalent to the radiant energy dissipated from the system (e.g.,

Joule heating), which cannot account for the observed effects.




Fig. 5 EM Momentum Generator schematics.

An experimental setup was accordingly built up which consists of mounting the device as a seismic mass atop a mechanical suspension. A supply of 6 A - AC (square wave) to the coils and 4 kV - DC to the capacitors allows for a total EM momentum (Minkowski’s formulation) of around 1.E-8 Ns (square wave), by using BaTiO₃ ceramic dielectrics. The alleged conversion of Minkowski’s EM momentum into mechanical momentum of the EMMG in turn generates forces acting upon this device. By means of a square wave activation of the device at a frequency close to the fundamental frequency of the seismic suspension, the supporting blade of the test fixture can be made to resonate so an amplified upper end displacement response is obtained.

Displacements in the range 10⁻⁸ - 10⁻⁷ m were to be expected. Piezoceramic strain transducers were devised to detect this range of displacements, taking into account technological as well as financial constraints. PZTs output voltages proportional to the strain level in a broad dynamic range, achieving sensitivities (seismic and acoustic threshold in controlled environments) up to 10⁻¹¹ m/m.²⁰ This is two orders of magnitude lower than the expected levels, as related to the sensing fixture shown in Fig. 6. However, the full signal includes ground and environment induced noise as observed in preliminary testing. This microseismic excitation can account for displacements comparable to those expected to be caused by the investigated effect, with a narrow band frequency response centered in the first natural frequency of the sensing fixture.

Another source of unwanted noise is the residual interaction between the coils and the Earth magnetic field, which can account for equally comparable total displacements, albeit with a deterministic distribution in the frequency domain. A third source of noise relates to the magnetic interaction between the moving and the fixed parts of the AC and DC circuits (self magnetic interaction), those belonging to the device atop the resonant blade and to the external power supply

respectively. It was also found to contribute to the displacements on practically the same foot as the formerly two mentioned sources but at twice the coil activation frequency.




Fig. 6 Micromotion sensing concept.

Other sources of noise have been considered, too, like air motion, electrostatic couplings, sound, radiometric effects, spherics, etc, which can have a degrading effect on the measurements quality, although to a lesser extent than the forementioned sources. The overall estimated effect amounts to $-60 \text{ dB} < \text{S/N} < -40 \text{ dB}$ at the transducer output and the need for further processing arises. To this aim, the analog transducer output signal is digitalized through a 12 Bit data acquisition board, making it available to PC based storage devices.

Test Implementation and Philosophy

Two series of tests were conducted during the period 1993 - 1997. Only one measurement channel was available along the first series of tests with no vibration isolation provisions. The second test series included, besides the main transducer measurement channel, a dummy seismic fixture with its transducer and measurement channel, a voltage supply measurement channel, and a vibration-free table. In both series data was acquired in 5000 samples sequences at a rate of 500 samples/sec. Power Spectral Density (PSD) using Welch's averaged periodogram method was estimated over a 2048 length frequency interval. The test philosophy was based upon comparison of results in the frequency domain, due to different excitation schemes. These were: A) Ground induced noise. B) Coils ON, capacitors OFF + (A). C) Coils ON, capacitors ON + (A). D) Coils OFF, capacitors ON + (A).

Following modeling and simulation activities, geomagnetic and self-magnetic interaction noises were expected to appear in (B) and (C) as compared to (A), while the influence of the capacitors should appear in (C) as compared to (B) if thrust by inertia manipulation is acting upon the device; no difference was expected to

arise between (D) and (A), since static electric fields alone cannot account for the vibratory behavior of the sensing fixture.

First Test Series

Results corresponding to the first test series are shown in Figs. 7-10 where, as expected, differences can be observed between the (A) and (B) spectra, mainly caused by geomagnetic noise. Differences can also be observed between the (A) and (C) spectra, but there are intriguing differences between the (B) and (C) spectra, while again, as expected, no difference appears between the (A) and (D) spectra.




Fig. 7 First test series PSD [V^2/Hz] – Case (A)




Fig. 8 First test series PSD [V^2/Hz] – Case (B)




Fig. 9 First test series PSD [V^2/Hz] – Case (C)




Fig. 10 First test series PSD [V^2/Hz] – Case (D)




Fig. 11 Second test series PSD [V^2/Hz] – Case (A)




Fig. 12 Second test series PSD [V^2/Hz] – Case (B)




Fig. 13 Second test series PSD [V^2/Hz] – Case (C)




Fig. 14 Second test series PSD [V^2/Hz] – Case (D)

SUSTAINED THRUST EXPERIMENTS

Third Test Series

Notwithstanding the improvements implemented on the second test series, uncertainties still remained which could account for the observed "positive" results, regarding especially to:

- a) Geomagnetic influence.
- b) Numerical artifacts as in Figs. 11, 14.
- c) Colored ground noise centered around the excitation frequency.
- d) Air motion (ionic wind).
- e) Power supply induced EMI.

In order to get rid of these interfering effects, the experiment was modified during 1999, according to an alternative formulation of the Abraham-Minkowski controversy, this time in terms of force densities. If dispersion is negligible and the medium is allowed to be spatially inhomogeneous and anisotropic, $D_i = \epsilon_{ik}E_k$, $B_i = \mu_{ik}H_k$. The force densities are given by^{3,8}

$$\mathbf{f}^M = \rho\mathbf{E} + \mathbf{j} \times \mathbf{B} - \frac{1}{2}E_iE_k\nabla\epsilon_{ik} - \frac{1}{2}H_iH_k\nabla\mu_{ik}, \quad (11)$$

$$\begin{aligned} \mathbf{f}^A = \mathbf{f}^M - \nabla \cdot (\mathbf{D}\mathbf{E} - \mathbf{E}\mathbf{D}) - \nabla \cdot (\mathbf{B}\mathbf{H} - \mathbf{H}\mathbf{B}) \\ + \frac{\partial}{\partial t} \left(\mathbf{D} \times \mathbf{B} - \frac{1}{c_0^2} \mathbf{E} \times \mathbf{H} \right). \end{aligned} \quad (12)$$

With the isotropic medium approximation Eqs. (11) and (12), reduce to

$$\mathbf{f}^M = \rho\mathbf{E} + \mathbf{j} \times \mathbf{B} - \frac{1}{2}E^2\nabla\epsilon - \frac{1}{2}H^2\nabla\mu, \quad (13)$$

$$\mathbf{f}^A = \mathbf{f}^M + \frac{\epsilon_r\mu_r - 1}{c_0^2} \frac{\partial}{\partial t} (\mathbf{E} \times \mathbf{H}). \quad (14)$$

These force densities clearly differ inside matter for generic fields; they are however identical for static fields, irrespective of the medium. If harmonic fields are considered, the force densities instantaneous values differ but their averaged values become identical and therefore useless for discriminating between the two formulations. This is the reason Walker&Walker's claim,¹⁸ favoring Abraham's one, is essentially wrong and the experiment remains inconclusive.

The EM inertia manipulation (EMIM) experiment was modified in such a way that both \mathbf{D} and \mathbf{B} fields were subjected to harmonic evolution. The averaged effect of the total EM force was sought after, so Eq. (13) must be used for its theoretical estimation as applied to the schematics of Fig. 5. This estimation can proceed

straightforwardly by neglecting capacitor edge effects, by considering the coils as a conducting "box" bearing a negligible voltage w.r.t. to the capacitor plates and by assuming that the polarization current within the dielectric contributes to the second term of Eq. (13).

As a result, electric self-interaction, represented by the first term of Eq. (13) and magnetic self-interaction represented by the second term of the same equation, simply cancel out. Since non-magnetic matter is involved, contribution to the total EM force acting upon the device comes from the third term. This contribution is non-zero through the boundaries of the dielectric filled volume. Since induced electric fields of magnitude E_i appear on these boundaries yielding an unbalancing effect on the electric field of magnitude E set by the capacitor, the integrated effect does not cancel out in the $\mathbf{E} \times \mathbf{H}$ direction and can be written as ($\epsilon_r \gg 1$)

$$F = 2\epsilon_r\epsilon_0EE_iA_L, \quad (15)$$

where A_L represents the lateral open surfaces of the dielectric. Application of the formula for parallel plates capacitor of width d , Lenz's Law and Ampere's Law for an infinite length solenoid of n turns, yields the following expression for the EM instantaneous thrust, as a function of the harmonic voltage $V \sin\omega t$ on the capacitor and the harmonic current $I \sin(\omega t + \phi)$

$$F = -\frac{\epsilon_r\omega nIVd}{c_0^2} \left(\frac{1}{2} \sin 2\omega t \cos \phi - \sin^2 \omega t \sin \phi \right). \quad (16)$$

The averaged value then results

$$\langle F \rangle = \frac{\epsilon_r\omega nIVd}{2c_0^2} \sin \phi, \quad (17)$$

with maxima at $\phi = \pm \pi/2$. The results obtained with Walker&Walker's experiment are consistent with this formulation and can, as the authors readily did, be interpreted in terms of the polarization current contribution to the Lorentz force.

However, Eq. (17) must be seen as a conflicting result if total momentum must be conserved, as stated previously. In fact, the standard treatment of the problem requires the polarization current to be excluded from the magnetic contribution to the Lorentz force, the self-magnetic interaction does not longer cancel out and Eq. (16) must be corrected as follows

$$F = 2 \varepsilon_r \varepsilon_0 E E_i A_L - P B V_C , \quad (18)$$

where V_C stands for the capacitor volume filled with the dielectric. By adopting the same assumptions as for the derivation of Eq. (16), the second term of Eq. (18) becomes

$$F_{(2)} = -\frac{\varepsilon_r \omega n I V d}{c_0^2} \left(\frac{1}{2} \sin 2\omega t \cos \varphi + \cos^2 \omega t \sin \varphi \right) \quad (19)$$

so the total "standard" force is

$$F^S = -\frac{\varepsilon_r \omega n I V d}{c_0^2} (\sin 2\omega t \cos \varphi + \cos 2\omega t \sin \varphi), \quad (20)$$

and the averaged value goes to zero.

Therefore, the modified EMIM experiment should allow to discriminate between the "standard" and the presently proposed formulation of the averaged EM force. To take advantage of the sensing device characteristics, the voltage supply is reversed at a frequency different from the supply frequency, so the seismic setup is put into vibratory motion if the "proposed" formulation is correct. By detecting this force, the interfering effect (a) becomes averaged out; direct detection also permits to overcome the interfering effect (b) since no Wiener filtering is necessary; if the voltage reversing frequency is different from the setup natural frequencies, the interfering effect (c) becomes less significant and air motion (d) being related to the power supply frequency averages out too. Uncertainties are expected to remain regarding interfering effect (e).

Experiments were performed according to the test philosophy of the preceding test series. A reversing frequency of 30 Hz was chosen. Propulsive effects show up only when the Caps ON – Coils ON condition holds. Furthermore, maxima are obtained for a voltage-current phase shift of nearly 90 degrees, as predicted by the proposed formulation. A comparison of case (C) results with the corresponding simulation results (dotted line) is shown in Fig. 15, where a close agreement is found for the response to the alleged EMIM averaged force at 30 Hz reversing frequency. A comparison between cases (A) and (C) is shown in Fig. 16. Phase shift dependence is shown in Fig. 17, where experimental PSD peak values at 27 Hz are plotted against the voltage-current shift angle.




Fig. 15 Third test series PSD [dB] – Case (C) experimental and simulation results.




Fig. 16 Third test series PSD [dB] – Cases (A)&(C).




Fig. 17 Third test series average PSD peak values [V^2/Hz] @ 27 Hz, with 90% confidence intervals.

Spreading of the PSD peak values for a given angle was initially thought being due to contributions of magnetic interactions between the capacitor and the coil circuits during non-synchronized voltage reversings and/or between the capacitor circuit and the geomagnetic field. However, order of magnitude estimations show those interactions to be unable to account for the observed deviations. Their source remains unknown, so far, except for fluctuations of ground noise components at the reversing frequency. Nevertheless, a slightly shifted squared sine trend can clearly be seen to emerge from the plotted data.

LAYMAN'S SUMMARY

Either to go to the stars or, more pragmatically, to substantially cut down space transportation costs, new propulsion mechanisms must be found which get rid of propellants and/or conventional external assistance, i.e., the mythical "space drive" must still be invented.

Recent theoretical works show that jetless–sailless–beamless–tetherless propulsion can be achieved by manipulating the spaceship inertia in a way analogous to a dancer who increases her angular velocity by manipulating her body moment of inertia. To do that, an "extended" spaceship including the fields it eventually generates must be considered; a thrust then appears on the "material" spaceship by means of momentum exchange with its "field" complement.

This research has been conducted with the goal of checking whether the electromagnetic field is suitable for the above-mentioned purpose, since it can be shown that a theoretical formalism (Minkowski's) gives credit to that possibility. Experiments were designed and performed which, in an exploratory phase, gave indirect evidence of Minkowski's approach being valid. In a second and ongoing phase, following slight changes of the experiment implementation, direct and clearer evidence of sustained thrust (as it should be for operational propulsion) is being found.

However, the sustained thrust must, in principle, be seen as an "anomalous" effect since the Law of Momentum Conservation seems to be violated. This casts doubts about the validity and/or interpretation of the results and further experimental work, including in-orbit testing, is required to arrive to safer conclusions. If the effect still remains, propellantless propulsion would have been achieved but additional theoretical work will be needed for the full understanding of the underlying physical principles.

CONCLUSIONS

Theoretical as well as experimental research about a propellantless propulsion concept has been presented. On the theoretical side, thrusting without reaction mass or beamed power, seems to be possible by EM inertia manipulation, provided existence conditions hold, as those consistent with Minkowski's Energy-Momentum tensor for the EM field. However, the validity of this formalism is presently arguable under the still-standing Abraham-Minkowski controversy, the main argument being the unsymmetrical nature of that tensor. Experimental elucidation of the controversy was sought after and instrumented around a so called EMIM force-producing device.

Tests performed during the period 1993 - 1997 produced results which after intensive data processing, consistently pointed to a mechanical vibration induced by mass/inertia tensor warping of the device, or matter-electromagnetic field momentum exchange, as predicted by Minkowski's formalism. However, no direct detection of the sought effect has been obtained up to now; the experimental setup overall detectability needs further improvements, jeopardized by several potentially interfering effects.

A third series of tests conducted during 1999 and 2000 on a redesigned experiment to get rid of most of the identified spurious effects, yield comparatively sharper and clearer evidence of EMIM induced effects, albeit in contradiction with null results predicted by the "standard" formulation of global EM forces. Moreover, the results seem to indicate that the system momentum is not conserved, casting severe doubts about their validity; however, a conclusive demonstration that this is absolutely forbidden by the laws of Physics has not yet been provided and future breakthroughs in understanding the nature of inertia and motion may lead to more optimistics prospects. On the other hand, the alternative formulation presented in this paper correctly predicts peer-reviewed experimental results.

Further work still remain to be done to confirm these results, especially from the viewpoint of power supply induced EMI on the measurement channels, sharing the same spectral signature with the pursued effect, which could be overcome by means of Interferometer/Doppler vibrometry techniques. Another noise source was identified as being related to self-magnetic interactions in wiring and windings of power-supply components. Although efforts have already been made to filter out its influence from the experimental data, safer results will be obtained by a substantial increase of power. This will allow to use a reversing voltage frequency close to the fundamental frequency of the sensing device, in

order for the alleged observable effect to show up well over the ground induced noise. In-orbit testing will indeed provide final answers, by simultaneously getting rid of all mentioned interferences.

ACKNOWLEDGMENTS

The author wishes to thank the U.S. Air Force Research Laboratory for supporting the presentation of this work under its Windows-on-Science Program, and to NASA BPP Project for procuring that help. Support from the Argentine National Agency for the Advancement of Science and Technology under Grant PICT-99-10-07107, and "Instituto Universitario Aeronautico" of Cordoba, Argentina, are also gratefully acknowledged.

REFERENCES

- ¹ Millis, M. G., "Challenge to Create the Space Drive," *Journal of Propulsion and Power* **5**, 577-582 (1997).
- ² Brito, H. H., "A Propulsion-Mass Tensor Coupling in Relativistic Rockets Motion," in *Proceedings of the Space Technology Applications International Forum (STAIF-98)*, edited by Institute for Space and Nuclear Power Studies, Albuquerque, NM, Part Three, 1998, pp. 1509-1515.
- ³ Brevik, I., "Definition of some Energy-Momentum Tensors," *Experiments in Phenomenological Electrodynamics and the Electromagnetic Energy-Momentum Tensor*, PHYSICS REPORT (Review Section of Physics Letters) **52**, No. 3, 1979, p. 139.
- ⁴ Antoci, S., Mihich, L., "A forgotten argument by Gordon uniquely selects Abraham's tensor as the energy-momentum tensor for the electromagnetic field in homogeneous, isotropic matter," *Nuovo Cimento B* **112 B**, 991-1007 (1997).
- ⁵ Johnson, F. S., Cragin, B. L., Hodges, R. R., "Electromagnetic momentum density and the Poynting vector in static fields," *American Jnl. of Physics* **62**, 33-41 (1994).
- ⁶ Jackson, J. D., "Time-Varying Fields, Maxwell's Equations, Conservation Laws," in *Classical Electrodynamics*, 2nd Ed., edited by John Wiley & Sons, Inc., New York, Publisher, 1962, pp. 169-202.
- ⁷ Portis, A. M., "Fuentes del campo electromagnético III - Cantidad de movimiento del campo," in *Campos Electromagnéticos*, Sp. ed., Vol. 2, edited by Ed. Reverté, Barcelona, Publisher, 1985, pp. 469-472.
- ⁸ Eu, B. C., "Statistical foundation of the Minkowski tensor for ponderable media," *Phys. Review A* **33**, 4121-4131 (1986).
- ⁹ Marchal, R., "Sur l'inertie électromagnétique," *Comptes Rendus* **268 A**, 299-301 (1969).
- ¹⁰ Brito, H. H., "Propellantless Propulsion by Electromagnetic Inertia Manipulation: Theory and Experiment," *AIP Conference Proceedings* **458**, American Institute of Physics, New York, 1999, pp. 994-1004.
- ¹¹ Furry, W. H., "Examples of Momentum Distributions in the Electromagnetic Field and in Matter," *American Jnl. of Physics* **37**, 621-636 (1969).
- ¹² Haisch, B., Rueda, A., Puthoff, H. E., "Inertia as a zero-point field Lorentz force," *Physical Review A* **49**, 678-694 (1994).
- ¹³ Lai, H. M., "Electromagnetic momentum in static fields and the Abraham-Minkowski controversy," *American Jnl. of Physics* **48**, 658-659 (1980).
- ¹⁴ Brevik, I., "Comment on 'Electromagnetic Momentum in Static Fields and the Abraham-Minkowski Controversy,'" *Physics Letters* **88 A**, 335-338 (1982).
- ¹⁵ Lai, H. M., "Reply to 'Comment on 'Electromagnetic Momentum in Static Fields and the Abraham-Minkowski Controversy,'" *Physics Letters* **100 A**, 177 (1984).
- ¹⁶ James, R. P., "Force on Permeable Matter in Time-Varying Fields," *Ph.D Thesis*, Dept. of Electrical Engineering, Stanford University, 1968.
- ¹⁷ Walker, G. B., Lahoz, D. G., Walker, G., "Measurement of the Abraham Force in Barium Titanate Specimen," *Canadian Jnl. of Physics* **53**, 2577-2586 (1975).
- ¹⁸ Walker, G. B., Walker, G., "Mechanical forces in a dielectric due to electromagnetic fields," *Canadian Jnl. of Physic* **55**, 2121-2127 (1977).
- ¹⁹ Lahoz, D. G., Graham, G. M., "Observation of Electromagnetic Angular Momentum within Magnetite," *Physical Review Letters* **42**, 137-1140 (1979).
- ²⁰ Forward, R. L., "Picostrain measurements with piezoelectric transducers," *Jnl. of Applied Physics* **51**, 5601-5603 (1980).
- ²¹ Widrow, B. et al, "Adaptive Noise Cancelling:Principles and Applications," *Proceedings of the IEEE*, Vol. 63, No. 12, Dec. 1975, pp. 1692-1716.
- ²² Proakis, J. G., Rader, C. M., Ling, F., Nikias, C. L., "Adaptive Filters," in *Advanced Signal Processing*, Macmillian Publishing Company, New York, 1992, pp. 315-399.
- ²³ "DSP Design and Simulation Using the SIMULINK DSP Blockset," *SIMULINK Technical Computing Brief*, The Mathworks Inc., Natick MA, 1996.

A01-34343



AIAA-2001-3657

**Experimental Results of Schlicher's
Thrusting Antenna**

Gustave C. Fralick

NASA Glenn Research Center
Cleveland, OH 44135

Janis M. Niedra
6052 Oakwood Circle
North Ridgeville, OH 44039

**37th AIAA/ASME/SAE/ASEE Joint Propulsion
Conference and Exhibit**
8-11 July 2001 / Salt Lake City, Utah

For permission to copy or republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA 20191-4344

EXPERIMENTAL RESULTS OF SCHLICHER'S THRUSTING ANTENNA*

Gustave C. Fralick
NASA Glenn Research Center
Cleveland, Ohio 44135

Janis M. Niedra
6052 Oakwood Circle
N. Ridgeville, OH 44039

Abstract

Experiments were conducted to test the claims by Rex L. Schlicher et.al. (Patent 5,142,861) that a certain antenna geometry produces thrust greatly exceeding radiation reaction, when driven by repetitive, fast rise and relatively slower decay current pulses. In order to test this hypothesis, the antenna was suspended by strings as a 3 m pendulum. Current pulses were fed to the antenna along the suspension path by a very flexible coaxial line constructed from loudspeaker cable and copper braid sheath. When driving the antenna via this cabling, our pulser was capable of sustaining 1200 A pulses at a rate of 30 per second up to a minute. In this way, bursts of pulses could be delivered in synch with the pendulum period in order to build up any motion. However, when using a laser beam passing through a lens attached to the antenna to amplify linear displacement by a factor of at least 25, no correlated motion of the beam spot could be detected on a distant wall. We conclude, in agreement with the momentum theorem of classical electromagnetic theory, that any thrust produced is far below practically useful levels. Hence within classical electrodynamics, there is little hope of detecting any low level motion that cannot be explained by interactions with surrounding structural steel and the Earth's magnetic field.

Introduction

This paper describes the attempt to measure the thrust of the "Non-Linear Electromagnetic Propulsion System" (NEMPS), patent no. 5,142,861, (Ref. 1) and described at the 31st Joint Propulsion Conference (Refs. 2-5). The device supposedly generates a mechanical

force when electric current is sent through it. Such a device, if it produced more than the minuscule thrust of a photon rocket, would be a boon for deep space, perhaps, even interstellar, missions. No longer would it be necessary to carry half the mass of the mission in fuel. Just connect your electric thruster to a power source, such as a radioisotope heat source, and off you are to the nearest star!

After an attempt to acquire a thruster from the developer, we decided to build our own, using the patent drawings and other published material (Refs. 1-5) as construction guidelines. We also built our own pulse power supply, again relying on the sketchy descriptions provided in the references. Since we received no direct help from the developer, the device is not a perfect reproduction, but likely works as well. The reasons for this statement are given at the end of this paper.

Principle of Operation of the NEMPS thruster

According to the device description given in Ref. 2, "the rigid three dimensional geometric asymmetries in the loop antenna structure cause a magnetic field density gradient internal to the antenna structure along a single axis. This magnetic field density gradient then causes an imbalance in the internal magnetic-mechanical forces that normally result from the interactions of the loop antenna's internal magnetic field with the current in the conductors of the loop antenna's structure, as described by the Lorentz force law." The thruster is powered by a pulsed power supply "designed to provide the proper waveform to the antenna structure at an impedance matching the load impedance of the antenna. The rise time and shape of the input current waveform is crucial to maximizing the production of reaction thrust." However, neither the pulse parameters nor the reason for their criticality are given.

*Copyright © 2001 The American Institute of Aeronautics, Inc. No copyright is asserted in the United States under Title 17, U.S. Code. The U.S. Government has a royalty-free license to exercise all rights under the copyright claimed herein for Government Purposes. All other rights are reserved by the copyright owner.

Description of Experiment

Conducting the experimental test of the claims of the NEMPS thruster turned out to present a different set of problems than was expected. Initially we thought that we would see evidence of some force being produced, and then having to eliminate any spurious or non-relevant forces, such as interaction with the earth's magnetic field, or with the structural steel of the building, or perhaps even the electrostatic precipitator effect. But we really observed no evidence of any force whatsoever. Thus we were reduced to setting an upper limit for the force produced by the thruster.

The first step in testing the NEMPS thruster was to build it. As can be seen from the drawing, (Fig. 1), and our implementation of it (Fig. 2) the device is a coaxial transmission line, the outer conductor of which blooms into a petal like arrangement. Four bundles of two conductors each lead from the petals back to the center conductor to complete the circuit. In our embodiment of the NEMPS thruster, the center conductor is a piece of copper rod, $1/4"$ in diameter and $30"$ long. The portion of the outer conductor between the input and the petals is a piece of copper pipe, $1\frac{1}{2}"$ in diameter. The distance from the petals to the input is $5\frac{1}{2}"$. The petals, there are eight of them, are made of $1/8"$ aluminum plate. They fan out at an angle of approximately 45 degrees to a diameter of $6"$, and then are parallel to the central axis for another $2\frac{3}{4}"$. The four conductor bundles run parallel to the axis for $10"$, then back to the center conductor at an angle of 50 degrees. The mass of the thruster is approximately 1.8 kg.

The second step in testing the thruster was to provide power to it. The power is fed through a pair of 14 gauge wires, surrounded with copper braid, effectively making a high current coaxial cable.

The shape of a typical current pulse when operating into a 0.05Ω resistive load is shown in Fig. 3. Here the pulse height is 2400 A, with rise and fall times of about $13 \mu s$ and $314 \mu s$, respectively. However, when driving the thruster via our made-up coaxial cables, the amplitude was reduced by the increased impedance to about 1200 amps peak, and the rise and fall times increased to about $20 \mu s$ and $400 \mu s$, respectively. The pulse repetition rate was 30 pulses per second. To the best of our ability, we duplicated the pulse shape shown in Fig. 4, which is a sketch from Ref. 1.

This brings us to actually measuring the force. After considering various schemes, we decided to hang the thruster as a pendulum. This provides a low friction

mount, and a convenient means for measuring the force. Any force produced by the thruster produces a displacement from its equilibrium position $d=L(F/W)$, where L is the pendulum length, F is the force produced, and W is the thruster's weight.

In order to increase our ability to detect any motion of the thruster, a simple lens (focal length = $4"$) was attached to the thruster, and the beam from a small laser was passed through the lens and onto an opposite wall. The displacement of the spot on the wall, D , is related to the thruster displacement d , by the relation $D = (R/f)d$, where R is the distance from the lens to the wall and f is the focal length of the lens. In our case $R = 9'4"$ and $f = 4"$, so the magnification factor $R/f = 28$.

With this setup, we could estimate the upper limit for the thrust produced by the antenna. Usually there was spot displacement of perhaps a centimeter upon initial energizing of the thruster. It was not repeatable during a test session, and we attributed it to momentary stiffening of the power cable. The power supply was equipped with a switch that allowed us to apply power to the thruster in bursts. We tried to apply power at the natural frequency of the pendulum, in hope of seeing a greater displacement, but to no avail.

Thus we (generously) take the maximum observable spot displacement as 1 cm. This translates to a pendulum displacement of $(10^{-2} \text{ m})/28 = 3.6 \times 10^{-4} \text{ m}$. The length of the pendulum was $88" = 2.2 \text{ m}$, so the ratio d/L is $(3.6 \times 10^{-4} \text{ m})/(2.2 \text{ m}) = 1.6 \times 10^{-4}$. The weight of the thruster is $1.8 \text{ kg} \times 9.8 \text{ ms}^{-2} = 17.6 \text{ newtons}$, so the upper limit for the thruster force is $17.6 \text{ N} \times 1.6 \times 10^{-4} = 2.8 \times 10^{-3} \text{ N} = 6.3 \times 10^{-4} \text{ lbf} = 1.0 \times 10^{-2} \text{ ozf}$. This is considerably less than the force expected by the authors: 0.03 to 0.3 N at 4 amps, 28 volt dc. (Ref. 5). But, in a sense, the amount of thrust produced by this device is irrelevant to spacecraft propulsion. That is the topic of the next section.

Is There Hope for a Space Drive Within Classical Electrodynamics?

We should recognize, and the earlier the better, that there is little hope for a propulsion scheme that appears as a space reaction drive to an external inertial observer and is based only on the exceedingly well established Maxwellian electrodynamics. The heart of the matter is that classical electrodynamics and mechanics requires conservation of momentum in flat spacetime. Without the existence of an ether-like medium that could sustain volume forces, the momentum of an isolated

electrodynamic system (matter plus the EM fields it generates) is strictly conserved. And of course, Maxwell's equations are incompatible with that kind of ether. Thus a scheme based on classical concepts can be at most a photon rocket.

Since space reaction schemes based just on EM fields keep being proposed, it may be worthwhile to review the derivation of the EM momentum theorem in order to appreciate its compelling nature. Taking the macroscopic view of material properties, formal manipulation of Maxwell's equations leads to the divergence relation

$$\nabla \cdot \bar{T} = \rho \bar{E} + \bar{J} \times \bar{B} + \frac{\partial}{\partial t} (\bar{D} \times \bar{B}),$$

where

$$\bar{T} \equiv \bar{E} \bar{D} + \bar{H} \bar{B} - \frac{1}{2} (\bar{E} \cdot \bar{D} + \bar{H} \cdot \bar{B}) \bar{I}$$

is the Maxwell stress tensor (in dyadic notation). This form of \bar{T} neglects the various internal stresses in matter, due to magneto- and electro-striction and nonuniformity of polarizabilities; however, these terms do not contribute to the volume integral of $\nabla \cdot \bar{T}$.

Although the terms in the $\nabla \cdot \bar{T}$ have the dimensions of force density and $\bar{F}_v \equiv \rho \bar{E} + \bar{J} \times \bar{B}$ is clearly the usual Lorentz force density on matter, nothing in the derivation requires that every term in $\nabla \cdot \bar{T}$ be interpreted as a force density. Forcing such an interpretation on $\frac{\partial}{\partial t} (\bar{D} \times \bar{B})$ suggests the dubious

notion of a force on EM fields or even space itself! It may be best to avoid such interpretations, with their ether-like notions, and to proceed with the integration leading to the momentum theorem, as per standard textbooks in electrodynamics (Ref. 6). The physical interpretation is then thrown back to well established concepts. Under the assumption that the Lorentz force is the only force on the matter in our electrodynamic system, the rate of change of its mechanical momentum is just

$$\frac{d\bar{p}}{dt} = \iiint_V \bar{F}_v dv,$$

where the volume of integration V encloses the system. Hence in this notation,

$$\frac{d}{dt} \left[\bar{p} + \iiint_V (\bar{D} \times \bar{B}) dv \right] = \iint_A \bar{T} \cdot \hat{n} dA .$$

Letting the surface A of V expand in all directions, the above surface integral vanishes rigorously. To see this, observe that the nonradiative components of the fields

decay faster than $1/r$ with distance, thus contributing terms to $\bar{T} \cdot \hat{n}$ that decay faster than $1/r^2$. And if the surface is sufficiently far away, then any radiative fields turned on at a finite time in the past have not yet reached the surface – all photons are caught within the surface and taken into account. The result is the well-known, and relativistically valid, momentum theorem:

$$\bar{p} + \iiint_V (\bar{D} \times \bar{B}) dv = \text{const.}$$

The above volume integral is easily interpreted as the total momentum in the EM field and for free space, the integrand reduces to the familiar $(\bar{E} \times \bar{H})/c^2$.

This momentum conservation law clearly shows that from the point of view of classical mechanics, the validity of Maxwell's equations in flat spacetime limits the space drive to at best a photon rocket. One might try to argue such things as what is the EM momentum in matter etc., but this is at the level of 'tweaking' and not a solution to practical interstellar transport. This severe bottom line strongly suggests that for practical, globally fast mass/energy transport, one must work around the classical limitations of momentum conservation by digging into the deeper layers of spacetime structure itself – the so called "spacetime engineering". Quite likely, that will bring one to deal with the vacuum zero-point fluctuations, the so-called virtual particle pairs and possibly other elements of the support of spacetime.

Summary and conclusions

No reproducible evidence was ever seen for any thrust when current pulses of 1200 A peak and 20 μ s risetime were applied to the antenna at a rate of 30 per second. The result was still null, even when bursts of these pulses were applied in synch with the period of the pendulum-like suspension of the antenna. Our experiment could detect a steady force as low as 3 mN.

The simplicity and import of the electromagnetic momentum theorem underscore the hopelessness of any space reaction scheme strictly within classical electrodynamics. This severe bottom line strongly suggests that for practical, globally fast mass/energy transport, one must work around the classical limitations of momentum conservation by digging into the deeper layers of spacetime structure itself – the so called "spacetime engineering". Quite likely, that will bring one to deal with the vacuum zero-point fluctuations, the so-called virtual particle pairs and possibly other elements of the support of spacetime.

Acknowledgements

The authors would like to thank Carl Lorenzo and Ken Weiland, who constructed the antenna, John Wrbanek, who helped with the figures, and Dr. Norm Wenger, who contributed to the analysis.

References

1. R. L. Schlicher, et al., United States Patent 5,142,861, Nonlinear Electromagnetic Propulsion System and Method, Sept. 1, 1992
2. R. L. Schlicher, et al., Nonlinear Electromagnetic Propulsion System and Method, 19th Power Modulation Symposium, IEEE, New York, 1990, p139-145.
3. R. L. Schlicher, A. W. Biggs, W. J. Tedeschi, Mechanical Propulsion from Unsymmetrical Magnetic Induction Fields, 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, AIAA 95-2643, July, 1995.
4. R. L. Schlicher, A. W. Biggs, W. J. Tedeschi, "Antenna Geometry" Mechanical Propulsion from Unsymmetrical Magnetic Induction Fields, 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, July, 1995.
5. R. L. Schlicher, A. W. Biggs, W. J. Tedeschi, "Theoretical Analysis", Mechanical Propulsion from Unsymmetrical Magnetic Induction Fields, 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, July, 1995.
6. W.K.H. Panofsky and M. Phillips, *Classical Electricity and Magnetism*, 2nd Edition, Addison-Wesley Publishing Co., 1962. pp. 181-184.




Figure 1. A typical NEMPS antenna, illustrated by R.L. Schlicher in his patent 5,142,861.




Figure 2. Our rendition of a NEMPS antenna, suspended from the ceiling. The coaxial Cable for current feed is visible on the right. Note also the lens for detection of motion by means of a laser beam.




Figure 3. Typical pulse shape at maximum capability: 2400 A peak into a 0.05Ω resistive load. When driving the thruster antenna through our coaxial cable, the pulse height was reduced to about 1200 A, at rise and fall times of about 20 μ s and 400 μ s, respectively.




Figure 4. Current pulse shape sketch shown in the Schlicher patent. No scales are given.



A01-34344

**AIAA 2001-3658
Specially Conditioned EM Radiation Research
with Transmitting Toroid Antennas**

H.D. Froning Jr.
Flight Unlimited
Flagstaff, AZ, USA

G.W. Hathaway
Hathaway Consulting Services
Toronto, ON, Canada

**37th AIAA/ASME/SAE/ASEE Joint Propulsion
Conference and Exhibit**
8-11 July, 2001
Salt Lake City, UTAH

For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

SPECIALLY CONDITIONED EM RADIATION RESEARCH WITH TRANSMITTING TOROID ANTENNAS

H. David Froning
Flight Unlimited
5450 Country Club Dr.
Flagstaff, AZ 86004, U.S.A.

George Hathaway
Hathaway Consulting Services
39 Kendal Ave.
Toronto, ON/M5R 1L5, Canada

Abstract

Experimental work to: (a) determine em field characteristics associated with em radiation created by alternating current flowing through toroidal coils at resonant frequencies, and (b) determine if the specially conditioned em fields associated with such radiation could cause a discernable gravity modification, is described. This experimental work was the result of collaboration between Flight Unlimited (FU) and Hathaway Consulting Services (HCS), performed at the laboratories of HCS in Toronto, Canada during a test period in 1998 and during a test period in 2000. Tested toroid configurations included: circular toroids with differing diameters and winding densities; and asymmetrical toroids for focusing em radiation into narrower and more intense beams. The toroid configurations and the AC power and instrumentation systems available at HCS limited the experimental work to the relatively low radio frequencies (400 kHz to 110 MHz) of the electromagnetic spectrum.

INTRODUCTION

Just as airflight was not revolutionized until propeller propulsion was superceded by a new mode of impulsion (jet propulsion), so spaceflight may not be revolutionized until jet propulsion is superceded by a new mode of impulsion (field propulsion). Field propulsion would develop thrust by actions and reactions of fields instead of by combustion and expulsion of mass. And field actions and reactions that would greatly reduce propellant (the major portion of rocketship mass) and engine thrust requirements would be those that would reduce the resistance of gravity and inertia to ship acceleration.

One conceivable way of reducing the resistance of gravity and inertia is by accomplishment of a favorable coupling between those fields which underlie

electromagnetism and gravity. But no significant coupling of ordinary em fields with those that give rise to gravity may be achievable because their essence is completely dissimilar. Yang (1) notes that "nonabelian" fields which probably give rise to gravity are of more intricate topology and higher internal symmetry than the "abelian U(1)" fields that underlie ordinary electromagnetism. In this respect, Barrett (2, 3) has identified two ways of transforming ordinary em fields into specially conditioned em fields of nonabelian form and higher than U(1) symmetry. One identified way of creating such fields is modulating the polarization of em wave energy emitted from microwave or laser transmitters. Such polarization modulation creates em fields of nonabelian form and SU(2) symmetry within beams of radiated power that can be focused into very narrow beams of very high energy density. Thus an experiment to detect possible gravity modifications within narrow polarization modulated laser beams has been submitted to the NASA Breakthrough Propulsion Physics (BPP) program. This experiment is described in (4).

Another way of transforming ordinary em fields into specially conditioned em fields of nonabelian form and SU(2) symmetry is with toroidal coils through which alternating current is flowing at resonant frequencies. Barrett (3) shows that such specially conditioned em radiation includes - not only electric and magnetic field energy - but A Vector potential field energy as well. Barrett predicts that A Vector field intensity maximizes at discreet resonant frequencies. Thus, if an A Vector potential field underlies the essence of gravitation, gravity modification might be possible in the vicinity of toroids transmitting at such frequencies.

Fabrication and testing costs were significant for polarization modulated laser beams. However, they were found to be relatively modest for toroidal coils

"Copyright© 2001 by H.D. Froning Jr.

Published by the American Institute of Aeronautics
and Astronautics, Inc. with Permission

configured for operation in the lower (radio-frequency) range of the em spectrum. Thus a cooperation between Hathaway Consulting Services (HCS) and Flight Unlimited (FU) was established to: test Barrett's hypotheses as to specially conditioned em radiation emitted from toroidal coils; and to determine if gravity modification could occur within such radiation. Probability of gravity modification by radio frequency radiation from inexpensive toroids was deemed to be very low. But it was hoped that the tests would reveal interesting electromagnetic phenomenon and extend our knowledge of electromagnetics.

TRANSMITTING TOROID ANTENNAS

EM wave propagation by transmitting toroid antennas has been examined by various investigators for more than a decade. Examples are U.S. Patent No. 4,751,515 awarded to Corum for an "electrically small, efficient electromagnetic structure that may be used as an antenna or waveguide probe" and U.S. Patent No. 5,442,369 awarded to Van Voorhies et al. for an antenna that "has windings that are contra wound in segments on a toroid form and that have opposed currents on selected segments". In this respect, Barrett (4) has shown that specially conditioned em fields of SU(2) symmetry and nonabelian form can be created by transmitting toroid antennas - as in Figure 1.




Fig. 1 – A Vector Potential Patterns

The magnetic and electric fields which encompass a transmitting toroid are accompanied by A Vector potential fields, and the alternating current flow

produces overlapping A Vector potential patterns which encircle the toroid ring, as shown in Figure 1. These A Vector patterns combine into "phase factor" waves which represent disturbances in A Vector potential. The maximum disturbance in A Vector potential occurs as phase factor wave intensity peaks at the resonant frequencies where A Vector potential patterns are exactly out-of-phase, and a predicted pattern of these disturbances is shown in Figure 2.




Fig. 2 – “Phase Factor” Wave Patterns

Resonant frequencies are determined by the shape and dimensions of the toroid, and by the propagating direction and speed of the alternating electric current thru its windings. And, if an A Vector potential field underlies the essence of gravitation, the probability of gravity modification in the toroid vicinity would be highest at resonant frequencies.

INITIAL TOROID EXPERIMENTS

Initial experimental work involved: (a) fabrication of transmitting toroid antennas that, according to (4), should emanate specially conditioned em radiation; and testing of the toroids at low power levels at the laboratories of HCS in Toronto. The general goals of this initial work, which was performed on March 6 and 7 during 1998, were: perfection of techniques for fabricating toroidal coils, detection of resonant phenomenon indicative of A Vector potential resonances with such coils; and identification of problems associated with operating toroidal coils over wide frequency ranges and at significant power levels.

Most of the goals of the initial work were achieved. Toroid antennas with conventional and caduceus windings were successfully fabricated, and although no instrumentation (such as Josephson Junction arrays) were available for directly detecting A Vector fields, measured resonances (reversals in phase and amplification of signal strength) were in good agreement, as indicated in Figure 3, with Barrett's predictions predicated on occurrence of A Vector fields. Heat generated by current flow within the relatively thin windings of the toroids and their relatively fragile styrofoam interiors limited input power to less than 100 watts in the initial experiments. This identified the need for thicker wires and stronger structures for higher toroid power and temperature.




Fig. 3 – Correlation of Theory and Experiment

FOLLOW-ON TOROID EXPERIMENTS

Because results of the initial toroid experiments were somewhat encouraging, it was decided to have a follow-on experimental program, which included toroids configured for much higher power levels, at HCS between June 9 and June 15, 2000. It included: (a) signal phase/amplitude tests to precisely determine

the resonant frequency characteristics of each different transmitting toroid configuration; (b) magnetic field measurements to map em field intensity in the vicinity of each toroid; (c) propagation characteristics of toroid radiation; and (d) limited gravimeter testing to search for a gravitational disturbance at a one location near one of the transmitting toroids.

Toroid Configurations Tested

To our knowledge, transmitting toroid antennas built and tested by most other investigators have been designed for communication purposes - with wires loosely wound (widely separated) around the toroid's ring in order to maximize far field intensity and range. By contrast, our tested toroids were "tightly wound" to maximize near-field intensity for possible gravity modification - not far field range for communication. Our tested toroids were "contra-wound" in a caduceus pattern to allow 2 types of modulation. One, in which current flowed in opposite directions in crossing wires, resulted in an "opposing" or "bucking" mode which caused opposing magnetic fields that cancel themselves along the toroid ring centerline. The other, in which current flowed in the same direction - resulted in an "adding" mode. Figure 4 shows the 4 different toroid configurations that were tested during the follow-on experimental program.

The loosely-wound toroid (upper left) was built for comparing its near-field intensity with that of the tightly wound toroid (lower left). Both toroids had similar cross sections (approximately 4.0 cm) and the same outer diameters (21cm) and wire size (No. 20). The greater winding density of the tightly wound toroid (350/333 inner/outer turns vs 26/25 inner/outer turns) resulted in greater near-field intensity for a given input power. The toroid in the upper right was configured with a larger outer diameter (31 cm.) than the lower left one and No. 20 wire size but its cross-section is the same. The larger diameter resulted in more windings (398/384 inner/outer turns of No. 14 wire). And the "tear drop" shaped toroid (lower right) was configured to focus radiation into more intense and elongated beams. Its length, breadth and thickness was 26.5, 18.0, 2.5 cm. It had a hole diameter of 7.3 cm; and 95/88 inner/outer turns of No. 14 wire. And, because of their stronger structure (hard maple wood) and larger wire diameter, the tear drop and larger diameter toroids could withstand the heating associated with 1.0 KW of radiated em power.





Fig. 4 – Toroid Configurations Tested



As in the first test series, resonant conditions (revealed by reversal in signal phase and rise in signal amplitude) were searched for at all ac frequencies between 400 KHZ and 110 MHZ. This was done for each toroid configuration for current opposing and current adding modes of operation. Equipment used for the resonance sweeps was an HP 4193 vector impedance analyzer. Figure 5 shows part of the test set-up for detecting resonant modes for each toroid configuration and each operating mode.

Although resonances were detected throughout almost the entire 400 KHZ to 110 MHZ frequency spectrum available at HCS, toroid radiation of significant power was only achievable in the 1.0 to 20 MHZ range. Resonant frequencies selected for measuring field characteristics of each toroid were therefore within this range. Selected resonant frequencies for the large diameter toroid were 2.36 and 17.30 MHZ for current-adding and current-opposing modes of operation, while those for the medium diameter toroid were 2.36 and 18.30 MHZ. Selected resonant frequencies for the

tear drop toroid was 5.66 and 3.94 MHZ for current-adding and current-opposing, while the current-opposing, resonant frequency selected for the loosely wound toroid was 19.70 MHZ.




Fig. 5 – Resonance Sweep Set-Up

Toroid Field Intensity Measurements

Magnetic field intensity was measured out to 50 cm from each toroid center, and along the upper and lower surface of each toroid as well. For an applied power of 10W, the magnetic field component of each toroid's radiation was measured by a small magnetic pick-up coil shown in Figure 6, which converted the actual magnetic field intensity into an equivalent electric field strength (in micro volts per meter).




Fig. 6 – Magnetic Field Probe




Fig. 7 –Toroid Range Test Set-Up

Variation of the large diameter toroid's field strength with range (out to 10 meters) was measured with various types of antennas outside the HCS facility with the test set up as indicated in Figure 7. Data consistent with expected near-field signal strength variation with range was measured when the toroid was radiating in a current-adding resonant mode at 1.20 MHZ. But measurements in a current-opposing resonant mode were anomalous - in that no significant signal strength variation with range was detected.

Search For Gravity Modification

Final toroid testing activity was searching for gravitational field modifications within the specially conditioned em field regions surrounding toroids radiating at resonant frequencies. Gravitational disturbances were searched for with a "Prospector Model 420" gravimeter, manufactured by W. Sodin Ltd, which is capable of detecting changes as small as one-millionth of one percent of ambient gravity. This gravimeter's stainless steel shell and aluminum base does not provide complete magnetic field shielding. But its dewar-enclosed, all-quartz mechanical balance system is not influenced by ordinary em emissions. Unfortunately, preceding test activities took longer than expected, leaving time to search for gravity modification for only one of the toroids (the large diameter one) at only one location with respect to the gravimeter. The limited time remaining also required a very rapid toroid/gravimeter set up. This was achieved by the positioning shown in Figure 8.




Fig. 8 –Toroid/Gravitometer Set-Up

TOROID TESTING RESULTS

Resonant frequencies between 400 KHZ and 110 MHZ were obtained for each toroid. And, for the purposes of mapping magnetic field intensity in each toroid's vicinity, one resonant frequency was selected for each operating mode for each toroid. Field intensity out to 10 m from the large diameter toroid was also measured together with the influences of magnetically shielded structures on its field intensity. Finally, the effect of large diameter toroid field intensity on gravity modification was explored. The results of these efforts are summarized in the following sections.

Toroid Resonance Determination

Resonant frequencies for current-adding and current-opposing operating modes were obtained for each toroid. Figures 9 and 10 show examples of the resonances obtained for the large diameter and tear-drop toroids throughout the 400 KHZ-110 MHZ radio frequency spectrum available at HCS.

Toroid Field Patterns

Magnetic field intensity variation with radial distance for the 3 circular toroids was similar with intensity maximizing near the inner surface of each toroid's ring. And, as would be expected, intensity diminished rapidly with increasing distance above and outside each toroid. Figure 11a and 11b show no definite trend with respect to the influence of toroid diameter. Higher magnetic field intensity is achieved by the smaller diameter toroid in a current-opposing mode of operation while higher magnetic field intensity is achieved by the larger diameter toroid in a current-aiding mode. Figure 11c shows a definite trend - with increased windings over a given toroid geometry resulting in increased magnetic field intensity.

Significant focusing of the electromagnetic energy radiated from the asymmetrical "tear drop" toroid was accomplished. Figure 11d shows that magnetic field intensity is enormously greater at given distances forward of the center of the toroid's hole than for the same distances aft of the hole center. Figure 9d also shows a top and front view of the tear drop toroid field pattern for a given magnetic field intensity. It is seen that more electromagnetic energy is focused into the forward direction than into the aft or side directions

and that the toroids flattened shape (its reduced thickness) causes less radiation to be dissipated in directions transverse to the toroid plane.

One interesting discovery was formation, in the circumferential direction, of standing em waves along the upper and lower surfaces of transmitting toroids. No standing wave measurements were made on the loosely wound toroid. However, numbers of magnetic field peaks and nodes measured circumferentially on the top and bottom surfaces of the other circular toroids were 8 for the medium diameter toroid and 10 for the large diameter one. And at least 6 magnetic field peaks and nodes were measured on the top and bottom surface of the tear drop toroid.

Attenuation of Toroid Field Intensity

Field strength attenuation with range from the center of the large diameter toroid is shown in Figure 12. It was about as expected when radiating at a resonant frequency of 1.2 MHZ in a current adding mode of operation, with a steep signal drop (greater than $(1/r)^3$) out to about 1.0 meter and with an expected near-field $(1/r)^2$ variation between 3 and 10 meters from the toroid. But measurements made with the toroid radiating at a resonant frequency of 17.3 MHZ in a current opposing mode indicated no significant variation in signal strength at distances 3 to 10 meters from the toroid. After continual measurements and re-measurements with various types of antennas, we have no definitive explanation for lack of signal strength reduction with increasing range - other than the possibility of operating slightly off resonance and a significant drop in signal strength.

Additional anomalous behavior may also have been observed for the large-diameter toroid - in that almost identical signal strength was measured at a given location and distance from the toroid when radiating in free space and when radiating from within a magnetically shielded (mu metal) enclosure. These results are considered inconclusive because stray signals were detected from power supply leads which were outside the shielded enclosure. Since measured signals were almost identical for both shielded and unshielded conditions, and since it is unlikely that almost all of the measured free space signal was from the power supply leads, it is conceivable that some of the toroid's em wave energy was propagated through the magnetically shielded mu metal walls.




Fig. 9 – Resonances for Large Diameter Toroid






Fig. 10 – Resonances for “Tear-Drop” Toroid

Fig. 11 – Magnetic Field Patterns in Toroid Vicinity






Fig. 12 - Attenuation of Toroid Signal with Range

Search for Gravity Modification

The possibility of gravity modification in the vicinity of transmitting toroid antennas was briefly investigated by use of the Prospector 420 gravimeter and the large diameter toroid radiating up to 0.5 KW of average power at the resonant frequencies associated with current-aiding and current-opposing operation modes. For these powers and operating modes, no discernable gravity modification was detected for the single toroid/gravimeter positioning that time allowed.

As previously mentioned, time limitations required a rapid toroid/gravimeter test set-up which resulted in the gravitational mass being located in a magnetic field region whose intensity was subsequently found to be much less than magnetic field intensity existing in other locations. For the current opposing mode of operation, measured magnetic field intensity at the gravimeter test mass location was only about 15 percent of the maximum intensity measured near the toroids inner diameter. And toroid magnetic field intensity at the gravimeter test mass location was only about 2 percent of the maximum measured magnetic field intensity for the current-adding mode.

One conceivable reason for non-discriminable gravity modification is, of course, 5 to 50 times less em field intensity at the single location probed by the gravimeter, as compared to locations of maximum intensity. But another reason could be dissimilarity in field topologies associated with toroid em emanations and gravity. And still another reason could be enormous possible differences in the frequencies and wavelengths characterizing gravitational fields and those that characterize electromagnetic fields created by transmitting toroid antennas.

SUMMARY AND CONCLUSIONS

Although interesting phenomenon are associated with em fields created by alternating current flowing at resonant frequencies through toroid coils, no discernable gravity modification (caused by coupling of these fields with those of gravity) was detected.

Interesting electromagnetic phenomenon were: (a) standing em waves along toroid surfaces; (b) em wave energy focused into more intense beams by asymmetrical toroid shapes; and (c) possibly, em wave propagation through magnetically shielded enclosures.

There might have been increased probability of detecting a discernable gravity modification if there had been time for gravimeter measurements in regions where toroid field strength was much greater.

The possibility of anomalous wave propagation should be confirmed or refuted by re-testing the large diameter toroid within a magnetically shielded structure that encloses both the toroid and its power leads.

Zero gravity modification within the radio frequency em fields surrounding transmitting toroid antennas should be confirmed by a gravimeter search throughout the entire vicinity of the large diameter toroid.

ACKNOWLEDGMENTS

This experimental work was motivated by theoretical work by Dr. Terence Barrett with respect to specially conditioned em fields created by transmitting toroid antennas. Dr. Barrett also contributed useful suggestions as to recommended toroid configurations and test procedures. The most recent experimental work was conducted with the considerable assistance of Mr. Blair Cleveland, who was intimately involved in every aspect of the test preparations conduction and data gathering portions of this work.

REFERENCES

- (1) Yang, C.N., "Gauge Theory", McGraw-Hill Encyclopedia of Physics, 2nd Edition, p483, (1993)
- (2) Barrett, T.W., "Electromagnetic Phenomenon not Explained by Maxwell's Equations", Essays on Formal Aspects of Electromagnetic Theory, p6, World Scientific Publ. Co., (1993)
- (3) Barrett, T.W., "Toroid Antenna as Conditioner of Electromagnetic Fields into (Low Energy) Gauge Fields", Proceedings of the: Progress in Electromagnetic Research Symposium 1998, (PIERS '98) 13-17 July, Nantes, France (1998)
- (4) Froning, H.D., Barrett, T.W., "Theoretical and Experimental Investigations of Gravity Modification by Specially Conditioned EM Radiation", Space Technology and Applications International Forum 2000, Editor: Mohamed S. El-Genk, Published by the American Institute of Physics (2000)

A01-34345



AIAA 2001-3660

An Experimental Investigation of the Physical Effects in a Dynamic Magnetic System

V. Roschin and S. Godin
Institute for High Temperatures
Russian Academy of Science
Moscow, Russia

**37th AIAA/ASME/SAE/ASEE
Joint Propulsion Conference and Exhibit
8-11 July 2001
Salt Lake City, Utah**

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

AN EXPERIMENTAL INVESTIGATION OF THE PHYSICAL EFFECTS IN A DYNAMIC MAGNETIC SYSTEM.

Vladimir Roschin, E-mail: rochtchin@mail.ru

Sergei Godin, E-mail: smgodin@online.ru

Institute for High Temperatures, Russian Academy of Science
Izhorskaya st. 13/19, Moscow 127412, Russia

Abstract

In the current paper the results of experimental research of magnetic-gravity effects are presented. Anomalous magnetic and thermal changes within a radius of 15 meters from the researched device were measured as well.

PACS: 41.20.-q; 44.60.+k; 76.50.+q

Introduction

We have experimentally studied the physical effects in a system based on rotating permanent magnets [1]. Below we describe the technology of manufacture, assembly, and the results of testing this experimental setup, which is referred to as the converter.

Received effects:

- Generation of mechanical energy in a self-governing mode of operations;
- Change of weight of the converter;
- Formation of a local magnetic and temperature fields as concentric cylinders around converter.

Description of the Experimental Setup

The basic difficulty arises in choosing the materials and maintaining the necessary magnetic pattern ("imprinting") on the plate and roller surfaces. To simplify the technology we decided to use a one-ring design with one-ring plate (stator) and one set of rollers (rotor). It is obvious, that it was necessary to strengthen the rollers on a rotor by the bearings and balance the rollers well. In the suggested design, air bearings were used which provided the minimum losses due to friction.

From the available description [1] it was not clear how it is possible to make and magnetize the stator with a diameter of about one meter. In order to make the stator from separate magnetized segments executed with rare earth magnets with the residual induction of 1T, the segments were magnetized in a usual way by discharging a capacitor-battery energizer through a coil. Afterwards the segments were assembled and glued together in a special iron armature, which reduced magnetic energy.

"Copyright © 2001 by Vladimir Roschin and Sergei Godin. Published by American Institute of Aeronautics and Astronautics, Inc. with permission."

To manufacture the stator 110 kg of rare earth magnets were used. To manufacture the rotor 115 kg of that material was used. High-frequency field magnetization was not applied. It was decided to replace imprinting technology described in [1] with cross-magnetic inserts having a flux vector directed at 90 degrees to the primary magnetization vector of the stator and rollers. For these cross-inserts the modified rare earth magnets with residual magnetization of 1.2 T and coercive force a little bit greater than in a base material were used. Fig.1 and Fig.2 show the joint arrangement of stator 1 and rotor, made up of rollers 2, and a way of their mutual gearing or sprocketing by means of cross magnetic inserts 19. Between the stator and roller surfaces air gap δ of 1 mm is maintained.




Fig. 1. Variant of one-ring converter.

No layered structure was used except a continuous copper foil of 0.8 mm thickness, in which the stator and rollers were wrapped up. This foil had a direct electrical contact to magnets of the stator and rollers. Distance between the inserts in rollers is equal to distance between the inserts on the stator. In other words $t_1 = t_2$ on a Fig.2.




Fig.2. Sprocket effect of magnetic stator and roller inserts.

The ratio of parameters of the stator 1 and the rotor 2 on Fig.2 is chosen so that the relation of stator diameter D to the roller diameter d is an integer equal to or greater than 12. Choosing such ratio allowed us to achieve a resonant mode between elements of a working body of the device.

The elements of magnetic system were assembled in a uniform design on an aluminum platform. Fig.3 displays the general view of the platform with the one-ring converter. This platform was supplied with springs, shock absorbers and had a possibility of moving vertically on three supports. The possible vertical displacement was 10mm and the induction meter 14 was used for measuring this displacement. Thus, the instantaneous change of platform weight was defined during the experiment in real time. Gross weight of the platform with magnetic system in an initial condition was 350 kg.

The stator 1 was mounted motionlessly and the rollers 2 were assembled on a mobile common rotor - separator 3, which is connected with the basic shaft 4 of the converter. Through this shaft the rotary moment was transferred. The basic shaft by the means of friction

muff 5 was connected to a starting engine 6, which accelerated rotor of the converter up to a mode of self-sustained rotation. An ordinary DC electrodynamics generator 7 also was connected to the basic shaft as a main loading of the converter. Adjacent to the rotor, electromagnetic inductors 8 with open cores 9 were located.

The magnetic rollers 2 crossed the open cores of inductors and closed the magnetic flux circuit through electromagnetic inductors 8, inducing an electromotive force in them, which acted directly on an auxiliary active load 10 - a set of incandescent lamps with total active power of 1 kW. The electromagnetic inductors 8 were equipped with an electrical drive 11 on supports 12. Driven coils were used for smooth stabilization of the rotor rpm. The speed of the rotor also could be adjusted by changing the main loading 10.

To study influence of high voltage on characteristics of the converter, a system for radial electrical polarization was mounted. On a periphery of rotor the ring electrodes 13 were set between the electromagnetic inductors 8 and had an air gap of 10 mm with the rollers 2. The electrodes are connected to a high-voltage source; the positive potential was connected to the stator, and the negative - to the polarization electrodes. The polarizing voltage was adjusted in a range of 0-20 kV. In the experiments, a constant value of 20 kV was used.

In case of emergency braking, a friction disk from the ordinary car braking system was mounted on a basic shaft of the rotor. The electrodynamics generator 7 was connected to an ordinary passive resistive load through a set of switches guaranteeing a step connection to the load from 1 kW to 10 kW - a set of ten ordinary electric water heaters.




Fig.3. The general view of the converter and its platform.

Converter under testing had in its inner core the oil friction generator of thermal energy 15 intended for directing a superfluous power (more than 10 kW) into the

thermo-exchange contour. But since the real output power of the converter in experiment has not exceeded 7 kW, the oil friction thermal generator was not used.




Fig.4. A diagram illustrating various operation regimes of the converter,
 $-G$, $+G$ changes in weight of the platform vs. rpm

Experimental results

The magnetic converter was mounted in a laboratory room on three concrete supports at a ground level. The ceiling height of lab room was 3 meter. A common working area of the laboratory was about 100 sq. meters. Besides a presence of an iron-concrete ceiling in the immediate proximity from the magnetic system there was a generator and electric motor, which contained tens of kilos of iron and could potentially deform the field structures.

The device was started by the electric motor, which accelerated rpm of the rotor. The revolutions were smoothly increased up to the moment the ammeter included in a circuit of the electric motor begin to show zero or a negative value of consumed current. The negative value indicated a presence of back current. This back current was detected at approximately 550 rpm under clockwise rotation. The displacement meter 14 starts to detect the change in weight of the whole installation at 200 rpm. Afterwards, the electric motor is completely disconnected by the electromagnetic muff and an ordinary electrodynamics generator is connected to the switchable 10 kW resistive load. The converter rotor continues self-accelerating and approaches the critical regime of 550 rpm when the weight of the whole installation quickly changes.

In addition to dependence on a speed of rotation, the common weight of device depends from output power on the load and from applied polarizing voltage

as well. As seen on Fig.4, under maximum output power is equal to 6-7 kW the change of weight ΔG of the whole platform (total weight is about 350 kg), reaches 35 % of the weight in an initial condition G_i . Applying a load of more than 7 kW results in a gradual decrease of rotation speed and an exit from self-sustained mode (right sides of the curves on Fig.4 for a 7 kW loading).

The net weight G_n of the platform can be controlled by applying high voltage to polarization ring electrodes located at a distance of 10 mm from external surfaces of the rollers. Under a high 20 kV voltage (electrodes have a negative polarity) the increase of tapped power of the basic generator more than 6 kW does not influence ΔG if rotation speed is kept above 400 rpm. "Tightening" of this effect is observed as well as the effect of hysteresis on ΔG at rotation of a rotor on a clockwise and counter-clockwise (a kind of "residual induction"). The experimental diagrams given on Fig.4 illustrate the $+G$ and $-G$ changes in weight of the converter vs. rotor rpm. The effect of a local change of the platform weight is reversible, relative to the direction of rotor revolution, and has the same hysteresis. A clockwise rotation causes the critical regime to occur in area of 550 rpm and a propulsion force against the direction of gravitation vector is created. Correspondingly, a counter-clockwise rotation causes the critical mode to occur in area of 600 rpm and a force in the direction of gravitation vector is created.

The difference in approach to a critical regime of 50 - 60 rpm was observed. It is necessary to mention that the most interesting region are situated above the critical area of 550 rpm, but due to of a number of circumstances the implementation of such research was not possible. Probably, there are also other resonant modes appropriate to higher rpm of a rotor and to the significant levels of useful loading and weight changing. Proceeding from the theoretical assumptions, the dependence of tapped mechanical energy from the pa-

rameters of magnetic system of the converter and rpm of a rotor has a non-linear character and the received effects are not optimum. From this point of view, the revealing of a maximal output power, of maximal change of weight and converter resource represents a large practical and scientific interest. In tested sample of the converter the using of higher rpm was inadmissible because of unsufficient mechanical durability of the magnetic system, which has been stuck together from separate pieces.




Fig.5 Diagrams of rotor accelerating and loading of the converter.

On Fig.5 the dependence of total weight of whole installation and its output power directed into active loading from rpm of a rotor of the converter is explained in detail. The diagrams are constructed for a case of HV polarization ON (top diagram) and HV polarization OFF (bottom diagram). The time from starting of the engine up to a mode of self-sustaining of the converter at the rotation of a rotor clockwise approximately is equal 1.5 minutes. Power of starting DC engine was about of 2 kW and reduction on a shaft of the converter was equal 1/10. At achievement of a critical mode (550 rpm.) the change of gross weight of a platform already achieves +/-30% from G_i . Under transition to resonant mode the revolutions with the large acceleration have increases up to 590 rpm and weight has changes up to +/-35% from G_i . This point on the diagram begins at once after a critical point (inclination of

a curve α_1). At achievement of 590 rpm the first stage of active loading 1 kW is connected to the electrodynamics generator. The revolutions have a sharply reducing and ΔG also is changing. As soon as the revolutions begin to grow again, the second switchable loading is connected and rotor rpm are stabilized at a level of 590-595 rpm. ΔG continues to change. The increasing of switchable loading occurs by steps on 1 kW up to total power of 6 kW. All intervals are equal approximately of 10-30 sec. Afterwards the short-term increasing of revolutions and then the full stabilization of the 6 kW output during of 12-15 min was observed.

More than 50 launches of the converter with an absolute repeatability within three months were carried out. It is necessary to note that revolutions will grow with acceleration shown on Fig.5 by angles $\alpha_1 \dots \alpha_5$, if do not switch on the next step of loading to the genera-

tor at rpm increasing. For returning to a previous rpm mode it is necessary to switch on a twice more loading.

The words above concern a mode with switched ON a high voltage polarization of 20 kV, "plus" is on a grounded stator. Without polarization voltage (lower curve on Fig.5) the diagram is approximately the same, but is well indicated the more soft character of a loading and faster changing of weight of a platform due to decreasing of rpm.

Other interesting effect is corona discharges, which was observed at the work of the converter in a dark room. At this, around the converter rotor a blue-pink glowing luminescence and a characteristic ozone smell were noticed. On Fig.6 the cloud of ionization covers area of the stator and rotor and is having accordingly a toroidal form.




Fig.6. Corona discharges around the converter.

On a background of luminescence glowing on the roller surfaces, we distinguished a separate «wave picture». A number of more vigorous strips of discharges around the rollers were observed. These discharges were of a white-yellow color but the characteristic sound for arc discharges was not audible. There were not noticed any erosive damages by arc discharges on surfaces of the stator and the rollers as well.

One more effect previously not mentioned anywhere was observed - the vertical concentric magnetic "walls" around the installation. We noticed and measured a permanent magnetic field around the converter within a radius of 15 meters. For this magnetic field measurement a Russian made magnetometer F4354/1 was used. Magnetometer had a Hall-effect sensor in a copper shielding. The zones of increased intensity of a magnetic flux 0.05 T, located concentrically from the center of the installation were detected. The direction of the magnetic field vector in these walls coincided with direction of magnetic field vector of rollers. The structure of these zones was like the Bessel function of zero order of two arguments. No any magnetic fields were registered between these zones by portable magnetometer. The layers of increased intensity were distributed practically without losses up to a distance of about 15 meters from a center of the converter and had a quick decreasing at a border of this zone. The thickness of each layer was approximately of 5 - 6 cm the border of each layer was very sharp. The distance between layers was about of 50 - 80 cm where the upper value is seen when moving from center of the converter. A stable picture of this field was observed as well as at a height of 6 m above the installation (on the second floor above the lab). Above the second floor, measurements were not carried out. The similar picture was observed and outside of a room of laboratory, directly in the street, on the ground. The concentric walls were strictly vertical and no had appreciable distortions. The Fig.7 illustrates the schematic placing of the converter in a room of laboratory and arrangement of concentric magnetic and thermal fields around the installation.




Fig.7. Schematic placing of the converter and field patterns in lab room.

An anomalous decrease of temperature in a vicinity of the converter was also found. While the common temperature background in laboratory was + 22°C ($\pm 2^\circ\text{C}$) the stable fall of temperature of 6-8°C was noticed. The same phenomenon was observed in concentric vertical magnetic walls as well as. The measurements of temperature inside the magnetic walls were carried out by an ordinary alcohol thermometer with an inertia of indication about 1.5 min. Inside the magnetic walls the temperature changes can even be distinctly observed by hand. When the hand is placed into this magnetic wall the cold is felt at once. A similar thermal picture was observed at height above the installation, i.e. on a second floor of the laboratory as well as despite the steel-reinforced concrete blocks of a ceiling and also on an open air outside of the laboratory.

Concentric magnetic walls and accompanied thermal effects begin to appear approximately from 200 rpm and have a linearly increasing with speeding up of revolutions up to a critical regime. The measurements above 600 rpm were not made because of fear of destruction of magnetic system. On Fig.8 the curve of intensity of magnetic field in mT and change of temperature in Celsius degrees due to rpm changing is represented.




Fig.8. The dependence of intensity of magnetic field and changes of temperature vs. rotor rpm of the converter.

Discussion

All the results we obtained are extremely unusual and require some theoretical explanation. Some theoretical approaches and phenomenological descriptions can be found in our book [2]. Unfortunately, interpretation of these results within the framework of conventional physical theory cannot explain all observed phenomena especially the change of weight. We can interpret change of weight as a local change of gravitation or

as some propulsion force like a propeller work in air. Only a role of a propeller the magnetic system is playing, and a role of air - the quantum environment - ether filled with a "dark matter and energy". Direct experiment, which would have confirming the presence of a propulsion force was not performed, but in any case both interpretations of the weight change do not correspond to the modern physics paradigm.

In conclusion we would like to emphasize that issues of biological influence and especially effects of unknown radiation around of the converter were not considered at all. Our own experience allows us to do only cautious assumption that the short-term stay (dozen minutes) in a working zone of the converter with fixed output power of 6 kW remains without observed consequences for exposed persons.

Bibliography

1. J. A. Thomas, Jr., ANTI-GRAVITY: the Dream Made Reality: the Story of John R. Searl (Direct International Science Consortium, London, 1994), Vol.VI, Iss.2.
2. V.F. Zolotarev, V.V. Roschin, S.M. Godin. About the structure of space-time and some interactions. (The theory and original experiments with extraction of an internal atomic energy). Moscow, 'Prest', 2000, 309p. with illustrations. (in Russian).



A01-34536

AIAA 2001-3907

Rapid Spacetime Transport and Machian Mass Fluctuations: Theory and Experiment

J. Woodward and T. Mahood

California State University

Fullerton, CA

P. March

Lockheed Martin Space Operations

Houston, TX

**37th AIAA/ASME/SAE/ASEE
Joint Propulsion Conference and Exhibit
8-11 July 2001
Salt Lake City, Utah**

For permission to copy or to republish, contact the copyright owner named on the first page.

**For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.**

RAPID SPACETIME TRANSPORT AND MACHIAN MASS FLUCTUATIONS: THEORY AND EXPERIMENT

James F. Woodward,* Thomas L. Mahood,* and Paul March†

ABSTRACT

The various schemes for achieving rapid spacetime transport currently in circulation are briefly reviewed, and then the methods based on Machian mass fluctuations are considered. Results of experimental work designed to detect a negative-definite Machian mass fluctuation expected on the basis of simple theory are then reported. They suggest that such mass fluctuations may in fact occur and have sufficient magnitude to be engineerable into practical devices.

INTRODUCTION

It is a commonplace that although rockets may serve to cover interplanetary distances, however inefficiently, when interstellar voyages in some reasonable fraction of a human lifetime are considered, conventional rocketry comes up seriously wanting. The fundamental problem, simply put, is: How does one accelerate a reasonable payload to some significant fraction of the speed of light in short order without having to lug idiotically large amounts of propellant along to push off of? Only a moment's reflection is required to see that, short of creating "wormhole" spacetime shortcuts with "exotic matter", there are two distinct, but related ways to tackle this problem. One is to reduce the mass of the payload. The other is to find some means of producing an accelerating force on the payload without lugging along all the propellant required by rockets. Conventional approaches of considerable ingenuity have been devised in both cases. But they still fall far short of the technical requirements for serious interstellar travel. The question then is, do the laws of physics allow the invention of unconventional spacecraft – starships – equal to interstellar voyages?

In the matter of payload reduction, ideally one would want to render the mass of a ship and its con-

tents as nearly equal to zero as possible. Were that possible, only minute forces would be needed to accelerate the ship to near lightspeed. Is this possible? To answer this question we must understand the cause of the inertial properties of matter; that is, why things push back on us when we try to accelerate them.

At the moment three putative causes of inertia are advocated. There are those who simply assert that inertia is an innate property of matter that is independent of all external circumstances. Others believe that inertia results from the local action of quantum mechanical "zero point" fluctuations of the electromagnetic field (and perhaps other non-gravitational fields) on things that are accelerated. And some are convinced that inertial reaction forces are caused by the gravitational action of chiefly distant matter in the universe. If inertia is an inalterable, innate property of matter, then the only way to reduce the mass of a ship is to find and collect some "exotic" matter (with negative mass) to offset the positive mass of the payload. Such material, naturally occurring, has yet to be discovered in nature.

Compelling reasons exist to set aside the electromagnetic "zero point" fluctuations explanation of inertia.^{1,2} But even were that not the case, it seems exceedingly unlikely that any means could ever be found to suppress zero point fluctuations over the volume of a starship (thereby supposedly rendering it essentially inertialess). The Casimir effect, often cited in connection with these arguments, does point up the fundamental problem here. It is the low frequency part of the vacuum fluctuation spectrum that is suppressed by material objects in close proximity that produces this effect. But, alas, it is the high frequency part of the fluctuation spectrum that is chiefly responsible for the induction of the vast bulk of inertia in this scheme. (We note here that Lorentz invariance requires that the spectral energy density of the vacuum fluctuations increase with the cube of the frequency thereof.)

Taking inertial forces to be due to the gravitational action of chiefly distant matter in the universe – as is the case in general relativity theory – leads to two possible approaches. One is to find some means of "shielding" the bulk of the starship from the action of the gravity field of the distant matter. This might be done either passively by finding a "gravity insulator" – H.G. Wells' favorite that is – or actively by devising some apparatus that generates a gravity field that can

* California State University, Fullerton, CA 92634;
email: jwoodward@fullerton.edu

† Lockheed-Martin Space Operations, 2400 NASA Rd.
1, MS: C18, Houston, TX 77058-3799

Copyright © 2001 by James F. Woodward. Published
by the American Institute of Aeronautics and
Astronautics, Inc. with permission.

be used to "null" the gravity field produced by distant matter. Cavorite, like naturally occurring "exotic" matter, remains to be discovered. Several active methods have been explored – notably spinning superconducting disks in suitable electromagnetic fields – so far to no avail. One troubling aspect of these schemes is that they do not follow from the widely accepted form of the laws of physics. And they usually lead to violations of fundamental conservation laws. The second approach consists of finding a way, within the structure of accepted physical laws, to transiently induce a reduced mass state, perhaps fleetingly even an "exotic" mass state, in normal matter by some technically feasible means. This may sound quite implausible, but it turns out that there may be a way to do this. We discuss this, along with some experimental results, below.

Turning to the issue of generating accelerating forces, we see that the fundamental problem, irrespective of the propulsion system being considered, is how to push off of the distant matter in the universe. In conventional rocket propulsion systems, this is achieved by using propellant as an intermediary. That is, when we push on propellant, the gravitational action of the chiefly distant matter in the universe causes the propellant to push back on us, causing an accelerating force on us. Since the prodigious propellant payload is the chief impediment to interstellar voyages, one might want to devise a propulsion scheme that involves, ideally at least, no lugged propellant whatsoever. In conventional systems this requirement translates to either "harvesting" propellant locally as one goes along, using beamed energy from home with "sails", or finding a way to couple locally to cosmic fields of one sort or another. None of these seem very realistic at present.

Alas, the strictly propellantless "break-through" schemes appear even less realistic than the conventional proposals, for they all seem to violate some fundamental, well-tested physical principle. Usually, but not always, it is the conservation of momentum that bites the vacuum. The best known of these schemes involves the aforementioned spinning superconducting disks. In these circumstances, however, the action of the disks is not interpreted as shielding or attenuating the gravitational field of other objects. Rather, it is regarded as generating a directed "gravity or inertia-like" field analogous to the "gravitomagnetic" field generated by spinning objects that is predicted by general relativity, but magnified in intensity by many orders of magnitude. This putative force does not act on the generating apparatus *per se*. But aimed at distant matter in the direction one wants to go, it acts on the distant matter and a mutual force is established that accelerates the spacecraft. No violation of momentum conservation is necessarily involved here.

But there is no compelling reason to believe that when electric charge carriers form Cooper pairs (as they do in the superconducting state), no matter how they are acted upon by external electro-magnetic fields, they acquire the magical ability to create from nothing the gargantuan "inertialike" fields claimed.

Another strictly propellantless propulsion scheme is based on the observation that electromagnetic circuits can be arranged so that they appear to generate self-forces – that is, forces appear in one part of the circuit that seem not to be cancelled by equal and opposite forces anywhere else in the circuit. Perhaps the best example of this type of scheme is the "Slepian" approach advocated by Dering, Corum, and several of their associates.³ The issue of "hidden momentum" in electrical circuits has a long history. Suffice it to say that the generation of a net force in an isolated electrical circuit capable of accelerating a spacecraft is a very dubious proposition. To be relativistically (and empirically) correct, the laws of electrodynamics must be invariant under the group of infinitesimal Lorentz transformations. This invariance insures that energy and momentum are conserved in isolated systems. Aside from the obvious violation of the conservation of momentum involved in the Slepian scheme, simple physical common sense suggests that it is not reasonable. Accelerations take place with respect to the distant matter in the universe. The likelihood that one could produce accelerations without coupling to that matter in any way is, on its face, pretty preposterous. Well, unless the quantum vacuum couples to the distant matter and can be acted upon in some way by the local electromagnetic fields. But that is not what is claimed in "Slepian" systems.

Should it prove possible to produce transient variations in the masses of material objects, another means of propulsion may be achievable. Here some local mass is used as a propellant. But instead of simply ejecting the matter once and for all, after it has been "ejected" one operates on it to reduce its mass so that it can be recovered without canceling the momentum imparted to the craft when it was earlier ejected. It may then be ejected again in its normal state, and the mass-reduced recovery repeated. And this cycle may be repeated over and over again, *ad nauseam*. This may sound like having one's cake and eating it too. But when we keep in mind that the gravity/inertia field couples the craft (and recycled propellant) to the distant matter in the universe, it is apparent that such a scheme merely amounts to producing a directed momentum flux in the gravitational/inertial field by the local manipulations matter. One may object that the propagating "momenergy" in the gravitational field produced by any reasonable concentration of local matter is so ridiculously small that even were such a scheme in principle possible, surely it would be utterly

impractical. But one ignores, in this view, the fact that inertial reaction forces are gravitational actions too. *They are enormous and instantaneous*. So there is hope, should it prove possible to induce a transient mass-reduced state in some propellant without violating the laws of physics.

THEORY

When inertial reaction forces are viewed as arising from the action of a locally Lorentz-invariant field (as they actually are), one finds that objects subjected to accelerations by external forces undergo changes in their restmasses during the accelerations.^{4, 5, 6, 7} This is a consequence of the four-vector nature of the reaction forces – and the fields producing them – and the four-dimensionality of the divergence operator that is applied to the four-field strength to obtain an expression for the local charge density. To see how this works, consider an object accelerated by an external force. An (equal and opposite) inertial reaction force arises in the object owing to the action of the local “inertial” field on it.^{8, 9} The field strength experienced by the object is just the inertial reaction four-force divided by the mass of the object. To get the local source density of this field we apply the four-dimensional generalization of Gauss’s divergence theorem. Taking account of the fact that the field is irrotational (so the three-force can be expressed as the gradient of a scalar potential) and translating charges into charge densities (as is customary in constructing field equations), this yields:

$$\nabla^2\phi - (\phi/\rho_0 c^2)(\partial^2 E_o/\partial t^2) - (\phi/\rho_0 c^2)^2(\partial E_o/\partial t)^2 = 4\pi G \rho_0, \quad (1)$$

where ϕ is the scalar potential of the inertial field, ρ_0 the local restmass density, c the vacuum speed of light, E_o the local rest-energy density ($=\rho_0 c^2$), and G Newton’s universal gravitational constant.

To transform this equation into a field equation of standard form we need merely note that inertial reaction forces are gravitational forces (Mach’s principle) and that since this is true, in this approximation we have that $\phi = c^2$ and the rest energy density E_o can be written as $E_o = \rho_0 \phi [= \rho_0 c^2]$. Using this expression for E_o in Equation (1) and rearranging we obtain:

$$\nabla \cdot \phi - (1/c^2)(\partial^2 \phi/\partial t^2) = \square \phi = 4\pi G \rho_0 + (\phi/\rho_0 c^2)(\partial^2 \rho_0/\partial t^2) - (\phi/\rho_0 c^2)^2(\partial \rho_0/\partial t)^2 - c^4(\partial \phi/\partial t)^2, \quad (2)$$

the inhomogeneous relativistically invariant wave equation for the scalar gravitational potential. Note the

transient matter source terms on the RHS of Equation (2). If real, they may make rapid spacetime transport technically feasible.

If we ignore $-c^4(\partial \phi/\partial t)^2$ since it is always minuscule given its c^4 coefficient (not compensated for by any factor of ϕ in the numerator, as in the other terms), and extract a factor $4\pi G$, we may write the total, time-dependent matter density as:

$$\rho(t) \approx \rho_0 + (1/4\pi G)[(\phi/\rho_0 c^2)(\partial^2 \rho_0/\partial t^2) - (\phi/\rho_0 c^2)^2(\partial \rho_0/\partial t)^2]. \quad (3)$$

The first time-dependent term – $(1/4\pi G)[(\phi/\rho_0 c^2)(\partial^2 \rho_0/\partial t^2)]$ – can be both positive and negative when ρ_0 undergoes periodic fluctuations. When effects arising from this term are convolved with a second periodic force in an object, this restmass fluctuation can lead to the production of stationary forces.^{10, 11, 7} One merely pushes on an object made more massive by this mass fluctuation, and then pulls it back when it is in a mass-reduced state to achieve a net accelerating force. For this reason we call this the “impulse engine” term in Equations (2) and (3). The second time-dependent term, unlike the first, is always negative, because the negative sign of the term is unaffected by the sign of ρ_0 since it appears exclusively in quadratic factors: $-(1/4\pi G)[(\phi/\rho_0 c^2)^2(\partial \rho_0/\partial t)^2]$. This term, if it can be made large, holds out the possibility of the transient formation of “exotic” matter – matter with negative mass – the material needed to achieve the most extreme types of rapid spacetime transport by the warping of spacetime. We resist the temptation to assign this term a romantic label, calling it merely the “negative mass” term in Equations (2) and (3).

As with the impulse engine term, when mass fluctuations driven by the negative mass term in some object are convolved with a second periodic force, stationary forces can be produced. So both the impulse engine and negative mass effect terms hold out the promise of realizable propellantless propulsion. Note that the transient mass terms in Equation (3) are enormous in comparison with, for example, the mass changes that occur from simply changing the energy content of some region in space-time. This is a consequence of the transient terms being “magnified” by the factor $(\phi/\rho_0 c^2)$ coefficients where, as a consequence of Mach’s principle, $\phi/c^2 \approx 1$, rather than being a very small number as one might otherwise expect when ϕ has a value of the scale normally assumed (that is, $\ll c^2$).

The stationary forces that result from both of these terms depend on the relative phase of the driven mass fluctuation and the second applied force. Their periodicities, as a function of the relative phase, how-

ever, are not the same because for a driven periodic fluctuation in ρ_0 , the resulting contributions from the impulse engine term have the same periodicity as the driving signal, but the negative mass term varies at twice the frequency of the driving signal because the time derivative of the driving signal is squared. Were the negative mass term much smaller than the impulse engine term, as simple calculations can lead one to believe, this would not be a matter of much moment. But, in fact, the negative mass term is comparable to the impulse engine term, as can be seen by inserting the *ansatz* $\rho_0 = \rho_0 \cos(\omega t)$ into Equation (2) and computing the resulting expressions for the impulse engine and negative mass terms. This procedure does not yield an exact solution of this non-linear equation. But it does reveal that the amplitudes of the two factors in the square brackets are the same, to factors of order unity at any rate. Thus, should Machian mass fluctuations actually occur, one should be prepared to see evidence of the negative mass term in any system sensitive enough to detect the impulse engine term.

Before turning to a discussion of experiments designed to test for the presence of the mass fluctuations just described, we mention general considerations that should inform experiments. We ask first, where might one expect to see the sort of mass fluctuations predicted by Equation (2) above? Since this equation follows from the consideration of the action of an accelerating force on matter, plainly some material must be present that is accelerated. The fact that the transient terms in Equation (2) both involve derivatives of the local rest matter density, ρ_0 , indicates that the local rest energy density, E_0 [$=\rho_0/c^2$], must also be changing. Evidently, the largest Machian mass fluctuation effects are to be expected in materials where, viewed at the microscopic level, the application of external electromagnetic fields produce large accelerations accompanied by rapid changes in stored internal energy. Thus, for example, if mass fluctuations are to be driven in dielectric materials in capacitors, then one will want to use material with the highest possible dielectric constant. In particular, ferroelectric substances. It is worth noting, nonetheless, that if the vacuum can be truly regarded as a polarizable substance, then one may expect to drive mass fluctuations in the vacuum itself. And should such fluctuations be convolved with an appropriate periodic force, stationary forces like those expected in material media should be producible. Such

forces, however, would be minuscule in comparison with those produced in materials with large dielectric constants. An experiment designed to test for mass fluctuations in the vacuum would make a good test of the reality of the "polarizable vacuum" model of GRT that has recently been elaborated by Puthoff.¹²

EXPERIMENT

Results of experiments designed to detect Machian mass fluctuations have already been reported.^{4, 5, 11, 13, 7} We report here current results of the ongoing experimental program underway at California State University, Fullerton (CSUF). For the past several years that program was focused on the detection of stationary forces in stacks of lead-zirconium-titanate [PZT] crystals driven with voltage signals of several hundred volts at several tens of kHz using sensitive torsion pendula, started by JFW and TLM in the course of TLM's graduate work. (See refs. 7, 13, and especially 14 for the details of this work.) Since fall 2000 the focus of this work has changed to an updated version of early experimental tests of mass fluctuations by looking for the effect as a weight fluctuation with force transducers.^{4, 5} *Approaching experimental detection of any mass fluctuation effect in this way has the great merit that only the mass fluctuation per se is detected (as a weight fluctuation), so all issues relating to the production of stationary forces by signals of several frequencies and phases are set aside (unless such forces are inadvertently produced by convolution of harmonics of the fundamental signal with mass fluctuations driven thereby that might affect weight determinations).*

The search we are conducting for Machian mass/weight fluctuations, for the moment at least, utilizes PZT stacks like that shown in Figure 1. This PZT stack consists of ten crystal disks, 1.91 cm in diameter made of EDO Ceramics EC-65 material




Figure 1: A. A picture of the device mounted on the force transducer. The thermistor and accelerometer connection are at the top center-left. B. A schematic diagram of the device showing the arrangement of the crystals and accelerometer in the PZT stack.

epoxied together along with suitable electrodes between the crystals. The stack is clamped by six machine screws between an aluminum cap and a 0.95 cm thick brass disk. The pair of crystal disks closest to the brass disk are 0.31 cm thick, whereas all but two of the crystal disks at the other end of the stack are 0.15 cm thick. Thinner disks, with higher stored energy for a given applied voltage to the stack, are used in the part of the stack near the aluminum cap where the largest accelerations, and thus predicted effects, are expected. Two of the crystal disks are very thin, being only 0.03 cm thick. This pair of crystals was placed between the two pairs of 0.15 cm thick crystals closest to the aluminum cap and independently wired so that they could be used as an accelerometer to monitor the behavior of this part of the PZT stack.

An aluminum mounting bracket is attached to the brass disk as shown in Figure 1. Note the thin rubber pad that isolates the bracket from the brass disk. This was included to suppress the transmission of vibration from the stack assembly to the weight sensors used. This pad, however, contributes significantly to improved performance of the stack. It also suppresses the flow of heat into the bracket. The temperature of the stack assembly is monitored with a thermistor glued onto a small aluminum ear that is also attached (without a rubber pad) to the brass disk. (This aluminum ear also carries a bimetallic strip thermometer used in earlier work.)

Two different weight detection systems were developed for the work we report here. The chief system employed a Unimeasure U-80 position sensor configured as a load cell. (See Figure 2.) This type of weigh system has been used extensively by one of us (JFW) in earlier work, and the chief results reported here were obtained with this system. The second weight detection system, still under development at the time of this writing, utilizes optical detection of displacements driven in a cantilevered beam to detect any weight fluctuations. (The optical/cantilevered beam technique was suggested by John Cramer.) Both weigh systems were enclosed in clear acrylic containers that were pumped down to vacua typically of 10 to 20 microns so as to eliminate several potential sources of spurious signals.

In earlier work of this sort, arrays of capacitors, rather than stacks of PZTs, were driven by a voltage signal at about half the mechanical resonance frequency of the load cell so that the mass fluctuation excited at the power frequency of the applied voltage signal would produce a periodic weight fluctuation that would be resonance amplified by mechanical properties of the load cell.^{4,5} Since the resonance frequency of the load cell was typically about 100 Hz, and the magnitude of the mass fluctuation scales with the square of the frequency of the voltage signal, only

minuscule weight fluctuations could be driven at the 100 Hz frequency, and mechanical resonance amplification and signal averaging were essential to see any effect at all. Nonetheless, this seemed the only path to




Figure 2: The device mounted on the U-80 (located in the steel shielding case) via a vibration isolator made from several layers of plastic and neoprene rubber. An accelerometer is affixed to the bottom of the aluminum stage that the isolator is bolted to. The stage is stabilized against lateral motion by tensioned fine steel wires.

detection of the mass fluctuation effect open at that time, for the presence and significance of the negative mass term in Equations (2) and (3) above was not then appreciated. Confined to the impulse engine term, which time-averages to zero, no other approach seems feasible.

In the present experiments, the cantilevered beam method of weight detection is problematical because of the rather large mass of the suspended device (more than 100 grams). In order to get sufficient sensitivity, the mechanical resonance frequency of the beam must be made rather low (less than ten Hz). This makes it difficult to detect a short, transient weight fluctuation. The situation with the U-80 load cell is better, as its mechanical resonance frequency is several tens of Hz – out of the range of low frequency acoustic and seismic noise sources. Moreover, this higher resonance frequency makes it possible to follow transient weight fluctuations with durations in the 100

to 200 millisecond range. This is especially important as, since the early work was done, the importance of the negative mass term in Equations (2) and (3) has come to light. Because the effect of this term is always to reduce the mass of the object in which it is excited (being negative definite), a stationary time-averaged mass shift arising from this effect can be produced by a voltage signal of much higher frequency than the mechanical resonance frequency of the load cell. Indeed, this effect can be optimized by driving the PZT stack at one of its mechanical resonance frequencies (typically about 75 kHz in devices like that in Figure 1). Since the time-average of this effect during an applied power interval will lead to a stationary weight reduction, one may simply observe any weight change directly should it be large enough. Impulse engine term effects, by themselves, do not lead to such stationary weight shifts since they time-average to zero.

Mass fluctuations arising from the negative mass term, in real circumstances, if present at all, may be much smaller than ideal considerations suggest. So, should one want to excite a mechanical oscillation of the weigh system near its resonant frequency to bring out a small effect, one need only pulse the high frequency voltage signal delivered to the PZT stack at some suitable low frequency. This procedure, however, proved unnecessary, as with reasonable signal averaging an effect of the sort sought could be obtained. Instead, the frequency of the applied voltage signal was swept through a predetermined range of values to allow for the fact that changing conditions in the PZT stack might cause the frequency of peak effect to change from test to test.

The experimental protocol actually followed was to take data during an interval of several seconds. During the first and last parts of this interval no appreciable voltage signal was applied to the device being weighed. For several seconds between these quiescent periods the voltage signal was applied to the device. By frequency modulation of the signal generator used, the frequency was swept from 70 to 78 kHz during this interval. This frequency interval was chosen because it includes the fundamental mechanical resonance frequency of the PZT stack where the effect sought was expected. Even though a surprisingly large weight fluctuation near the resonance was found, because of a rather poor signal to noise ratio in the weigh signal, we found it necessary to average the data for typically 50 to 100 or more of these data cycles together in order to extract a clear, statistically significant signal.

Aside from the weigh systems *per se*, other components of the apparatus used in this experiment, indicated in Figure 3, are: a power amplifier (a DBX 150 Watt linear amplifier), a 5-to-1 toroidal stepup and isolation transformer (TRANS), voltage and current

sense resistors in the high voltage secondary transformer circuit so that the voltage and current signals could be multiplied (4 QUAD MULT.) to get the instantaneous power delivered to the PZT stack, and a rectification and filtering circuit (POWER METER) so that the power could be read as a voltage in real time. Additional instrumentation allows measurement of acceleration of the PZT stack by the embedded accelerometer and the temperature of the device was recorded with the aforementioned thermistor.




Figure 3: A schematic diagram of the chief electronic circuits used in this experiment. The DAC and ADC are computer controlled.

While the chief parts of the associated circuitry and instrumentation are displayed schematically in Figure 3, several other signal detection features are not displayed. For example, a Picoscope AD212 was also used to monitor various signals at various times. Its software provides FFT power spectrum display capability that is quite helpful for analyzing harmonics present in those signals. Toward the end of the acquisition of data for this paper it became clear that more detailed knowledge of the thermal behavior of the device than that provided by the thermistor was desirable, so two types of non-contact infrared (IR) detection – one a simple thermo-couple device and the other an IR imaging camera – were introduced. A thermo-couple vacuum gauge was used to monitor the vacuum in the vacuum vessels. Operation of the system and data acquisition and processing is computer controlled via a Canetics PCDMA data acquisition board. Data acquisition and post-acquisition data processing is done with software written specifically for that task (adapted from software used in earlier experiments).

The core apparatus in use in the load cell version of this experiment is shown in Figure 2. The weight transducer is located in the 1 cm thick steel shielding case visible in the figure. The load cell vacuum chamber is a small box about 22 cm on a side 26 cm high made of 3 cm thick acrylic plastic (not shown in Figure 2). As in earlier realizations of

experiments using the U-80 load cell, *with sufficient averaging of data from individual cycles*, a mass (or weight) shift of a few milligrams could be resolved – assuming that the static calibration of the load cell with a one gram mass is applicable to signals registered in conditions where high frequency vibration is present in the PZT stack. This last assumption is open to question, for the manufacturer of the U-80 stipulates that the sensitivity of the Hall probe used changes unpredictably in the presence of significant vibration with frequencies above 1 kHz.

The obvious way to deal with this potential problem is to mount the PZT device on the weigh stage of the U-80 with high frequency vibration isolation. (Low frequency isolation is neither needed nor desirable. Indeed, hard coupling at low frequencies is required to make the weight measurements that are the objective of this experiment.) This was done in the earlier phase of this experiment by progressively introducing more and more vibration isolation components. Changes in the weight response of the U-80 disappeared before the isolator configuration seen in Figure 2 was implemented. That is, the isolator shown constitutes over-kill. Nonetheless, to be certain that appreciable high frequency vibration was not communicated to the U-80, an accelerometer (made with thin chips of PZT material) was affixed to the bottom of the weigh stage, and the weight signal was scrutinized with the Pico-scope run in power spectrum mode. Both the accelerometer signals and the power spectrum analysis of the weight signal confirmed that no significant high frequency vibration was present in the U-80 during the operation of the PZT device.

RESULTS

The chief result of this experiment is displayed in Figure 4. It shows the average of several hundred cycles of data taken in a variety of different circumstances. The data displayed is “net” data. While the actual data were being acquired, “null” data where the power to the device in each cycle was manually shut off as soon as the power was automatically activated. The average of the null data has been subtracted from the raw data average. In each cycle data was taken for a period of 14 “seconds” at a rate of 50 acquisitions per “second”. In fact one “second” was 0.65 actual seconds, the shift having been made by the adjustment of a

delay loop in the data acquisition program. Power was not applied to the device until four “seconds” into each cycle. When power was turned on, the frequency of the applied sinusoidal signal was swept from 70 kHz to 78 kHz during the ensuing six “seconds”. After the power was switched off, data continued to be taken for an additional four “seconds”. Three channels of data in addition to the weight sensor were acquired. One channel recorded the rectified and filtered (10 millisecond time-constant) power signal. This is displayed with the weight signal in panel A of Figure 4. Another channel recorded the rectified and filtered signal from the accelerometer embedded in the PZT stack; and yet another channel recorded the temperature measured by the thermistor epoxied to the aluminum ear bolted to the back of the brass disk on which the PZT stack was mounted. The data from these channels are displayed with the weight signal in panel B of Figure 4.

Simple inspection of Figure 4 reveals that a pronounced weight decrease occurs near 74 kHz. The magnitude of the decrease, about 60 ADC counts, corresponds to roughly 0.1 gram for a peak applied power of about 105 Watts. This is nearly the fluctuation expected on the basis of a naïve, ideal calculation based on Equation (3) above (within a factor of three or so). Note that the weight fluctuation is correlated to spikes in the power and accelerometer traces that signal the occurrence of resonant behavior in the PZT stack. Moreover, the temperature trace recorded by the thermistor shows an inflection in the heating rate that also correlates to the weight shift at 74 kHz.

The thermistor, of course, does not promptly measure the temperature of the PZT stack. Heat evolved in the PZT stack as power is applied is localized in the end of the stack where, by design, the maximum activity is induced – as can be seen in the far infrared (7.5 to 13.3 micron) image of the device




Figure 4: The weight fluctuation in the device shown in Figures 1 and 2 was swept with a sinusoidal signal. The weight is recorded in ADC counts, 66 counts scaling to 0.1 gram. The power trace shown in panel A peaks at 105 Watts. The corresponding thermistor (temperature) and accelerometer traces shown in panel B were not absolutely calibrated.

during power application presented in Figure 5 (taken with an Indigo Alpha Microbolometer camera). Prompt heating of the thermistor, evidently, is produced by vibration driven by the activated PZT stack. And the inflection in the thermistor trace shows that this vibration at the thermistor drops dramatically just before and during the onset of the power and accelerometer resonances.

This device and several others like it were actually designed to behave in this way, though they are so cranky that the preload on the PZT stack and the operating temperature must be "just so" to get them to work as intended. Indeed, the device used to obtain the data displayed showed such crankiness. When optimal operating conditions were not present, the inflection in the thermistor trace was markedly diminished. So thermistor trace inflection was used as a selection criterion for the inclusion of cycle data in the averaged result. (Roughly one cycle in four was rejected using the thermistor trace inflection criterion. The cycle rejection rate improved [decreased] with knowledge of the operation of the system during the course of data acquisition.)




Figure 5: A far infrared image of the device during a power cycle. As expected, the bulk of the energy is delivered to the end of the PZT stack near the aluminum cap. Note too the prompt heating of the thermistor and rubber pads. The temperature rise in the stack during a power cycle is about 5 degrees Celsius.

One may ask: Is the weight fluctuation correlated with the power, thermistor, and accelerometer events real, or just a statistical fluke? Inasmuch as the number of cycles averaged is about 250, and taking into consideration the weight trace away from the fluctuation, it seems quite plain that the correlated weight fluctuation is *most unlikely* to be a mere statistical fluke. More elaborate statistical analyses of the data are possible. For example, a formal error can be obtained by selective suppression of parts of the data

and computing the mean and standard deviation of the spread of the resulting weight fluctuations. This, however, doesn't really shed more light on the significance of the result contained in Figure 4. Given that a weight fluctuation of the sort expected is seen in Figure 4, the question then is: Is the observed weight fluctuation caused by a real Machian effect? Or is it the consequence of the operation of some spurious signal source? Keeping in mind that, "extraordinary claims require extraordinary evidence," we turn now to the tests we have conducted to see if the signal in Figure 4 can be claimed to be real.

SPURIOUS SIGNALS AND CHECK PROTOCOLS

If we assume that the weight fluctuation signal in Figure 4 is *not* due to the predicted Machian effect, then, evidently, it must be a consequence of either the electromagnetic signals present during activation of the device or the mechanical vibration induced by those signals, for no other possibilities worth considering exist. Of these two classes of potential spurious signals, those due to electromagnetic effects are much easier to deal with. All one has to do is recreate the electromagnetic signals without driving the electro-mechanical response in the PZT stack to test for their effects. This is easily done by shorting out the PZT stack while applying the same voltage signal to another, similar device wired into the circuit far from the weigh system. This way all of the currents and voltages in the device can be emulated, but the electro-mechanical response that purportedly produces the effect is not excited. Figure 6 displays the results of




Figure 6: The averages of the current and voltage emulation weight traces. Note the absence of any weight fluctuation correlated with the peak of the power resonance.

this test. Very obviously, no weight fluctuation of the sort seen in Figure 4 is present, so the effect seen cannot be attributed to this source. A small systematic

effect is seen in Figure 6 however. It is caused by the heating of the power leads to the device. (This heating effect is not seen in Figure 4 because data for two lead configurations were averaged together to cancel it out.)

The fact that the current and voltage emulation test does not *exactly* replicate all of the conditions present in the actual data runs makes possible the elimination of another potential source of spurious signals. When the device is actually run, one of the power leads is "hot" while the other is at ground potential. In emulation runs either both leads are hot, or both are not. A moment's reflection should convince one that the "both not" configuration more nearly approximates actual running conditions. Nonetheless, emulations were done in both configurations, and averaged together in Figure 6 because there was no significant difference in the results for the two configurations. In the "hot" configuration, however, since the entire device is (periodically) driven to high voltage, dielectrics in proximity to the device (the vacuum case in particular) will be polarized by the electric field. Since the field may be expected to have large gradients, these polarization charges might interact with the field, and thus the device. Roughly half of the data averaged in Figure 6 being for the "hot" configuration, it shows that no such effects are present.

Other tests of electromagnetic effects were carried out; notably, static magnetic fields of a few Gauss were applied to the system during its operation. No detectable effect was found in any of these tests that would lead one to suspect the effect seen in Figure 6 to be spurious. Mechanical vibration, the other leading candidate for spurious effects, was dealt with chiefly by aggressive vibration suppression and careful monitoring of vibration in the U-80 transducer. In addition, a test involving a unique signature of the negative mass effect was carried out. Note that the negative mass term in Equation (3) above depends on the square of the time derivative of the matter density. Thus, it also depends on the square of the energy density, and ultimately the square of the applied power (and their time derivatives). Normal effects driven by mechanical vibrations, however, may be expected to scale linearly with the applied power.

Although simple in principle, this test is difficult to carry out in practice for several reasons, chiefly secular evolutionary changes in the system during the extensive data runs needed to get even plausible results. Nevertheless, we present results of such a test in Figure 7. They cannot be claimed to be

compelling, especially given the noise in the low power weight trace. Moreover, the differences in the power curves for the two power levels casts further doubt on simple comparisons of the two responses. But notwithstanding these caveats, the predicted power scaling appears to be present.

So far we have considered voltage and current emulation, the induction of electric dipoles in surrounding media, static magnetic fields, the vibration measurements of the mounting stage accelerometer, and the power scaling behavior of the system. And all results are consistent with a real Machian mass fluctuation as the cause of the signals in Figures 4 and 7. It would seem that but a few potential sources of spurious results remain to be considered. Perhaps the most important of these are corona and sonic wind. Both of these depend strongly on the density of the air in which the device is run. So they may be explored by noting any change in the behavior of the system as the pressure in the vacuum vessel is changed. To insure that these effects were not responsible for the signals seen, the device was run at atmospheric pressure, 150 microns, and the normal operating pressure of roughly




Figure 7: The weight fluctuations recorded for two different power levels. Note that whereas the ratio of the powers is 1.27, the ratio of the weight fluctuations is about 1.6, suggesting that the weight fluctuation scales with the square of the applied power.

15 microns. No noticeable change in behavior with pressure was observed, allowing us to set aside these possible sources of signals. (Nonetheless, the device was operated at the pump limit of about 15 microns except during these tests.)

In light of the results of the tests we report here, perhaps the most important test remaining is the replication of the main result using a completely different weigh system – in particular, a system like the cantilevered beam mentioned above. Time has not permitted us to complete such a test by the time of this writing. But we hope to have one done shortly.

CONCLUSION:

As noted above, the Saganism that extraordinary claims require extraordinary evidence is certainly true. Extraordinarity, however, like beauty, is in some measure in the eye of the beholder. So whether the evidence we present here rises to that level is a matter of individual judgment. Our judgment is that the evidence suggests that a real effect is present; although, being quite cranky, it is far more difficult to produce and manipulate than we would like. Partly, perhaps, these difficulties can be attributed to the very modest effort involved. Greater effort, we think, is warranted.

But where, beyond further confirmation and exploration of the effects reported here, does one go from here? Well, from the point of view of rapid spacetime transport two issues seem to be the most important. First, can the effects reported here be scaled to useful levels? A mass fluctuation of a tenth of a gram or so is only of laboratory interest. It is worth noting that the negative mass term effect, everything else held constant, scales with the square of the frequency and with the square of the power. So, in principle, very large mass fluctuation effects should be producible with only relatively modest power levels. We hasten to add that there are subtleties to the production of this effect that are not yet fully explored. For example, generation of the effect seems to depend quite critically on the production of higher harmonics in the PZT stacks, for sinking a lot of power into a stack at a mechanical resonance of the stack is not, itself, sufficient to yield an effect. The details of the processes that lead to large mass fluctuations need to be studied and *understood* in some detail before serious engineering is undertaken.

Second, from the point of view of propellantless propulsion, the creation of impulse engines requires that objects with induced mass fluctuations be acted upon by another periodic force to extract a stationary force. In the torsion pendulum experiments that preceded those sketched here, two of us (JFW and TLM) were only able to produce minuscule forces many orders of magnitude smaller than those expected on the basis of naive calculations. Likely this can in part be attributed to subtleties involving the yet to be explored details of the production of mass fluctuations in real non-linear systems. These details will have to be sorted out before the dream of propellantless propulsion can be fully realized.

In this regard we note that it may be worth exploring systems other than simple stacks of PZT crystals driven with complex voltage waveforms (like those already investigated by TLM and JFW). For instance, one might expect to see impulse engine effects in the system described by Corum, *et al.* where a sinusoidal voltage is applied to a circuit with an

inductor and capacitor in series.³ The inductor is arranged so that the periodic magnetic field it generates permeates the dielectric between the plates of the capacitor, orthogonal to the electric field therein. Taken as a strictly electromagnetic system, the conservation principles mentioned at the outset of this paper require that any forces produced in the dielectric be balanced by equal and opposite forces elsewhere in the circuit. So there is no reason to expect to see strictly electromagnetic propulsive forces in this type of system.

But note that if Machian mass fluctuations actually occur, then the acceleration of the charge carriers that comprise the displacement current in the dielectric in the capacitor, driven by the periodic electric field, will induce a mass fluctuation at the power frequency of the voltage frequency – that is, at twice the frequency of the applied voltage signal. Now consider the action of the magnetic field. The motion of the displacement current charge carriers will interact with the period magnetic field to yield a force on the charge carriers via the magnetic part of the Lorentz force: $(q/c)(v \times B)$. Since the velocity of the charge carriers, v , and the magnetic field, B , are both periodic with a frequency equal to the applied voltage frequency, their cross product will yield a force with twice this frequency – that is, the frequency of the mass fluctuation putatively being driven by the action of the electric field in the dielectric. Should the relative phase of the mass fluctuation and the magnetic part of the Lorentz force be auspicious, a stationary force should result.

Such a force, if present, is a result of the mass fluctuation that arises from the inertial coupling of the constituents of the dielectric to the rest of the universe. It is not due to a local violation of the conservation of momentum in a purely electrodynamical system. It is worth remarking that a stationary force in a system of this sort may be expected even if the “substance” between the plates of the capacitor is the vacuum. If the charged particle pair production in the vacuum of quantum lore actually takes place, the pairs should experience the same effects as material dielectric media. So exploration of this sort of arrangement of circuit elements has scientific value (as a test of the “polarizable vacuum” conjecture), as well as potential technological implications.

Finally, we note that the nearly 0.1 gm mass shift that seems to have taken place in the PZT stack approaches a percent of the total mass of the PZT material (about 40 gm), and this with an applied power on the order of 100 Watts. If this effect is real, and should no physical mechanism operate that enjoins true negative total mass states, then it seems that we may be able to actually generate, however fleetingly, truly exotic matter. Should this prove possible, the most extreme sorts of rapid spacetime transport may be

feasible. Experiments designed to probe these possibilities, however, should be undertaken with some care.

ACKNOWLEDGEMENT:

The work now in progress at CSUF has profited significantly from important contributions of others. They have contributed in several veins. Kirk Goodall helped with the construction of parts of the apparatus, participated in many discussions about the experiment, and did some of the testing of various systems. Probing questions and comments from John Cramer, Michael Dornheim, Edward Harris, and Graham O'Neil, among others, have made us think carefully about several important aspects of Machian mass fluctuations and their application to rapid space-time transport. And the careful critiques of colleagues at CSUF, especially Ronald Crowley and Stephen Goode have been essential. Indeed, those critiques are largely responsible for the appreciation of the significance of the negative mass term in Equations (1) and (2) that underlies the work reported here.

REFERENCES:

1. Woodward, J.F. and Mahood, T.L. (1999), "What is the Cause of Inertia?", *Foundations of Physics* **29**, 899 – 930.
2. Woodward, J.F. (2001), "Gravity, Inertia, and Quantum Vacuum Zero Point Fields," *Foundations of Physics* **31**, 819 – 835.
3. Corum, J.F., Dering, J.P., Pesavento, P., and Donne, A. (1999), "EM Stress-Tensor Space Drive," *Space Technology and Applications International Forum-1999* (ed. M.S. El-Genk, AIP Press, 1999), pp. 1027-1032.
4. Woodward, J.F. (1990), "A New Experimental Approach to Mach's Principle and Relativistic Gravitation," *Foundations of Physics Letters* **3**, 497 – 506.
5. Woodward, J.F. (1991), "Measurements of a Machian Transient Mass Fluctuation," *Foundations of Physics Letters* **4**, 407 – 423.
6. Woodward, J.F. (1995), "Making the Universe Safe for Historians: Time Travel and the Laws of Physics," *Foundations of Physics Letters* **8**, 1 – 39.
7. Mahood, T.L. (1999), "A Torsion Pendulum Investigation of Transient Machian Effects," California State University, Fullerton masters thesis, available at <http://www.serve.com/mahood/thesis.pdf>.
8. Sciama, D. (1953), "On the Origin of Inertia," *Monthly Notices of the Royal Astronomical Society* **113**, 34 – 42.
9. Sciama, D.W. (1964), "The Physical Structure of General Relativity," *Reviews of Modern Physics* **36**, 463 – 469.
10. Woodward, J.F. (1992), "A Stationary Apparent Weight Shift from a Transient Machian Mass Fluctuation," *Foundations of Physics Letters* **5**, 425 – 442.
11. Woodward, J.F. (1996), "A Laboratory Test of Mach's Principle and Strong-Field Relativistic Gravity," *Foundations of Physics Letters* **9**, 247 – 293.
12. Puthoff, H. E. (1999), "Polarizable-Vacuum representation of general relativity," available at <http://xxx.lanl.gov/abs/gr-qc/9909037>.
13. Woodward, J.F. (2000), "Mass Fluctuations, Stationary Forces, and Propellantless Propulsion," *Space Technology and Applications International Forum 2000* (American Institute of Physics/Springer Verlag, New York, 2000), pp. 1018 – 1025.
14. Woodward, J.F. and Mahood, T.L. (2000), "Mach's Principle, Mass Fluctuations, and Rapid Spacetime Transport," unpublished conference paper available at: <http://chaos.fullerton.edu/jimw/staif2000.pdf>.



A01-34539

AIAA 2001-3910

Global Monopoles and the Bondi-Forward Mechanism

**C. Van Den Broeck
The Pennsylvania State University
University Park, PA**

**37th AIAA/ASME/SAE/ASEE
Joint Propulsion Conference and Exhibit
8-11 July 2001
Salt Lake City, Utah**

For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

GLOBAL MONOPOLES AND THE BONDI-FORWARD MECHANISM

Chris Van Den Broeck

*Department of Physics, The Pennsylvania State University, 104 Davey Laboratory, University Park, PA 16802, USA
e-mail: vdbroeck@gravity.phys.psu.edu*

Abstract. Based on work by Bondi, Forward considered the possibility of using putative negative masses for a propellantless propulsion mechanism. Recent advances in the study of black holes in the presence of fields may provide an explicit mechanism to mimick the effects of negative mass without the need to invoke exotic quantum effects, or to postulate the existence of matter with negative mass in the classical domain. We evaluate the feasibility of using such objects in a Bondi-Forward type propulsion mechanism.

1. Introduction

Bondi¹ was the first to consider the effects of putative negative masses within classical general relativity. As pointed out by Forward², negative mass could be used to create a propellantless propulsion mechanism.

The underlying principle can most easily be understood in Newtonian mechanics. Consider a point particle 1 with gravitational mass $m > 0$ and a particle 2 with mass $-m$, both moving along the x -axis of a Cartesian coordinate system. Let \bar{e}_x be the unit vector pointing in the positive x -direction, and suppose that initially $x_2 < x_1$, where x_1 and x_2 are the coordinates of particle 1 and particle 2, respectively. Particle 2 exerts a force on particle 1 equal to

$$\bar{F}_1 = -\frac{G(-m)m}{(x_1 - x_2)^2} \bar{e}_x = \frac{Gm^2}{(x_1 - x_2)^2} \bar{e}_x, \quad (1)$$

while particle 1 exerts a force on particle 2 given by

$$\bar{F}_2 = \frac{Gm(-m)}{(x_1 - x_2)^2} \bar{e}_x = -\frac{Gm^2}{(x_1 - x_2)^2} \bar{e}_x. \quad (2)$$

Note that the forces are equal in magnitude but have opposite direction, in accordance with Newton's third law. To get the accelerations, we now have to divide the forces by the respective *inertial* masses. We will assume these to be equal to the gravitational masses, including the sign. To fully understand why this is a reasonable assumption, one would have to take recourse to general relativity¹; we will also come back to this issue in Section 3. Dividing \bar{F}_1 by m and \bar{F}_2 by $-m$, we find the accelerations for particle 1 and particle 2 to be

$$\bar{a}_1 = \bar{a}_2 = \frac{Gm}{(x_1 - x_2)^2} \bar{e}_x. \quad (3)$$

Copyright (c) 2001 The American Institute of Aeronautics and Astronautics Inc. All rights reserved.

This means that both particles will undergo an acceleration in the same direction, and this acceleration will persist until the relativistic regime is reached. (Note that in this special case the magnitudes of the accelerations are also equal to each other; this is just because we chose the absolute values of the masses to be the same.) Again taking into account the signs of the masses, it will be clear that both the (Newtonian) total momentum and total kinetic energy are conserved. (In fact, in this case both are identically zero for all times.)

Negative masses have never been observed in Nature, although it is generally assumed that quantum field theory effects can give rise to negative energy densities (a very readable introduction can be found in Visser's book³). In this paper, we will describe a possible mechanism to mimick the effects of negative mass, starting from purely classical physics. Our tools will be general relativity, together with the classical theory of the weak and strong nuclear interactions.

As it is well-known, black holes can carry an electric charge, and when a charged particle is moving in such a spacetime, it will feel the effect of both the charge Q of the hole and its mass M . Since energy is stored in the electric field generated by the charge Q , it will contribute to the curvature of spacetime in the vicinity of the hole. Because of the fact that in general relativity, the gravitational force is a result of the curvature of spacetime, this contribution will make it seem as if the hole has a mass different from M . Consequently, even the motion of an uncharged particle in such a spacetime will be different from what it would have been if the hole had only carried a mass M , and no charge. In particular, the particle will appear to be moving in the gravitational field of a mass that is smaller than M .

It is a generic feature of general relativity that objects appear to have a mass that is smaller than their actual mass content (see e.g. Wald⁴). Consider the example of a spherical, non-rotating star. According to Birkhoff's theorem, the geometry outside the star is that of Schwarzschild spacetime, with some characteristic mass \tilde{M} . This is the mass outside observers will perceive the star to have, for example by releasing test particles and observing their motion. As it turns out, \tilde{M} will always be slightly smaller than the 'proper' mass content M of the star (i.e., the mass that went into the star when it formed). If we define $\Delta M = M - \tilde{M}$, then $E_B = \Delta M c^2$ may be interpreted as the gravitational binding energy.

It needs to be stressed that the latter is merely a

heuristic statement, since gravitational binding energy has no general definition. Even so, one could ask the following question: Can a self-gravitating object be so strongly bound that its apparent mass is negative (i.e., can the contribution to the mass due to binding energy be larger than the proper mass content)? Preliminary investigations seem to point towards a positive answer. The purpose of this paper is to describe under what circumstances this could come about, and to evaluate its relevance for the Bondi–Forward (BF) propulsion mechanism.

The paper is structured as follows. In the next Section, we will introduce solutions to Einstein gravity in the presence of additional, non-gravitational fields, which for all practical purposes have a negative gravitational mass, even though the energy stored in the fields, as well as any proper mass that may be present, are positive. In Section 3, we first of all evaluate to what extent the objects described may be considered realistic, or physically acceptable, and we consider the problems that may be expected when incorporating them in BF propulsion. Conclusions are presented in Section 4. Although we will assume that the reader has some knowledge of basic relativistic physics, we will review the theory of the Higgs field and its coupling to gravity in the Appendix, as it will be the basis for the more promising class of objects discussed in Section 3.

Throughout this note we will use units such that $G = c = \hbar = 1$, unless stated otherwise. The signature of spacetime metrics is chosen to be $(-, +, +, +)$.

2. Black holes with negative mass

The fact that in the presence of fields, black holes can appear to have negative mass has been known for quite some time; it is apparent in the solutions discovered by Barriola and Vilenkin⁵. Recently, two classes of black holes with negative effective mass were studied in detail. In what follows, we will talk about proper mass and proper charge; for definiteness, let us first define these rigorously. The spacetimes we will consider all have a metric of the form

$$ds^2 = -\sigma^2 N dt^2 + N^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (4)$$

where σ and N are functions of r . Define $u(r) = r^2 N(r)$ and expand this function near $r = 0$:

$$u = Q^2 - 2Mr + \dots \quad (5)$$

By definition, Q is the proper charge and M the proper mass. This definition is motivated by the fact that in the case of a charged black hole, Q and M are indeed the amounts of charge and mass that end up in the hole as it is formed. These should be distinguished from the effective charge and mass, which are the values measured by an outside observer, e.g. using test particles.

So far we have only talked about black holes carrying the electric charge associated to electromagnetism. However, holes can also have proper charges associated to the weak and strong nuclear forces. Just like electromagnetism, these interactions are described by gauge theories, which are characterized by some internal group of invariances, or symmetries. (In the case of electromagnetism, this is the invariance of the electric and magnetic fields when adding to the vector potential the gradient of an arbitrary smooth function.) The symmetry group inherent in electromagnetism is called $U(1)$, and those of the weak and strong interactions are $SU(2)$ and $SU(3)$, respectively. The weak interaction is somewhat special, in that its symmetry is broken in Nature, presumably by interaction with the yet to be discovered Higgs field. For an elementary introduction into the physics involved, we refer the reader to the textbook by Aitchison and Hey⁶.

Zotov⁷ studied the appearance of black holes with negative effective mass in the context of unbroken $SU(2)$ gauge theory, called Yang–Mills theory. Although this model is not realized in Nature, it can be considered as a useful ‘toy model’ for the full $SU(3)$ theory describing the strong interaction. It is generally believed that the qualitative features of solutions of Yang–Mills theory coupled to gravity will usually carry over to $SU(3)$. Zotov’s black holes have a proper mass, and also a proper charge. The latter is not the familiar charge of electromagnetism, but its $SU(2)$ equivalent. Numerical results suggest that Yang–Mills theory admits black hole like solutions with negative effective mass but positive proper mass. However, the solutions studied by Zotov do not have an event horizon; their singularities are naked. Although a proof is lacking, it is generally assumed that this cannot happen in Nature (this is the famous *Cosmic Censorship Conjecture*⁸), so we will not discuss these solutions any further.

Instead, we turn to the slightly earlier work of Nuemann and Sudarsky⁹. They studied black hole solutions with an event horizon, arising from a model with three scalar fields having a broken global $O(3)$ symmetry. Here ‘global’ merely means that, unlike gauge symmetries, the transformations under which the theory is invariant are not allowed to depend on the position in spacetime. The broken $O(3)$ theory should be a good model for the Higgs interaction. It is assumed that there exists more than one Higgs. The way they interact among themselves will involve a global symmetry, which needs to be broken itself in order to break the $SU(2)$ symmetry of the weak interaction. The breaking occurs when the Higgs particles form a condensate, meaning that the Higgs fields acquire a non-zero vacuum expectation value.

A peculiar feature of the solutions presented by Nuemann and Sudarsky is that they carry neither proper mass nor proper charge. Instead, they are ‘topological defects’, usually called *global monopoles*. A useful picture is that of a field which has been tied into a knot.

The energy stored in such a ‘knotted’ field can give rise to a spacetime geometry which is similar to that of a black hole with an event horizon. (Note that the term ‘topological defect’ refers to the gauge field itself, *not* the topology of the spacetime it gives rise to.) Henceforth, we will refer to the particular solutions found by Nucamendi and Sudarsky as *NS monopoles*.

In general, the equations describing the interaction between the fields and gravity in the model of Nucamendi and Sudarsky are too complex to solve analytically. As a consequence, the spacetime metrics for black hole like solutions can usually not be written in closed form. However, at large distances, NS monopoles have a metric that looks like

$$ds^2 \simeq - \left(1 - \frac{2\tilde{M}}{r} \right) dt^2 + \left(1 - \frac{2\tilde{M}}{r} \right)^{-1} dr^2 + (1 - \alpha) r^2 (d\theta^2 + \sin^2\theta d\phi^2), \quad (6)$$

where $\alpha = 8\pi\eta^2$, with η the magnitude of the Higgs field condensate. The equations of motion for uncharged test particles with positive mass moving in the vicinity of the monopole are given by the geodesic equation. A particle that is initially at rest at sufficiently large distances will be subject to a radial proper acceleration

$$a_r \simeq \frac{-\tilde{M}}{r^2 - 2\tilde{M}r}. \quad (7)$$

As numerical results show, it is possible to have solutions with $\tilde{M} < 0$, so that $a_r > 0$: The central object is repelling the particle. For all practical purposes, the global monopole behaves as an object with negative mass.

It is worth pointing out that although the monopole behaves as if it carries negative mass/energy, all of the energy conditions of general relativity are satisfied because of the way the fields couple to gravity¹⁰. Therefore, no observer will ever measure negative energy density in this spacetime; the monopole indeed only *mimicks* the effect.

Note that the metric (6) is not asymptotically flat: In the limit $r \rightarrow \infty$, one does not retrieve the usual uncurved (or Minkowski) spacetime. Rather, the geometry is ‘asymptotically flat up to a deficit angle’. This means that if we were to measure the surface area of a sphere with very large radius R centered on the black hole, we would not find the usual value of $4\pi R^2$, but rather $4\pi(1 - \alpha)R^2$. We will come back to this important fact in the next section.

3. Global monopoles and the BF mechanism

Although the NS monopoles satisfy most of the criteria to be physically acceptable solutions of the broken $O(3)$ theory coupled to gravity, it is not known how they

could be created in Nature. The relevant literature is very recent, and it deals only with static solutions, so that at this time it would be inappropriate to speculate on how global monopoles with negative effective mass might come about. This makes it impossible to assess how much energy is needed to generate these objects. As we mentioned, the proper mass is zero; all energy is stored in the fields. In general relativity there is no generic scheme to compute the energy stored in a particular fields configuration, especially for solutions that are not asymptotically flat.

The failure of the metric (6) to be asymptotically flat is a possibly problematic feature from a physical point of view. The deficit angle will create tidal stresses. As we mentioned in the previous section, the magnitude of α is determined by the magnitude of the Higgs field condensate. The magnitude of the stresses on a compact object of fixed size will depend on distance; even though the metric is not asymptotically flat globally, it still approaches flatness locally. A good analogy is to consider the surface of a cone. The spheres of radius R we talked about in the previous section are now circles on the cone in a plane perpendicular to the cone’s axis. R should be interpreted as the distance of a circle from the apex of the cone, so that their circumference is $2\pi(1 - \beta)R$, with $2\pi\beta$ some deficit angle. But now consider a disk of radius ρ stuck on the surface of the cone. If the disk is close to the apex, its surface area will be smaller than $\pi\rho^2$. However, as one slides the disk away from the apex while keeping ρ fixed, the curvature of the cone’s surface becomes less noticeable and the area of the disk will tend to $\pi\rho^2$ asymptotically. Similarly, in the spacetime (6), a spherical object with radius ρ will have a volume smaller than $4\pi\rho^3/3$ when close to the central singularity, but as it is moved away from it while keeping ρ fixed, the volume will tend to that value.

The magnitude of the Higgs condensate is expected to be $\eta \simeq 250 \text{ MeV/m}^3$; in units where $G = c = \hbar = 1$, this becomes $\eta \simeq 5 \times 10^{-126}$, so that the deficit angle $4\pi\alpha = 32\pi^2\eta^2$ will be of no consequence in the regime where the approximation (6) is valid.

NS monopoles have two characteristic length scales associated to them: The radius of the event horizon r_h , and the monopole’s ‘core radius’ r_c . The latter is completely fixed by the mass the Higgs bosons themselves acquire due to symmetry breaking. Different NS monopole solutions are characterized by different values of r_h . Negative effective masses occur when $r_h \ll r_c$, so that for these solutions the metric (6) is a good approximation for $r \gg r_c$. In units where $G = c = \hbar = 1$, one has $r_c = 1/(\sqrt{2}m_H)$. Although the Higgs mass is as yet unknown, it is generally assumed to be approximately $m_H \simeq 200 \text{ GeV}$. In SI units, this gives $r_c \sim 10^{-35} \text{ m}$. For all practical purposes, the monopole can be considered as a point particle.

The smallness of the core radius raises an important question. So far, we have been discussing purely classical

field theory, both on the gravitational and on the Higgs side. However, r_c is comparable to the Planck length. This means that realistic NS monopoles are not only essentially quantum mechanical objects; quantum gravity effects can be expected to influence their structure in a major way. As yet, no quantum gravity theory has been verified either experimentally or observationally, so that it is impossible to say what these effects will do to the monopole. However, although the Higgs mass is a parameter in the Standard Model of elementary particle physics, it is not generally considered to be a fixed constant of Nature. This is because it is not only determined by the coupling strength of the Higgs self-interaction, but also by the magnitude of the Higgs condensate. The latter is expected to be a consequence of the particular conditions that existed shortly after the Big Bang¹¹. It is not unimaginable that its value could be changed locally, although any statement as to how this could be done would again be pure speculation. For now, I will confine myself to saying that it is justifiable to continue the discussion on the basis of the qualitative results from the classical theory.

Without going through the full numerical computation for the spacetime geometry, from looking at the evolution equations in the paper by Nucamendi and Sudarsky⁹ one can immediately tell that if $\alpha \simeq 0$, the effective mass will also be exceedingly small. Nevertheless, one can think of schemes to use NS monopoles for BF propulsion. For instance, consider a very large number of them contained in a spherical shell of positive mass. Using Newtonian arguments again, the spherical shell will not influence the effectively negative mass inside; as it is well-known, such a shell acts like the gravitational equivalent of a Faraday cage. On the other hand, negative mass particles will repel both each other and the surrounding shell. One can therefore conceive of a thin, spherical, positive mass shell filled homogeneously with negative mass. If the amount of negative mass inside is much larger in absolute value than the mass of the shell, the net gravitational field outside will be that of a negative point mass. Still using Newtonian arguments, for an amount of negative mass in the order of, say, a large asteroid mass ($M \sim -10^{20}$ kg), the outward pressure on the spherical shell will be comparable to the inward pressure felt by the matter in a positive mass asteroid (this is of course assuming the dimensions are comparable). Close to the shell, the repulsive gravitational force will be the same in magnitude as the (attractive) force caused by an asteroid with the same (positive) mass and dimensions. This will then be the force acting on a positive 'payload' mass placed close to the shell.

A possible implementation of the BF mechanism would consist of a container of NS monopoles together with a positive payload mass, which would act as the positive mass particle in the system. Now a new question arises: Although we know that the container will repel the pay-

load, can we be sure that the payload will attract the container? Phrased differently, is the inertial mass of NS monopoles equal to the negative effective mass? Recall from the Introduction that it was precisely the fact that we divided the force on the particle with negative gravitational mass by $-m$ that led to an acceleration in the same direction as that of the positive mass particle. The inherent assumption was the equality of gravitational and inertial mass. So far, we have tacitly assumed that this equality holds for NS monopoles. Are we justified in doing so?

When computing the inertial mass of e.g. a proton, the binding energy of its quarks due to the strong nuclear interaction needs to be taken into account to arrive at the correct result. Similarly, if the negative effective mass of NS monopoles is to be understood as a consequence of gravitational binding energy (cf. the example of the spherical non-rotating star), then one expects a positive answer.

Since gravitational binding energy is but a heuristic concept, a more fruitful way of looking at the problem is to consider the metric caused by an isolated payload mass. Supposing the payload mass is spherically symmetric, the metric of the geometry outside will be Schwarzschild. If the negative mass container is sufficiently far away, it can be considered as a point particle. Alternatively, one could look at the motion of just a single NS monopole, which for all practical purposes should behave as a test particle in the technical sense: A point particle with negligible mass. It is reasonable to assume that a single monopole will move on a geodesic, and that this will also be approximately true for a container of macroscopic extent. Since the geodesic equation is insensitive to the mass of the point particle of which it describes the motion, in that case a monopole, and a negative mass container, would certainly be attracted by the payload.

The geodesic equation can be derived from Einstein's equations by considering the motion of an extended body in curved spacetime and taking the limit where its spatial extent goes to zero. The derivation is subtle, as was made clear most recently by the work of Mino, Sasaki and Tanaka¹², and Quinn and Wald¹³. In the case of a small and light NS monopole superposed on a curved background, the analysis will be extremely complicated, partly because of the internal structure of a monopole, and especially because of the fact that the spacetime geometry is not asymptotically flat. A direct proof that NS monopoles will move on geodesics can be expected to be very difficult.

Because of its importance for the BF mechanism, the issue of inertia calls for further investigation. To evaluate the inertial mass rigorously, one needs a detailed study of the behavior of NS monopoles in a dynamical situation. One possibility would be to construct a binary solution of two NS monopoles connected by a

'strut'. Such constructions have been carried out for binary black hole solutions (a recent reference is Letelier and Oliveira¹⁴). If NS monopoles have negative inertial mass as well as negative effective mass, what we would expect for a monopole pair is that the connecting strut would tend to be stretched. Alternatively, one could consider perturbations of the NS monopole metric with non-zero momentum and study the recoil effects, as has been done by Andrade and Price¹⁵ for ordinary black holes.

Assuming that the system composed of the negative mass container and the payload mass can not go superluminal, it is easy to guess what will happen as it approaches relativistic speeds. The neutral positive matter can only radiate gravitationally. Since the Higgs interaction is much stronger than the gravitational one, the NS monopoles inside the shell will start radiating 'Higgs field waves' before the positive matter loses energy through gravitational wave emission. The Higgs radiation will carry off positive energy: The positive energy stored in the fields composing the monopoles. The strength of those fields then diminishes, and the effective negative mass becomes less negative even though positive energy is being radiated away. We reiterate that although the monopoles generate a repulsive gravitational force indistinguishable from that of negative mass, the true energy density is positive everywhere.

4. Summary and conclusions

We have described a possible implementation of the Bondi-Forward propulsion mechanism on the basis of global monopoles arising in a spontaneously broken $O(3)$ Higgs model coupled to gravity. Clearly, there are several open problems, each of them crucial to the relevance of NS monopoles in the context of the BF scheme.

The typical length scale of a monopole with negative effective mass is the core radius, which is determined by the Higgs mass and turns out to be of the order of the Planck scale. In that case, the classical description is inappropriate, and a description in terms of quantum gravity is called for. Since no quantum gravity theory has ever been tested, it is impossible to say at this point how this will effect the properties of the monopole. However, the Higgs mass depends both on the coupling strength of the Higgs self-interaction and the magnitude of the condensate. Both are parameters in the Higgs potential (see the Appendix), and standard lore¹¹ has it that this potential is dynamically generated by whatever physics goes on at very high energies. Consequently, it is possible that these parameters can be changed in a physical process, but the theoretical framework to even conceive of a mechanism to bring this about is not yet in place.

So far, the study of solutions to the theory of the Higgs coupled to gravity has been confined to static ones. As a result, no physical mechanism is known in which NS

monopoles might be created. Since these objects do not have a proper mass or charge, this makes it hard to even estimate how much energy would be involved.

The most pressing problem is the question whether NS monopoles with negative effective mass also have negative inertial mass. One would expect a positive answer, but this should not be taken for granted. We have indicated a number of ways in which it could be tested computationally. It seems unrealistic to hope for a direct proof that the motion of NS monopoles is approximately geodesic, but we have proposed other schemes which have already been worked out for less exotic objects.

It will be clear that a full evaluation of the possibility of using NS monopoles to implement the BF mechanism would require a protracted effort on the theory side. Even then, it seems unlikely that a definite statement could be made on the basis of what we currently know about high energy physics. But at least there is some hope; even though negative mass presumably does not exist in its 'pure' form, Nature may have given us a loophole.

Appendix

It is generally assumed that the breaking of the $SU(2)$ gauge symmetry of the electroweak interaction is a consequence of the breaking of a *global* symmetry associated with the dynamics of the Higgs fields. Nucamendi and Sudarsky considered a theory with three Higgs fields ϕ^a , $a = 1, 2, 3$, interacting with each other according to a potential

$$V(\phi^a) = V(\phi^1, \phi^2, \phi^3) = \frac{\lambda}{4}(\phi^2 - \eta^2)^2, \quad (8)$$

where ϕ^2 is shorthand for $\sum_{a=1}^3 \phi^a \phi^a$. Note that if one would set e.g. ϕ^3 to zero and consider only ϕ^1 and ϕ^2 , a diagram of the potential versus ϕ^1 , ϕ^2 would resemble a Mexican hat. Indeed, $V(\phi^a)$ then has a local maximum at the origin of the (ϕ^1, ϕ^2) -plane given by $V(0, 0, 0) = (\lambda/4)\eta^4$ (the top of the 'hat'). As one moves away from the origin, the potential decreases until the valley defined by $\phi^2 = \eta^2$ is reached, where $V = 0$ (the rim of the hat). Moving still further away from the origin, V increases without bound.

What cosmologists generally expect to have happened during the early phases of the Universe can be very roughly described as follows. Shortly after the Big Bang, when the temperature of the Universe was still very high, η was equal to zero, so that the potential had a unique minimum at the origin and increased without bound as one moved away from the origin. The equilibrium value for the classical fields ϕ^a was then $\phi^a = 0$ for $a = 1, 2, 3$, since the potential is then at its minimum, $V = 0$. As the Universe cooled down, η increased and the potential formed a 'bump' near the origin, leading to the Mexican hat shape. The values $\phi^a = 0$ now corresponded to an

unstable equilibrium point: The top of the ‘hat’. Consequently, the fields ϕ^a acquired non-zero values throughout spacetime, namely values that minimized the new potential. But note that for $\eta \neq 0$, (8) does not have a unique minimum; all values of ϕ^a for which $\phi^2 = \eta^2$ are a minimum of the potential. Which minimum the fields ended up in was determined by the small perturbations causing the field configuration to be pushed away from the unstable equilibrium point, and may be considered arbitrary for all practical purposes. The situation is comparable to that of a pencil balanced on its tip: The pencil could fall in any direction. This process is called *spontaneous symmetry breaking*. The result is that the classical values of the Higgs fields are no longer zero; in the jargon one says that the particles associated to the fields have formed a *condensate* ϕ^a such that $\phi^2 = \eta^2$. η is the magnitude of the condensate; presumably it is determined by the specific conditions existing in the early Universe, together with the dynamics of the as yet unknown high-energy physics driving the symmetry breaking. The terminology stems from a very analogous process in condensed matter physics, the Meissner effect, which is responsible for superfluidity and superconductivity.

In a curved spacetime, the field equations governing the classical dynamics of the Higgs fields are given by

$$\nabla_\mu \nabla^\mu \phi^a = \frac{\partial V}{\partial \phi^a}, \quad (9)$$

where the ∇_μ ($\mu = 0, \dots, 3$) are the covariant derivatives associated to the metric; in flat spacetime these would simply be the partial derivatives with respect to the coordinates. (Raising and lowering of indices is done using the metric and its inverse, and as always summation over repeated indices is implied.) Using the expression for the potential (8), the field equations (9) can be written in more detail as

$$\nabla_\mu \nabla^\mu \phi^a + \lambda \eta^2 \phi^a - \lambda \phi^2 \phi^a = 0. \quad (10)$$

As usual, the prefactor of the term that is linear in the fields and carries no derivatives gives us the square of the mass⁷, so that the Higgs mass is equal to $m_H = (\lambda \eta^2)^{1/2}$. Apart from that, there is a cubic interaction of the Higgs among themselves, with coupling strength λ .

The presence of the fields affects the geometry of spacetime through the Einstein equations,

$$R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R = 8\pi T^{\mu\nu}. \quad (11)$$

Here $R^{\mu\nu}$ is a tensor formed from the metric $g_{\mu\nu}$ and its first and second derivatives, $R = R^{\mu\nu} g_{\mu\nu}$, and $T^{\mu\nu}$ is the stress-energy tensor of the Higgs fields. One has

$$T^{\mu\nu} = \nabla^\mu \phi^b \nabla^\nu \phi^b - g^{\mu\nu} \left[\frac{1}{2} (\nabla^\kappa \phi^b)(\nabla_\kappa \phi^b) + V(\phi^a) \right], \quad (12)$$

where summation over the indices b is assumed. For derivations we refer to Aitchison and Hey⁷ and Hawking and Ellis¹⁰.

To solve the equations (9) and (11), Nucamendi and Sudarsky started with the ansatz

$$\phi^a = \eta f(r) \frac{x^a}{r} \quad (13)$$

for the fields, and

$$ds^2 = - \left(1 - \frac{2m(r)}{r} \right) e^{-2\delta(r)} dt^2 + \left(1 - \frac{2m(r)}{r} \right)^{-1} dr^2 + (1 - \alpha) r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (14)$$

for the spacetime metric. The x^a are spatial asymptotic Cartesian coordinates, obtained from (r, θ, ϕ) in the same way as Cartesian coordinates in Euclidean space. The functions $f(r)$, $m(r)$, and $\delta(r)$ are to be obtained from the field equations (9) and the Einstein equations (11), with suitable boundary conditions. We note in passing that (14) is indeed of the form (4).

The natural boundary condition for f as $r \rightarrow \infty$ is $f(r) \rightarrow 1$, since one wants the Higgs fields (13) to tend to a minimum of the potential, i.e., $\phi^2 \rightarrow \eta^2$. As was shown by Barriola and Vilenkin⁵, the asymptotic behavior (6) then leads to a globally well-defined solution for both the fields and the metric (apart from a central singularity). However, in numerically solving (9) and (11), Nucamendi and Sudarsky found it more convenient to impose boundary conditions at the origin $r = 0$:

$$\begin{aligned} f(0) &= 0, \\ m(0) &= 0, \\ \frac{\partial m}{\partial r}(0) &= -\frac{\alpha}{2(1-\alpha)}. \end{aligned} \quad (15)$$

The fact that one needs $f \rightarrow 1$ as $r \rightarrow \infty$ then imposes a unique value for $\partial f / \partial r$ at $r = 0$, which can only be determined by a ‘shooting method’: Different values are tried out until a consistent solution for fields and metric is obtained. For more details, the reader is encouraged to read the original paper⁹.

Using the conditions (15), it is easy to convince oneself that the proper mass and charge of an NS monopole as defined by the expansion (5) are indeed zero. The function $u(r)$ is

$$u = r^2 \left(1 - \frac{2m(r)}{r} \right) = -2m(r)r + r^2. \quad (16)$$

Near $r = 0$ there is no zeroth order term in r , so $Q = 0$, and since $m(0) = 0$ the first order term vanishes as well, which gives $M = 0$.

Acknowledgement

It is a pleasure to thank Jorge Pullin for very helpful discussions.

¹ H. Bondi, *Review of Modern Physics* **29** (1957) 423–428

² R.L. Forward, *Journal of Propulsion and Power* **6** (1990) 28–37

³ M. Visser, *Lorentzian Wormholes*, American Institute of Physics, Woodbury, New York, 1995

⁴ R.M. Wald, *General Relativity*, University of Chicago Press, Chigaco, 1984

⁵ M. Barriola and A. Vilenkin, *Physical Review Letters* **63** (1989) 341

⁶ I.J.R. Aitchison and A.J.G. Hey, *Gauge Theories in Particle Physics*, 2nd ed., Institute of Physics Publishing, London, 1989

⁷ M.Y. Zotov, unpublished; LANL preprint gr-qc/0011065

⁸ R. Penrose, in *General Relativity, and Einstein Centenary Survey*, Eds. S.W. Hawking and W. Israel, Cambridge University Press, Cambridge, 1979

⁹ U. Nucamendi and D. Sudarsky, *Classical and Quantum Gravity* **17** (2000) 4051–4058

¹⁰ S.W. Hawking and G.F.R. Ellis, *The large scale structure of spacetime*, Cambridge University Press, Cambridge, 1973

¹¹ An excellent overview of current ideas in cosmology is *COSMO-99: Proceedings of the Third International Workshop on Particle Physics and the Early Universe*, Eds. U. Cotti, R. Jeannerot, G. Senjanović, and A. Smirnov, World Scientific, London, 2000

¹² Y. Mino, M. Sasaki, and T. Tanaka, *Physical Review D* **55** (1997) 3457–3476

¹³ T.C. Quinn and R.M. Wald, *Physical Review D* **56** (1997) 3381–3394

¹⁴ P.S. Letelier and S.R. Oliveira, *Classical and Quantum Gravity* **15** (1998) 421–433

¹⁵ Z. Andrade and R. Price, *Physical Review D* **56** (1997) 6336–6350

A01-34540



**AIAA 2001-3911
Induction and Amplifications of
Non-Newtonian Gravitational Fields**

M. Tajmar

Austrian Research Centers Seibersdorf
Austria

and C.J. de Matos
ESA-ESTEC
The Netherlands

**37th AIAA/ASME/SAE/ASEE Joint Propulsion
Conference and Exhibit**
8-11 July 2001
Salt Lake City, Utah, USA

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

Induction and Amplification of Non-Newtonian Gravitational Fields

M. Tajmar*

Austrian Research Centers Seibersdorf, A-2444 Seibersdorf, Austria

C. J. de Matos†

ESA-ESTEC, Directorate of Scientific Programmes, PO Box 299, NL-2200 AG Noordwijk, The Netherlands

One obtains a Maxwell-like structure of gravitation by applying the weak-field approximation to the well accepted theory of general relativity or by extending Newton's laws to time-dependent systems. This splits gravity in two parts, namely a gravitoelectric and gravitomagnetic (or cogravitational) one. Both solutions differ usually only in the definition of the speed of propagation, the lorentz force law and the expression of the gravitomagnetic potential energy. However, only by extending Newton's laws we obtain a set of Maxwell-like equations which are perfectly isomorphic to electromagnetism. Applying this theory to explain the measured advance of the mercury perihelion we obtain exactly the same prediction as starting from general relativity theory. This is not possible using the weak-field approximation approach. Due to the obtained similar structure between gravitation and electromagnetism, one can express one field by the other one using a coupling constant depending on the mass to charge ratio of the field source. This leads to equations e.g. of how to obtain non-Newtonian gravitational fields by time-varying magnetic fields. Unfortunately the coupling constant is so small that using present day technology engineering applications for gravitation using electromagnetic fields are very difficult. Calculations of induced gravitational fields using state-of-the-art fusion plasmas reach only accelerator threshold values for laboratory testing. Possible amplification mechanisms are mentioned in the literature and need to be explored. We review work by Henry Wallace suggesting a very high gravitomagnetic susceptibility of nuclear half-spin material as well as coupling of charge and mass as shown by e.g. torque pendulum experiments. The possibility of using the principle of equivalence in the weak field approximation to induce non-Newtonian gravitational fields and the influence of electric charge on the free fall of bodies are also investigated, leading to some additional experimental recommendations.

1. Introduction

The control and modification of gravitational fields is a dream pursued by propulsion engineers and physicists around the world. NASA's Breakthrough Propulsion Physics Project is funding exploratory research in this area to stimulate possible breakthroughs in physics that could drastically lower costs for access to space¹. Although not commonly known, Einstein's well accepted general relativity theory, which describes gravitation in our macroscopic world, allows induction phenomena of non-Newtonian gravitational fields similar to Faraday induction in electromagnetic fields by moving heavy masses at high velocities.

The basis for such phenomena are even dating back before general relativity theory when

Oliver Heaviside² in 1893 investigated how energy is propagated in a gravitational field. Since energy propagation in electromagnetic fields is defined by the Poynting vector – a vector product between electric and magnetic fields – Heaviside proposed a gravitational analogue to the magnetic field. Moreover he postulated that this energy must also be propagating at the speed of light. Another approach to the magnetic part of gravity is to start from Newtonian gravity and add the necessary components to conserve momentum and energy³. This leads to the same magnetic component and a finite speed of propagation, the speed of light.

Heaviside's gravitomagnetic fields are hidden in Einstein's Tensor equations. Alternatively, general relativity theory can be written as linear perturbations of Minkowski spacetime. Forward⁴ was the first to show that these perturbations can be rearranged to assemble a Maxwell-type structure which splits gravitation into a gravitoelectric (classical Newtonian gravitation) and a gravitomagnetic (Heaviside's prediction) field. The magnetic effects in gravitation are more commonly known as the Lense-Thirring or frame dragging effect describing precision forces of rotating masses orbiting each other. NASA's mission Gravity Probe B will look for experimental evidence of this effect. Similar to electrodynamics, a variation in gravitomagnetic fields induces a gravitoelectric (non-Newtonian) field and hence provides the possibility to modify gravitation.

* Research Scientist, Space Propulsion,
Phone: +43-50550-3142, Fax: +43-50550-3366,
E-mail: martin.tajmar@arcs.ac.at

† Staff Member, Science Management
Communication Division,
Phone: +31-71-565 3460, Fax: +31-71-565 4101,
E-mail: clovis.de.matos@esa.int

Copyright © 2001 by the American Institute of
Aeronautics and Astronautics Inc. All rights reserved.

Since both gravitation and electromagnetism have the same source, the particle, the authors recently published a paper evaluating coupling constants between both fields⁵ based on the charge-to-mass ratio of the source particle. This paper will review the coupling between gravitation and electromagnetism and point out the limits of present day technology and the expected order of magnitude of non-Newtonian gravitational fields that can be created by this method. Possible amplification mechanism such as ferro-gravitomagnetism and more speculative work published in the literature will be reviewed.

The principle of equivalence in the limit of weak gravitational fields (the gravitational Larmor theorem) will be explored and a possible new effect (the gravitomagnetic Barnett effect) recently suggested by the authors is discussed⁶. However the direct detection of this effect is pending on the possibility to have materials with high gravitomagnetic susceptibility. Nevertheless we show that the principle of equivalence in the weak field approximation together with the gravitational Poynting vector associated with induced non Newtonian gravitational fields (through angular acceleration) account properly for the conservation of energy in the case of cylindrical mass with angular acceleration. This is an encouraging result regarding the possible detection of macroscopic non-Newtonian gravitational fields induced through the angular acceleration of the cylinder in the region located outside the rotating cylinder. The detection of these non-Newtonian gravitational fields outside the cylinder would represent an indirect evidence of the existence of the gravitomagnetic Barnett effect.

Finally the free fall of a massive cylinder carrying electric charge is studied. It is shown that in order to comply with the law of conservation of energy, and with the equivalence principle, the acceleration with which the cylinder will fall depends on its electric charge, its mass and its length.

If the last two effects exposed above are experimentally detected, a technology that can control the free fall of bodies with mass in the laboratory is at hand. If the result is negative, a better empirical understanding of Einstein's general relativity theory in the limit of weak gravitational fields and when extended to electrically charged bodies, would have been achieved, which is a significant scientific result as well.

2. Maxwell Structure of General Relativity Theory

Einstein's field equation⁷ is given by

$$R_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} R = \frac{8\pi G}{c^4} T_{\alpha\beta} \quad (1)$$

During the linearization process, the following limitations are applied:

1. all motions are much slower than the speed of light to neglect special relativity
2. the kinetic or potential energy of all bodies being considered is much smaller than their mass energy to neglect space curvature effects
3. the gravitational fields are always weak enough so that superposition is valid
4. the distance between objects is not so large that we have to take retardation into account

We therefore approximate the metric by

$$g_{\alpha\beta} \approx \eta_{\alpha\beta} + h_{\alpha\beta} \quad (2)$$

where the greek indices $\alpha, \beta = 0, 1, 2, 3$ and $\eta_{\alpha\beta} = (+1, -1, -1, -1)$ is the flat spacetime metric tensor, and $|h_{\alpha\beta}| \ll 1$ is the perturbation to the flat metric. By proper substitutions and after some lengthy calculations⁵ (the reader is referred to the literature for details), we obtain a Maxwell structure of gravitation which is very similar to electromagnetics and only differs due to the fact that masses attract each other and similar charges repel:

$\text{div } \vec{E} = \frac{\rho}{\epsilon_0}$	$\text{div } \vec{g} = -\frac{\rho_m}{\epsilon_g}$
$\text{div } \vec{B} = 0$	$\text{div } \vec{B}_g = 0$
$\text{rot } \vec{E} = -\frac{\partial \vec{B}}{\partial t}$	$\text{rot } \vec{g} = -\frac{\partial \vec{B}_g}{\partial t}$
$\text{rot } \vec{B} = \mu_0 \rho \vec{v}$	$\text{rot } \vec{B}_g = -\mu_g \rho_m \vec{v}$
	$+ \frac{1}{c^2} \frac{\partial \vec{E}}{\partial t}$
	$+ \frac{1}{c^2} \frac{\partial \vec{g}}{\partial t}$

Maxwell Equation Maxwell-Einstein Equation
(Electromagnetism) (Gravitation)

where \vec{g} is the gravitoelectric (or Newtonian gravitational) field and \vec{B}_g the gravitomagnetic field. The gravitational permittivity ϵ_g and gravitomagnetic permeability μ_g is defined as:

$$\epsilon_g = \frac{1}{4\pi G} = 1.19 \times 10^9 \frac{\text{kg} \cdot \text{s}^2}{\text{m}^3} \quad (4)$$

$$\mu_g = \frac{4\pi G}{c^2} = 9.31 \times 10^{-27} \frac{\text{m}}{\text{kg}} \quad (5)$$

by assuming that gravitation propagates at the speed of light c . Although not unusual, this assumption turns out to be very important. Only if gravity propagates at c the Maxwell-Einstein equations match the ones obtained from adding necessary terms to Newtonian gravity to conserve momentum and energy³.

Moreover, the authors could show that with this set of equations, the advance of the Mercury perihelion – one of the most successful tests of general relativity – can be calculated giving the exact prediction than without linearization⁷. This is a very surprising result because the advance of Mercury's perihelion is attributed to a space curvature in general relativity (Schwarzschild metric) which we neglected in our linearization process. The assumption of c as the speed of gravity propagation also implies that the Lorentz force law and the gravitomagnetic potential energy differ from their electromagnetic counterparts by a factor of four⁸. Therefore some authors⁴ use $c/2$ as the speed of gravity propagation to get a gravity Lorentz force law similar to electromagnetics.

The Einstein-Maxwell equations allow to clearly see the gravitomagnetic component of gravitation and the possibility to induce non-Newtonian gravitational fields. Their close relation to electrodynamics allow to transform electromagnetic calculations into their gravitational counterparts⁹.

3. Coupling of Electromagnetism and Gravitation in General Relativity

By comparing gravitation and electromagnetism in Equation (3), we see that both fields are coupled by the e/m ratio of the field source and we can write:

$$\begin{aligned}\ddot{g} &= \kappa \cdot \ddot{E} \\ \ddot{B}_g &= \kappa \cdot \ddot{B}\end{aligned}\quad (6)$$

using the coupling coefficient κ

$$\kappa = -\frac{m \mu_g}{e \mu_0} = -\frac{m \epsilon_0}{e \epsilon_g} = -7.41 \times 10^{-21} \frac{m}{e} \quad (7)$$

Obviously, this coefficient is very small and gravitational effects associated with electromagnetism have never been detected so far¹⁰. By combining Equation (6) with Equation (3), we see how both fields can induce each other:

$$\begin{aligned}\text{rot } \ddot{E} &= -\frac{1}{\kappa} \frac{\partial \ddot{B}_g}{\partial t} \\ \text{rot } \ddot{B} &= \frac{e}{m} \mu_0 \rho_m \ddot{v} + \frac{1}{\kappa c^2} \frac{\partial \ddot{g}}{\partial t}\end{aligned}\quad (8)$$

Coupled Maxwell-Einstein Equations
(Gravitation→Electromagnetism)

$$\begin{aligned}\text{rot } \ddot{g} &= -\kappa \frac{\partial \ddot{B}}{\partial t} \\ \text{rot } \ddot{B}_g &= -\frac{m}{e} \mu_g \rho \ddot{v} - \kappa \frac{1}{c^2} \frac{\partial \ddot{E}}{\partial t}\end{aligned}\quad (9)$$

Coupled Maxwell-Einstein Equations
(Electromagnetism→Gravitation)

For an electron in a vacuum environment $\kappa=4.22 \times 10^{-32}$ kg/C. For example, let us consider an infinitely long coil as shown in Figure 1.




Figure 1 Magnetic Field Induced in a Coil

The magnetic field induced in the center line is then

$$B = \mu_0 I n \quad (10)$$

where I is the current and n is the number of coil winds per length unit. For a current of 10,000 Ampère and 1,000 winds per meter, the magnetic field would be $B=12.56$ T which is state of the art. The corresponding gravitomagnetic field is then $B_g=5.3 \times 10^{-31}$ s⁻¹. Even using a coil with 100,000 winds to induce an electric field, the amplitude of the resulting gravitational field would only be in the order of $g=10^{-26}$ ms⁻². This is much too small to be detected by any accelerometers having measurement thresholds of 10^{-9} ms⁻². By using heavy ions in a plasma instead of electrons we can increase the m/e ratio by 6 orders of magnitude, however, the magnetic fields to contain such a plasma transmitting a similar current of 10,000 Ampère are out of reach.

Nevertheless, although the induced gravitational fields are very small, in principle it is possible to create non-Newtonian gravitational fields along the same principles as we are used to in electromagnetism.

4. Amplification Mechanisms

Since all these electromagnetic-gravitational phenomena are so small, how can we amplify the coupling coefficient in order to obtain measurable non-Newtonian fields?

4.1 Gravitation-Magnetism

Similar to para-, dia-, and ferro-magnetism, the angular and spin momentums from free electrons in material media could be used to obtain a gravitomagnetic relative permeability μ_g , which increases the gravitomagnetic field \ddot{B}_g . Since an alignment of magnetic moments causes also an alignment of gravitomagnetic moments, the gravitomagnetic susceptibility will be the same as the magnetic susceptibility in a magnetized material⁵

$$\chi_g = \chi \quad (11)$$

For our example of the coil in Figure 1, a ferromagnetic core would accordingly increase the gravitomagnetic field and induced non-Newtonian gravitational field by three orders of magnitude. Although significant, the resulting fields are still too low to be detected.

4.2 Coupling of Charge and Mass

All our discussions up to now are based on a coupling at the source particle by the e/m ratio. However, an additional coupling between charge and mass of the source itself might exist and provide a significant amplification mechanism.

Well accepted peer-review journals like Nature and Foundations of Physics featured articles on this topic describing experiments that suggest a coupling between charge and mass in combination with rotation (or acceleration, movement in general).

Dr. Erwin Saxl published an article¹¹ reporting a period change of a torque pendulum if the pendulum was charged. A positive charge caused the pendulum to rotate slower than when it was charged negatively. Figure 3 shows his observations with a small asymmetry of the period change between positive and negative potentials applied to the pendulum. The period is expressed by

$$T = \text{Constant} \cdot \sqrt{\frac{m}{g}} \quad (12)$$

where m is the mass of the pendulum and g the Earth's gravitational acceleration. Assuming that g is not changed (it is highly improbable that the whole Earth is affected), Saxl's measurement can be interpreted as a change of the pendulum's mass by applying an electric potential to it.

Prof. James Woodward from the University of California reported experiments of accelerating masses that, on the other hand, charged up according to their mass and speed of rotation. His experiments were done both for rotating masses¹² as well as for linear accelerated test bodies¹³. Published in the Foundations of Physics and General Relativity and Gravitation, he suggested a broader conservation principle including mass, charge and energy. Results of a test body hitting a target and inducing a charge are shown in Figure 4. His results follow

$$q' \equiv \text{Constant} \cdot m \cdot a \quad (11)$$

where q' is the induced charge, m the test mass and a the acceleration (from rotation or calculated from the impact velocity).

Hence, both Saxl and Woodward experimentally reasoned a relationship between charge, mass and acceleration. A combination of all these factors to reduce/increase the weight of a body is described in a patent by Yamashita and Toyama¹⁴. A cylinder was rotated and charged using a Van der Graff generator. During operation the weight of the rotating cylinder was monitored on a scale. The setup is shown in Figure 5. If the cylinder was charged positively, a positive change of weight up to 4 grams at top speed was indicated. The same charge negative produced a reduction of weight of about 11 grams (out of 1300 grams total weight). This is an asymmetry similar to the one mentioned by Saxl¹¹. Also the relationship between charge, rotation and mass is similar to Saxl and Woodward. The experimentors note that the weight changed according to the speed of the cylinder ruling out electrostatic forces, and that it did not depend on the orientation of rotation ruling out magnetic forces. The reported change of weight (below 1 %) is significant and indicates a very high order of magnitude effect.





Figure 3 Change of Torque Pendulum Period vs. Applied Potential¹¹




Figure 4 Charged Induced by Body Hitting Target¹³




Figure 5 Setup of Charged Rotating Cylinder on Scale¹⁴

4.3 Alignment of Nuclear Spins

Henry Wallace, an engineer from General Electric, holds three patents on a method to produce a macroscopic gravitomagnetic and gravitational field by aligning nuclear spins due to rotation¹⁵⁻¹⁷. He claims that if materials with a net nuclear half-spin (one neutron more than protons in the nucleus) are rotated, this nuclear spin is aligned and produces a macroscopic gravitational effect. This is in fact similar to the Barnett effect where a metal rod is rotated and magnetisation of the material is observed. However, macroscopic magnetism in electromagnetism is caused by spin alignment of electrons, nuclear magnetism plays a very minor role due to the much higher mass of a proton or neutron compared to the electron. In a gravitational context the difference in mass is no major drawback anymore and nuclear magnetism should be on the same order of magnitude than electron magnetism. Usually, very low temperatures in the order of nano Kelvin are required to align nuclear spins, simple rotation would be much more easy.

The contribution of neutron spins to gravitomagnetic fields is theoretically on the order of

ferromagnetism⁵. However, since Wallace claims to have measured at least the induction of nuclear spin alignment in a rotating detector material – by what he thinks a gravitomagnetic field, possible unknown amplification mechanisms (quantum gravity, nuclear strong force interaction) could cause much higher order of magnitude effects.

His setup is shown in Figure 6. A generator assembly (test mass rotating in 2 axis) is mounted on the left side and a detector assembly (similar to generator) is mounted on the right side with the possibility of rotation in the plane of the paper. A laser is monitoring the oscillations of this detector assembly. If both are rotated in the same orientation and counter wise, the laser detected a difference (Figure 7) which Wallace attributed to a force field. Since it only depended on the nuclear spin (e.g. Iron did not work but is a strong ferromagnetic material), Wallace ruled out magnetism as the origin of the force. In a different setup he showed that the field generated could constructively reduce the vibrational degrees of freedom of the crystal structure resulting in a change of its electrical properties (Figure 8).




Figure 6 Setup of Rotating Test Mass (2 Axis) and Generator (Left) and Detector (Right) Position¹⁵




Figure 7 Oscillations of Detector Assembly¹⁵




Figure 8 Change in Thermal Vibration of Crystal Lattices¹⁵

Hence, there is quite some experimental evidence for an amplification mechanism through nuclear magnetism to generate non-Newtonian gravitational fields using effects predicted by general relativity theory.

5. Principle of Equivalence and High-Order of Magnitude Non-Newtonian Gravitation

We explored the limits of inducing non-Newtonian gravitation using general relativity theory as well as looking at possible and speculative amplification mechanisms. Let us go back to the foundation of gravitation itself and explore the principle of equivalence in the limit of weak gravitational fields.

Einstein based his thoughts of gravitation on a famous Gedankenexperiment explaining the principle of equivalence: An observer can not distinguish between being inside a falling elevator or in a uniform gravitational field. Based on this equivalence, he developed the geometrical structure of general relativity. In the limit of weak gravitational fields, this simple Gedankenexperiment however is not complete as it covers only gravitoelectric fields and not the magnetic component of gravitation. According to the Larmor theorem of electromagnetics, a magnetic field can be replaced locally by a rotating reference frame with the Larmor frequency

$$\omega_L = \frac{1}{2} \frac{e}{m} B \quad (14)$$

The same argument applies for gravitation and a rotating reference frame rotating with the Larmor frequency can replace a gravitomagnetic field

$$\omega_{Lg} = -B_g \quad (15)$$

independent of the particle mass. The principle of equivalence¹⁸ for weak gravitational fields (neglecting space curvature) also called gravitational Larmor theorem (GLT) should then be:

An observer can not distinguish between a uniformly accelerated (\ddot{v}) reference frame rotating with the gravitational Larmor frequency (ω_{Lg}) and a reference frame at rest in a corresponding gravitational field ($\vec{\gamma} = -\dot{\vec{v}}$, $\omega_{Lg} = -\vec{B}_g$).

But what happens if the speed of rotation of the elevator changes? According to the GLT, this would correspond to a change of a gravitomagnetic field flux and therefore induce a non-Newtonian gravitational component according to the gravitational Faraday law:

$$\begin{aligned} \epsilon_g &= \oint_{\Gamma} \vec{\gamma} \cdot d\vec{l} = -\frac{d\phi_{gm}}{dt} = -\frac{d}{dt} \iint_{\Sigma} \vec{B}_g \cdot d\vec{\sigma} \quad (16) \\ &= \frac{d}{dt} \iint_{\Sigma} \vec{\Omega} \cdot d\vec{\sigma} \end{aligned}$$

where $\vec{\gamma}$ is the non-Newtonian gravitational field, Γ and Σ are respectively the contour and surface of integration, ϕ_{gm} is the gravitomagnetic flux, \vec{B}_g is the gravitomagnetic field, and $\vec{\Omega}$ is the angular velocity of the reference frame. If the observer measures this additional gravitational field the principle of equivalence holds and he can not distinguish between the elevator and the gravitational field. If he does not observe this effect, the gravitational Larmor theorem is not valid, as a weak field approximation to Einstein's general relativity theory. We will show later that these "induced" non-Newtonian gravitational fields contribute to account for the mechanical energy absorbed (dissipated) by a rotating body during the phase of angular acceleration (deceleration).

Suppose the gravitational Larmor theorem holds, every rotation corresponds to a gravitomagnetic field, which is many orders of magnitude higher than the gravitomagnetic field responsible for the precession forces in the classical Lense-Thirring effect.

5.1 Gravitomagnetic Barnett Effect

The authors discussed such rotational effect described as the gravitational Barnett effect⁶. In 1915 Barnett¹⁹ observed that a body of any substance set into rotation becomes the seat of a uniform intrinsic magnetic field parallel to the axis of rotation, and proportional to the angular velocity. If the substance is magnetic, magnetization results, otherwise not. This physical phenomenon is referred to as *magnetization by rotation* or as the *Barnett effect*.

If a mechanical momentum with angular velocity Ω is applied to a substance, it will create a force on the elementary gyrostats (electrons orbiting the nucleus) trying to align them. This is equivalent to the effect of a magnetic field in this substance B_{equi} and we can write:

$$B_{equi} = -\frac{1}{g_I} \frac{2m}{e} \Omega \quad (17)$$

where g_I is the Landé factor for obtaining the correct gyromagnetic ratio. We can now apply the same argument to the gravitational case and postulate an equivalent gravitomagnetic field $B_{g equi}$ which counteracts the mechanical momentum:

$$B_{g equi} = -\frac{2}{g} \Omega \quad (18)$$

For an electron, $g=2$ and we see that physical rotation is indeed equivalent to a gravitomagnetic field. From Equation (18) we can compute the gravitomagnetization acquired by the rotating material:

$$\tilde{M}_g = \frac{\chi_g}{\mu_0 g} \tilde{B}_{g\,equi} = -\frac{\chi_g}{\mu_0 g} \frac{2}{g} \tilde{\Omega} \quad (19)$$

where χ_g is the gravitomagnetic susceptibility.

Taking into account the coupling between gravitation and electromagnetism presented above we can demonstrate the general result:

$$\tilde{M}_g = -\frac{\chi}{\mu_0 g} \left(\frac{m}{e} \right)^2 \tilde{\Omega} \quad (20)$$

where χ is the magnetic susceptibility of the material⁶. This indicates that the gravitomagnetic moment associated with the substance will be extremely small. Therefore we can not use this gravitomagnetic moment to induce macroscopic non-Newtonian gravitational fields. However we can show, following our discussion on the equivalence principle, that if the field of rotation in Equation (18) can not be distinguished from gravitomagnetism, it must be a real field which we can use to induce non-Newtonian gravitational fields. The detection of such fields would represent an indirect proof of the existence of the gravitomagnetic Barnett effect.

5.2 Gravitational Poynting Vector and Gravitational Larmor Theorem in Rotating Bodies with Angular Acceleration

The gravitational Poynting vector, defined as the vectorial product between the gravitational and the gravitomagnetic fields, $\tilde{S}_g = \frac{c^2}{4\pi G} \tilde{\gamma} \times \tilde{B}_g$, provides a mechanism for the transfer of gravitational energy to a system of falling objects (we will consider in the following a cylindrical mass m , with radius a and length l). It has been shown²⁰ that using the gravitational Poynting vector, the rate at which the kinetic energy of a falling body increases is completely accounted by the influx of gravitational field energy into the body. Applying the gravitational Larmor Theorem (GLT) to a body with angular acceleration. We get that a time varying angular velocity flux will be associated with a non-Newtonian gravitational field proportional to the tangential acceleration. The gravitational electromotive force produced in a gyrogravitomagnetic experiment can be calculated using the gravitational Faraday induction law as given in Eqn (16). Together with the GLT expressed through Equation (15) we get

$$\varepsilon_g = \frac{d}{dt} \iint_{\Sigma} \tilde{\Omega} \cdot d\vec{\sigma} \quad (21)$$

The induced non-Newtonian gravitational field associated with this gravitational electromotive force is at the surface of the cylinder is:

$$\oint_{\Gamma} \tilde{\gamma} \cdot d\vec{l} = \frac{d}{dt} \iint_{\Sigma} \tilde{\Omega} \cdot d\vec{\sigma} \quad (22)$$

$$\tilde{\gamma}_\theta = \frac{1}{2} a \dot{\Omega} \hat{e}_\theta \quad (23)$$

From this non-Newtonian gravitational field and the gravitomagnetic field produced by the rotating mass current, we can compute a gravitational poynting vector

$$\tilde{S}_{g\Omega} = \frac{c^2}{4\pi G} \tilde{\gamma}_\theta \times \tilde{B}_{g\Omega} = \frac{1}{4\pi} \left(\frac{a}{a + \ell} \right) \Omega \dot{\Omega} m \hat{n} \quad (24)$$

which will also provide an energy transfer mechanism to explain how massive bodies acquire rotational kinetic energy when mechanical forces are applied on them²¹. The rate at which the rotational kinetic energy of a body increases (or decreases) due to the application of external mechanical forces on that body, is completely accounted by the influx (out-flux) of gravitational energy into (outward) the body.

$$\left. \frac{dU}{dt} \right|_{\Omega} = S_{g\Omega} (2\pi a^2 + 2\pi a \ell) = \frac{d}{dt} \left(\frac{1}{2} I \Omega^2 \right) \quad (25)$$

where I is the moment of inertia of the cylinder. This demonstrates the validity of the gravitational Larmor theorem, and shows how the transfer of mechanical work to a body can be interpreted as a flux of gravitational energy associated with non-Newtonian gravitational fields produced by time varying angular velocities. This is an encouraging result regarding the possible detection of macroscopic non-Newtonian gravitational fields induced through the angular acceleration of the cylinder in the region located outside the rotating cylinder. The non-Newtonian gravitational field outside the cylinder is given by:

$$\gamma = \frac{1}{2} \frac{a^2}{r} \dot{\Omega} \quad (26)$$

where $r > a$ is the distance from the cylinder's longitudinal axis. For $r \leq a$ we have, $\gamma = 0.5 \times a \dot{\Omega}$. For the following values of $r=1$ m, $a=0.1$ m, $\dot{\Omega} = 200$ Hz/s, γ will have the value of 1 ms^{-2} . We recommend that experiments shall be performed with the aim of evaluating Equation (26).

Is it possible to use fluxes of radiated electromagnetic energy to counteract the effect of absorbed fluxes of gravitational energy? That is a question Saxl, Woodward and Yamashita tried to evaluate empirically. These empirical approaches shall be complemented in the following by a

theoretical analysis of the net energy flow associated with the free fall of an electrically charged cylindrical mass.

5.3 Free Fall of a Cylindrical Mass Electrically Charged

A cylindrical mass electrically charged in free fall must comply with the law of conservation of energy and with the principle of equivalence²². During the free fall the cylindrical mass will absorb gravitational energy, which is described by the following gravitational Poynting vector:

$$\tilde{S}_g = \frac{c^2}{4\pi G} \vec{\gamma} \times \vec{B}_g = \frac{mv\gamma}{2\pi a\ell} \hat{n}_{in} \quad (27)$$

where v is the speed of the cylinder while it is falling, γ is the Earth gravitational field, m , a , ℓ are respectively the mass, length and radius of the cylinder and \hat{n}_{in} is a unit vector orthogonal to the surface of the cylinder and Poynting inwards. The cylinder due to its electric charge will also radiate electromagnetic energy according to the following electromagnetic Poynting vector:

$$\tilde{S}_{em} = \frac{1}{\mu_0} \vec{B} \times \vec{E} = \frac{\mu_0}{8\pi^2} \frac{Q^2 v \dot{v}}{a\ell(a+\ell)} \hat{n}_{out} \quad (28)$$

where Q is the electric charge carried by the cylinder, μ_0 is the magnetic permeability of vacuum and \hat{n}_{out} is a unit vector orthogonal to the surface of the cylinder and Poynting outwards. The principle of equivalence states that if the cylinder is at rest with respect to a reference frame which is uniformly accelerating upwards (with respect to the laboratory) with acceleration $\dot{v} = \gamma \hat{e}_z$, the cylinder will radiate (with respect to the laboratory) according to the following Poynting vector:

$$\tilde{S}_{em} = \frac{1}{\mu_0} \vec{B} \times \vec{E} = \frac{\mu_0}{8\pi^2} \frac{Q^2 v \gamma}{a\ell(a+\ell)} \hat{n}_{out} \quad (29)$$

Therefore to comply with the principle of equivalence, we shall take in Equation (28) $\dot{v} = \gamma$ ²³.

The sum of both energy fluxes in Equations (27) and (29) must comply with the law of conservation of energy. Therefore the Sum of gravitational incoming flux and the radiated electromagnetic energy flux must be equal to the rate at which the kinetic energy of the body varies in time.

$$S_{em}(2\pi a^2 + 2\pi a\ell) + S_g 2\pi a\ell = \frac{d}{dt} \left(\frac{1}{2} mv^2 \right) \quad (30)$$

From Equation (30) we deduce that the acceleration with which the electrically charged cylindrical mass will fall is:

$$\dot{v} = \gamma \left(1 - \frac{\mu_0 Q^2}{4\pi m \ell} \right) \quad (31)$$

Equation (31) shows that the free fall of an electrically charged body would violate the law of Galilean free fall, because the acceleration of fall would depend on the electric charge, size and mass of the falling body. The fact that the acceleration of fall depends on the square of the electric charge rules out the possibility to explain with the present analysis, the observations of Saxl and Yamashita, regarding the increase of mass for positively charged bodies and the decrease of mass for negatively charged bodies. Notice that following the rational which leads to equation (31), the phenomenon described by this equation should happen either in a reference frame at rest in an external gravitational field or inside a uniformly accelerated reference frame, therefore we are not able to use this phenomenon to distinguish between both situations. Consequently equation (31) do not violate the principle of equivalence. To test equation (31) we propose to measure the time of fall of charged cylindrical capacitors, and compare it with the time of fall of similar uncharged capacitors. For $m=10$ grams, $\ell=10$ cm, $Q=100$ C, we will have $\dot{v}=0$. For these values the cylinder would not be able to fall! However to avoid disruption currents for such a high value of electric charge is a technological challenge.

Conclusion

In the present work we did an extensive review of possible "classical ways" to induce non-Newtonian gravitational fields from electromagnetic phenomena or by using the principle of equivalence in the limit of weak gravitational fields. If the experiments performed by Saxl, Yamashita, Woodward and Wallace were reproducible this would represent a breakthrough in the possibility to control gravitational phenomena at the laboratory scale. The understanding of the principle of equivalence for electrically charged bodies and in the limit of weak gravitational fields is crucial to evaluate respectively:

- the possibility of directly convert gravitational energy into electromagnetic energy during the free fall of an electrically charged body.
- the possibility of inducing non-Newtonian gravitational fields through the angular acceleration we might communicate to solid bodies.

The experimental confirmation of such phenomena would be a dramatic step forward in the technological control of free fall. The non detection of the presented phenomena could lead to a better

empirical understanding of Einstein's general relativity theory in the limit of weak gravitational fields and when extended to electrically charged bodies, which is a significant scientific result as well. These experiments could also contribute to decide which approach to weak gravity is the correct one, i.e. linearized general relativity or the extension of Newton's laws to time dependent systems.

References

¹Millis, M., "Breakthrough Propulsion Physics Research Program", NASA TM 107381, Lewis Research Center, 1997

²Heaviside, O., "A Gravitational and Electromagnetic Analogy", *The Electrician*, **31**, 1893, pp. 281-282 and 359

³Jefimenko, O.D., "Causality, Electromagnetic Induction and Gravitation," Electret Scientific Company, 1992

⁴Forward, R.L. "General Relativity for the Experimentalist", *Proceedings of the IRE*, **49**, 1961, pp. 892-586

⁵Tajmar, M., and de Matos, C.J., "Coupling of Electromagnetism and Gravitation in the Weak Field Approximation," *Journal of Theoretics*, **3**(1), 2001

⁶De Matos, C.J., and Tajmar, M., "Gravitomagnetic Barnett Effect", *Indian Journal of Physics*, 2001, submitted

⁷Misner, C.W., Thorne, K.S., and Wheeler, J.A., "Gravitation", W.H.Freeman, San Francisco, 1973

⁸De Matos, C.J., and Tajmar, M., "Advance of Mercury Perihelion Explained by Cogravity", Proceedings of the Spanish Relativity Meeting, 2000

⁹Forward, R.L., "Guidelines to Antigravity", *American Journal of Physics*, **31**, 1963, pp. 166-170

¹⁰Braginski, V.B., Caves, C.M., Thorne, K.S., "Laboratory Experiments to Test Relativity Gravity", *Physical Review D*, **15**(5), 1977, pp. 2047-2068

¹¹Saxl, E.J., "An Electrically Charged Torque Pendulum", *Nature*, **203**, 1964, pp. 136-138

¹²Woodward, J.F., "Electrogravitational Induction and Rotation", *Foundations of Physics*, **12**(5), 1982, pp. 467-479

¹³Woodward, J.F., "An Experimental Reexamination of Faraday Electrogravitational Induction", *General Relativity and Gravitation*, **12**, 1980, pp. 1055-1067

¹⁴Yamashita, H., and Toyama, T., "Machine for Acceleration in a Gravitational Field", European Patent EP 0 486 243 A2, 1982

¹⁵Wallace, H.W., "Method and Apparatus for Generating a Secondary Gravitational Force Field", US Patent 3.626.605, 1971

¹⁶Wallace, H.W., "Method and Apparatus for Generating a Dynamic Force Field", US Patent 3.626.606, 1971

¹⁷Wallace, H.W., "Heat Pump", US Patent 3.823.570, 1974

¹⁸Mashhoon, B., "Gravitoelectromagnetism", Los Alamos Archive, gr-qc/0011014, 2000

¹⁹Barnett, S.J., "Magnetization by Rotation", *Physical Review*, **4**(4), 1915

²⁰Krumm, P., Bedford, D., "The Gravitational Poynting Vector and Energy Transfer", *American Journal of Physics*, **55**(4), 1987, pp. 362-363

²¹De Matos, C. J., Tajmar, M., "Gravitational Poynting Vector, Gravitational Larmor Theorem and Rotational Kinetic Energy", submitted to Los Alamos Archive, 2001

²²De Matos, C. J., Tajmar, M., "Free Fall of a Cylindrical Mass Electrically Charged", submitted to Los Alamos Archive, 2001

²³Soker, N., "Radiation from an Accelerated Charge and the Principle of Equivalence", NASA Breakthrough Propulsion Physics Workshop Proceedings, NASA / CP - 1999-208694, 1999, pp. 427-440



A01-34541

AIAA 2001-3912

TOWARDS THE CONTROL OF MATTER WITH GRAVITY

Dr D Burton, Dr S Clark, Prof T Dereli, Dr J Gratus
Mr W Johnson, Prof R W Tucker, Dr C Wang

Department of Physics,
Lancaster University,
Lancaster LA1 4YB, UK

37th Aerospace Sciences Meeting and Exhibit
8-11 July 2001
Salt Lake City, Utah

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

TOWARDS THE CONTROL OF MATTER WITH GRAVITY

Dr D Burton, Dr S Clark, Prof T Dereli, Dr J Gratus
Mr W Johnston, Prof R W Tucker, Dr C Wang

Department of Physics, Lancaster University, Lancaster LA1 4YB, UK

d.burton@lancaster.ac.uk, t.dereli@lancaster.ac.uk, j.gratus@lancaster.ac.uk
robin.tucker@lancaster.ac.uk, c.wang@lancaster.ac.uk

Abstract

This article summarises some of the current activities of the gravitational research group above. After surveying the role of gauge fields as carriers of the basic forces in physics and the momentum conservation constraints placed on current propulsion technology a number of alternative strategies, under active investigation, are briefly discussed. The essential theme is the detection of hitherto undetected components of gravitation that may have implications for the motion of matter. Such components include gravito-magnetic fields and radiation modes as well as new scalar interactions resulting from various proposals for the unification of gravitation with the other basic forces. Experiments are suggested that may shed light on the existence of novel gravitational interactions. These include the detection of transient gravitational fields by means of orbiting Cosserat antennae and the effects of torsional forces on the perihelion shift of planets.

1 Forces and Physics

The effective exploration of interplanetary systems and beyond demands advances in propulsion technology. Our visionaries tell us that the pace of development will be interrupted by sudden advances analogous to the past discoveries of manned flight and reciprocating combustion, gas turbine and rocket engines. Besides "technological breakthroughs" there is also a hope that new "conceptual breakthroughs" in the foundations of physics will offer new perspectives as guides in the search for new technologies. Harnessing the power of the electromagnetic field had to wait for Faraday and Maxwell and the development of nuclear power sprang from an understanding of the quantum nature of matter and the relativistic structure of space-time. History teaches one that major developments in science and engineering are not always predictable within the context of the prevailing wisdom of the time.

Modern physics continues to power technology on many fronts. Much of this development springs

from the quantum nature of matter and radiation. By contrast gravitation defies a unique and consistent quantum formulation. Its most elegant formulation is in terms of the classical geometrical properties of space-time and it is related to the other interactions in nature at the level of classical physics. Here classical fields are the primary structures and matter is modelled as "crystallised field energy". Such concentrations of energy owe their existence to the basic quantum laws of nature but classical laws are often adequate to describe the interactions of such matter with fields on a macroscopic level. These interactions depend on a small number of physical attributes such as inertial mass and interaction strengths ("charges") that enter into the classical field equations. These are expressed as tensor equations on a dynamic space-time in general. Thus electromagnetic fields couple to electric charge, nuclear force fields to colour charge, the forces responsible for radioactivity to weak-hypercharge and the mass-energy of all fields to gravity. The significance of different force fields in any particular situation is in large measure determined by the relative sizes of these couplings. This follows since most basic laws follow remarkable mathematical similarities. See Figure 1. They are all associated with symmetries of mathematical patterns (Lie groups) with field

Copyright ©2001 by Robin Tucker. Published by the American Institute of Aeronautics and Astronautics, Inc with permission

Interaction	Mediating Quanta	Lie Group	Connection	Dynamics
Electromagnetic: Electric charge	QED: Photon	U(1)	\mathbf{A}_6	$\mathbf{F} = d\mathbf{A}_6$ $d \star \mathbf{F} = \mathbf{J}[\Phi, \mathbf{A}_6]$
Electroweak: Hypercharge	QAD: W and Z Bosons	$SU(2) \times U(1)$	\mathbf{A}_4	$\mathbf{F} = d\mathbf{A}_4 + [\mathbf{A}_4, \mathbf{A}_4]$ $D \star \mathbf{F} = \mathbf{J}[\Phi, \mathbf{A}_4]$
Strong: Colourcharge	QCD: Gluons	SU(3)	\mathbf{A}_3	$\mathbf{F} = d\mathbf{A}_3 + [\mathbf{A}_3, \mathbf{A}_3]$ $D \star \mathbf{F} = \mathbf{J}[\Phi, \mathbf{A}_3]$
Gravity: Mass-energy and spin	QG: Graviton	Spin(3,1)	\mathbf{A}_5	$\mathbf{F} = d\mathbf{A}_5 + [\mathbf{A}_5, \mathbf{A}_5]$ $E[g, \mathbf{A}_5] = T[g, \Phi, \mathbf{A}_6, \mathbf{A}_5, \mathbf{A}_3, \mathbf{A}_4]$

Figure 1: This table classifies the known force fields in terms of their attributes. Each can be associated with a mathematical symmetry (Lie group) that determines how the fields interact with themselves, other force fields and matter at each event in space-time. The precise coupling to matter is determined by different kinds of charge. Thus electrically charged particles couple to the electromagnetic field, colour charged particles to the chromodynamic nuclear field and weak-hypercharged particles to the electro-weak fields responsible for radioactive decay. At the quantum level these fields are mediated by field quanta; photons, gluons and W-Z bosons respectively. All matter and fields with mass-energy couple to gravity. It is expected that the quanta of gravitational waves (gravitons) will be responsible for a theory of quantum gravity. The dynamics of all these fields have many similarities (they arise from “connections associated with Lie groups”) although gravitation occupies a unique position in classical physics since it is intimately related to the structure of space and time itself.

strengths defined in terms of directions (“connections”) in collections of mathematical structures at each event in space-time. The gravitational field is similarly determined by a field strength associated with a Lie group although according to Einstein its field equation breaks the mould set by the other force fields in nature. This difference is accentuated by the recognition that relativistic gravitation may be formulated in terms of space-time geometry, an idea that has had profound implications for cosmogony and cosmology.

1.1 Propulsion Constraints

Given a propulsion unit, its effectiveness can usually be gauged from the laws of *classical* physics. In most cases forces on matter are prescribed and Newtonian dynamics is in principle adequate to describe the subsequent motion. For propulsion in the vicinity of the earth the earth’s gravitational field (along with lift and drag of the air) is omni-present and propulsion forces are derived from the rate of change of momentum between “fuel” and “propelled system”. If propellant is to be ejected as radiation or systems achieve speeds that are an appreciable fraction of the speed of light c , Newtonian dynamics is inadequate and must be replaced by the more accurate relativistic equations of motion.

A great deal can be learnt about the demands of enhanced propulsion proposals by elementary

considerations based on well established physical principles¹. It is inconceivable that the constraints imposed by such principles will require dramatic refinement by future scientific developments.

It is useful to distinguish between propulsion systems that must operate in the vicinity of the earth’s surface and those in which Newtonian gravitational fields are negligible. With current technology expensive multi-stage chemical propellants are required to place structures into earth orbit and beyond. For a single propulsion system of mass $M(t)$ and speed $V(t)$ at time t , lifting vertically off the surface of a non-rotating earth with zero initial velocity and ejecting mass at a speed u_0 relative to the propelled system, $V(t)$ is given by

$$V(t) = -u_0 \log \frac{M(0)}{M(t)} - gt \quad (1)$$

in a region where the acceleration due to gravity g may be considered constant. If the rate of mass ejection is constant one may eliminate the burn time $t = t_b$ given the specific thrust ratio $\mathcal{R} = 1 + \frac{A}{g}$ where A is the initial acceleration of the system at lift off. Then

$$\frac{V(t_b)}{V_{esc}} = \frac{u_0}{V_{esc}} \left\{ \log \frac{M(t_b)}{M(0)} + \frac{1}{\mathcal{R}} \left(1 - \frac{M(t_b)}{M(0)} \right) \right\}$$

where V_{esc} is the escape velocity for the earth. Figure 2 shows $V(t_b)/V_{esc}$ versus $M(t_b)/M(0)$ for three values of u_0/V_{esc} appropriate for chemical propellants for both $\mathcal{R} = 1$ and $\mathcal{R} = 2$. These

estimates set a benchmark for any new terrestrial propulsion system that claims to improve current multi-staging and chemical propellant technology.




Figure 2: These plots indicate the mass ratio $M(t_b)/M(0)$ required to reach a non-relativistic speed V_b in the presence of the earth's (uniform) gravitational field. V_{esc} is the escape velocity for the earth. The upper three (dotted) curves correspond to a specific thrust parameter $R = 2$. The lower three (full) curves correspond to $R = 1$. In each triplet the ejection speed is chosen to vary according to three typical chemical propellants.

1.2 Relativistic Rockets

In an extra-terrestrial environment where Newtonian gravitational fields are negligible the system dynamics are different. To accommodate relativistic mass-energy ejection one has from consideration of rectilinear relativistic impulse dynamics in an inertial frame:

$$\lim_{\Delta t \rightarrow 0} \frac{p_{t+\Delta t} - p_t}{\Delta t} = 0 \quad (2)$$

where the 4-momenta of the composite system are respectively

$$p_t = \Pi(V, M), \quad (3)$$

and

$$p_{t+\Delta t} =$$

$$\Pi(V + dV, M + dM) + \Pi \left(\frac{V + dV - u_0}{1 - \frac{u_0(V+dV)}{c^2}}, dm \right) \quad (4)$$

for

$$\Pi(V, M) = \frac{M}{(1 - \frac{V^2}{c^2})^{1/2}} [c, V, 0, 0]. \quad (5)$$

Taking the limit in (2) and eliminating dm yields:

$$\frac{1}{M(t)} \frac{dM(t)}{dt} = - \frac{dV(t)}{dt} \frac{1}{u_0(1 - \frac{V^2}{c^2})} \quad (6)$$

where $V(t)$ is the speed of the system relative to an inertial frame with clock time t . Here u_0 is the mass-energy ejection speed relative to the system at any ejection event. If the initial speed of the system is $V(0)$ in the inertial frame and u_0 is constant then at any time $t > 0$ where mass-energy ejection continues it follows that:

$$\frac{V(t)}{c} = \frac{1 - \alpha \mu(t)^\lambda}{1 + \alpha \mu(t)^\lambda} \quad (7)$$

where $\alpha = \left(1 - \frac{V(0)}{c}\right) / \left(1 + \frac{V(0)}{c}\right)$, $\mu(t) = M(t)/M(0)$, $\lambda = 2u_0/c$. This reduces to (1) for $g = 0$ when $c \mapsto \infty$. For chemical propellants with $u_0 \approx 10 \text{ km/sec}$, $M(t) \approx 1 \text{ ton}$ and for a final speed $V(t) \approx c/2$ this implies $M(0) \approx 10^{7000} \text{ tons!}$ In Figure 3, V/c is plotted against the mass ratio μ for any t for $V(0) = 0$ and varying values of u_0/c .




Figure 3: These plots indicate the mass-energy ratio $M(t)/M(0)$ required to reach a relativistic speed ratio V/c in the absence of any gravitational field. Each curve has a definite ejection speed ratio u_0/c relative to the propulsion system. These are $1/20, 1/10, 1/5, 3/10, 1/2, 7/10$ and $9/10$ in order starting from the bottom of the plot. The advantage of relativistic mass-energy ejection is apparent from these curves.

The advantages of high speed ejection $u_0 \mapsto c$ over non-relativistic mass ejection are apparent in this picture. To establish propulsion efficiency in such cases one needs to know how the ejection

mass ratio μ varies with proper time τ (i.e. the time measured by an on board clock). This permits one to integrate equation (7) for the motion of the system. Even for on-board robots a consideration of non-destructive proper accelerations is clearly desirable. Writing $\mu(\tau) = \exp(-\rho(\tau))$ the rectilinear relativistic equations of motion (2) of the system become:

$$\frac{dx(\tau)}{d\tau} = c \sinh \frac{\lambda\rho(\tau)}{2} \quad (8)$$

$$\frac{dt(\tau)}{d\tau} = \cosh \frac{\lambda\rho(\tau)}{2}. \quad (9)$$

The second equation (9) relates the on-board passage of time measured by τ to a clock fixed in the inertial frame measuring t . An attractive solution can be found that gives the propulsion system a constant proper acceleration A :

$$x(\tau) = \frac{c^2}{A} \left\{ \cosh \frac{A}{c}\tau - 1 \right\} \quad (10)$$

$$t(\tau) = \frac{c}{A} \sinh \frac{A}{c}\tau. \quad (11)$$

To achieve such a uniform proper acceleration the proper mass ejection rate is required to satisfy

$$\frac{d\rho(\tau)}{d\tau} = \frac{A}{u_0}. \quad (12)$$

If a "photon" rocket ($u_0 = c$) could be produced to achieve an on-board mass-energy ejection system with $A = g$ (9.8 m/sec/sec) for an interval of time τ_b then $\mu(\tau_b) = \exp(-\frac{g\tau_b}{c})$. One readily finds that $M(0) = M(\tau_b) \exp(\kappa)$ if τ_b is taken as κ years. These considerations have powerful implications given the scales of space and time compared to those of human existence. For a round trip to a neighbouring star by such a photon rocket with $\kappa \approx 9$ one has $\exp(9) \approx 8000$. Even a mass ratio on this scale in a compact unit raises severe issues of heat dissipation.

2 Alternative Strategies

An alternative approach is to restrict mass-energy ejection mechanisms to local control programmes and seek dynamical propulsion in terms of interactions between ambient fields and their associated charges. Thus the system may be regarded as spending most of its time "coasting" or "surfing" in space. Standard "coasting" in a gravitational field is often referred to as "free-fall" and the motion of some point in the system under such conditions can be used as a reference of zero proper acceleration. It would be a rare occurrence for space travel to be based entirely on free fall in an ambient gravitational field. Departure

from such motion is often associated with non-gravitational forces. Like any surfer or swimmer, adjusting body configuration to the environment can optimise transport. Such manoeuvring require a local expenditure of energy. However it may be possible to mine this energy from the environment itself. A natural source in the solar system is the sun itself or power transmitted by microwave from orbiting space-stations. When these sources are not available recourse to other fields including the gravitational field itself may be necessary. Thus it is important to understand the interaction of fields and gravity with matter, particularly if the latter has an extended structure that can respond to the geometry of space-time.

2.1 Gravcraft

To illustrate a "swimming" mechanism one may explore the concept of the gravcraft. In 1968 Beletsky and Givertz ^{2, 3, 4} introduced this notion, powered by actively controlling the dimensions of an extended body in orbit. Effective rates of orbital elevation above the earth demanded eccentric orbits and tethered structures on the order of several thousand kilometers in extent. Similar systems ^{5, 6, 7, 8} have been proposed to enhance maneuverability without the intervention of chemical propellants and offer an attractive means of cargo transfer between orbiting stations and energy generation. However, most theoretical estimates of the transfer speed between orbits have remained disappointingly low despite the obvious advantages of relying on environmental energy sources such as solar radiation and gravitational tidal forces*. Gravcraft motion relies on its coupling to the tidal component of a gravitational field. For static gravitational fields the spatial dimension of a gravcraft is determined by the scale over which the field varies. We have recently devised⁹ a new method of gravcraft propulsion by actively varying its instantaneous moment of inertia in a quasi-circular orbit. A primitive craft is composed of two massive bodies connected via a light tether whose length can vary between l_{max} and l_{min} by the use of powered winches inside the bodies. By varying the moment of inertia of such a rotating system it is possible to transfer energy between the spin and orbital angular momentum of the craft. If the cycle of tether length variation is chosen appropriately it is possible for the entire structure to gain lift in the gravitational field. In

*Another attractive feature of gravcraft structures controlled by tethers in the ionosphere is their potential for generating power from ambient magnetic fields. In 1996 NASA used a conducting tether connected to the space shuttle in order to generate electricity.

the static field of a spherical mass M an estimate of the average rate of height gain at radius r is given by:

$$\dot{r} = \frac{3(GM)^{1/2}}{4\pi r^{5/2}} (l_{max}^2 - l_{min}^2) \quad (13)$$

where G is the Newtonian gravitational coupling constant. For a gravicraft in low earth orbit with a tether length varying between 50 and 10 km this yields $\dot{r} \simeq 300m/hr$ and if its mass is 10^3 kg it expends about 425 watts. This ascension rate is an order of magnitude greater than any previous gravicraft mechanism known to the authors. Although primitive, such a mechanism illustrates how one can in principle manoeuvre matter in an ambient gravitational field by exploiting its interaction with the field gradient. It also shows that the smallness of G demands a significant sustained gradient field for this mechanism to offer a viable means of long term propulsion. It may however offer a viable means to transport cargo between space-stations in distinct orbits by taking energy from the sun.

2.2 Gravito-electromagnetism

Several authors have noticed that a subset of the Einstein equations when perturbed about flat spacetime can be written in a form that looks remarkably similar to Maxwell's equations with the Newtonian gravitational field corresponding to the *gravito-electric* field and mass-currents playing the role of electric currents. Since the laws of electromagnetism are well studied and understood this analogy has proved quite fruitful in the *gravito-electromagnetic* context particularly in astrophysical applications. Extended "astrophysical jet-structures" are now thought to have their origin in *gravito-electromagnetic* forces.

In¹⁰ we have considered the *most useful way* to define the *gravito-electromagnetic* fields in terms of the perturbed components of the space-time metric. Different choices are often responsible for the location of odd factors of 4 that permeate the *gravito-electromagnetic* equations compared with the Maxwell equations. Such choices also have implications for the form of the induced *gravito-electromagnetic* Lorentz force (and torque) in terms of the *gravito-electromagnetic* fields that enter into the equation for the motion of a massive point (spinning) particle. By analogy with the covariant laws of electromagnetism in space-time *gravito-electromagnetic* potentials and fields are then defined to emulate electromagnetic gauge transformations under substitutions belonging to the gauge symmetry group of perturbative gravi-

tation. These definitions have the advantage that on a flat background, with the aid of a covariantly constant time-like vector field, a *subset* of the linearised gravitational field equations can be written in a form that is *fully analogous to Maxwell's equations* (without awkward factors of 4 and extraneous tensor fields). It is shown how the remaining equations in the perturbed gravitational system restrict the time dependence of solutions to these equations and thereby prohibit the existence of propagating vector fields. The induced *gravito-electromagnetic* Lorentz force on a test particle is evaluated by geodesic perturbation in terms of these fields together with the torque on a small gyroscope. It is concluded that the analogy of perturbative gravity to Maxwell's description of electromagnetism can be valuable for (quasi-) stationary gravitational phenomena but that the analogy has its limitations. The relevance of this work for propulsion is that the equation of motion for any massive particle with velocity \mathbf{v} in a pure *gravito-electromagnetic* field (\mathcal{E}, \mathcal{B}) takes the form

$$\frac{d\mathbf{v}}{dt} = \frac{1}{4}\mathcal{E} + \mathbf{v} \times \mathcal{B} \quad (14)$$

in close correspondence with the Lorentz force on electrically charged particles in the electromagnetic field (\mathbf{E}, \mathbf{B}). In the static field of the earth the term $\mathcal{E}/4$ is responsible for the Newtonian gravitational acceleration field. The accompanying gravito-magnetic force determined by \mathcal{B} on mass currents has to date escaped detection, but along with propagating gravitational waves is thought to exist. A gyroscope in such a field with velocity \mathbf{v} is predicted to precess at a rate $(\frac{3}{4}\mathbf{v} \times \mathcal{E} - \mathcal{B})/2$ independent of its spin. In 2002 the NASA satellite Gravity Probe B will be launched to measure this effect. Although such *gravito-electromagnetic* forces and torques are small in a terrestrial environment they can be made to accumulate and their detection could herald a new avenue of gravitational precision technology. In addition to the \mathcal{E} and \mathcal{B} *gravito-electromagnetic* fields the equations of Einsteinian gravitation involve a second degree symmetric tensor Σ that has no electromagnetic analogue. The *gravito-electromagnetic* fields couple to this tensor and produce non-Maxwellian terms in the *gravito-electromagnetic* force and torque equations. This suggests that the equations of motion for massive spinning particles in gravitational fields deserve closer scrutiny.

2.3 Orbiting Cosserat Detectors

We have begun an intensive study of the behaviour of slender visco-elastic structures in space.

Such structures can be used to convert tidal gravitational fields into heat and hence extract energy from gravitational fields.

They are also being studied with a view to using them as antennae for the detection of gravitational radiation. The orbital dynamics of slender loop structures, several km in length, are ideally suited to analysis by the simple theory of *Cosserat rods*¹¹. Such a description offers a clean conceptual separation of the vibrations induced by bending, shear, twist and extension and the coupling between eigen-modes due to tidal accelerations can be reliably estimated in terms of the constitutive properties of each structure.

The proposal is to use (several) orbiting Cosserat structures in order to detect gravitationally induced displacement correlations from among unwanted sonic signals in the system. Each continuum structure can be tuned to the entire acceleration field of a gravitational wave and resonant response to circumferential excitations tuned to optimise power absorption from continuous or pulsed gravitational excitations. Different topologies (e.g. open spirals) yield alternative broadband non-resonant responses.

The detection of gravitational signals in the 1 Hz region would provide vital information about stochastic backgrounds in the early Universe and the relevance of super-massive black holes to the processes that lead to processes in the centre of galaxies. Such information can only enhance our understanding of the interaction of gravity and matter and test the predictions of many proposed generalisations of Einstein's description of gravitation.

2.4 Scalar Fields and Mass Generation

In section (1) it was stated that the structure of the basic interactions was related to a pattern of symmetry groups. For gravity the symmetry group of relevance encodes the invariance of the speed of light in space-time and the behaviour of particles with mass and intrinsic spin. One of the great advances in recent years has been the recognition that all these symmetries arise as approximations from a description of extended structures in higher dimensions and that space-time is a classical approximation to a more general quantum description that unites all the fields shown in Figure 1. Although this picture is incomplete there are indications that Einstein's original description of pure gravitation may need supplementing with additional fields. In particular, remnants of a higher symmetry group may persist on scales accessible to experiment. In this regime additional scalar fields

seem inevitable. The role of such scalar fields can modify the short range behaviour of gravitation. It is therefore of importance to explore whether such fields can be detected experimentally and assess their significance on the behaviour of other matter.

One approach is to examine the manner in which residual symmetries for gravity are broken. We have recently shown¹² that the breakdown of local Weyl symmetry in a theory of gravity can be accommodated in the context of the standard model of particle interactions. A natural setting for this mechanism is a space-time geometry described by a connection with dynamical torsion and a metric that is not covariantly constant. Together with a scalar field such a connection encodes new gravitational interactions that can be reformulated in terms of the standard description of Einsteinian gravity. The emergence of space-time torsion, dependent on the gradient of the dynamic scalar field, is responsible for the appearance of the so-called *improved stress-energy tensor*. In the broken phase in which electro-weak phenomenology is discussed the theory gives rise to a Higgs particle with mass M_η and a new electrically neutral vector boson with mass M_Q such that

$$\frac{M_\eta^2}{M_Q^2} = \frac{8\lambda v^2}{\gamma}$$

in terms of the couplings in the theory. It is of interest to note that a number of grand unified models predict a new neutral vector boson and experimental data are now detailed enough to check for its existence. It appears that such data are better described if the presence of such a boson is assumed. The theory above has been analysed in a broken phase in which normal gravitation (based on metric perturbations about Minkowski space-time) is negligible. The mass generation mechanism has been connected with a component of non-Einsteinian gravitation associated with the Weyl connection Q . This forms part of the natural space-time geometry determined from an action principle and may be expected to give rise to new kinds of force on matter. If the neutral boson described by the excitation of the Weyl potential Q in Minkowski space-time can be observed in current electro-weak data it may signal that a new component of gravitation can influence phenomenology at energies well above the Planck scale.

2.5 Scalar Fields and Torsion Forces

In the scalar-tensor theory of gravitation formulated by Brans and Dicke¹³, the motion of a test particle was originally assumed to be a "Levi-

Civita geodesic" associated with the metric derived from the Brans-Dicke field equations (even though the scalar field could vary in space-time). Later Dirac in ¹⁵ showed that in a Weyl invariant generalisation it was more natural to generate the motion of a test particle from a Weyl invariant action principle and that such a motion in general differed from a Brans-Dicke Levi-Civita geodesic. Although Dirac was concerned with the identification of electromagnetism with aspects of Weyl geometry even for neutral test particles it turns out that test particles would follow "auto-parallels" of a connection with torsion. We have shown¹⁶ that the theory of Brans and Dicke¹³ can be reformulated as a field theory on a space-time with dynamic torsion determined by the gradient of the Brans-Dicke scalar field Φ . Of course no new physics of the fields can arise from such a reformulation, although it does clarify certain issues relating to the conformal structure of the theory and its couplings to matter with intrinsic spin. However, the behaviour of spin-less particles in such a geometry with torsion depends on the choice made from two possible alternatives. One may assert that their histories are either geodesics associated with auto-parallels of the Levi-Civita connection or the auto-parallels of the non-Riemannian connection with torsion. Since one may find a spherically symmetric, static solution to the Brans-Dicke theory (in either formulation), it is possible to compare these alternatives for the history of Mercury about the Sun by regarding it as a spin-less test particle as in General Relativity.

To summarise the relevant equations of motion the language of covariant differentiation is appropriate. Differentiation involves a limiting procedure in which objects are compared at the same point. In ordinary calculus space has an "affine vector space structure" and there is a natural parallelism which permits vectors at different points to be transported to common points for comparison. Gravitation is formulated on a more general manifold. A manifold does not in general have a vector space structure and such a "natural parallelism" is absent. A new structure[†] called a "space-time connection", ∇ , is required to trans-

[†]Consider a curve C with tangent vector \dot{C} and an arbitrary vector field X . An affine connection at a point p in space-time is a map that takes the pair (\dot{C}, X) to a new vector $\nabla_{\dot{C}}X$ at p , which is linear in \dot{C} , and satisfies

$$\nabla_{\dot{C}}(X + Y) = \nabla_{\dot{C}}X + \nabla_{\dot{C}}Y$$

$$\nabla_{\dot{C}}(\phi X) = \dot{C}(\phi)X + \phi \nabla_{\dot{C}}X$$

for all scalar fields ϕ and vector fields X, Y . Due to the \dot{C} linearity $(\nabla X)_p$ is a $(1,1)$ tensor at p with $(\nabla X)_p(\dot{C}, \alpha_p) = \alpha_p(\nabla_{\dot{C}}X)$ for all covectors α_p at p . It is called the covariant differential of X at p .

port vectors from one point in the manifold to another, along an arbitrary curve.

The equation for a time-like auto-parallel is

$$\nabla_{\dot{C}}\dot{C} = 0$$

where the 4-velocity \dot{C} is normalised with respect to the space-time metric g :

$$g(\dot{C}, \dot{C}) = -c^2. \quad (15)$$

By expressing ∇ in terms of the Levi-Civita connection $\tilde{\nabla}$ with $\tilde{V} = g(V, -)$ for any vector V one may write this as

$$\widetilde{\nabla}_{\dot{C}}\dot{C} = -\frac{1}{2\Phi}i_{\dot{C}}(d\Phi \wedge \tilde{C})$$

(the operator $i_{\dot{C}}$ denotes contraction of the 2-form with the vector \dot{C}) and interpret the right hand side as a torsion acceleration field analogous to the Lorentz force on electrically charged particles. Note however that the torsion force produces the same acceleration on all massive test particles. If \dot{C} is parameterised in terms of proper time τ in any coordinates $x^\mu(\tau)$, the above is:

$$\begin{aligned} \frac{d}{d\tau} \left(\Phi^{1/2} \frac{dx^\mu}{d\tau} \right) + \Phi^{1/2} \{^\mu_{\nu\lambda}\} \frac{dx^\nu}{d\tau} \frac{dx^\lambda}{d\tau} \\ = -g^{\mu\nu} \frac{\partial_\nu \Phi}{2\Phi^{1/2}}. \end{aligned} \quad (16)$$

To illustrate the effects of this equation of motion we have recomputed¹⁷ the classical shift in the perihelion rate of Mercury's orbit about the Sun in terms of a static spherically symmetric solution of the vacuum Brans-Dicke field equations ¹³. Taking into account that the speed of Mercury is non-relativistic and that its Newtonian orbit is much larger than the Schwarzschild radius $r_s = 2GM/c^2$ of the Sun, one finds the perihelion shift per revolution Δ of the orbit:

$$\Delta = \frac{3\omega + 5}{3\omega + 6} \delta_\omega \quad (17)$$

where $\delta_\omega = 3\lambda_\omega\pi$, $\lambda_\omega = r_s/\hat{r}_0$ and ω is the Brans-Dicke coupling parameter. Using the Kepler period

$$\hat{T} = 2\pi(\hat{r}_0^3/(1-\epsilon^2)GM)^{1/2},$$

the shift δ_ω may be expressed in terms of \hat{T} and ϵ . This may be compared with the result based on the assumption that Mercury's orbit follows from a geodesic of the torsion-free Levi-Civita connection. In this case one finds

$$\Delta = \frac{3\omega + 4}{3\omega + 6} \delta \quad (18)$$

where $\delta = 3\lambda\pi$ (with $\lambda = r_s/r_0$) is the perihelion shift per revolution of the orbit based on the Schwarzschild solution for the metric in General Relativity.




Figure 4: Behaviour of Δ/δ as a function of the Brans-Dicke coupling parameter ω . The full curve corresponds to a precession rate of Mercury's orbit under both metric and torsional acceleration. The lower dotted line corresponds to the original prediction of the Brans-Dicke theory. The experimental observations are consistent with Δ/δ lying between the full horizontal lines centered on $\Delta/\delta = 1$.

Given the prominent role played by the motion of test particles in many astrophysical phenomena and the possibility of new gravitational interactions mediated by scalar fields we feel that, with the enhanced technology now available to modern space science, the possible relevance of scalar field induced space-time torsion should not be ignored. Should the departure of spin-less particles from Levi-Civita geodesic motion be detected in some purely gravitational environment it would indicate that matter has additional "gravitational charge" in addition to its mass and to its electromagnetic, weak and strong couplings to other fields in nature.

3 Summary

This article summarises some of the current activities of the gravitational research group above. Having surveyed the role of gauge fields as carriers of the basic forces in physics and the momentum conservation constraints placed on cur-

rent propulsion technology a number of alternative strategies, under active investigation, have been briefly discussed. The essential theme is the detection of hitherto undetected components of gravitation that may have implications for the motion of matter. Such components include expected configurations of Einsteinian gravitation (including gravito-magnetic and radiation modes) as well as new scalar interactions resulting from various proposals for the unification of gravitation with the other basic forces. It would be premature to rank these strategies in terms of potential for the advancement of propulsion technology. International efforts are well advanced for the detection of gravitational radiation and the STEP experiment¹⁸ is expected to place firm experimental bounds on the foundations of General Relativity. We have suggested above that there are also less expensive experiments that may also shed light on the existence of novel gravitational fields and that if scalar field gravitational interactions with matter can be detected they may offer unsuspected avenues for development.

4 Acknowledgement

The authors are grateful to BAe-Systems for the support of this research and to Dr R Evans (BAe-Systems UK) for stimulating discussions.

References

- [1] W Rindler, *Introduction to Special Relativity*, Clarendon Press, Oxford,
- [2] V V Beletsky, M E Givertz, *On the motion of a Pulsating Dumbbell in the Gravitational Field*, Kosmicheskie Issledovaniya **6** 304-306 (1968) (in Russian)
- [3] V V Beletsky, *Essays on Motions of Space Bodies*, Moscow, Nauka 1997 (in Russian)
- [4] V V Beletsky, E M Levein, *Dynamics of Space Tether Systems*, Advances in the Astronautical Sciences, Vol 83, American Astronautical Society Publications 1993
- [5] NASA/AIAA/PSN International Conference on Tethers, Arlington, VA. USA. 1986 , Advances in the Astronautical Sciences, Vol 62, American Astronautical Society Publications 1986. Eds. P M Bainum et al.
- [6] S Kalantzis, V J Modi, A K Misra, S Pradhan, *Dynamics and Control of Multibody Tethered Systems*, Acta Astronautica **42** (9) 503-517 (1998)

- [7] K Kumar, K D Kumar, *Variable Attitude Maneuver via Tethers for a 'Drifting' Twin Satellite System in Elliptic Orbits*, Acta Astronautica **45** (3) 135-142 (1999)
- [8] J Ashenberg, E C Lorenzini, *Active Gravity-Gradient Stabilization of a Satellite in Elliptic Orbits*, Acta Astronautica **45** (10) 619-627
- [9] J Gratus, R W Tucker. An Improved Method of Gravicraft Propulsion, Submitted for Publ. 2000.
- [10] S Clark, R W Tucker Gauge Symmetry and Gravito-electromagnetism Class. Quantum Grav. **17** 4125-4157 (2000)
- [11] R Tucker, C Wang, A Cosserat Detector for Dynamic Geometry, International Conference on Geometrical Methods in Continuum Mechanics, University of Torino, 2000.
- [12] T Dereli, R W Tucker, A Broken Gauge Approach to Gravitational Mass and Charge, Lancaster University Preprint, Submitted for Publ. 2000.
- [13] C H Brans, R Dicke, Phys.Rev. **124** 925 (1961)
- [14] A Einstein, L Infeld, B Hoffmann, Ann. Math. **39** 65 (1938)
- [15] P A M Dirac, Proc. Roy. Soc **A333** 403 (1973)
- [16] T Dereli, R W Tucker, Phys. Letts **110** 206 (1982)
- [17] T Dereli, R W Tucker, On the Detection of Scalar Field Induced Spacetime Torsion, gr-qc/0104050.
- [18] R Reinhard, Y Jafry, R Laurance, STEP-A Satellite Test of the Equivalence Principle, ESA Journal-European Space Agency **17**: (3) 251-263 (1993)



A01-34542

**AIAA 2001-3913
An Asymmetric Gravitational Wave
Propulsion System**

**J. Cameron
Huntsville, AL**

**Joint Propulsion Conference
8-11 July 2001
Salt Lake City, Utah**

**For permission to copy or to republish, contact the copyright owner named on the first page.
For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.**

AIAA 2001-4335

AN ASYMMETRIC GRAVITATIONAL WAVE PROPULSION SYSTEM

Jeffery A. Cameron, member, Chief Scientist

Transdimensional Technologies, Inc., Huntsville, Alabama 35801

Abstract

Gravitational wave radiation is generated by the quadrupole moment of matter that is in motion. An analytical model is constructed to investigate the radiation pattern and radiated power of a single resonant vibrator cylinder, as compared to that of a phased linear array. The linear array is then evaluated in terms of phase relationship in order to create an asymmetry in the radiated pattern and hence a directional force. This is compared to the force of a solar sail in the earth orbital plane.

Introduction

Einstein¹ derived the weak-field solution of the gravitational wave in accordance with the general and special theories of relativity. This paper uses this scientific background to describe a revolutionary new approach to propulsion. Programmable laser diodes in conjunction with semiconductor materials will be used to generate a highly directional transverse wave gravitational wave (TWGW) radiator. This asymmetric TWGW radiator will create a directional force through the center of mass of the radiating system, thus forming a propellantless propulsion system.

Theoretical work indicates the ability to generate gravitational wave radiation through the quadrupole moment of matter under stress and strain. The radiation pattern is symmetric about the center of mass, and the direction of the pattern is at right angles to the stress and strain vector. The radiation pattern looks like a torus. The radiated power is very small (10^{-29} watts); however, when a linear sequence of radiators is put together and their patterns are allowed to superimpose, the total radiated power can approach kilowatts. The asymmetry of the radiated pattern is produced by adjusting the phase of the radiators. The resulting power imbalance will produce a force through the center of mass of the radiators. In order to accomplish this task, high peak power laser diodes will be used to photo-acoustically drive thin-film resonators. It becomes

imperative to understand the energy flow between the laser diode driving the resonator and the result in gravitational radiation from the radiator.

Theory

In order to establish an analytical background consider the following. Let $\eta_{\mu\nu}$ be a Lorentz metric, then the Riemannian metric is expressed as

$$g_{\mu\nu} = \eta_{\mu\nu} \sqrt{K} h_{\mu\nu} \quad (1)$$

as a first approximation under the weak-field assumptions, K is Einstein's constant. The potential of the field can be expressed as

$$\phi_{\mu\nu} = h_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} h \quad (2)$$

resulting in the form for 4-space

$$\square \phi^{\alpha\beta} = 2\sqrt{K} T^{\alpha\beta} \quad (3)$$

with retarded potential solutions of the form

$$\phi^{\mu\nu}(x) = (\sqrt{K}/2\pi) \int T^{\mu\nu}(x', x^0) \quad (4)$$

This form will enable the definition of the energy-momentum complex of the gravitational field in order to evaluate the radiation energy and directivity of the gravitational wave.

The Poynting vector of the gravitational wave can be expressed as

$$U_o^l = \frac{1}{8} (-2\phi^{\rho\sigma,l}\phi_{\rho\sigma,o} + \phi^l\phi_{,o}) \quad (5)$$

The derivatives of the potential fields ϕ with respect to time and space coordinates are expressed by the second and third derivatives of the mass tensor,

$$mc^2 \equiv \iiint T^{00}(x^1, x_G^0) d^3x^1 \quad (6)$$

momentum tensor,

$$cP^K = \delta_0 x^K \iiint T^{00}(x', x_G^0) d^3 x' \quad (7)$$

and the quadruple moment tensor,

$$I^K = x' x^K \iiint T_{00}(x', x_G^0) d^3 x' \quad (8)$$

The derivative forms for the potential fields are substituted into the expression for the Poynting vector, thus giving the expression for the radiated energy per unit time, or power, within a solid angle $d\Omega$, as follows

$$P_o^t = U_o^t d\sigma_t = \frac{1}{8} \left(\frac{\sqrt{K}}{4\pi} \right)^2 f(\theta, \phi, x_G^0) d\Omega \quad (9)$$

Here the factor

$$f(\theta, \phi, x_G^0) \quad (10)$$

represents the directivity of the gravitational wave radiator.

Of particular interest for this paper is the resonance vibrator, conceptually similar to what Weber^{2,3,4} used as a gravitational wave detector. The resonance vibrator is a cylinder (Figure 1), which is placed under stress and strain.




Figure 1: Resonant Vibrator Cylinder

At this point assume the following dynamic variables:

$$\text{Displacement } \varepsilon = A_l \sin\left(\frac{\omega}{V_s}\right) \cos \omega t \quad (11a)$$

$$\text{Particle velocity } v = \frac{d\xi}{dt} = -V_p \sin\left(\frac{\omega}{V_s}\right) z \sin \omega t \quad (11b)$$

$$\text{Strain } \varepsilon = \frac{d\xi}{dz} = \frac{V_p}{V_s} \cos\left(\frac{\omega}{V_s}\right) z \cos \omega t \quad (11c)$$

$$V_p = A_l \omega \quad V_s = \left(\frac{B_s}{\rho}\right)^{\frac{1}{2}} \quad (11d)$$

$B_s \rightarrow$ Young's Modulus

The directivity can now be expressed as follows (Figure 2).

$$f \sim \sin^4 \theta \quad (12)$$




Figure 2: Resonant Vibrator/Gravitational Wave Radiation

The pattern resembles a torus or "donut" mode.

It can be seen that the maximum radiation occurs in the plane perpendicular to the vibrating z axis (Figure 2).

Consider a linear array of resonant vibrators (Figure 3).




Figure 3: Resonant Vibrator Linear Array

Let high peak power laser radiation from laser diodes be injected along the z axis to induce acoustic stress in the material (Figure 4).




Figure 4: Laser Diodes and Linear Array

The stresses will generate a weak gravitational wave along the x axis. The gravitational wave generated from a number of "cells" along the x axis can be added in phase. The resultant gravitational "beam" along the x axis is extremely intensified compared to a single resonant vibrator "cell." The linear arrangement will be referred to as a traveling wave (TW)gravitational wave (GW) radiator, or TWGW radiator.

The directivity of the TWGW radiator can be expressed as (Figure 5).

$$f \approx \left(\frac{\sin(\frac{\pi\omega}{V_s}(1-\cos\Theta))}{(\frac{\pi\omega}{V_s})(1-\cos\Theta)} \right)^2 \cos^4 \Theta \quad (13)$$




Figure 5: TWGW Radiator Symmetric Radiation Pattern

The graphical result of Figure 5 is based on the geometry of Figure 3 and the constants as follows.

Material: Quartz

$$b = 3 \times 10^{-5} m$$

$$a = 3 \times 10^{-3} m$$

$$\ell = 0.3 m$$

$$\omega = 6.28 \times 10^{12} \text{ radian / sec}$$

(resonant frequency)

$$V_s = 5 \times 10^3 m / \text{sec} (\text{wave speed})$$

$$\rho_o = 2650 kg / m^3 (\text{density})$$

As can be seen in Figure 5, the TWGW radiation is very directional. This brings to question what type of power levels could be estimated. To begin, consider the single resonant vibrator cell (Figures 2 and 3). The radiated power can be estimated by the following expression.

$$P_{RV} = (5.7 \times 10^{-2}) G \rho_o^2 \pi^2 a^4 e_m^2 V_s \left(\frac{V_s}{C} \right)^5 \quad (14)$$

Where

$$G = 6.67 \times 10^{-11} \frac{Nt \cdot m^2}{kg}$$

(Gravitational Constant)

$$\epsilon_M = 4 \times 10^{-3} (\text{Strain})$$

Then

$$P_{RV} = 2.14 \times 10^{-29} \text{ watt} \quad (15)$$

As anticipated, this is a very small number. However, for the TWGW radiator, the radiated power can be estimated by the following expression.

$$P_{TW} = (7 \times 10^{-5}) G \rho_o^2 a^2 \ell^2 e_m^2 V_s \left(\frac{V_s}{C} \right)^5 \left(\frac{b\omega}{V_s} \right)^6 \Omega W \quad (16)$$

here

$$\Omega(\pm 5.3^\circ) = 2.7 \times 10^{-2} \text{ sr}$$

(Solid angle of the radiation)

Then

$$P_{TW} = 1.66 \times 10^3 \text{ watt} \quad (17)$$

This shows how important the phase relationships between individual resonant vibrators are!

At this point it is important to realize that the radiated gravitational wave carries energy and momentum with it. This is expressed as

$$P = \frac{\epsilon}{C} = \frac{Pt}{C} \quad (18)$$

Where P is the radiated power, t is time, ϵ is the energy of the gravitational wave, and P is the momentum.

The resulting reaction force on the TWGW structure is expressed as

$$F = \frac{P}{t} = \frac{P}{C} \quad (19)$$

Referring to Figure 5, it can be seen that the radiation pattern is symmetric about the center of the TWGW structure. Therefore any reaction force is balanced! However, consider a variation in the phasing of the laser diodes (Figure 4), where the lobes become asymmetric (Figure 6).




Figure 6: TWGW Radiator Asymmetric Radiation Pattern

This can be accomplished by pulse timing, variation in pulse rate, laser diode intensity, alternate materials, and geometry. Here there is a net force in the direction of the least intense gravitational wave (Figure 7).




Figure 7: Reaction Force for Asymmetric Radiator TWGW

The reaction force is expressed as

$$\begin{aligned} F_{net} &= F_{RL1} - F_{RL2} \\ &= \frac{P_{RL1} - P_{RL2}}{C} \end{aligned} \quad (20)$$

As an example, let $P_{RL1} = 500$ watt and $P_{RL2} = 3.5 \times 10^3$ watt, then

The net force, although small, raises the possibility of a propellantless propulsion concept, utilizing current technologies.

$$F_{net} = 10 \times 10^{-6} Nt \quad (21)$$

Further Investigations

Mathematical analysis indicates that gravitational wave propulsion is possible. Further investigation will entail the following objectives.

- 1) Investigate the efficiency of converting laser light into acoustic stress within TWGW generating material candidates. The acoustic stresses are responsible for generating the quadrupole moment needed to give rise to TWGW radiation. It will be imperative to address questions on how material uniformity, temperature, geometry, laser pulse width, repetition rate, and wavelength will impact performance.
- 2) Investigate phasing techniques of the TWGW elements in order to create various asymmetries in the radiating pattern. The directivity and intensity of the TWGW radiator is key to the success of generating a net propulsive force. How energy and momentum transports are affected by individual radiator phasing are of utmost importance. The question of whether harmonics distribute energy into other "modes" or are negligible must be considered.

The TWGW system must be analytically modeled as part of the above investigations. The results will lead to study of a test article that will be used to demonstrate the use of the gravitational wave Poynting vector imbalance as a means of generating a propulsive force through the center of mass of the TWGW system. Conceptually a scale version of the system could provide propellantless propulsion into the outer regions of the solar system and/or orbital transfer missions.

Space Flight Application

An interesting example is a comparison of the TWGW system to that of a solar sail at the earth orbital plane. Let the TWGW system be 1000 meters in length. The resulting net propulsive force would be about 61 Nt. For a square solar sail with a perfect reflectivity experiencing a solar flux of 1.3×10^3 watts/m², the required area would be around 1.4×10^7 m² for a force of 61 Nt. This is a perimeter length of approximately 3.7×10^3 m and must be normal to the solar disk to experience the maximum momentum transfer. Also, as the sail increases its distance from the solar disk, the intercepted flux decreases with the square of the distance. The TWGW system is not dependent on the solar disk for operation.

Conclusion

An analytical model has been created to investigate the nature of the radiated TWGW pattern with respect to phase relationships between individual "cell" radiators. Future models will investigate the magnitude of side lobes or harmonics to determine whether asymmetries cause less than desirable effects. These results will be used for evaluation of the generation of a net propulsive force.

References

¹ Einstein, A., *Naherungsweise Integration der Feldgleichungen der Gravitation* (Berlin Sitzungsberichte, 1916), pp. 688-696.

² Weber, J., *Physics Review*, 117, 1960, p. 306.

³ Weber, J. *General Relativity and Gravitational Waves* (Interscience, New York, 1961).

⁴ Weber, J., *Physics Review Letters*, 22, 1969, p. 1320.

Acknowledgements

The author wishes to express sincere thanks to Remigius Shatas and Robert Asprey of 2C Computing, Huntsville, Alabama for their encouragement and support.



A99-31034

AIAA 99-2147

**Fabrication of Large Bulk Ceramic Superconductor
Disks for Gravity Modification Experiments and
Performance of YBCO Disks Under e.m. Field
Excitation**

**R. Koczor and D. Noever
NASA Marshall
Huntsville, AL**

**35th AIAA/ASME/SAE/ASEE Joint Propulsion
Conference and Exhibit
20-24 June 1999
Los Angeles, California**

ABSTRACT

We have previously reported on investigations pertaining to the measurements of gravitational field changes in the vicinity of a large rotating Type II superconducting disk. Published reports indicate that test masses have been observed to lose up to 2% of their weight when in the vicinity of such rotating disks. We have produced 30 cm (nominal) diameter disks using $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ and BiSCCO. In addition we have performed tests on the interactions between smaller YBCO disks and AC levitation fields and RF fields, using a sensitive gravimeter to monitor and record the local gravity field above the superconductors. We have not yet duplicated the reported experimental protocols, including rotation and levitation of a large two-layered YBCO disk. Static disk experiments to date have uncovered no measurable and repeatable gravity effect for the conditions tested to within the sensitivity of the gravimeter (which has a noise level on the order of 10 nano-G or less).

FABRICATION OF LARGE BULK CERAMIC SUPERCONDUCTOR DISKS FOR GRAVITY MODIFICATION EXPERIMENTS AND PERFORMANCE OF YBCO DISKS UNDER EM FIELD EXCITATION

Ronald J. Koczor, NASA, Marshall Space Flight Center, AL,
David A. Noever, NASA, Marshall Space Flight Center, AL

Background

Attempts to unify theories of gravity with the other forces of nature have a long and checkered history. Some of the greatest minds the world of physics has known have believed that the connection is there. Clearly at the astronomical scale, gravity and electromagnetics interact through the mechanism of gravitational warping of space around massive bodies. Nonetheless, all credible attempts to experimentally demonstrate that connection on a laboratory scale have remained elusive. It is clear that if a manipulative method is found to treat gravity, extensive changes would occur in our everyday life, just as our lives changed once electromagnetism was understood.

In 1992, E. Podkletnov and researchers at the University of Tampere, Finland, reported¹ demonstration of what he termed "gravity shielding."

The experiment consisted of spinning a Type 2 superconductor (yttrium, barium, copper oxide, YBCO) in the presence of electromagnetic levitation and rotation fields and measuring weight changes in test masses suspended above the spinning disk. The effect reported was small, varying from tenths of a percent to as much as 2% of the test mass' weight. A subsequent paper was distributed giving more details of the experiment.²

While this is a small effect, it can be recalled that some of the most telling and seminal experiments in electromagnetics in the early 19th century were similarly of small magnitude. Their value was in demonstrating the basic phenomenon; useful amplifications and extensions came from others, once the basic idea had been demonstrated and accepted.

In 1997 researchers at the Marshall Space Flight Center, Huntsville, Alabama, and colleagues began to investigate the Podkletnov experiment. The goal was to understand various phenomena related to the experiment and to replicate it. Marshall has a historical connection to the YBCO material, having been intimately involved in the first fabrication and characterization experiments over a decade ago.

While there is general information available about the experimental set up, several critical

"Copyright c 1999 by the American Institute of Aeronautics and Astronautics, Inc. No copyright is asserted in the United States under Title 17, U.S. Code. The US Government has a royalty-free license to exercise all rights under the copyright claimed herein for Government Purposes. All other rights are reserved by the copyright owner."

details are unclear. However, as we have reported,^{3,4,5} progress is being made and our understanding of bulk ceramic superconductors and their interactions with various electromagnetic fields grows. Also, a significant step forward in the measurement of small changes in gravity fields was made with the application of a highly sensitive and stable gravimeter to the experiment. This gravimeter is capable of measuring changes in the local gravity field as small as 5×10^{-9} G. In this paper we will discuss our progress in fabricating large bulk superconductors and various preliminary experiments to understand how these materials interact with AC fields levitation.

Large Bulk YBCO and BiSCCO Disks

The central component in the Podkletnov experiments is a composite structure annular disk of YBCO superconductor. The YBCO material normally exhibits a superconducting transition temperature of around 94K, making it an exciting superconducting material since it is useable with liquid nitrogen cooling. Podkletnov fabricated a dual layered disk, having one layer of YBCO that became superconducting around 94K and another layer that became superconducting around 60K. It is his belief that this composite structure is a mandatory feature to see the effect. Our results to date do not dispute that assertion.

While the ceramic superconductors offer much potential, their well-known characteristics have led present day research with YBCO from bulk to thin film configurations. There are no published reports of fabrication of large bulk ceramic superconductors, in part due to the enormous challenges facing anyone who attempts to fabricate them. Since it is central to our goals, we undertook to fabricate such a disk. Working with colleagues at the University of Alabama in Huntsville and at Martin Technologies, Inc. (also of Huntsville), and after much trial and error, we succeeded in fabricating several single layer YBCO disks (on the order of 30 cm in diameter and ranging in thickness from 0.5 to 1 cm). Recently with our colleagues at Tomorrowtools, Huntsville, we fabricated a 32 cm diameter annular disk made from BiSCCO (bismuth, strontium, calcium, copper, oxide) superconducting material with a nominal superconducting transition temperature between 100 and 110 K. Figure 1 shows the YBCO disk (left) and the BiSCCO disk (right) fabricated at Marshall.

Successful fabrication of such large superconductors resulted from attention to 3 process steps, material selection, mechanical pressing, and thermal processing. Details of the process are being

reviewed for patent applicability and so cannot yet be discussed. However, the process results in disks having excellent mechanical structure and moderate superconductive performance as expected from sintered bulk materials. Characterization of the resulting disks indicates transition temperatures for YBCO in the 92K to 94K range and critical currents on the order of 20 to 100 amps/cm². The BiSCCO disk is presently undergoing characterization and results are not available.

Fabrication of the YBCO disk used pre-processed YBCO superconducting powders; The process used for the BiSCCO disk was based upon the sol-gel process prepared with precursor nitrates in an autoignition process. Subsequent pressing and heat treating created the correct BiSCCO superconducting formulation.

Only one em field test was run using these large disks, as we are presently completing their characterization. This test involved a preliminary attempt to induce radio frequency energy into the disk and measure any resultant gravity field changes. This test is reported below. Other experiments have been underway using smaller, commercially available YBCO disks to characterize their interactions with various electromagnetic fields. These experiments are also reported below.

Small Disk Interactions with AC Levitation Fields

One series of tests was run to determine how the superconductors react to alternating current, AC, levitation fields. It is generally known that these materials react differently to AC fields than DC. And, again, while much information is available for thin films and very small bulk samples, we believe these tests are the first reported for large pieces.

The experiments were performed using two commercially available melt-textured crystal seeded YBCO disks. One was a single layer disk having a diameter of about 13 cm and a thickness of about 1 cm. The other disk was a 10 cm diameter two layer disk. The levitation field was generated by a 3 solenoid assembly driven by Elgar model 1001SL variable voltage, variable frequency power supplies with a Model 9023 variable frequency oscillator for frequency control. These supplies are capable of generating up to 30 amps from 50 Hz to 4500Hz. Magnetic field characteristics were measured using an F.W. Bell model 9500 gaussmeter. The experimental protocol was to cool the superconductor, place it over the solenoids and turn on the field.

Results show that the 13 cm diameter single layered disk levitated with an 860g field at 45 Hz, 750g at 300 Hz, and 650g at 600 Hz. Levitation was

not achieved at higher frequencies due to the inductive reactance of the solenoids and the subsequent current limitations in them. Another test revealed that the 10 cm two-layer disk levitated with 600 g at 45 Hz.

Given these data and the fact that a sintered 30 cm disk will be both heavier and have lower critical current than the melt textured disks above, we expect significantly stronger fields will be required to levitate the large disks. On the other hand, the coils had small field cross sections. Larger field cross sections as well as stronger fields will assist levitation.

For one series of tests, we placed the levitating disk approximately 1 meter below the gravimeter. The disk and levitation fields were in a Faraday cage with the gravimeter above and outside the cage. Tests without the disk in place demonstrated that the gravimeter reading was not influenced by the levitation fields. For this test a 45 Hz field was applied to the disk using the 3-solenoid system described above for levitation while the gravimeter was recording. Figure 2 gives the results for one run using the 10 cm two layered disk. Each division on the vertical scale is 10 nano-G; each division on the horizontal scale is 72 seconds in elapsed time. In all cases reported here, the gravimeter is set for a 2 second sample integration time. The results of this test show no evidence of any effect of the levitating disk on the gravimeter above the noise level of the gravimeter. Other frequencies were run with similar results. At higher frequencies, fields were applied even though they were insufficient to levitate the disk. In some cases changes in the gravity reading are seen; however, they are all on the order of a 100 nano-Gs or less and are not repeatable.

Figure 3 (having the same scale as Figure 2) exhibits data from a run in which the 10 cm composite structure disk was exposed to 9 pulses of 45 Hz field. For this test field intensity was 1100 gauss and the disk did levitate with each pulse. Pulse rate was one pulse per minute with a 50% duty cycle. As can be seen, there is no significant change visible in the gravimeter reading during the experiment although there are minor transients that may be related to the pulses. The experimental setup was as described above.

It has been suggested that when a superconductor transitions from normal to superconducting mode, a possible transient gravity signal is produced. Figure 4 exhibits data from a run in which gravimeter readings were taken during cooldown of the 2 layer disk using the experimental setup described above. The figure shows where the liquid nitrogen was first poured and when rapid

boiling of the liquid stopped. A small magnet levitated when placed over the disk at the cessation of boiling, indicating that the disk was superconducting.

In summary, no conditions were found in which excitation of a small YBCO disk by a low frequency levitation signal produced repeatable changes in the gravimeter reading.

Large YBCO disk RF Field Interactions

A series of experiments was run to characterize the effects of megahertz-range rf on a 30 cm YBCO disk. The experiments used a tunable rf generator and amplifier capable of 100 watts output between 1 and 100 MHz. Most tests focused on the 3 to 20 MHz range. Rf was applied through a 12 turn coil wound toroidally around one section of the disk. An impedance matching network was used to maintain a constant load to the amplifier.

Initial tests demonstrated that the gravimeter was susceptible to the rf. This susceptibility manifested itself in lockup of the gravimeter data system at high rf levels (25 watts or more). This susceptibility was eliminated using standard rf shielding procedures on the gravimeter and by enclosing the YBCO disk and rf coil in a grounded Faraday cage.

Figure 5 is representative of these tests. In this case a series of four 3.5 MHz, 100 watt pulses were applied to the disk while it was superconducting. As can be seen, there are changes in the gravimeter reading appearing to coincide with the pulses, having amplitudes on the order of 10 nano-G.

Other tests performed include application of rf during cooldown of the disk and measurement of the gravity field with various rf frequencies applied. These data are presently being evaluated and will be reported later. In general, there are more repeatable and consistent gravimeter variations observed with the rf excitation than with the low frequency AC levitation excitation. However, all changes seen to date are less than 100 nano-G in amplitude.

Summary

We have succeeded in manufacturing large (30 cm diameter) superconducting disks using the YBCO and BiSCCO materials. These disks, weighing between 3 and 4 Kg, depending on thickness, are being used to investigate reported interactions between em fields and the local gravity field.

The reported interactions occur in an experimental protocol that we have not yet duplicated. Our initial attempts are to individually

characterize possible interactions under static (non-rotating) conditions. We have completed these initial characterizations, data is being analyzed, and preliminary results were reported here.

All observed interactions were on the order of nano-Gs and considerable care is being taken to assure that these observations are not artifacts of the experimental set up or interactions between the em fields and the gravimeter.

We are currently developing experimental hardware to test the large disks now being routinely fabricated. We expect to achieve levitation of the 30 cm disks under em field excitation later this year, with rotation soon to follow. Further experiments are also underway to optimize disk fabrication for increased critical current and mechanical stability and to develop a process to fabricate a two layered 30 cm disk.

The authors wish to acknowledge the indefatigable support provided by Tony Robertson, of MSFC's Space Transportation Directorate, Rick Roberson, of Tomorrowtools, Inc., Robert Hiser, of Maartin Technologies, Greg Jerman, of the MSFC Engineering Directorate, and Charles Sisk of the MSFC Science Directorate. We also wish to acknowledge and thank Ning Li and her colleagues at the University of Alabama in Huntsville, for their participation in the formative stages of this investigation.

References

1. "A possibility of gravitational force shielding by Bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ superconductor" E. Podkletnov and R. Nieminen, Physica C 203 (1992) 441-444.
2. "Weak gravitational shielding properties of composite bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ superconductor below 70K under an e.m. field" E. Podkletnov, MSU-chem-95-cond-mat/9701074 5 Feb 1997 preprint.
3. "Static test for a gravitational force coupled to type II YBCO superconductors," N. Li, et al, Physica C 281 (1997) 260-267.
4. "Superconductor-mediated Modifications of gravity? AC motor experiments with bulk YBCO disks in rotating magnetic fields," D. Noever, R. Koczor, AIAA 98-3139, Proceedings, 1998 AIAA/AASME/SAE/ASEE Joint Propulsion Conference, Cleveland, OH, July 13-15, 1998.
5. "Granular superconductors and gravity," D. Noever, R. Koczor, NASA/CP1999-208694, NASA Breakthrough Propulsion Physics Workshop Proceedings, August 12-14, 1998, Lewis Research Center, Cleveland, OH.




Figure 1 30 cm YBCO superconducting disk (left) and 32 cm BiSCCO superconducting disk (right) fabricated at NASA/MSFC.




Figure 4 Gravimeter reading during cool down of a 10 cm YBCO superconducting disk.




Figure 5 Gravimeter reading during rf excitation of a 30 cm YBCO superconducting disk.




Figure 2 Gravimeter reading with 600 gauss, 45 Hz levitation field applied to a 10 cm YBCO superconducting disk.




Figure 3 Gravimeter reading with 9 pulses of 1100 gauss, 45 Hz levitation field applied to 10 cm YBCO superconducting disk.



A99-31033

AIAA 99-2146

Large Scale Sakharov Condition

D. Noever and C. Bremner

NASA Marshall Space Flight Center

Huntsville, AL

**35th AIAA/ASME/SAE/ASEE Joint Propulsion
Conference and Exhibit
20–24 June 1999
Los Angeles, California**

ABSTRACT

Recent far-reaching theoretical results have used the quantum vacuum noise as a fundamental electromagnetic radiation field to derive a frequency (ω) -dependent version of Newton's gravitational coupling term, $G(\omega)$. This paper reconciles the cutoff frequency with the observed cosmological constant, then briefly puts forward a realizable laboratory test case in the 10-100 MHz frequency range. One analogy is drawn between the classical vacuum energy experiments with attraction between two closely spaced plates (Casimir cavity) and the arbitrarily dense material boundaries possible in Bose condensates, such as irradiation at MHz frequencies of superfluid helium or superconductors.

LARGE-SCALE SAKHAROV CONDITION

David Noever and Christopher Bremner

NASA Marshall Space Flight Center, Space Sciences Laboratory, Mail Code: ES76, Huntsville, AL 35812

Theoretical Background

Zel'dovich [1] first suggested that gravitational interactions could lead to a small disturbance in the (non-zero) quantum fluctuations of the vacuum and thus give rise to a finite value of Einstein's cosmological constant in agreement with astrophysical data. Using dimensional analysis and the suggestion by Zel'dovich, Sakharov [2] derived a value for Newton's gravitational constant, G , in only one free parameter, frequency, ω :

$$G \sim c^5/h \int \omega d\omega \sim 1/\int \omega d\omega$$

where c is the speed of light and h is the Planck constant. The free parameter in frequency when integrated over all values from zero to high frequencies, must contain the usual integration cutoff value (Planck frequency on observable electromagnetic phenomenon).

Puthoff [3] and others [4-5] have extended Sakharov's condition in a relativistically consistent model to determine

"Copyright 1999 by the American Institute of Aeronautics and Astronautics, Inc. No copyright is asserted in the United States under Title 17, US Code. The US government has a royalty-free license to exercise all rights under the copyright claimed herein for government purposes. All other rights are reserved by the copyright owner."

constants of proportionality. His model derives an acceleration term in first order expansion (in flat space-time), then equates inertial and gravitational mass (by the equivalence principle) to make contact with the gravitational constant, G , directly as:

$$G = (\pi c^5/h \omega_c^2) \sim 1/\int \omega d\omega$$

which is the Sakharov condition [2,3]. This paper revisits the meaning of the cutoff frequency, ω_c , for radiation interactions, of which the quantum vacuum [6-10] and Planck frequency are only the leading terms, and for which linear combinations of forces can introduce other plausible frequencies. One purpose of this reexamination is whether the resulting gravitational coupling constant, G , can be reconciled with the anticipated energy density of the universe [11] without resorting to extreme space-time curvature and thus yield enough critical density to contain the expansion of the universe. Finally we particularize the case to the high density fluctuations possible in Bose condensates [12], a potential experimental test case for how the effects of vacuum noise might manifest observably.

One far-reaching consequence of the vacuum energy model is the attractive force of gravity becomes reducible to the radiative interaction between oscillating charges, e.g. the zero point field (ZPF) applied to subatomic charges. Mass and inertia arise from the fundamentally electromagnetic ZPF oscillations.

This random background gives the usual quantum mechanical energy spectrum from particle-field effects:

$$\rho(\omega) d\omega \sim \omega^3 d\omega,$$

a very important dimensional relationship, since the third power in frequency avoids anomalous Doppler shifts from velocity boosts, or stated alternatively is the correct spectra for a Lorentzian (non-accelerated) invariant radiation field [13].

More specifically, the energy spectrum [3] can be written as:

$$\begin{aligned} \rho(\omega) d\omega &= [\omega^2 / \pi^2 c^3] [\hbar\omega/2] d\omega \\ &= \hbar\omega^3 / 2\pi^2 c^3 d\omega \sim \omega^3 d\omega \end{aligned}$$

which is an expression in the first parenthesis of the density of the normal modes and in the second parenthesis of the average energy per mode. When this energy density is integrated over all frequencies, the ω^3 divergence produces well-known infinities in the integration limit of high frequencies, thus an assumed cutoff frequency (appropriate to experimental observation limits at the Planck frequency), is usually introduced:

$$\omega_p = (c^5/\hbar G)^{1/2}$$

For mass, m , moving in an accelerated reference frame, $g = -a = Gm/r^2$, the resulting energy spectrum includes a gravitational spectral shift [3],

$\Delta\rho'(\omega) d\omega = \hbar\omega / 2\pi^2 c^5 [Gm/r^2]^2 d\omega \sim 1/r^4 d\omega$ a kind of short-range ($1/r^4$) gravitational energy shift, but electromagnetic in origin when zero point fluctuations are included. (N.B. To account for equal gravitational mass effects in neutrons and protons, the ZPF oscillations must involve subatomic charges, or ‘parton’ effects. The assumption derives from high-frequency interactions of ZPF wherein these subatomic particles are asymptotically free to oscillate as independent or free particles as quantum noise].

A further far-reaching consequence [3] is mass itself becomes interpretable as a dependent quantity derived from a damped (with decay constant, Γ) oscillation driven by random ZPF:

$$m = \Gamma c^3 / G = 2h \Gamma / \pi^2 c^3 \int \omega d\omega,$$

with the only two free parameters, the damping factor Γ , and again the frequency, ω . The internal kinetic energy of the system contributes to the effective mass.

This leads to an overall average spectral density, written in terms of mass as:

$$\Delta\rho'(\omega) = m^2 c^5 \omega / 2h\omega_c^4 r^4$$

for the electromagnetic field distribution near ($1/r^4$) to the mass, m , which in detail is half electric and half magnetic.

One additionally attractive feature is the correspondence between this derivation and the view of gravity as a dynamical scale-invariance-breaking model (e.g. symmetry breaking near the Planck mass energy [14]). A final result includes the force calculation between two ZPF radiation oscillators, of the correct form yielding Newton’s average force law:

$$\langle F \rangle = -Gm^2/r^2$$

Thus, for a Newtonian force to first order in a flat space-time, Sakharov [2] could be credited for proposing gravity as not a fundamentally separate force and Puthoff [3] and co-workers [4-5] applied the vacuum electromagnetic field to equate gravity to a long-range radiation force (e.g. van der Waals-like force). Higher order oscillatory gravity modes vary as $(\sin[\omega_\alpha/\omega_c])^2$.

To first order, a weak G coupling constant, $G = (\pi c^5 / \hbar\omega_c^2)$, appears for high frequency cutoffs at the Planck scale. A corollary in analogy to electromagnetic shielding by ordinary matter can be rationalized as the problem of frequency mismatch at high Planck frequencies, e.g. ZPF cannot be fundamentally shielded. In other words, frequency mismatch precludes gravity shielding by matter.

The purpose here is to revisit the only free parameter, the frequency cutoff, more in the spirit of a mass resonant frequency. The motivation for this approach can be summarized as: 1) the generality of other complementary radiation effects without relying on ZPF alone (e.g. other isotropic, homogeneous radiation sources); 2) the weak coupling constant, G , yields a vastly smaller than observed size of the universe (e.g. too small cosmological constant) when the Planck frequency is used as a cutoff value; and (3) the particle mass, $m = \Gamma c^3 / G$, can be viewed as a renormalized or ‘dressed’ mass with a resonant interaction potential that is frequency dependent in its coupling constant, G , and with ‘bare’ mass that is large, $m_b \sim (m_p^2/m)$, where the experimentally unobservable, $m_p = (hc/G)^{1/2}$ is the Planck mass.

In particular, why this large ‘bare’ mass does not generate a large gravitational field is not a unique anomaly in the Sakharov

derivation, since similarly large vacuum point energies are common to field theories. The important point is that the derivation $G(\omega)$ is general however to any isotropic radiation field with the Lorentz-invariant energy spectra [$\rho(\omega) \sim \omega^3$], thus the candidates for the cutoff frequency of the particular radiation source can be interpreted as a Planck scale only if the rest mass, m_o , is not composed of many terms, rather than just the ZPF leading term. Since the ZPF is akin to a van der Waals force [3-5], polarizability (in charge and mass) must be considered, but without also excluding any number of linear combinations that might have alternative cutoff frequencies, ω_c , or damping terms, Γ , 'ala particle physics interpretations for resonant masses during renormalization. In other words, once a gravitational energy spectrum, $\rho(\omega)$, is postulated that is Lorentzian invariant, many fundamental sizes (or corresponding frequency values) are smeared (or dressed) by any number of characteristic frequencies between zero and the high frequency electromagnetic (Planck) cutoff, ω_p . Quite simply, is the expression, $\omega_c = \omega_p$, a requirement for all radiation sources?

Many types of particle oscillations may satisfy the general requirements of a Sakharov condition, each having a characteristic mass (and energy) as in calculating the mass of any fundamental particle at its resonant frequency (including underlying partial charges or dense bosons). This brings the calculation to a consideration of the high density fluctuations characteristic of a Bose condensate [15-19]. While the high density variation may intrinsically be of interest, the exploration has more to do with reconciling the ZPF interpretation of the Sakharov condition with the observed cosmological constant [14].

A 'top-down' view of calculating the cutoff frequency imposes the self-consistency test for the cosmological constant, Λ , from the outset. To calculate, Λ , for ZPF, the total frequency-integrated energy density of the universe must be included:

$$\rho(E) = \int \rho(E) dE = h\omega_c^4 / 8\pi^2 c^3$$

which must have a mass equivalent, contribute to the universe's curvature, and thus have a fundamental relation to the critical density to contain the expansion of the universe [14-15]. The mass-equivalent ZPF to reach the universe's critical density [15], $\rho \sim 10^{-29} \text{ g cm}^{-3}$ would

necessarily limit the cutoff frequency for gravity to the value, $\omega_c < 7 \times 10^7 \text{ s}^{-1}$, or between 10-100 MHz.

A higher frequency greatly overshoots the cosmological constant, Λ , and induces extreme curvature in the universe. This problem has been cited frequently and stated most bluntly as either ZPF or the cosmological constant requires revision. The relevance here arises from similarly large positive coupling terms in quantum gravity [15], which also generate a local gravitational instability for typical upper limits on the cosmological constant, $\Lambda / 8\pi G < 10^{12} \text{ cm}^{-4}$.

Rather than to dwell on the inconsistencies that plague attempts to reconcile quantum gravity, we particularize the problem to a case where the restriction to Planck scale becomes less clear, namely the high density fluctuations and universal scaling introduced in a Bose condensate. A Bose condensate, such as superfluid helium or superconductors [15-19], becomes of potential interest, mainly because of its arbitrarily dense boundaries and the classic Casimir experiment [20-22] which allows such dense material boundaries (two closely spaced conducting plates), if available, to modulate the background quantum fluctuation of ZPF. In other words, the matter-ZPF interaction becomes measurable by the observed attraction between two material boundaries. What dense boundaries might generate in Bose condensates remains a subject of great interest.

The significant case to investigate is whether Casimir-like interactions [20-22] will not only couple to ZPF radiation at a scale comparable to the quantum noise (or other radiation field), but also alter the value imposed by the Sakharov condition for G . It remains an open question whether this potential coupling interaction shares, as in ordinary critical phenomenon, the density correlation function, Φ , that is both independent of the coupling strength (or universal in renormalization) and consistent with the observed average energy density of the visible universe.

Thus the purpose here has been to restate the Sakharov condition in the gravitational coupling constant, G , based on its only free parameter, a frequency cutoff, ω_c . Any potential relevance arises from similarly large values for the positive coupling term in quantum gravity, which generate conditions for a local

gravitational instability for typical upper limits on the cosmological constant, $\Lambda/8\pi G < 10^{12} \text{ cm}^{-4}$.

To restate the Sakharov condition, matter in the vacuum provides boundaries for reduced ‘Casimir-like’ modes available for otherwise isotropic radiation from quantum fluctuations (broad spectral noise). That this view reproduces Einstein gravity has been examined, including the full relativistic derivation [4-5]. The details of the appropriate mass, however, remain buried in the kinetic energy of general internal particle (‘parton’) motion [3]. Any appeal to a specific parton representation is limited only by essentially free particles with high-frequency interactions, including underlying partial charges or dense bosons. The basis of considering arbitrarily high density fluctuations in Bose condensate in analogy to the ZPF-Casimir experiment remains both an empirical and theoretical case to examine. There exist laboratory scale cases [15-19] where resonant radiation in the required 10-100 MHz range appear to produce anomalous effects for such Bose condensates as superconductors, but further work to confirm these results would be needed. In other contexts, these effects have been discussed as the Schiff-Barnhill effect for superconductors interacting with a gravitational field [23], but for the static rest mass rather than an effective mass in a conduction band.

Experimental Propositions

J. Weber [24,25] proposed the use of a superconducting Bose condensate for gravity wave detection, principally because of its potentially higher signal-to-noise ratio in carrying electrical signals upon length dilations in a relativistic framework for gravity waves travelling near the speed of light. W. Weber and Hickman [26] derived an experimentally testable relation based on torquing of a charged capacitor parallel to a gravity field, with

$$\tau = 2E_g/\pi [\alpha / (1 - \alpha)^{1/2}],$$

where the capacitor will rotate relative to the gravity vector, for $\alpha = 2GM/r c^2$, r is Schwarzschild radial coordinate [$dR = dr(1 - \alpha)^{1/2}$], E_g is dependent on the capacitor charge and geometry of the plates, $E_g = Q^2 d / [2\epsilon WL(1 - \alpha)^{1/2}]$, for a plate separation, d , and radial dimensions, W and L , charge Q , and ϵ the permittivity of free space. For plate separations

of 2 mm on Earth, the maximum torque is approximately $\tau = 10^{-12} \text{ Nm}$, when charged to 2/3 dielectric breakdown. While not entirely promising for detection of such low torques, the large separation (2 mm) distance between capacitative plates naturally prompts generalization to the classic Casimir force [21] experiments only recently confirmed experimentally [20]. In particular, we rewrite the torque values to include the frequency terms derived with the Sakharov condition [$G = (\pi^5 / h\omega_c^2)$]:

$$\alpha = 2 M \pi c^3 / h\omega_c^2 r$$

The appeal of this formulation is that a frequency dependent torque is derived, which further makes contact with proposals to modulate the Casimir capacitative plates for continuous extraction of energy [27]. This result requires further investigation experimentally, particularly to compare with previous reports for anomalies in AC-tuned electrical capacitors [28].

References

- [1] Zel'dovich, Ya. B. JETP Letters, 6, 345, 1967.
- [2] Sakharov, A. Vacuum quantum fluctuations in Curved Space and the Theory of Gravitation, Sov. Phys. Doklady, 12, 1968, 1040-1041.
- [3] Puthoff, H. E. (1989) Gravity as a zero-point-fluctuation force, Physical Review A, 39(5):2333-2342, March 1, 1989.
- [4] Haisch, B., Rueda, A., and Puthoff, H. E. (1994) Inertia as a Zero-Point Field Lorentz Force, Physical Review A, 49:678-694.
- [5] Haisch, B., A. Rueda, H.E. Puthoff, “Inertia as a Zero-Point Field Force” Physical Review A 49, Nr 2, 678 (1994).
- [6] Ambjørn, J. and Wolfram, S. (1983) Properties of the Vacuum, 1. Mechanical and Thermodynamic, and Properties of the Vacuum, 2. Electrodynamical, Annals of Physics,
- [7] Ambjørn, J. and Wolfram, S. (1983) Properties of the Vacuum. I. Mechanical and Thermodynamic, Annals of Physics, 147:1-32.
- [8] Fulcher et al., "The Decay of the Vacuum," Sci. Am., vol. 241, p. 150, Dec. 1979
- [9] Puthoff, H.E. “Source of Vacuum Electromagnetic Zero-Point Energy” Physical Review A 40, 4857 Nov 1 (1989); Errata and Comments, Physical Review A 41, March

- [1](1990); Physical Review A 4 4, 3382, 3385 (1991).
- [10] Senitzky, I. R., "Radiation-Reaction and Vacuum Field Effects in Heisenberg - Picture Quantum Electrodynamics," Phys. Rev. Lett. 31(15), 955 (1973). As pointed out by Puthoff [3] the relativistic results for the Sakharov condition have so far been encouraging, while the consequences for nuclear interactions in all coordinate frames have not been fully explored.
- [11] da Costa, L. N., Freudling, W., Wegner, G., Giovanelli, R., Haynes, M. P., and Salzer, J. J. (1996) The Mass Distribution in the Nearby Universe, Astrophysical Journal Letters, 468: L5-L8 and Plate L1
- [12] Modanese, G. (1996) Theoretical analysis of a reported weak gravitational shielding effect, Europhy. Lett., 35(6):413-418.
- [13] Shupe, M.A. The Lorentz-invariant vacuum media, Am. J. Phys. 53, 122 (1985). A cautionary note is that lower frequency cutoffs can violate Lorentzian invariance, thus allowing a moving detector to reveal absolute motion by recording Doppler-shifted frequencies. Standard methods might treat such effects like the cancellation of terms that remove anomalous ZPF infinities from field theories, but these topics remain to be explored.
- [14] Zee, A.Phys. Rev. Lett., 42,417 (1979); Phys. Rev. D. 23, 858, (1981).
- [15] Torr, D. G. and Li. N. (1993) Gravitoelectric-Electric Coupling vVia Superconductivity, Foundations of Physics Letters, 6(4):371-383.
- [16]Unnikrishnan, C. S. (1996) Does a superconductor shield gravity?, Physica C, 266:133-137.
- [17] Podkletnov, E. and Nieminen, R. (1992) A Possibility of Gravitational Force Shielding by Bulk YBa₂Cu₃O_{7-x} Superconductor, Physica C, C203:441-444.
- [18]Li, N. and Torr, D. G. (1992) Gravitational effects on the magnetic attenuation of superconductors, Physical Review B, 46(9):5489-5494. A simple consequence of the Sakharov condition, $G=(\pi c^5/\hbar\omega_c^2)\sim 1/\int \omega d\omega$, can be written for the gravitomagnetic permeability as:
 $\mu_g=4\pi G/c^2=4\pi^2 c^3/(h \int_0^\infty \omega d\omega) \sim 1/\int \omega d\omega$
which suggest that the same frequency resonance implied by the ZPF derivation will share similar consequences for vector gravity effects. See also, DeWitt, B.S. Superconductors and Gravitational Drag, Phys. Rev. Lett. 16, 1092 (1966).
- [19] Li, N., Noever, D., Robertson, T., Koczor, R., and Brantley, W. (1997) Static Test for a Gravitational Force Coupled to Type II YBCO Superconductors, Physica C, 55, 287.
- [20] Lamoreaux, S. K. (1997) Demonstration of the Casimir Force in the 0.6 to 6 μm Range, Phys. Rev. Letters, 78:5-8.
- [21] Milonni, P.W. et al., "Radiation pressure from the vacuum: Physical interpretation of the Casimir force", Phys. Rev. A, Vol. 38, No. 3, 1621 August 1988.
- [22] Milonni, P. W. (1994) The Quantum Vacuum, Academic Press, San Diego, CA.
- [23] Schiff, L.I. and Barnhill, M.V. Bull. Am. Phys. Soc. 11, 96, (1966) and refn. 18.
- [24]Weber, J. (1960), Detection and Generation of Gravitational Waves, Physical Review, 117(1):306-313.
- [25]Weber, J. (1966) Gravitational Shielding and Absorption, The Physical Review (The American Physical Society), 146(4):935-937.
- [26] Weber, W. and Hickman, H. (1997)A possible interaction between gravity and the electric field, Spec. Science Tech. 20, 133-136
- [27]Forward, R.L. "Extracting electrical energy from the vacuum by cohesion of charged foliated conductors" Phys. Rev. B, Vol. 30, No. 4, 1700 August 1994
- [28]Woodward, J. F. (1992) A Stationary Apparent Weight Shift From a Transient Machian Mass Fluctuation, Foundations of Physics Letters, 5:425-442.

Breakthrough Propulsion Physics Research Program

Marc G. Millis
*Lewis Research Center
Cleveland, Ohio*

Prepared for the
Space Technology and Applications International Forum
cosponsored by NASA, DSWA, U.S. DOE, and USAF
Albuquerque, New Mexico, January 26–30, 1997



National Aeronautics and
Space Administration

BREAKTHROUGH PROPULSION PHYSICS RESEARCH PROGRAM

Marc G. Millis
NASA Lewis Research Center
21000 Brookpark Rd., Cleveland, OH 44135
(216) 977-7535
marc.millis@lerc.nasa.gov

Abstract

In 1996, a team of government, university and industry researchers proposed a program to seek the ultimate breakthroughs in space transportation: propulsion that requires no propellant mass, propulsion that can approach and, if possible, circumvent light speed, and breakthrough methods of energy production to power such devices. This Breakthrough Propulsion Physics program, managed by Lewis Research Center, is one part of a comprehensive, long range Advanced Space Transportation Plan managed by Marshall Space Flight Center. Because the breakthrough goals are beyond existing science, a main emphasis of this program is to establish metrics and ground rules to produce near-term credible progress toward these incredible possibilities. An introduction to the emerging scientific possibilities from which such solutions can be sought is also presented.

INTRODUCTION

In 1996, Marshall Space Flight Center (MSFC) was tasked to formulate a comprehensive strategic plan for developing space propulsion technology for the next 25 years. This “Advanced Space Transportation Plan” spans the nearer-term launcher technologies all the way through seeking the breakthroughs that could revolutionize space travel and enable interstellar voyages.

New theories and phenomena have emerged in recent scientific literature that have reawakened consideration that such breakthroughs may be achievable. To establish a program to address these visionary possibilities, a “Product Definition Team” of researchers was assembled. This team, led by NASA Lewis Research Center, consisted of 19 individuals from various NASA Centers, other government laboratories, industries, and academia (listed in acknowledgments section). Most team members are part of an existing informal network that had already recognized the potential of the emerging science and had conducted preparatory research on how to apply these prospects to the goal of creating revolutionary propulsion.

To anchor the program in real and tangible terms, the team configured the program to produce near-term, credible, and measurable progress toward determining how and if such breakthroughs can be achieved – credible progress to incredible possibilities. There is no guarantee that the desired breakthroughs are achievable, but it is possible to produce progress toward a goal without first proving it is achievable. This paper introduces how this program aims to answer these challenges as well as giving a brief introduction to the emerging physics which reawakened interest in these visionary ambitions.

SPECIFYING GOALS AND SCOPE

To focus the program, the first step is to specify what breakthroughs are genuinely required to revolutionize space travel. A NASA precedent for systematically seeking revolutionary capabilities is the “Horizon Mission Methodology” (Anderson 1996). This method forces paradigm shifts beyond extrapolations of existing technologies by using *impossible* hypothetical mission goals to solicit new solutions. By setting impossible goals, the common practice of limiting visions to extrapolations of existing solutions is prevented. The “impossible” goal used in this exercise is to enable practical interstellar travel. Three major barriers exist to practical interstellar travel; propellant mass, trip time, and propulsion energy. To conquer these hurdles the following three propulsion breakthroughs are sought. These are the goals of the Breakthrough Propulsion Physics program:

- (1) Eliminate or dramatically reduce the need for rocket propellant. This implies discovering fundamentally new ways to create motion, presumably by manipulating inertia, gravity, or by any other interactions between matter, fields, and spacetime.
- (2) Dramatically reduce trip time to make deep space travel practical. This implies discovering a means to move a vehicle at or near the actual maximum speed limit for motion through space or through the motion of spacetime itself. If possible, this means circumventing the light speed limit.
- (3) Discover fundamentally new on-board energy production methods to power propulsion devices. This third goal is included in the program since the first two breakthroughs could require breakthroughs in energy generation to power them, and since the physics underlying the propulsion goals is closely linked to energy physics.

The scope of this program only covers seeking the genuinely needed breakthroughs rather than seeking refinements to existing solutions. As such, existing concepts that are based on firmly established science, such as light sails, magnetic sails, beamed energy, nuclear rockets, and antimatter rockets, are not part of this program. These concepts are being explored in other programs.

SCIENTIFIC FOUNDATIONS

New possibilities have emerged in recent scientific literature that have reawakened interest toward conquering the goals described above. These include theories that suggest that gravity and inertia are electromagnetic side effects of vacuum fluctuations (Haisch 1994 and Puthoff 1989), anomalous experimental evidence suggesting a possible gravity altering affect from spinning superconductors (Podkletnov 1992), theories suggesting that faster-than-light transport may be possible using wormholes (Morris 1988) or using warp drives (Alcubierre 1994), and a theory suggesting that sonoluminescence is evidence of extracting virtual photons from vacuum fluctuation energy (Eberlein 1996). These are in addition to older theories about creating propulsive effects without rockets (Bondi 1957 and Forward 1963).

In addition, there have been workshops (Bennett 1995, Evans 1990, and Landis 1990), recent surveys (Cravens 1990, Forward 1990, and Mead 1989), suggested research approaches (Cramer 1994, Forward 1984 and 1996, and Millis 1996), and even some exploratory experiments (Millis 1995, Schlicher 1995, and Talley 1991) on this subject. And recently, a non-profit society, the Interstellar Propulsion Society, was established to provide a collaborative forum to accelerate advancements toward these goals (Hujšak 1995).

PROGRAM CHALLENGES

Since the scientific principles do not yet exist from which to engineer the technological solutions to these challenges, new scientific principles are sought. Seeking such visionary and application-focused physics is not a usual activity for aerospace institutions, so this program faces both the technical challenge of discovering the desired breakthroughs and the programmatic challenges of how to conduct this work. To answer these challenges, the program will develop the research solicitation and selection criteria and the metrics for quantifying progress to meet the technical and programmatic challenges.

The technical challenges include: (1) focusing emerging theories and experiments to answer NASA's propulsion needs, (2) finding the shortest path to developing the breakthroughs amidst several, divergent and competing approaches, and (3) balancing the imagination and vision necessary to point the way to breakthroughs with the credible, systematic rigor necessary to make genuine progress.

The programmatic challenges include: (1) advocating such long range research amidst dwindling resources and stiff competition from nearer-term, more conservative programs, (2) creating confidence that research funded today will lead to the necessary breakthroughs, (3) selecting the most promising research tasks from the large number of divergent and competing approaches, and (4) conducting meaningful and credible research economically.

PROGRAM PRIORITIES

To simultaneously focus emerging sciences toward answering the needs of space travel and to provide a programmatic tool for measuring progress and relevance, this program will develop prioritization criteria suitable for breakthrough-seeking research. These criteria help potential researchers to focus their work, and provide NASA with the means to quantify the relative benefit of competing research proposals. Examples of these criteria, which are still evolving, are presented below:

- **APPLICABILITY:** Research proposals that directly address a propulsive effect are given preference over those addressing basic science or supporting technologies.
- **EMPIRICISM:** Research proposals for experimental tests are given preference over proposals for analytical or theoretical work. Empiricism is considered to be a more direct indication of physical phenomena and experimental hardware is considered to be closer to becoming technology than pure theory.
- **TARGETED GAIN:** Research proposals that address large potential improvements in propulsive abilities are given preference over those of lesser potential improvements. This comparison can only be made between concepts that are of similar developmental maturity. Targeted gains include reducing trip time, reducing non-payload vehicle mass, reducing energy requirements, and reducing development cost.
- **ACHIEVABILITY:** Research proposals whose subjects are closer to becoming a working device are given preference over longer range developments. This comparison can only be made between concepts that are of similar propulsive ability.
- **IMPACT:** The more likely the research results will be used by others, the higher it is ranked.

Another criteria to be used is a “Traceability Tree.” This tree is the reverse of a fault tree (a common tool used in accident risk assessments). Instead of branching out all the ways that a given accident can occur, the Traceability Tree branches out the conceivable paths to reach a propulsion breakthrough, Figure 1. This tool provides a means to measure how a given research task is linked to the end goals of discovering a propulsion breakthrough.




FIGURE 1. Illustrative Example of a Traceability Tree.

Another philosophy behind the NASA program is to support multiple small tasks rather than a single, larger task. Since the search for breakthrough propulsion physics is at an early stage, there are a large number of divergent and competing approaches. It is still too soon to tell which approaches will provide the shortest path to success. To cover the bases, a diverse program of multiple, small, and short-term projects is preferred.

To balance vision and imagination with credible systematic rigor, the NASA program is open to perspectives that go beyond or appear to contradict conventional theory, with the conditions that (1) the perspectives must be completely consistent with credible empirical evidence, (2) the utility of the alternate perspective must be clearly delineated, and (3) the make-or-break issues must be clearly identified to suggest the next-step research objectives.

There is another facet to this “beyond conventionalism.” Institutions such as NASA routinely get numerous unsolicited submissions from individuals who claim to have a breakthrough device or theory. Frequently these submissions are too poorly conceived or incomplete, or too complex to be easily evaluated. To distribute this workload to other credible organizations and to keep an open mind to the possibility that one of these maverick inventors may have actually made a breakthrough, the following approach is suggested. Any individual who thinks they have a breakthrough device or theory is strongly advised to collaborate with a university or other educational institute to conduct a credible test of their claim. The university can propose the test as an educational student project, where the students will learn first-hand about the scientific method and how to apply systematic rigor and open-mindedness in conducting a credible test of an incredible claim. In such collaborations it is suggested that the inventors retain full intellectual property rights to their devices or theories, and that the universities make the proposal and receive all the funds to conduct the student projects. With this procedure, if the device or theory works, then supporting evidence would be established in a credible fashion and the originator would retain the intellectual property rights. If the device or theory does not work, then at least the students would have had a meaningful educational experience, and the concept’s originator can work on another idea.

MEASURING PROGRESS

One of the challenges to this program is to demonstrate that research conducted today is making measurable progress directly toward the long-range targeted breakthroughs. The traceability tree and selection criteria presented earlier provides a means to demonstrate that a given research approach is linked to the end goals, but another metric is needed to quantify that progress is being made. To provide this measure, a more explicit version of the Scientific Method is used for measuring the level of advancement of a given scientific approach, similar to the way that the “Technology Readiness Levels” (Hord 1985) are used for quantifying technological progress. A draft of these scientific readiness levels is below:

- **Pre-Science:** Suggests a correlation between a desired effect and an existing knowledge base, or reports observations of unexplained anomalous effect.
- **Scientific Method Level 1, Problem Formulation:** Defines a problem specifically enough to identify the established knowledge base and the remaining knowledge gaps. A “problem” is an explicit statement of a desired goal (e.g. the 3 Breakthrough Propulsion Physics program goals) or explicit observations of an anomalous effect which cannot be explained using the established knowledge base.
- **Scientific Method Level 2, Data Collection:** Compiles relevant information to address a specific problem by experiment, observation, or mathematical proof.
- **Scientific Method Level 3, Hypothesis:** Suggests a mathematical representation of an effect or the relation between physical phenomena.
- **Scientific Method Level 4, Test Hypothesis:** Empirically tests a hypothesis by comparison to observable phenomena or by experiment.
- **Technology Readiness Level 1, Basic Principles Observed and Reported:** Equivalent to established science, where an effect has been observed, confirmed, and modeled sufficiently to produce a mathematical description of its operation.
- **Technology Readiness Level 2, Conceptual Application Designed.**
- **Technology Readiness Level 3, Conceptual Design Tested Analytically or Experimentally.**

STATUS AND DIRECTION

A government steering group containing members from various NASA centers, DOD and DOE laboratories has been established to develop the research solicitation and selection criteria to meet the programmatic and technical challenges. These criteria are to be in place by mid 1997, in time for the next program milestone; a kick-off workshop. The invitation-only workshop will examine the relevant emerging physics and will produce a list of next-step research tasks. If the workshop successfully demonstrates that promising and affordable approaches exist, funding may be granted to begin conducting the step-by-step research that may eventually lead to the breakthroughs.

CONCLUSIONS

New theories and laboratory-scale effects have emerged that provide new approaches to seeking propulsion breakthroughs. To align these emerging possibilities toward answering the propulsion needs of NASA, a research program has been established that focuses on producing near term and credible progress that is traceable to the breakthrough goals.

Acknowledgments

Special thanks is owed to the Product Definition Team who helped shape this program: Team Leader; Marc G. Millis, NASA Lewis Research Center; Team Members: Dana Andrews, Boeing Defense and Space Group; Gregory Benford, University of California, Irvine; Leo Bitteker, Los Alamos National Labs; John Brandenburg, Research Support Instruments; Brice Cassenti, United Technologies Research Center; John Cramer, University of Washington, Seattle; Robert Forward, Forward Unlimited; Robert Frisbee, NASA Jet Propulsion Lab; V.E. Haloulakos, McDonnell Douglas Aerospace; Alan Holt, NASA Johnson Space Flight Center; Joe Howell, NASA Marshall Space Flight Center; Jonathan Hujas, Interstellar Propulsion Society; Jordin Kare, Lawrence Livermore National Labs; Gregory Matloff, New York University; Franklin Mead Jr., Phillips Labs, Edwards Air Force Base; Gary Polansky, Sandia National Labs; Jag Singh, NASA Langley Research Center; Gerald Smith, Pennsylvania State University; and to the Lewis Research Center volunteers; Michael Binder, Gustav Fralick, Joseph Hemminger, Geoffrey Landis, Gary S. Williamson, Jeffrey Wilson, and Edward Zampino.

References

- Alcubierre, M. (1994) "The Warp Drive: Hyper-fast Travel Within General Relativity", *Classical and Quantum Gravity*, 11:L73-L77.
- Anderson, J. L. (1996) "Leaps of the Imagination: Interstellar Flight and the Horizon Mission Methodology," *Journal of the British Interplanetary Society*, 49:15-20.
- Bennett, G., Forward, R. L. and Frisbee, R. (1995) "Report on the NASA/JPL Workshop on Advanced Quantum/Relativity Theory Propulsion" AIAA 95-2599, 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference.
- Bondi, H. (1957) "Negative Mass in General Relativity," *Reviews of Modern Physics*, 29:423-428.
- Cramer J., Forward, R. L., Morris, M., Visser, M., Benford, G. and Landis, G. (1994) "Natural Wormholes as Gravitational Lenses," *Physical Review D*, 15 March 1995:3124-3127.
- Cravens D. L. (1990) "Electric Propulsion Study," Report # AL-TR-89-040, Air Force Astronautics Lab (AFSC), Edwards AFB, CA
- Eberlein, C. (1996) "Theory of quantum radiation observed as sonoluminescence," *Phys Rev A*, 53:2772-2787.

- Evans, R. A., ed. (1990) *BAe University Round Table on Gravitational Research*, Report on Meeting held in Preston UK, March 26-27, 1990, Report # FBS 007, British Aerospace Limited, Preston, Lancashire, UK.
- Forward, R. L. (1963) "Guidelines to Antigravity", *American Journal of Physics*, 31:166-170.
- Forward, R. L. (1984) "Extracting Electrical Energy from the Vacuum by Cohesion of Charged Foliated Conductors", *Physical Review B*, 39(15 AUG 1984):1700-1702.
- Forward, R. L. (1990) "21st Century Space Propulsion Study," Report # AL-TR-90-030, Air Force Astronautics Lab (AFSC), Edwards AFB, CA.
- Forward, R. L. (1996) "Mass Modification Experiment Definition Study," Report # PL-TR-96-3004, Phillips Lab, Edwards AFB, CA
- Foster, R. N. (1986) *Innovation; The Attacker's Advantage*, Summit Books.
- Haisch, B., Rueda, A., and Puthoff, H. E. (1994) "Inertia as a Zero-Point Field Lorentz Force", *Physical Review A*, 49:678-694.
- Hord, R. M. (1985) *CRC Handbook of Space Technology: Status and Projections*, CRC Press, Inc. Boca Raton, FL.
- Hujsak, J. T. and Hujsak, E. (1995) "Interstellar Propulsion Society," Internet address: <http://www.tyrian.com/IPS/>.
- Landis, G. L., ed. (1990) "Vision-21: Space Travel for the Next Millennium" Proceedings, NASA Lewis Research Center, April 3-4 1990, NASA CP 10059, NASA Lewis Research Center.
- Mead, F. Jr. (1989) "Exotic Concepts for Future Propulsion and Space Travel", *Advanced Propulsion Concepts, 1989 JPM Specialist Session*, (JANNAF), CPIA Publication 528:93-99.
- Millis, M. G. and Williamson, G. S. (1995) "Experimental Results of Hooper's Gravity-Electromagnetic Coupling Concept," NASA TM 106963, Lewis Research Center.
- Millis, M. G. (1996) "The Challenge to Create the Space Drive", NASA TM 107289, Lewis Research Center.
- Morris, M. and Thorne, K. (1988) "Wormholes in Spacetime and Their Use for Interstellar Travel: A Tool for Teaching General Relativity," *American Journal of Physics*, 56:395-412.
- Podkletnov, E. and Nieminen, R. (1992) "A possibility of gravitational force shielding by bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ superconductor" *Physica*, C203:441-444.
- Puthoff, H. E. (1989) "Gravity as a zero-point-fluctuation force", *Phys Rev A*, 39:2333-2342.
- Schlischer R. L., Biggs, A. W., and Tedeschi, W. J. (1995) "Mechanical Propulsion From Unsymmetrical Magnetic Induction Fields", AIAA 95-2643, 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference.
- Talley, R. L. (1991) "Twenty First Century Propulsion Concept," Report # PL-TR-91-3009, Phillips Laboratory, Air Force Systems Command, Edwards AFB, CA.

REPORT DOCUMENTATION PAGE			<i>Form Approved OMB No. 0704-0188</i>
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE December 1996	3. REPORT TYPE AND DATES COVERED Technical Memorandum	
4. TITLE AND SUBTITLE Breakthrough Propulsion Physics Research Program		5. FUNDING NUMBERS WU-242-74-40	
6. AUTHOR(S) Marc G. Millis			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Lewis Research Center Cleveland, Ohio 44135-3191		8. PERFORMING ORGANIZATION REPORT NUMBER E-10569	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA TM-107381	
11. SUPPLEMENTARY NOTES Prepared for the Space Technology and Applications International Forum cosponsored by NASA, DSWA, U.S. DOE, and USAF, Albuquerque, New Mexico, January 26-30, 1997. Responsible person, Marc G. Millis, organization code 5340, (216) 977-7535.			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified - Unlimited Subject Category 70 This publication is available from the NASA Center for AeroSpace Information, (301) 621-0390.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) In 1996, a team of government, university and industry researchers proposed a program to seek the ultimate breakthroughs in space transportation: propulsion that requires no propellant mass, propulsion that can approach and, if possible, circumvent light speed, and breakthrough methods of energy production to power such devices. This Breakthrough Propulsion Physics program, managed by Lewis Research Center, is one part of a comprehensive, long range Advanced Space Transportation Plan managed by Marshall Space Flight Center. Because the breakthrough goals are beyond existing science, a main emphasis of this program is to establish metrics and ground rules to produce near-term credible progress toward these incredible possibilities. An introduction to the emerging scientific possibilities from which such solutions can be sought is also presented.			
14. SUBJECT TERMS Space propulsion; Physics; General relativity; Gravity; Antigravity; Special relativity; Quantum physics			15. NUMBER OF PAGES 08
			16. PRICE CODE A02
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT

MEASUREMENT OF REPULSIVE QUANTUM VACUUM FORCES

Jordan Maclay, Principal Scientist, Member, Quantum Fields LLC, Richland Center WI 53581

Jay Hammer, Senior Engineer, MEMS Optical Inc., 205 Import Circle, Huntsville AL 35806

Michael A. George, Assoc. Professor, Lelon Sanderson, Research Assistant, Department of Chemistry, University of Alabama at Huntsville, Huntsville, AL 35899

Rob Ilic, Research Assistant, Department of Applied Physics, Cornell University, 212 Clark Hall, Ithaca NY 14853

Quinn Leonard, Laboratory Manager, Univ.of Wisc.-Madison, Center for NanoTechnology, Stoughton, WI 53558

Rod Clark, President, MEMS Optical Inc., 205 Import Circle, Huntsville AL 35806

ABSTRACT

Quantum electrodynamics predicts that empty space (the quantum vacuum) contains a large amount of energy that corresponds to the lowest energy state (energy >0) of the electromagnetic field. Surfaces in the vacuum can experience forces that arise from the disturbance in the vacuum energy. The presence of attractive "Casimir" forces between uncharged, parallel, metal plates has been accurately verified in the last several years. Theoretical calculations have suggested the presence of repulsive vacuum forces for certain geometrical configurations. Here we describe an experiment in progress that is designed to determine if repulsive vacuum forces exist. In the experiment we measure the force exerted on a 200 um diameter metallized sphere mounted on an Atomic Force Microscope (AFM) that is placed very close to an array of gold microcavities. Observing a repulsive force on the sphere would verify the existence of repulsive forces. The ability to create attractive and repulsive vacuum forces by means of the geometry of the surfaces may permit the construction of devices that use ubiquitous vacuum energy in ways that assist with the space travel mission of NASA.

INTRODUCTION

Understanding the nature of vacuum forces and vacuum energy and how to manipulate this energy to obtain desired forces is a prerequisite to using these ubiquitous natural resources in any space application¹. The theory for vacuum forces and quantum vacuum energy comes from Quantum Electrodynamics (QED), the theory of the interaction of matter and light². **The role of the quantum vacuum is pervasive in modern physics. For example, it is involved in the calculation of atomic energy levels, the magnetic moment of the electron, the mass of elementary particles, spontaneous emission, dispersion forces between molecules, the large-scale structure of space-time.**

The experiment discussed in this paper is part of a three-year effort to begin to build, step by step, the knowledge base necessary for vacuum engineering. Our objective is to develop theoretical models of elementary systems that utilized vacuum forces and energy, to understand how these models behave, and then to explore some of these models experimentally. Since the critical dimensions required for these devices are typically micron to submicron, the experimental research utilizes microfabrication technology, and the methods developed for MicroElectro-Mechanical Systems (MEMS).

In space applications the application of vacuum energy systems might be power generation, propulsion itself, or the manipulation of the metric of space-time itself by the creation of regions of positive and negative energy density³. It is too early to determine if such developments are possible or to be able to clearly determine the role of vacuum energy in future space applications. If we can develop technologies for space travel that utilize vacuum energy, it is very convenient since this energy is pervasive throughout the universe.

Fifty years ago, Casimir predicted that the modifications to the vacuum energy arising from the presence of two uncharged, parallel, metal plates would cause the plates to attract each other. This attractive Casimir force varies as the inverse fourth power of the separation. At a separation of 10 nm the force/area is about 1 atm. In 1997 the prediction of Casimir was verified for the first time. In 1998 precision measurements corroborated the predictions of an attractive vacuum force between neutral parallel plates to an accuracy of several percent. **In March 2001, scientists at Lucent Technology used attractive parallel plate vacuum forces (Casimir forces) to actuate a MEMS torsion device⁴. Other MEMS devices using vacuum energy have been proposed.⁵**

*Copyright ©2001 Quantum Fields LLC.

Published by American Institute of Aeronautics and Astronautics, Inc. with permission.

Recent calculations have indicated that forces due to the quantum vacuum predicted by QED depend very strongly on the geometry of surfaces. For certain rectangular metal cavities, QED predicts the existence of repulsive forces on the walls of the cavity⁶. In this paper we describe the current status of the first experiment specifically designed to measure repulsive forces due to modifications in vacuum energy density achieved by using metal surfaces. The vacuum force is measured by means of an Atomic Force Microscope using a 200 um diameter metallized ball placed on the end of a calibrated cantilever. Our model suggests that a repulsive force on the ball would be observed when it approaches within 10's of nanometers from the top of an array of rectangular cavities, each of which is 100nm across and 1 um deep, patterned in gold using x-ray photolithography (Figure 2). For small separations between the surface of the sphere and the top of the cavity array, we are approximating an array of closed cavities, which, according to a QED calculation, exhibit repulsive forces. The force between the sphere and the cavity array is modeled numerically, with heuristic approximations, to be compared to the measured force. It is important to note that no rigorous method has yet been developed to calculate the vacuum force between any two non-planar conducting surfaces using QED. Only the parallel plate problem has been solved. Indeed there is some disagreement that a repulsive Casimir force should ever be present between two separate bodies⁷. Measurement of repulsive forces between separate conducting bodies may be expected to stimulate new developments in QED.

If our experiment verifies the existence of repulsive vacuum forces, then it may be possible to utilize repulsive forces as well as attractive vacuum forces in microelectro-mechanical systems (MEMS). The existence of attractive and repulsive Casimir forces might permit the development of a variety of novel MEMS devices of potential use to NASA.

Quantum Electrodynamics (QED), Vacuum Energy and Casimir Forces

Quantum Electrodynamics (QED), the theory of the interaction of electromagnetic fields and matter, has made predictions of atomic energy levels and electron magnetic moments that have been verified to 1 part in 10^{12} , which makes QED the most precisely verified theory in science^{8,9}. In order to achieve this accuracy, QED predictions have to include the

interaction of matter with “empty space” or, more accurately, the quantum vacuum¹⁰.

Some predictions of QED are less enthusiastically received by the physics community than others. One of the confounding predictions of QED is an energy density in empty space that is many orders of magnitude greater than the energy density of matter itself¹¹. For years this feature of QED was dismissed as of no physical significance. However, observable forces can result when surfaces are present that alter this vacuum energy density. About 50 years ago, Phillips Laboratory physicist H.G.B. Casimir predicted the presence of an attractive quantum vacuum force between neutral, parallel, metal plates¹². In the last three years, experiments have accurately confirmed this prediction of QED for the first time, verifying the existence of attractive vacuum forces between conductive surfaces^{13,14,4}. The parallel plate Casimir force goes as the inverse fourth power of the separation between the plates. At a separation of 100 nm the predicted force/area is equivalent to about 10^{-4} atm.; at 10 nm it is about 1 atm.

BACKGROUND

Since most aerospace researchers do not have backgrounds in quantum systems, we provide a brief background to motivate our study. It is certainly not obvious that there should be any energy at all in empty space, much less a very large amount! Nor is obvious why there should be forces due to the vacuum fluctuations. The evidence for this theoretical conclusion lies in numerous well verified experiments on atomic energy levels, the magnetic moment of the electron, the behavior of liquid helium, and the scattering of elementary particles¹⁵.

Vacuum energy is a consequence of the quantum nature of the electromagnetic field, which is composed of photons. A photon of frequency ω has energy $E = \hbar\omega$, where \hbar is Planck's constant. The quantum vacuum can be interpreted as the lowest energy state (or ground state) of the electromagnetic (EM) field that occurs when all charges and currents have been removed, and the temperature has been reduced to absolute zero. In this state no ordinary photons are present. Nevertheless, because the electromagnetic field is a quantum system, like an atom, which has internal motion even at absolute zero, the energy of the ground state of the EM field is NOT zero. Although the average value of the electric field $\langle E \rangle$ vanishes in the ground state, the Root Mean Square

(RMS) of the field $\langle E^2 \rangle$ is **not** zero. Similarly the RMS of the ground state magnetic field $\langle B^2 \rangle$ is not zero. Therefore the electromagnetic energy in the ground state, which from classical electrodynamics is proportional to $\langle E^2 \rangle + \langle B^2 \rangle$, is not zero. A detailed theoretical calculation tells us that the electromagnetic energy in each mode of oscillation with frequency ω is $\frac{1}{2}\hbar\omega$, which equals one half of the amount of energy that would be present if a single “real” photon of that mode were present. Adding up $\frac{1}{2}\hbar\omega$ for all possible modes of the electromagnetic field gives a very large number for the vacuum energy E_0 in the quantum vacuum:

$$E_0 = \frac{1}{2} \sum_i \hbar\omega_i \quad (0.1)$$

The resulting vacuum energy E_0 is infinity unless a high frequency limit is used.

Inserting surfaces into the vacuum causes the modes of the EM field to change. This change in the modes that are present occurs since the electromagnetic field must meet the appropriate boundary conditions at each surface¹⁶. **Surfaces alter the modes of oscillation and therefore the surfaces alter the energy density corresponding to the lowest state of the EM field.** In actual practice, the modes with frequencies above the plasma frequency do not appear to be significantly affected by the metal surfaces since the metal becomes transparent to radiation above this frequency. In order to avoid dealing with infinite quantities, the usual approach is to compute the finite change in the energy of the vacuum λ_0 due to the presence of the surfaces¹⁷:

$$\Delta E_0 \left[\begin{array}{l} \text{change in vacuum} \\ \text{energy due to surfaces} \end{array} \right] = E_0 \left[\begin{array}{l} \text{energy in} \\ \text{empty space} \end{array} \right] - E_s \left[\begin{array}{l} \text{energy in space} \\ \text{with surfaces} \end{array} \right] \quad (0.2)$$

where the definition of each term is given in brackets. This equation can be expressed as a sum over the corresponding modes:

$$\Delta E_0 (\text{due to surfaces}) = \frac{1}{2} \sum_n \hbar\omega_n - \frac{1}{2} \sum_i \hbar\omega_i \quad (0.3)$$

The quantity ΔE_0 , which is the change in the vacuum energy due to the presence of the surfaces, can be computed for various geometries. The forces F due to the quantum vacuum are obtained by computing the change in the vacuum energy for a small change in the geometry. For example, consider a hollow conducting rectangular cavity with sides a_1, a_2, a_3 . Let $\delta E(a_1, a_2, a_3)$ be the change in the vacuum energy due to the cavity, then the force F_1 on the side perpendicular to a_1 is:

$$F_1 = -\frac{\partial \delta E}{\partial a_1} \quad (0.4)$$




Figure 1. Geometry of rectangular cavity.

Equation (0.4) also represents the conservation of energy when the wall perpendicular to a_1 is moved infinitesimally¹⁸:

$$\delta E = -F_1 \delta a_1 \quad (0.5)$$

Thus if we can calculate the vacuum energy as a function of the dimensions of the cavity we can compute derivatives which give the forces on the surfaces. For uncharged parallel plates with a large area A , very close to each other, this equation predicts an attractive or negative force between the plates:

$$F_{att} = -\frac{\pi^2}{240} \frac{\hbar c}{d^4} A \quad (0.6)$$

This force is called the parallel plate Casimir force, which was measured in three different experiments in the last three years^{13,14,4}. The

Casimir force has only been computed and measured for this very large parallel plate geometry.

QED makes some unexpected predictions about Casimir forces in other geometries that have never been verified. For conductive rectangular cavities, the vacuum forces on a given face can be repulsive (positive), attractive (negative), or zero depending on the ratio of the sides⁶. **We are particularly interested in measuring these repulsive Casimir forces. Verifying the existence of such forces would have important implications in quantum electrodynamics and would be an important step to utilizing Casimir forces in a variety of MEMS devices.**

DESCRIPTION OF THE EXPERIMENT

Atomic Force Microscope

It is not practical to directly measure the force on one wall of a submicron metallic cavity. Hence another approach is needed. We chose to use an Atomic Force Microscope (AFM), which can provide a very sensitive measure of forces into the piconewton range (10^{-12} Newton). The AFM employs a 300 μm long micromachined silicon nitride cantilever with a 200 μm metallized sphere on the end that can be used to probe the vacuum energy density in the neighborhood of a rectangular micromachined cavity with no top surface¹⁹ (Figure 2). When the sphere experiences a force, the cantilever is deflected. The deflection of the sphere is measured by shining a laser diode onto the reflective surface of the cantilever. The reflected

light is collected in a photodiode that is divided into two adjacent regions. As the spot of light moves during a deflection, the ratio of current from the two regions changes, giving a sensitive quantitative measure of the cantilever deflection. It is possible to measure deflections of several nanometers in this manner. The cantilever is calibrated by determining the cantilever deflection for a known electrostatic force. The high precision of this experiment is made possible by the use of a Molecular Imaging AFM system that was specially developed at the University of Alabama at Huntsville for vacuum operation. With the very small distance (much less than the mean free path of the molecules) between the sphere and cavity, gas molecules can become effectively trapped, taking hours to remove under vacuum. For the most reliable measurements it is necessary to remove the trapped molecules and operate at a sufficiently low vacuum. Trapped molecules may result in a squeeze film damping force because the cantilever is always vibrating slightly.

The AFM stage was connected to a vacuum flange with the necessary feedthroughs. The sample is mounted and aligned when the AFM is in air (Figure 3a). Then the AFM is inserted into the vacuum chamber and the flange bolted in place using copper gaskets (Figure 3b). The vacuum without the AFM is 10^{-8} torr; inserting the AFM reduces the vacuum to about 10^{-4} torr. The system is pumped with a turbo molecular pump and an ion pump. The AFM is housed in a clean room environment.




Figure 2. Schematic of Atomic Force Microscope measurement of the vacuum force between a metallized sphere on a cantilever and a rectangular cavity etched in gold (not to scale).

The force constant of the AFM cantilever is measured by using electrostatic forces. A know

potential is applied between the test surface and the cantilever, and the deflection of the cantilever due to

this potential is measured. The corresponding force is calculated using a finite element classical electrodynamic calculation. The system was tested by




Figure 3. (a) shows the AFM stage below the vacuum flange, connected to the vacuum feedthroughs above. (b) shows the AFM mounted inside a small, stainless steel, vacuum chamber supported by elastic cords to reduce vibration.

Cavity Design


Numerical computations were done of the change in the vacuum energy and vacuum forces for a variety of rectangular cavities using QED methods^{1,6}. The goal was to determine a cavity geometry that 1) would yield a large, detectable, repulsive force, 2) the repulsive force would change slowly with distance, and 3) that could be fabricated. The second requirement was deemed advisable to insure that the repulsive vacuum force would not vary too rapidly as the distance between the sphere and the opening of an etched rectangular cavity changed. The final cavity

making measurements on the attractive Casimir force between the metallized sphere and a flat gold region and comparing these results to known values.

design selected was $0.1 \text{ um} \times 100 \text{ um} \times 1 \text{ um}$ (width x length x depth), with walls that are 0.1 um thick.

Wisconsin Center for X-Ray Lithography. The cavity arrays fabricated are $100\text{um} \times 100 \text{ um}$ square, with cavity walls 0.5 um deep, with thickness t between 250 and 300 nm thick, and cavity widths w between 125 and 150 nm . Calibration surfaces for the AFM were also included in the design. The overall test pattern design is shown in Figure 4, and a portion of one of the cavity arrays closest to the target design is show in Figure 5.

Figure 4. A SEM photograph of a portion of the test die, showing two 500 um square calibration patterns on the left, and two 500 um square cavity array regions on the right side. The center of each calibration pattern is a flat gold surface at the same level as the bottom of the cavities, surrounded by a gold surface at the level of the top of the array. The white regions on the right are the $100 \times 100 \text{ um}$ cavity arrays. The rectangular regions below the arrays indicate the nominal cavity width and wall thickness (50 x).






Figure 5. SEM photograph of portion of the gold cavity array. Each cavity is actually 100 μm long. The entire array is 100 μm wide (magnification 37,000; 15kV).

Theoretical Modeling

As mentioned previously, no QED method has been developed to compute the vacuum forces between separate conducting surfaces. There is no theoretical model for such a configuration of two separate surfaces in the literature; no QED calculation of Casimir forces have been done except for planar or slightly rough planar surfaces. Hence we developed

a heuristic model in which we assume the force on the sphere arises from two effects: 1) the attractive force due to the proximity of the sphere to the flat top surfaces of the cavity walls (parallel plate attractive Casimir force Eq 1.6), and 2) the repulsive forces on the sphere due to the cavity. The geometry of the experiment is shown in Figure 4.




Figure 6. Distance definitions, illustrating actual cavity depth (C_{CAV}), individual cavity width (w), cavity wall thickness (t), separation distance (d), local sag (s), and local separation distance (δ).

The sphere has radius R . The closest point or tangent point of the sphere is located a distance d above the tops of the cavities. The local separation distance is $\delta = d + s$, where s is the local sag of the spherical surface. The quantity δ can be given as a function of r , the horizontal distance from the tangent point of the sphere. The expression for the attractive Casimir force Eq. 0.6 is actually derived for an infinite parallel plate geometry in which the lateral dimensions of the plates are much bigger than the spacing between the plates. Since this condition is not met for spacing $d > t$ (width of cavity wall), we applied a correction factor to the attractive force, obtaining the expression for the attractive force:

$$F_{att}(r) = \frac{\pi^2}{240} \frac{\hbar c}{\delta(r)^4} A_w \left(\frac{t}{t + \delta(r)} \right) \quad 0.7$$

where we used the local separation distance $\delta(r)$, t is the thickness of the cavity wall, and A_w is the area of the tops of the cavity walls. The QED calculation of the repulsive Casimir force was for a closed, rectangular metallic box. Therefore we need a method to correct for the experimental geometry in which there is a gap at the top of the box. For the repulsive cavity force, we used the computed force for a closed cavity of width “w” with a depth equal to the actual depth (C_{CAV}), and multiplied it by an

approximate correction factor $K(r)$ suggested by theoretical analysis²⁰:

$$K(r) = \left(\frac{C_{CAV}}{C_{CAV} + \delta(r)} \right)^3 \left(\frac{w}{w + \delta(r)} \right) \quad 0.8$$

Eqs. 0.7 and 0.8 predict that both forces decay rapidly as the separation distance is increased. Rather than sum over individual cavities, we used an effective pressure distribution, which is an area-weighted combination of the cavity force and wall force. This provides a pressure distribution $p(r)$ over the surface of the sphere, where r is the radial coordinate. This pressure distribution depends on the geometry, including the separation distance, d .

To obtain the total force on the sphere, we integrated the pressure $p(r)$ on the bottom of the sphere due to the sum of the forces, i.e.,

$$F = \int_0^{R/2} 2\pi r p(r) dr \quad 0.9$$

Figure 7 shows a plot of the force F as a function of the separation d for a cavity array with 1) the target dimensions, namely $w = 0.1$ um wide with $t=0.1$ um thick sidewalls and 1.0 um deep, and 2) the best actual cavity dimensions fabricated, namely $w=0.125$ um wide cavities with $t=0.250$ um sidewalls, 0.5 um deep.




Figure 7. Force vs. distance d for a 100 um square cavity array with 1) the target dimensions, cavity width $w = 0.1$ um, wall thickness $t=0.1$ um, depth $C_{CAV} = 1.0$ um deep; 2) the best fabricated cavity, width $w=0.125$ um, thickness 0.250, depth 0.5 um deep. The two uppermost curves show the force for the sphere above the cavity arrays. The solid curve shows the attractive force for a flat surface, with no cavities. Conductivity corrections are included. The sphere diameter is 210 um.

Also shown for comparison, is the calculated force (labeled Flat in the figure) for the case of a sphere over a flat surface, i.e. with no cavities at all in the gold. The known correction factors for finite conductivity for the force between parallel plates were used except for the curve labeled infinite conductivity. For the lack of any other theory, the same conductivity correction factors were applied to the cavity geometry.

For the parallel plate case, the force is always negative or attractive, and the force decreases rapidly (Eq 1.7 integrated over the hemisphere, or approximately as $-1/\delta^4$) with the separation δ . The component with upward curvature in the cavity array curve is due to the repulsive force (Eq 1.8 integrated over the hemisphere, which goes approximately as $1/\delta^3$). Because of the more rapid inverse variation of the attractive force, the attractive force dominates at very small separations d , going to zero more rapidly than the attractive force. Hence the repulsive force dominates at larger separations, above about 0.1 um. From QED we expect that smaller cavities would give larger repulsive forces; and thinner walls give smaller attractive forces. The predictions of the model show this desired behavior, and the model calculations appear to go smoothly into these known curves for flat surfaces. It appears that the repulsive force is on the edge of detectability for the cavities fabricated.

CONCLUSIONS

Quantum electrodynamics, which has made predictions which have been verified to 1 part in 10^{12} , predicts the existence of a large, ubiquitous, zero-point vacuum energy density in empty space. The question arises: Can we make use of this energy in some way to facilitate space travel, such as energy generation, propulsion, or creation of wormholes? It is too early to determine if such developments are possible or to be able to clearly determine the role of vacuum energy in future space applications. Our investigation begins with what we do know about vacuum energy, and extends those boundaries. We know that QED predicts that as a consequence of this energy, an attractive force will exist between uncharged, parallel, metal plates. In the last few years, accurate measurements have confirmed the existence of this force. QED also predicts the existence of repulsive forces in small rectangular metal boxes in which one dimension is much less than at least one of the other two dimensions. Although no one has done a rigorous calculation, it appears probable, based on theory, that a repulsive

force should exist between two separate surfaces that closely approximate such a closed box. Vacuum forces that are repulsive because of the geometry have never been observed.

We have designed an experiment to measure repulsive vacuum forces. A model has been developed to predict the vacuum force on a metallized sphere attached to a cantilever on an AFM when the sphere is brought to within nanometer distances of an array of gold cavities. An AFM that operates in vacuum at 10^{-4} torr has been constructed to perform the experiment. Based on our model calculations, it appears we should be able to measure repulsive vacuum forces using the AFM, provided the cavities have small enough dimensions. Based on our model calculations (Figure 7) the cavities fabricated to date (125 nm width, 245 nm wall thickness) have dimensions that may be too large to provide a clear indication of a repulsive force. We need to utilize cavities with dimensions of approximately 100 nm width, 100 nm wall thickness in order to have a clear indication of repulsive forces. The University of Wisconsin Center for NanoTechnology is currently upgrading one of its synchrotron exposure systems in order to provide features of this size.

If we can obtain repulsive as well as attractive vacuum forces by a suitable choice of geometry, we are one step closer to being able to design a variety of novel MEMS devices using vacuum energy that could assist in attaining some of the NASA objectives for space travel.

Acknowledgements: GJM would like to thank Marc Millis and the NASA Breakthrough Propulsion Physics Program, MEMS Optical Inc., and Quantum Fields LLC for their support of this program, and Robert Forward, Peter Milonni, Carlos Villarreal, Gabriel Barton, and Michael Serry for helpful conversations. MG and LS would like to thank Molecular Imaging Inc. for their support of the development of a vacuum AFM. We would like to thank Hui Liu for taking SEM photographs and Jeff Meier for assistance with microfabrication.

¹ J. Maclay, "A Design Manual for Micromachines using Casimir Forces: Preliminary Consideration," **PROCEEDINGS OF STAIF-00** (Space Technology and Applications International Forum-2000, Albuquerque, NM, January, 2000), edited by M.S. El-Genk, AIP Conference Proceedings, American Institute of Physics, New York 2000. Published in hardcopy and CD-ROM by AIP.

-
- ² P. Milonni, *The Quantum Vacuum* (Academic Press, San Diego, CA, 1994).
- ³ M. Visser, pp 81-87, *Lorentzian Wormholes: From Einstein to Hawking*, (American Institute of Physics, New York, 1996).
- ⁴ Chan H B, Aksyuk V A, Kleiman R N, Bishop D J, and Capasso F, Quantum mechanical actuation of microelectromechanical systems by the Casimir force Science **291**, 1941-44 (2001).
- ⁵ M. Serry, D. Walliser, J. Maclay, “The anharmonic Casimir oscillator (ACO)- the Casimir effect in a model microelectromechanical system,” J. Microelectromechanical Systems **4**, 193-205 (1995).
- ⁶ J. Maclay, “Analysis of zero-point electromagnetic energy and Casimir forces in conducting rectangular cavities,” Phys. Rev. A, 61, 052110(2000)
- ⁷ G. Barton, “Perturbative Casimir energies of dispersive spheres, cubes, and cylinders,” J. Phys. A.: Mathematical and General **34**, 4083-114 (2001). Also personal communication from Dr. Barton, Boston 6/01.
- ⁸ R. Van Dyck, Jr., P. Schwinberg, and H. Dehmelt, Phys Rev. Lett. **59**, 26 (1987).
- ⁹ P. Milonni, p. 108, *The Quantum Vacuum* (Academic Press, San Diego, CA, 1994).
- ¹⁰ J. Bjorken, and S. Drell, *Relativistic Quantum Fields*, McGraw-Hill, New York (1965)
- ¹¹ C. Misner, K. Thorne, J. Wheeler, p. 1203, *Gravitation*, W.H.Freeman, SanFrancisco (1973).
- ¹² [30] E.Elizalde and A.Romero, “Essentials of the Casimir effect and its computation,” Am. J. Phys. **59**, 711-719 (1991). Also see reference 2, p.54.
- ¹³ S. Lamoroux., “ Measurement of the Casimir force between conducting plates,” Physics Review Letters, **78**, 5-8 (1997).
- ¹⁴ Mohideen, U., Anushree, Roy, “Precision Measurement of the Casimir Force from 0.1 to 0.9 micron”, Physical Review Letters, **81**, 4549 (1998).
- ¹⁵ See ref 2 for a discussion. Also it should be mentioned that these phenomena can be interpreted in an equivalent way in which we postulate that the effects are due to the fluctuational energy in the atoms involved in these experiments.
- ¹⁶ J.D. Jackson, *Classical Electrodynamics*, Wiley, New York (1962)
- ¹⁷ P. Plunian, B. Muller, W. Greiner, “The Casimir Effect,” Physics Reports (Review Section of Physics Letters) **134**, 2&3, pp. 87-193 (1986).
- ¹⁸ L. Brown, and J. Maclay, “Vacuum Stress between Conducting Plates: an image solution,” Phys. Rev. **184**, 1272-1279 (1969). We assume absolute zero temperature.
- ¹⁹ J. Maclay, R. Ilic, M. Serry, P. Neuzil, “Use of AFM (Atomic Force Microscope) Methods to

Measure Variations in Vacuum Energy Density and Vacuum Forces in Microfabricated Structures,” NASA Breakthrough Propulsion Workshop, Cleveland, Ohio, May, 1997.

²⁰ D.Deutsch and P. Candelas, “Boundary effects in quantum field theory,” Phy. Rev. D **20**, 3063-3080 (1979).

SEARCH FOR EFFECTS OF ELECTRIC POTENTIALS ON CHARGED PARTICLE CLOCKS

Harry I. Ringermacher^{*}
 Mark S. Conradi and Caleb D. Browning[†]
 Brice N. Cassenti[‡]

ABSTRACT

Results of experiment to confirm a theory that links classical electromagnetism with the geometry of space-time will be described. The theory, based on the introduction of a Torsion tensor into Einstein's equations and following the approach of E. Schrödinger, predicts effects on clocks attached to charged particles, subject to intense electric fields, analogous to the effects on clocks in a gravitational field. We show that in order to interpret this theory, one must re-interpret all clock changes – both gravitational and electromagnetic – as arising from changes in potential energy and not merely potential. The clock is provided naturally by proton spins in hydrogen atoms subject to Nuclear Magnetic Resonance trials. No frequency change of clocks was observed to a resolution of 6×10^{-9} . A new “Clock Principle” was postulated to explain the null result. There are two possible implications of the experiments: (a) The Clock Principle is invalid and, in fact, no metric theory incorporating electromagnetism is possible; (b) The Clock Principle is valid and it follows that negative rest mass cannot exist.

INTRODUCTION

The goal of the present work is to investigate an electromagnetic alternative to exotic physics for the purpose of coupling matter to space-time. Electromagnetic forces have distinct advantages. They are 10^{40} times stronger than gravity. They can be manipulated at will. Resources to create intense fields of virtually any geometry are readily available. However, there is currently no accepted theory linking electrodynamics directly with the geometry of space-time other than to curve it via extremely high energy densities. The mainstream approach taken is “bottom up”, attempting to unite all forces in the context of quantum gauge field theory which has to-date been successful in unifying the weak and electromagnetic

forces and describing the strong force in what is known as “the standard model”. Gravity and therefore space-time geometry remains isolated from the internal geometry of gauge theory. If such a theory could be found and even its simplest predictions tested and verified, then there would be hope that electromagnetic coupling to space and time might be possible. This could lead to new interpretations and possibly new effects in gravitation and electromagnetism.

The experiments described in this work measure the predictions of a theory¹, linking space-time geometry and electrodynamics. This is grounded upon E. Schrödinger’s later works on gravitation theory². In his work, Schrödinger attempted to link electromagnetism to geometry through a non-symmetric affine connection (Torsion tensor). He failed at the attempt, primarily because of an error of oversight. The theory upon which the present work is based corrects this error³, resulting in the definition of a new type of affine connection – an electrodynamic connection – that precisely matches Schrödinger’s concepts.

THEORY

We describe a simplified theory and shall only write the new field equations and their solutions for the present case. The theory is summarized in the BPP final report⁴. The governing equations are:

$$G_{\mu\nu} = -\frac{\kappa}{2} u^\sigma (F_{v\sigma;\mu} + F_{\mu\sigma;v}) \quad (1)$$

$$F^{\mu\tau}_{;\tau} = 0 ; \quad F_{\mu\nu;\sigma} + F_{v\sigma;\mu} + F_{\sigma\mu;v} = 0 \quad (2)$$

$G_{\mu\nu}$ is the Einstein tensor. $F_{\mu\nu}$ is the Maxwell electromagnetic field tensor. u^λ is the test particle 4-velocity and $\kappa = -e/mc^2$, the charge/mass ratio of the test particle. Greek indices range from 0, the time index, to 1,2,3, the space indices. Equation (1) is the modified Einstein equation including Electrodynamic Torsion. Equations (2) are the usual covariant Maxwell equations. Selecting a metric for an appropriate

^{*} KRONOTRAN Enterprises LLC, Delanson, NY 12053

[†] Washington University, St. Louis, MO 63130

[‡] United Technologies Research Center, E. Hartford, CT 06108

Copyright ©2001 by Harry I. Ringermacher. All rights reserved.
 Published by the American Institute of Aeronautics and Astronautics
 with permission

⁴ General Electric Corp. R&D Center, Schenectady, NY 12301

geometry results in a set of solvable differential equations coupling the metric and electromagnetic field variables.

Ideal Experiment

The theory predicts that a particle of charge e , and mass m , immersed in a suitable electric field but unshielded and supported will see, in its rest frame, a time differing from the proper time of an external observer arising from the electrostatic potential at its location. In general, from the theory, the time shift in a clock interval is related to the potential difference between two points:

$$\frac{d\tau_2}{d\tau_1} = 1 + 2\kappa(\varphi_2 - \varphi_1) \quad (3)$$

This equation is exactly analogous to that for the gravitational red shift.

One possible clock for such a test is Nuclear Magnetic Resonance. A proton placed in an intense electric field within a radio-frequency transverse field "H₁" coil aligned orthogonally to a uniform magnetic field, H₀, is resonant at the Larmor frequency, $\omega = \gamma H_0$, where the gyromagnetic ratio, γ , for the proton spin is proportional to e/m . We thus have a natural clock. From eq. (3) we expect the proton's clock frequency to depend on its relative positions r₁ and r₂:

$$\omega(r_2) = \omega(r_1)[1 + 2\kappa(\varphi_2 - \varphi_1)] \quad (4)$$

The Larmor field distribution is then given by

$$H(r_2) = H(r_1)[1 + 2\kappa(\varphi_2 - \varphi_1)] \quad (5)$$

From this it is straightforward to calculate the NMR lineshape and shift that will result when the electric field is turned on as compared to the field off. Under ideal circumstances, for a supported proton in a 8T magnetic field with a 5kV/cm electric field, a line shift and broadening of approximately five parts per million is expected. The expected NMR lines are modeled in the "Experiments" section.

Present Approach

In practice it is experimentally difficult to "support" a charged particle. Generally, this can be accomplished electromagnetically, but then, by definition, the electric field and force at the particle location must vanish since the charge does not accelerate. The approach we have chosen uses the proton in a hydrogen atom. It is supported electromagnetically. The consequences of this approach

will be the main subject of the conclusions of this work. A detailed description of the three experiments performed follows.

EXPERIMENTS

Three experiments were performed: A temporally and spatially constant potential applied to the proton in the hydrogen atom; A time-varying but spatially constant potential applied to the proton in the hydrogen atom; A hydrogen atom physically displaced through an intense electric field.

Experiment 1 – Constant potential

A 354 MHz, 8.4 Tesla NMR system was chosen for the experiments. This magnet has a field homogeneity of 0.1 ppm or about 35 Hz, more than sufficient to resolve small effects. The proton sample was Benzene. The initial experiment was a simple free induction decay (FID) with the E-field on vs. off. The sample was enclosed in a 2mm thick aluminum can (Fig. 1) placed at high potential. Thus the E-field will vanish in the interior of the can at the sample but there will remain a constant potential. The voltage terminal (sample chamber) could be set + or - with respect to ground and the NMR FID was monitored.




Figure 1. NMR "can" arrangement. External magnetic field, H₀, is perpendicular to the radio frequency field, H₁, applied through a coil wrapped around the proton sample, S. The 5kV electric field is applied outside the can leaving a constant 5kV potential inside.

A field homogeneity of 10 ppb was achieved for the 8.4T field, sufficient to resolve the smallest predicted effects. The observed line shift, was $\Delta\nu/\nu \leq 0 \pm 1.0 \times 10^{-9}$, consistent with a null prediction of both classical E&M and the present theory. The null result is consistent with both classical theory and the present one.

Experiment 2 - Time-varying potential

The second experiment, a time-varying potential applied to the proton in the hydrogen atom, was initially expected to produce an effect. However, it was found during the course of this program that this theory was also invariant under a pure time-varying potential⁵, a result consistent with the classical “Lorentz gauge condition” which is the relativistic generalization of the Coulomb gauge condition. That is, it states that the potential is also arbitrary to an additive time-derivative of a scalar. Thus, a null result was also expected and was observed. A “spin-echo” experiment was performed to observe any phase shift introduced by the time varying electric field. Extreme care was taken to calibrate the system to ensure that any small phase shift could be identified and that stray currents would not affect the data. Upon application of a 5kV step-function (20 ms risetime) between the aluminum can and ground in both turn-on and turn-off modes, the observed shift in the NMR line was $\Delta\nu/\nu \leq 0 \pm 1.0 \times 10^{-11}$.

Experiment 3 - Physical displacement of Hydrogen atom through high electric field

For the third and final experiment, hydrogen in benzene at room temperature was gravity-flowed between two electrodes, an upper one at ground potential and a lower one at +5000 V, both situated in the NMR coil in the external 8.4T magnetic field while NMR was performed with the electric field on and off. The electrodes were copper discs placed in the 3mm I.D. of glass tubing and connected to a high voltage source through glass/epoxy seals. The electrode spacing was 1.0 cm giving an average electric field of 5000V/cm. The 2-turn NMR coil diameter was 1.5 cm ensuring that the HV region was inside the coil. The coil was untuned to avoid radiation damping since the signal was already very large. Figure 2 shows the NMR coil system arrangement and a photo of the open chamber.

An NMR FID experiment was performed with and without flow. A 20-30 ms T_2 was obtained by careful adjustment of the B-field shim coils. The NMR line and effects of flow without the presence of an E-field were modeled. The T_2 value gives a line width of approximately 17 Hz ($\Delta f = 1/\pi T_2$). Figure 3 shows the measured NMR line with voltage off as a function of flow velocity through the coil varying from 3 cm/s to 50 cm/s. Since the FID has a time constant of 20-30 ms, the proton must stay in the H1 field at least this long in order to contribute a significant time-shift signal arising from the maximum 5kV potential change. This corresponds to a flow speed of 15-30 cm/s, the mid-range of the chosen flows. Note that the measured shift with zero volts is approximately 10 Hz, from 0 to 50 cm/s flow rate, in approximate agreement with the theoretical calculation, Figure 3. Figure 4 shows the predicted lines for a maximum potential of 5000 Volts. The shift is at least ten times that for zero volts.

For the flow experiment, the NMR line was obtained for voltage off, voltage on, and a dummy voltage on (voltage on but HV cable disconnected). The probe voltage was discharged when the dummy experiment was performed. Figure 5 shows data for flows of 0, 3, 6, 12, 25 and 50 cm/s with the voltage applied .

We note in Figure 5 that there is no change in the line positions greater than the FFT resolution of ± 2 Hz corresponding approximately to a line shift of $\Delta\nu/\nu \leq 0 \pm 6.0 \times 10^{-9}$, at least a hundred times smaller than the predicted shift of Figure 4. Figure 3 is therefore representative of the results with applied field as well.

When the experiment was completed, the probe was disassembled and carefully inspected to ensure that all voltage connections were secure.




Figure 2. Diagram and photo of NMR coil and flow arrangement for E-field experiment




Figure 3. Measured NMR lines (left) and calculated (0,12,25,50 cm/s) for zero E-field and flows from 0 - 50 cm/s.




Figure 4. Calculated NMR lines for 5000V/cm E-field for flows of 0, 12, 25 and 50 cm/s.




Figure 5. Measured variation in NMR line at 0, 6, 25 and 50 cm/s flow with applied voltage.

CONCLUSIONS

Experimental Conclusions

All three experiments showed null results. The first experiment, constant potential applied to the proton in the hydrogen atom, produced a null result as expected and explained earlier. The second experiment, a time-varying potential applied to the proton in the hydrogen atom, also produced a null result as expected and explained. From the second experiment we have learned that the present theory is relativistically Gauge Invariant in the context of Maxwell's equations – as it should be.

The third experiment, physical displacement of a proton in a hydrogen atom through an electric field, was expected to produce a frequency shift of 5ppm in the NMR line. No change was observed to high precision. This could be explained as follows. When a hydrogen atom is placed in an electric field, classically, only one effect is produced – an induced dipole moment of the atom resulting from the stretching of the electron orbital. Otherwise, no work is done on the hydrogen atom (except for the exceedingly small amount involved in stretching) since the work done on the proton is cancelled by the opposite work done on the electron(due to its negative charge) in the electric field. Furthermore, it is straightforward to show⁴ that no work is actually done on the proton itself since the electric

force on it is precisely balanced by an opposing force, in the field direction, from the electron.

Implied Conclusions

If we substitute potential energy change and therefore work done rather than simple potential in the metric, then the fact that no work is done on the proton because of the electron's influence can explain the null result. We now discuss the relation of work to the metric.

Relation of the Metric to the Lagrangian and Work Done

Let \tilde{L} be defined as the specific Lagrangian or Lagrangian per unit rest energy,

$$\tilde{L} = \frac{(T - V)}{mc^2} \quad (6)$$

where T , V and m are the kinetic energy, potential energy and rest mass of the test particle respectively.

Suppose we are at rest at some height z in a gravitational field, so that $T=0$ and $V=mgz$, then $\tilde{L}=gz/c^2$. We have shown for weak fields and nonrelativistic speeds that the proper time element can be written in terms of the Specific Lagrangian and coordinate time element as:

$$d\tau = \sqrt{1 - 2\tilde{L}} dt \quad (7)$$

This describes the behavior of clocks. Consider two positions in the gravitational field z_1 and $z_2 = z_1 + h$. Assume a proper time interval $d\tau_1$ at z_1 and $d\tau_2$ at z_2 . Then, for weak fields:

$$\frac{d\tau_2}{d\tau_1} = 1 - \frac{gh}{c^2} \quad (8)$$

We can rewrite this in terms of potential energy and Lagrangian rather than potential;

$$\frac{d\tau_2}{d\tau_1} = 1 - \frac{mgh}{mc^2} = 1 - (\tilde{L}_2 - \tilde{L}_1) = 1 - \Delta\tilde{L}. \quad (9)$$

In general, a variation in the Specific Lagrangian results in a change in the clock rate. Referring more concisely to the metric for a single particle, p , we may express the Lagrangian change more clearly in terms of the net conservative work, W_c , done on p .

$$\Delta\tilde{L}_p = 1 - \frac{(\Delta T - \Delta V)}{m_p c^2} = \frac{\Delta T_p + W_c}{m_p c^2} \quad (10)$$

The weak-field, non-relativistic, metric for a given particle, p , acted upon by forces and hence net conservative work, W_{cp} , done upon it by all other particles in the field is given by

$$d\tau_p^2 = \left(1 - \frac{2W_{cp}}{mc^2}\right) c^2 dt_p^2 - dx_p^2 - dy_p^2 - dz_p^2, \quad (11)$$

We have rewritten the metric in this way because potential is not well-defined except through potential energy and work, where it is defined as work per unit mass in gravitation and work per unit charge in electromagnetism.

The Clock Principle

In the previous section we found a simple Lagrangian formulation that places gravitation on an equal footing with our theory in regards to changes in the temporal portion of the metric. The Lagrangian formulation deals with kinetic and potential energy changes. Clocks raised in a gravitational field are at rest in the two positions and can be slowly moved between them. Thus the Lagrangian becomes simply the negative change of potential energy (work done on) of the clock, moved from the lower position to the upper. However, we must be careful and can no longer use the word "clock" loosely. When we refer to "clock" henceforth, we mean the *mechanism of the clock*. Thus we mean that work is performed on the *mechanism*. Clearly all clock mechanisms are driven by energy changes. What is not as obvious is that the mechanism of any clock must reflect the proper time variations in a gravitational field. For example, the mass-spring mechanism of a simple clock must somehow change with different heights in a gravitational field. Similarly, a pendulum clock must exhibit changes in its mechanism. Even an atomic clock is subject to this consideration. This brings us to the *first clock postulate*:

I) Every clock has a mechanism which must be held accountable for observed changes in its measurement of time.

We shall now examine the relationship between work and changes in clock time in a gravitational field. We use the notation "nc" to mean nonconservative and "c" for conservative forces.

Einstein Rocket -- Equivalence Principle for clock changes in a Gravitational Field

We shall employ the famous Einstein rocket “gedanken-experiment” to demonstrate our concepts. Consider a rocket lifting a mass, m , by a stiff wire in a gravitational field. Let us suppose that we adjust its thrust to precisely oppose the gravitational pull on the mass. The rocket-mass system is now balanced and hovers, for example, at the surface of the earth. An arbitrarily small external force, $\vec{\epsilon}$, may now raise the system to a height, h . This is shown in Figure 7. The force $\vec{\epsilon}$ does no work on the system since it can be made arbitrarily small. However, upon rising a height h , due to the infinitesimal assistance of the guiding force, the rocket does work on the mass m . The potential energy of the mass is V_1 at the surface and V_2 at height h .




Figure 7. Balanced force configuration of a rocket-mass system

In this situation, the nonconservative force is that of the rocket. This is equal and opposite to gravity. We make the following definitions:

$$z_2 - z_1 = h \quad ; \quad d\bar{s} = dz\hat{k} \quad ; \quad \vec{g} = -g\hat{k} \quad (12)$$

Then the non-conservative work performed on the mass m by the rocket is

$$W_{nc} = \int_{z_1}^{z_2} \vec{F}_{nc} \cdot d\bar{s} = mgh \quad (13)$$

and the conservative work done on m by gravity, opposing the rocket is:

$$W_c = -mgh = -(V_2 - V_1) = -\Delta V \quad (14)$$

Einstein rocket replaced by negative mass

Let us now reconsider the rocket of Figure 7. We can replace the rocket by negative mass equal in magnitude to the lower positive mass (Fig. 8). Earth’s gravity repels negative mass and thus an amount ($-m$) will precisely balance the force of attraction on m , a situation equivalent to the rocket.




Figure 8. Balanced force configuration for rocket replaced by negative mass

We evaluate the work done in this situation. There are two forces acting on m , \vec{F}_g and \vec{F}_{-m} . This time *both* forces are conservative. The total work done on the mass m is zero because there is no change in the kinetic energy, since $\vec{F}_{-m} = -\vec{F}_g$.

$$W_{total} = \int \vec{F}_g \cdot d\bar{s} + \int \vec{F}_{-m} \cdot d\bar{s} = \Delta T = 0 \quad (15)$$

However, now the total work done on m is conservative since, unlike with the rocket, there are no non-conservative forces acting. The rigid wire transmits the conservative force from $-m$. Thus we find:

$$W_c = W_g + W_{-m} = 0 \quad (16)$$

This model of a massive neutral dipole is exactly equivalent to an electric neutral dipole such as the proton and electron of a hydrogen atom in an electric field. In the latter case, the two forces acting on the proton, the external electric field and the opposing force of the electron are both conservative. The total conservative work on the proton is zero when the atom is moved a distance h through a known potential difference. That is precisely our experiment. There was no clock change observed.

This leads us to the **second and third Clock Postulates**:

2) External conservative work done on a clock mechanism is responsible for changes in its rate when the motion of the clock can be neglected.

3) When the clock is not at rest, the change in its rate arises more generally from the change in the Specific Lagrangian for the mechanism.

The third postulate takes into account the change when there is kinetic energy present.

These postulates reconcile the observations and considerations of the present experiment with those observed for gravity. Therefore one should not expect to see a clock change in the imagined massive dipole

experiment since there was no net conservative work done on m .

Inconsistency between General Relativity and existence of a neutral mass dipole

Now consider a clock on the mass m attached to the rocket. The mass m might itself be part of a clock mechanism. Upon reaching height h , the clock will change. But the mass m cannot distinguish between the pull of the rocket on the wire from that of the negative mass. In the case of the negative mass, the Clock Principle states that since no net conservative work is done on m , no clock change should be observed. So, there is a conflict here. It follows that:

Since we assume GR to be true, either the Clock Principle is false or a mass dipole cannot exist as the analogue of a charge dipole. Thus, if the clock principle is true, negative mass cannot exist. If the clock principle is false, then work and energy change are not related in general to time change. It follows that a metric theory of forces other than gravitation cannot be constructed.

Clock Conclusions

It is generally agreed that in order for a clock to indicate the passage of time, energy change in the mechanism is required. Thus, it appears that space is coupled to time through energy. When a clock is raised in a gravitational field, external work is performed on the mechanism. This permits a time change. If $W_c = 0$ so that applied forces opposing conservative forces are also conservative, then there will be no observed clock change.

We have shown that if the clock principle is correct then **negative mass cannot exist**. This does not preclude the existence of negative energy since the general energy-momentum relation has two roots,

$$E = \pm \sqrt{p^2 c^2 + m^2 c^4}$$

of which the positive root, in the rest frame of a particle of mass m ($p=0$), is the famous Einstein energy-mass relation.

ACKNOWLEDGEMENTS

This work was supported by the NASA Glenn Research Center under Contract NAS3-00094. We wish to thank Marc Millis for having the daring and fortitude to envision and create the BPP program and support this effort as part of it. We would like to thank Prof. Larry Mead for supportive discussions revolving around this new view of General Relativity. We also

wish to thank Dan Leopold for his advice and support of the experiment.

REFERENCES

1. Harry I. Ringermacher and Brice N. Cassenti, *Search for Effects of an Electrostatic Potential on Clocks in the Frame of Reference of a Charged Particle*, Breakthrough Propulsion Physics Workshop, (NASA Lewis Research Center, Cleveland, August 12-14, 1997, NASA publ. CP-208694 , 1999, Millis & Williamson, ed).
2. E. Schrödinger, *Space-Time Structure*, (Cambridge University Press, 1986).
3. H. I. Ringermacher, *Classical and Quantum Grav.*, **11**, 2383 (1994).
4. NASA-GRC final report, Contract NAS3-00094.
5. “Mathematica” was used to test for invariance of the Modified Einstein equations in this theory under a purely time-dependent potential in g_{00} . The solutions to the equations were found to be invariant under an additive time-dependent potential.

Tests of Mach's Principle with a Mechanical Oscillator^{*}

John G. Cramer[†], Damon P. Cassissi, and Curran W. Fey

Department of Physics, Box 351560, University of Washington, Seattle, WA 98195-1560, USA

Abstract

James F. Woodward has made a prediction, based on Sciama's formulation of Mach's Principle in the framework of general relativity, that in the presence of energy flow the inertial mass of an object may undergo sizable variations, changing as the 2nd time derivative of the energy. We describe an attempt to observe the predicted effect for a charging capacitor, using a technique that does not require a reactionless force or any local violation of Newton's 3rd law of motion. We attempt to observe the effect of the mass variation on a driven harmonic oscillator with the charging capacitor as the oscillating mass. Positive and negative phase shifts in the oscillator motion with respect to the driving force are predicted to result from appropriately programmed inertial mass variations. The phase shift is constant, so that data may be accumulated over a very large number of oscillation cycles to insure high precision in the phase shift determination. We report on the predicted effect and the design and implementation of the measurement apparatus. At this time, however, we will *not* report on observations of the presence or absence of the Woodward effect.

Introduction

This is a status report on a new experiment to test a prediction based on general relativity and Mach's Principle, which has been supported by the Breakthrough Propulsion Program of NASA.

Einstein's Principle of Equivalence, a cornerstone of general relativity, asserts the exact universal identity of inertial mass and gravitational mass. However, the origins of inertia and its connection to gravitational mass remain obscure. Mach's Principle, the idea that inertia originates in the gravitational interaction of massive objects with the distant matter of the universe, is an attempt to unify gravitational and inertial mass, but it is not a part of general relativity. Dennis Sciama [1,2] attempted to improve this situation by showing that, for sufficiently symmetric and homogeneous universes, the gravitational interaction of massive objects with distant matter leads to an acceleration-dependent force, i.e., inertia.

James F. Woodward [3,4] extended Sciama's calculations by introducing energy flow (e.g., the

energy flowing to a charging capacitor) into the gravitating system. He demonstrated that the equations acquire extra transient contributions in Sciama's inertia term that are proportional to $1/G$ (Newton's gravitational constant) and therefore are quite large. The implications of this work are: (a) that it may be possible to modify inertia, and (b) that it may be possible to demonstrate the validity of Mach's Principle with a tabletop experiment.

Woodward and his students [4-7] have attempted to observe the predicted inertia-variation effect by accelerating a mass-varying object so that it produces a reactionless force. To illustrate this, assume that an inertia-varying test mass is accelerated to the right when it has low inertia and to the left when it has high inertia. In this circumstance, it is argued, the reaction forces of the two accelerations are unequal and one might expect the net reactionless force to "row" the system to the right. Woodward's group reports [7] using a sensitive torsion balance to observed small reactionless forces at magnitudes that are near the limits of their sensitivity and about five orders of

Copyright ©2001 by John G. Cramer. All rights reserved. Published by the American Institute of Aeronautics and Astronautics with permission.

* Supported in part by the National Aeronautics and Space Administration.

† E-mail address: cramer@phys.washington.edu

magnitude smaller than the predicted effect (see the calculations below.)

Unfortunately, this scheme for observing the predicted inertia variation appears to be at odds with the relativistically invariant form of Newton's 2nd law of motion:

$$\vec{F} = d\vec{p}/dt = m d\vec{v}/dt + \vec{v} dm/dt \quad (1)$$

Since the inertial mass m of the test body is expected to vary with time, the last term of Eqn. (1) cannot be ignored. It is not surprising, in view of Newton's 3rd law of motion, that for any closed cycle of acceleration and variation of the inertial mass around a central value, the force contribution from the $\vec{v} dm/dt$ term is found to precisely cancel the supposed "reactionless force" arising from the $m dv/dt$ term, leading to a net force of zero for the overall system.

From this simple calculation, it appears that reactionless force searches are *not* good tests of the proposed effect. There remains the question of whether the Woodward inertia variation is indeed present in a system with energy flow. We have found, as will be described below, that a mechanical oscillator, driven at resonance, with its mass programmed to vary at the drive frequency, shows sensitive variations in drive-to-response phase and amplitude, depending on the relative phase between the mass variation and the oscillator drive.

Theory

Woodward has shown [5] that the relativistically invariant wave equation, in the simplest approximation and expressed as a function of an overall scalar gravitational potential ϕ , has the form:

$$\nabla \cdot \phi - (1/c^2)(\partial^2 \phi / \partial t^2) = \phi = 4\pi G \rho_0 + (\phi / \rho_0 c^2)(\partial^2 \rho_0 / \partial t^2) - (\phi / \rho_0 c^2)^2 (\partial \rho_0 / \partial t)^2 \quad (2)$$

where G is Newton's gravitational constant, ρ_0 the rest mass density, and c the speed of light. This field equation is obtained only if one assumes, as suggested by Mach's Principle, that the local energy density of matter is equal to the matter density times ϕ . Since Mach's Principle demands that $\phi = c^2$ when measured locally, this constraint is equivalent to asserting that $E = mc^2$. Additional terms would be present in this equation were it not for the fact that, as a consequence of Mach's principle in this approximation, $\phi = c^2$.

In writing Eqn. (2), Woodward neglects a term of the form $c^{-4}(\partial \phi / \partial t)^2$ because it is always small, given its c^{-4} coefficient that is not compensated for by any factor of ϕ in the numerator. Combining the last three terms of Eqn. (2) into an effective mass density $\rho(t)$ and solving for this quantity gives the time-dependent effective mass density as:

$$\rho(t) \approx \rho_0 + (1/4\pi G)[(\phi/\rho_0 c^2)(\partial^2 \rho_0 / \partial t^2)] - (1/4\pi G)[(\phi/\rho_0 c^2)^2 (\partial \rho_0 / \partial t)^2]. \quad (3)$$

The second term in Eqn. (3) has the form $(1/4\pi G)[(\phi/\rho_0 c^2)(\partial^2 \rho_0 / \partial t^2)]$. This time-dependent fluctuation in the inertial mass can be both positive and negative when ρ_0 undergoes periodic time variations, e.g., when a varying flow of mass-energy is present. This is the inertia-varying term of interest.

In the present work we will ignore the last time-dependent or "wormhole" term, which has the form $-(1/4\pi G)[(\phi/\rho_0 c^2)^2 (\partial \rho_0 / \partial t)^2]$. This mass term is always negative or zero and for sinusoidal variations is about 0.1% or less of the other terms.

If a capacitance C is driven by a voltage source with time dependent potential $V(t) = V_0 \sin(\omega t)$, then the energy in the capacitor, assuming dissipative and inductive effects can be neglected, is $U(t) = 1/2 C V(t)^2 = 1/2 C V_0^2 \sin^2(\omega t)$. The second time derivative of this stored energy divided by c^2 (to convert it to a mass) is $d^2 U / dt^2 = C V_0^2 \omega^2 \cos(2\omega t) / c^2$. This is the $(\partial^2 \rho_0 / \partial t^2)$ factor in Woodward's Eqn. (3). The corresponding time dependent variation in inertial mass, assuming that $\phi = c^2$, is then:

$$dm(t) = 1/(4\pi \rho_0 G c^2) C V_0^2 \omega^2 \cos(2\omega t). \quad (4)$$

We will use this form for the variation in inertial mass in the analysis that follows.

To give this prediction a scale, let us assume that $c = 2.998 \times 10^8$ m/s, $G = 6.672 \times 10^{-11}$ m³/kg s², $\rho_0 = 2,000$ kg/m³, $C = 9.3 \times 10^{-9}$ F, $V_0 = 2,000$ V, and $\omega = 2\pi \times 1,000$ Hz. With these values, we find that:

$$dm(t) = 9.7 \text{ mg} \times \cos(2\pi \times 2,000 \text{ Hz} \times t). \quad (5)$$

In other words, under these conditions, which should be realizable in the experiment described here, the inertial mass of the capacitance is predicted to vary by about ± 10 milligrams at *twice* the capacitor charging frequency, or 2,000 Hz. If the mass of the capacitor and its holder were about 1 g, this would represent a mass variation of about $\pm 1\%$. Such a mass variation would have large observable consequences. However,

we note that Woodward [4] has made arguments involving mobile charges to explain why the actual variation in the inertial mass may be orders of magnitude smaller than that predicted by simple calculations and more consistent with his reported observation of very small reactionless forces.

It is also of interest to consider the maximum current flow that is necessary to charge the capacitor in the manner assumed above. The charge on the capacitor is $q=C V(t)$, so the charging current is $i(t) = C dV/dt = C V_0 \omega \cos(\omega t) = 116 \text{ mA} \times \cos(\omega t)$ for the specified conditions. It turns out that a high voltage power supply/amplifier capable of delivering an audio-frequency peak current of a few hundred millamps at a few kilovolts is very expensive (~\$14,000) and represent the most costly component required for the present test of the Woodward effect.

A Driven Mass-Varying Oscillator

We test for the presence of the Woodward effect by using the capacitor as the mass in a system that forms a driven mechanical mass-and-spring oscillator with an undriven resonant frequency of ω_0 . Such an oscillator is shown schematically in Fig. 1




Figure 1. Schematic mass-and-spring mechanical oscillator with time varying mass $m+dm(t)$ and restoring-force spring constant k . The system is assumed to have a dissipative damping force of $-b dx/dt$.

The oscillator is driven at its resonant frequency $\omega_d = (\omega_0^2 - b/2m)^{1/2}$ with a voice coil actuator and audio amplifier. At the same time, we charge the capacitor sinusoidally, using approximately the parameters

specified above, at a frequency of $\omega_0/2$ so that, in the presence of the Woodward effect, the capacitor's inertial mass should vary at frequency ω_d .

The inhomogeneous non-linear differential equation describing such a system is:

$$F_d \cos[\omega_d t] = kx(t) + [b + \mu'(t)]x'(t) + [m + \mu(t)]x''(t), \quad (6)$$

where $x(t)$ is the motion of the capacitor, F_d is the magnitude of the driving force, ω_d is the angular frequency of the driving force, k is the Hooke's law restoring force constant, b is the damping constant representing dissipative forces in the system, m is the average mass of the capacitor and associated structure, and $\mu(t)$ is the time-dependent mass variation due to the Woodward effect. Note that the $\mu'(t)x'(t)$ term in Eqn. (6) arises from the $v dm/dt$ term in Eqn. (1).

We can replace the spring constant k with $m \omega_0^2$ and replace $\mu(t)$ with $\mu_0 \cos(\omega_d t + \phi_m)$, which assumes that we have arranged the mass variation to be at the same frequency as the driving force but shifted in phase by ϕ_m . With these substitutions, Eqn. (6) becomes:

$$F_d \cos(\omega_d t) = m\omega_0^2 x(t) + [b - \mu_0 \omega_d \sin(\omega_d t + \phi_m)]x'(t) + [m + \mu_0 \cos(\omega_d t + \phi_m)]x''(t) \quad (7)$$

This non-linear differential equation has no analytic solutions and must be solved numerically. Fig. 2 shows the results of such numerical solutions of Eqn. (7), assuming that $F_d/m=0.01$, $b/m=0.01$, and $\mu_0/m=0.001$. The latter assumption represents only about 10% of the predicted 10 mg mass variation.




Figure 2. Response phase shifts of the system to variable mass. The central line is the system response with $\mu_0=0$. The other two lines represent $\mu_0=0.001$ with $\phi_m=-\pi/2$ (low), and $\phi_m=+\pi/2$ (high). The phase shifts shown are about ± 0.04 radians= ± 2.3 degrees.

We find that when the mass variation has a relative phase of $\pm\pi/2$ with respect to the driving force, it causes a positive or negative phase shift in the response motion by shifts, using the values listed above, of several degrees. Other phases near 0 or π can cause an increase or decrease in the amplitude of oscillation. The experiment we have constructed is designed in an attempt to observe these phase-shift effects.

Experimental Apparatus

Fig. 2 below shows a top view of the mechanical oscillator arrangement, which we call the “Mach Guitar”. The barium titanate capacitor test mass is suspended between pairs of tensioned wires, with the tension adjusted so that the resonant vibration frequency for vertical oscillations is about 1-2 kHz.




Figure 2 Top view of “Mach Guitar” arrangement. The capacitor is suspended between pairs of tensioned wires that provide the restoring force for the mechanical oscillator. Capacitor drive voltage is supplied through the wires.




Figure 3 Laser-optics table with oscillator removed, showing voice-coil drive (below) and laser position monitor (above).

Electrical connections for the capacitor drive voltage are supplied through the tensioned wire pairs. The capacitor and its support structure have a net mass of about 1 g.

The restoring force provided by the tensioned wire pairs is $F=-(8T/L)x$, where T is the tension in a given wire, L is the overall length, from bridge to bridge, of the system, and x is the vertical displacement of the capacitor. Therefore, neglecting the mass of the wire, the resonant frequency of the oscillator is $\omega_0=(8T/mL)^{1/2}$. If $m=1$ g, $L=0.5$ m and $\omega_0=2\pi \times 1000$ Hz, then the required tension is 553 lb. This tension can be reached with 13 gauge steel wire.

Fig. 3 shows a view of the laser-optics table (with oscillator removed) that is the foundation of the experiment. A pre-drilled aluminum laser-optics base plate supports the general-purpose aluminum beam structures, on which are mounted (below) the voice-coil drive (a modified audio speaker) for the mechanical oscillator, and (above) the laser position-measuring device shown in Fig. 4.

The Mach Guitar is mounted on the laser-optics base, which provides “bridges” to support for the wires

and their tensioning mechanism. Electrical connections to the capacitance are made through the support wires. Below the oscillator is an audio speaker, which drives the oscillator through a light spring. Above the oscillator is a commercial laser position detector, which measures the vertical position of the capacitor's upper surface by electronic triangulation. The laser position sensor is shown in Fig. 4



Fig. 4 Laser position measurement device.

The mass-varying object used in the measurements is a low-loss and low-mechanical-movement barium titanate capacitor with a capacitance of about 9 nF and a voltage rating of 3 kV. This oscillator mass is suspended between pairs of 13 gauge steel wires (0.25 m long on each side) that have been tensioned to about 500 lb to provide a system resonant frequency of about 1000 Hz.



Fig. 5 Trek Model PO923A HV Power Amplifier, used for driving the capacitor at 2 kV and 400 mA.

As previously mentioned, the most challenging problem presented by the present experiment is driving the capacitor to high voltages at audio frequencies. The reason is that all high-voltage amplifiers driving capacitive loads are severely limited by the charging current that they must deliver. We have selected a Trek Model PO923A High Voltage Power Amplifier, shown in Fig. 5, as the capacitor driver. It can drive at voltages up to 2 kV with a peak charging current of up to 400 mA.

The Mach's Principle test employs a Pentium-2 850 MHz computer system with a Windows 98 operating system for experiment control, using control software based on LabView. It consists of controls for the mechanical oscillator driver and the capacitance driver, a data collection system that records the drive signal and the position measurements, a data recording and retrieval system, and analysis software for processing the data and extracting the phase information of interest.




Figure 6. LabView control panel and display for system drivers.

The LabView control panel and display for the experiment is shown in Fig. 6 above. The system generates sine waves with adjustable phases and amplitudes at two frequencies, normally set to differ by a factor of two. The low frequency signal provides the input to the high voltage amplifier that drives the capacitor. The high frequency provides input to an audio amplifier connected to a voice coil that drives the mechanical oscillator.




Figure 7. LabView data acquisition and display.

Figure 7 shows the LabView panel for the data collection system. The system samples the mechanical drive voltage and the capacitor position measurement of the mechanical oscillator as separate data streams. These are sampled for real-time display and also recorded on the system hard disk. These data streams can be read back and re-analyzed. The data are analyzed by integration over a long time period to extract the relative phase of the drive and response signals for a given setting of the capacitor drive phase with respect to the mechanical driver.

The processed quantity that will be accumulated in the analysis is the cosine of the relative phase between the driving signal $D(t)$ and the mechanical response signal $R(t)$. There are a variety of ways of extracting this signal, but the one we will use initially is:

$$\text{Cos}(\phi) = \frac{\int_0^T [D(t) + R(t)]^2 dt - \int_0^T [D(t) - R(t)]^2 dt}{4 \sqrt{\int_0^T D(t)^2 dt \times \int_0^T R(t)^2 dt}} \quad (8)$$

Here T is an arbitrary integration time that increases as data is collected and the running integrals are accumulated. The values of $\text{Cos}(\phi)$, which is near 0 because ϕ is approximately $\pi/2$ on resonance, will be compared for the two most extreme settings of the phase of the capacitor drive, which should produce phase shifts like those shown in Fig. 2. We estimate that with a data collection cycle of a few hours, $\text{Cos}(\phi)$ can be determined to an accuracy of a few parts in 10^5 . This should enable us to determine the shift in ϕ to similar accuracy, providing a fairly stringent test of the presence or absence of the Woodward mass variation.

Experiment Status

The experiment is presently being reconfigured on the laser-optic table. The initial cantilever arrangement is being replaced with the new "Mach guitar" mechanical oscillator system described above.

The previous mechanical oscillator, which used a capacitor mass suspended at the free end of an aluminum cantilever, was tested and found to present three serious problems for the experiment: (1) its resonant frequency was fixed by the length and mass of the cantilever and was not easily adjustable, (2) The cantilever mass dominated that of the capacitor, greatly reducing the magnitude of the predicted effect and (3) it was not capable of achieving resonant frequencies above a few hundred Hz. Since the size of the predicted mass-variation effect increases as ω_0^2 , this was a serious limitation. However, initial experience with this cantilever oscillator provided valuable experience in operating and testing the position measuring device and the data collection system.

The new Mach Guitar oscillator provides significant improvements over the cantilever in reduced mass and increased operating frequencies, and it offers the additional advantage that it is easily tunable through simple adjustments of the wire tension.

We expect to begin data collection with the new configuration in the next few weeks.

Conclusion

The test of Mach's principle and the Woodward effect described above is not yet completed, but it shows promise of providing an independent test of the predictions that does not depend on the possibility of a reactionless force. The experiment in the present configuration is not as sensitive as the torsion-balance measurements recently reported by Woodward [7]. However, since it is not based on a reactionless force, it may not need that sensitivity.

If the Woodward Effect is observed, it will have important implications for general relativity and cosmology, for validating Mach's Principle, for control of inertia, and possibly for propulsion. If the Woodward Effect is not observed at the sensitivity limit of the experiment, this will also be worth knowing.

[6] James F. Woodward, "A Laboratory Test of Mach's Principle and Strong-Field Relativistic Gravity," *Foundations of Physics Letters* **9**, 247 – 293 (1996).

[7] James F. Woodward, "Mass Fluctuations, Stationary Forces, and Propellantless Propulsion," *Space Technology and Applications International Forum 2000* (American Institute of Physics/Springer Verlag, New York, 2000), pp. 1018 – 1025 (2000).

Acknowledgements

The authors are grateful to Marc G. Millis and the NASA Breakthrough Propulsion Program for providing encouragement and funding for the present experiment. We thank James F. Woodward for providing the very high quality barium titanate capacitors used in the experiment, along with much useful advice.

Bibliography

[1] D. Sciama, "On the Origin of Inertia," *Monthly Notices of the Royal Astronomical Society* **113**, 34–42 (1953).

[2] D. Sciama, "The Physical Structure of General Relativity," *Reviews of Modern Physics* **36**, 463–469 (1964).

[3] Woodward, J.F., "A New Experimental Approach to Mach's Principle and Relativistic Gravitation," *Foundations of Physics Letters* **3**, 497 – 506 (1990).

[4] James F. Woodward, "Measurements of a Machian Transient Mass Fluctuation," *Foundations of Physics Letters* **4**, 407–423 (1991).

[5] James F. Woodward , "A Stationary Apparent Weight Shift from a Transient Machian Mass Fluctuation," *Foundations of Physics Letters* **5**, 425–442 (1992).

SUPERLUMINAL BUT CAUSAL WAVE PROPAGATION

Mohammad Mojahedi, Kevin J. Malloy

Center for High Technology Materials, Department of Electrical and Computer Engineering,
University of New Mexico, 1313 Goddard SE, Albuquerque NM, 87106, (email:
mojahed@chtm.unm.edu).

Raymond Chiao

Department of Physics, University of California at Berkeley, Berkeley, CA 94720.

Abstract: A series of experiments in recent years have shown that under carefully designed circumstances the group velocity, or even more surprisingly the energy velocity can exceed the speed of light in vacuum or become negative (abnormal velocities). These abnormal results have led some researchers to question the validity of special relativity, or at least cast doubt on the relevance of these principles to the aforementioned experiments. In this work series of experiments with single electromagnetic pulses measured in both time and frequency domain are described. It is seen that while these experiments verify the aforementioned abnormal velocities, they are not in contradiction with the principles of special relativity (Einstein causality). In this regard, the important concept of “front” or “Sommerfeld forerunner” is reintroduced, and it is argued that the only physical velocity required to obey the Einstein causality is the “front velocity.”

I. Introduction

The fact that the group velocity of an electromagnetic wave packet (pulse) can exceed the speed of light in vacuum (become superluminal) has been demonstrated in many experiments using single photons^{1, 2}, at optical frequencies³, and using microwaves⁴⁻¹⁰. As a starting point, an interested reader may consult the review by Chiao and the references therein¹¹. Despite one’s initial impression, the

superluminal group or even energy velocities (defined as the ratio of the Poynting vector to the stored electromagnetic energy) are not at odds with the requirements of relativistic causality (Einstein causality), and indeed it can be shown that they must exist as the natural consequence of the Kramers-Kronig relations, which in themselves are a statement of the system linearity and causality¹²⁻¹⁵.

The point that in the regions of anomalous dispersion, group velocity can become superluminal was first considered by Sommerfeld and his student Brillouin¹⁶. In their studies, they examined a sinusoidally modulated step-function propagating through a medium with Lorentzian dispersion. They identified five different velocities: phase, group, energy, Sommerfeld forerunner (“front”) and Brillouin forerunner velocities[†]. However, with the passage of time, and for reasons unknown to the authors, while the first three velocity terms have received much attention in both undergraduate and graduate books, the latter two have not enjoyed the same status. This is even more surprising since, among the above velocities, it is only the

[†] To be more rigorous the term “front” refers to the onset of Sommerfeld forerunner propagation.

[‡] To be complete one has to add the term “signal velocity” defined as the velocity of the half maximum point to the list. However, by their own admission such a definition is arbitrary¹⁶ and as discussed in Ref. (4) can become superluminal.

velocity of the “front” that must satisfy the requirements of Einstein causality under all circumstances. In other words, it is rather a naïve understanding of Einstein causality to equate the group velocity with the velocity of information transfer under all circumstances, particularly when one is concerned with the propagation of “attenuated traveling waves[§].”

Our objective here is to discuss the phenomenon of superluminal and negative group and energy velocities which generically is referred to as the abnormal velocities. In Sec. II a time-domain experiment used to detect the superluminal group velocities in the case of a one dimensional photonic crystal (1DPC) is described. Section III discusses a frequency-domain experiment used to demonstrate the same superluminal phenomenon. The case of superluminal or negative group and energy velocities for an inverted medium (medium with gain) or in the case of medium with negative index of refraction is considered in Sec. IV. Section V is intended to put the reader’s mind at ease by providing some general arguments on why the abnormal velocities discussed in the previous sections are not in contradiction with the requirements of special relativity. Section VI is our condensed attempt is addressing the issue of superluminality in the limit of very weak light (very few photons). Our final remarks and a discussion for the general public can be found in Sec. VII.

II. Time-domain Experiment

Consider the problem of electromagnetic wave propagation through a periodic structure. Figure 1 shows an experimental setup used to detect the superluminal group velocity for a

microwave wave packet tunneling through a 1DPC. A backward wave oscillator (BWO) was used to generate the pulse, and a mode converter (MC) was used to convert the TM_{01} mode of the BWO to a TE_{11} . The pulse was then radiated via a conical horn




Fig. 1: Time-domain experimental setup

antenna (CHA). The frequency output of the source was tuned to the mid-gap frequency of the 1DPC at 9.68 GHz (FWHM of 100 MHz) and was detected by two HP 8470-B, Schottky diode detector (provided in pairs). The CHA radiation intensity was sampled along two distinct directions (paths), referred to as “side” and “center”. A series of microwave pulses were fired in order to measure and then remove the time difference between the two paths due to the differences in cable lengths, internal detection of the oscilloscopes (Tektronix SCD 5000) and other incompatibilities. This measured time difference was electronically compensated such that the peaks corresponding to the pulses traveling through the “center” path and “side” path in the absence of the 1DPC arrived at the same time. At this point, the 1DPC was inserted along the “center” path and series of single pulse were fired. Figure 2 shows the result. It is seen that the peak of the wave packet propagating along the “center” path and tunneling through the 1DPC arrives (440 ± 20) ps earlier than the accompanying pulse propagating through free-space along the “side” path. This

[§] We have used the term “attenuated traveling waves” in the same sense as in Ref. (17), although, sometimes the term evanescent is used to signify the same thing.

advancement in time for the tunneling pulse can easily be translated to a measure of the pulse group velocity, indicating that the tunneling wave packet propagated with a




Fig. 2: Superluminal propagation for the tunneling pulse

group velocity (2.38 ± 0.15) c .

Furthermore, the traditional view of pulse propagation through a region with high attenuation (regions of anomalous dispersion) held that the extreme




Fig. 3: A measure of the pulse broadening due to tunneling through the 1DPC. The two pulses have propagated along the same path (“center”); one in the free-space and the other through the 1DPC.

attenuation, coupled with the dispersion, would distort the signal to such an extent that the originally well defined wave packet and its peak would not be recognizable upon the emergence from such a medium^{17 16}. Figure 3 shows that in contrast to this common belief, the tunneling wave packet of Fig. (2) suffered minimal dispersion such

that the FWHM of the pulse after tunneling was only increased by 1.5%. In obtaining this figure the tunneling wave packet was manually moved to later times as to make the comparison between the two pulses easier. A full description of the above experiment can be found in Ref. (4).

III. Frequency-Domain Experiment

In the last section the feasibility of measuring superluminal group velocities directly in time-domain was discussed. This abnormal behavior can also be demonstrated in frequency-domain. Figure 4 shows the




Fig. 4: Frequency-domain experimental setup

free-space setup used to detect the superluminal group velocities in frequency-domain. The setup consists of two K-band standard horn antennas (SHA) configured a transmitter and receiver and connected to ports 1 and 2 of an HP 8722D vector network analyzer (VNA). The radiated quasi continuous waves are collimated using two microwave lenses and the setup is enclosed in a anechoic chamber to reduce stray signals.

The essence of the approach is to measure *accurately* and *reliably* the transmission phase associated with the 1DPC. Once this quantity is measured, the group delay (τ_g) and group velocity (V_g) can be calculated according to

$$\tau_g = -\partial \phi / \partial \omega, \quad (1)$$

$$\frac{v_g}{c} = \frac{L_{pc}}{c \tau_g} = \frac{-L_{pc}}{c (\partial \phi / \partial \omega)}, \quad (2)$$

where ϕ is the transmission phase, and L_{pc} is the physical length of the 1DPC.

Fortunately, recent advances in non-coaxial (free-space) calibration techniques for VNA such as the “Thru-Line-Reflect” (TRL),^{18, 19} make it possible to measured the transmission coefficient accurately and reliably. After calibrating the system (without the 1DPC), a reference plane of unit magnitude for transmission magnitude and zero phase for ϕ is established midway between the two SHAs. At this point, the 1DPC is inserted and the receiver horn is moved back exactly by a length equal to the thickness of the 1DPC (L_{pc}).

Figure 5 is the calculated (solid line)




Fig 5: The unwrapped transmission phase for the 1DPC with various number of Eccostock® dielectric slabs.

and measured (dotted line) unwrapped phase for a 1DPC with four, three, two and one dielectric slabs (the spacer is always air). The theoretical calculations are based on the diagonalization of one period matrix, and is presented in Ref. (5).

According to Eqs. (1) and (2), to ascertain the group delay and group velocity the data presented in Fig. 5 must be differentiated. However, differentiating a noisy data amplifies the noise and may lead to spurious effects. To avoid the arbitrariness associated with smoothing, a best nonlinear least square fit of the experimental phase data is obtained. The parameters used in fitting the experimental data all match the actual variables very well

and for the sake of brevity are not given here, but can be found in Ref. (5).

Figure 6 shows the result of the least square fit to the phase data of Fig. 5 together




Fig. 6: Measured and calculated group delay for the 1DPC. The parameters used to obtain the fitted curves (measurement) and the calculated curves (theory) are given in Ref (5).

with Eq. (1), in order to determine the group delay in a 1DPC with one, two, three, and four dielectric slabs. Consequently, the normalized group velocity given by Eq. (2)




Fig. 7. Normalized group velocity for the 1DPC. The dotted curves are the measured results obtained from the fitted curves in Fig. 6 and Eq. 2. The solid curves are theoretically calculated.

can be obtained from the Fig. 6 and is shown in Fig. 7. Along with the velocities derived from the fit (dotted curves), the theoretical group velocities calculated from the measured values of thicknesses and indices are also shown (solid curves). As Fig. 7 indicates, in the case of $N=4$ and $N=3$, a

maximum superluminal group velocities 3 and 2.1 times c is observed. The results depicted in this figure can be interpreted as the following. If one is to construct a pulse entirely composed of the frequency components for which the superluminal behavior is predicted, then the pulse is expected to propagate with group velocity exceeding c , similar to the situation discussed in the Sec. II.

IV. Abnormal Velocities in Inverted Medium and Medium with Negative Index

The circumstances under which the group or even energy velocity are abnormal are not limited to the evanescent wave propagation discussed so far. In this section three situations are described which exhibit the aforementioned abnormal behavior.

First, for an inverted medium (medium with gain) described by a Lorentz-Lorenz dispersion, the index of refraction is given by

$$n(\omega) = \left(1 - \frac{\omega_p^2}{\omega_0^2 - \omega^2 - i\gamma\omega}\right)^{1/2} \quad (3)$$

where ω_0 is the resonance frequency, γ is a small damping factor, and ω_p is the effective plasma frequency. Note that a negative sign precedes the second term under the square root due to the population inversion of the medium. A plot of the index of refraction for both the inverted and non-inverted medium is shown in Fig. 8. From the figure it is clear that for an inverted medium, in the limit of very low frequencies, the index is less than one implying that the phase velocity is superluminal. More importantly, at the low frequencies, the group velocity given by

$$\begin{aligned} V_g(0) &= \frac{c}{[n(\omega) + \omega dn/d\omega]_{\omega \rightarrow 0}} = \\ &= \frac{c}{n(0)} = V_p(0) > c \end{aligned} \quad (4)$$

is also superluminal. Under these circumstances the energy velocity (V_e), given by the ratio of Poynting vector (S), to the stored energy density (u), is also equal to the phase and group velocity and exceeds the speed of light in vacuum.

$$V_e = \frac{S}{u} = \frac{c}{n(0)} = V_p(0) > c \quad (5)$$

The equivalence of the above three




Fig. 8: The real part of the index of refraction for an inverted medium (solid curve) and non-inverted medium (dashed curve.)

velocities is merely a statement of the fact that in the limit of low frequencies the medium is transparent and dispersionless¹³.

Second, it is also possible to observe




Fig. 9: Two cells of equal length containing inverted medium and vacuum.

negative group velocities for electromagnetic (EM) pulses tuned slightly away from a gain line of an inverted medium.^{14, 20-22}. The physical meaning of a negative group velocity can be explained as the following. Consider two cells of physical length L containing an inverted medium and vacuum as shown in Fig. 9.

The time difference between two well behaved identical EM pulses propagating through the lower (inverted) and the upper cell (vacuum) is given by

$$\Delta t = \tau_g - t_{vac} = \frac{L}{V_g} - \frac{L}{c} = \frac{L}{c} (n_g - 1) \quad (6)$$

where n_g is the group index. From the above equation it is clear that if the group index is zero the time difference between the two pulses is given by the negative of L/c . In other words, when one of the EM pulses is at the exit face of the lower cell the other pulse is about to enter the upper vacuum cell. Stating this point differently, if one only consider a single cell containing the inverted medium, for a negative group index, the peak of the transmitted wave packet leaves the cell prior to the peak of the incoming wave packet entering the medium. It must be pointed out that as shown by Landauer²³⁻²⁵ it is naïve to regard the peak of the outgoing pulse as the causal response of the medium to the peak of the incident pulse. The theoretical prediction by one of us²² regarding the feasibility of detecting negative group velocity was recently verified in an experiment by Wang²⁶ in which the inverted medium was a cell containing Cesium vapor.

Finally, let us consider a situation for which the medium effective index is a negative value. A point worth emphasizing is the fact that for these media it is the effective index and not the actual material index which is negative. In other words, the wavelength of the incident wave is many times larger than the physical size of the components comprising the media, allowing one to characterize the over all response of the media in terms of an effective index.

The first theoretical work in this area was done by Veselago^{27, 28}, and the more recent interest in the subject was reignited by the work of Pendry^{29, 30} and Smith et. al.^{31, 32}, which demonstrated the possibility of manufacturing these media at microwave

frequencies. Figure 10, shows the dispersion relation for a negative index




Fig. 10: The dispersion curves for a medium with negative effective index

medium, borrowed from Ref. (32). From the figure it is clear that for a certain frequency range, the derivative of the curve depicted in Fig. 10, (i.e. the group velocity) is negative. Even more surprising is the fact that the energy velocity, given by Eq. (7), is also negative, since in these media both permittivity and permeability are negative parameters.

$$V_e \propto \frac{\vec{E} \times \vec{H}}{\epsilon |\vec{E}|^2 + \mu |\vec{H}|^2} \quad (7)$$

The presence of negative group and energy velocities for the above media can be understood in the following manner. The




Fig. 11: Transmission magnitude and phase for the LRC circuit shown in the inset.

negative index medium, considered by Smith et. al.³¹, is in essence a distributed

LRC transmission line that its response can be approximated by a lumped LRC circuit. The inset in Fig. 11 is a typical LRC circuit that exhibits negative group delay in the region of maximum attenuation^{**}. Once again, if one is to construct an EM pulse mostly composed of frequency components having negative group delays, it is expected that the group and energy velocities for this EM pulse to be negative. We currently are pursuing the detection of the aforementioned abnormal velocities in the negative index media. We end this section by pointing out that in addition to negative velocities, the negative index medium has many other interesting properties such as inverted Doppler shift, Cherenkov radiation, and Snell's law.

V. Superluminal Velocities and Einstein Causality

In so far we have discussed situations for which the phase, group, and energy velocities are abnormal (superluminal or negative). The reader may began to wonder whether or not these abnormal velocities are in contradiction with the requirements of relativistic causality. The short answer to this question is that under no circumstances the so called "front velocity" may exceed the speed of light in vacuum, and in fact under all circumstances the "front velocity" is exactly luminal. In other words, the requirement of Einstein causality that no "signal" (information) can be transmitted superluminally is satisfied in all cases, since the "front velocity" is always luminal. This means that the presence of the genuine information should not be associated with the pulse maximum, half maximum, or the envelope, but indeed is contained within the singularities (points of

^{**} In obtaining Fig. 11, in contrast to the curves depicted in Fig. 5, a time dependency of $e^{-i\omega t}$ in place of $e^{j\omega t}$ was used. In other word, the group delay has the opposite sign of that shown in Eq. (1).

non-analyticity) of the pulse. Because of the important role played by the "front" in satisfying the requirements of special relativity, let us briefly discuss some of the most general ideas associated with this concept. The interested reader may consult the Ref. (4) for more detailed analysis.

The essential point to remember is the fact that any *physically realizable* signal is restrictively time-limited. In other words, any electromagnetic signal created and later propagated through free-space or a 1DPC must be generated at a point in time and space. One can then always point to a time prior to which the signal did not exist. This point in time, or more precisely the transient "turn on times," are points of non-analyticity for which the amplitude of the pulse or its first or higher derivative are discontinuous.

The importance of these points of non-analyticity becomes clear when considering the following. While the future behavior of a truly analytical signal such as a Gaussian wave packet can be completely predicted by means such as a Taylor expansion (or a Laurent expansion for functions that are holomorphic in an annular region), the presence or arrival of the singularities do not yield themselves to such an extrapolation. Moreover, as discussed in the above, no *physically realizable* signal can be presented by an *entire function*^{††} hence, any communication of information must involve the transmission of the "front". To summarize, there is no more information in a pulse peak or envelope that is not already contained within the earliest parts of the signal.

The mathematical proof that no signal (information) may be detected sooner than $t_0 = x/c$ can be seen via contour integration of an expression such as Eq. (8). Equation (8) describes the field at the position x and time t for a wave packet

^{††} An *entire function* is the one that is analytical everywhere in the complex domain.

impinging at normal incident on a medium characterized by an index of refraction, n ,³³

$$u(x, t) = \int_{-\infty}^{+\infty} \frac{2}{1+n(\omega)} A(\omega) e^{ik(\omega)x-i\omega t} d\omega, \quad (8)$$

$$A(\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} u(x=0, t) e^{i\omega t} d\omega. \quad (9)$$

Transforming the integral in Eq. (8) into the complex domain and closing the contour over the upper-half-plan, along with requiring that the medium characterized by n to be causal, and that the incident wave packet has a “front,” are sufficient conditions to show that the value of the integral is identically zero for $t \leq t_0 = x/c$ or equally for velocities, $V = x/t > c$. The condition that the medium characterized by n is causal, means that for this medium the effect can not proceed the cause. Mathematically this is expressed as $G(\tau) = 0$ for $\tau < 0$, where $G(\tau)$ is the susceptibility kernel given by

$$G(\tau) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} [\epsilon(\omega)/\epsilon_0 - 1] e^{-i\omega\tau} d\omega. \quad (10)$$

For times immediately after t_0 , ($t \approx t_0$) the earliest part of the signal known as the Sommerfeld forerunner or precursor can be detected. The frequency of oscillation and the field amplitude for the Sommerfeld forerunner are discussed by Mojahedi et. al.^{4, 34}. To summarize those results, the frequency of oscillation is given by

$$\omega_s = \sqrt{G'(0)} / \sqrt{2 \left(\frac{t}{t_0} - 1 \right)}, \quad (11)$$

where $G'(0)$ is the time derivative of the susceptibility kernel³³ evaluated at $t = 0$. Furthermore, for the incident wave packet proportional to t^m (m is an integer) the Sommerfeld forerunner is described by a Bessel function of order m according to

$$u(x, t) \approx a \left(\frac{t-t_0}{\gamma} \right)^{m/2} J_m \left(2 \sqrt{\gamma(t-t_0)} \right); \quad (12)$$

$$\gamma = \frac{G'(0) t_0}{2}; \quad \text{for } t > t_0.$$

From the above discussion it is clear that, for a given medium if the quantity $G'(0)$ is known, the calculation of the Sommerfeld forerunner frequency of




Fig. 12: Effective index for a 1DPC. The structure parameters are: $d_i = 1.76$ cm, $d_j = 1.33$ cm, $n_i = 1$, $n_j = 3.40$

oscillation and functional form is relatively straight forward. In other words, if one is capable of calculating $\epsilon(\omega)/\epsilon_0 - 1 = n^2 - 1$ for a 1DPC, undersized waveguide, or any other photonic barrier used in the superluminal experiments, then one can perform the inverse Fourier transform and the differentiation operation to obtain $G'(0)$. For example, let us consider the case of 1DPC used in the experiments discussed in Sections. II and III.

At the normal incidence the dispersion relation (K vs. ω) can be obtained from

$$\cos(K\Lambda) = \cos\left(\frac{\omega n_i d_i}{c}\right) \cos\left(\frac{\omega n_j d_j}{c}\right) - \frac{1}{2} \left(R + R^{-1}\right) \sin\left(\frac{\omega n_i d_i}{c}\right) \sin\left(\frac{\omega n_j d_j}{c}\right), \quad (13)$$

where R is the ratio of the indices given by $R = n_i/n_j$, Λ is the one-period length $\Lambda = d_i + d_j$, and K is the Bloch wave vector. The above equation can be used to solve for the real and imaginary parts of the Bloch wave vector, and Eqs. (14)-(16), below, can in turn be used to transform the photonic crystal spatial dispersion [Eq. (13)] to a more manageable temporal dispersion

$$\text{Re}(n_e) = n'_e = \frac{c}{\omega} \text{Re}[K(\omega)], \quad (14)$$

$$\text{Im}(n_e) = n''_e = \frac{c}{\omega} \text{Im}[K(\omega)], \quad (15)$$

$$|n_e| = \left[(n'_e)^2 + (n''_e)^2 \right]^{1/2} = \frac{c}{\omega} |K(\omega)|. \quad (16)$$

The results are shown in Fig. (12) which displays our first attempt in obtaining an effective index for a 1DPC, with 1, 2, ..., and infinite number of dielectric slabs. The next step is to perform the Fourier transform indicated by Eq. (10), followed by the differential operation evaluated at time equal to zero. Having obtained the quantity $G'(0)$, the frequency of oscillation and the functional form of the Sommerfeld forerunner in a 1DPC can be arrived at with the help of Eqs. (11) and (12).

VI. Superluminal Propagation and Quantum Noise in the Limit of Very Weak Pulse.

The question of superluminality in the limit of very weak pulse (one or few photons) was considered in a recent work ³⁵. For the sake of brevity, we refer the interested reader to Ref. (36) for a complete and detailed analysis of the situation. Here, we suffice to mention that according to Aharonov et. al. ³⁶ in the limit of few

photons, signal must be exponentially large in order to distinguish it from the quantum noise. In other words, the signal-to-noise ratio becomes vanishingly small. In Ref. (36) this assertion is investigated and it is seen that if the condition stated by Aharonov et. al. is replaced by a weaker condition, the signal-to-noise ratio can exceed unity even for one photon pulse. It is worth mentioning that the original experiment by Chiao and Steinberg, ² although involved the detection of single photon, but the results were interpreted in terms of statistics of many photons.

VII. Concluding Remarks: A Discussion for General Public

A simple yet interesting description of superluminal propagation can be found at web link

<http://www.abqjournal.com/scitech/180964scitech11-19-00.htm>.

This article written by John Fleck, the science writer for Albuquerque Journal, tries to explain our newly published paper in *Physical Review E* to the general public. To use John's analogy consider two dragsters competing against each other, driving the same exact cars and traveling the same exact distances. However, whereas one of the dragsters travels through air (vacuum if you like) with the maximum allowable speed, the other driver travels through a series of barriers normally thought to slow his car. The question is then the following: What does the referee at the end line observe? The answer depends on the referee detection equipment. If the referee is well equipped with the most sensitive and expensive detection systems he or she will observe that the front bumpers of the two cars arrive at the finishing line at exactly the same instance. The referee will also observe that the bulk (the main body, the cockpit and the driver) of the dragster's car who tunneled through the barriers reaches the end line

sooner than his challenger. Interestingly, if the race is decided by arrival of the cars main body or if the referee is not equipped with the most sensitive detection apparatus, he or she will invariably call the race for the tunneling dragster.

Bibliography

- 1 A. M. Steinberg and R. Y. Chiao, Physical Review a **51**, 3525-3528 (1995).
- 2 A. M. Steinberg, P. G. Kwiat, and R. Y. Chiao, Physical Review Letters **71**, 708-711 (1993).
- 3 C. Spielmann, R. Szipocs, A. Stingl, *et al.*, Physical Review Letters **73**, 2308-2311 (1994).
- 4 M. Mojahedi, E. Schamiloglu, F. Hegeler, *et al.*, Physical Review E **62**, 5758-5766 (2000).
- 5 M. Mojahedi, E. Schamiloglu, K. Agi, *et al.*, Ieee Journal of Quantum Electronics **36**, 418-424 (2000).
- 6 A. Ranfagni, P. Fabeni, G. P. Pazzi, *et al.*, Physical Review E **48**, 1453-1460 (1993).
- 7 A. Ranfagni, D. Mugnai, P. Fabeni, *et al.*, Applied Physics Letters **58**, 774-776 (1991).
- 8 D. Mugnai, A. Ranfagni, and L. Ronchi, Physics Letters a **247**, 281-286 (1998).
- 9 A. Enders and G. Nimtz, Physical Review B **47**, 9605-9609 (1993).
- 10 A. Enders and G. Nimtz, Journal De Physique I **2**, 1693-1698 (1992).
- 11 R. Y. Chiao and A. M. Steinberg, Progress in Optics **37**, 345-405 (1997).
- 12 E. L. Bolda, R. Y. Chiao, and J. C. Garrison, Physical Review A **48**, 3890-3894 (1993).
- 13 R. Y. Chiao, Physical Review a **48**, R34-R37 (1993).
- 14 E. L. Bolda, J. C. Garrison, and R. Y. Chiao, Physical Review a **49**, 2938-2947 (1994).
- 15 R. Y. Chiao and J. Boyce, Physical Review Letters **73**, 3383-3386 (1994).
- 16 L. Brillouin, *Wave propagation and group velocity* (Academic Press, New York,, 1960).
- 17 L. D. Landau, E. M. Lifshitz, and L. P. Pitaevski, *Electrodynamics of continuous media* (Pergamon, Oxford [Oxfordshire] ; New York, 1984).
- 18 (Hewlett Packard, 1997), p. 1-60.
- 19 (Hewlett Packard, 1992), p. 1-24.
- 20 R. Y. Chiao, A. E. Kozhekin, and G. Kurizki, Physical Review Letters **77**, 1254-1257 (1996).
- 21 A. M. Steinberg and R. Y. Chiao, Physical Review a **49**, 2071-2075 (1994).
- 22 R. Y. Chiao, *Amazing light : a volume dedicated to Charles Hard Townes on his 80th birthday* (Springer, New York, 1996).
- 23 T. Martin and R. Landauer, Physical Review a **45**, 2611-2617 (1992).
- 24 M. Buttiker and R. Landauer, Physica Scripta **32**, 429-434 (1985).
- 25 R. Landauer, Nature **365**, 692-693 (1993).
- 26 L. J. Wang, A. Kuzmich, and A. Dogariu, Nature ; **406**, 277-279 (2000).
- 27 V. G. Veselago, Soviet Physics-Solid State **8**, 2854-2856 (1967).
- 28 V. G. Veselago, Soviet Physics USPEKHI **10**, 509-514 (1968).
- 29 J. B. Pendry, A. J. Holden, D. J. Robbins, *et al.*, Ieee Transactions On Microwave Theory and Techniques **47**, 2075-2084 (1999).
- 30 J. B. Pendry, Physical Review Letters **85**, 3966-3969 (2000).

- 31 D. R. Smith, W. J. Padilla, D. C. Vier, *et al.*, Physical Review Letters **84**, 4184-4187 (2000).
- 32 R. A. Shelby, D. R. Smith, and S. Schultz, Science **292**, 77-79 (2001).
- 33 J. D. Jackson, *Classical electrodynamics* (Wiley, New York, 1998).
- 34 M. Mojahedi, in *EECE* (University of New Mexico, Albuquerque, 1999).
- 35 B. Segev, P. W. Milonni, J. F. Babb, *et al.*, Physical Review a **62**, 022114-1-15 (2000).
- 36 Y. Aharonov, B. Reznik, and A. Stern, Physical Review Letters **81**, 2190-2193 (1998).



A01-34135

**AIAA 2001-3360
Geometrodynamics, Inertia and the
Quantum Vacuum**

Bernard Haisch
California Institute for Physics and Astrophysics
366 Cambridge Ave., Palo Alto, CA 94306

Alfonso Rueda
Department of Electrical Engineering, ECS Building
California State University, 1250 Bellflower Blvd.
Long Beach, California 90840

**37th AIAA/ASME/SAE/ASEE Joint
Propulsion Conference and Exhibit**
8-11 July 2001
Salt Lake City, Utah

For permission to copy or to republish, contact the copyright owner named on the first page.

For AIAA-held copyright, write to AIAA Permissions Department,
1801 Alexander Bell Drive, Suite 500, Reston, VA, 20191-4344.

Geometrodynamics, Inertia and the Quantum Vacuum

Bernard Haisch

California Institute for Physics and Astrophysics, 366 Cambridge Ave., Palo Alto, CA 94306
<www.calphysics.org>
Associate Fellow, AIAA — haisch@calphysics.org

Alfonso Rueda

Department of Electrical Engineering, ECS Building
California State University, 1250 Bellflower Blvd., Long Beach, California 90840
arueda@csulb.edu

(Presented at the AIAA/ASME/SAE/ASEE Joint Propulsion Conference, Salt Lake City, July 8–11, 2001)

ABSTRACT

Why does \mathbf{F} equal $m\mathbf{a}$ in Newton's equation of motion? How does a gravitational field produce a force? Why are inertial mass and gravitational mass the same? It appears that all three of these seemingly axiomatic foundational questions have an answer involving an identical physical process: interaction between the electromagnetic quantum vacuum and the fundamental charged particles (quarks and electrons) constituting matter. All three of these effects and equalities can be traced back to the appearance of a specific asymmetry in the otherwise uniform and isotropic electromagnetic quantum vacuum. This asymmetry gives rise to a non-zero Poynting vector from the perspective of an accelerating object. We call the resulting energy-momentum flux the *Rindler flux*. The key insight is that the asymmetry in an accelerating reference frame in flat spacetime is identical to that in a stationary reference frame (one that is not falling) in curved spacetime. Therefore the same Rindler flux that creates inertial reaction forces also creates weight. All of this is consistent with the conceptualization and formalism of general relativity. What this view adds to physics is insight into a specific physical process creating identical inertial and gravitational forces from which springs the weak principle of equivalence. What this view hints at in terms of advanced propulsion technology is the possibility that by locally modifying either the electromagnetic quantum vacuum and/or its interaction with matter, inertial and gravitational forces could be modified.

INTRODUCTION

Why does \mathbf{F} equal $m\mathbf{a}$ in Newton's equation of motion, $\mathbf{F} = m\mathbf{a}$?

How does a gravitational field produce a force?

Why are inertial mass and gravitational mass the same?

These are questions that are usually thought to be more appropriate for philosophers than for physicists, since the apparent facts of nature addressed in these questions are generally regarded as axiomatic.[1] If one assumes that one plus one equals two, plus a limited number of additional axioms, one can develop a self-consistent system of mathematics, but one has to start with such assumptions. In the realm of geometry, the discovery of non-Euclidean geometry in the 19th century taught us that other, non-intuitive assumptions are possible, and indeed, Riemannian geometry became the basis for physics in Einstein's general relativity (GR). Alternate foundational assumptions can lead to new insights.

Since 1991 we have been engaged in investigations involving the nature of inertia based on the foundational assumption that the electromagnetic quantum vacuum, also called the zero-point field or zero-point

fluctuations, is a real and underlying universal sea of energy capable of interacting with matter, and that this interaction may be described using the techniques of stochastic electrodynamics (SED). [2] Following a successful multiyear NASA-funded investigation at Lockheed Martin and California State University at Long Beach, the privately-funded California Institute for Physics and Astrophysics (CIPA) was established in 1999 specifically to study the electromagnetic quantum vacuum and its effects. A group of five postdoctoral fellows with expertise in quantum electrodynamics, superstring and M-brane theory, general relativity, plasma physics and Casimir effects are engaged in theoretical efforts to come to a deeper understanding of foundational questions in physics and to examine the possible nature and degree of electromagnetic quantum vacuum-matter interactions both within and beyond the SED approximations. CIPA has also funded projects by experts at several universities directed toward the general goal of explaining the origin of inertia and certain aspects of gravitation and other relevant physical and astrophysical effects as due to the electromagnetic quantum vacuum.

It appears that all three of the seemingly axiomatic foundational questions posed above have an answer involving an identical physical process: interaction between the electromagnetic quantum vacuum and the fundamental charged particles (quarks and electrons) constituting matter. All three of these effects and equalities can be traced back to the appearance of a specific asymmetry in the otherwise uniform and isotropic electromagnetic quantum vacuum. The key insight is that the asymmetry in an accelerating reference frame in flat spacetime is identical to that in a stationary reference frame in curved spacetime.

It was shown by Unruh [3] and by Davies [4] that a uniformly-accelerating detector will experience a Planckian-like heat bath whose apparent "temperature" is a result of quantum vacuum radiation. A tiny fraction of the (enormous) electromagnetic quantum vacuum energy can emerge as real radiation under the appropriate conditions. The theoretical prediction of Unruh-Davies radiation is now generally accepted and SLAC physicist P. Chen has recently proposed an experiment to detect and measure it. [5] Rueda and Haisch (hereafter RH) [6] analyzed a related process and found that as perceived by an accelerating object, an energy and momentum flux of radiation emerges from the electromagnetic quantum vacuum and that the strength of this momentum flux is such that the radiation pressure force on the accelerating object is proportional to acceleration. Owing to its origin in an accelerating reference frame customarily called the *Rindler frame*, and to the relation of this flux to the existence of a *Rindler event horizon*, we call this flux of energy and momentum that emerges out of the electromagnetic quantum vacuum upon acceleration the *Rindler flux*.

If the Rindler flux is allowed to electromagnetically interact with matter, mainly but perhaps not exclusively at the level of quarks and electrons, a reaction force is produced that can be interpreted as the origin of Newton's $F = ma$. In this view, which we call the *quantum vacuum inertia hypothesis*, matter resists acceleration not because of some innate property of inertia, but rather because the quantum vacuum fields provide an acceleration-dependent drag force. In future attempts we and coworkers intend to examine the possible contributions of other components of the quantum vacuum besides the electromagnetic. (This is relevant to the issue of possible neutrino mass, which could not be due to the electromagnetic quantum vacuum, but might possibly be due to the vacuum fields of the weak interaction.)

GR declares that gravity can be interpreted as spacetime curvature. Wheeler coined the term geometrodynamics to describe this: the dynamics of objects subject to gravity is determined by the geometry of four-dimensional spacetime. What geometrodynamics actually specifies is the family of geodesics — the shortest four-dimensional distances between two points in spacetime — in the presence of a gravitating body. Freely-falling objects and light rays follow geodesics. However when an object is prevented from following a geodetic trajectory, a force is experienced: the well-known force called weight. Where does this force come from? Or put another way, how does a gravitational field exert a force on a non freely-falling, fixed, object, such as an observer standing on a scale on the Earth's surface? *This proves to be the identical physical process as described in the quantum vacuum inertia hypothesis, due to a non-zero Rindler flux.*

In the SED approximation, the electromagnetic quantum vacuum is represented as propagating electromagnetic waves. [7,8] These should follow geodesics. It can be shown that propagation along curved geodesics creates the identical electromagnetic Rindler flux with respect to a stationary fixed object as is the case for an

accelerating object.[9] This is perfectly consistent with Einstein's fundamental assumption of the equivalence of gravitation and acceleration. An object fixed above a gravitating body will perceive the electromagnetic quantum vacuum to be accelerating past it, which is of course the same as the perception of the object when it is doing the accelerating through the quantum vacuum. Thus in the case of gravity, it would be the electromagnetic Rindler flux acting upon a fixed object that creates the force known as weight, thereby answering the second question. The answer to the third question then immediately follows. Since the same flux would be seen by either a fixed object in a gravitational field or an accelerating object in free space, the force that is felt would be the same, hence the parameters we traditionally call inertial and gravitational mass must be the same. This would explain the physical origin of the weak principle of equivalence.

All of this is consistent with the mathematics of GR. What this view adds to physics is insight into a specific physical process creating identical inertial and gravitational forces. What this view hints at in terms of advanced propulsion technology is the possibility that by locally modifying either the quantum vacuum fields and/or their interaction with matter, inertial and gravitational forces could be modified and possibly one day freely controlled.

THE ELECTROMAGNETIC QUANTUM VACUUM

The quantization of the electromagnetic field in terms of quantum-mechanical operators may be found in various standard textbooks, e.g. Loudon [10]: "The electromagnetic field is now quantized by the association of a quantum-mechanical harmonic oscillator with each mode \mathbf{k} of the radiation field." This can be understood as follows: Application of the Heisenberg uncertainty relation to a harmonic oscillator requires that its ground state have a non-zero energy of $h\nu/2$. This reflects the fact that quantum mechanically a particle cannot simultaneously be exactly at the bottom of its potential well and have exactly zero momentum. There exists the same $h\nu/2$ zero-point energy expression for each mode of the electromagnetic field as for a mechanical oscillator. (Formally, mode decomposition yields that each mode can be mathematically made into a harmonic oscillator in the sense that the same differential equation is obeyed as for a mechanical oscillator.)

Summing up the energy over the modes for all frequencies, directions, and polarization states, one arrives at a zero-point energy density for the electromagnetic fluctuations, and this is the origin of the electromagnetic quantum vacuum. An energy of $h\nu/2$ per mode of the field characterizes the fluctuations of the quantized radiation field in quantum field theory. In the semi-classical representation of SED the quantum vacuum is represented by propagating electromagnetic plane waves, \mathbf{E}^{zp} and \mathbf{B}^{zp} , of random phase having this average energy, $h\nu/2$, in each mode.

The volumetric density of modes between frequencies ν and $\nu + d\nu$ is given by the density of states function $N_\nu d\nu = (8\pi\nu^2/c^3)d\nu$. Each state has a minimum $h\nu/2$ of energy, and using this density of states function and this minimum energy — that we call the zero-point energy — per state one gets the spectral energy density of the electromagnetic quantum vacuum:

$$\rho(\nu)d\nu = \frac{8\pi\nu^2}{c^3} \frac{h\nu}{2} d\nu. \quad (1)$$

Writing this zero-point radiation together with ordinary blackbody radiation, the energy density is:

$$\rho(\nu, T)d\nu = \frac{8\pi\nu^2}{c^3} \left(\frac{h\nu}{e^{h\nu/kT} - 1} + \frac{h\nu}{2} \right) d\nu. \quad (2)$$

The first term (outside the parentheses) represents the mode density, and the terms inside the parentheses are the average energy per mode of thermal radiation at temperature T plus the zero-point energy, $h\nu/2$, which has no temperature dependence. Take away all thermal energy by formally letting T go to zero, and one is still left with the zero-point term. The laws of quantum mechanics as applied to electromagnetic

radiation force the existence of a background sea of electromagnetic zero-point energy that is traditionally called the electromagnetic quantum vacuum.

The spectral energy density of eqn. (1) was thought to be no more than a spatially uniform constant offset that cancels out when considering energy fluxes, but it was discovered in the mid-1970's that the quantum vacuum acquires special characteristics when viewed from an accelerating frame. Just as there is an event horizon for a black hole, there is an analogous event horizon for an accelerating reference frame. Similar to radiation from evaporating black holes proposed by Hawking [11], Unruh [3] and Davies [4] determined that a Planck-like radiation component will arise out of the quantum vacuum in a uniformly-accelerating coordinate system having constant proper acceleration \mathbf{a} (where $|\mathbf{a}| = a$) with what amounts to an effective "temperature"

$$T_a = \frac{\hbar a}{2\pi c k}. \quad (3)$$

This "temperature" characterizing Unruh-Davies radiation does not originate in emission from particles undergoing thermal motions.^a As discussed by Davies, Dray and Manogue [12]:

One of the most curious properties to be discussed in recent years is the prediction that an observer who accelerates in the conventional quantum vacuum of Minkowski space will perceive a bath of radiation, while an inertial observer of course perceives nothing. In the case of linear acceleration, for which there exists an extensive literature, the response of a model particle detector mimics the effect of its being immersed in a bath of thermal radiation (the so-called Unruh effect).

This "heat bath" is a quantum phenomenon. The "temperature" is negligible for most accelerations. Only in the extremely large gravitational fields of black holes or in high-energy particle collisions can this become significant. Recently, P. Chen at the Stanford Linear Accelerator Center has proposed using an ultra high intensity laser to accelerate electrons violently enough to directly detect Unruh-Davies radiation.[5]

Unruh and Davies treated the electromagnetic quantum vacuum as a scalar field. If a true vectorial approach is considered there appear additional terms beyond the quasi-thermal Unruh-Davies component. For the case of no true external thermal radiation ($T = 0$) but including the acceleration effect (T_a), eqn. (1) becomes

$$\rho(\nu, T_a) d\nu = \frac{8\pi\nu^2}{c^3} \left[1 + \left(\frac{a}{2\pi c \nu} \right)^2 \right] \left[\frac{h\nu}{2} + \frac{h\nu}{e^{h\nu/kT_a} - 1} \right] d\nu, \quad (4)$$

where the acceleration-dependent pseudo-Planckian Unruh-Davies component is placed after the $h\nu/2$ term to indicate that except for extreme accelerations (e.g. particle collisions at high energies) this term is negligibly small. While these additional acceleration-dependent terms do not show any spatial asymmetry in the expression for the spectral energy density, certain asymmetries do appear when the momentum flux of this radiation is calculated, resulting in a non-zero Rindler flux.[6] This asymmetry appears to be the process underlying inertial and gravitational forces.

ORIGIN OF THE INERTIAL REACTION FORCE

Newton's third law states that if an agent applies a force to a point on an object, at that point there arises an equal and opposite reaction force back upon the agent. In the case of a fixed object the equal and opposite reaction force can be traced to interatomic forces in the neighborhood of the point of contact which act to resist compression, and these in turn can be traced to electromagnetic interactions involving orbital electrons of adjacent atoms or molecules, etc.

^a There is likely to be a deep connection between the fact that the spectrum that arises in this fashion due to acceleration and the ordinary blackbody spectrum have identical form.

Now a similar experience of an equal and opposite reaction force arises when a non-fixed object is forced to accelerate. Why does acceleration create such a reaction force? We suggest that this equal and opposite reaction force also has an underlying cause which is at least partially electromagnetic, and specifically may be due to the scattering of electromagnetic quantum vacuum radiation. RH demonstrated that from the point of view of the pushing agent there exists a net flux (Poynting vector) of quantum vacuum radiation transiting the accelerating object in a direction opposite to the acceleration: the Rindler flux.[6] Interaction of this flux with the quarks and electrons constituting a material object would create a back reaction force that can be interpreted as inertia. One simply needs to assume that there is some dimensionless efficiency factor, $\eta(\omega)$, that in the case of particles corresponds to whatever the interaction process is (e.g. dipole scattering). In the case of elementary particles we suspect that $\eta(\omega)$ contains one or more resonances — and in the Appendix discuss why these resonances likely involve Compton frequencies of relevant particles forming a composite particle or object — but this is not a necessary assumption.

The RH approach relies on making transformations of the \mathbf{E}^{zp} and \mathbf{B}^{zp} from a stationary to a uniformly-accelerating coordinate system (see, for example, §11.10 of Jackson for the relevant transformations [13]). In a stationary or uniformly-moving frame the \mathbf{E}^{zp} and \mathbf{B}^{zp} constitute an isotropic radiation pattern. In a uniformly-accelerating frame the radiation pattern acquires asymmetries. There appears a non-zero Poynting vector in any accelerating frame, and therefore a non-zero Rindler flux which carries a net flux of electromagnetic momentum. The scattering of this momentum flux generates a reaction force, \mathbf{F}_r , proportional to the acceleration. RH found an invariant scalar with the dimension of mass quantifying the inertial resistance force of opposition per unit of acceleration resulting from this process. We interpret this scalar as the inertial mass,

$$m_i = \frac{V_0}{c^2} \int \eta(\nu) \rho_{zp}(\nu) d\nu, \quad (5)$$

where ρ_{zp} is the well known spectral energy density of the electromagnetic quantum vacuum of eqn. (1). In other words, the amount of electromagnetic zero point energy instantaneously transiting through an object of volume V_0 and interacting with the quarks, electrons and all charges in that object is what constitutes the inertial mass of that object in this view. It is change in the momentum of the radiation field that creates the resistance to acceleration usually attributed to the inertia of an object.

Indeed, not only does the ordinary form of Newton's second law, $\mathbf{F} = m_i \mathbf{a}$, emerge from this analysis, but one can also obtain the relativistic form of the second law: [6]

$$\mathcal{F} = \frac{d\mathcal{P}}{d\tau} = \frac{d}{d\tau} (\gamma_\tau m_i c, \mathbf{p}). \quad (6)$$

The origin of inertia, in this picture, becomes remarkably intuitive. Any material object resists acceleration because the acceleration produces a perceived flux of radiation in the opposite direction that scatters within the object and thereby pushes against the accelerating agent. Inertia in the present model appears as a kind of acceleration-dependent electromagnetic quantum vacuum drag force acting upon electromagnetically-interacting elementary particles. The relativistic law for "mass" transformation involving the Lorentz factor γ — that is, the formula describing how the *inertia* of a body has been calculated to change according to an observer's relative motion — is automatically satisfied in this view, because the correct relativistic form of the reaction force is derived, as shown in eqn. (6).

ORIGIN OF WEIGHT AND THE WEAK EQUIVALENCE PRINCIPLE

Einstein introduced the *local Lorentz invariance* (LLI) principle in order to pass from special relativity to GR. It is possible to use this principle immediately to extend the results of the quantum vacuum inertia hypothesis to gravitation (details discussed in two forthcoming papers [9]).

The idea behind the LLI principle is embodied in the Einstein elevator thought experiment. He proposed that a freely-falling elevator in a gravitational field is equivalent to one that is not accelerating and is far

from any gravitating body. Physics experiments would yield the same results in either elevator, and therefore a freely-falling coordinate frame in a gravitational field is the same as an inertial Lorentz frame. (This is rigorously only true for a “small elevator” since a gravitational field around a planet, say, must be radial, hence there are inevitably tidal forces which would not be the case for an ideal acceleration.) The device Einstein used to develop general relativity was to invoke an infinite set of such freely falling frames. In each such frame, the laws of physics are those of special relativity. The additional features of general relativity emerge by comparing the properties of measurements made in freely-falling Lorentz frames “dropped” one after the other.

This approach of Einstein is both elegant and powerful. The LLI principle immediately tells us that an object accelerating through the electromagnetic quantum vacuum is equivalent to an object held fixed in a gravitational field while the electromagnetic quantum vacuum is effectively accelerating (falling) past it. The prediction of GR that light rays deviate from straight-line propagation in the presence of a gravitating body — which Eddington measured in 1919 thereby validating GR — translates into acceleration (falling) of the electromagnetic quantum vacuum. An object accelerating through the electromagnetic quantum vacuum experiences a Rindler flux which causes the inertia reaction force. A fixed object past which the electromagnetic quantum vacuum is accelerating, following the laws of GR, experiences the same Rindler flux and the resulting force is what we call weight. That is why $m_g = m_i$ and is the basis of the weak equivalence principle.

CONCLUSIONS

Geometrodynamics is an elegant theoretical structure, but there is a very fundamental physics question that geometrodynamics has never satisfactorily addressed. If an object is forced to deviate from its natural geodesic motion, a reaction force arises, i.e. the weight of an object. Where does the reaction force that is weight come from? That same force would also be the enforcer of geodesic motion for freely falling objects. GR specifies the metric of spacetime from which geodesics can be calculated, but is there a physical mechanism to keep freely-falling objects from straying from their proper geodesics? Geometrodynamics does not provide a physical mechanism for this. It can only claim that deviations of an object from its proper geodesic motion results in an inertial reaction force. This is true but uninformative. The quantum vacuum inertia hypothesis provides a physical process generating inertia and weight.

Quantum physics predicts the existence of an underlying sea of zero-point energy at every point in the universe. This is different from the cosmic microwave background and is also referred to as the electromagnetic quantum vacuum since it is the lowest state of otherwise empty space. This sea of energy fills all of space and is absolutely the same everywhere as perceived from a constant velocity reference frame. But viewed from an accelerating reference frame, the radiation pattern of the energy becomes minutely distorted: a tiny directional flow is experienced by an accelerating object or observer, the Rindler flux. Importantly, the force resulting from that energy-momentum flow turns out to be proportional to the acceleration. When this energy-momentum flow — that arises automatically when any object accelerates — interacts with the fundamental particles constituting matter (quarks and electrons) a force arises in the direction opposite to the acceleration. This process can be interpreted as the origin of inertia, i.e. as the basis of Newton’s second law of mechanics: $\mathbf{F} = m\mathbf{a}$ (and its relativistic extension).[6]

It has now been discovered that exactly the same distortion of the radiation pattern occurs in geometrodynamics when the metric is non-Minkowskian.[9] The curved spacetime geodesics of geometrodynamics affect the zero-point energy in the same way as light rays (because the zero-point energy is also a mode of electromagnetic radiation). The gravitational force causing weight and the reaction force causing inertia originate in an identical interaction with a distortion in the radiation of the zero-point energy field. Both are a kind of radiation pressure originating in the electromagnetic quantum vacuum. The underlying distortion of the radiation pattern is due to an event horizon-like effect and is related to Unruh-Davies radiation and Hawking radiation.

What the quantum vacuum inertia hypothesis accomplishes is to identify the physical process which is the enforcer of geometrodynamics or general relativity. The quantum vacuum inertia hypothesis appears

to provide a link between light propagation along geodesics and mechanics of material objects. Moreover, since the distortion of the zero-point energy radiation pattern is the same whether due to acceleration or being held stationary in a gravitational field, this explains a centuries old puzzle: why inertial mass and gravitational mass are the same: both are due to the same non-zero Rindler flux. This gives us a deeper insight into Einstein's principle of equivalence.

APPENDIX: INERTIA AND THE DE BROGLIE WAVELENGTH

Four-momentum is defined as

$$\mathbf{P} = \left(\frac{E}{c}, \mathbf{p} \right) = (\gamma m_0 c, \mathbf{p}) = (\gamma m_0 c, \gamma m_0 \mathbf{v}), \quad (A1)$$

where $|\mathbf{P}| = m_0 c$ and $E = \gamma m_0 c^2$. The Einstein-de Broglie relation defines the Compton frequency $\hbar\nu_C = m_0 c^2$ for an object of rest mass m_0 , and if we make the de Broglie assumption that the momentum-wave number relation for light also characterizes matter then $\mathbf{p} = \hbar\mathbf{k}_B$ where $\mathbf{k}_B = 2\pi(\lambda_{B,1}^{-1}, \lambda_{B,2}^{-1}, \lambda_{B,3}^{-1})$. We thus write

$$\frac{\mathbf{P}}{\hbar} = \left(\frac{2\pi\gamma\nu_C}{c}, \mathbf{k}_B \right) = 2\pi \left(\frac{\gamma}{\lambda_C}, \frac{1}{\lambda_{B,1}}, \frac{1}{\lambda_{B,2}}, \frac{1}{\lambda_{B,3}} \right) \quad (A2)$$

and from this obtain the relationship

$$\lambda_B = \frac{c}{\gamma v} \lambda_C \quad (A3)$$

between the Compton wavelength, λ_C , and the de Broglie wavelength, λ_B . For a stationary object λ_B is infinite, and the de Broglie wavelength decreases in inverse proportion to the momentum.

Eqn. (5) is very suggestive that quantum vacuum-elementary particle interaction involves a resonance at the Compton frequency. De Broglie proposed that an elementary particle is associated with a localized wave whose frequency is the Compton frequency. As summarized by Hunter [14]: "...what we regard as the (inertial) mass of the particle is, according to de Broglie's proposal, simply the vibrational energy (divided by c^2) of a localized oscillating field (most likely the electromagnetic field). From this standpoint inertial mass is not an elementary property of a particle, but rather a property derived from the localized oscillation of the (electromagnetic) field. De Broglie described this equivalence between mass and the energy of oscillational motion... as '*une grande loi de la Nature*' (a great law of nature)."

This perspective is consistent with the proposition that inertial mass, m_i , may be a coupling parameter between electromagnetically interacting particles and the quantum vacuum. Although De Broglie assumed that his wave at the Compton frequency originates in the particle itself (due to some intrinsic oscillation or circulation of charge perhaps) there is an alternative interpretation discussed in some detail by de la Peña and Cetto that a particle "is tuned to a wave originating in the high-frequency modes of the zero-point background field." [8] The de Broglie oscillation would thus be due to a resonant interaction with the quantum vacuum, presumably the same resonance that is responsible for creating a contribution to inertial mass as in eqn. (5). In other words, the electromagnetic quantum vacuum would be driving this ν_C oscillation.

We therefore suggest that an elementary charge driven to oscillate at the Compton frequency, ν_C , by the quantum vacuum may be the physical basis of the $\eta(\nu)$ scattering parameter in eqn. (5). For the case of the electron, this would imply that $\eta(\nu)$ is a sharply-peaked resonance at the frequency, expressed in terms of energy, $\hbar\nu_C = 512$ keV. The inertial mass of the electron would physically be the reaction force due to resonance scattering of the electromagnetic quantum vacuum radiation, the Rindler flux, at that frequency.

This leads to a surprising corollary. It has been shown that as viewed from a laboratory frame, a standing wave at the Compton frequency in the electron frame transforms into a traveling wave having the de Broglie wavelength for a moving electron.[8,14,15,16] The wave nature of the moving electron (as measured in the Davisson-Germer experiment, for example) would be basically due to Doppler shifts associated with its Einstein-de Broglie resonance at the Compton frequency. A simplified heuristic model shows this, and

a detailed treatment showing the same result may be found in de la Peña and Cetto [8]. Represent a quantum vacuum-like driving force field as two waves having the Compton frequency $\omega_C = 2\pi\nu_C$ travelling in equal and opposite directions, $\pm\hat{x}$. The amplitude of the combined oppositely-moving waves acting upon an electron will be

$$\phi = \phi_+ + \phi_- = 2 \cos \omega_C t \cos k_C x. \quad (A4)$$

But now assume an electron is moving with velocity v in the $+x$ -direction. The wave responsible for driving the resonant oscillation impinging on the electron from the front will be the wave seen in the laboratory frame to have frequency $\omega_- = \gamma\omega_C(1 - v/c)$, i.e. it is the wave below the Compton frequency in the laboratory that for the electron is Doppler shifted up to the ω_C resonance. Similarly the zero-point wave responsible for driving the electron resonant oscillation impinging on the electron from the rear will have a laboratory frequency $\omega_+ = \gamma\omega_C(1 + v/c)$ which is Doppler shifted down to ω_C for the electron. The same transformations apply to the wave numbers, k_+ and k_- . The Lorentz invariance of the electromagnetic quantum vacuum spectrum ensures that regardless of the electron's (unaccelerated) motion the up- and down-shifting of the laboratory-frame spectral energy density will always yield a standing wave in the electron's frame.

It can be shown [8,15] that the superposition of these two oppositely-moving, Doppler-shifted waves is

$$\phi' = \phi'_+ + \phi'_- = 2 \cos(\gamma\omega_C t - k_B x) \cos(\omega_B t - \gamma k_C x). \quad (A5)$$

Observe that for fixed x , the rapidly oscillating "carrier" of frequency $\gamma\omega_C$ is modulated by the slowly varying envelope function in frequency ω_B . And *vice versa* observe that at a given t the "carrier" in space appears to have a relatively large wave number γk_C which is modulated by the envelope of much smaller wave number k_B . Hence both timewise at a fixed point in space and spacewise at a given time, there appears a carrier that is modulated by a much broader wave of dimension corresponding to the de Broglie time $t_B = 2\pi/\omega_B$, or equivalently, the de Broglie wavelength $\lambda_B = 2\pi/k_B$.

This result may be generalized to include quantum vacuum radiation from all other directions, as may be found in the monograph of de la Peña and Cetto [8]. They conclude by stating: "The foregoing discussion assigns a physical meaning to de Broglie's wave: it is the *modulation* of the wave formed by the Lorentz-transformed, Doppler-shifted superposition of the whole set of random stationary electromagnetic waves of frequency ω_C with which the electron interacts selectively."

Another way of looking at the spatial modulation is in terms of the wave function: the spatial modulation of eqn. (A5) is exactly the $e^{ipx/\hbar}$ wave function of a freely moving particle satisfying the Schrödinger equation. The same argument has been made by Hunter [14]. In such a view the quantum wave function of a moving free particle becomes a "beat frequency" produced by the relative motion of the observer with respect to the particle and its oscillating charge.

It thus appears that a simple model of a particle as an electromagnetic quantum vacuum-driven oscillating charge with a resonance at its Compton frequency may simultaneously offer insight into the nature of inertial mass, i.e. into rest inertial mass and its relativistic extension, the Einstein-de Broglie formula and into its associated wave function involving the de Broglie wavelength of a moving particle. If the de Broglie oscillation is indeed driven by the electromagnetic quantum vacuum, then it is a form of Schrödinger's *zitterbewegung*. Moreover there is a substantial literature attempting to associate spin with *zitterbewegung* tracing back to the work of Schrödinger [17]; see for example Huang [18] and Barut and Zanghi [19].

REFERENCES

- [1] M. Jammer, Concepts of Mass in Contemporary Physics and Philosophy, Princeton Univ. Press (2000)
- [2] B. Haisch, A. Rueda, and H. E. Puthoff, Phys. Rev. A **49** 678 (HRP) (1994)

- [3] W. G. Unruh, Phys. Rev. D, **14** 870 (1976)
- [4] P. C. W. Davies, J. Phys. A, **8** 609 (1975)
- [5] P. Chen, reported at American Astronomical Society meeting, June 6, 2000.
- [6] A. Rueda and B. Haisch, Found. Phys. **28** 1057 (1998); A. Rueda and B. Haisch, Phys. Lett. A **240** 115 (1998)
- [7] T. H. Boyer, Phys. Rev. D, **11** 790 (1975)
- [8] L. de la Peña, and A. M. Cetto, *The Quantum Vacuum: An Introduction to Stochastic Electrodynamics*, Kluwer Acad. Publ. (1996)
- [9] A. Rueda, B. Haisch and R. Tung, in prep. (2001); R. Tung, B. Haisch and A. Rueda, in prep. (2001)
- [10] R. Loudon, *The Quantum Theory of Light* (2nd ed.) Clarendon Press, Oxford (1983)
- [11] S. Hawking, Nature, **248** 30 (1974)
- [12] P. C. W. Davies, T. Dray, and C. A. Manogue, Phys. Rev. D **53** 4382 (1996)
- [13] J. D. Jackson, *Classical Electrodynamics* (3rd ed.) (1999)
- [14] G. Hunter, in *The Present Status of the Quantum Theory of Light*, S. Jeffers et al. (eds.), (Kluwer Acad. Publ.), chap. 12 (1996)
- [15] A. F. Kracklauer, Physics Essays **5** 226 (1992); for a formal derivation and further illuminating discussion see Chap. 12 of [4]
- [16] B. Haisch and A. Rueda, Phys. Lett. A, **268**, 224 (2000)
- [17] E. Schrödinger, Sitz. Ber. Preuss. Akad. Wiss. Phys.-Math. Kl, **24**, 4318 (1930)
- [18] K. Huang, Am. J. Phys. **20** 479 (1952)
- [19] A. O. Barut and N. Zanghi, Phys. Rev. Lett. **52** 209 (1984)

[Home](#)[Publications](#)[Search](#)[My Alerts](#)[My Profile](#)[Help](#)

Quick Search:

within

(?) [Search tips](#)

1 of 3

[results list](#)[◀ previous](#)[next ▶](#)

[Physics Letters A](#)

[Volume 294, Issue 2](#), 18 February 2002, Pages 79-83DOI: [10.1016/S0375-9601\(01\)00672-7](https://doi.org/10.1016/S0375-9601(01)00672-7)

PII: S0375-9601(01)00672-7

Copyright © 2002 Elsevier Science B.V. All rights reserved.

This Document

- [SummaryPlus](#)
- [Full Text + Links](#)
- [PDF \(70 K\)](#)

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

Interplay between gravity and quintessence: a set of new GR solutions

Arthur D. Chernin^{a, b, c}, David I. Santiago^{a, d}, and Alexander S. Silbergleite^a Sternberg Astronomical Institute, Moscow University, Moscow, 119899, Russia^b Astronomy Division, Oulu University, Oulu, 90401, Finland^c Tuorla Observatory, University of Turku, Piikkiö, 21 500, Finland^d Department of Physics, Stanford University, Stanford, CA 94305-4060, USA^e Gravity Probe B, W.W. Hansen Experimental Physics Laboratory, Stanford University, Stanford, CA 94305-4085, USA

Received 15 June 2001; revised 21 August 2001; accepted 8 October 2001 Communicated by P.R. Holland Available online 24 January 2002.

Abstract

A set of new exact analytical general relativity (GR) solutions with time-dependent and spatially inhomogeneous quintessence demonstrate (1) a static non-empty space-time with a horizon-type singular surface; (2) time-dependent spatially homogeneous 'spheres' which are completely different in geometry from the Friedmann isotropic models; (3) infinitely strong **anti-gravity** at a 'true' singularity where the density is infinitely large. It is also found that (4) the GR solutions allow for an extreme 'density-free' form of energy that can generate regular space-time geometries.

PACS classification codes: 98.80.Cq; 04.25.Dm; 04.60.-m

Article Outline

- [1. Introduction](#)
 - [2. Solution S \$\pm\$](#)
 - [3. Solution S-1](#)
 - [4. Solution S-2](#)
 - [5. Solution S-3](#)
 - [References](#)
-

1. Introduction

The idea of quintessence (Q) as a dynamic, time-dependent and spatially inhomogeneous energy with negative pressure-to-density ratio ($p=w\rho$, $-1 < w < 0$) [1] provides new 'degrees of freedom' in cosmology [1 and 2] and extends the variety of modern field models to include extreme forms of energy [1 and 3]. It may also stimulate a better understanding of the fundamental problem of interplay between gravity and the known field theories (for a review of this problem, see [4]).

The equation of state with $w=-1$ represents vacuum-type quintessence (VQ) which is phenomenologically described by the cosmological constant [5]; VQ is known to induce the dynamic effect of cosmological acceleration, if its density is positive. Recent data on the brightness of distant SN Ia [6] (as well as the evidence coming from the cosmic age, large scale structure, and cosmic microwave background anisotropy combined with the cluster dynamics) most probably indicate that the observed cosmological expansion is indeed accelerating if the universe is homogeneous. The physical reason for the cosmic acceleration may generally be attributed not only to VQ, but to any form of Q with $w < -1/3$, $\rho > 0$ or $w > -1/3$, $\rho < 0$, as seen from the Friedmann equations. In the original paper [1], positive energy for Q is preferred; in contrast with that, the whole range $-\infty < \rho < +\infty$ is considered below, for completeness and to examine whether any restrictions on the sign of energy could come from GR, if $w=\text{const}$. Note that the 'inertial energy dominance condition', $\rho + p > 0$, which is not met in the basic case of VQ, $w=-1$, is not satisfied also for some other forms of energy discussed below.

The special case $w=-1/3$ represents the only form of energy (with the linear equation of state) that can generate gravity with zero acceleration or deceleration effect in a homogeneous and isotropic universe, regardless the sign of the energy density. Gravity of this origin reveals itself only in the curvature of the four-dimensional space-time, it has no Newtonian analogs (unlike all other forms of Q, including VQ), and its nature is completely due to general relativity (GR) physics. Historically, Q with $w=-1/3$ was the first energy component that appeared in GR cosmology: in the Einstein static cosmological solution of 1917 the dynamic balance of anti-gravity of the cosmological constant and gravity of pressure-free matter gives rise to an effective equation of state with $w=-1/3$. This special type of Q with $w=-1/3$ may be called *Einstein quintessence* (EQ).

In the present Letter, some special properties of gravity produced by EQ, as well as by other extreme forms of Q in static and time-dependent, spatially homogeneous and inhomogeneous space-times are studied by means of a set of new exact analytical GR solutions. The metric of the solutions has 3D spherical symmetry,

$$ds^2 = A(r,t) dt^2 - B(r,t) d\Omega^2 - C(r,t) dr^2; \quad (1)$$

the components of the metric tensor $A(r,t), B(r,t), C(r,t)$ are functions of the radial coordinate r and time t , and $d\Omega^2 = \sin^2\theta d\varphi^2 + d\theta^2$.

Four major new results are reported in the Letter. (1) The interplay between gravity and Q is capable of creating a static

non-empty space-time with a horizon-type singular surface. (2) This interplay is also revealed in the formation of time-dependent spatially homogeneous 'spheres' which are completely different from the Friedmann isotropic models and have no analogs in Newtonian gravity. (3) Q can induce infinitely strong anti-gravity at a 'true' singularity where the density is infinitely large. (4) GR allows for an extreme 'density-free' form of energy that can generate regular space-time geometries. We use units with $G=c=1$.

2. Solution S \pm

All 'static' (no dependence on t in the metric coefficients (1)) solutions to Einstein's equations (see, for instance, [7]) with EQ equation of state ($w=-1/3$) are given by

$$\begin{aligned} B(r) &= r^2, & C &= \frac{r^4}{a^2} \frac{(A')^2}{A}, \\ 8\pi\rho &= \frac{3a^2}{r^3} \frac{A}{(A')^2} \left(\frac{A''}{A'} + \frac{2}{r} \right), \end{aligned} \quad (2)$$

where $A(r)$ is determined from a quadratic equation

$$kA^2 + \frac{2}{a^2}A + b + \frac{1}{r^2} = 0, \quad (3)$$

with three arbitrary constants k , b and $a^2>0$. For $k\neq 0$ the two ($S\pm$) solutions thus are

$$\begin{aligned} A(r) &= \frac{1}{ka^2} \left[-1 \pm \sqrt{K(r)} \right], \\ C(r) &= \frac{-1 \mp \sqrt{K(r)}}{(br^2+1)K(r)}, \end{aligned} \quad (4)$$

$$\begin{aligned} 8\pi\rho(r) &= -\frac{3(1-ka^4b)}{ka^4} \left[-1 \pm \sqrt{K(r)} \right], \\ K(r) &= 1 - ka^4(b + 1/r^2). \end{aligned} \quad (5)$$

To describe the parameter cases, it is convenient to introduce $q=1-ka^4b$, so that $K(r)=q-ka^4/r^2$. Evidently, $\sqrt{K(r)}$ should be positive at least for some interval of r , which leads to the following:

- (A) For $q<0$, $k>0$ there are no (physical) solutions.
- (B) For $q<0$, $k<0$ we have 'interior' solutions, $r^2\leq ka^4/q$; $S+$ does not have an event horizon for $q<-1/2$, its horizon coincides with the boundary $r^2=2|k|a^4$ for $q=-1/2$, there is a horizon inside the space-time for $-1/2<q<0$; apparently, $S-$ always has no horizon.
- (C) For $q=0$, $k=0$ $S+$ is the Schwarzschild exterior solution, $S-$ is not physical because it is also Schwarzschild but with a negative central mass.
- (D) For $q>0$, $k<0$ both solutions are 'global' (space-time extends to any r), $S+$ has a horizon.
- (E) For $q>0$, $k>0$ both are 'exterior' solutions, $r^2\geq ka^4/q$, $S+$ has no horizon for $0<q<1/2$, its horizon coincides with the boundary $r^2=2ka^4$ for $q=1/2$, there is a horizon inside the space-time for $q>1/2$.

Note that whenever $r=0$ is within the space-time, solutions $S\pm$ have there a true (non-coordinate) singularity: both the Ricci curvature and the density are infinite at $r=0$. For $S-$ this is a naked singularity, with no horizon around it. Note also that in the limit $k\rightarrow 0$ $S+$ goes to the solution $S-1$, which is obtained from Eq.(3) with $k=0$ and is discussed immediately below, while $S-$ is singular.

3. Solution S-1

If $k=0$ in Eq. (3), we have a particular solution for EQ of an especially simple form:

$$A(r) = A_0 \left(1 - \frac{1}{\alpha r^2}\right), \quad B(r) = r^2, \\ C(r) = \frac{2}{\alpha r^2} \left(1 - \frac{1}{\alpha r^2}\right)^{-1}. \quad (6)$$

$$8\pi\rho = \frac{3}{2}\alpha \left(\frac{1}{\alpha r^2} - 1\right). \quad (7)$$

where $\alpha=-b$ and $A_0=a^2\alpha/2$. To keep the proper signature of the metric as $r\rightarrow\infty$, we require $A_0>0$, so that $\alpha>0$; without loss of generality, we set $A_0=1$, i.e., $a^2=2/\alpha=-2/b>0$. Let us discuss this solution in more detail, setting, for simplicity, $\alpha=1$.

(A) The space-time of the solution S-1 has a horizon at $r=1$ where the components of the metric tensor $g_{00}=A(r)$ and $g_{11}=-C(r)$ change their signs. The horizon separates (or connects) a static spatially inhomogeneous exterior space (E-space) at $r>1$ and a 'hidden' interior object (I-object) at $r<1$. These two regions are similar to R- and T-regions, respectively, as described by Novikov [8] for the Schwarzschild solution. The true singularity with the infinite Ricci curvature and density is at $r=0$. The density in S-1 may be regarded as consisting of two components with the same equation of state: one is uniform and negative, $8\pi\rho_-=-3/2<0$, the other is non-uniform ('isothermal' law) and positive, $8\pi\rho_+(r)=3/2r^2>0$. The first one dominates the E-space, and the other one dominates the I-object; the total density changes its sign at the horizon.

(B) The 'energy equation' introduced for the metric of Eq. (1) by Lemaître (see Ref. [5]) may be used to find an analog of the Newtonian gravitational potential as one of the physical characteristics of gravity produced by EQ in S-1. The equation has a form of energy conservation relation, $(1/2)(dB^{1/2}/dt)^2-m(r,t)/r=E$, where $E=C^{-1}(dB^{1/2}/dr)^2-1$ and $m(r,t)$ is the Newtonian analog of gravitating mass. In S-1, the Newtonian mass proves to be $m(r)=(3/4)r[1-(1/3)r^2]$, and the Newtonian potential $u(r)=-m/r=-(3/4)[1-(1/3)r^2]$ describes 'the Newtonian component' of gravity produced by EQ in S-1. The potential u is finite at the true singularity ($u(0)=-3/4$) (unlike that in the Schwarzschild solution) and goes to $+\infty$ at spatial infinity where the density becomes constant (see below). On the singular surface, $m(1)=1/2$, and so $2|u(1)|=1$ (like that in the Schwarzschild solution).

(C) EQ gravity in S-1 is not reduced to Newtonian gravity completely; in particular, the dynamic effect of acceleration is essentially different in S-1 from what may be produced by the Newtonian potential u . The total accelerating effect can be evaluated in terms of the effective potential, $U(r)$, which may be derived using the fact that S-1 describes EQ in a state of hydrostatic equilibrium: its pressure and self-gravity are balanced in the E-space. The pressure gradient produces radial acceleration $F_p=-(dp/dr)(\rho+p)^{-1}=[8\pi(r^3-r)]^{-1}$, which is positive. So the gravity acceleration is $F=F_p-[8\pi(r^3-r)]^{-1}$; it is negative, and therefore gravity is attracting. The effective gravitational potential responsible for this attraction is $U=(1/2)\ln[r^2/(r^2-1)]$.

(D) S-1 can be rewritten in the E-space as

$$ds^2 = \tanh^2 \left(\chi/\sqrt{2} \right) dt^2 - \cosh^2 \left(\chi/\sqrt{2} \right) d\Omega^2 - d\chi^2. \quad (8)$$

$$8\pi\rho = -\frac{3}{2} \tanh \left(\chi/\sqrt{2} \right). \quad (9)$$

with the new spatial coordinate χ related to r by $r = \cosh(\chi/\sqrt{2})$; its range is from zero to infinity. The density decreases monotonically from zero at $\chi=0$ to $-3/2$ at $\chi=\infty$. This form of S-1 may possibly be considered both in relation to and independently of the I-object. In the limit $r\rightarrow\infty$, where the density is spatially homogeneous, the

metric of S-1 takes the form $ds^2 = dt^2 - r^2 d\Omega^2 - 2r^2 dr^2$, or $ds^2 = dt^2 - \exp(\sqrt{2}\chi) d\Omega^2 - d\chi^2$. Since the general static solution for EQ in the homogeneous and isotropic 3-space has the form $ds^2 = dt^2 - r^2 d\Omega^2 - (1 - kr^2/a_0^2)^{-1} dr^2$, $8\pi\rho = -24\pi$ $\rho = 3k/a_0^2$ ($k=1, 0, -1$ is the sign of 3-curvature, a_0 is an arbitrary scaling constant), one can see that the metric of (2) and (4) reduces to the isotropic metric with $k=-1$ and $a_0=1/2$ in the limit $r \rightarrow \infty$. For comparison: $k=1$, $a_0=1$ in the Einstein static solution.

(E) Since $A(r)$ and $C(r)$ in S-1 change their signs at the singular surface, the signature of the metric of the I-object is (- - +). This means that the coordinate r becomes time-like, and the coordinate t becomes space-like inside the I-object. The same is true for the Schwarzschild space-time [8]. Because of that, the density of the I-object depends only on the time-like coordinate, thus it is spatially homogeneous. However, in contrast with the Schwarzschild space-time, a formal replacement of g_{00} with g_{11} and vice versa, which transforms the signature of the I-object to the 'ordinary' type (+-- -), does not work for S-1, because the metric resulting from Eqs. (2) after this transformation is not a solution to the GR equations.

4. Solution S-2

GR equations for the same metric of Eq. (1) allow for the following time-dependent particular solution

$$\begin{aligned} A(t) &= \frac{1}{4} (n^2 - 4) = \text{const} > 0, \\ B(t) &= t^2, \quad C(t) = t^n, \end{aligned} \tag{10}$$

$$\begin{aligned} 8\pi\rho &= \frac{n(n+4)}{n^2 - 4} \frac{1}{t^2}, \quad p = w\rho, \\ n &= -\frac{4w}{1+w} = \text{const}. \end{aligned} \tag{11}$$

The parameter range of this solution apparently consists of two parts: $n>2$ ($-1 < w < -1/3$) and $n<-2$ ($|w|>1$; this is not Q as defined above). Note that both VQ ($w=-1$) and EQ ($w=-1/3$) are outside this range.

(A) The most striking feature of this solution (S-2) is that A , C and even B do not depend on r , and are functions of the time only. One may see here a similarity to the I-object of S-1 whose metric also depends on the time-like coordinate only. A closer similarity may be recognized with the 'T-sphere' found by Ruban [9] for pressure-free matter and the same S-2 symmetry of 3-space with $B=B(t)$. The density and pressure are spatially homogeneous (functions of the time only) in both the S-2 and Ruban's T-sphere. Time t varies in S-2 from $-\infty$ to $+\infty$, and the density varies from zero at $t \rightarrow \pm\infty$ to infinity in the (true) singularity $t=0$ which has the same character as that in homogeneous and isotropic cosmological models. However, there are no coordinate transformations that could reduce S-2 and Ruban's T-sphere to the FRW metric with the isotropic 3-space.

(B) Unlike the Friedmann solutions, gravity in S-2 and Ruban's T-sphere does not have any Newtonian analogs. Rather, the space-time of this special type has common features with anisotropic spatially homogeneous cosmological models (cf. [9]).

(C) The pressure, $p=-n^2/[8\pi(n^2-4)t^2]$, is negative in S-2, while the density may be either positive or negative. The density is positive if w is in 'the Q range' $-1 < w < -1/3$ ($n>2$). (Note that in the isotropic 3-space Q with w in this range produces positive acceleration, see Section 1.) The density is also positive for $w>1$ (i.e., $-4 < n < 2$), but it is negative when $w<-1$ ($n<-4$).

(D) The density turns to zero for $w=\infty$ ($n=-4$). This case describes a form of energy with $\rho=0$, $p<0$, which is perhaps the most extreme form of Q. As we see, GR does not exclude the equation of state with $w=\infty$, and gives a regular solution for it, $A(r)=3$, $B(r)=t^2$, $C=t^{-4}$. 'Density-free' energy may also be in a state of hydrostatic equilibrium (see below).

5. Solution S-3

A power law particular static solution, which is a counterpart to S-2, is also allowed by the GR equations for the metric of Eq.(1):

$$\begin{aligned} A(r) &= r^n, & B(r) &= r^2, \\ C(r) &= \frac{1}{4} (4 + 4n - n^2) = \text{const} > 0, \end{aligned} \quad (12)$$

$$\begin{aligned} 8\pi\rho &= \frac{n(4-n)}{4+4n-n^2} \frac{1}{r^2}, & p &= w\rho, \\ n &= \frac{4w}{1+w}. \end{aligned} \quad (13)$$

The parameter range here looks even more peculiar: $-2(\sqrt{2}-1) < n < 2(\sqrt{2}+1)$, which corresponds to the two intervals for the values of w , $w > -(\sqrt{2}-1)/(\sqrt{2}+1) > -1/3$ and $w < -1$. Neither VQ nor EQ are within this range, and the solution relates to Q only when $-(\sqrt{2}-1)/(\sqrt{2}+1) < w < 0$.

(A) Similar to S-1 and S-2, solution S-3 depends on one coordinate only, and the true singularity of all the three solutions is at the origin of this coordinate. The positive component of density in S-1, and the total densities in S-2 and S-3 follow the inverse square law. Both S-1 and S-3 describe Q in a state of hydrostatic equilibrium.

(B) The pressure, $p=n^2/[8\pi(4+4n-n^2)r^2]$, is positive in S-3 for any $n \neq 0$. At $n=0$ the pressure and density are both identically zero. In this case, $A=1$, $C=1$, and the metric of S-3 turns to the Lorentz metric of empty space-time. The density in S-3 is positive for $0 < n < 4$ and negative otherwise.

(C) The density vanishes also for $n=4$, $w=\pm\infty$. This is the 'density-free' ($\rho=0, p>0$) form of energy, and S-3 (same as S-2 above) gives a regular metric for this case: $A=1$, $B(r)=r^2$, $C(r)=r^4$. The fact that this form of energy is in the state of hydrostatic equilibrium shows that 'density-free' energy has its inertial mass (per unit volume) $\rho_i=p$ which is equal to its passive gravitational mass, and that the active mass density is positive.

(D) For $n \neq 0$, the sign of the active gravitating mass depends on the sign of n . If n is positive, the mass is positive; but if n is negative (and thus w is negative), the active mass is negative also. In terms of Newtonian physics, the negative mass produces a positive acceleration, which goes to infinity as $r \rightarrow 0$, and so anti-gravity is infinitely strong at the singularity. Whether it is capable, under these conditions, of producing 'auto-emission' of particles from the singularity (where the density is infinite), and/or enhance quantum evaporation of particles from the singularity, should be discussed separately.

(E) S-3 describes hydrostatic equilibrium of not only Q with negative w , but also, for instance, of ultra-relativistic fluid ($w=1/3, n=1, C=7/4$) and of Zeldovich ultra-stiff fluid ($w=1, n=2, C=2$).

References

1. R.R. Caldwell, R. Dave and P.J. Steinhard *Phys. Rev. Lett.* **80** (1998), p. 1582. [Abstract-INSPEC](#) | [APS full text](#) | [Full Text via CrossRef](#)
2. R.R. Caldwell, R. Dave and P.J. Steinhard *Appl. Space Sci.* **261** (1998), p. 303. [Full Text via CrossRef](#)
L. Wang and P.J. Steinhard *Astrophys. J.* **508** (1998), p. 483. [Full Text via CrossRef](#)
- C.-P. Ma, R.R. Caldwell, P. Bode and L. Wang *Astrophys. J.* **521** (1999), p. L1. [Full Text via CrossRef](#)
- A.R. Cooray and D. Huterer *Astrophys. J.* **513** (1999), p. L95. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)
- J.S. Alcaniz and J.A.S. Lima *Astron. Astrophys.* **349** (1999), p. 729. [Abstract-INSPEC](#)
- I.S. Zlatev and P.J. Steinhard *Phys. Lett. B* **459** (1999), p. 570. [Abstract](#)

L. Hui *Astrophys. J.* **519** (1999), p. L9. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

I. Zlatev, L. Wang and P.J. Steinhard *Phys. Rev. Lett.* **82** (1999), p. 896. [Abstract-INSPEC](#) | [APS full text](#) | [Full Text via CrossRef](#)

P.J.E. Peebles and A. Vilenkin *Phys. Rev. D* **590** (1999), p. 811.

M. Giovannini *Phys. Rev. D* **601** (1999), p. 277.

L. Wang, R.R. Caldwell, J.P. Ostriker and P.J. Steinhard *Astrophys. J.* **550** (2000), p. 17. [Full Text via CrossRef](#)

G. Efstathiou *MN RAS* **310** (2000), p. 842.

P.P. González-Díaz *Phys. Lett. B* **481** (2000), p. 353. [Abstract](#)

J.D. Barrow, R. Bean and J. Magueijo *MN RAS* **316** (2000), p. L41. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

3. S.M. Carroll *Phys. Rev. Lett.* **81** (1998), p. 3067. [Abstract-INSPEC](#) | [APS full text](#) | [Full Text via CrossRef](#)

S.M. Barr *Phys. Lett. B* **454** (1999), p. 92. [Abstract](#)

R.S. Kalyana *Phys. Lett. B* **457** (1999), p. 268.

Ch. Kolda and D.H. Lyth *Phys. Lett. B* **459** (1999), p. 570.

P. Binétruy *Phys. Rev. D* **600** (1999), p. 80.

R. Horvat *Mod. Phys. Lett. A* **14** (1999), p. 2245. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

T. Chiba *Phys. Rev. D* **601** (1999), p. 4634.

P.H. Brax and J. Martin *Phys. Lett. B* **468** (1999), p. 40. [Abstract](#)

A.B. Kaganovich *Nucl. Phys. B Proc. Suppl.* **87** (1999), p. 496.

O. Bertolami and P.J. Martins *Phys. Rev. D* **610** (2000), p. 7.

Y. Nomura, T. Watari and T. Yanagida *Phys. Lett. B* **484** (2000), p. 103. [Abstract](#)

S.C.C. Ng *Phys. Lett. B* **485** (2000), p. 1. [Abstract](#) | [Full Text via CrossRef](#)

I.G. Dynnikova *Phys. Lett. B* **472** (2000), p. 33.

A. Hebecker and C. Wetterich *Phys. Rev. Lett.* **85** (2000), p. 3339. [Abstract-Compendex](#) | [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

N. Arkani-Hamed, L.J. Hall, C. Colda and H. Murayama *Phys. Rev. Lett.* **85** (2000), p. 4434. [Abstract-Compendex](#) | [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

C. Armendariz-Picon, V. Mukhanov and P.J. Steinhard *Phys. Rev. Lett.* **85** (2000), p. 4438. [Abstract-Compendex](#) | [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

4. S. Weinberg *Rev. Mod. Phys.* **61** (1989), p. 1. [Abstract-INSPEC](#) | [MathSciNet](#) | [Full Text via CrossRef](#)

5. G. Lemaître *Rev. Mod. Phys.* **21** (1949), p. 357. [MathSciNet](#)

E.B. Gliner *JETP* **22** (1966), p. 378.

E.B. Gliner *Sov. Phys. Dokl.* **6** (1970), p. 559.

6. A.G. Riess *et al.* *Astron. J.* **116** (1998), p. 1009. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

S. Perlmutter *et al.* *Astrophys. J.* **517** (1999), p. 565.

J. Cohn (astro-ph/9807128).

S. Carol (astro-ph/0004075).

7. L.D. Landau and E.M. Lifshitz *The Classical Field Theory*, Pergamon Press, London (1959).

8. I.D. Novikov *Comm. Sternberg Astron. Inst.* **132** (1963), p. 43.

9. V.A. Ruban *JETP* **29** (1969), p. 1027. [MathSciNet](#)

 Corresponding author; email: david@small.stanford.edu

This Document

- [SummaryPlus](#)
- ▶ Full Text + Links
- [PDF \(70 K\)](#)

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

1 of 3

[results list](#)

◀ [previous](#)

[next](#) ▶

[Home](#)

[Publications](#)

[Search](#)

[My Alerts](#)

[My Profile](#)

[Help](#)

SCIENCE  DIRECT

Send [feedback](#) to ScienceDirect

Software and compilation © 2002 ScienceDirect. All rights reserved.

ScienceDirect® is an Elsevier Science B.V. registered trademark.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.

Quick Search:

within

(?) [Search tips](#)

2 of 3

[results list](#)[◀ previous](#)[next ▶](#)

[Physics Letters A](#)

[Volume 283, Issues 1-2, 7 May 2001, Pages 62-70](#)[DOI:](#) 10.1016/S0375-9601(01)00229-8

PII: S0375-9601(01)00229-8

Copyright © 2001 Elsevier Science B.V. All rights reserved.

This Document

- [SummaryPlus](#)
- ▶ [Full Text + Links](#)
- [PDF \(111 K\)](#)

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

Classical analogous of quantum cosmological perfect fluid models

A. B. Batista✉, J. C. Fabris✉, S. V. B. Gonçalves✉ and J. Tossa✉, 1

Departamento de Física, Universidade Federal do Espírito Santo, 29060-900, Vitória, Espírito Santo, Brazil

Received 16 January 2001; revised 15 March 2001; accepted 26 March 2001 Communicated by P.R. Holland Available online 2 May 2001.

Abstract

Quantization in the minisuperspace of a gravity system coupled to a perfect fluid, leads to a solvable model which implies singularity free solutions through the construction of a superposition of the wavefunctions. We show that such models are equivalent to a classical system where, besides the perfect fluid, a repulsive fluid with an equation of state $p_Q = -\rho_Q$ is present. This leads to speculate on the true nature of this quantization procedure. A perturbative analysis of the classical system reveals the condition for the stability of the classical system in terms of the existence of an **anti-gravity** phase.

PACS classification codes: 04.20.Cv; 04.20.Me

Article Outline

● [Acknowledgements](#)**● [References](#)**

The existence of an initial singularity is one of the major drawbacks of the so-called standard cosmological model. It is a general belief that such problem can be solved through the employment of a quantum theory of gravity. Indeed, near the singularity sub-Planckian scales are reached and a classical description of the Universe under this situation is not appropriate. However, there is no consistent quantum theory of gravity until now, and in this sense the problem of the initial singularity remains of actuality. On the other hand, it is possible to construct a quantum model for the Universe as a whole, through the Wheeler-DeWitt equation, based in the ADM decomposition of the gravity sector, which leads to a Hamiltonian

formulation of general relativity, from which a canonical quantization procedure can be applied. This gives birth to quantum cosmology [1 and 2].

Quantum cosmology is not free from problems. First, it can be applied only to geometries where a foliation is possible. Moreover, the Hamiltonian formalism leads to a breakdown of general covariance, and the notion of time is lost [3]. There are some recent proposals by which this notion of time can be recovered. One of these proposals is based in the coupling of the gravity sector to a perfect fluid. Using the Schutz's formulation of a perfect fluid [4], a quantization procedure is possible. The canonical momentum associated with the perfect fluid appears linearly in the Wheeler-DeWitt equation, permitting to rewrite this equation in the form of a Schrödinger equation and a time coordinate associated with the matter field can be identified.

Solutions based on this approach reveal that a superposition of the wavefunctions which are solutions of the resulting Schrödinger equation leads to a singularity-free Universe [5, 6, 7 and 8]. The behaviour of the scale factor may be determined in two different ways: calculating the expectation value of the scale factor, in the spirit of the many worlds' interpretation of quantum mechanics; evaluating the Bohmian trajectories in the ontological formulation of quantum mechanics. The results are essentially the same in both approaches and the scale factor display a bounce, the singularity never being reached. It must be noted that even if the two procedures are technically equivalent, they are conceptually very different from each other; there are claims that from the conceptual point of view only the ontological formalism can be consistently applied to quantum cosmology [2].

The existence of a bounce indicates that there is a repulsive effect, of quantum origin, when the scale factor approaches the singularity. In this Letter we study more in detail such scenario. It is shown that the quantum scenario can be reproduced exactly by a very simple classical model where a repulsive fluid is added to the normal perfect fluid. It is surprising that the repulsive fluid is always the same, given by a stiff matter equation of state $p_Q=\rho_Q$, independently of the content of the normal fluid. The existence of this classical analogous of the quantum model leads us to ask questions on the true nature of the quantization in this case. Under which conditions the features of a quantum system can be exactly reproduced by a classical system? Our analysis is restricted to a perfect fluid coupled to gravity system, where the notion of time is recovered. But we sketch some considerations on other situations where gravity is coupled to matter fields.

The existence of a classical analogous of the quantum model allows us to perform a perturbative analysis establishing under which conditions the repulsive phase near singularity may be stable or not. In fact, when the mechanism responsible for the avoidance of the singularity implies the violation of the strong energy condition only, we can expect this configuration to be stable; but, our analysis suggests that the quantum effects are due to a real anti-gravity phase, which can lead to instabilities under certain conditions.

In the perfect fluid formulation developed by Schutz, the degrees of freedom associated with the fluid are given by five scalar potentials in terms of which the four velocity is written:

$$u_\mu = \frac{1}{\mu}(\phi_{,\nu} + \zeta\beta_{,\nu} + \theta S_{,\nu}), \quad (1)$$

where μ is the specific enthalpy. The four velocity is subjected to the condition

$$u^\nu u_\nu = 1 \quad (2)$$

what enables us to express the specific enthalpy in terms of the other five potentials. From now on, we work in natural units, for which $G=c=\hbar=1$. The action is then given by

$$S = \int_M d^4x \sqrt{-g}R + 2 \int_{\partial M} d^3x \sqrt{h}h_{ij}K^{ij} + \int_M d^4x \sqrt{-g}p. \quad (3)$$

where h_{ij} is the metric on the spatial section, K_{ij} is the extrinsic curvature and p is the pressure of ordinary matter. Action (3) is apparently non-covariant because of the pressure term. But, in fact, the constraints intrinsic to this formalism permit to recover the covariance.

We will consider from now on the Robertson-Walker flat geometry (the spatial section must be compact in order to be

consistent with the boundary conditions)

$$ds^2 = N^2 dt^2 - a(t)^2 h_{ij} dx^i dx^j. \quad (4)$$

We assume a barotropic equation of state $p = \alpha \rho$. Analyzing the conjugate momentum, and eliminating non-physical degrees of freedom, we can reduce action (3) to [6]

$$S = \int \left\{ \dot{a} p_a + \dot{\phi} p_\phi + \dot{S} p_S - NH \right\}. \quad (5)$$

where

$$H = -\frac{p_a^2}{24a} + p_\phi^{\alpha+1} a^{-3\alpha} e^S. \quad (6)$$

There are three cases which will interest us here: $\alpha = -1$, $\alpha = 1/3$ and $\alpha = 0$. The dynamical vacuum ($\alpha = -1$) has been studied in [6], the radiative fluid ($\alpha = 1/3$) in [5, 8 and 9], while the dust case ($\alpha = 0$) was treated in [7].

Following the Schutz formalism for the description of perfect fluid, and specializing it to a dust fluid, with $p = 0$, we obtain the following Lagrangian:

$$L = \dot{a} p_a + \dot{\phi} p_\phi - NH, \quad (7)$$

where ϕ is the dust variable, with p^ϕ being its conjugate momentum, and H is the Hamiltonian

$$H = -\frac{p_a^2}{24a} + e^S p_\phi. \quad (8)$$

Classically, this system admits, for the flat case, the well-known dust solution $a \propto t^{2/3}$, where t is the proper time, or equivalently $a \propto \eta^2$, where η is the conformal time.

Now, we perform the canonical transformation

$$\tau = p_S e^{-S} p_\phi^{-1}, \pi \tau = p_\phi e^S, \quad (9)$$

reducing the Lagrangian to the form

$$H = -\frac{p_a^2}{24a} + p_\tau. \quad (10)$$

Imposing the quantization rules

$$p_a \rightarrow -i \frac{\partial}{\partial a}, \quad p_\tau \rightarrow -i \frac{\partial}{\partial \tau}. \quad (11)$$

and considering that the Hamiltonian becomes an operator which acts on the wavefunction annihilating it,

$$\bar{H} \Psi = 0, \quad (12)$$

we obtain the following partial differential equation governing the behaviour of the wavefunction:

$$\frac{1}{24} \frac{\partial^2 \Psi}{\partial a^2} - ia \frac{\partial \Psi}{\partial \tau} = 0. \quad (13)$$

The fact that the conjugate momentum p_τ associated to the dust fluid variable appears linearly in the Hamiltonian, implies that the Wheeler-DeWitt equation in the minisuperspace assumes a form similar to the Schrödinger equation with τ playing the role of time. It is possible to show, through the canonical transformations, that in this cases it is the cosmic time.

Performing the redefinitions [10]

$$\alpha = \frac{R}{\sqrt{12}}, \quad \tau = \frac{t}{\sqrt{12}}. \quad (14)$$

we end up with the equation

$$\frac{1}{2} \frac{\partial^2 \Psi}{\partial R^2} = iR \frac{\partial \Psi}{\partial t}. \quad (15)$$

We solve Eq. (15) using the method of separation of variables. It leads to the following decomposition of $\Psi(R,t)$:

$$\Psi(R,t) = \xi(R)e^{iEt}, \quad (16)$$

where E is a (positive) constant, and $\xi(R)$ obeys the equation

$$\xi'' + 2RE\xi = 0. \quad (17)$$

The prime means derivative with respect to R . The solution for (17) is under the form of Bessel functions:

$$\xi(R) = \sqrt{R} \left(c_1 J_{1/3}(\beta R^{3/2}) + c_2 J_{-1/3}(\beta R^{3/2}) \right), \quad (18)$$

with $\beta = \sqrt{8E/9}$. The condition for the Hamiltonian operator to be self-adjoint leads to two possible boundary conditions:

$$\xi(0) = 0 \text{ or } \xi'(0) = 0. \quad (19)$$

The final results is insensitive to which boundary condition we employ. Hence, we will work with the first one, but all results are essentially recovered if the second condition is used.

The general solution is a superposition of (18). In order to have analytical expressions, we will use the following superposition:

$$\Psi(R,t) = \sqrt{R} \int_0^\infty \beta^{4/3} e^{-\gamma\beta} e^{i(9/8)\beta^2 t} J_{1/3}(\beta R^{3/2}) d\beta. \quad (20)$$

where γ is a kind of "Gaussian factor" determining the rapidity the wavefunction goes to zero as β increases. Its solution is [11]

$$\Psi(R,t) = \frac{R}{(2A)^{4/3}} e^{-R^3/4A}, \quad (21)$$

where $A = \gamma - i(9/8)t$.

It may be asked which predictions such model makes for the behaviour of the scale factor. Using the many worlds' interpretation, this mounts to evaluate the expectation value of the scale factor. It must be stressed that essentially the same result is achieved by calculating the Bohmian trajectories [2]. The measure employed in the expression is imposed again by the self-adjoint condition, and the expression for the expectation value reads

$$\sqrt{12} \langle \alpha(t) \rangle = \langle R \rangle = \frac{\int_0^\infty R^2 \Psi(R,t)^* \Psi(R,t) dR}{\int_0^\infty R \Psi(R,t)^* \Psi(R,t) dR}. \quad (22)$$

Using (21), the expectation value for the scale factor can be calculated and the final result is

$$\langle \alpha(t) \rangle = \frac{1}{\sqrt{12}} \left(\frac{2}{\gamma} \right)^{1/3} \frac{\Gamma(5/3)}{\Gamma(4/3)} \left(\gamma^2 + \frac{81}{64} t^2 \right)^{1/3}. \quad (23)$$

The parameter t can be identified as the cosmic time due to the canonical transformations employed before. Notice that the

classical behaviour is recovered for $t \rightarrow \infty$. But, in general, the quantum model predicts a non-singular model exhibiting a bounce: when the singularity is approached, quantum effects leads to a repulsive effect, which leads to a regular transition from a contracting to an expanding phase.

A dynamical vacuum and a radiative fluid can be analyzed through the same lines as before. These problem were treated in [5, 6 and 8] and we just present the final results.

The dynamical vacuum is realized through the equation of state $p = -\rho$. Using again the Schutz formalism for this particular case, we rewrite the Hamiltonian (6), with $\alpha = -1$, as

$$H = -\frac{p_a^2}{24a} + \alpha^3 p_T \quad (24)$$

through the canonical transformations $p_T = e^S$ and $T = -e^{-S} p_S$ [6]. Since p_T appears linearly in the Hamiltonian, it can be identified with a time variable as before, in this case in the gauge $dt = dT/a^3$, where t is the cosmic time [6]. Finding the solutions of the corresponding Wheeler-DeWitt equation, evaluating the expectation value of the scale factor, it results

$$\langle a(T) \rangle = \frac{\Gamma(4/3)}{\Gamma(7/6)} \left[\frac{64\gamma^2 + 9T^2}{8\gamma} \right]^{1/6}. \quad (25)$$

Asymptotically the classical solution for a cosmological constant is recovered if we choose, in the classical equations of motion, the time gauge $N = a^{-3}$ (the identification of the time coordinates can be justified rigorously [6] using the canonical transformations). In terms of the proper time, the solution (25) can be rewritten as

$$\langle a(t) \rangle = \frac{\Gamma(4/3)}{\Gamma(7/6)} \sqrt{8\gamma} \left\{ \cosh \left[\frac{3}{\sqrt{8\gamma}} \left(\frac{\Gamma(4/3)}{\Gamma(7/6)} \right)^{1/3} t \right] \right\}^{1/3}. \quad (26)$$

For the radiative case, Hamiltonian (6) reads [5 and 9], for $\alpha = 1/3$ and after the canonical transformations $T = p_S e^{-S} \phi^{-4/3}$ and $p_T = p \phi^{4/3} e^S$,

$$H = -\frac{p_a^2}{24a} + \frac{p_T}{a}. \quad (27)$$

Again, the conjugate momentum associated to the matter variable appears linearly and consequently we can identify this matter variable with a time variable in the gauge $dt = a d\eta$, with is justified through the canonical transformation as in the preceding cases. Hence this time variable is the conformal time. The wavefunctions can be determined through the Green's function method. Fixing the wavefunction at $t=0$ as [9]

$$\Psi(a) = \left(\frac{8\sigma_1}{\pi} \right)^{1/4} e^{-(\sigma_1 + i\sigma_2)a^2} \quad (28)$$

and using the harmonic oscillator Green's function, integrable expression can be obtained [5 and 9]. The scale factor expectation value is given by

$$\langle a(\eta) \rangle = \frac{1}{12} \sqrt{\frac{2}{\pi\sigma_1}} \sqrt{\sigma_1^2 \eta^2 + (6 - \sigma_2 \eta)^2}. \quad (29)$$

where η is the conformal time, σ_1 and σ_2 being real parameters. Again, this solution represents a non-singular eternal Universe which coincides asymptotically with the classical radiative solution $a \propto \eta$.

It must be stressed that in all cases, the classical behaviour is recovered for large values of the proper time. Also, all solutions are singularity-free, with a bounce. Near the bounce repulsive effects appear which, in the ontological formulation, are connected with the quantum potential which corrects the classical equations of motion.

A general feature of the quantum models developed previously is the appearance of a repulsive phase for small values of the scale factor, leading to the avoidance of the singularity. In [12], it was shown that a repulsive gravity single fluid model can lead to consistent cosmological models if the curvature is negative; however, its stability is not assured in the absence of ordinary (attractive) matter. Another way of implementing a repulsive phase in classical cosmology is to consider two fluids, one that acts attractively, and the other that acts repulsively. In this case, we may have consistent solutions with flat spatial section. It is desirable that the repulsive fluid dominates for small values of the scale factor, whereas the attractive fluid dominates for large values of the scale factor. Hence, in the general, considering just the flat spatial section, we may obtain possible consistent models from

$$\left(\frac{\dot{a}}{a}\right)^2 = 8\pi G(\rho_M - \rho_Q) = \frac{C_1}{a^m} - \frac{C_2}{a^n}, \quad (30)$$

where $\rho_M = \alpha_M \rho_M$, $\rho_Q = \alpha_Q \rho_Q$, $m=3(1+\alpha_M)$ and $n=3(1+\alpha_Q)$. The subscripts M and Q stand for "normal" matter component and for "quantum" repulsive component.

Ordinarily, normal matter may be a cosmological constant, dust or a radiative fluid, corresponding to $\alpha_M=-1, 0, 1/3$, respectively. Since it is desirable that the repulsive component dominates at small values of a , then $\alpha_Q > 1/3$. We choose then a repulsive stiff matter fluid $\alpha_Q=1$, what leads to $n=6$. Hence, we will solve Eq. (30) with $n=6$ and $m=0, 3$ and 4 . The solutions are the following:

$$\alpha_M=-1, \alpha_Q=1:$$

$$a(t) = \left(\frac{C_2}{C_1}\right)^{1/6} \cosh^{1/3} 3\sqrt{C_1}t, \quad (31)$$

$$\alpha_M=0, \alpha_Q=1:$$

$$a(t) = \left(\frac{C_2}{C_1}\right)^{1/3} \left[\frac{9C_1^2}{4C_2} t^2 + 1 \right]^{1/3}. \quad (32)$$

$$\alpha_M=1/3, \alpha_Q=1:$$

$$a(\eta) = \sqrt{\frac{C_2}{C_1}} \left[\frac{C_1^2}{C_2} (\eta - \eta_0)^2 + 1 \right]^{1/2}. \quad (33)$$

The comparison of the above solutions with those obtained through the construction of a superposition of the wavefunctions resulting from the Wheeler-DeWitt equation in the minisuperspace with only the ordinary perfect fluid, reveals that they are the same provided that C_1 and C_2 are suitable functions of the parameters of the wavepackets. For the dust case, for example, we can fix

$$\begin{aligned} C_1 &= \frac{1}{12^{3/2}} \frac{9}{8} \frac{1}{y} \left(\frac{\Gamma(5/3)}{\Gamma(4/3)} \right)^3, \\ C_2 &= \frac{1}{12^3} \frac{9}{4} \left(\frac{\Gamma(5/3)}{\Gamma(4/3)} \right)^6. \end{aligned} \quad (34)$$

For the other cases, the expressions for these constants are quite similar. But notice that, in the radiative case, we have introduced a phase η_0 , which can be zero only if α_2 is also zero. Hence, wavepackets constructed from a quantum model where, besides the scale factor, there is a perfect fluid matter degree of freedom (which leads to a time coordinate) are equivalent to a classical model where gravity is coupled to the same perfect fluid plus a repulsive fluid with a stiff matter equation of state $\rho_Q = \rho_Q$. It is really surprising that the repulsive fluid, in the classical model, is the same irrespective of the normal fluid employed in the quantum model.

The question of reproducing the classical equations of motion from the quantum ones appears already in the ordinary quantum mechanics, and they are expressed in the so-called Ehrenfest's theorem. According to this theorem, the center of the wavepacket may follow a classical trajectory under certain conditions. Explicitly, taking the expectation values of the

Heisenberg's equations for the position and momentum operators for a particle of mass m in a potential $V(\vec{r})$, we find [13]

$$\langle \dot{\vec{r}} \rangle = \frac{\vec{p}}{m}, \quad (35)$$

$$\langle \dot{\vec{p}} \rangle = -\langle \nabla V(\vec{r}) \rangle. \quad (36)$$

These relations coincide with the classical one only if $\langle \nabla V(\vec{r}) \rangle = \nabla V(\langle \vec{r} \rangle)$. This happens only for very special forms of the potential, the harmonic oscillator being an example. Only in these special cases, we may say that the center of the wavepacket follows a classical trajectory.

However, the situation discussed here is somehow different from that analyzed in the Ehrenfest's theorem. In fact the expectation value of the scale factor of a quantum model derived from gravity and a perfect fluid of attractive nature is reproduced by a classical model where another fluid, of repulsive nature, appears. This seems somehow mysterious.

Some insights into what is happening in this case may come from the employment of the ontological interpretation of quantum mechanics. In this case, the problem of time is solved in any situation (not only when a matter field is present). In fact, the ontological interpretation predicts that the system follows a real trajectory given by the equations

$$p_q = S_{,q}, \quad (37)$$

where the subscript q designates one of the degrees of freedom of the system, and S is the phase of the wavefunction, which is written as $\Psi = R e^{iS}$, R and S being real functions. The equation of motion (37) is governed not only by a classical potential V but also by a quantum potential $V_Q = \nabla^2 R / R$.

These considerations suggest that the quantum potential has, at least in the case of the quantization of perfect fluid systems, a very clear behaviour which can be classically reproduced by a repulsive stiff matter fluid. However, we must stress that even in this case we must find first the wavefunction, through the Wheeler-DeWitt equation, determining than its phase, from which the Bohmian trajectories is computed. In the quantum models studied previously, it is not possible, in principle, to identify a classical and quantum potential in terms of the scale factor from the beginning. Moreover, even if this would be possible, the classical analogous we have determined are completely independent of the Wheeler-DeWitt equation.

We may obtain more informations on the meaning of this classical analogous of a quantum system by studying the dispersion of the wavepacket. We will treat again the dust case where the wavepacket is given by (21). The calculation is standard and we obtain the following expression for the dispersion of the wavepacket:

$$\Delta a(t) = \left[\langle a^2(t) \rangle - \langle a(t)^2 \rangle \right]^{1/2} = \frac{1}{\sqrt{12}} \left(\frac{2}{\gamma} \right)^{1/3} \left[\frac{\Gamma(2)}{\Gamma(4/3)} - \frac{\Gamma(5/3)^2}{\Gamma(4/3)^2} \right]^{1/2} \left(\gamma^2 + \frac{81}{64} t^2 \right)^{1/3}. \quad (38)$$

Hence, the wavepacket disperse as time evolves; its minimal dispersion is reached in the origin $t=0$, exactly where the classical and quantum behaviours (both with just the dust fluid) do not coincide. Notice that the ratio of $\Delta a(t)$ to $\langle a(t) \rangle$ is constant, and as the wavepacket becomes more localized, the scale factor approaches the singularity. A localized wavepacket means a small value for γ . Notice that as γ becomes small, the coefficient C_2 becomes negligible compared with C_1 (Eqs. (34)), and the two-fluid model tends to approach the one fluid model.

The situation described above is very similar to what happens with a free particle in ordinary quantum mechanics: the wavepacket has its minimum width at origin and disperse as time evolves. In this case the question of the classical limit is somehow subtle. If an ensemble of classical particles, with some initial dispersion in their velocities, is settled in the origin, their trajectories do not reproduce the possible results of measurement of the free quantum particle initially; as time evolves, the dispersion of the wavepacket tends to coincide with the dispersion of the trajectories of the classical particles. This does not mean that notion of classical trajectory is recovered [13 and 14].

In quantum cosmology, however, the scenario is more complex. In this case, we may use the many worlds' interpretation [15] or the ontological interpretation [16] of quantum mechanics. In any case, the notion of trajectory is recovered. In this sense, the above results indicate that, for large values of time, the quantum trajectory coincides with the corresponding

classical trajectory with suitable initial conditions, while near the origin these trajectories do not coincide. But, our analogous model, with two fluids, reproduce the quantum trajectory even in the vicinity of the origin, keeping the asymptotical agreement untouched.

Since we have a classical analogous of the bounce models determined through the Wheeler-DeWitt equation, we can investigate if the repulsive effect leading to the avoidance of the singularity may spoil the stability of the model. First of all we define what we understand here by instability. A cosmological model is considered unstable if the perturbative variables diverge when all background quantities are finite. Here, we will consider a weaker condition: the model is unstable if the perturbed quantities takes very large values in comparison with the background quantities, even if they are not divergent. This is due to the fact that, if this happens, the hypothesis of homogeneity and isotropy, employed in the definition of the background, are compromised.

Let us consider our non-singular classical system. It can be written as

$$R_{\mu\nu} = 8\pi G \left[\frac{M}{T_{\mu\nu}} - \frac{1}{2} g_{\mu\nu} \frac{M}{T} \right] - 8\pi G \left[\frac{Q}{T_{\mu\nu}} - \frac{1}{2} g_{\mu\nu} \frac{Q}{T} \right]. \quad (39)$$

$$\frac{\frac{M}{T}{}^{\mu\nu}}{T_{;\mu}} = 0, \quad (40)$$

$$\frac{\frac{Q}{T}{}^{\mu\nu}}{T_{;\mu}} = 0. \quad (41)$$

We perturb these equations in the usual way, introducing the quantities

$$\begin{aligned} g_{\mu\nu} &= \frac{0}{g_{\mu\nu}} + h_{\mu\nu}, & \rho_M &= \frac{0}{\rho_M} + \delta\rho_M, \\ \rho_Q &= \frac{0}{\rho_Q} + \delta\rho_Q. \end{aligned}$$

The computation of the perturbed equations is quite standard [17], and we just present the final equations, at linear level:

$$\ddot{h} + 2\frac{\dot{a}}{a}\dot{h} = \frac{1}{\alpha_M - \alpha_Q} \left[-(1+3\alpha_M) \left(2\frac{\ddot{a}}{a} + (1+3\alpha_Q)\frac{\dot{a}^2}{a^2} \right) \Delta_M + (1+3\alpha_Q) \left(2\frac{\ddot{a}}{a} + (1+3\alpha_M)\frac{\dot{a}^2}{a^2} \right) \Delta_Q \right], \quad (42)$$

$$\dot{\Delta}_M + (1+\alpha_M) \left(\Psi - \frac{h}{2} \right) = 0, \quad (43)$$


$$(1+\alpha_M) \left[\dot{\Psi} + (2-3\alpha_M)\frac{\dot{a}}{a}\Psi \right] - \frac{n^2}{a^2} \alpha_M \Delta_M = 0, \quad (44)$$

$$\dot{\Delta}_Q + (1+\alpha_Q) \left(\theta - \frac{h}{2} \right) = 0, \quad (45)$$


$$(1+\alpha_Q) \left[\dot{\theta} + (2-3\alpha_Q)\frac{\dot{a}}{a}\theta \right] - \frac{n^2}{a^2} \alpha_Q \Delta_Q = 0. \quad (46)$$

In these expressions, we have used the following definitions: $h=h_{kk}/a^2$, $\Delta_M=\delta\rho_M/\rho_M$, $\Delta_Q=\delta\rho_Q/\delta\rho$, $\Psi=\delta u_M^i$, $\theta=\delta u_Q^i$, where δu_M^i and δu_Q^i are the perturbations on the four velocity of the normal and repulsive fluid, respectively.


The perturbed equations presented above do not seem to admit analytical solutions for the background solutions (31), (32) and (33). Hence, we are obliged to perform a numerical integration. In Fig. 1, Fig. 2 and Fig. 3 we display the evolution of density perturbations for the exotic fluid for the cosmological constant, dust and radiative cases, respectively, in the long wavelength limit $n \rightarrow 0$. The other perturbed quantities exhibits essentially the same features.



(2K)

Fig. 1. Behaviour of $\Delta_Q(t)$ for $n=0$ with cosmological constant.

(2K)

Fig. 2. Behaviour of $\Delta_Q(t)$ for $n=0$ with a dust fluid.

(2K)

Fig. 3. Behaviour of $\Delta_Q(t)$ for $n=0$ with a radiative fluid.

From the numerical study performed we can expose the following conclusions. When the cosmological constant is coupled to the repulsive fluid, the bounce model is unstable. Approaching the minimum of the scale factor (which in all three cases occurs in the origin), the perturbations diverge. However, when the dust or radiative fluid is coupled to the repulsive fluid, the perturbations behave regularly during all the evolution of the Universe. Hence, these models are stable. The different behaviours for the cosmological constant and the other cases may be easily understood. Indeed, in [12] a stability study was performed for the same repulsive fluid (represented there by a free scalar field); the curvature was taken to be negative. Instabilities were found in the absence of ordinary matter. In the present cosmological constant case, as it happens also in [12], the perturbation of ordinary matter decouples completely and the metric perturbation is coupled only to the perturbation of the repulsive fluid. In contrast, for the radiative and dust cases all perturbed variables are coupled. More important, in the present cosmological constant case, and in the model studied in [12], the energy conditions are violated during all the evolution of the Universe, while for the dust and radiative cases studied here, this violation occurs only near the bounce. The violation of the energy conditions leads to a repulsive gravity effect which contributes to a very fast growing of perturbations. If it occurs for all times, this lead to divergence.

In spite of the fact that we display the results for $n=0$ only, for other values of n the features are very similar. For very large values of n , the perturbations exhibit strong oscillations, and they become divergent near the minimum of the scale factor for the cosmological constant case. Of course, we have studied the stability of the classical analogous of the quantum model. But, this study leads to some insights to what happens in the original framework. Indeed, the considerations made above about the expectation value of the scale factor and the dispersion of the wavepacket permits to take seriously the classical perturbative study as indication of the behaviour of the quantum model, mainly if we take the point of view that in quantum cosmology the notion of trajectory is essential. The only important drawback concerns the fact that a complete quantum perturbation should take into account excitations of degrees of freedom that were frozen in the minisuperspace approach. Other important point is what we mean by "near the bounce", where instabilities may appear. In fact, we consider it as the region where the quantum effects are dominant and all energy conditions are violated.

The main point of the present work is that the quantum model, derived from the Wheeler-DeWitt equation for gravity plus

perfect fluid through the Schutz's formalism, has a classical analogous. In this classical analogous system, the perfect fluid is coupled to another perfect fluid, with a fixed equation of state $p_Q = -\rho_Q$ which appears with a "wrong" sign for the gravitational coupling. The existence of this classical analogous for all equations of state of the normal fluid studied in this work, rises doubts about the true quantum nature of the original quantum cosmological scenario. It is not clear to us how to solve these doubts for the moment. But, the existence of a classical system reproducing different quantum models may indicate that the quantization of a gravity system in the minisuperspace may be not a real quantization and a more careful analysis of this problem is deserved.

In the analysis performed previously, it has been considered a specific superposition of the solutions of the Wheeler-DeWitt equation. Since these solutions are not square integrable, a superposition of them is in fact a necessity in those models. It can be argued that other types of superpositions are possible which may not be in agreement with the classical analogous treated here; all the richness of the original quantum model would not be reproduced by the classical model. However, the superposition procedure must agree with physical requirements as, for example, the localization of the wavepacket, what is the case of the preceding examples. We may guess that other possible superpositions, satisfying the same physical requirements, will lead to essentially the same results. These comments can also be applied to the question of the boundary conditions. Perhaps, the choice of other boundary conditions could lead to different scenarios. But, the boundary conditions used here are motivated by physical requirements, like the regularity of the wavefunction and the self-adjointness of the Hamiltonian operator. We may guess that any physically reasonable boundary condition would lead to similar scenarios to those presented here.

A perturbative study was performed in the classical model. It can happens that a bounce model, where the avoidance of the singularity is obtained through an anti-gravity phase, may not be a stable model. We have verified that when the normal fluid decouples from the other perturbed equations, in such a way that the metric perturbation are coupled to the repulsive fluid only, the background model is unstable. Otherwise, we can obtain stable singularity-free models with an anti-gravity phase.

We must stress that the classical analogous model reveals that the "quantum effects" exhibit an anti-gravity behaviour. That is, the singularity is avoided with conditions much more stronger than the simple violation of the strong energy condition, as it happens in many others singularity-free models [18]. It must also be emphasized that all considerations have been done for a perfect fluid quantum model. It should be important to verify if the correspondence found here remains when gravity are coupled to matter fields. In [19] the case of a free scalar field was analyzed. However, a free scalar field is equivalent to stiff matter. Consequently, the classical analogous (if still valid for this case) would contain two kinds of stiff matter, an attractive one and a repulsive one. Perhaps, the strange behaviour found in [19], with the quantum phase being recovered for large values of the scale factor, is due to this fact. This specific case deserves to be analyzed carefully.

Acknowledgements

We have benefited of many enlightfull discussions with N. Pinto-Neto and Nivaldo Lemos. A.B.B., J.C.F. and S.V.B.G. thank CNPq (Brazil) for financial support. J.T. thanks CAPES (Brazil) for financial support and the Department of Physics of the Universidade Federal do Espírito Santo for hospitality.

References

- [1.](#) J.A. Halliwell In: S. Coleman, J.B. Hartle, T. Piran and S. Weinberg, Editors, *Quantum Cosmology and Baby Universes*, World Scientific, Singapore (1991).
- [2.](#) N. Pinto Neto In: M. Novello, Editor, *Cosmology and Gravitation II* (1996), p. 229.
- [3.](#) C.J. Isham (gr-qc/9210011).
- [4.](#) B.F. Schutz *Phys. Rev. D* **2** (1970), p. 2762. [Abstract-INSPEC](#) | [MathSciNet](#) | [Full Text via CrossRef](#)
B.F. Schutz *Phys. Rev. D* **4** (1971), p. 3559. [Abstract-INSPEC](#) | [Full Text via CrossRef](#)

- [5.](#) N.A. Lemos *J. Math. Phys.* **37** (1996), p. 1449. [Abstract-INSPEC](#) | [MathSciNet](#) | [Full Text via CrossRef](#)
- [6.](#) F.G. Alvarenga and N.A. Lemos *Gen. Rel. Grav.* **30** (1998), p. 681. [Abstract-INSPEC](#) | [MathSciNet](#)
- [7.](#) M.J. Gotay and J. Demaret *Phys. Rev. D* **28** (1983), p. 2402. [Abstract-INSPEC](#) | [MathSciNet](#) | [Full Text via CrossRef](#)
- [8.](#) J. Acacio de Barros, N. Pinto-Neto and M.A. Sagioro-Leal *Phys. Lett. A* **241** (1998), p. 229.
- [9.](#) V.G. Lapchinskii and V.A. Rubakov *Theor. Math. Phys.* **33** (1977), p. 1076.
- [10.](#) F.G. Alvarenga and N.A. Lemos *Gen. Rel. Grav.* **31** (1999), p. 1743.
- [11.](#) I.S. Gradshteyn and I.M. Rizhik *Tables of Integrals, Series and Products*, Academic Press, New York (1980).
- [12.](#) A.B. Batista, J.C. Fabris and S.V.B. Gonçalves (gr-qc/0009040) to appear in *Class. Quantum Grav.*.
- [13.](#) C. Cohen-Tannoudji, B. Diu and F. Laloë *Quantum Mechanics* **1**, Wiley (1977).
- [14.](#) A. Messiah *Quantum Mechanics* **1**, North-Holland, Amsterdam (1961).
- [15.](#) F.J. Tipler *Phys. Rep.* **137** (1986), p. 231. [Abstract-INSPEC](#) | [MathSciNet](#)
- [16.](#) P.R. Holland *The Quantum Theory of Motion*, Cambridge University Press, Cambridge (1993).
- [17.](#) S. Weinberg *Gravitation and Cosmology*, Wiley, New York (1972).
- [18.](#) C.P. Constantinidis, J.C. Fabris, R.G. Furtado and M. Picco *Phys. Rev. D* **61** (2000), p. 043503.
- [19.](#) R. Colistete, Jr., J.C. Fabris and N. Pinto Neto *Phys. Rev. D* **57** (1998), p. 4707. [Abstract-INSPEC](#) | [MathSciNet](#) | [APS full text](#) | [Full Text via CrossRef](#)

¹ Permanent address: IMSP, Université Nationale du Bénin, Porto Novo, Benin.

 Corresponding author; email: fabris@cce.ufes.br

[Physics Letters A](#)

[Volume 283, Issues 1-2, 7 May 2001, Pages 62-70](#)

This Document

- [SummaryPlus](#)
- Full Text + Links
- [PDF \(111 K\)](#)

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

2 of 3

[results list](#) | [◀ previous](#) | [next ▶](#)

[Home](#)

[Publications](#)

[Search](#)

[My Alerts](#)

[My Profile](#)

[Help](#)

SCIENCE @ DIRECT

Send [feedback](#) to ScienceDirect

Software and compilation © 2002 ScienceDirect. All rights reserved.

ScienceDirect® is an Elsevier Science B.V. registered trademark.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.

Quick Search:

within

(?) [Search tips](#)

3 of 3

[results list](#)[◀ previous](#)[next ▶](#)

[Physics Letters B](#)

[Volume 489, Issues 1-2, 14 September 2000, Pages 203-206](#)[DOI:](#) 10.1016/S0370-2693(00)00917-5

PII: S0370-2693(00)00917-5

Copyright © 2000 Published by Elsevier Science B.V. All rights reserved.

Gravity and antigravity in a brane world with metastable gravitons

Ruth Gregory^a, Valery A. Rubakov^b and Sergei M. Sibiryakov^b^a Centre for Particle Theory, Durham University, South Road, Durham, DH1 3LE, UK^b Institute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary prospect, 7a, Moscow 117312, Russia

Received 26 July 2000; accepted 11 August 2000. Editor: P.V. Landshoff. Available online 7 September 2000.

This Document[► Abstract](#)**Actions**

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

Abstract

In the framework of a five-dimensional three-brane model with quasi-localized gravitons we evaluate metric perturbations induced on the positive tension brane by matter residing thereon. We find that at intermediate distances, the effective four-dimensional theory coincides, up to small corrections, with General Relativity. This is in accord with Csaki, Erlich and Hollowood and in contrast to Dvali, Gabadadze and Porrati. We show, however, that at ultra-large distances this effective four-dimensional theory becomes dramatically different: conventional tensor gravity changes into scalar **anti-gravity**.

[Physics Letters B](#)

[Volume 489, Issues 1-2, 14 September 2000, Pages 203-206](#)**This Document**[► Abstract](#)**Actions**

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

3 of 3

[results list](#)[◀ previous](#)[next ▶](#)

Weak gravitation shielding properties of composite bulk $YBa_2Cu_3O_{7-x}$ superconductor below 70 K under e.m. field.

E.E. Podkletnov

Moscow Chemical Scientific Research Centre

113452 Moscow - Russia

Abstract

A high-temperature $YBa_2Cu_3O_{7-x}$ bulk ceramic superconductor with composite structure has revealed weak shielding properties against gravitational force while in a levitating state at temperatures below 70 K. A toroidal disk with an outer diameter of 275 mm and a thickness of 10 mm was prepared using conventional ceramic technology in combination with melt-texture growth. Two solenoids were placed around the disk in order to initiate the current inside it and to rotate the disk about its central axis. Samples placed over the rotating disk initially demonstrated a weight loss of 0.3-0.5%. When the rotation speed was slowly reduced by changing the current in the solenoids, the shielding effect became considerably higher and reached 1.9-2.1% at maximum.

74.72.-h High- T_c cuprates.

1 Introduction.

The behavior of high- T_c ceramic superconductors under high-frequency magnetic fields is of great interest for practical applications. Crystal structure seems to be the key factor determining all physical properties of bulk superconductors, and the interaction of this structure with

external and internal e.m. fields might result in quite unusual effects. Despite a large number of studies [1, 2, 3] the nature of these interactions still remains unresolved.

Our recent experimental work [4] clearly indicated that under certain conditions single-phase bulk, dense $YBa_2Cu_3O_{7-x}$ revealed a moderate shielding effect against gravitational force. In order to obtain more information about this unusual phenomenon, a new installation was built, enabling operation with larger disks (275 mm in diameter), in magnetic fields up to 2 T and frequencies up to 10⁸ Hz at temperatures from 40 to 70 K. A new experimental technique was employed to modify the structure of the ceramic superconductor. All these efforts yielded a larger value of the shielding effect (up to 0.5% in stationary conditions and to 2.1% for shorter periods), providing good hopes for technological applications.

A gravitational shielding effect of this strength has never been previously observed, and its implications present serious theoretical difficulties (see [11] for references and an analysis of some hypotheses). Thus, great attention was devoted to the elimination of any possible source of systematic errors or of spurious non-gravitational effects. The small disturbances due to air flows pointed out by some authors [9, 10] were eliminated by weighing the samples in a closed glass tube (see Section 4). The entire cryostat and the solenoids were enclosed in a stainless steel box. But probably the best evidence for the true gravitational nature of the effect is that the observed weight reductions (in %) were independent of the mass or chemical composition of the tested samples (Section 6).

This work is organized as follows. Sections 2 and 3 describe our experimental setup. Section 2 summarizes the main steps in the sinterization of the composite ceramic disk and contains information about the final properties of the disk (T_c for the two layers, J_c for the upper layer, etc.) and about the microscopic structure of the material. Section 3 describes how we obtain and control the levitation and rotation of the disk, up to an angular speed of about 5000 rpm. In Section 4 we describe the weight measurement procedure and analyze in detail possible error sources and parasitic effects. Several checks were performed to exclude any influence of spurious factors (Section 5). In Section 6 we give the maximum % shielding values obtained in dependence on the rotation speed of the disk and on the frequency of the applied magnetic field. Finally, Sections 7 and 8 contain a short discussion and our conclusions.

According to public information, a NASA group in Huntsville, Alabama, is now attempting to replicate our experiment. This is a difficult task, especially in view of the sophisticated

technology involved in the construction of the large ceramic disk and in the control of its rotation. We are also aware, through unofficial channels, that other groups are working on similar experiments with smaller disks.

2 Construction of the disk.

The shielding superconducting element was made of dense, bulk, almost single-phase $YBCO$ and had the shape of a toroidal disk with an outer diameter of 275 mm, an inner diameter of 80 mm, and a thickness of 10 mm. The preparation of the 123-compound and fabrication of the disk involved mixing the initial oxides, then calcining the powder at 930⁰ C in air, grinding, pressing the disk at 120 MPa, and sintering it in oxygen at 930⁰ C for 12 hours with slow cooling to room temperature. After that, the disk was put back in the furnace at 600⁰ C, and the upper surface was quickly heated to 1200⁰ C using a planar high-frequency inductor as shown in Figure 1. During this last heating, the gap between the disk and the inductor was chosen precisely so that heating would occur only in the top 2 mm-thick layer of the disk, although the material's high heat conductivity caused some heating below this region. Finally, the disk was slowly cooled down to room temperature in a flow of oxygen and treated mechanically in order to obtain good balance during rotation. A thin (1 mm) foil of magnetic material was attached (without electric contact) to the upper surface of the disk, using hot-melt adhesive, to facilitate rotating the disk as described below, especially at the initial stages of rotation.

The phase and crystal structure of the superconductor were studied using X-ray diffraction analysis (XRD) and a scanning electron microscope (SEM) equipped with an energy dispersive spectral (EDS) analyzer. The samples were cut layer by layer from the bulk ceramic disk.

The analysis of the cross-section of the ceramic $YBa_2Cu_3O_{7-x}$ disk revealed the existence of two zones with different crystal structures. The upper part of the disk (6-7 mm thick) had an orthorhombic structure typical of the quench and melt growth process [5, 6] and consisted mainly of single-phase orthorhombic 123-compound. This material was dense, with uniformly fine grain boundaries, i.e. no impurities or secondary phases were found between the grains. Inter-grain boundaries were barely visible, indicating that there were good electrical contacts between the particles of the superconducting body and that the sintering of the material had

produced a nearly perfect polycrystal lattice with no apparent defects.

The grains were less than $2 \mu m$ wide and were oriented (about 75%) with c-axis parallel to the surface of the disk. The transition temperature T_c for this region of the disk was found by direct measurements to be $94 K$, with a width of $1.5\text{--}2 K$. T_c was determined from the resistive transition in a variable temperature cryostat, under zero magnetic field, using an AC current and sputtered golden contacts.

The lower part of the disk, which was in contact with a water-cooled base during the high-frequency heat treatment, had a markedly different structure: randomly oriented grains, with typical grain sizes between 5 and $15 \mu m$. The porosity of this zone varied from 5 to 9% and the material contained about 40% of the tetragonal phase. The transition temperature T_c was equal to $60 K$, with a width of ca. $10 K$. EDS analysis showed the presence of small inclusions of Y_2BaCuO_5 in the lower layer.

Crystal lattice parameters for these two layers, as calculated from XRD, are listed below. These are dimensions in nm:

Upper layer: $a=0.381$; $b=0.387$; $c=1.165$;

Lower layer: $a=0.384$; $b=0.388$; $c=1.170$ (orthorhombic phase);

$a=0.387$; $c=1.183$ (tetragonal phase).

The critical current density was measured for samples cut from the top of the superconducting disk. Measurements of J_c were carried out at $75 K$ using an AC current, four-probe method, and direct transport measurements. The accuracy for J_c determination was defined as $1 \mu V/cm$ in a zero magnetic field, with the sample immersed in liquid nitrogen. It turned out that J_c exceeded $15000 A/cm^2$. The value of J_C for the lower part of the disk was not measured, since it is not superconducting at the temperature of operation.

We also estimated the current density in the upper part of the disk while subjected to the magnetic fields usually applied during the measurements. To measure this we made a thin radial cut through the sample disk and attached electric contacts to an ampermeter, with a technique allowing fast on/off switching. We calibrated the currents in the driving coils that correspond to the currents inside the disk. These currents are slightly different for each new disk as the thickness of the SC part is not the same in every new sintered ceramic toroid, but we estimate their density to range between 5000 and $7000 A/cm^2$.

3 Operation of the apparatus.

Two identical solenoids were placed around the superconductor using fibreglass supports, as shown in Figures 2, 4, 5. The gaps between these solenoids and the disk were large enough for it to move about 20 mm in any lateral direction. The toroidal disk was placed inside a cryostat equipped with a set of three coils (Fig. 3) that could keep it levitating when it reached the superconducting state. The angle β was between 5 and 15 degrees. This helped to keep the rotating disk in a stable position, otherwise it tended to slip aside.

A schematic of the electrical connections is shown in Fig. 6. High-frequency electric current (10^5 Hz) was first sent to the two main solenoids around the disk to initiate an internal current in the ceramics while the disk was still at room temperature. Then the system was slowly cooled down to 100 K by liquid nitrogen, and then quickly cooled by liquid helium vapors. We estimate the temperature of the disk to be lower than 70 K in stationary conditions. Thus the upper layer of the disk is superconducting in these conditions, while the lower layer is not. The main solenoids were switched off. After this, the high-frequency current was sent to the coils below the disk, and the superconductor raised up (at least 15 mm; see Section 6).

Then a small current (10^5 Hz) was sent to the main solenoids, causing the disk to begin rotating counter-clockwise with increasing speed. The rotation speed was increased up to 5000 rpm. At this point the current in the rotating coils was of the order of 8-10 A. (The diameter of the wire of these coils is 1.2 mm). This current was supplied by powerful high-frequency generators usually employed for induction heating and quenching of metals.

Most weight measurements for various objects were taken in these conditions, which can be maintained in a stable way for quite long periods (10 minutes or more). Next, the disk's rotational speed was slowly reduced by changing the current in the main solenoids (Fig. 9). The speed of rotation was regulated by means of laser beam reflections off a small piece of plastic light-reflecting foil attached to the disk.

The frequency of the e.m. field was varied from 10^3 to 10^8 Hz . Samples made of various materials were tested, including metals, glass, plastic, wood and so on. All the samples were hung over the cryostat on a cotton thread connected to a sensitive balance. The distance between the samples and the cryostat varied from 25 to 3000 mm.

4 Weight measurements. Error sources.

To measure the weight loss of the samples, we used a Dupont balance that is a part of the standard equipment for DTA and TGA (differential thermal analysis, thermo-gravimetric analysis). One of the two arms of the balance, holding the sample to be weighed, was lying within the vertical projection of the HTC disk (we call this region the "shielding cylinder"), while the other arm was well outside. The arms of the balance were up to 220 cm long. The sensitivity of the balance for masses of 10-50 g, like those employed in the measurements, is on the order of 10^{-6} g, which was sufficient to detect the observed weight loss. Three different balances were used for verification and are described below (see "Checks", Section 5).

The main error sources in the weight measurements were the following.

1. *Buoyancy and air flow.* The presence of the cryostat perturbs the air and causes weak local flows. Moreover, a much larger ascending flux is caused by the weight loss of the air in all the shielding cylinder; this also produces a pressure drop in the shielding cylinder (see Section 6). The error introduced by this effect in the weight measurements has been substantially reduced by enclosing the samples to be weighed in a long glass tube, closed at the bottom. The samples had the form of elongated objects (like big pencils). In order to set an upper limit on the buoyancy effect, some flat samples were weighed too, without enclosing them in the glass tube, in vertical and horizontal positions, and the results for the two cases were compared. It turned out that the weight loss in horizontal position was larger (by approx. 10% of the total loss) than in vertical position.
2. *Hydrostatic force.* This introduces a slight dependence of the weight loss on the density of the sample. For a material with density 1 Kg/dm^3 , the Archimede pull amounts to about the 0.1% of the weight. Since the air density is slightly lower in the shielding cylinder (see above and Section 6), the Archimede pull is lower itself, but this effect can be disregarded, being of the order of 0.001% of the total weight or less.
3. *Diamagnetism.* It is known that molecular diamagnetism produces a small levitating force on samples placed in a magnetic field gradient. For a large class of materials, this force is essentially independent on the chemical composition of the material and is proportional to its weight. For instance, a standard value for the diamagnetic levitating force is the

following: in a field gradient of 0.17 T/cm the force exerted on a 1 g sample of NaCl , SiO_2 , S or diamond is ca. equal to 16 dyne (about 0.016 g , or 1.6% of the weight).

Since the force is proportional to the square of the field gradient, one easily finds that the value of the field gradient corresponding to a percentage weight loss of 0.1% is about 0.04 T/cm (for comparison, we recall that the maximum weight loss we observed in stationary conditions was 0.5%). A mapping of the static field produced by our apparatus yielded in all cases smaller values of the field gradient near the disk, and much smaller values at a height of 50 cm or more above the disk, where the observed weight loss stays the same (compare Section 6). We thus conclude that the effect of molecular diamagnetism can be completely disregarded.

4. *RF fields.* The possibility of a slight levitating effect from an RF magnetic field cannot be excluded completely. However, such disturbance was attenuated, if not eliminated, by placing thick metal screens between the cryostat and the samples. Copper, aluminum, and steel screens were tested separately and in many different combinations. The individual screens had a diameter of 300 mm and a thickness of 50 mm . The presence of the screens never altered the effect.

5 Checks.

We did several checks in order to correlate more clearly the appearance of the effect to specific experimental conditions.

1. *Substituting a metal disk or a disk made totally from superconducting ceramic.* In all these cases, the shielding effect was not observed. This confirms, in our opinion, that the origin of the effect resides in the interaction between the upper (superconducting) part of the disk and the lower part, where considerable resistive phenomena take place.
2. *Measurements in vacuum, or in different gases, or in a fluid.* These measurements show that the effect exists in these conditions, too, and has the same magnitude as in air. For these cases, however, we cannot furnish data as precise as the measurements taken in air yet, because the experimental conditions are more difficult and the necessary statistics

are still being accumulated. Measurements in vacuum or in gases other than air (N_2 , Ar) are hampered by the fact that the samples and the analytical balance must be enclosed in a sealed container. For the measurements in fluids (H_2O , C_2H_5OH) Archimede pull is more relevant and thus complicates measurements.

3. *An AC field is indispensable.* The shielding effect was completely absent when only static magnetic fields were employed.
4. *Other weighing techniques.* Although the most accurate measurements have been taken with the Dupont balance for precision differential thermogravimetric analysis, we employed for verification three other balances. In the latest runs, using heavier samples with weights varying from 100 g to 250 g, a standard analytical balance was used. Given these masses and the balance's accuracy (0.01 g), weight losses of 0.01% were easy to detect. In addition, we used two other types of balances, sketched in Fig. 8. The first one is a torsion balance, whose oscillation period depends on the tension of the wire and thus on the weight of a suspended sample. This period can be measured with high accuracy through a laser beam reflected by a mirror attached to the wire. Finally, we employed a spring balance whose movements are detected by an induction transducer.

6 Results.

The levitating disk revealed a clearly measurable shielding effect against the gravitational force even without rotation. In this situation, the weight-loss values for various samples ranged from 0.05 to 0.07%. As soon as the main solenoids were switched on and the disk began to rotate in the vapors of liquid helium, the shielding effect increased, and at 5000 rpm, the air over the cryostat began to rise slowly toward the ceiling. Particles of dust and smoke in the air made the effect clearly visible. The boundaries of the flow could be seen clearly and corresponded exactly to the shape of the toroid.

The weight of various samples decreased no matter what they were made from. Samples made from the same material and of comparable size, but with different masses, lost the same fraction of their weight. The best measurement gave a weight loss of 0.5% while the disk was spinning at 5000 rpm, with typical values ranging from 0.3 to 0.5%. Samples placed above the

inner edge of the toroid (5-7 *mm* from the edge) were least affected, losing only 0.1 to 0.25% of their weight. The external boundary of the shielding cylinder was quite clear (no more than 2 *cm*). The maximum values of weight loss were obtained when the levitation height of the disk was at its maximum value, about 30-35 *mm* over the magnets. This condition cannot be reached above 70 *K*, although the disk had become superconducting already at 94 *K*.

During the time when the rotation speed was being decreased from 5000 to 3500 *rpm*, using the solenoids as braking tools, the shielding effect reached maximum values: the weight loss of the samples was from 1.9 to 2.1%, depending on the position of the sample with respect to the outer edge of the disk. These peak values were measured during a 25-30 seconds interval, when the rotational speed was decreasing to 3300 *rpm*. Because of considerable disk vibration at 3000-3300 *rpm*, the disk had to be rapidly braked in order to avoid unbalanced rotation, and further weight measurements could not be carried out.

The samples' maximum weight loss was observed only when the magnetic field was operating at high frequencies, on the order of 3.2 to 3.8 *MHz*. The following tables show how the shielding effect varied in response to changes in the disk's rotation speed or the current frequency.

At constant frequency of 2 MHz:

Rotation speed (*rpm*) Weight loss (%)

4000	0.17
4200	0.19
4400	0.20
4600	0.21
4800	0.22
5000	0.23

At constant rotation speed of 4300 rpm:

Frequency (*MHz*) Weight loss (%)

3.1	0.22
3.2	0.23
3.3	0.24
3.4	0.26
3.5	0.29
3.6	0.32

Remarkably, the effective weight loss was the same even when the samples, together with the balance, were moved upward to a distance of 3 *m*, but still within the vertical projection of the toroidal disk. No weight loss at all was observed below the cryostat.

We also observed a slight diminution of the air pressure inside the shielding cylinder. The difference between the external and the internal pressure was up to 5 *mm* of mercury in stationary conditions (disk rotating at 4000-5000 *rpm*) and increased up to 8 *mm* of mercury during the "braking" phase. Pressures were measured through a vacuum chamber barometer. We believe that the diminution of the pressure inside the shielding cylinder is originated by the pseudo-convective motion of the air, which being lighter tends to raise. This phenomenon is favoured by the fact that an entire cylinder of air get lighter at the same time and thus the ascending motion is amplified.

7 Discussion.

The interaction of a superconducting ceramic body with the gravitational field is a complicated process and cannot be characterized by one single law or physical phenomenon. Also, a comprehensive explanation of the mechanism responsible for high-temperature superconductivity has not yet been found. Still, these facts do not make the observed phenomenon less interesting.

In our previous work [4] the weight loss of samples over the levitating superconductor was smaller, varying from 0.05 to 0.3%. At that time it was difficult to exclude entirely any influence from the radio-frequency field because the sample was separated from the disk and the magnets by a thin plastic film. Now, the superconductor is situated in a stainless steel cryostat, so this influence should be negligible (see also Section 5).

The modification of the superconductor's crystalline structure produced a composite body with a dense and highly oriented upper layer and a porous lower layer with random grain orientation. The upper layer is superconducting at the operation temperature and is able to carry high J_c current under considerable magnetic field, while the lower layer cannot conduct high currents. The boundary between the two layers is likely to constitute a "transition" region in which the supercurrents, that are completely free to move in the upper layer, begin to feel some resistance.

It is also expected that a complex interaction between the composite ceramic body and the external magnetic field takes place. This interaction depends on the coherence length, the flux pinning, the field frequency and the field force, the penetration depth, and the parameters of the crystal lattice. These characteristics are interrelated in a complex way. According to the experimental data (compare also [10], where only a static field was applied), a levitating superconductor does not reveal any unusual shielding if it has no contact with the high frequency AC magnetic field.

As analyzed in [7, 8], pinning centers with different origins may exist inside the superconducting disk, and fluxes will be trapped at some of them. Fluxes trapped at weak centers will begin to move first, while those trapped at strong centers will not move until the Lorentz force exceeds the pinning force. The overall current will be composed of the superposition of flux motions with different speeds.

There are no grounds to claim that the rotation momentum of the disk interacts with

gravitation force, but it seems that fast rotation is favorable for stabilization of the shielding effect.

Finally, it is worth noting that the experimental equipment described above has much in common with magneto-hydro-dynamic (MHD) generators.

The first attempt at a theoretical explanation of the effect has been offered by G. Modanese [11, 12]. Further investigations now in progress may help to prove, change, or complete the present understanding of the observed phenomenon.

8 Conclusions.

A levitating superconducting ceramic disk of $YBa_2Cu_3O_{7-x}$ with composite structure demonstrated a stable and clearly measurable weak shielding effect against gravitational force, but only below 70 K and under high-frequency e.m. field. The combination of a high-frequency current inside the rotating toroidal disk and an external high-frequency magnetic field, together with electronic pairing state and superconducting crystal lattice structure, apparently changed the interaction of the solid body with the gravitational field. This resulted in the ability of the superconductor to attenuate the energy of the gravitational force and yielded a weight loss in various samples of as much as 2.1%.

Samples made of metals, plastic, ceramic, wood, etc. were situated over the disk, and their weight was measured with high precision. All the samples showed the same partial loss of weight, no matter what material they were made of. Obtaining the maximum weight loss required that the samples be oriented with their flat surface parallel to the surface of the disk. The overall maximum shielding effect (2.1%) was obtained when the disk's rotational speed and the corresponding centrifugal force were slightly decreased by the magnetic field.

It was found that the shielding effect depended on the temperature, the rotation speed, the frequency and the intensity of the magnetic field. At present it seems early to discuss the mechanisms or to offer a detailed analysis of the observed phenomenon, as further investigation is necessary. The experimentally obtained shielding values may eventually prove to have fundamental importance for technological applications as well as scientific study.

8.1 Acknowledgments.

The author is grateful to the Institute for High Temperatures at the Russian Academy of Sciences for their help in preparing the unique superconducting ceramic disks and for being permitted to use their laboratory equipment. The effect was first observed and studied at Tampere University of Technology. The author would like to thank G. Modanese, F. Pavese and O. Port for advice and support.

References


- [1] Riise A.B., Johansen T.H., Bratsberg H. and Yang Z.J., *Appl. Phys. Lett.* **60** (1992) 2294.
- [2] Brand E.H., *Am. J. Phys.* **58** (1990) 43.
- [3] Lofland S., Huang M.X. and Bhagat S.M., *Physica C* **203** (1992) 271.
- [4] Podkletnov E. and Nieminen R., *Physica C* **203** (1992) 441.
- [5] Murakami M., Morita M., Doi K., Miyamoto K. and Hamada H., *Jpn. J. Appl. Phys.* **28** (1989) 399.
- [6] Murakami M., Morita M. and Koyama N., *Jpn. J. Appl. Phys.* **28** (1989) 1125.
- [7] Takizawa T., Kanbara K., Morita M. and Hashimoto M., *Jpn. J. Appl. Phys.* **32** (1993) 774.
- [8] Xu J.H., Miller J.H. Jr. and C.S. Ting, *Phys. Rev. B* **51** (1995) 424.
- [9] Bull M., De Podesta M., *Physica C* **253** (1995) 199.
- [10] Unnikrishnan, C.S., *Physica C* **266** (1996) 133.
- [11] Modanese G., *Europhys. Lett.* **35** (1996) 413.
- [12] Modanese G., *Phys. Rev. D* **54** (1996) 5002; Modanese G. and Schnurer J., *Possible quantum gravity effects in a charged Bose condensate under variable e.m. field*, report UTF-391/96, November 1996, Los Alamos database nr. gr-qc/9612022.

FIGURE CAPTIONS.

1. Schematic cross-section of the furnace for high-temperature treatment of the ceramic disk with planar high-frequency inductors.
2. General magnets and cryostat setup.
3. Typical geometry and position of the disk over supporting solenoids.
4. Schematic design of rotating solenoids. a, b: various configurations.
5. Typical configuration of the tested set-up for the rotating solenoids.
6. Block scheme of the electrical connections.
7. Schematic design of the cryogenic system for the refrigeration of the superconducting disk.
8. General configuration of the equipment for the weight loss measurements.
9. Typical design of the three-point disk-braking system.

High T_c materials, fabrication

|general, fig. 21/3|



Created using Super. Tech. Top Draw 1/25/97 7:20:55 PM

Fig. 1





FIG.2

Created using Super. Tech. Top Draw 1/25/97 7:31:36 PM

Supporting solenoids

|general, fig. 21/5|




Fig. 3

Created using Super. Tech. Top Draw 1/25/97 7:33:14 PM

Rotating solenoids

|general, fig. 31/3|






Fig. 4
Created using Super. Tech. Top Draw 1/25/97 7:34:00 PM

Rotating solenoids

|general, fig. 31/4|




Fig. 5

Created using Super. Tech. Top Draw 1/25/97 7:34:52 PM

Electric circuits

|general, fig. 41/3|




Fig. 6

Created using Super. Tech. Top Draw 1/25/97 7:35:28 PM

Cryogenic system

|general, fig. 51/5|




Fig. 7

Created using Super. Tech. Top Draw 1/25/97 7:36:13 PM

Weight & pressure measurements

|general, fig. 61/4|




Fig. 8

Created using Super. Tech. Top Draw 1/25/97 7:36:43 PM

Disk braking system

|general, fig. 71/3|





Fig. 9

Created using Super. Tech. Top Draw 1/25/97 7:37:17 PM

MPI-PhT/95-44
May 1995
hep-th/9505094

Theoretical Analysis of a Reported Weak Gravitational Shielding Effect.

Giovanni Modanese*

Max-Planck-Institut für Physik

Werner-Heisenberg-Institut

Föhringer Ring 6, D 80805 München (Germany)

To appear in: Europhys. Lett.

Abstract

Under special conditions (Meissner-effect levitation in a high frequency magnetic field and rapid rotation) a disk of high- T_c superconducting material has recently been found to produce a weak shielding of the gravitational field. We show that this phenomenon has no explanation in the standard gravity theories, except possibly in the non-perturbative Euclidean quantum theory.

04.20.-q Classical general relativity.

04.60.-m Quantum gravity.

74.72.-h High- T_c cuprates.

*A. Von Humboldt Fellow.

e-mail: modanese@science.unitn.it

In two recent experiments [1, 2], Podkletnov and co-workers have found indications for a possible weak shielding of the gravitational force through a disk of high- T_c superconducting material. In the first experiment a sample made of silicon dioxide of the weight of ca. 5 g, was found to lose about 0.05% of its weight when placed 15 mm above the disk. The diameter of the disk was 145 mm and its thickness 6 mm. The disk was refrigerated using liquid helium and was levitating over a solenoid due to the Meissner effect. When the disk was set in rotation by means of lateral alternating magnetic fields, the shielding effect increased up to 0.3%. When the disk was not levitating, but was placed over a fixed support, no shielding was observed.

In the second experiment the disk had the form of a toroid with the outer diameter of 275 mm and was enclosed in a stainless steel cryostat. Samples of different composition and weight (10 to 50 g) were placed over the disk and the same percentual weight loss was observed for different samples, thus enforcing the interpretation of the effect as a slight diminution of the gravitational acceleration. While the toroid was rotating (at an angular speed of 5000 rpm) the weight loss was of 0.3-0.5%, like in the first experiment, but it reached a maximum of 1.9-2.1% when the speed was slowly reduced by varying the current in the solenoids.

In both experiments, the magnetic fields were produced by high frequency currents and the maximum effect was observed at frequencies of the order of ca. 1 MHz. Measurements were effected also in the vacuum, in order to rule out possible buoyancy effects. The dependence of the shielding value on the height above the disk was very weak. Within the considered range (from a few cm to 300 cm) no sensible variation of the shielding value was observed. This weak height dependence is a severe challenge for any candidate theoretical interpretation, as it violates an intuitive vectorial representation of the shielding. We have analyzed in detail this issue in [3].

Independent repetitions of the experiment have already been undertaken, stimulated by scientific and especially technological interest. We would like to stress here the importance of precise measurements. In particular, it is essential to obtain exact spatial field maps and information about the transient stages. It is also crucial to use a different kind of balance from that used by the authors of [1, 2] and possibly a gravity gradiometer [4]. If the effect turned out to be of non-gravitational nature, its fundamental interest would be strongly reduced. On the contrary, if the effect is really a gravitational shielding its theoretical explanation calls for new and non-trivial dynamical mechanisms, as we argue in the following.

For clarity we shall organize our analysis as follows. Considering two masses m_1 and m_2 (which represent the Earth and the sample) and a medium between them (the disk), we shall

evaluate their potential energy in these alternatives:

- (1) the *medium* can be regarded as a classical system, as a quantum system, or as a Bose condensate with macroscopic wave function;
- (2) the *gravitational field* can be regarded as classical or quantized, and in both cases as weak (perturbations theory) or strong.

In which of these approximations could a shielding effect arise? We can immediately observe that the possibilities of Point (1) are severely restricted by a large body of experimental evidence. Namely a sensible gravitational shielding has never previously been observed. Several experiments, starting with the classical measurements of Q. Majorana, have shown that the gravitational force is not influenced by any medium, up to one part in 10^{10} or less (for a very complete list of references see [5]). For a “classical” medium the reason for this is essentially the absence of charges of opposite sign which, by shifting or migrating inside the medium, might generate a field which counteracts the applied field. On the other hand when the medium is regarded as a quantum system, it is easy to check that the probability of a (virtual) process in which a graviton excites an atom or a molecule of the medium and is absorbed is exceedingly small, essentially due to the smallness of the gravitational coupling at the atomic level (see for instance [6]). It is then clear that the reported shielding can only be due to the Bose condensate present in the high- T_c superconductor.

Coming to Point (2), first we regard the gravitational field as classical. It is readily realized that in general neither the superconducting disk nor any other object of reasonable density, if placed close to the sample mass, can influence the local geometry so much as to modify its weight by the observed amount. To check this one just needs to write the Einstein equations (or even some generalizations of them) and impose suitable conditions on the source $T_{\mu\nu}$:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -8\pi GT_{\mu\nu}; \quad (1)$$

$G \sim 10^{-66} \text{ cm}^2$ in natural units;

$$|T_{\mu\nu}| < \dots$$

Let us be more explicit: according to Einstein equations any apparatus with mass-energy comparable to that of Ref.s [1, 2], if placed far away from any other source of gravitation, is unable to produce a gravitational field of the intensity of ca. 0.01 g [†]. The shielding effect, if

[†]When examining this possibility one should hypothesize that the disk produces a repulsive force. But it is known that arguments in favour of ”antigravity” are untenable [7] and that local negative energy densities

true, must then consist of some kind of "absorbtion" of the Earth's field in the superconducting disk.

Having thus excluded any possibility of shielding for a classical gravitational field, we need now an expression for the gravitational potential energy of two masses m_1 and m_2 which takes into account quantum field effects, possibly also at non-perturbative level. This is given in Euclidean quantum gravity by the functional integral [13]

$$E = \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{\int d[g] \exp \left\{ -\hbar^{-1} \left[S_g + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{g_{\mu\nu}[x_i(t)] \dot{x}_i^\mu(t) \dot{x}_i^\nu(t)} \right] \right\}}{\int d[g] \exp \left\{ -\hbar^{-1} S_g \right\}} \quad (2)$$

$$\equiv \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \left\langle \exp \left\{ -\hbar^{-1} \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} ds_i \right\} \right\rangle_{S_g}. \quad (3)$$

where S_g is the gravitational action

$$S_g = \int d^4x \sqrt{g} \left(\frac{\Lambda}{8\pi G} - \frac{R}{8\pi G} + \frac{1}{4} a R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right). \quad (4)$$

The R^2 term in S_g (important only at very small scale) is necessary to ensure the positivity of the action. The trajectories $x_i(t)$ of m_1 and m_2 are parallel with respect to the metric g ; let L be the distance between them, corresponding to the spatial distance of the two masses. An evaluation of (3) in the non-perturbative lattice theory has been carried out recently by Hamber and Williams [14].

In perturbation theory [15] the metric $g_{\mu\nu}(x)$ is expanded in the traditional way as the sum of a flat background $\delta_{\mu\nu}$ plus small fluctuations $\kappa h_{\mu\nu}(x)$ ($\kappa = \sqrt{8\pi G}$). The cosmological and the R^2 terms are dropped, leaving the pure Einstein action. Eq. (2) is rewritten as

$$E = \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{\int d[h] \exp \left\{ -\hbar^{-1} \left[S_{\text{Einst.}} + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{1 + h_{00}[x_i(t)]} \right] \right\}}{\int d[h] \exp \left\{ -\hbar^{-1} S_{\text{Einst.}} \right\}}, \quad (5)$$

where the trajectories $x_1(t)$ and $x_2(t)$ are two parallel lines in flat space. Expanding (5) in powers of κ one obtains to lowest order the Newton potential [13], and to higher orders its relativistic and quantum corrections [16].

are strongly constrained in Quantum Field Theory (see for instance [8]). There remains only the possibility of gravitomagnetic and gravitoelectric effects, which are however usually very small [9]. In [10] it is argued using the Maxwell-like approximated form of Einstein equations that the gravitoelectric field produced by a superconductor could be abnormally strong. In our opinion this conclusion contrasts with the full equations (2). For a comparison, consider the strength of the "gravitational Meissner effect": in a neutron star with density of about 10^{17} kg/m^3 the gravitational London penetration depth is ca. 12 km [11]. An experimental check disproving the hypothesis of a repulsive force is the measurement of g below the disk. Preliminary measurements [12] do not show any variation of g .

The Bose condensate composed by the Cooper pairs inside the superconductor is described by a bosonic field ϕ with non-vanishing vacuum expectation value $\phi_0 = \langle 0|\phi|0\rangle$. Using the notation $\phi = \phi_0 + \tilde{\phi}$, the action of such a field coupled to the gravitational field has the form

$$S_\phi = \int d^4x \sqrt{g(x)} \left\{ \partial_\mu [\phi_0(x) + \tilde{\phi}(x)]^* \partial_\nu [\phi_0(x) + \tilde{\phi}(x)] g^{\mu\nu}(x) \right. \\ \left. + \frac{1}{2}m^2 |\phi_0(x)|^2 + \frac{1}{2}m^2 [\phi_0^*(x)\tilde{\phi}(x) + \phi_0(x)\tilde{\phi}^*(x)] + \frac{1}{2}m^2 |\tilde{\phi}(x)|^2 \right\}, \quad (6)$$

where m is the mass of a Cooper pair. In order to describe the interaction we insert S_ϕ into (2) and include $\tilde{\phi}$ into the integration variables, while ϕ_0 is considered as an external source, being determined essentially by the structure of the superconductor and by the external e.m. fields. In the following we shall disregard in S_ϕ the terms containing $\tilde{\phi}$, as they give rise to emission-absorption processes of gravitons which we know to be irrelevant.

Perturbatively, the interaction of $h_{\mu\nu}(x)$ with the condensate $\phi_0(x)$ is principally mediated by the vertex

$$\mathcal{L}_{h\phi_0} = \kappa \partial_\mu \phi_0^*(x) \partial_\nu \phi_0(x) h^{\mu\nu}(x). \quad (7)$$

This produces corrections to the gravitational propagator, which are however practically irrelevant, because they are proportional to powers of $\kappa \sim 10^{-33} \text{ cm}$. It is straightforward to compute the corresponding corrections to eq. (5). We do not need to investigate in detail the signs of these corrections or their dependence from ϕ_0 : they are in any case too small (by several magnitude orders) to account for the reported shielding effect.

Looking at the total action $S = S_g + S_\phi$ we recognize besides the familiar vertex (7) a further coupling between $g_{\mu\nu}(x)$ and $\phi_0(x)$. Namely, the Bose condensate contributes to the cosmological term. We can rewrite the total action (without the R^2 term for gravity) as

$$S = S_g + S_\phi = \int d^4x \sqrt{g(x)} \left\{ \left[\frac{\Lambda}{8\pi G} + \frac{1}{2}\mu^2(x) \right] - \frac{R}{8\pi G} \right\} + S_{h\phi_0} + S_{\tilde{\phi}}, \quad (8)$$

where

$$\frac{1}{2}\mu^2(x) = \frac{1}{2} [\partial_\mu \phi_0(x)]^* [\partial^\mu \phi_0(x)] + \frac{1}{2}m^2 |\phi_0(x)|^2, \quad (9)$$

$$S_{h\phi_0} = \int d^4x \sqrt{g(x)} \mathcal{L}_{h\phi_0} \quad (10)$$

and $S_{\tilde{\phi}}$ comprises the terms which contain at least one field $\tilde{\phi}$ and are thus irrelevant, as we mentioned above.

We see from (8) that the condensate $\phi_0(x)$ and its four-dimensional gradient give a positive contribution to the intrinsic cosmological term $\Lambda/8\pi G$. It is known that a positive cosmological

constant turns Einstein gravity into an unstable theory, as it corresponds in the action to a mass term with negative sign [15]. In the case we are considering here, the cosmological term is spacetime dependent. The situation is thus quite complicated and we shall just sketch it briefly in the following; a more complete account can be found in [17].

Since in any four-dimensional region Ω in which $\mu^2 > |\Lambda|/8\pi G$ the mass term of the gravitational field is non-zero and negative, in this region the field can grow without limit, at least classically (suppose to minimize the action (4) by trial functions which are vacuum solutions ($R = 0$), disregarding the R^2 term). In fact there will be some physical cut-off; thus within Ω the gravitational field will be forced to some fixed value, independent of the external conditions. This is similar to what happens in electrostatics in the presence of perfect conductors: the electric field is constrained to be zero within the conductors. In that case the physical origin of the constraint is different (a redistribution of opposite charges); but in both cases the effect on the field propagator turns out to be that of a shielding.

At this point we would need an estimate for both $|\Lambda|/8\pi G$ and μ^2 . It is known that the cosmological constant observed at astronomical scales is very small; a typical upper limit is $|\Lambda|G < 10^{-120}$, which means $|\Lambda|/8\pi G < 10^{12} \text{ cm}^{-4}$. To estimate μ^2 we can assume an average density of Cooper pairs in the superconductor of $\sim 10^{20} \text{ cm}^{-3}$. Remembering that the mass of a pair is $\sim 10^{10} \text{ cm}^{-1}$ in natural units, we find that $\phi_0 \sim 10^5 \text{ cm}^{-1}$ and thus $m^2|\phi_0|^2 \sim 10^{30} \text{ cm}^{-4}$. If ϕ_0 varies over distances of the order of $10^{-8} \div 10^{-7} \text{ cm}$, as usual in high- T_c superconductors, the gradient $(\partial\phi_0/\partial x)$ can be of the order of 10^{12} cm^{-2} .

These values support our hypothesis that the total cosmological term is positive in the superconductor. Actually the positive contribution of the condensate is such that one could expect the formation of gravitational singularities in any superconductor, subjected to external fields or not – a fact which contrasts with the observations. This can be avoided if we take the point of view, supported by lattice quantum gravity with a fundamental length [14, 17, 18], that the effective intrinsic cosmological constant depends on the momentum scale p like

$$|\Lambda|G \sim (\ell_0 p)^\gamma, \quad (11)$$

where $\ell_0 \sim \kappa$ is the fundamental lattice length (Planck length) and γ is a critical exponent which up to now has been computed only for small lattices. The sign of Λ is negative. This provides a well defined flat space limit for the non-perturbative Euclidean theory based on the action (4), and ensures that the signs in our discussion of the minimization of the total action are correct.

We hypothesize that at the length scale of $10^{-8} \div 10^{-6}$ cm Λ could be of the same order of the average of μ^2 , so that the competition between the two terms would lead to singularities only in those regions of the superconductor where the condensate density is larger than elsewhere. From this hypothesis we deduce that $|\Lambda| \sim 10^{-36}$ cm $^{-2}$, a value well compatible with the conventional experimental data at that scale \ddagger .

In conclusion, a shielding effect of the reported magnitude cannot be explained by classical General Relativity, nor by the usual perturbation theory of quantum gravity coupled to the Cooper-pair density $\phi_0(x)$ through the vertex (7). We have then considered a further possible coupling mechanism: the term $\mu^2(x) = [\partial^\mu \phi_0^*(x)][\partial_\nu \phi_0(x)] + m^2 \phi_0^2(x)$ in the condensate's action may act as a positive contribution to the effective gravitational cosmological constant. This may produce localized gravitational instabilities and thus an observable effect, in spite of the smallness of G , which makes the coupling (7) very weak. In the regions where the gravitational field becomes unstable it tends to take "fixed values" independent of the neighboring values. This reminds of the situation of electrostatics in the presence of perfect conductors. The effect on the field propagator and on the static potential is that of a partial shielding.

According to this picture, the magnetic fields which keep the disk levitating and rotating and produce currents inside it play the role (together with the microscopic structure of the HTC material) of determining the condensate density $\phi_0(x)$. The energy necessary for the partial "absorbtion" of the gravitational field is supplied by the high-frequency components of the magnetic field.

The autor would like to thank D. Maison and the staff of MPI for the kind hospitality in Munich, and the A. von Humboldt Foundation for financial support. Useful suggestions of G. Fiore, G. Fontana, F. Pavese, V. Polushkin, D. Saam and R. Stirnimann are acknowledged too.

References

- [1] E. Podkletnov and R. Nieminen, Physica C **203** (1992) 441.

\ddagger It corresponds to a graviton mass of $\sim 10^{-18}$ cm $^{-1}$ and thus to a (unobservable) range of gravity of $\sim 10^{18}$ cm. Note that a similar argument allows to restrict the exponent γ in eq. (11) to $\gamma > 2$. In fact, for $\gamma = 2$ we would have $|\Lambda| \sim p^2$, in contradiction with the fact that in order to be unobservable the range of gravity must be much larger than the length scale p^{-1} which we are considering.

- [2] E. Podkletnov and A.D. Levit, *Gravitational shielding properties of composite bulk $YBa_2Cu_3O_{7-x}$ superconductor below 70 K under electro-magnetic field*, Tampere University of Technology report MSU-chem, January 1995.
- [3] G. Modanese, *Updating the theoretical analysis of the weak gravitational shielding experiment*, report UTF-367/96, January 1996 (supr-con/9601001).
- [4] H.A. Chan and H.J. Paik, Phys. Rev. **D 35** (1987) 3551; H.J. Paik and J.M. Lumley, *Superconducting gravity gradiometers on STEP and GEM*, U. of Maryland and Oxford Instruments report, January 1996.
- [5] G.T. Gillies, Metrologia **24** (Suppl.) (1987) 1.
- [6] S. Weinberg, *Gravitation and cosmology: principles and applications of the general theory of relativity*, J. Wiley, New York, 1972.
- [7] M.M. Nieto, T. Goldman, Phys. Rep. **205** (1991) 221.
- [8] L.H. Ford, Phys. Rev. **D 48** (1993) 776 and references.
- [9] J.M. Nitschke and P.A. Wilmarth, Phys. Rev. Lett. **64** (1990) 2115; J.E. Faller and W.J. Hollander, Phys. Rev. Lett. **64** (1990) 825.
- [10] D.G. Torr and Ning Li, Found. Phys. Lett. **6** (1993) 371.
- [11] R.P. Lano, *Gravitational Meissner effect*, report U.ofIowa 96-4, March 1996 (hep-th/9603077).
- [12] E. Podkletnov, private communication, April 1996.
- [13] G. Modanese, Phys. Lett. **B 325** (1994) 354; Nucl. Phys. **B 434** (1995) 697; Riv. Nuovo Cim. **17**, n. 8 (1994).
- [14] H.W. Hamber and R.M. Williams, Nucl. Phys. **B 435** (1995) 361, and references therein.
- [15] M.J.G. Veltman, *Methods in field theory*, Les Houches Summer School 1975, ed.s R. Balian and J. Zinn-Justin (North-Holland, Amsterdam, 1976); E. Alvarez, Rev. Mod. Phys. **61** (1989) 561.
- [16] I.J. Muzinich, S. Vokos, Phys. Rev. **D 52** (1995) 3472.

- [17] G. Modanese, *Role of a "local" cosmological constant in Euclidean quantum gravity*, report
UTF-368/96, January 1996 (hep-th/9601160).
- [18] G. Modanese, Phys. Lett. **B 348** (1995) 51.

Proposal
for the

Experimental Detection of Gravitomagnetism in the
Terrestrial Laboratory

By

Robert E. Becker
131 Old County Road, #168, Windsor Locks, CT, 06096
(860)627-9116

Paul A. Smith
Dept. of Physics, Coe College, Cedar Rapids, IA 52402

Jeffrey Bertrand
Huntsville, AL

Abstract

A. Introduction

One of the predictions of the General Theory of Relativity, unveiled eighty years ago, is the existence, in a linear approximation, of a gravitational analog of the magnetic field, known as the gravitomagnetic (GM) or prorotational field. In the form of the Lense-Thirring Effect, this aspect of the gravitational field has been studied extensively since then. Indeed, there is a large-scale space mission, Gravity Probe-B, planned for launch near the turn of the millennium, designed to detect the exceedingly weak GM field of the earth.

It is the very weakness of the GM field of the entire earth that makes it so difficult, if not outright impossible, to measure the GM field in a terrestrial laboratory. Indeed, it has required extraordinary engineering ingenuity and precision to design gyroscopes sensitive enough to detect the 10^{-15} Hz precession induced by the earth at the orbit of Gravity Probe-B. Likewise, artificially generating a detectable GM field in the laboratory is clearly hopeless, though one of us (Becker) has elsewhere proposed a variant of a magnetic resonance experiment utilizing scanning probe microscopies and nanotechnologies, to attempt this.

The present experiment proposes to overcome the problem of the weakness of the GM field by taking advantage of the unique properties of macroscopically coherent systems

such as superconductors (SC). It is by now well known that the magnetic flux threading a superconductor is quantized with a quantum number n . We will show that the GM field enters superconducting flux relations in the same manner as the magnetic field, and that the GM flux should also be quantized. Furthermore, if the value of magnetic flux quantum number is fixed for a given physical configuration, that same quantum number value governs the GM flux quantum as well. And this GM flux quantum, for sufficiently high n , should be macroscopically detectable. This experiment is designed to detect the GM field represented by this quantized GM flux.

One obvious question that must be asked is "If the GM field of even a body as massive as the earth is so tiny, how is it conceivable that a small SC can generate a field large enough to be readily measurable in the laboratory?" To address this, first note that in the presence of an external GM field, there is an additional GM term in the usual quantized magnetic fluxoid relation. This GM contribution to the magnetic flux is correctly taken to be negligible because the external GM field is so tiny.

Similarly, there is a magnetic term in the quantized GM flux relation. For substantial magnetic fields, such as those that would result in a high magnetic flux quantum n , this term is not small. Thus, a large GM flux quantum may be detected due to a magnetically induced GM field in the SC. The exact physical mechanism responsible for this "amplification" is still nebulous. One of the objectives of this experiment is to help elucidate such a mechanism. One of us (Becker, M.S. Thesis, Un. Alabama, Huntsville, 1994) has taken initial steps in identifying possible theoretical mechanisms.

It should be emphasized that a GM field large enough to be detectable in a terrestrial laboratory is predicted to occur only in macroscopic coherent systems such as superconductors or superfluids. (Though the effect should also occur in superfluids with a toroidal topology, the supercarriers in superfluids are much more massive than the electrons in SC, leading to a much smaller GM flux quantum than can be found in a SC.)

Ultimately, the effect to be studied, as with so many other unusual phenomenon, owes its postulated existence to the primacy of quantum mechanics, and the even more special properties of macroscopic, coherent systems.

Non-detection of the effect would actually raise equally important theoretical questions, such as: "If the Maxwell approximation to general relativity is correct, why is there not a GM flux quantum just as there is for the magnetic field?" After all, both vector potentials appear "symmetrically" in the quantum mechanical canonical momentum. "Could the U(1) gauge field and topological description of the GM field in the Maxwell approximation be wrong, even if the Maxwell approximation to the Einstein equations is correct? What is the fundamental topological relationship between magnetism, GM, and rotation?"

B. Theoretical Background

B.1 Gravitomagnetism Background

Gravitomagnetism (also known as the prorotational field) is the gravitational analog of magnetism. The gravitational analog of the electric field is none other than our familiar Newtonian gravitational field, E_G . GM, on the other hand, has never been detected because it is so small. It is the result of general relativistic effects (GR) related to the curvature of space-time, and even the GM field of the entire earth is unimaginably weak. It can be visualized as the "dragging" of space-time by the rotating.

Despite the difficulties, there have been numerous experiment proposals for detection of GM over the last 30 years. The first one is still ongoing, Gravity Probe B, a spacecraft which will use an extraordinarily accurate and sensitive SC gyroscope to attempt to detect the GM field of the earth. Others exploit the properties of SC or superfluids (SF) to try to detect the earth's or laboratory generated fields. These have not, to the best of our knowledge, been carried out, because the predicted effects are still too minute to detect with current technology.

Though GM is not quite a new force of nature, it is nonetheless a new aspect of gravitation, and its detection would be regarded as extremely important to science, particularly if it can be done in a terrestrial laboratory.

An understanding of the expected physical effects of GM can best be gained by first revisiting electromagnetism. The ordinary electromagnetic (EM) magnetic flux density B (we will see why \mathbf{B} , also called the magnetic inductance,

is called the flux density; a related quantity, \mathbf{H} , is called the magnetic field, but for our purposes will be regarded as one and the same) can be defined by its torque on a magnetic moment:

$$(1) \quad \mathbf{N} = \mathbf{M} \times \mathbf{B}$$

where \mathbf{N} is the torque and \mathbf{M} the magnetic moment. Since torque is defined to be the time rate of change of angular momentum \mathbf{L} , and since \mathbf{M} is related \mathbf{L} in many classical systems by

$$(2) \quad \mathbf{M} = (q/2mc)\mathbf{L}$$

where q is the charge and m the mass of the particle, we can rewrite (1) as

$$(3) \quad d\mathbf{L}/dt = (q/2mc)\mathbf{L} \times \mathbf{B} = \mathbf{L} \times (q/2mc)\mathbf{B}.$$

It is well known that (3) represents the precession of the angular momentum \mathbf{L} about the direction of \mathbf{B} at a Larmor frequency of

$$(4) \quad \Omega = -qB/2mc.$$

Just as a spinning magnetic moment generates a magnetic field, solutions of the GR Einstein equations show that a spinning mass generates a dipolar GM field \mathbf{B}_g . In this case, the GM equivalent of the magnetic moment is the spin angular momentum \mathbf{s} , so that the GM torque on a spinning body can be written as

$$(5) \quad \mathbf{N} = (1/2c)\mathbf{s} \times \mathbf{B}_g.$$

Comparing (5) with (3) we conclude

$$(6) \quad d\mathbf{s}/dt = (1/2c)\mathbf{s} \times \mathbf{B}_g = \mathbf{s} \times (1/2c)\mathbf{B}_g$$

and thus there is a GM precessional frequency of

$$(7) \quad \Omega_g = -B_g/2c.$$

It is Ω_g which we will attempt to detect in our experiment. Unlike mechanical quantities, but like EM, both the dimensions of, and the equations for, \mathbf{B}_g depend on the unit system chosen. In cgs, which is what will be used here, \mathbf{B}_g has units of acceleration. In mks, it actually has units of frequency. Given this discordance, it will be best to translate final answers from one

system of units to another rather than expressing the basic equations in several systems.

This frequency is a function only of the source field \mathbf{B}_g , and not of the nature of the test particle. The nature of the source field outside a spinning mass in a dipole approximation is such that

$$(8) \quad \mathbf{B}_g = GS/2c^2r^3,$$

where S is the spin angular momentum of the source body. This is similar to the dipole magnetic field outside a magnetized body.

Even for the earth, (8) is extremely small, on the order of 10^{-15} Hz at the surface.

Another similarity between EM and GM is in the Maxwell equations. In EM we can express \mathbf{B} in terms of a magnetic vector potential

$$(9) \quad \mathbf{B} = \mathbf{curl}\mathbf{A}.$$

Similarly,

$$(10) \quad \mathbf{B}_g = \mathbf{curl}\mathbf{A}_g.$$

A Lorentz force equation also holds:

$$(11) \quad d\mathbf{v}/dt = \mathbf{E}_g + (\mathbf{v}/c) \times \mathbf{B}_g.$$

A mass current generates a GM field just like a charged current generates a \mathbf{B} field. One can also write down GM Maxwell equations, but we will not need them here.

B.2 Superconductors

SC have many curious characteristics. They are perfect conductors, so that once a current is started for any reason, it will continue. But they also obey the Meissner effect by which a magnetic field is excluded from the interior of a bulk SC, even in an applied \mathbf{B} field. Physically, this comes about because the \mathbf{B} field induces surface (not bulk) currents, which set up a field opposing the applied field in the interior; the net field is $B=0$. These currents only flow in a layer, the London penetration depth, in which \mathbf{B} and the current \mathbf{J} fall-off exponentially with distance into the SC.

All the strange properties of SC ultimately derive from the quantum mechanical (QM) state of the system. Below a certain critical temperature, it becomes energetically favorable for conduction electrons in the material to form pairs such that their center of mass momentum is zero; these are Cooper pairs. At absolute zero, all the electrons in the SC form Cooper pairs, and so condense into the same QM state of zero momentum.

Since electrons have QM spin of 1/2, and since opposite spin electrons are paired, a Cooper pair is QM object of integral spin, called a boson. Unlike fermions, which obey the Pauli Exclusion Principle, any number of bosons can occupy the same QM state. So below the critical temperature, T_c , the SC can be said to be in a macroscopically occupied QM state. In order to change the state, a physical phenomena would have to break all the Cooper pairs and remove the electrons from this state, known as a condensate. This requires much more energy than just breaking a single Cooper pair, and so changing QM states becomes exceedingly unlikely.

This explains properties like persistent currents. They persist, because once the QM state is set-up, it can't easily be changed. There are complications to the simple picture just described. For example, persistent London screening currents must be generated by an applied field, and then the Cooper pairs do not have zero center of mass momentum. But they do have zero QM canonical momentum, which is the correct vanishing quantity in this case.

At $T > 0$, not all electrons form Cooper pairs, but the ground state is still macroscopically occupied. For our purposes, this will suffice. Another complication is the existence of several types of SC. The first to be discovered was Type I SC, which obey the Meissner effect for all B and all $T < T_c$. Next came Type II SC, which obey the Meissner effect only for $B < B_{c1}$, the Lower Critical Field. Between B_{c1} and a B_{c2} (Upper Critical Field), an applied magnetic field can penetrate the SC and form current vortices around the field lines. Finally, there are the famous high- T_c SC, which we will actually use. High- T_c SC do not obey the same microscopic QM theory as do standard Type I and II SC (s-wave BCS theory), but for our purposes can be regarded as Type II SC: they have a B_{c1} . We shall actually be interested in the Meissner regime of the Type II SC.

In the Type II regime, the critical property is that the magnetic field can penetrate in a vortex. The vortex core

is not SC, and is composed of normal material, that is, electrons not paired into Cooper pairs. Topologically, this has enormous importance, because it means the SC material is no longer simply-connected; it now has a hole in it. The same thing can happen to any SC if one drills a hole into an otherwise bulk SC, creating a ring or doughnut shaped object.

Whereas screening, diamagnetic, currents are always set-up around the outer periphery of a SC to cancel an applied field in the interior, if a hole is present, another current, called the paramagnetic current, is set-up around the inner circumference of the hole. The paramagnetic current ensures that once a SC is cooled below T_c in an applied field, the magnetic flux penetrating the hole remains constant even if the applied field is removed. And this flux through the hole obeys a quantization condition. It is this condition, when amended by GM, which our experiment will investigate.

B.3 SC Fluxoid Quantization

To see where these conditions come from, we can delve into QM briefly. A Cooper pair moving in an external magnetic field of vector potential \mathbf{A} has a momentum given not by the ordinary momentum $m\mathbf{v}$, but by the canonical momentum

$$(12) \quad \mathbf{P} = 2m\mathbf{v} + 2q\mathbf{A}.$$

Note that \mathbf{A} is an external or applied potential on, not the field generated by this moving charge, q . In QM, any particle can be represented by a wavepacket, and if the particle has a finite momentum, the wavepacket will also have finite wavelength. This means there will be a phase change across the wavepacket as the particle moves by, just as the phase of a sine wave sweeps from 0 to 2π as it moves past. The phase change is proportional to the line integral of the canonical momentum. Over a closed path, this must be $2\pi n$, where n is the number of times around the path. Recognizing that in a SC, the supercurrent \mathbf{J} is proportional to the (ordinary) momentum of the Cooper pairs of charge $2e$ and mass $2m$, we have

$$(13) \quad 2\pi n = (4\pi m/hde) \int \mathbf{J} \cdot d\mathbf{l} + (4\pi e/h) \int \mathbf{A} \cdot d\mathbf{l}$$

where h is the Planck constant and d in the denominator is the SC number density, while $d\mathbf{l}$ denotes a line integral around a closed contour. This is the Phase

Quantization condition. We can apply the Stokes theorem for vectors related by (9),

$$(14) \int \mathbf{A} \cdot d\mathbf{l} = \int \mathbf{curl A} \cdot d\mathbf{a} = \int \mathbf{B} \cdot d\mathbf{a},$$

to (13), where $d\mathbf{a}$ denotes a surface integral over a closed contour. If we also divide both sides of (13) by $4\pi e/h$, we obtain

$$(15) nh/2e = n\Phi = (m/de^2) \int \mathbf{J} \cdot d\mathbf{l} + \int \mathbf{B} \cdot d\mathbf{a}$$

This equation defines the Magnetic Fluxoid Quantum Φ on the LHS. The LHS is called the fluxoid rather than the flux because it is the sum of a flux term and another term. Since the first term on the RHS is a line integral, and since the current \mathbf{J} vanishes deep inside the SC, for any contour in the bulk SC, this term can be neglected, and we can say that the magnetic flux, defined by the second term RHS, is quantized.

Notice that it is the flux that is quantized, not the field.

Mathematically, the vanishing of the first term of (15) makes perfect sense. However, it is a little more difficult to interpret in terms of physical measurements. To do that, we will rewrite the first term by the Stokes theorem (14) to obtain

$$(16) nh/2e = n\Phi = (m/d^2) \int \mathbf{curl J} \cdot d\mathbf{a} + \int \mathbf{B} \cdot d\mathbf{a}$$

The screening layer with the current is contained in the surface integral when the contour is in the bulk SC, so we interpret (16) to mean that an experiment measures the TOTAL magnetic field due to the flux through the hole (2nd term) PLUS the magnetic flux generated by the London current layer around the hole (1st term). (In a SC, $\mathbf{curl J}$ is proportional to magnetic field.) Since that screening current is precisely the current necessary to establish and maintain the flux through the hole at a quantized value according to theory, experiments confirmed the theory when they detected a quantum of flux given by the LHS. Indeed, if the applied field had a flux of, say, $(n+1/4)\Phi$, a current is set up to move the trapped flux in the hole to an even quantized value of $n\Phi$. There is also a very small magnetic flux contribution from the penetration of the external field into the London layer around the hole (from which it is not screened, unlike

the rest of the SC). This would show up as a slight deviation from the quantized value.

If the SC is very thin, such that its thickness is no greater than the penetration depth, effectively, there is no bulk SC, and the contour goes right through the London layer around the hole. The interpretation of this in light of (16) is that the London current layer is no longer inside the integration surface area and so cannot contribute. Therefore, the only term left is the 2nd, with the result that a measurement now sees a deficit in the flux, since the 1st term has effectively been excluded. This deficit was also observed in experiments.

B.4 GM Quantization

The importance of the discussion in the last section is that the GM field simply adds another term to the RHS of each of the equations in that section. Therefore, if what is quantized and measured is the SUM of ALL terms on the RHS, there is, a good chance for some sort of GM quantum which could be measured.

To include the contribution of GM, one need only add appropriate terms of $2m\mathbf{A}_g/c$ or $2m\mathbf{B}_g/c$ to equations (12-16). The phase equation becomes

$$(17) \quad 2\pi n = (4\pi m/hde) \int \mathbf{J} \cdot d\mathbf{l} + (4\pi e/h) \int \mathbf{A} \cdot d\mathbf{l} + (4\pi m/ch) \int \mathbf{A}_g \cdot d\mathbf{l}$$

and the magnetic fluxoid equation,

$$(18) \quad nh/2e = n\Phi = (m/de^2) \int \mathbf{curl J} \cdot d\mathbf{a} + \int \mathbf{B} \cdot d\mathbf{a} + (m/ec) \int \mathbf{B}_g \cdot d\mathbf{a}.$$

If our interpretation in the last section holds up, we can expect the 3rd term of (18) to make a contribution to the measured magnetic flux such that the total for a bulk SC is the LHS. This term will be extremely small, not only because the applied GM is so small, but because m is the mass of an electron.

Now we may use the phase equation (17) to obtain a GM fluxoid relation by dividing both sides by $(4\pi m/ch)$:

$$(19) \quad nhc/2m = (c/de^2) \int \mathbf{curl J} \cdot d\mathbf{a} + (ec/m) \int \mathbf{B} \cdot d\mathbf{a} + \int \mathbf{B}_g \cdot d\mathbf{a}.$$

We have transformed the quantization relation into a form for which the LHS has dimensions of GM-flux (that is, acceleration-flux in cgs). Carrying over the interpretation of the previous section, we interpret the RHS as being the total GM flux in the system, which is

quantized. The first term is that GM flux from the moving currents in the London layer; the third term is the "pure" GM flux from the external field; the second term we interpret to be the GM flux induced by the applied magnetic field. Note that the GM quantum number is the same as the magnetic one for the same physical situation. Quantization of one establishes the other. And we could have just as easily reversed the process by deriving the GM flux quantization relation from the phase quantization relation first, and then derived the magnetic flux quantization relation from the GM relation.

Since the third term is so small, the important question is whether the interpretation of the second term is valid; this term will not be small in general, if for no other reason than the factor of $1/m$ is so large. Objections can be made to the concept that the entire RHS is measurable on the grounds that there is no known mechanism to explain the size of the second term if it is understood to be an induced GM field. Exotic mechanisms can be imagined, such as coupling of the magnetic and GM field lines through the topology of vortices, perhaps via a form of coherent zero point motion (Becker, Thesis, UAH 1994; or the possibility that it might be ions that are also responding to the GM field, in which case m would be at least 7500 times larger than the electron mass entered in the above equations. Another interpretation is that the second term represents the acceleration-flux induced by the magnetic field that could be felt by a magnetic body, if one were present. In this case, the RHS would not be entirely GM in origin. However, this last interpretation is not consistent with the interpretation of the GM term in the magnetic fluxoid as a GM induced magnetic field. All these ideas are pure speculation, however.

So the experiment hinges on that second term: if it does not physically contribute to a GM flux, then the other two terms are far too small to be measurable. On the other hand, if the three terms and the fluxoid itself are GM, the effect may be very large and definitely noticeable given the right circumstances.

B.5 GM Experiment Concept

The actual experiment consists of suspending a small high- T_c SC from a balancing-torsion mechanism akin to that used in the Cavendish experiment. A hole is scooped out of the bulk SC, such that a small, rotating cylinder of non-magnetic material can be placed inside. The

cylinder has its long axis oriented in the horizontal plane, and spins along that axis.

The cylinder is supported by a mechanism which allows rotation around the long axis, but resists rotation around any other. The SC itself, immersed in liquid nitrogen, is free to rotate in the horizontal plane. Cylinder rotation is maintained by a small gas jet. There is a small mirror or other such rotation sensor on the SC suspension assembly.

The SC is projected to be a YBa₂Cu₃O₇ compound, a high T_c, Type II SC, with T_c=90-98°K, and a lower critical field H_{c1} = 200G. Its configuration might be in the realm of 23mm in diameter and 4mm in thickness. The hole size will have to be chosen, in part, by considerations of fitting the cylinder suspension apparatus inside.

Before the hole is drilled into the SC, it will be tested to see if there are any detectable GM effects when the SC is magnetized above the lower critical field, whereupon vortices may enter the SC. Each vortex has n=1, which may produce a torque in its vicinity which can be calculated from the equations in section B.6.

If a strong enough total GM fluxoid quantum is present in the hole, due to a large applied magnetic field, the cylinder ought to precess about an axis perpendicular to the horizontal plane. However, the cylinder is constrained from precessing by the fixture which holds it in place against the air stream. Since the SC to which the cylinder is, we believe, coupled by virtue of the GM field, is free to rotate in the precession plane, the cylinder's precessional angular momentum is transferred to the SC, which twists about its suspension, permitting a measurement of the degree of turn from the sensor on the suspension. Alternatively, a direct measurement of the cylinder's precession can be attempted.

To calibrate the apparatus and ensure that there are no lingering magnetic effects in the cylinder, the cylinder set-up experiment should first be run with no SC present, with just the applied field. Next, the experiment should be repeated with the SC, but at a T > T_c, where there should be no quantization effects. Finally, the full SC plus cylinder tests can be run, varying the applied magnetic field as desired, as well as other parameters. A magnetized cylinder should be substituted for the non-magnetic test body, not only to verify the basic experimental set-up, but in case no GM effect is seen, to

verify that an experimental effect is present even in the well known case of the SC magnetic field.

B.6 Experiment Calculations

In this section we will outline the calculations needed to compute the forces that are expected to arise from the GM-induced precession of the cylinder based on the theory of section B.4 is correct.

For a given applied magnetic field, the fluxoid quantum number is obtained from

$$(20) \quad n = \pi r^2 H / 2.07 \times 10^{-7}$$

where H is the applied field, r is the radius of the hole and the magnetic fluxoid quantum has the value $2.07 \times 10^{-7} \text{ Gcm}^2$. Since n is an integral quantum number, it should be rounded to the nearest integer.

Since it is believed that the GM field can penetrate the SC, unlike the magnetic field, the area that enters the GM fluxoid is that of the whole SC, not the hole. From the LHS of (19) we estimate the GM field as an average over its penetration area. This may just be an estimate because, at least in the case of the magnetic field, the field is not uniform in a flat cylindrical SC as will be used for this experiment. Conversely, if the GM field is not expelled from the bulk SC, its lines should not deviate from linearity to any degree, making the average field estimate from the GM flux a better estimate than one for \mathbf{B} itself.

$$(21) \quad B_g = (nhc/2m)/\pi R^2 = (Hhc/2m)(r/R)^2/2.07 \times 10^{-7}$$

where R is the radius of the SC.

If B_g is given by (21), the precession velocity of the cylinder is therefore

$$(22) \quad \Omega = B_g/2c = (Hh/4m)(r/R)^2/2.07 \times 10^{-7}.$$

The torque the GM field will attempt to apply to the cylinder so as to result in the precession velocity (22) is given by (6):

$$(23) \quad \mathbf{N} = \mathbf{s} \times \mathbf{B}_g/2c = \mathbf{s} \times (Hh/4m)(r/R)^2/2.07 \times 10^{-7}.$$

Rigid body spin angular momentum is given by

$$(24) \quad \mathbf{s} = \mathbf{I}\boldsymbol{\omega}$$

where ω is the spin angular velocity (not the same as Ω). A cylinder rotating around its long axis has moment of inertia I of

$$(25) \quad I = M\alpha^2/2$$

where M is the mass of the cylinder and α is its radius. Putting both (24) and (25) into (23), we obtain (assuming right angles)

$$(26) \quad N = (M\omega\alpha^2/2)(Hh/4m)(r/R)^2/2.07 \times 10^{-7}.$$

Collecting factors, (26) can be rewritten as

$$(27) \quad N = (M\omega Hh/8m)(r\alpha/R)^2/2.07 \times 10^{-7}.$$

Since the cylinder will be fixed in the precessional rotation direction, but the SC will be free, the back coupling of the cylinder to SC should transfer the applied torque to the SC, resulting in the torque, and therefore force, on the SC suspension mechanism. The torque is given by (assuming right angles)

$$(28) \quad \mathbf{N} = \mathbf{R} \times \mathbf{F}.$$

So, assuming right angle relative to the apparatus,

$$(29) \quad F = N/R = (M\omega Hh/8Rm)(r\alpha/R)^2/2.07 \times 10^{-7}.$$

We can also write the force equation in terms of n , the quantum number, by reinserting (20) into the RHS of (29):

$$(30) \quad F = (nM\omega h/8\pi Rm)(\alpha/R)^2.$$

For the following suggested estimated parameter values

M - Mass of Test body - 2.0 g

w - Angular velocity of Test body - 10.0 rad/s

R - Radius of SC - 0.11 cm

r - Radius of SC hole - 0.05 cm

α - Radius of Test Body - 0.03 cm

m - Electron mass - 9.11×10^{-28} g

h - Planck Constant - 6.63×10^{-27} erg•s

the ratio of force to quantum number is

$$(31) F/n = 3.916 \text{ dynes}/n.$$

The value of n at the lower critical field (200 G) for the material proposed for this experiment, with r as above, is from (20)

$$(32) n = 7.58 \times 10^6.$$

This would make the force

$$F = 2.97 \times 10^7 \text{ dyne}!$$

For a more reasonable magnetic field, like the earth's (0.3 G) we get

$$n = 1.14 \times 10^4 \text{ dyne}$$

and the force is

$$F = 4.46 \times 10^4 \text{ dyne}.$$

It is essential to use small-area holes in the SC. While it is the flux which is quantized, it is the field which induces the precession. Too large a hole means that the flux can be dominated by the hole area, rather than the field, with the result that precessional effects could be quite small, despite a large applied magnetic field. Conversely, it is not quite customary to suspend small test bodies in small holes in SC, which may explain why these effects have not been noticed before.

C. Experiment Design

The proposed experiment concept is a combination of a variant of the classical Cavendish experiment for the measurement of the gravitational constant G , and the classic experiments demonstrating the quantization of magnetic flux in SC systems.

The Cavendish-like assembly will be designed to detect even small torques induced by the GM field in a high-density cylinder rotating inside a SC toroidal ring in the superconducting regime in trapped magnetic flux (see Fig. TBD). A persistent (paramagnetic) supercurrent can be established by lowering the temperature of a SC through the critical temperature in an applied magnetic field, and then removing the field. Quantized magnetic flux through the toroidal hole in the SC is thereby also

established. For ease of experimental design and operation at liquid nitrogen temperatures, the SC is chosen to be a High Tc ceramic oxide (TBD), operated in magnetic fields below its lower critical field of (TBD) to avoid the complication of flux vortices being induced in the SC.

The SC and the rotating cylinder are enclosed in a liquid nitrogen dewar made of transparent Pyrex glass about 1.0 m high and 0.15 m in diameter (see Fig. TBD). The dewar has two standard cryogenic silvered walls with a vacuum between them to insulate the liquid nitrogen inside from the external environment. The dewar has a vertical slit of un-silvered glass to permit some viewing of the interior.

A clear Pyrex glass vacuum chamber approximately 0.9 m high and 0.10 m in diameter is gently lowered by a manipulating mechanism into the dewar filled with liquid nitrogen. It has a vacuum flange at the top to permit the removal of a vacuum lid through which electrical wires and the manipulator arm pass to facilitate adjustments inside the chamber under cryogenic and vacuum conditions. Gas may be pumped out of the vacuum chamber through the vacuum lid with the aid of a high quality mechanical pump. Pressure in the chamber can be lowered to a fraction of a torr, thus reducing extraneous torques on the apparatus in the chamber due to convective flows. Helium or hydrogen gas jets can be passed in a highly controlled manner through the vacuum lid. These jets exert micro-torques on components of the apparatus in the chamber. The temperature inside the vacuum chamber are monitored via thermocouples.

Hanging from the vacuum lid is a thin and narrow flat strip of non-magnetic metal as in a Cavendish Gravitational Balance. Hanging on the end of the metal strip is a pure teflon yoke holding a pure glass rod which passes through holes in the yoke, so that the rod can turn freely with minimal friction, but without sliding back and forth in the yoke.

Centered on the glass rod, and concentric with it, is a precision cylinder (dimensions TBD) made of a high density substance. The dense cylinder is free to rotate at high speed around its axis with very little friction. It rotates precisely along one of the principle axes of its moment of inertia.

To act as experimental controls, eliminating the possibility of extraneous magnetic effects in the

cylinder, and for experiment operation above T_c , where there should be no macroscopic GM effect, cylinders of different compositions, but similar mass properties, will be capable of being changed out. The GM effect should be invariant to changes in cylinder composition, so long as the mass properties are kept the same, and so a variety of cylinder compositions will be used: relatively non-magnetic materials such as copper, lead, cesium, lead-glass, gold, silver, platinum with permeability TBD; paramagnetic materials such as TBD; and ferromagnetic materials such as iron, nickel, or magnetic composites (permeability TBD). Cavendish Balance measurements of any induced torques will be made at non-cryogenic temperatures, under external magnetic fields resulting in a flux equivalent to that which will be trapped in the SC, to calibrate and characterize the behavior of the sample cylinders in applied magnetic fields.

Pointed horizontally at the upper and lower surfaces of the cylinder, and from opposite sides of the cylinder, are four micro-tubes blowing helium or hydrogen gas against the cylinder, inducing it to spin in both possible directions without exerting on it any significant net force or torque parallel to the vertical axis. Opposite each of the four flow tubes is a "getter funnel" through which the gas which has just passed the cylinder is caught and pumped out.

There is a safety catch to hold the yoke securely any time the vacuum chamber is moved. The catch very gently releases the yoke once the vacuum chamber is securely in position. There is TBD provision for reducing the amplitude of any accidental oscillations of the yoke to zero by use of tiny jets of helium gas blowing against the yoke from opposite sides. The jets exert tiny, controlled torques of reversing sign applied out of phase with the yoke's natural frequency oscillations about the vertical axis. When not in use, the jets will be secured so as not to leak and exert accidental torques. Accidental vibrational and impulsive (bump) disturbances will also be minimized by TBD.

Potential thermal gradients in the vacuum chamber due to boil-off and heat influx can be reduced by first pumping-down the liquid nitrogen to a temperature below the atmospheric pressure level boiling point, and then letting the pressure build up as the temperature rises.

Visibility into the chamber can be temporarily increased by countering potential thermal gradients by the introduction of thermal baffles which can be raised and

lowered into the chamber, alternatively covering and uncovering the silvered and non-silvered portion of the vacuum chamber. The baffles might include many cylindrically concentric thin silver cylinders surrounding the vacuum chamber held together by thermally insulating spacers at separations of about 1.0 mm. Under conditions of thermal equilibrium they would help to keep heat energy from radiating into the vacuum chamber through the un-silvered portions of the dewar.

Using helium or hydrogen gas jets, the high density cylinder spins with its horizontal axis passing through the hole in the SC ring around which the supercurrent flows. While maintaining the basic toroidal topology, the SC could be varied in thickness from disk-like to cylinder-like. The latter, more solenoidal, dimensions may be preferable for ensuring uniformity of magnetic field lines, and presumably GM field lines, in the hole.

The SC ring is rigidly attached to the vacuum lid so any torque which may be exerted on the cylinder will be associated with a reaction in the SC, causing the yoke suspended from the metal strip to rotate around the vertical axis.

A convex mirror of about 2.0 m focal length is mounted on the teflon yoke holding the cylinder and reflects a horizontal beam from an external laser (the beam of which passes into and out of the vacuum chamber and liquid nitrogen dewar through the optically smooth glass surface). The horizontally reflected/focused beam lands as a point of light on a millimeter scale about 2.0 m away. Any significant rotation of the yoke and cylinder will cause shifts in the position of the light focus of more than 0.1 mm.

Three external Helmholtz coils are oriented so as to cancel the ambient magnetic field of the earth to below 0.1% of its normal value in three orthogonal directions, and are used to apply the external magnetic fields resulting in the quantized magnetic flux. Magnets of TBD-type are used to apply the external magnetic fields leading to the quantized magnetic flux. The apparatus can be placed in a standard TBD-cage to minimize unwanted magnetic fields of external origin when it is so desired.

The experimental apparatus will be pre-calibrated for sensitivity to Newtonian torques exerted by nearby mobile masses in the laboratory. Since these torques can also be generated by the presence of human experimenters, an

experimental protocol will be designed to minimize any such disturbance.

While the various sources of disturbance torques are postulated to be significantly lower magnitude than the predicted GM induced torques in strong magnetic fields, differential and resonance torsional measurement can be employed to further reduce any contamination.

Flexibility in the experiment design is sufficient to permit variations in the following experimental parameters:

1. The axis of the SC can be moved smoothly and without induced vibrations in or causing collisions between the rotating cylinder and the SC. This can be done under cryogenic and vacuum conditions.
2. The speed and direction of cylinder rotation can be changed by varying the directions and flow rates of the helium or hydrogen jets directed at the top and bottom of the cylinder, under cryogenic and vacuum conditions.
3. The magnitude of the trapped magnetic flux can be adjusted after each run of the experiment sequence by changing the applied magnetic field.
4. The composition of the cylinder and its mass properties (e.g. moment of inertia) are adjustable by switching out cylinder samples.

The following Tables illustrate schematically the way parameter space can be scanned to test whether there is or is not a macroscopic quantized GM flux separate and distinguishable from magnetic flux. In Table 1, it is assumed there is an applied magnetic field equivalent to some integer multiple of the magnetic flux quantum. In Table 2, it is assumed that some non-integer (say, $n+1/4$) of the magnetic flux quantum is applied. Each entry in the Tables contains an "equivalent" magnetic (for a magnetic test body) or GM (for a non-magnetic test body) flux strength that would be felt by the test body as a torque, in the different temperature regimes. If there is no macroscopic GM flux in the superconducting regime, then all entries in the second column would be zero in both Tables.

	Magnetic Test Body	Non-magnetic Test Body
--	--------------------	------------------------

$T < T_c$	n	n
$T > T_c$	n	0

Table 1: Integer Quantized Applied Magnetic Flux

	Magnetic Test Body	Non-magnetic Test Body
$T < T_c$	n	n
$T > T_c$	$(n+1/4)$	0

Table 2: Non-integer Quantized Applied Magnetic Flux

Revision History:

Original Draft - 9/23/95

Revision 1 - 11/17/95

Revision 2 - 9/4/97

Gravitational Meissner Effect

R.P. Lano

Department of Physics and Astronomy
The University of Iowa
Iowa City, IA, 52242

March 10, 1996

Abstract

The gravitational analogue of the electromagnetic Meissner effect is investigated. Starting from the post-Newtonian approximation to general relativity we arrive at gravitational London equations, predicting a gravitational Meissner effect. Applied to neutron stars we arrive at a London penetration depth of 12km, which is about the size of a neutron star.

PACS number(s): 04.60, 74.25.H, 97.60.J, 04.70

I Introduction

There is strong evidence that neutron stars are superfluid and maybe superconducting [1], indicating that macroscopic quantum effects play a significant role in neutron stars. In addition neutron stars are very strongly gravitating objects. We therefore would like to investigate the possibility of macroscopic quantum gravity effect in the form of a gravitational Meissner effect.

In the post-Newtonian approximation Einstein's theory of general relativity, gravity looks like a U(1) theory and is very similar to the electromagnetic Maxwell theory [2]. This allows us to follow closely the approach of London [3, 4] in deriving gravitational London equations, predicting a gravitational Meissner effect. Not knowing the underlying microscopic theory of quantum gravity we have to resort to this semi-classical approach. We find the gravitational London penetration depth to be about 12 km for a typical neutron star of radius 15 km. Furthermore, as the density of the neutron star increases, the London penetration depth decreases, suggesting that quantum gravity aspects may become significant as the neutron star approaches collapse.

Macroscopic quantum aspects are best described by a BCS type theory[5]. However, since BCS needs a quantum theory of the underlying interaction, and since we do not have a microscopic theory of quantum gravity at this moment,

a gravitational BCS type theory is not within our reach. Despite this, we can certainly make a plausible argument for what happens at the microscopic scale. Inspired from the Cooper-pair forming electrons in the BCS theory of superconductivity, the Interacting Boson Model (IBM) [6, 7] assumes nucleons to form pairs inside nuclei. These pairs are bosons and by only looking at the interaction of those bosons among themselves, one can explain a wealth of nuclear data to a high degree of accuracy. Neutron stars consist mostly of neutrons and some protons and for low enough temperatures ($< 10^9$ K) they are expected to pair up. Experimental evidence for the pairing comes from the theory of cooling of neutron stars, where the reduction of the specific heat due to extensive pairing is essential [8]. In addition, bose-condensation can occur and superfluidity as well as superconductivity are consequences. Evidence for superfluidity can be found in the glitches in the timing history of pulsars [1, 9]. Therefore, although we do not have a microscopic theory of quantum gravity, we have enough evidence justifying a phenomenological investigation of macroscopic quantum gravity effects following London's approach.

Symmetry breaking is a very convenient way to look at macroscopic quantum effects. To break a symmetry one needs a bose-condensate. What causes the bose-condensation and which symmetry is broken are two entirely unrelated problems. Any mechanism leading to an attraction between the fermions at the Fermi surface is equally well suited for producing the bose-condensate. For example, in the standard BCS theory of superconductivity, the attraction is caused by the phonon interaction, the symmetry broken is the electromagnetic U(1) symmetry [10]. In the theory of superfluid helium 3 [11], the attraction is caused by van der Waals forces, the symmetries broken are translational [12] and rotational in spin and orbital space as well as gauge symmetry. In the theory of the Interacting Boson Model, the attraction is caused by the strong nuclear force, the symmetry broken is related to angular momentum and Runge-Lenz vector [6]. In the theory of superconductivity of protons inside neutron stars, the attraction is caused by the strong nuclear force, the symmetry broken is the electromagnetic U(1) symmetry. Therefore, we see no reason why to exclude the scenario, where the attraction is caused by the strong nuclear force and the symmetry broken is gravitational.

This work is also an attempt to probe certain aspects of quantum gravity. The microscopic approach to quantum gravity has been poised with serious problems, and no satisfactory theory has emerged as of yet. The macroscopic approach seems much more tractable, and maybe following the theory of superconductivity from London to Ginzburg and Landau to Bardeen, Cooper and Schrieffer will give us another route to learn more about quantum gravity.

II Gravitational Maxwell Equations

Let us review the post-Newtonian approximation [13, 14]. For weak gravi-

tational fields and low velocities, i.e.,

$$U/c^2 \sim (v/c)^4 \sim (p/\rho c^2)^2 \sim (\Pi/c^2)^2 \ll 1,$$

Einstein's field equation can be written in a form resembling Maxwell's equations of electrodynamics. We are mostly interested in magnetic-type gravity, and therefore we shall use the truncated and rewritten version of the parametrized-post-Newtonian (PPN) formalism given by Braginsky et. al. [15]. Scalar and vector potentials are defined as

$$\phi = -(U + 2\Psi) \quad \text{and} \quad A_j = -\frac{7}{2}V_j - \frac{1}{2}W_j, \quad (1)$$

where U, Ψ, V_j , and W_j are defined in chapter 39 of Misner, Thorne and Wheeler [14]. This allows us to define a 'gravitoelectric' field \mathbf{g} and a 'gravitomagnetic' field \mathbf{H} through

$$\mathbf{g} = -\nabla\phi - \frac{1}{c}\frac{\partial\mathbf{A}}{\partial t} \quad \text{and} \quad \mathbf{H} = \nabla \times \mathbf{A}. \quad (2)$$

In this weak field, slow motion expansion of general relativity, \mathbf{g} contains mostly first order corrections to flat space-time, and \mathbf{H} contains second order corrections. For the solar system, \mathbf{g} is just the normal Newtonian gravitational acceleration, whereas \mathbf{H} is related to angular momentum interactions and effects due to \mathbf{H} are about 10^{12} times smaller than those due to \mathbf{g} .

In the above approximation, Einstein's field equations can be rewritten in a form very much resembling that of Maxwell's equations

$$\nabla \cdot \mathbf{g} = -4\pi G\rho_0 + \mathcal{O}(c^{-2}), \quad \nabla \times \mathbf{g} = -\frac{1}{c}\frac{\partial\mathbf{H}}{\partial t} \quad (3)$$

$$\nabla \cdot \mathbf{H} = 0, \quad \nabla \times \mathbf{H} = 4\left(-4\pi G\rho_0 \frac{\mathbf{v}}{c} + \frac{1}{c}\frac{\partial\mathbf{g}}{\partial t}\right), \quad (4)$$

where ρ_0 is the density of rest mass in the local rest frame of the matter and \mathbf{v} is the ordinary velocity of the rest mass relative to the PPN coordinate frame. The specific internal energy Π and the pressure p have been neglected since they only contribute $\mathcal{O}(c^{-2})$ effects to the electric field.

The equation of motion of an uncharged particle is identical to the electromagnetic Lorentz force law [2, 15],

$$\frac{d\mathbf{v}}{dt} = \mathbf{g} + \frac{1}{c}\mathbf{v} \times \mathbf{H} + \mathcal{O}(c^{-2}) \quad (5)$$

III Gravitational London Equations

With these gravitational Maxwell equations we can derive the equivalent of London's equations for gravity. The derivation is completely analogous to the electromagnetic case [4, 5, 16, 17]. The Lorentz force law tells us that given a gravitational electric field there will be a resultant acceleration. Multiplying by ρ_0 , the matter density, and defining the matter current $\mathbf{j} = \rho_0 \mathbf{v}$, we arrive at

$$\frac{d\mathbf{j}}{dt} = \rho_0 \mathbf{g}. \quad (6)$$

This is sometimes referred to as the first London equation. Now substitute this into the right equation of (3), the equivalent of Faraday's law, we obtain

$$\frac{\partial}{\partial t} \left(\frac{1}{\rho_0} \nabla \times \mathbf{j} + \frac{1}{c} \mathbf{H} \right) = 0. \quad (7)$$

One possible solution to the above equation is obtained by integrating and setting the integration constant to zero. This leads to the second (and most famous) London equation

$$\nabla \times \mathbf{j} = -\frac{\rho_0}{c} \mathbf{H}, \quad (8)$$

which predicts the Meissner-Ochsenfeld effect [18].

In the Ginzburg-Landau theory of superconductivity, it is shown that setting the integration constant in equation (7) to zero is equivalent to the assumption that the significant spatial variation of the order parameter $\psi = |\psi| e^{i\phi}$ is through the phase ϕ , and not the magnitude $|\psi|$ [16, 19]. For the density of Cooper pairs this means that it cannot vary appreciably from its uniform thermal equilibrium value. For a superfluid neutron star in thermal equilibrium, where the pairs can flow, but not accumulate or be destroyed, this should be the case.

To obtain the London penetration depth, we take the static part of the right equation of (4)

$$\nabla \times \mathbf{H} = -\frac{16\pi G}{c} \mathbf{j} \quad (9)$$

and take the curl of it. Using the second London equation (8) we arrive at

$$\nabla^2 \mathbf{H} = 16\pi \frac{G\rho_0}{c^2} \mathbf{H}, \quad (10)$$

where we immediately identify the gravitational London penetration depth

$$\Lambda_L = \left(\frac{c^2}{16\pi G\rho_0} \right)^{1/2} = 5.177 \times 10^{12} \rho_0^{-1/2}, \quad (11)$$

where the density ρ_0 is measured in kg/m^3 and Λ_L is given in meters.

IV Coherence Length

For our derivation of the London equations to be valid we must require that the London penetration depth Λ_L is much larger than the coherence length ξ_0 [17]:

$$\Lambda_L \gg \xi_0, \quad (12)$$

where the coherence length is defined through

$$\xi_0 = \frac{\hbar v_F}{\pi \Delta}, \quad (13)$$

with v_F the Fermi velocity and 2Δ the pairing energy or energy gap. We are only interested in a rough estimate of the upper limit of ξ_0 . An upper limit for the Fermi velocity obviously is the speed of light. If the pairing is caused by the strong nuclear force, as we assumed, then a lower limit on Δ is a few hundred thousand keV [20]. With these numbers we find for the coherence length

$$\xi_0 \leq 10^{-13} \text{m}, \quad (14)$$

which is larger than the separation between two nucleons, but much smaller than the London penetration depth. Therefore, our derivation of the London equations was justified in the given approximation.

V Discussion

We notice that for regular densities occurring in the universe the gravitational London penetration depth will be a very large number. However, interesting enough, for neutron stars with densities of about $\rho_{NS} \cong 2 \times 10^{17} \text{kg/m}^3$, we obtain a London length of 12km, which is slightly smaller than the radius of a neutron star with that density. The London length is inversely proportional to the density, meaning with increasing density the London penetration depth decreases.

For neutron stars this means that the onset of a gravitational Meissner effect is predicted. In essence it implies that the gravitational magnetic field caused by the huge angular momentum of the neutron star, will be expelled from the center of the neutron star. This could be accomplished through induced matter-supercurrents in the outer layers of the neutron stars, creating counter-magnetic fields to expel the gravitational magnetic field from its interior, in analogy to the electromagnetic case.

A more detailed description is not possible in the approximation used, since the post-Newtonian approximation is only a rough approximation if used for neutron stars. We are currently working on the second order post-Newtonian approximation and numerical simulations [21].

VI Conclusion

More significant than the expulsion of the gravitational magnetic field, however, may be the fact that the Meissner effect is a macroscopic quantum effect, and in essence, the *gravitational* Meissner effect is a macroscopic quantum *gravity* effect. There is experimental evidence that neutron stars are superfluid as well as superconducting. This indicates that macroscopic quantum effects do play a significant role for neutron stars. Furthermore, neutron stars are the strongest gravitational objects observed with certainty. Under the assumptions used here, we find that macroscopic quantum gravity effects may play a significant role for neutron stars.

Since the work of Oppenheimer and Volkoff [22], we know that neutron stars will be stable only up to a certain point. After that the gravitational pressure overcomes the pressure of the neutron Fermi gas and continued collapse is predicted. This leads to increasing densities, and in our model to decreasing London penetration lengths, indicating that gravitational quantum effects might become more and more important. Therefore, we would like to emphasize that the prediction of Oppenheimer and Volkoff is classical in nature, and our work indicates, that a quantum gravitational treatment may lead to new results.

VII Acknowledgment

I would like to thank Professor V.G.J. Rodgers for spawning my interest in this topic and for his insightful guidance throughout this work.

References

- [1] D. Pines et. al., eds., The Structure and Evolution of Neutron Stars, Addison-Wesley Publishing Company, 1992
- [2] K.S. Thorne in J.D. Fairbank et. al., eds., Near Zero: New Frontiers of Physics, W.H. Freeman and Company, 1988
- [3] F. London, Phys. Rev. 24, 562 (1948)
- [4] F. London, Superfluids Vol.1, John Wiley & Sons, 1950
- [5] J.R. Schrieffer, Theory of Superconductivity, Addison-Wesley Publishing Company, 1964
- [6] F. Iachello, Rev. Mod. Phys. 65, 569 (1993)
- [7] M. Mukerjee and Y. Nambu, Ann. Phys. 191, 143 (1989)
- [8] D. Page, Astrophys. J. 428, 250 (1994)

- [9] J.A. Saul in H. Ögelman et. al., eds., Timing Neutron Stars, Kluwer Academic Publishers, 1989
- [10] L.H. Ryder, Quantum Field Theory, Cambridge University Press, 1985
- [11] D. Vollhardt and P. Wölfle, The Superfluid Phases of Helium 3, Taylor & Francis, 1990
- [12] N.D. Mermin in J. Ruvalds et. al., eds., Quantum Liquids, North-Holland Publishing Company, 1978
- [13] S. Weinberg, Gravitation and Cosmology, John Wiley & Sons, 1972
- [14] C.W. Misner, K.S. Thorne and J.A. Wheeler, Gravitation, W.H. Freeman and Company, 1973
- [15] V.B. Braginsky, C.M. Caves and K.S. Thorne, Phys. Rev. D 15, 2047 (1977)
- [16] N.W. Ashcroft and N.D. Mermin, Solid State Physics, CBS Publishing Asia LTD, 1976
- [17] P.G. de Gennes, Superconductivity of Metals and Alloys, Addison-Wesley Publishing Company, 1989
- [18] W. Meissner and R. Ochsenfeld, Naturwissenschaften 21, 787 (1933)
- [19] V.L. Ginzburg and L.D. Landau, Zh. Eksp. Teor. Fiz. 20, 1064 (1950)
- [20] J.W. Clark, R.D. Davé and J.M.C. Chen in D. Pines et. al., eds., The Structure and Evolution of Neutron Stars, Addison-Wesley Publishing Company, 1992
- [21] R.P. Lano, in preparation
- [22] J.R. Oppenheimer and G. Volkoff, Phys. Rev. 55, 374 (1939)

UTF-367/96
Jan 1996

Updating the Theoretical Analysis of the Weak Gravitational Shielding Experiment.

Giovanni Modanese¹

*I.N.F.N. – Gruppo Collegato di Trento
Dipartimento di Fisica dell’Università
I-38050 POVO (TN) - Italy*

Abstract

The most recent data about the weak gravitational shielding produced recently through a levitating and rotating HTC superconducting disk show a very weak dependence of the shielding value ($\sim 1\%$) on the height above the disk. We show that whilst this behaviour is incompatible with an intuitive vectorial picture of the shielding, it is consistently explained by our theoretical model. The expulsive force observed at the border of the shielded zone is due to energy conservation.

74.72.-h High- T_c cuprates.

04.60.-m Quantum gravity.

The measurements of Podkletnov et al. of a possible weak gravitational shielding effect [1, 2] have been repeated several times and under different conditions by that group, with good

¹e-mail: modanese@science.unitn.it

reproducibility, including results in the vacuum. In the forthcoming months other groups will hopefully be able to confirm the effect independently. While the Tampere group was mainly concerned with obtaining larger values for the shielding, studying its dependence on numerous experimental parameters and testing new materials for the disk, in the future measurements it will be important to obtain more exact data, including detailed spatial field maps. The theoretical model suggested by us [3] is still evolving, although at a fundamental level; a more detailed account appears elsewhere [4].

Let us recall in short the main features of the experiment. A HTC superconducting disk or toroid with diameter between 15 and 30 *cm*, made of $YBa_2Cu_3O_{7-x}$, is refrigerated by liquid helium in a stainless steel cryostat at a temperature below 70 *K*. The microscopic structure of the material, which plays an important role in determining the levitation properties and the amount of the effect, is described in details in the cited works.

The disk levitates above an electromagnet and rotates by the action of lateral alternating e.m. fields. Samples of different weight and composition are placed over the disk, at a distance which can vary from a few *cm* to 1 *m* or more (see below). A weight reduction of about 0.05% is observed when the disk is levitating but not rotating; the weight loss reaches values about 0.5% when the disk rotates at a frequency of ca. 5000 *rpm*. If at this point the rotating fields are switched off, the sample weight remains decreased till the rotation frequency of the disk decreases. On the other hand, if the rotation frequency is decreased from 5000 to 3500 *rpm* using the solenoids as breaking tools, the shielding effect reaches maximum values from 1.9 to 2.1%, depending on the position of the sample with respect to the outer edge of the disk.

This effect, if confirmed, would represent a very new and spectacular phenomenon in gravity; namely, as well known, there has never been observed any conventional gravitational shielding up till now, up to an accuracy of one part in 10^{10} , and General Relativity and perturbative Quantum Gravity exclude any measurable shielding [3]. Our temptative theoretical explanation is based on some properties of non-perturbative quantum gravity. We have shown that the density field $|\phi_0|^2$ of the Cooper pairs inside the superconductor or, more likely, the squared gradient $(\partial_\mu\phi_0)^*(\partial^\mu\phi_0)$ may act as localized positive contributions to the small negative effective gravitational cosmological constant Λ ; if the sum turns out to be positive in a certain four-dimensional region, a local gravitational singularity arises there, affecting the gravitational propagators and thus the interaction potential (between the Earth and the samples, in this

case).

To sketch our model – although not rigorously – we could say that there is an "anomalous coupling" between the mentioned density $|\phi_0|^2$ or the squared gradient $(\partial_\mu\phi_0)^*(\partial^\mu\phi_0)$ and the gravitational field, and that the net result is to partly "absorb" the field. We expect that only in some regions of the superconductor the density $|\phi_0|^2$ or the squared gradient will be strong enough and that the inhomogeneities of the material and the pinning centers will be crucial in determining such regions. Since the gravitational field is attractive, its "absorption" requires energy from the outside. This means that there must be some mechanism external to the disk which *forces* $|\phi_0|^2$ or $(\partial_\mu\phi_0)^*(\partial^\mu\phi_0)$ to take high values. This is caused in the experiment by the action of the external electromagnetic field and by the disk rotation.

The dependence of the shielding effect on the height, at which the samples are placed above the superconducting disk, has been recently measured up to a height of ca. 3 m [5]. No difference in the shielding value has been noted, with a precision of one part in 10^3 . It is also remarkable that during the measurement at 3 m height the sample was placed in the room which lies above the main laboratory, on the next floor; in this way the effect of air flows on the measurements was greatly reduced. For the used 500 g sample the weight loss was ca. 2.5 g.

Such an extremely weak height dependence of the shielding is in sharp contrast with the intuitive picture, according to which the gravitational field of the Earth is the vectorial sum of the fields produced by each single "portion" of Earth. In the absence of any shielding, the sum results in a field which is equivalent to the field of a pointlike mass placed in the center of the Earth; this can be checked elementarily by direct integration or invoking Gauss' theorem and the spherical symmetry. But if we admit that the superconducting disk produces a weak shielding, the part of the Earth which is shielded lies behind the projection of the disk as seen from the sample, i.e., within an angle θ about the vertical direction, such that $\tan\theta = h$, where h is the sample height over the disk measured in units of the disk radius. (For simplicity we suppose now the sample to be centered above the disk.)

In order to obtain the shielding effect as a function of h , taking into account this geometrical factor, one must integrate the Newtonian contribution $\cos\phi/R^2$ over the intersection between the Earth and the cone defined by $\phi < \theta$. We have done this for the values $h = 1, 2, 3, 4, 6, 8, 10$, through a Montecarlo algorithm. We took into account the higher density of the Earth's core ($\rho_{core} \sim 2\rho_{mantle}$; $r_{core} \sim (1/2)r_{Earth}$; it is straightforward to insert more accurate values, but

the final results change very little); we also computed analytically the contribution of the tip of the cone, from the Earth's surface to the Earth's core, in order to reduce the fluctuations in the Montecarlo samplings for small R .² The resulting values were the following:

h	shielding/maximum-shielding
<hr/>	
1	0.62 +/- 0.02
2	0.34 +/- 0.01
3	0.18 +/- 0.01
4	0.102 +/- 0.003
6	0.050 +/- 0.002
8	0.029 +/- 0.001
10	0.018 +/- 0.001

This strong height dependence is clearly incompatible with the mentioned experimental data, which instead seem to indicate that in the shielding process all the mass of the Earth behaves effectively as if it would be concentrated in one point.

In our theoretical model this property arises in a natural way. We employ a quantum formula which expresses the static gravitational interaction energy of two masses m_1 and m_2 in terms of an invariant vacuum expectation value, namely [6]

$$E = \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{\int d[g] \exp \left\{ -\hbar^{-1} \left[S[g] + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{g_{\mu\nu}[x_i(t)]} \dot{x}_i^\mu(t) \dot{x}_i^\nu(t) \right] \right\}}{\int d[g] \exp \left\{ -\hbar^{-1} S[g] \right\}} \quad (1)$$

$$\equiv \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \left\langle \exp \left\{ -\hbar^{-1} \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} ds_i \right\} \right\rangle_S \quad (2)$$

where g has Euclidean signature and S is the gravitational action of general form

$$S[g] = \int d^4x \sqrt{g} \left(\lambda - kR + \frac{1}{4} a R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right). \quad (3)$$

²For the detailed algorithm and figures please ask the author at the e-mail address above.

The constants k and λ are related – in general as “bare quantities” – to the Newton constant G and to the cosmological constant Λ : k corresponds to $1/8\pi G$ and λ to $\Lambda/8\pi G$. The trajectories $x_i(t)$ of m_1 and m_2 are parallel with respect to the metric g ; let R be their distance.

In the weak-field approximation, eq. (1) reproduces to lowest order the Newton potential and can be used to find its higher order quantum corrections [7], or implemented on a Regge lattice to investigate the non-perturbative behaviour of the potential at small distances [9]. The addition to the gravitational action (3) of a term which represents a localized *external* Bose condensate³ mimics a shielding effect which is absent from the classical theory and which we take as our candidate model for the observed shielding.

The feature of eq. (1) which is of interest here is that if the two masses m_1 and m_2 are not pointlike, the trajectories $x_1(t)$ and $x_2(t)$ must be those of their centers of mass. (This also makes irrelevant the question – actually ill-defined in general relativity – whether they are pointlike or not.) Thus, when applying eq. (1) to the Earth and the sample, we only need to consider the centers of mass of those bodies. In this way we reproduce the observed behaviour for the shielding as well as for the regular interaction. The ensuing apparent failure in the “local transmission” of the gravitational interaction does not contrast with any known property of gravity (compare [6, 8], and references about the problem of the local energy density in General Relativity and [6] about the non-localization of virtual gravitons. One should also keep in mind that (1) holds only in the static case.)

Finally, if we describe the shielding effect as a slight diminution of the effective value of the gravitational acceleration g , and remember that the gravitational potential energy $U = -\frac{Gm_{Earth}}{r_{Earth}} = -gr_{Earth}$ is negative, it follows that the energy of a sample inside the shielded zone is larger than its energy outside. This means in turn that the sample must feel an expulsive force at the border of the shielded region. Such a force has been indeed observed [5], although precise data are not available yet. From the theoretical point of view it is however not trivial to do any prevision about the intensity of the force. In fact, the shielding process absorbs energy from the experimental apparatus and thus any transient stage is expected to be highly non-linear, especially for heavy samples.

³This means that the density of the condensate is not included into the functional integration variables.

References

- [1] E. Podkletnov and R. Nieminen, Physica **C 203** (1992) 441.
- [2] E. Podkletnov and A.D. Levit, *Gravitational shielding properties of composite bulk $YBa_2Cu_3O_{7-x}$ superconductor below 70 K under electro-magnetic field*, Tampere University of Technology report MSU-95 chem, January 1995.
- [3] G. Modanese, *Theoretical analysis of a reported weak gravitational shielding effect*, report MPI-PhT/95-44, hep-th/9505094, May 1995.
- [4] G. Modanese, *Role of a "local" cosmological constant in Euclidean quantum gravity*, to appear as report UTF and on hep-th@xxx.lanl.gov, January 1996.
- [5] E. Podkletnov, private communication, October 1995.
- [6] G. Modanese, Phys. Lett. **B 325** (1994) 354; Nucl. Phys. **B 434** (1995) 697; Riv. Nuovo Cim. **17**, n. 8 (1994).
- [7] I.J. Muzinich and S. Vokos, Phys. Rev. **D 52** (1995) 3472.
- [8] D. Bak, D. Cangemi, R. Jackiw, Phys. Rev. **D 49** (1994) 5173.
- [9] H.W. Hamber and R.M. Williams, Nucl. Phys. **B 435** (1995) 361.

UTF-368/96
Jan 1996

Role of a "Local" Cosmological Constant in Euclidean Quantum Gravity.

Giovanni Modanese¹

*I.N.F.N. – Gruppo Collegato di Trento
Dipartimento di Fisica dell'Università
I-38050 POVO (TN) - Italy*

Abstract

In 4D non-perturbative Regge calculus a positive value of the effective cosmological constant characterizes the collapsed phase of the system. If a local term of the form $S' = \sum_{h \in \{h_1, h_2, \dots\}} \lambda_h V_h$ is added to the gravitational action, where $\{h_1, h_2, \dots\}$ is a subset of the hinges and $\{\lambda_h\}$ are positive constants, one expects that the volumes V_{h_1}, V_{h_2}, \dots tend to collapse and that the excitations of the lattice propagating through the hinges $\{h_1, h_2, \dots\}$ are damped. We study the continuum analogue of this effect. The additional term S' may represent the coupling of the gravitational field to an external Bose condensate.

04.20.-q Classical general relativity.

04.60.-m Quantum gravity.

¹e-mail: modanese@science.unitn.it

Since the first perturbative formulations of quantum gravity it was realized that the addition of a cosmological term $\frac{\Lambda}{8\pi G} \int d^4x \sqrt{g(x)}$ to the pure Einstein action gives the graviton a mass, which is positive if $\Lambda < 0$ and negative – that means, the theory becomes unstable – if $\Lambda > 0$ [1].

Nevertheless, whilst the vacuum fluctuations of the quantum fields should in principle produce a very large value of the cosmological constant, there is no observable hint of it. Namely, neither the Newtonian potential shows any finite range up to solar system distances, nor pure gravity exhibits any instability in the weak field case; finally the observations on cosmological scale set an upper limit on $|\Lambda|$ as small as $|\Lambda| < 10^{-120}G^{-1}$ in natural units. This discrepancy is known as the "cosmological constant problem" and several possible escapes have been suggested [2]. Treating the problem at a fundamental level requires a non-perturbative approach, since one should be able to explain why the actual large-scale geometry of spacetime is flat just from dynamic considerations. It is therefore not surprising that a definitive and generally accepted solution of the paradox of the cosmological constant is still remote. We shall not make any attempt to a new explanation here.

We shall consider 4D pure gravity in the Euclidean approach, with special reference to quantum Regge calculus [3]. In this model the results of the numerical non-perturbative simulations compose the following picture of the behaviour of the cosmological constant: while its "bare" value λ is generally nonzero, the effective value Λ depends on the energy scale μ and vanishes at large distances like $|\Lambda| \sim G^{-1}(\mu l_0)^\gamma$, where l_0 is the lattice spacing and γ a critical exponent. This means that the quantum geometry fluctuates on small scales, but reproduces flat space at macroscopic distances. The sign of Λ is negative, thus the flat limit is well defined.

Our aim is to study the interaction of this gravitational system with a particular external source, namely a Bose condensate described by a scalar field $\phi(x) = \phi_0(x) + \tilde{\phi}(x)$. We assume that the vacuum density $\phi_0(x)$ is forced from the outside to a certain value, as it can happen for instance in a superconductor subjected to external electromagnetic fields. The positivity of the Euclidean action of ϕ ensures that the terms $[\partial^\mu \phi_0(x)]^* [\partial_\mu \phi_0(x)]$ and $m_\phi^2 |\phi_0(x)|^2$ act like positive cosmological contributions to the gravitational action, possibly inducing local gravitational instabilities.

The structure of the paper is the following. In Section 1 we recall the main results of Euclidean 4D quantum Regge calculus concerning the effective cosmological constant. We

mention its scale behaviour and two possible interpretations of the lattice spacing l_0 . In Section 2 we consider in the continuum theory, at distances much larger than Planck scale, a weak Euclidean gravitational field $g_{\mu\nu}(x) = \delta_{\mu\nu} + \kappa h_{\mu\nu}(x)$, whose action includes an infinitesimal effective cosmological term with $\Lambda < 0$. Such a theory can be regarded as the continuum limit of the lattice theory described in Section 1. We then couple minimally $g_{\mu\nu}(x)$ to the mentioned scalar field $\phi(x)$, with vacuum density $\phi_0(x)$ determined from the outside, and write in detail the various terms of the action. It turns out that the infinitesimal graviton mass $m_g^2 \propto |\Lambda|$ receives a local negative contribution, which we denote by $-\mu^2(x)$. This means, as we show explicitly for the simpler case of an almost-massless scalar field χ , that if there exist some four-dimensional regions in which $\mu^2(x) > m_\chi^2$, it is possible to find field configurations $\chi(x)$ which make the Euclidean action unbounded from below. The field tends to develop singularities in those regions, or – if a cut-off mechanism comes into play – it tends to assume constant extremal values which are independent from those in the neighboring regions. We suggest that these "constraints" should be inserted in the equation for the propagator of the field.

In Section 3 we remind the formula which gives the static potential energy in Euclidean quantum gravity as a functional average [4, 5]. We illustrate its statistical meaning in the case of a weak field through an analogy with a simple 2D Ising model, which we also treat numerically with an elementary simulation. We show numerically that the insertion of local supplementary constraints due to an external field (in analogy to the singularities of the continuum case) damps in a sensible way the spin-spin correlations, and thus the interaction they represent in the model.

1 Regge calculus.

In the last years discretized quantum gravity on the Regge lattice has led, through the MonteCarlo numerical simulations of Hamber and Williams, to a better comprehension of the non-perturbative behaviour of the Euclidean gravitational field in four dimensions. Some of the features which emerge from their results [3] depend on the specific model, but several others are quite general. In this approach the physical quantities are extracted from functional

averages. The partition function is written as

$$Z = \int_{\text{Geometries}} d[g] e^{-\hbar^{-1} S[g]} \quad (1)$$

where the geometries are described by piecewise flat simplicial manifolds. The integral (i.e. the Montecarlo sampling) runs over the lengths of the links which define a field configuration. The action in (1) has the form

$$S = \int d^4x \sqrt{g} \left(\lambda - kR + \frac{1}{4}aR_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \right) \quad (2)$$

or, in discretized version,

$$S[l] = \sum_{\text{hinges } h} V_h \left[\lambda - k \frac{A_h \delta_h}{V_h} + a \frac{A_h^2 \delta_h^2}{V_h^2} \right] \quad (3)$$

where A_h is the area of an hinge, V_h its volume and δ_h the defect angle (see the original papers for the definitions of lattice quantities and the functional measure).

In the following we shall set $\hbar = 1$. The constants k and λ are related, as "bare" quantities, to the Newton constant G and to the cosmological constant Λ : k corresponds to $1/8\pi G$ and λ to $\Lambda/8\pi G$. It is important, however, to keep distinct the physical values G and Λ from k and λ . The latter are entered as parameters at the beginning, and then a second order transition point for the statistical system described by Z is found by Montecarlo simulation. Actually, there is a line of transition, since one can also vary the adimensional parameter a , which does not have a macroscopic counterpart. On this line in the parameter space the theory admits a continuum limit. Unlike in perturbation theory, where a flat background is introduced by hand, here the flat space appears dynamically; namely, the average value of the curvature is found to vanish on the transition line, which separates a "smooth phase", with small negative curvature, from a "rough", unphysical phase, with large positive curvature. In this way the effective, large scale cosmological constant

$$\Lambda = \frac{\langle \int \sqrt{g} R \rangle}{\langle \int \sqrt{g} \rangle} \quad (4)$$

vanishes in the continuum quantum theory.

More precisely, the dependence of the effective (or "running") cosmological constant Λ_{eff} on the scale is the following. If we compute (4) on small volumes, the curvature fluctuates wildly. At larger distances the average curvature decreases, because the fluctuations tend to

average out. If μ denotes the energy scale, close to the critical point the adimensional quantity $|\Lambda|G$ behaves like

$$(|\Lambda|G)(\mu) \sim (l_0\mu)^\gamma, \quad (5)$$

where l_0 is the average spacing of the dynamical lattice and $\gamma \sim 1.56$ in the first simulations. The sign of Λ is negative, as mentioned. Since the μ -dependence of G is quite weak, the effective cosmological constant decreases approximately like a power law as the length scale grows.

Eq. (5) admits two different physical interpretations, depending on the role we attribute to the average lattice spacing $l_0 = \sqrt{\langle l^2 \rangle}$. In the usual lattice theories, l_0 is sent to zero in order to obtain the continuum limit. In this first interpretation, eq. (5) shows the way the lattice theory reproduces flat space in the physical limit. No real physical meaning is assigned to the effective Λ which, at a fixed scale μ , is simply proportional to a positive power of the regulator.

On the other hand, we can believe that in quantum gravity l_0 has an intrinsic minimum value of the order of Planck length $L_{Planck} \sim 10^{-33} \text{ cm}$. This hypothesis arises independently from several operational models (for a review see [6]) or from more complex quantum theories (see for instance [7] and references). In the framework of Regge calculus, it is possible to fix l_0 by imposing that the effective Newton constant G computed non-perturbatively [5] is of the same magnitude order of the observed value; in this way one finds one more time that l_0 has to be of the order of Planck length [8].

In this second case, that is $l_0 \sim L_{Planck}$, the interpretation of eq. (5) is different: it means that the effective cosmological constant tends to zero on large scale, while it is nonvanishing, in principle, on small scale (here and in the following we mean by "large" scale the laboratory or atomic scale, and by "small" scale the Planck scale). This interpretation does not necessarily have observable physical consequences, since in fact Λ could be far too small. Namely, the exponent gamma has been computed only for small lattices; an evaluation for lattices of "macroscopic" size is of course technically impossible, and thus only the experiments could tell us whether the law (5) keeps true for large distances, and with which exponent. The fact that on astronomical scale we have $|\Lambda|G < 10^{-120}$ for $\mu \sim 10^{-30} \text{ cm}^{-1}$ constrains γ to be approximately larger than 2. But the vanishing could be much more rapid, so that we could disregard Λ at any physically relevant scale.

Another property of Regge calculus which shows an intrinsic feature of quantum gravity is the instability of the "rough" phase with positive average curvature, i.e. with positive effective

cosmological constant. This phase does not admit any continuum limit. Its fractal dimension is small, which denotes that the geometry is collapsed.

The non-perturbative instability properties of the phase with positive average curvature extend the validity of known considerations based on perturbation theory. Namely, in the weak field approximation a positive cosmological constant in the gravitational lagrangian produces a negative mass for the graviton (compare [1] and the next Section). Instabilities arise in the perturbative theory on a De Sitter background too [9, 10]. On the contrary, a small negative cosmological constant generally does not imply any instability, but a small mass m_g for the graviton, of the order of $\Lambda^{1/2}$ in natural units.

In conclusion, at a scale large enough with respect to the lattice scale it is possible to regard the discretized gravitational functional integral (1) as describing an almost flat mean field plus fluctuations. If the fixed point of the lattice theory is approached from the physical, smooth phase, the effective cosmological constant Λ (i.e. the average of the scalar curvature R) is very small and negative, and the large scale fluctuations of R are small too. The system is stable, because the field configurations with larger volume, in which the links are as stretched as possible, are preferred to the collapsed configurations, since the Euclidean action depends on the volume like $S \sim \frac{\Lambda}{8\pi G} \int d^4x \sqrt{g} \sim \frac{\Lambda}{8\pi G} \sum_h V_h$. (On the contrary, a positive value of Λ would favour the collapsed configurations with smaller volume.) In this picture Λ can be regarded either as a purely formal regulator, which goes to zero in the physical limit $l_0 \rightarrow 0$, or as a physical quantity, though possibly extremely small (in the second interpretation, in which $l_0 \sim l_{Planck}$).

Keeping these properties in mind, it is interesting to consider the case in which the coupling of the Euclidean gravitational field with an external source gives in some four-dimensional regions a positive contribution to the effective cosmological constant. In the next Section we shall analyse this phenomenon in the continuum case, that is, on large scale (in the meaning of "large scale" we precised above).

On the lattice, such a coupling would correspond in the action (3) to an additional term of the form

$$\sum_{\{h_1, h_2, \dots\}} \lambda_h V_h \quad (6)$$

where $\{h_1, h_2, \dots\}$ is a subset of the hinges and λ_h are fixed positive constants.

We expect that when the Montecarlo algorithm chooses for the random variation a link which ends in a hinge $h_i \in \{h_1, h_2, \dots\}$, the favoured variation will be that for which the volume V_{h_i} decreases. Thus the volumes $\{V_{h_1}, V_{h_2}, \dots\}$ will tend to collapse and the lattice excitations propagating through the hinges $\{h_1, h_2, \dots\}$ will be damped.

2 Continuum case.

We have seen that the results of quantum Regge calculus can be interpreted as leading at distances large compared to L_{Planck} to an effective Euclidean action for pure gravity of the form

$$S_{eff} = \int d^4x \sqrt{g(x)} \left[\frac{\Lambda}{8\pi G} - \frac{1}{8\pi G} R(x) \right] \quad (7)$$

where the curvature fluctuations around flat space are small and the effective cosmological constant Λ is negative and very small too. (As we saw in the previous Section, Λ is scale-dependent; we suppose here to stay at some fixed scale.) From the geometrical point of view, the small negative value of Λ stabilizes the system, preventing it from falling into small-volume, collapsed configurations.

In the naive perturbation theory around the flat background the Λ -term represents a small mass for the graviton ². Namely, setting $g_{\mu\nu}(x) = \delta_{\mu\nu} + \kappa \tilde{h}_{\mu\nu}(x)$, with $\kappa = \sqrt{8\pi G}$, the determinant g of $g_{\mu\nu}$ can be expanded as

$$g = 1 + \kappa \tilde{h}^{(1)} + \kappa^2 \tilde{h}^{(2)} + \dots, \quad (8)$$

where $\tilde{h}^{(1)}, \tilde{h}^{(2)}, \dots$ denote expressions which are linear, quadratic etc. in $\tilde{h}_{\mu\nu}$. The linear "tadpole" term $\tilde{h}^{(1)}$ is usually disregarded, since it is proportional to the trace \tilde{h}_μ^μ , which vanishes on physical states. The term $\tilde{h}^{(2)}$ takes the form of a graviton mass term, such that the mass is positive when $\Lambda < 0$ (compare [1]). We thus have

$$\begin{aligned} S_{eff} = \int d^4x \{ & \left[m_g^2 \tilde{h}^{(2)}(x) - \tilde{R}^{(2)}(x) \right] + \Lambda \left[\kappa \tilde{h}^{(3)}(x) + \kappa^2 \tilde{h}^{(4)}(x) + \dots \right] + \\ & - \left[\kappa \tilde{R}^{(3)}(x) + \kappa^2 \tilde{R}^{(4)}(x) + \dots \right] \} \end{aligned} \quad (9)$$

²As remarked in [1], this widespread belief is not rigorously true. It has also been proved [11] that all theories of a massive tensor field in Minkowski space which satisfy the usual QFT postulates are incompatible with General Relativity in the limit of vanishing mass. In our reasoning it is not essential to regard the cosmological term as a graviton mass term, but for simplicity we stick to this terminology.

where m_g is proportional to $\Lambda^{1/2}$ and $\tilde{R}^{(2)}, \tilde{R}^{(3)} \dots$ denote the parts of R which are quadratic, cubic etc. in $\tilde{h}_{\mu\nu}$.

The first bracket contains the quadratic part of the action. The third bracket contains the familiar self-interaction vertices of the graviton, involving respectively 1, 2, ... derivatives of $h_{\mu\nu}$. The second bracket contains self-interaction vertices which are peculiar of the theory with $\Lambda \neq 0$ and do not involve derivatives. We are however not interested in the self-interaction vertices of $h_{\mu\nu}$ in the following.

Now we would like to consider the interaction of \tilde{h} with a scalar field ϕ having non-vanishing vacuum expectation value ϕ_0 . We suppose ϕ_0 to be spacetime dependent and denote $\phi(x) = \phi_0(x) + \tilde{\phi}(x)$; $\phi_0(x)$ is regarded as a quantity determined from the outside, that is, as a source term. In this way, the field $\phi(x)$ describes a Bose condensate with ground state density $\phi_0(x)$ fixed by external conditions (like, for instance, the Cooper pairs density in certain superconductors under external e.m. field).

The total action is

$$S = S_{eff} + \int d^4x \sqrt{g(x)} \left\{ \partial_\mu [\phi_0(x) + \tilde{\phi}(x)]^* \partial_\nu [\phi_0(x) + \tilde{\phi}(x)] g^{\mu\nu}(x) + \frac{1}{2} m_\phi^2 |\phi_0(x)|^2 + \frac{1}{2} m_\phi^2 [\phi_0^*(x) \tilde{\phi}(x) + \phi_0(x) \tilde{\phi}^*(x)] + \frac{1}{2} m^2 |\tilde{\phi}(x)|^2 \right\} \quad (10)$$

We can rewrite it as

$$S = \int d^4x \sqrt{g(x)} \left\{ \left[\frac{\Lambda}{8\pi G} + \frac{1}{2} \mu^2(x) \right] - \frac{1}{8\pi G} R(x) \right\} + S_1 + S_2, \quad (11)$$

where

$$\frac{1}{2} \mu^2(x) = \frac{1}{2} [\partial_\mu \phi_0^*(x)][\partial^\mu \phi_0(x)] + \frac{1}{2} m_\phi^2 |\phi_0(x)|^2; \quad (12)$$

$$S_1 = \frac{1}{2} \int d^4x \sqrt{g(x)} \partial_\mu \phi_0^*(x) \partial_\nu \phi_0(x) \kappa \tilde{h}^{\mu\nu}(x); \quad (13)$$

$$S_2 = \frac{1}{2} \int d^4x \sqrt{g(x)} \left\{ m_\phi^2 |\tilde{\phi}(x)|^2 + m_\phi^2 [\phi_0^*(x) \tilde{\phi}(x) + \phi_0(x) \tilde{\phi}^*(x)] + [\partial_\mu \tilde{\phi}^*(x) \partial_\nu \tilde{\phi}(x) + \partial_\mu \phi_0^*(x) \partial_\nu \tilde{\phi}(x) + \partial_\mu \tilde{\phi}^*(x) \partial_\nu \phi_0(x)] g^{\mu\nu}(x) \right\} \quad (14)$$

For brevity, we have not expanded here \sqrt{g} and R like in eq. (9). Let us first look at the terms S_1 and S_2 . The term S_1 describes a process in which gravitons are produced by the source $\phi_0(x)$. The term S_2 contains the free action of the field $\tilde{\phi}(x)$, which describes the excitations of the condensate, and several vertices in which the graviton field $\tilde{h}_{\mu\nu}(x)$ and $\tilde{\phi}(x)$

interact between themselves and possibly with the source. All these interactions are not of special interest here and are generally very weak, due to the smallness of the coupling κ . The relevant point is that the purely gravitational cosmological term $\frac{\Lambda}{8\pi G}$ receives a "local" positive contribution $\frac{1}{2}\mu^2(x)$ which depends on the fixed external source $\phi_0(x)$. According to eq.s (9), (7) and to our discussion of the sign of Λ , this amounts to a negative mass contribution and could lead to instabilities.

Let us study the effect of such a local negative mass contribution in the simpler case of a scalar field in flat space. We consider a scalar field χ with very small mass m_χ and add to its free Euclidean action a source term of the form $-\frac{1}{2}\mu^2(x)\chi^2(x)$, which represents a "localized negative mass". The action becomes

$$S_\chi = \frac{1}{2} \int d^4x \left\{ [\partial^\mu \chi(x)][\partial_\mu \chi(x)] + m_\chi^2 \chi^2(x) - \mu^2(x)\chi^2(x) \right\} \quad (15)$$

To fix the ideas, let us suppose that $\mu^2(x)$ is different from zero in certain four-dimensional regions Ω_i , where it takes the constant values μ_i . Outside these regions let $\mu^2(x)$ go rapidly to zero. The solution of the classical field equation for χ is obtained by minimizing the action. In the absence of the source term we would have of course $\chi(x) = \text{const.} = 0$, because this minimizes at the same time the gradient term $\frac{1}{2}[\partial^\mu \chi(x)\partial_\mu \chi(x)]$ and the mass term $\frac{1}{2}m_\chi^2 \chi^2(x)$ (both positive-defined).

In the presence of the source term $-\frac{1}{2}\mu^2(x)\chi^2(x)$, the action decreases when χ^2 becomes large within the regions Ω_i . The growth of χ in these regions is limited only by the fact that outside them χ must go to zero (due to the term m_χ^2) and that the gradient in the transition region cannot in turn be too large. Let us suppose for instance that there is only one region Ω_0 , with the shape of a 4-sphere of radius r_0 . Let $\mu(x)$ take the value μ_0 inside Ω_0 and zero outside. We consider for χ the trial function $\chi(r) = \chi_0 f(r)$. The action becomes (we admit spherical symmetry)

$$S_\chi = \pi^2 \chi_0^2 \int_0^\infty dr r^3 \left\{ [f'(r)]^2 + m_\chi^2 [f(r)]^2 - \mu_0^2 \theta(r - r_0) [f(r)]^2 \right\} \quad (16)$$

We see that if the integral is positive, the value of χ_0 which minimizes the action is still $\chi_0 = 0$. On the contrary, if the integral is negative, the action is not bounded from below as χ_0 grows.

We choose the following explicit form of $f(r)$: for $r < r_0$ let $f(r) = 1$, i.e., $\chi(r) = \chi_0$; for

$r > r_0$ let be $f(r) = \exp[-(r - r_0)/\delta]$. We thus have

$$S_\chi = \pi^2 \chi_0^2 \left\{ \left(\frac{1}{\delta^2} + m_\chi^2 \right) \int_{r_0}^\infty dr r^3 e^{-2(r-r_0)/\delta} - \frac{1}{4} (\mu_0^2 - m_\chi^2) r_0^4 \right\} \quad (17)$$

$$= \pi^2 \chi_0^2 \left\{ \left(\frac{1}{\delta^2} + m_\chi^2 \right) \left(\frac{3}{8} \delta^4 + \frac{3}{4} r_0 \delta^3 + \frac{3}{4} r_0^2 \delta^2 + \frac{1}{2} r_0^3 \delta \right) - \frac{1}{4} (\mu_0^2 - m_\chi^2) r_0^4 \right\} \quad (18)$$

It is easy to check that for suitable choices of the parameters the expression within the braces in eq. (18) turns out to be negative. For instance, if m_χ can be disregarded with respect to μ_0 and $\delta \sim r_0$, the expression is negative provided $\mu_0 \gg \delta^{-1}$. If $\delta \ll r_0$ the expression is negative provided $\mu_0 \gg r_0^{-1}$, etc. Thus the system is unstable. For the graviton the instability is even worse, because the kinetic term $\tilde{R}^{(2)}$ is not positive-definite.

Physically, we might of course invoke some additional "regularizing" process which comes into play for large values of χ_0 and cuts the action. Thus the value of $\chi^2(x)$ inside the region is forced by the source to a certain maximum, and this will affect the propagation of the field. The differential equation for the propagator of $\chi(x)$ in the presence of the source must now satisfy additional boundary conditions on each region Ω_i . This amounts to a very complex mathematical problem; the physical consequence will be a "damping" of the correlations of $\chi(x)$. One can check this numerically in some model; we shall do this shortly in the next Section, referring to a toy bidimensional spin model.

3 The formula for the static potential energy.

In this Section we remind the formula which gives the static potential energy in Euclidean quantum gravity as a functional average [4, 5]. We illustrate its statistical meaning in the case of a weak field through an analogy with a simple 2D Ising model, which we also treat numerically with an elementary simulation. We show numerically that the insertion of local supplementary constraints due to an external field (in analogy to the singularities of the continuum case) damps in a sensible way the spin-spin correlations, and thus the interaction they represent in the model.

Let us first consider, in Euclidean quantum field theory, a scalar field $\Phi(x)$ with action $S_0[\Phi] = \int d^4x L(\Phi(x))$. In the presence of the external source $J(x)$, the ground state energy of

the system can be expressed as

$$\begin{aligned} E &= \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{\int d[\Phi] \exp \left\{ -\hbar^{-1} [\int d^4x L(\Phi(x)) + \int d^4x \Phi(x) J(x)] \right\}}{\int d[\Phi] \exp \left\{ -\hbar^{-1} \int d^4x L(\Phi(x)) \right\}} \\ &= \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \left\langle \exp \left\{ -\hbar^{-1} \int d^4x \Phi(x) J(x) \right\} \right\rangle, \end{aligned} \quad (19)$$

where it is assumed that the source vanishes outside the interval $(-T/2, T/2)$ and that the coupling between J and Φ is linear. More generally, a formula similar to (19) holds when we are dealing with more fields Φ_A and corresponding sources J^A , and when the coupling between fields and sources is not linear.

As an useful application of eq. (19), we can write the interaction energy $E(L)$ of two static pointlike sources of the field, kept at a fixed distance L . We just need to insert the suitable expression for J . The trajectories of the two sources are in flat space

$$x_1^\mu(t_1) = (t_1, 0, 0, 0); \quad x_2^\mu(t_2) = (t_2, L, 0, 0). \quad (20)$$

In ordinary gauge theories we may re-obtain in this way the Wilson formula for the static quark-antiquark potential. In quantum gravity we are led to the following equation for the static potential [4]

$$E(L) = \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{\int d[g] \exp \left\{ -\hbar^{-1} \left[S[g] + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{g_{\mu\nu}[x_i(t)] \dot{x}_i^\mu(t) \dot{x}_i^\nu(t)} \right] \right\}}{\int d[g] \exp \left\{ -\hbar^{-1} S[g] \right\}} \quad (21)$$

$$\equiv \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \left\langle \exp \left\{ -\hbar^{-1} \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} ds_i \right\} \right\rangle_S \quad (22)$$

where S is the euclidean action ³.

In eq.s (21), (22) the lines $x_1(t)$ and $x_2(t)$ must be parallel with respect to the dynamic metric $g_{\mu\nu}$ and thus they should in principle be re-traced for each field configuration of the functional integral. In practice, it is extremely difficult to compute a functional integral defined in such a formal way. Let us then limit ourselves to consider weak fluctuations of the gravitational field about flat space. The trajectories of the sources may be defined with respect to the background

³Notice that this formula applies also to two masses which are not pointlike ("pointlike particle" is actually an ill-defined concept in General Relativity), provided we can disregard the internal degrees of freedom. Namely, their action is still equal to $\sum_i \int ds_i$, where the integrals are taken along the trajectories of the centers of mass.

metric like in (20). It is straightforward to reproduce in this way the Newton potential energy [4, 5]; higher order corrections have been computed too [12]. The geodesic distance between the trajectories $x_1(t)$ and $x_2(t)$ is now equal to L only on the average; in fact such an approximation is not without physical meaning, since in any realistic source the fixed distance at which the two masses are kept can only be an average value. Also in the non-perturbative evaluations of eq. (22) in quantum Regge gravity [5], the distance L is evaluated *a posteriori* as the mean value of the geodesic distance on all configurations.

Let us consider the almost-flat metric $g_{\mu\nu} = \delta_{\mu\nu} + \eta_{\mu\nu}$ in a fixed gauge. We obtain

$$\int_{-T/2}^{T/2} ds_i \simeq \int_{-T/2}^{T/2} dt_i \sqrt{1 + h_{11}[x(t_i)]} \simeq T + \frac{1}{2} \int_{-T/2}^{T/2} dt_i h_{11}[x(t_i)] \quad (23)$$

and we see that to a first approximation the fluctuations of h have the effect of making each line "shorter" or "longer". Let us call $\alpha[h]$ the (gauge-invariant) difference between the lengths of the two lines in a field configuration h , and assume for simplicity that the masses of the two sources are equal: $m_1 = m_2 = m$. We may expand the exponential in eq. (22), finding (note that $\langle \alpha[h] \rangle$ obviously vanishes by symmetry)

$$E = 2m + \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \left\langle \exp\{-\hbar^{-1} m \alpha[g]\} \right\rangle = 2m + \lim_{T \rightarrow \infty} -\frac{m^2}{2\hbar T} \langle \alpha^2[g] \rangle + \dots \quad (24)$$

This equation exhibits an interesting relation between the vacuum fluctuations of the geometry and the static gravitational potential. To illustrate better its "statistical" meaning we would like now to introduce a toy analogy with the 2D Ising model.

Let us consider a planar spin system with periodic boundary conditions and the local coupling $H = -J \sum_{i,j} s_{ij} s_{i'j'} (J > 0; (i', j') \text{ neighbours of } (i, j))$. Let us then consider two columns j_1 and j_2 at a distance of L lattice spacings (see fig. 1).

We can regard this system as the analogue of a discretized configuration of a 4D gravitational field on the plane between the two parallel lines of eq. (20). The spin variables ± 1 represent fluctuations of the metric. At the transition temperature, the fluctuations of the spin variables along the two lines are correlated, approximately like $1/L$.

Going back to eq. (23) and making the correspondence $s \leftrightarrow h_{11}$, we see that the analogue of $\alpha[g]$ is the difference between spin sums taken along the columns j_1 and j_2 . The analogue of eq. (24) is

$$\langle \alpha_{Ising}^2 \rangle = \left\langle \left(\sum_i s_{ij_1} - \sum_i s_{ij_2} \right)^2 \right\rangle =$$

$$= \left\langle \left(\sum_i s_{ij_1} \right)^2 + \left(\sum_i s_{ij_2} \right)^2 - 2 \sum_i s_{ij_1} \sum_k s_{kj_2} \right\rangle \quad (25)$$

We are interested only in the term which depends on the distance L between the two columns, that is, the product term

$$p_{12} = \left\langle \sum_i s_{ij_1} \sum_k s_{kj_2} \right\rangle. \quad (26)$$




Fig. 1 - Spin sums taken along two columns of a 2D Ising system.

A numerical simulation with a simple 10×12 system has given, as expected, the following results at the critical temperature:

L (lattice spacings)		$p_{\{12\}}$	
=====			
2	1.15	+/-	0.13
3	0.74	+/-	0.09
4	0.48	+/-	0.09
5	0.42	+/-	0.08
6	0.41	+/-	0.08

We see that the L^{-1} law is approximately verified also for the correlation between the spin sums taken along the two columns. Only for $L = 6$ there is a deviation, which can be explained

as due to the periodic boundary conditions.

We observe that, due to the nature of the spin system, the correlation between two spin sums along j_1 and j_2 is necessarily positive. Thus, the part of $\langle \alpha_{Ising}^2 \rangle$ which depends on the distance L is negative. In perturbative quantum gravity one finds instead that the correlation $\langle h_{11}(x)h_{11}(y) \rangle$ is always negative, which leads to the correct negative sign for the potential energy in eq. (24).




Fig. 2 – Insertions of singular points "*" between the columns j_1 and j_2 .

We now introduce some supplementary conditions, in order to simulate the case in which the spin variables assume on certain sites a fixed value. This could be due, like in the gravitational case to which we are interested, to the localized action of an external field. In the spin model we may imagine that an external magnetic field localized on certain sites forces spin-flips. With reference to Fig. 2, let us suppose that the spin-flips occur on the sites marked with a star and placed between the two columns j_1 and j_2 whose correlation we are measuring. (The two columns are denoted by "x", while all the remaining sites are denoted by "0".) To prevent an

uncontrolled "driving" of the total magnetization, we associate to each spin-flip an opposite flip on a neighboring site. The flips occur at each Montecarlo step; since the mean frequency at which the regular sites are flipped during the simulation is 120 times smaller, the resulting effect is to force a zero at the "*" sites.

The precise positions of the sites at which the spin-flips happen are almost irrelevant; we find in all cases, as appears from the following table, a sensible diminution of the correlations between the spin sums taken along the columns j_1 and j_2 :

Number of "*" sites	$p_{\{12\}}$ for $L=5$
<hr/>	
3 (Fig. 2.a)	0.19 +/- 0.04
4 (Fig. 2.b)	0.13 +/- 0.03

It appears therefore that the insertion of variables which are driven by an external field damps the correlations in the system, and that this mechanism is of a quite general nature, although we are not able to give a precise analytical description yet.

4 Conclusive remarks.

We have investigated in this paper an unusual interaction mechanism between gravity and a macroscopic quantum system driven by external fields. This idea was originally suggested by a possible phenomenological application [13, 14], but the mechanism is interesting also from the purely theoretical point of view and deserves further numerical and analytical investigation.

We have shown that under certain conditions the gravitational field becomes unstable and may develop singularities, but we have not tried to find a physical regularization and to compute the effect of the regularized singularities yet. Simple physical analogies show however that they generally reduce the gravitational long-range correlations. Our next task will be the explicit estimation of the changes in the correlation functions in terms of the squared density $|\phi_0(x)|^2$ of the Bose condensate and of its squared gradient $[\partial_\mu \phi_0(x)]^* [\partial^\mu \phi_0(x)]$.

References

- [1] M.J.G. Veltman, in *Methods in field theory*, Les Houches Summer School 1975, ed.s R. Balian and J. Zinn-Justin, North-Holland, Amsterdam, 1976.
- [2] S. Weinberg, Rev. Mod. Phys. **61** (1989) 1; J. Greensite, Phys. Lett. **B 291** (1992) 405 and references.
- [3] H.W. Hamber and R.M. Williams, Nucl. Phys. **B 248** (1984) 392; **B 260** (1985) 747; Phys. Lett. **B 157** (1985) 368; Nucl. Phys. **B 269** (1986) 712. H.W. Hamber, in *Les Houches Summer School 1984, Session XLIII*, North-Holland, Amsterdam, 1986. H.W. Hamber, Phys. Rev. **D 45** (1992) 507; Nucl. Phys. **B 400** (1993) 347.
- [4] G. Modanese, Phys. Lett. **B 325** (1994) 354; Nucl. Phys. **B 434** (1995) 697; Riv. Nuovo Cim. **17**, n. 8 (1994).
- [5] H.W. Hamber and R.M. Williams, Nucl. Phys. **B 435** (1995) 361.
- [6] L. Garay, Int. J. Math. Phys. **A 10** (1995) 145.
- [7] C. Rovelli, L. Smolin, Nucl. Phys. **B 442** (1995) 593.
- [8] G. Modanese, Phys. Lett. **B 348** (1995) 51.
- [9] N.C. Tsamis and R.P. Woodard, Phys. Lett. **B 301** (1993) 351; Comm. Math. Phys. **162** (1994) 217; Ann. Phys. (N.Y.) **238** (1995) 1.
- [10] A.D. Dolgov, M.B. Einhorn, V.I. Zakharov, Phys. Rev. **D 52** (1995) 717.
- [11] H. Van Dam and M. Veltman, Nucl. Phys. **B 22** (1970) 397; L.H. Ford and H. Van Dam, Nucl. Phys. **B 169** (1980) 126.
- [12] I.J. Muzinich, S. Vokos, Phys. Rev. **D 52** (1995) 3472.
- [13] G. Modanese, *Theoretical analysis of a reported weak gravitational shielding effect*, report MPI-PhT/95-44, May 1995 (hep-th/9505094); *Updating the analysis of Tampere's weak gravitational shielding experiment*, report UTF-367/96, Jan 1996 (supr-con/9601001).
- [14] E. Podkletnov and R. Nieminen, Physica **C 203** (1992) 441; E. Podkletnov and A.D. Levit, *Gravitational shielding properties of composite bulk $YBa_2Cu_3O_{7-x}$ superconductor below*

70 K under electro-magnetic field, Tampere University of Technology report MSU-chem,
January 1995.

LETTER TO THE EDITOR

The ac magnetic response in type-II superconductors

Yoshihisa Enomoto[†] and Kazuma Okada[‡]

[†] Department of Physics, Nagoya Institute of Technology, Gokiso, Nagoya 466, Japan

[‡] Department of Applied Physics, Nagoya University, Nagoya 464-01, Japan

Received 13 May 1996, in final form 1 July 1996

Abstract. The ac response of type-II superconductors to an alternating magnetic field is numerically studied on the basis of the time-dependent Ginzburg–Landau equations. We examine the temperature dependence of the ac susceptibility associated with a small ac magnetic field in the absence of a bias dc field. It is shown that with increasing temperature the in-phase component of the fundamental susceptibility exhibits a step-like change from a negative constant value to zero, while the out-of-phase component of the fundamental susceptibility and the third-harmonic component have a peak at a certain temperature near the superconducting transition temperature. These results are in qualitative agreement with those of recent experiments on high- T_c superconductors.

Magnetization measurements using alternating fields have been widely employed in the study of type-II superconductors [1]. This is because in these experiments the effective time window can be easily changed by varying the frequency of the applied ac field. Thus, the study of the ac magnetic response of type-II superconductors provides direct information on flux dynamics in these materials. Recently, several authors have measured the ac susceptibility in high- T_c materials as functions of various physical parameters such as temperature, and the frequency and amplitude of the applied ac field [2–5].

Various models have been proposed to explain experimental data for the ac susceptibility of type-II superconductors, and especially of high- T_c materials [6–12]. Although these models have partially succeeded in explaining the ac magnetic response of type-II superconductors, they are at rather macroscopic and/or phenomenological levels, and thus are still incomplete. In this letter, to complement the previous study, we attack the above problem by using a different type of approach, that is, the numerical approach of the time-dependent Ginzburg–Landau (TDGL) equations [13, 14]. The main advantage offered by this computer simulation study is the ability to visualize the dynamical processes of magnetization and thus to directly obtain information on the dynamics of the magnetic flux structure without making any of the *ad hoc* electrodynamical assumptions used in previous models (e.g., the field dependence of the critical current in the critical state model). Here, performing numerical calculation of the TDGL equations, we examine the ac magnetic response of type-II superconductors to a small alternating magnetic field in the absence of the steady bias field. In particular, we discuss the temperature dependence of the ac susceptibility with the amplitude and frequency of the ac field being fixed.

The TDGL equations are composed of two partial differential ones for the complex order parameter $\psi(\mathbf{r}, t)$ and the vector potential $\mathbf{A}(\mathbf{r}, t)$ at time t and position \mathbf{r} [13–15]:

$$\frac{\hbar^2}{2mD} \left(\frac{\partial}{\partial t} + i \frac{e\phi}{\hbar} \right) \psi = -\frac{\delta F}{\delta \psi^*} + f(\mathbf{r}, t) \quad (1)$$




Figure 1. Time variation of the magnetization $M(t)$ for $T/T_c = 0.8, 0.91, 0.93$, and 0.97 .

$$\sigma \left(\frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} + \nabla \phi \right) = -\frac{\delta F}{\delta \mathbf{A}} \quad (2)$$

with the covariant time derivative $(\partial/\partial t + ie\phi/\hbar)$ and a scalar potential ϕ , where ψ^* denotes the complex conjugate of ψ . These equations are invariant under the local U(1) gauge transformation for ψ , \mathbf{A} , and ϕ . Here, D and σ are the diffusion constant and the conductivity, associated with the normal phase, respectively, and they have the relation [16]

$$\sigma = \frac{c^2 \xi^2}{48\pi \lambda^2} \frac{1}{D} \quad (3)$$

with the coherence length ξ and the magnetic penetration depth λ . The last term of the r.h.s. of equation (1), $f(\mathbf{r}, t)$, denotes the thermal noise with zero mean, i.e. $\langle\langle f \rangle\rangle = 0$, and the correlation

$$\langle\langle f^*(\mathbf{r}', t') f(\mathbf{r}, t) \rangle\rangle = 12\xi_0^{-4} t_0 k_B T \left(\frac{H_c(0)^2}{8\pi} \right)^{-1} \delta(\mathbf{r}' - \mathbf{r}) \delta(t' - t) \quad (4)$$




Figure 2. Temperature dependences of χ_1' , χ_1'' , and V_3 .

where $\langle\langle \dots \rangle\rangle$ denotes the ensemble average, ξ_0 is the coherence length at zero temperature, and $t_0 \equiv \pi\hbar/(96k_B T_c)$ with the superconducting transition temperature T_c at zero field. The Ginzburg–Landau (GL) free-energy functional $F[\psi, \mathbf{A}]$ is given by

$$F = \int d\mathbf{r} \left[\frac{1}{2m} |\mathbf{D}\psi|^2 + \alpha(T) |\psi|^2 + \frac{\beta}{2} |\psi|^4 + \frac{1}{8\pi} (\nabla \times \mathbf{A})^2 \right] \quad (5)$$

with the covariant derivative $\mathbf{D} \equiv -i\nabla - (e/c)\mathbf{A}$ and the local magnetic flux density $\mathbf{b}(\mathbf{r}, t) \equiv \nabla \times \mathbf{A}$. Under the assumption that $\alpha(T) = \alpha(0)(T/T_c - 1)$, and $\alpha(0)$ and β are positive constants in equation (5), the upper and lower critical fields are given by $H_{c2}(T)/H_{c2}(0) = 1 - T/T_c$ and $H_{c1}(T)/H_{c2}(0) = (\ln \kappa/(2\kappa^2))(1 - T/T_c)$, respectively, with the GL parameter κ . The other notation is conventional [16]. These equations are supplemented with boundary conditions, $\mathbf{D}\psi|_n = 0$ and $\nabla \times \mathbf{A}|_s = \mathbf{H}_e$, where the index n denotes the normal direction on the sample boundary and the index s denotes the sample boundary with an applied magnetic field \mathbf{H}_e .

We here consider a type-II superconductor in the x – y plane with a size $L_x \times L_y$. The




Figure 3. See facing page.




Figure 3. (Continued) Time variations of profiles of the z -component of the local magnetic flux density, b_z , in units of $H_{c2}(0)$ along the x -axis at $y = 32\xi_0$ for $T/T_c = 0.8$ (a), 0.91 (b), and 0.97 (c). Time goes from the left-hand column to the right-hand one for one cycle of the ac field.

sample is assumed to be infinite in the z -direction, and the problem is reduced to two dimensions neglecting all derivatives along z . The external ac magnetic field is applied to the sample as $\mathbf{H}_e = H(t)\hat{z}$ with $H(t) = H_{ac} \sin \omega t$ where H_{ac} and ω denote the amplitude and angular frequency of the ac field, respectively, and \hat{z} is the unit vector along the z -axis. In actual simulations the TDGL equations are transformed into the dimensionless discretized equations on a two-dimensional lattice by introducing link variables for the vector potential and the gauge fixing such that the scalar potential is set to zero. Since these procedures are the same as those in references [13, 14], we will not discuss the numerical procedures any further.

In the following simulations, we set $L_x = L_y = 64\xi_0$ and $\kappa = 2$. The amplitude and frequency of the ac field are fixed to be $H_{ac} = 0.05H_{c2}(0)$ and $\omega t_0/2\pi = 0.25 \times 10^{-3}$ (that is, the period of the ac field is $4000t_0$), respectively. We also take the lattice spacing and time step for numerical calculations to be $0.5\xi_0$ and $0.0125t_0$, respectively. These values are chosen for the computational reason of obtaining efficient results within our computer availability. As the initial state we choose the zero-field-cooling state.

In figure 1 the magnetization, $M(t)$, of the sample is plotted against time for various values of temperature. The magnetization $M(t)$ is defined as $4\pi M(t) = \langle B \rangle(t) - H(t)$, where the magnetic induction $\langle B \rangle(t)$ is obtained from the sample average of the z -component, $b_z(r, t)$, of the local magnetic flux density $\mathbf{b}(r, t)$. At $T = 0.8T_c$, the magnetization is sinusoidal in nature, according to the external ac field $H_{ac} \sin \omega t$. With increasing temperature, the magnetization deviates from the sinusoidal character with a

decrease of its magnitude, and simultaneously a phase-shift phenomenon occurs in the t - $M(t)$ curve. Further increase of temperature results in the large degree of phase shift. Indeed, at $T = 0.97T_c$, the magnetization becomes almost zero at times when the applied ac field is maximum (minimum). Note that even for no bulk pinning case, the phase-shift phenomena take place due to the sample boundary effect, as has been discussed in our previous work [17].

To give details of the ac magnetic response of the system, we study the ac susceptibility defined by the Fourier transformation of the magnetization $M(t)$:

$$M(t) = H_{ac} \sum_{n=1}^{\infty} (\chi'_n \sin n\omega t - \chi''_n \cos n\omega t) \quad (6)$$

where χ'_n and χ''_n ($n = 1, 2, \dots$) denote the in-phase and out-of-phase components of the n th-harmonic susceptibility, respectively, with the n th harmonics $V_n \equiv \sqrt{\chi'^2_n + \chi''^2_n}$. In calculating the ac susceptibility, simulation data during the first period have been discarded to avoid transient effects. In the present case we have numerically checked that only odd harmonics are generated [8]. In figure 2, χ'_1 , χ''_1 , and V_3 are plotted as functions of temperature; they have been often measured experimentally. It is shown that with increasing temperature, χ'_1 exhibits a step-like change from a negative constant value ($=-1/(4\pi)$ theoretically [6]) to zero, while χ''_1 initially rises from zero, goes through a maximum at $0.91T_c$ (called the peak temperature, denoted by T_p), and then returns to a small value near T_c . Note that $H_{c1}(T_p) \simeq 0.008 < H_{ac} = 0.05 < H_{c2}(T_p) = 0.09$ for $T_p = 0.91T_c$ in units of $H_{c2}(0)$. The third harmonic V_3 is also found to have a similar temperature dependence to χ''_1 with the peak temperature $T_p = 0.93T_c$. These results are qualitatively consistent with those of recent experiments on high- T_c superconductors [2–5]. At present it is unclear whether the slight difference between the peak temperature estimated from χ''_1 and that from V_3 is physically meaningful or not.

Now we relate the above macroscopic behaviour of the system to the spatio-temporal structure of the local magnetic flux. In figures 3(a)–3(c) the time evolution of the profile of the local magnetic flux density b_z along the x -axis is shown at $y = 32\xi_0$ for a complete cycle of the ac field for $T/T_c = 0.8$ (a), 0.91 (b), and 0.97 (c), respectively. In figure 4 the time evolution of the profile of the y -component of the current density is shown along the x -axis at $y = 32\xi_0$ for $T = 0.91T_c$ during one half of the period of the ac field, as well as b_z . The current density is given by $(c/4\pi)\nabla \times \nabla \times \mathbf{A}$ in units of $j_0 \equiv cH_{c2}(0)/(4\pi\xi_0)$. No definite magnetic vortex structures are observed in figures 3(a)–3(c). It is also found that the step in χ'_1 is due to the transition from near-perfect screening (figure 3(a)) to complete penetration ($\langle B \rangle(t) \simeq H(t)$) of the ac field impinging into the whole sample (figure 3(c)). Moreover, we have numerically checked that the smallest temperature at which penetrating magnetic flux (current) reaches to the centre of the sample is $0.90T_c$ in the present case. Thus, the peak in χ''_1 (and maybe the peak in V_3) corresponds to the first penetration of the flux (current) to the centre of the sample (figure 3(b) and figure 4). Although such an interpretation has been already proposed by several authors [7, 9, 10], this is the first simulation study to discuss the ac magnetic response of type-II superconductors from the point of view of the local magnetic flux dynamics without making any of the *ad hoc* electrodynamical assumptions used in previous models. Finally, we remark that the magnetic relaxation phenomenon can be seen in figure 3(b) and figure 4. Indeed, even when the external field changes to a decreasing stage from the initial ramp-up phase, the magnetic flux near the centre of the sample still increases for a while. An importance of this relaxation behaviour has recently been pointed out as one of possible causes for the frequency dependence of the ac susceptibility [3, 12]. Such magnetic relaxation effects on the ac susceptibility will be




Figure 4. Time variations of profiles of the y -component of the current density (dashed lines) in units of $j_0 = cH_{c2}(0)/(4\pi\xi_0)$ along the x -axis at $y = 32\xi_0$ for $T/T_c = 0.91$ during one half of the period of the ac field. Profiles of b_z are also shown for comparison, as solid lines.

discussed elsewhere.

In conclusion, we have studied the ac magnetic response of type-II superconductors to the alternating magnetic field on the basis of numerical calculation of the TDGL equations. In particular, the temperature dependence of the ac susceptibility in the absence of the dc field with the amplitude and frequency of the ac field being fixed has been discussed. We have found a step-like change in χ'_1 , and a peak in χ''_1 and also in the third harmonic V_3 at a certain temperature near T_c . These results are in qualitative agreement with recent experimental data on high- T_c superconductors. Moreover, we have briefly discussed the relationship between these macroscopic behaviours of the system and the spatio-temporal behaviour of the local magnetic flux.

However, we should mention that there can be a big distance between simulations and experiments. This is because there are the following faults in the present model. Firstly, in real systems the ac response is strongly affected by pinning and thermal fluctuations, which are not in the model. Secondly, we have neglected the z -dependence of the problem. The present results, therefore, apply only to films, not bulk materials where the entanglement of flux lines may be important. These points will be discussed in future work.

Nevertheless, the present approach based on computer simulations of TDGL equations has been found to be potentially rich for throwing novel light on the problem studied here. For instance, the present model is applicable for discussing the validity of assumptions used in macroscopic and/or phenomenological models. Since the present study is still in a primitive stage, detailed simulations are now under way to allow discussion of a quantitative comparison of simulation results with theoretical results and experimental data obtained by not only the ac method but also the Hall probe method [18]. Moreover, many interesting problems still remain open, such as the dependence of the ac response on various physical parameters (e.g., the amplitude and frequency of the ac field, the dc field, bulk pinning and the sample geometry), and also the universal behaviour of the ac susceptibility. These problems are also now under consideration.

The authors are grateful to Professor S Maekawa for a number of valuable discussions.

References

- [1] For a review, see
Hein R A, Francavilla T L and Liebenberg D H (ed) 1991 *Magnetic Susceptibility of Superconductors and Other Spin Systems* (New York: Plenum)
- [2] Ghatak S K, Mitra A and Sen D 1992 *Phys. Rev. B* **45** 951
- [3] Wolfus Y, Abulafia Y, Klein L, Larkin V A, Shaulov A, Yeshurun Y, Konczykowski M and Feigel'man M 1994 *Physica C* **224** 213
- [4] Ding S Y, Wang G Q, Yao X X, Peng H T, Peng Q Y and Zhou S H 1995 *Phys. Rev. B* **51** 9107
- [5] Kumaraswamy B V, Lal R and Narlikar A V 1996 *Phys. Rev. B* **53** 6759
- [6] Bean C P 1964 *Rev. Mod. Phys.* **36** 31
- [7] Chen D X, Nogues J and Rao K V 1989 *Cryogenics* **29** 800
- [8] Ji L, Sohn R H, Spalding G C, Lobb C J and Tinkham M 1989 *Phys. Rev. B* **40** 10936
- [9] Geshkenbein V B, Vinokur V M and Fehrenbacher R 1991 *Phys. Rev. B* **43** 3748
- [10] van der Beek C J, Geshkenbein V B and Vinokur V M 1993 *Phys. Rev. B* **48** 3393
- [11] Shatz S, Shaulov A and Yeshurun Y 1993 *Phys. Rev. B* **48** 13871
- [12] Prozorov R, Shaulov A, Wolfus Y and Yeshurun Y 1995 *Phys. Rev. B* **52** 12541
- [13] For a review, see
Enomoto Y, Kato R and Maekawa S 1993 *Studies of High Temperature Superconductors* vol 11, ed A V Narlikar (New York: Nova Science) p 309
- [14] Fram H, Ullah S and Dorsey A T 1991 *Phys. Rev. Lett.* **66** 3067
Liu F, Mondello M and Goldenfeld N 1991 *Phys. Rev. Lett.* **66** 3071

- Kato R, Enomoto Y and Maekawa S 1991 *Phys. Rev. B* **44** 6916; 1993 *Phys. Rev. B* **47** 8016
- [15] Schmid A 1969 *Phys. Rev.* **180** 527
- [16] Tinkham M 1975 *Introduction to Superconductivity* (New York: McGraw-Hill)
- [17] Enomoto Y, Ishikawa Y and Maekawa S 1996 *Physica C* **263** 21
- [18] Zeldov E, Majer D, Konczykowski M, Larkin A I, Vinokur V M, Geshkenbein V B, Chikumoto N and Shtrikman H 1995 *Europhys. Lett.* **30** 367

UTF-391-96

Nov 1996

Possible quantum gravity effects in a charged Bose condensate under variable e.m. field.

Giovanni Modanese¹

I.N.F.N. – Gruppo Collegato di Trento

Dipartimento di Fisica dell'Università

I-38050 Povo (TN) - Italy

and

John Schnurer

Director Applied Sciences

Physics Engineering

P.O. Box CN 446, Yellow Springs, Ohio 45387-0466 - U.S.A.

Abstract

In the weak field approximation to quantum gravity, a “local” positive cosmological term $\mu^2(x)$ corresponds to a local negative squared mass term in the Lagrangian and may thus induce instability and local pinning of the gravitational field. Such a term can be produced by the coupling to an external Bose condensate. In the functional integral, the local pinning acts as a constraint on the field configurations. We discuss this model in detail and apply it to a phenomenological analysis of recent experimental results.

04.20.-q Classical general relativity.

04.60.-m Quantum gravity.

74.72.-h High- T_c cuprates.

Key words: General Relativity, Quantum Gravity, Experimental Gravitation, Bose Condensates, High-Tc Superconductors.

¹e-mail: modanese@science.unitn.it

The behavior of a Bose condensate – or more specifically of the Bose condensate of Cooper pairs within superconducting materials – in an external gravitational field has been the subject of some study in the past [1]. The presence in a superconductor of currents flowing without any measurable resistance suggests that a superconductor could be used as a sensitive detection system, in particular for gravitational fields. The possible back-reaction of induced supercurrents on the gravitational field itself has been studied too, in analogy with the familiar treatment of the Meissner effect. As one would expect, it turns out that the “gravitational Meissner effect” is extremely weak: for instance, it has been computed that in a neutron star with a density of the order of 10^{17} kg/m^3 , the London penetration depth is ca. 12 km [2].

Such a penetration depth shows that the coupling of supercurrents to the classical gravitational field is extremely weak. The reason, of course, stems from the smallness of the coupling between gravity and the energy-momentum tensor of matter $T_{\mu\nu}$. One might wonder whether in a quantum theory of gravity – or at least in an approximation of the theory for weak fields – the Bose condensate of the Cooper pairs, due to its macroscopic quantum character, can play a more subtle role than as a simple contribution to the energy-momentum tensor.

In other words, we wonder if the macroscopic quantum coherence of the condensate can be taken into account at a fundamental level when computing the interaction between the superconductor and the external gravitational field. Results might then differ from those obtained from the gravitational coupling of “regular”, incoherent matter. A similar approach has been earlier proposed by Weber in his studies of gravitational waves and neutrino fluxes detection [3].

In a quantum-field representation the condensate is described by a field with non-vanishing vacuum expectation value ϕ_0 , possibly depending on the spacetime coordinate x . It is interesting to insert the action of this field, suitably “covariantized”, into the functional integral of gravity, expand the metric tensor $g_{\mu\nu}$ in weak field approximation, and check if the only effect of $\phi_0(x)$ is to produce gravity/condensate interaction vertices proportional to powers of $\kappa = \sqrt{16\pi G}$. One finds that in general this is not the case; the quadratic part of the gravitational action is modified as well, by receiving a negative definite contribution. It can thus be expected that the condensate induces localized instabilities of the gravitational field, in a sense which we shall describe precisely in Section 2.

The present paper is based on the letter [4] and originates in part from previous theoretical work [5] and in part from recent experimental results ([6]; see Sections 3, 4) which show the possibility of an anomalous interaction, in special conditions, between the gravitational field

and a superconductor. In Section 1 the theoretical model is introduced. In Section 1.1 we recall the role of a global cosmological term in the gravitational action. In Section 1.2 we show how a “local” positive cosmological term appears in the gravitational action due to the minimal coupling to a scalar field $\phi(x)$ with x -dependent vacuum expectation value $\phi_0(x)$, representing a Bose condensate with variable density. We define the “critical regions” of spacetime as those in which the overall sign of the cosmological term is positive. In Section 2 we discuss the gravitational instability induced by the condensate in the critical regions. We note that the “kinetic term” $R(x)$ in the gravitational lagrangian admits zero modes, and discuss qualitatively the effect of the instabilities in the Euclidean functional integral for the gravitational static potential energy [5]. In Section 3 we summarize the experimental findings reported in [6], trying to focus on the essential elements of an experimental setup which is quite complex. We add some remarks on the possible theoretical interpretations of these observations, according to the model with “anomalous” coupling between $h(x)$ and $\phi_0(x)$ introduced in the previous Sections. In Section 4 we describe our own experimental results and their interpretation. Finally, in Section 5 we discuss some issues of elementary character – but important from the practical point of view – concerning the overall energetic balance of the shielding process.

1 Bose condensate as “local” cosmological term in perturbative quantum gravity.

In this Section we show that a scalar field with a x -dependent vacuum expectation value, coupled to the gravitational field, gives a positive x -dependent contribution to the cosmological term. Our argument follows that given in [7].

1.1 Global cosmological term.

Let us consider the action of the gravitational field $g_{\mu\nu}(x)$ in its usual form:

$$S_g = \int d^4x \sqrt{g(x)} \left[\frac{\Lambda}{8\pi G} - \frac{1}{8\pi G} R(x) \right], \quad (1)$$

where $-R(x)/8\pi G$ is the Einstein term and $\Lambda/8\pi G$ is the cosmological term which generally can be present.

It is known that the coupling of the gravitational field with another field is formally obtained by “covariantizing” the action of the latter; this means that the contractions of the Lorentz

indices of the field must be effected through the metric $g_{\mu\nu}(x)$ or its inverse $g^{\mu\nu}(x)$ and that the ordinary derivatives are transformed into covariant derivatives by inserting the connection field. Moreover, the Minkowskian volume element d^4x is replaced by $d^4x \sqrt{g(x)}$, where $g(x)$ is the determinant of the metric. The insertion of the factor $\sqrt{g(x)}$ into the volume element has the effect that any additive constant in the Lagrangian contributes to the cosmological term $\Lambda/8\pi G$. For instance, let us consider a Bose condensate described by a scalar field $\phi(x) = \phi_0 + \hat{\phi}(x)$, where ϕ_0 is the vacuum expectation value and $m_\phi |\phi_0|^2$ represents the particles density of the ground state in the non-relativistic limit (compare eq.s (8)-(10)). The action of this field in the presence of gravity is

$$S_\phi = \frac{1}{2} \int d^4x \sqrt{g(x)} \left\{ [\partial_\mu \hat{\phi}(x)]^* [\partial_\nu \hat{\phi}(x)] g^{\mu\nu}(x) + m_\phi^2 |\hat{\phi}(x)|^2 + m_\phi^2 [\phi_0^* \hat{\phi}(x) + \hat{\phi}^*(x) \phi_0] + m_\phi^2 |\phi_0|^2 \right\}. \quad (2)$$

Considering the total action ($S_g + S_\phi$), it is easy to check that the “intrinsic” gravitational cosmological constant Λ receives a contribution $\frac{1}{2}m_\phi^2|\phi_0|^28\pi G$.

Astronomical observations impose a very low limit on the total cosmological term present in the action of the gravitational field. The presently accepted limit is on the order of $|\Lambda|G < 10^{-120}$, which means approximately for Λ itself, that $|\Lambda| < 10^{-54} \text{ cm}^{-2}$. (See [8] and references therein. We use natural units, in which $\hbar = c = 1$ and thus Λ has dimensions cm^{-2} , while G has dimensions cm^2 ; $G \sim L_{\text{Planck}}^2 \sim 10^{-66} \text{ cm}^2$). This absence of curvature in the large-scale universe rises a paradox, called “the cosmological constant problem” [9]. In fact, the Higgs fields of the standard model as well as the zero-point fluctuations of any quantum field, including the gravitational field itself, generate huge contributions to the cosmological term – which, somehow, appear to be “relaxed” to zero at macroscopic distances. In order to explain how this can occur, several models have been proposed [10]. No definitive and universally accepted solution of the paradox seems to be at hand, if only because that would require a complete non-perturbative treatment of gravity – which for the moment, at least, is not feasible [11].

A model in which the large scale vanishing of the effective cosmological constant has been reproduced through numerical simulations is the Euclidean quantum gravity on the Regge lattice [12]. From this model emerges a property that could turn out to be more general than the model itself: if we keep the fundamental length L_{Planck} in the theory, the vanishing of the effective cosmological constant $|\Lambda|$, depending on the energy scale p , follows a law of the form $|\Lambda|(p) \sim G^{-1}(L_{\text{Planck}} p)^\gamma$, where γ is a critical exponent [13, 14]. We find no reason to exclude that this behavior of the effective cosmological constant may be observed in certain

circumstances (see [7]). Furthermore, the model predicts that in the large distance limit, Λ goes to zero while keeping negative sign. Also, this property probably has a more general character, since the weak field approximation for the gravitational field is “stable” in the presence of an infinitesimal cosmological term with negative sign; conversely, it becomes unstable in the presence of a positive cosmological term (see [12] and Section 2).

1.2 Local cosmological term.

Summarizing, there appears to exist, independent of any model, a dynamical mechanism that “relaxes to zero” any contribution to the cosmological term. This makes the gravitational field insensitive to any constant term in the action of other fields coupled to it. Nevertheless, let us go back to the previously mentioned example of a Bose condensate described by a scalar field $\phi(x) = \phi_0 + \hat{\phi}(x)$. If the vacuum expectation value ϕ_0 is not constant but depends on the spacetime coordinate x , then a positive “local” cosmological term appears in the gravitational action S_g , and this can have important consequences. Let us suppose that $\phi_0(x)$ is fixed by external factors. Now, let us decompose the gravitational field $g_{\mu\nu}(x)$ as usual in the weak field approximation, that is, $g_{\mu\nu}(x) = \delta_{\mu\nu} + \kappa h_{\mu\nu}(x)$, where $\kappa = \sqrt{8\pi G} \sim L_{Planck}$. The total action of the system takes the form

$$S = \int d^4x \sqrt{g(x)} \left\{ \left[\frac{\Lambda}{8\pi G} + \frac{1}{2}\mu^2(x) \right] - \frac{1}{8\pi G} R(x) \right\} + S_{h\phi_0} + S_{\hat{\phi}}, \quad (3)$$

where

$$\frac{1}{2}\mu^2(x) = \frac{1}{2}[\partial_\mu\phi_0^*(x)][\partial^\mu\phi_0(x)] + \frac{1}{2}m_\phi^2|\phi_0(x)|^2; \quad (4)$$

$$S_{h\phi_0} = \frac{1}{2} \int d^4x \sqrt{g(x)} \partial^\mu\phi_0^*(x)\partial^\nu\phi_0(x)\kappa h_{\mu\nu}(x); \quad (5)$$

$$\begin{aligned} S_{\hat{\phi}} = & \frac{1}{2} \int d^4x \sqrt{g(x)} \left\{ m_\phi^2 |\hat{\phi}(x)|^2 + m_\phi^2 [\phi_0^*(x)\hat{\phi}(x) + \phi_0(x)\hat{\phi}^*(x)] + \right. \\ & \left. + [\partial_\mu\hat{\phi}^*(x)\partial_\nu\hat{\phi}(x) + \partial_\mu\phi_0^*(x)\partial_\nu\hat{\phi}(x) + \partial_\mu\hat{\phi}^*(x)\partial_\nu\phi_0(x)] g^{\mu\nu}(x) \right\}. \end{aligned} \quad (6)$$

In the action S , the terms $S_{h\phi_0}$ and $S_{\hat{\phi}}$ represent effects of secondary importance. The term $S_{h\phi_0}$ describes a process in which gravitons are produced by the “source” $\phi_0(x)$. The term $S_{\hat{\phi}}$ contains the free action of the field $\hat{\phi}(x)$, describing the excitations of the condensate and several vertices in which the graviton field $h_{\mu\nu}(x)$ and $\hat{\phi}(x)$ interact between themselves and possibly with the source. None of these interactions is of special interest here; they are generally very weak, due to the smallness of the coupling κ . The relevant point (eq.s (3), (4)) is that the

purely gravitational cosmological term $\frac{\Lambda}{8\pi G}$ receives a local positive contribution $\frac{1}{2}\mu^2(x)$ that depends on the external source $\phi_0(x)$.

We shall use “critical regions” to designate those regions of spacetime where the following condition is satisfied:

$$\left[\frac{\Lambda}{8\pi G} + \frac{1}{2}\mu^2(x) \right] > 0. \quad (7)$$

As we shall see in Section 2, in these regions the gravitational lagrangian is unstable and the field tends to be “pinned” at some fixed values.

It is important to give a numerical estimate of the magnitude order of $\mu^2(x)$ in the case of a superconductor. To this end we recall that the Hamiltonian of a scalar field ϕ of mass m_ϕ is given by

$$H = \frac{1}{2} \int d^3x \left\{ \left| \frac{\partial \phi(x)}{\partial t} \right|^2 + \sum_{i=1}^3 \left| \frac{\partial \phi(x)}{\partial x^i} \right|^2 + m_\phi^2 |\phi(x)|^2 \right\}. \quad (8)$$

In our case ϕ describes a system with a condensate and its vacuum expectation value is $\langle 0 | \phi(x) | 0 \rangle = \phi_0(x)$. Then in the Euclidean theory μ^2 is positive definite and we have

$$\langle 0 | H | 0 \rangle = \frac{1}{2} \int d^3x \mu^2(x). \quad (9)$$

Thus μ^2 is connected to the coherent vacuum energy density. A prudent estimate gives, for a superconductor

$$\mu^2 > 10^{24} \text{ cm}^{-4}. \quad (10)$$

As we saw in Section 1.1, a typical upper limit on the intrinsic cosmological constant observed at astronomical scale is $|\Lambda|G < 10^{-120}$, which means $|\Lambda|/8\pi G < 10^{12} \text{ cm}^{-4}$. This small value, compared with the above estimate for $\mu^2(x)$, supports our assumption that the total cosmological term can assume positive values in the superconductor, and that the criticality condition can be satisfied in some regions. But in fact the positive contribution of the condensate is so large that one could expect the formation of gravitational instabilities in any superconductor or superfluid, subjected to external e.m. fields or not – a conclusion in contrast with the observations.

According to our discussion in Section 1.1, we may then hypothesize that the value of $\Lambda/8\pi G$ at small scale is larger than that observed at astronomical scale, and negative in sign, such that it would represent a “threshold” of the order of $\sim 10^{30} \text{ cm}^{-4}$ for anomalous gravitational couplings. As we noted in Section 1.1, a negative intrinsic cosmological constant is allowed in models of quantum gravity containing a fundamental length. Given the small magnitude cited above, this threshold would not influence any other known physical process.

2 Gravitational instability and field “pinning” in the critical regions.

In this Section we discuss the gravitational instability induced by the condensate in the critical regions (7). We note that the “kinetic term” $R(x)$ in the gravitational lagrangian admits zero modes. Finally, we discuss qualitatively the effect of the instabilities in the Euclidean functional integral for the gravitational static potential energy [5].

Let us first consider the Einstein action with cosmological term (1) in the weak field approximation. We recall [15] that after the addition of a harmonic gauge-fixing, the quadratic part of the action takes the form

$$S_g^{(2)} = \int d^4x h_{\mu\nu}(x) V^{\mu\nu\alpha\beta} (-\partial^2 - \Lambda) h_{\alpha\beta}(x), \quad (11)$$

where $V^{\mu\nu\alpha\beta} = \delta^{\mu\alpha}\delta^{\nu\beta} + \delta^{\mu\beta}\delta^{\nu\alpha} - \delta^{\mu\nu}\delta^{\alpha\beta}$.

It is seen from (11) that in this approximation a cosmological term amounts to a mass term for the graviton, positive if $\Lambda < 0$ and negative (with consequent instability of the theory) if $\Lambda > 0$ [16]. Since in the presence of a condensate the sign of the total cosmological term depends on the coordinate x (eq.s (3)-(4)), we expect some “local” instabilities to form in the critical regions (7) in that case.

Such local instabilities were studied in [7], for the case of an Euclidean scalar field χ . Let us summarize very briefly the conclusions. If the squared mass term of the field χ is negative in some “critical” regions, then inside these regions χ tends to grow in order to minimize the action. However, this growth is limited by the gradient squared term in the lagrangian. One concludes that the field strength diverges in the critical regions only if the product of the imaginary mass by the size of the critical regions exceeds a certain constant.

In the case of perturbative quantum gravity, the situation is even more favourable to the formation of local instabilities, since the kinetic term $R(x)$ in Einstein action admits zero modes. Among these are not only the field configurations $h_{\mu\nu}(x)$ such that $R(x) = 0$ everywhere, but also those which satisfy Einstein equations

$$R_{\mu\nu}(x) - \frac{1}{2}g_{\mu\nu}(x)R(x) = -8\pi GT_{\mu\nu}(x), \quad (12)$$

with $T_{\mu\nu}$ obeying the condition

$$\int d^4x \sqrt{g(x)} \text{Tr } T(x) = 0. \quad (13)$$

(Note that for solutions of (12) one has $R(x) = 8\pi G \text{Tr } T(x)$).

Condition (13) can also be satisfied by energy-momentum tensors that are not identically zero, but only if they have a balance of negative and positive signs, such that their total integral is zero. Of course, they do not represent any acceptable physical source, but the corresponding solutions of (12) exist nonetheless, and are zero modes of the action. (One can consider, for instance, the static field produced by a “mass dipole”.) Clearly, there is no obstacle to the growth of the zero modes in the critical regions.

We shall admit that the growth of the gravitational field within the critical regions is limited by a positive higher order term in the action, that can be otherwise disregarded. The situation reminds that of a “ $\phi^4 - \phi^2$ ” model (see Fig. 1), with the important difference that in the present case the field depends on x and is “pinned” only locally, in the critical regions [17].

How does this local pinning of the field influences the gravitational interaction of two masses m_1 and m_2 at rest? We recall the general formula for the static potential in Euclidean quantum gravity [5, 13, 18]:

$$U(L) = \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{1}{Z} \int d[g] \exp \left\{ -\hbar^{-1} \left[S_g + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{g_{\mu\nu}[x_i(t)] \dot{x}_i^\mu(t) \dot{x}_i^\nu(t)} \right] \right\}, \quad (14)$$

where S_g is the gravitational action (1) and Z a normalization factor. The trajectories $x_1(t)$ and $x_2(t)$ of m_1 and m_2 are parallel with respect to the metric g . L is the distance between the trajectories, corresponding to the spatial distance of the two masses. (Of course, in the case we have in mind, m_1 should represent the Earth and m_2 a weighed sample, or conversely.)

Eq. (14) means that the interaction energy $U(L)$ of the two masses depends on the correlations between the values of the gravitational field on the “Wilson lines” $x_1(t)$ and $x_2(t)$. This dependence can be verified explicitly in the weak field approximation or through numerical simulations.

We can restate eq. (14) in the presence of the Bose condensate as

$$\begin{aligned} U[L, \mu^2(x)] &= \lim_{T \rightarrow \infty} -\frac{\hbar}{T} \log \frac{1}{Z} \int d[g] \int d[\hat{\phi}] \\ &\quad \exp \left\{ -\hbar^{-1} \left[S[g, \hat{\phi}, \phi_0] + \sum_{i=1,2} m_i \int_{-\frac{T}{2}}^{\frac{T}{2}} dt \sqrt{g_{\mu\nu}[x_i(t)] \dot{x}_i^\mu(t) \dot{x}_i^\nu(t)} \right] \right\}, \end{aligned} \quad (15)$$

where $S[g, \hat{\phi}, \phi_0]$ is the total action defined in (3). If there are critical regions between the two Wilson lines, the field correlations are modified and in general the field “pinning” reduces $|U|$ (see also [7]). A detailed computation is in progress.

3 Experimental findings.

A recent experiment [6] has shown an unexpected interaction between the gravitational field and a superconductor subjected to external e.m. fields. In this Section we summarize the reported observations, trying to focus on the essential elements of an experimental setup which is quite complex. We add some remarks on the possible theoretical interpretations of these observations, according to the model with “anomalous” coupling between $h(x)$ and $\phi_0(x)$ introduced in the previous Sections. Attempts at an independent replication of the experiment have been started recently [19].

3.1 Summary of results with rotating disks.

The core of the experimental apparatus is a toroidal disk, 27 *cm* in diameter and made of high critical temperature (HTC) superconducting material. The disk is kept at a temperature below 70 *K*; it levitates above three electromagnets and rotates (up to ca. 5000 *rpm*) due to the action of additional lateral magnetic fields. All electromagnets are supplied with variable-frequency AC current.

Within certain frequency ranges one observes a slight decrease in the weight of samples hung above the disk, up to a maximum “shielding” value of ca. 1% (observed when the disk rotates at ca. 5000 *rpm*). A smaller effect, of the order of 0.1% or less, is observed if the disk is only levitating but not rotating. The percentage of weight decrease is the same for samples of different masses and chemical compositions. One can thus describe the effect as a slight diminution of the gravity acceleration g_E above the disk.

This effect does not seem to diminish with increased elevation above the disk: there appears to be a “shielding cylinder” over the disk (compare also [21, 20]), that extends upwards for at least 3 meters. The resulting field configuration is clearly non conservative (Fig. 2). An horizontal force at the border of the shielding cylinder has occasionally been observed, but it is far too small to restore the usual “zero circuitation” property of the static field. No weight reduction is observed under the disk.

The disk has a composite structure: the upper part has been treated with a thermal process which partially melts the grains of the HTC material and makes it a good superconductor, with high T_c (ca. 92 *K*) and high J_c (ca. 7000 A/cm^2). On the contrary, the lower part remains more granular and has a lower critical temperature (ca. 65 *K*). The aim of this double structure is to obtain good levitation properties, while also creating a layer in which considerable resistive

effects can arise. Both characteristics appear to be necessary for the effect to take place: (1) the disk must be able to support intense super-currents; (2) a resistive layer must be present.

Another crucial feature of the experimental apparatus is the frequency spectrum of the applied e.m. field. Regardless of the reason for which the external e.m. field was originally employed, it seems clear to us that it plays a fundamental role in supplying the energy necessary for the apparent “absorption” of the gravitational field in the disk (compare energetic balance considerations in Section 5). Experimentally one observes [6] that the maximum shielding value is obtained when the coils are supplied with high frequency current (of the order of 10 MHz).

Transferring power to the disk is inherently not very efficient. This represents one of the most serious problems to overcome in order to achieve a stable shielding effect, especially for heavy samples. Otherwise, large amounts of refrigerating fluid will be needed to keep the disk from overheating and exiting the superconducting state. Remember that the maximum shielding values have been observed under conditions which seem to trigger the production of heat. This makes clear why a disk made of HTC material is important: although a low-temperature superconductor could also reach the critical density conditions, its specific heat is probably too small to maintain the essential conditions while allowing adequate power transfer.

3.2 Interpretation.

We have already stressed in our analysis [4] that an interpretation of the reported effect in the framework of General Relativity, as due to repulsive post-newtonian fields produced by the super-currents [22], is untenable, since the magnitude order of the effect is far too large [23].

Our interpretative model of the experimental results is based on the “anomalous” coupling between Bose condensate and gravitational field described in the previous Sections. In this model the essential ingredient for the shielding is the presence of strong variations of the Cooper pairs density in the disk: *we assume that such variations produce small regions with higher density, where the criticality condition (7) is satisfied*. It is thus necessary to understand, from this point of view, how the experimental parameters summarized above combine to produce such critical conditions.

A correct theoretical interpretation should also suggest alternative experimental settings and possibly lead to a simplified version of the experiment, still giving an observable shielding effect. Some steps toward such a partial replication have been taken by the authors (see below, Section 4, and the Appendix).

We recall that the HTC toroidal disk has a composite structure, with two layers as described in Section 3.1. At the operation temperature (slightly below 70 K) the supercurrents, which circulate without any resistance and take the high value $J_c \sim 7000 \text{ A/cm}^2$ in the upper part of the disk, are essentially zero in the lower part. The interface represents a sort of boundary for a fluid motion: the intensity of the supercurrents, which flow parallel to the interface, falls to zero at the interface itself (see Fig. 3).

For our purposes we can regard the supercurrents as a perfect fluid and apply the corresponding equations of fluid dynamics. (Phenomenological theories of superconductivity based on the two-fluids model are largely independent from the detailed microscopic mechanism which produces superconductivity and which is not well known yet in the case of HTCs.) The fluid density – i.e., the density of the Bose condensate – is subjected to local variations, or the fluid is compressible. In other words, while the *average* fraction of superconducting charge carriers in the material depends only on the ratio T/T_c and is small at temperatures about 70 K, usually not exceeding a few %, locally there can be for some reason a concentration of superconducting charge carriers, up to a maximum density fixed by the total charge density of the lattice.

Let now v represent the local velocity of the superconducting fluid, ρ its local density and m the mass of a superconducting charge carrier. If the gradient of T can be disregarded and the flow is irrotational (as usually assumed for superfluid motion) the following differential equation holds [24]

$$\operatorname{grad} \left(\frac{1}{2} v^2 + \frac{1}{m} V \right) = -\frac{T}{m} \operatorname{grad}(\log \rho), \quad (16)$$

where V is the external potential, which can be set to zero in the following, since it is almost constant over small distances and in the present case we apply eq. (16) to a very thin region above the interface.

Assuming that ρ and v depend only on the coordinate z , the solution of (16) is

$$\rho(z) = \rho_0 \exp \left[-\frac{1}{T} \left(\frac{1}{2} m v^2(z) \right) \right]. \quad (17)$$

In order to find ρ as a function of the distance z from the interface we need to know the dependence $v(z)$. Far above the interface, v takes an approximately constant value v_{max} . We shall assume that v vanishes at the interface, due to ohmic resistance, and that v grows, close to the interface, as a power of z . The behavior of $\rho(z)$ is then shown in Fig. 4.

It is seen that the condensate density ρ takes an higher value at the interface than far above it, in the bulk of the “good” superconducting material. This fact is rather intuitive and could have been predicted without use of the fluid equation. Given a “threshold” value ρ_c of the

density, there will be a thin layer $0 < z < z_c$ close to the interface in which $\rho_c < \rho(z) < \rho_0$. This layer corresponds to a critical region as defined in Section 1.2. Below the interface ($z < 0$), ρ vanishes. (In practice however, since the interface is not very clear-cut, the curve $\rho(z)$ will go smoothly to zero for $z < 0$.)

The relative velocity of the supercurrent with respect to the lower part of the disk is essentially determined by the rotation frequency of the disk [25], because the average velocity of the charge carriers in the supercurrent is on the order of 10^{-5} m/sec , while the tangential velocity due to rotation is much higher, on the order of 10 m/sec . Let us work out this estimate in more detail. The critical current density in the upper layer of the disk is $\sim 10^8 \text{ A/m}^2$. Taking a density of charge carriers of 10^{32} m^{-3} , with an elementary charge of $\sim 10^{-19} \text{ C}$ we find $v = J/\rho \sim 10^{-5} \text{ m/sec}$. On the other hand, when the disk is rotating at angular frequency $\omega \sim 5000 \text{ rpm} \sim 500 \text{ sec}^{-1}$, a point at distance 10 cm from the rotation axis has tangential velocity $v = \omega r \sim 50 \text{ m/sec}$.

We shall briefly discuss in Section 4.1 how conditions similar to those described above, which are present in “stationary form” at the interface in the rotating composite superconducting disk used in [6], are reproduced for short time intervals in our own experiment.

4 Our experiment.

One of us (J.S.) has recently succeeded in partially reproducing the weak gravitational shielding effect described in the previous Section. The experimental setup was suggested by the theoretical model outlined in Sections 1-3. It was also designed to eliminate, as far as is possible, any non-gravitational disturbance, and to show a precise temporal correspondence between actions taken on the HTC disk and the weight reduction of the sample/proof mass. Although the observed weight reduction was large enough to be clearly distinguished from the noise (of the order of 1 % or more), it was detected only in transient form, lasting up to ca. 3 s. This happened because the weight reduction was coincident with the superconducting transition of the HTC disk, which occurred quite rapidly when the disk warmed up over its critical temperature.

Our earlier experimental setup is described in the Appendix. An improved setup allowed us to run more than 400 trials with an heavier proof mass (a glass rods bundle – total weight 63 g), accumulating better statistics. Approximately 10 % of the trials gave positive result, i.e., a clear transient reduction in the weight of the proof mass. This apparently random behavior signals that some of the conditions which are necessary to trigger the effect are not well under

control yet. In particular, we found that the duration of the superconducting transition is very crucial: if the transition is too quick or too slow with respect to certain criteria, no effect is observed. The duration of the transition, in turn, depends strongly on the thermal conditions of the disk, and the latter can be controlled only with limited accuracy. A closer discussion of this point can be found in Section 4.1.

Our improved setup did not comprise the AC generator anymore. Thus error analysis is further simplified. One only needs to consider the parasitic effects of buoyancy and molecular diamagnetism. For both error sources, upper limits can be set according to the discussion in [6] and these limits are much smaller (by a factor 20 at least) than the observed effect.

As mentioned above, the transient reduction in the weight of the proof mass was always coincident with the thermal transition of the YBCO disk from the superconducting to the non-superconducting state. This was checked as follows.

A dewar flask with an inside diameter of about 3 cm and 14 cm deep allows to observe the experiment from the side by virtue of the fact that about 2.5 cm of the silvering, inside and out, has been removed when manufactured (see Fig. 5). The top of a hollow cylinder of polyethylene is flush with the bottom of the removed silvering. This cylinder supports, through two parallel bamboo sticks, a 2.5 cm by 1.3 cm , 0.3 cm thick samarium cobalt permanent magnet with an $MGOe$ factor of approximately 18. The magnet is even with the bottom of the removed silvering.

The dewar contains liquid nitrogen, as to half fill the clear observation area. This leaves about 1.3 cm of empty dewar with the permanent magnet at the bottom. After filling, all is allowed to cool down. A LEVHEX, almost-single-crystal, “pinning type” YBCO hexagon (2.0 cm from side to side) is tied to a cotton string and lowered into a second dewar to cool. When the LEVHEX is chilled, it is lowered into the first dewar. The levitation effect is pronounced and a piece of bamboo is used to push the hexagon down and leave it at approximately 0.6 cm above the magnet, as opposed to ca. 2.0 cm of its highest equilibrium position. The hexagon is well pinned and the familiar “tied to little springs” effect is well in evidence.

The detection system (see Fig. 6) is quite similar in principle to that described in the Appendix. Namely, a 33 cm by 51 cm glass plate on ring stand clamps is interposed between the dewar and a 63 g proof mass made of a hexagonal bundle of 7 rods, each 10 cm long and 0.6 cm in diameter. The string is tied to a dry 1 cm by 1 cm , 200 cm long wooden beam. The beam acts as a balance and has a 3 mm hole drilled in the middle. A 1 mm stainless steel pin in an aluminum stirrup serves as a pivot in the middle. The purpose of the beam is to place a

“Mettler 300” scale (300 g full scale, 0.01 g resolution) at least 180 cm from the dewar. The proof mass is directly over the levitating hexagon and the far end of the beam has an excess weight (aluminum blocks tied by cotton string) which rests on the scale. Once the whole is set up the scale is tared to leave a weight of 44.0 g . In order to eliminate any electrostatic artifact, an aluminum screen and a brass screen are taped to the glass plate placed between the dewar and the proof mass.

As the liquid nitrogen boils away the temperature of the LEVHEX rises. We see this evidenced by a slight reduction in the height of levitation at first, and, finally, there is no levitation at all. Exactly during this phase, which may last typically between 5 and 10 s in the different runs, we observe a transient reduction in the weight of the proof mass as indicated by the counterweight on the balance. In a “positive” run the measurement on the scale generally goes from 44.0 to 46-47 g and then returns to baseline after 2-3 s . This corresponds to a reduction in the apparent weight of the proof mass.

4.1 Interpretation.

A mixed state with locally-enhanced condensate density, similar to that present at the interface of the composite rotating disk (Section 3.2) forms, for a short time interval (a few seconds), also when a good-quality, single-phase HTC superconducting disk warms up above its critical temperature and goes from the superconducting to the normal state.

It is known that at the transition to the normal state some small non-superconducting regions first appear in the superconducting material, and then they gradually grow. Now, if strong circulating supercurrents had been previously induced in the material, these currents will rapidly decay at the transition. However, there will be during the transition a short intermediate phase in which part of the supercurrents still circulate, through those regions in which the material is still superconducting (Fig. 7). During this phase the flow of the supercurrents can still be described by a perfect fluid model like in Section 3.2. The non-superconducting regions act as boundaries for the flow and the resulting pattern for the condensate density will exhibit roughly similar features to those we found earlier: namely, the condensate density will be higher near the resistive obstacles, where the velocity must be smaller. Clearly if the superconducting transition is “too slow”, and the system is allowed to reach equilibrium, the critical density may not be achieved anywhere.

Unlike in the case of the rotating toroid, the whole bulk of the HTC disk is involved in

this case. It is crucial for the occurrence of the effect that the supercurrent present in the disk at the transition be very high and that the transition width ΔT_c be small. This requires an excellent HTC ceramic material.

The application of AC magnetic fields may allow some control on the transition rate. We are presently improving our earlier apparatus (see the Appendix) in this direction. The method described above, however, can be used to demonstrate the effect in a very clean way.

5 Energetic balance.

It is necessary now to discuss some issues of elementary character – but important from the practical point of view – concerning the overall energetic balance of the shielding process.

In general energy must be supplied in order to reduce the weight of an object, because the potential gravitational energy of the object has negative sign and is smaller, in absolute value, in the presence of shielding. However, since the field is non-conservative, it is certainly wrong to compute the difference in the potential energy of an object between the interior and the exterior of the shielding cylinder by evaluating naively the difference (which turns out to be huge) between an hypothetical “internal potential”

$$U = -\frac{GM_E(1-\alpha)}{R_E} = -Mg_ER_E(1-\alpha), \quad (18)$$

and an “external potential” $U = -Mg_ER_E$. (Here M is the mass of the object, R_E the Earth radius, M_E the Earth mass and α the “shielding factor”, $\alpha \sim 0.01$.)

Moreover, the gravitational fields with which we are most familiar, being produced by very large masses, are relatively insensitive to the presence of light test bodies; thus, it makes sense in that case to speak of a field in the usual meaning: while a body falls down, we do not usually worry about its reaction on the Earth. But in the present case, the interaction between the shielded object and the external source (that is, the system [Bose condensate+external e.m. field], which, by fixing the constraints on the gravitational potential h , produces the shielding), is very important [26].

Let us then ask a provocative question, suggested by the experimental reality: if the superconducting disk is in a room and the shielding effect extends up to the ceiling, should we expect the disk and all the shielding apparatus to feel a back reaction? (And possibly an even stronger one if the ceiling is quite thick or if there are more floors above?) The most reasonable answer is this: since the ceiling is very rigid, the experimental apparatus is not able to exert

any work on it and thus does not feel its presence.

To clarify this point, imagine that we hang over the superconducting disk a spring of elastic constant k and holding a body of mass M at rest. Then we operate on the disk with proper e.m. fields and produce the shielding effect with factor α . Now, the gravity acceleration over the disk becomes $g_E(1-\alpha)$. If the shielding effect is obtained quickly, the mass should begin to oscillate, otherwise it should rise by an height of $\Delta x = \alpha g_E M k^{-1}$, while remaining in equilibrium. In any case, since the kinetic energy and the potential energy of a harmonic oscillator in motion have the same mean value, it is legitimate to conclude that the work done by the shielding apparatus on the system [mass+spring] is on the order of

$$\Delta E \sim k(\Delta x)^2 \sim (\alpha g_E M)^2 k^{-1}. \quad (19)$$

This example shows that the work exerted by the apparatus on a sample being “shielded” will depend in general on the response of the sample itself. Only if the object being shielded can respond to the effect will work be done.

At this point we can estimate how much energy is needed to achieve critical density in one region of the condensate of cross section σ . If σ_{sample} is the projection of the sample on the disk, this energy is given by

$$\Delta E_\sigma = \frac{\sigma}{\sigma_{sample}} \Delta E \sim \frac{\sigma}{\sigma_{sample}} (\alpha g_E M)^2 k^{-1}. \quad (20)$$

This energy must be supplied by the external variable e.m. field.

In conclusion, we must expect in general an interaction between the partially shielded samples and the shielding apparatus. The energy needed to shield a sample depends on the mass of the sample itself and on the way it is constrained to move. In particular, we deduce from eq. (19) that if we want to detect the shielding effect by measuring the deformation of a spring (and do this with the minimum influence on the shielding apparatus) we should use, as far as is allowed by the sensitivity of the transducer, a spring with a high rigidity coefficient k .

Acknowledgments.

We are very grateful to E. Podkletnov for advice and encouragement. We are very much indebted to J.R. Gaines, Vice President of Superconductive Components, Columbus, Ohio, USA, who supplied the HTC disk.

References

- [1] D.K. Ross, J. Phys. **A** **16** (1983) 1331. J. Anandan, Class. Q. Grav. **11** (1994) A23 and references.
- [2] R.P. Lano, *Gravitational Meissner effect*, report U.ofIowa 96-4, March 1996, to appear in Phys. Rev. **D**.
- [3] J. Weber, Phys. Rev. **D** **38** (1988) 32; *Detection of weakly interacting particles by nearly perfect single crystals with high Debye temperatures*, in *Common Problems and Trends of Modern Physics*, p. 91, ed.s T. Bressani, S. Marcello and A. Zenoni (World Scientific, 1992). G. Preparata, Mod. Phys. Lett. **A** **5** (1990) 1; *QED coherence in matter* (World Scientific, Singapore, 1995).
- [4] G. Modanese, Europhys. Lett. **35** (1996) 413.
- [5] G. Modanese, Phys. Lett. **B** **325** (1994) 354; Nucl. Phys. **B** **434** (1995) 697.
- [6] E. Podkletnov and R. Nieminen, Physica **C** **203** (1992) 441. E. Podkletnov, *Weak gravitational shielding properties of composite bulk $YBa_2Cu_3O_{7-x}$ superconductor below 70 K under electro-magnetic field*, report MSU-chem 95, (cond-mat/9701074).
- [7] G. Modanese, Phys. Rev. **D** **54** (1996) 5002.
- [8] For recent data and references see L.A. Kofman, N.Y. Gnedin and N.A. Bahcall, Astrophys. J. **413** (1993) 1; L.M. Krauss and M.S. Turner, Gen. Rel. Grav. **27** (1995) 1137.
- [9] S. Weinberg, Rev. Mod. Phys. **61** (1989) 1; *Theories of the Cosmological Constant*, report UTTG-10-96 (astro-ph/9610044).
- [10] See, for instance, J. Greensite, Phys. Lett. **B** **291** (1992) 405 and references.
- [11] Nevertheless, some authors argue that the relaxation process could be inherently infrared and thus unaffected by the bad ultraviolet behavior of quantum gravity. See N.C. Tsamis and R.P. Woodard, Commun. Math. Phys. **162** (1994) 217; Ann. Phys. **238** (1995) 1.
- [12] H.W. Hamber and R.M. Williams, Nucl. Phys. **B** **248** (1984) 392; **B** **260** (1985) 747; Phys. Lett. **B** **157** (1985) 368; Nucl. Phys. **B** **269** (1986) 712. H.W. Hamber, in *Les Houches Summer School 1984, Session XLIII* (North-Holland, Amsterdam, 1986). H.W. Hamber, Phys. Rev. **D** **45** (1992) 507; Nucl. Phys. **B** **400** (1993) 347.

- [13] H.W. Hamber and R.M. Williams, Nucl. Phys. **B** **435** (1995) 361.
- [14] G. Modanese, Phys. Lett. **B** **348** (1995) 51.
- [15] M.J.G. Veltman, *Methods in field theory*, Les Houches Summer School 1975, ed.s R. Balian and J. Zinn-Justin (North-Holland, Amsterdam, 1976); E. Alvarez, Rev. Mod. Phys. **61** (1989) 561.
- [16] There are some possible objections to this naive statement. First, it is known that there is no consistent gravitational field theory including a massive graviton [H. Van Dam and M. Veltman, Nucl. Phys. **B** **22** (1970) 397; L.H. Ford and H. Van Dam, Nucl. Phys. **B** **169** (1980) 126. For a recent discussion see M. Visser, *Mass for the graviton*, gr-qc/9705051]. Moreover, when a cosmological term is present in the gravitational lagrangian, it is not correct to consider the weak field expansion around a flat background and one should in principle expand around a De Sitter or anti-De Sitter background [11]. In our case, however, we consider only localized modifications to the lagrangian. Thus neither objection is relevant.
- [17] Short-distance regulators or higher-order terms, analogous to a “ ϕ^4 ” term in the scalar field case, are present in several models of quantum gravity [10,12]. Note that the size of our hypothesized critical regions can be regarded as macroscopic with respect to the scale where conformal fluctuations of the metric or R^2 -terms become important. For this reason, we believe that an approach based upon the Euclidean weak field approximation + regulator is appropriate.
- [18] I.J. Muzinich, S. Vokos, Phys. Rev. **D** **52** (1995) 3472.
- [19] Ning Li, D. Noever, T. Robertson, R. Koczor and W. Brantley, Physica **C** **281** (1997) 260.
- [20] C.S. Unnikrishnan, Physica **C** **266** (1996) 133.
- [21] G. Modanese, *Updating the theoretical analysis of the weak gravitational shielding experiment*, report UTF-367/96.
- [22] D.G. Torr and Ning Li, Found. Phys. Lett. **6** (1993) 371.
- [23] The work [20] shows that the contribution from the super-currents to the static component g_{00} of the post-newtonian gravitational field over the disk is not only much smaller than the

observed effect (by several magnitude orders), but it is attractive like the newtonian field of the Earth. Even taking into account perturbative quantum corrections to the Newton potential one reaches the same negative conclusion [4].

- [24] K. Huang, *Statistical Mechanics*, Ch. 5 (J. Wiley, New York, 1963).
- [25] We assume that the superconducting charge carriers are not able to follow the rotation of the disk, or only partially: those very close to the interface co-rotate with the disk, those farther away follow slower, with a relative velocity v_{max} . This behavior is also confirmed by the observations relating to the “braking phase”, during which the rotation frequency of the disk is suddenly reduced [6].
- [26] Because of this, considerations involving the local energy density of the gravitational field, which can be properly defined for weak fields, are not helpful in the present case.

Appendix: earlier setup.

Our first experimental setup consisted of (a) a hexagon-shaped YBCO 1" (2.5 cm) superconducting disk, 6 mm thick; (b) a magnetic field generator producing a $600\text{ gauss}/60\text{ Hz}$ e.m. field; (c) a beam balance with suspended sample.

The beam is made of bamboo, without any metal part, coming to a point on one end (24.6 cm long, weight 1.865 g). The sample is made as follows. A cardboard rectangle (16 mm by 10 mm by 0.13 mm) is suspended from the balance with 2.8 cm of cotton string. A polystyrene "pan" (7.2 cm by 8.7 cm by 1.7 mm) is attached with paper masking tape to the cardboard rectangle. The total sample assembly (with string, cardboard, tape) weighs 1.650 g .

The balance is suspended from the end of a 150 cm wood crossbeam by ca. 30 cm of monofilament fishing line attached to the balance's center of mass (5.5 cm from the end where the sample is attached). The other end of the crossbeam is firmly anchored by a heavy steel tripod. Thermal and e.m. isolation is provided by a glass plate (15 cm by 30 cm , 0.7 cm thick) with a brass screen attachment. This plate-and-brass-screen assembly is held about 4.5 cm below the sample by a "3-finger" ring stand clamp. A straightsided, 14 cm diameter, 25 cm deep dewar with ca. 10 cm of liquid nitrogen is used to cool the superconducting disk below its critical temperature, and is removed from the experiment area before the trial.

The experimental procedure comprises the following steps.

1. The YBCO superconductor is placed in a liquid nitrogen bath and allowed to come to liquid nitrogen temperature (as indicated when the boiling of the liquid nitrogen ceases). The superconductor will remain below its critical temperature (about 90 K) for the duration of the trial (less than 20 seconds).
2. The disk is then removed from the bath and placed on a strong NdFeB magnet to induce a supercurrent. The Meissner effect is counteracted by a wooden stick. The superconducting disk has a cotton string attached to it to assist handling.
3. The disk and wooden stick assembly is placed on the AC field generator, about 33 cm below the isolation plate and about 40 cm below the sample. The AC field generator is then cycled for ca. 5 seconds with 0.75 sec equal-time on/off pulses. Prior to a run the sample is centered to be over the middle of where the disk will finally be, on the AC field generator.

One observes that while AC current is flowing through the generator the balance pointer dips 2.1 mm downward. When the AC field generator is pulsed with no superconductor present, there is no measurable pointer deflection. Also air flows do not cause any measurable deflection. The whole procedure is well reproducible.

The weight difference required to raise the sample by 2.1 mm was then found to be 0.089 g . This was measured taking advantage of the fact that the suspension wire produces a small torque on the balance beam toward the equilibrium position: the balance pointer was found to raise 2.1 mm upward when a weight of 0.089 g was placed above the sample.

FIGURE CAPTIONS

1. Typical double-well potential of a $(\phi^4 - \phi^2)$ field system.

The curve A represents a familiar “ ϕ^2 ” potential, corresponding to a field with positive squared mass term. The curve B represents the same “ ϕ^2 ” potential, to which a “ ϕ^4 ” term has been added. The lowest energy excitations are nearly the same for A and B. In C, the sign of the “ ϕ^2 ” term has been reversed ($(\phi^4 - \phi^2)$ potential). Now, in the lowest excitations the field is pinned to finite values distinct from zero.

2. Non-conservative character of the “modified” field above the superconducting disk.

The ”modified field” described by Podkletnov, with an infinite cylinder-like shielding zone over the disk, is clearly non-conservative. This means that if a proof mass goes along a closed path (up within the shielding zone, and down outside), the gravitational field exerts a net positive work on it.

From the formal point of view, this means that the observed static field is not the gradient of a gravitational potential, even though the Einstein equation for the connection ($\text{div}\Gamma=0$) still holds true.

Physically, this property is due to the (“anomalous”) coupling of the gravitational field with the external e.m. field, coupling that is mediated by the Bose condensate in the superconducting disk. The mechanical energy that a proof mass gains when it goes along the closed circuit is furnished by the external e.m. field.

3. Schematic diagram representing the velocity distribution of the superconducting charge carriers in the upper layer of the toroidal disk (B) during rotation.

The lower layer (A) is not superconducting at the temperature of operation (slightly below 70 K). The *relative* velocity of the superconducting charge carriers with respect to the interface between the two layers is represented by thick arrows. It grows from zero to some velocity v_{max} according to a power law (compare Fig. 4).

This is a side view. In stationary conditions, the relative velocity has no component along the vertical direction z . We suppose that the disk is rotating clockwise, so the short arrow means in reality that the superconducting charge carriers near the interface are almost co-rotating with the disk; the carriers farther away from the interface “follow slower”, thus with larger relative velocity, up to v_{max} .

4. Variation of the condensate density in the upper layer of the disk as a function of the distance from the interface, according to the perfect-fluid model.

For the definition of the coordinate z and a qualitative description of the velocity pattern compare Fig. 3.

The curve $v(z)/v_{max}$ shows for illustration purposes a quasi-linear growth of the velocity of the superconducting charge carriers with increasing distance from the interface, followed by saturation to the value v_{max} . The curves A, B, C represent the behavior of the local condensate density $2\rho/\rho_0$ (the factor 2 is for graphical reasons) according to eq. (17), for the following specific cases:

$$A: v(z)/v_{max} \sim z^{1/2}$$

$$B: v(z)/v_{max} \sim z$$

$$C: v(z)/v_{max} \sim z^{3/2}$$

It is seen that the condensate density achieves its maximum value for $v = 0$ and its minimum for $v = v_{max}$.

The line D represents the threshold condensate density ρ_c necessary for anomalous coupling. Z_c represents the thickness of the critical region of the condensate if the velocity depends on z like in A (the generalization to the cases B and C is obvious). This means that in the thin layer $0 < z < z_c$ close to the interface one has $\rho_c < \rho(z) < \rho_0$, or the layer is a “critical region”. Note that the value of the ratio ρ_c/ρ_0 as depicted above and the z -scale are just for illustration purposes.

5. Scheme of our [magnet+LEVHEX] setup.

A: Dewar flask (height 14 cm, inside diameter 3 cm).

B: Part of the dewar (height 2.5 cm) where the silvering has been removed to allow side observation of the levitating YBCO hexagon (E).

C: Polyetilene tube supporting the magnet (D) through two parallel bamboo sticks.

D: samarium cobalt permanent magnet (2.5 cm by 1.3 cm, 0.3 cm thick).

E: LEVHEX “pinning type” YBCO hexagon (2.0 cm from side to side).

6. *Detection system for the demonstration experiment of transient shielding effect at the superconducting transition.*

A: Dewar flask (see details in Fig. 5) with LEVHEX levitating above magnet.

B: Glass screen, with brass and aluminum foils, to shield proof mass from air flows and electrostatic fields.

C: Proof mass: bundle of 7 glass rods, each 10 cm long and 0.6 cm in diameter; total weight 63 g.


D: Wooden balance beam, 1x1x200 cm.

E: Mettler 300 balance, 0.01 g resolution.

7. *Velocity and density pattern of the superconducting charge carriers in a bulk disk at the transition (dark: critical regions).*

At the transition to the normal state some small non-superconducting regions first appear in the superconducting material, and then they gradually grow. If strong circulating supercurrents had been previously induced in the material (top), these currents will rapidly decay at the transition. However, there will be during the transition a short intermediate phase in which part of the supercurrents still circulate, through those regions in which the material is still superconducting (bottom).

During this phase the flow of the supercurrents can still be described by a perfect fluid model like in Section 3.2. The non-superconducting regions act as boundaries for the flow and the resulting pattern of the condensate density will exhibit roughly similar features to those we found earlier: namely, the condensate density will be higher near the resistive obstacles, where the velocity must be smaller.



Engineering Analysis of the Podkletnov Gravity Shielding Experiment

[Peter L. Skeggs](#)

7 Nov 1997

Abstract

Eugene Podkletnov's recent work [1] involving the supposed shielding of gravity by means of a rotating superconductor in a high frequency magnetic field has attracted much attention. At least one group, Delta G at NASA Marshall Space Flight Center and University of Alabama Huntsville, is currently working on a reproduction of his experiment. However, this is quite difficult due to the lack of engineering detail. Here, the author pulls together important information from the text and a figure of Podkletnov's paper, then makes estimates of the magnetic field configuration of the coils used to levitate the superconducting disk via the Meissner effect. An upper limit magnetic flux density of 850 gauss near the poles of the coils is calculated. Finally, this estimate for the magnetic flux density is used to calculate the expected acceleration on a test sample due to the gravitomagnetic and gravitoelectric fields postulated by Li and Torr [6] -- at most 145×10^{-6} cm/s². Although not large enough to explain Podkletnov's results, it should be detectable by NASA's gravimeter.

Summary of Salient Points

1. Section 1, second paragraph: "a new installation was built... magnetic fields up to 2 T and frequencies up to 10^8 Hz..."
1. Section 2, first paragraph: "The shielding superconducting element was made of dense, bulk, almost single-phase YBCO and had the shape of a toroidal disk with an outer diameter of 275 mm, an inner diameter of 80 mm, and a thickness of 10 mm... A thin (1 mm) foil of magnetic material was attached (without electrical contact) to the upper surface of the disk, using hot-melt adhesive, to facilitate rotating the disk as described below, especially at the initial stages of rotation."
2. Section 3, first paragraph: "The toroidal disk was placed inside a cryostat equipped with a set of three coils (Fig. 3) that could keep it levitating when it reached the superconducting state."
3. Section 3, second paragraph: "High-frequency electric current (10^5 Hz) was first sent to the two main solenoids around the disk to initiate an internal current in the ceramics while the disk was still at room temperature." He cools the disk, then "the main solenoids were switched off. After this, the high-frequency current [*implying there is just one high frequency current in question, which is 10^5 Hz*] was sent to the coils below the disk, and the superconductor raised up (at least 15 mm..."
4. Section 3, third paragraph: "Then a small current (10^5 Hz) was sent to the main solenoids, causing the disk to begin rotating counter-clockwise with increasing speed. The rotation speed was increased up to 5000 rpm. At this point the current in the rotating coils was of the order of 8-10 A. (The diameter of the wire of these coils is 1.2 mm). This current was supplied by powerful high-frequency generators usually employed for induction heating and quenching of metals."

Wire Size Analysis

At first, it seems that 1.2 mm diameter wire would be too small to carry such large currents. A wire table [2] was consulted. A 1.2 mm diameter wire, which is the same as a diameter of 47.2 mils or around 16 gauge, is recommended to carry at most 1.7 A. Of course, this is a conservative limit to prevent heat buildup and resistive losses.

Another data point is what the NEC [3] recommends for 14 gauge wire; it's allowed for circuits carrying 15 A, 60 Hz, at 115 VAC in residential wiring applications. This would be equivalent to 9 A for 16 gauge wire. So perhaps 1.2 mm is not unreasonable.

A third data point is to calculate the skin depth [4] of copper wire at 10^5 Hz:

$$d_p = (2 / (w * u * s))^{0.5} \quad (1)$$

where $w = \omega = 2\pi \times 10^5$, $u = \mu_0 = \text{magnetic permeability of copper} = 4\pi \times 10^{-7}$, and $s = \sigma = \text{conductivity of copper} = 5.8 \times 10^7 \text{ mho/m}$. This gives $d_p = 0.21 \text{ mm}$. A rule of thumb is to use a conductor that is 5 times the skin depth, so a 1.05 mm wire should suffice. Because there is an inversely proportional relationship between frequency and skin depth, this is also acceptable up to the maximum operating frequency of 10^8 Hz.

Analysis of Levitation Coils

Unfortunately, Podkletnov's text gives no data as to how many turns of wire are in each coil, nor the dimensions of the coil, so a calculation of it's inductance or magnetic field strength seems difficult to do. However, if one assumes that fig. 3 of his paper is accurate, then dimensions of the coils can be inferred. Giovanni Modanese has suggested as much to the author previously [\[5\]](#).




Figure 3 from Podkletnov's cond-mat/9701074.

Fig. 3 shows that the 3 coils are actually each made of 2 coils connected to a roughly U-shaped core which positions one coil at the outer diameter of the disk and the other coil at the inner diameter. This forms a magnetic circuit that one might suppose has one pole (say the north) emitting flux at the outside diameter and the other pole collecting flux at the inner diameter; the poles switch direction every half cycle of applied AC. Because the flux would be partially or mostly expelled from the superconductor, the flux lines would travel radially along the bottom face of the toroid.

Since the outer diameter of the toroid is 275 mm, and the inner is 80 mm, I estimate (by measuring a printout of fig. 3, where 1 mm on paper = 6.25 mm in real life) the centers of the two coils in each of the 3 supporting solenoid structures to be 97.5 mm apart. Each coil body is 40.6 mm in diameter on the outside, and perhaps 12.5 mm inside (the diameter of the vertical piece of the magnetic core material). Each coil body is also ~25 mm tall.


From these dimensions, plus the diameter of the wire, one can estimate the number of turns. The cross-sectional area of one half of a coil is $(40.6/2 - 12.5/2) * 25 = 351 \text{ mm}^2$. The cross-sectional area of a 1.2 mm wire is 1.13 mm^2 . So, assuming optimal packing, there could be at most 310 turns in each coil.

The current density through one half of the coil is:

$$J = I * n / A \quad (2)$$


where I is the current equal to 10 A, n is the number of turns, and A is the cross sectional area, resulting in a current density $J = 8.83 \text{ A/mm}^2$, or $8.83 * 10^6 \text{ A/m}^2$. This number was entered into QuickField (a finite element analysis program for electric and magnetic fields), assuming that the current points in to the page for the left half of the left coil, out for the right half of the left coil, out for the left half of the right coil, and in for the right half of the right coil (so the two B fields generated are of opposite direction), the core is steel with a relative permeability of 5000, for the geometry consisting of 2 coils and a connecting bar of core material, and using dimensions of each part estimated from Podkletnov's fig. 3.

The program found that the B field generated 15 mm above a pole is a maximum of about 400 gauss, with H about 32700 A/m; even immediately adjacent to the core material at the top center of a coil, the maximum B is 850 gauss, with H about 67900 A/m. Note that these results are using QuickField in the magnetostatic or DC condition.



Magnetic Flux Lines around 1/2 of toroidal disk created by one solenoid assembly. Note the expulsion of flux from the superconductor.

Flux Density
B (T)



Map of magnetic flux density B around the assembly. Note the zero B field inside the superconductor.

Gravitomagnetic Force Estimate

Note that the magnetic flux density applied across the radius of the superconductor is above the lower critical field (B_c) for YBCO which is ~ 60 gauss. This implies that flux vortices should be formed which run horizontally through the radius of the disk. This also implies using the Li/Torr theory [6] for the generation of a gravitomagnetic field by the coherent rotation of lattice ions, that a B_g field with a radial component should be generated. Assuming that this is large outside the disk (which is highly speculative), and that there exists a gravitational analog of the Lorentz force [7]

$$F = m * (E_g + v \times B_g) \quad (3)$$

where m is the mass of a test sample, E_g is the normal gravitational pull of the Earth plus possibly an induced gravitoelectric field, v is the velocity of the test sample relative to the superconductor, and B_g is the gravitomagnetic field generated by the rotating ions. This should generate a vertically oriented force on a test sample, since v points in the direction of a tangent to the circumference of the disk, B_g points radially, and thus their cross product points vertically. This is exactly what Podkletnov has reported: the vertical change in apparent weight of an object.

The problem with this explanation is the magnitude of B_g . Li/Torr predict a value (eq. 3.8, p. 381) that is approximately

$$B_g = 10^{-9} B_o \quad (4)$$

where B_o is the applied magnetic field. The velocity v at the outer radius of the 275 mm disk rotating at 5000 rpm, is 7200 cm / s. With B_o of 0.24 Tesla (400 gauss for each of 6 coils), B_g is $2.4 * 10^{-10}$ / s. This results in an acceleration term due to the gravitomagnetic field (the right hand inner term of eq. 3) of only $1.7 * 10^{-6}$ cm / s², compared to the acceleration due to gravity of the Earth of approximately 980 cm / s². So, assuming the 10^{-9} coupling term is accurate, this model does not explain Podkletnov's claimed 0.05% to 2.1% loss of weight in his samples. However, the magnitude of this change of $1.7 * 10^{-6}$ cm / s² is well within the measurement capability of NASA's gravimeter [8], so there is hope of detecting it.

One could also estimate the magnitude of a directly induced gravitoelectric field (Li/Torr eq. 3.7, p. 381), under the assumption that the vector points in the proper direction. Its maximum instantaneous value is given by

$$E_g^{ind} = 10^{-9} dA_0 / dt = (10^{-9}) * (6 * 10^{-5} \text{ Wb/m}) * (2\pi * 3.8 * 10^6) = 143 * 10^{-6} \text{ cm/s}^2 = 0.143 \text{ mGal} \quad (5)$$

where the peak magnitude of the vector potential $A = 6 * 10^{-5}$ Wb/m was estimated using QuickField in the harmonic magnetic mode, and the term dA_0 / dt is calculated using:

$$d(A_{peak} \sin(w t)) / dt = A_{peak} * w \cos(w t) \quad (6)$$

whose maximum value is simply $A_{peak} * w$, and for which an operating frequency of 3.8 MHz was assumed. This still does not explain Podkletnov's result, but is much larger than the purely B_g acceleration. However, it is not clear what direction the E_g^{ind} vector would point in.

Conclusion

The wire size used in the coils seems to be adequate.

The levitation coils consist of 3 assemblies, each of which is made up of 2 coils connected with a U-shaped core, presumably of steel. This generates flux lines that travel horizontally under the disk, with some flux wrapping up and around the disk through the hole in the center.

Each coil could contain up to 310 turns of approximately 16 gauge wire, at 10 A maximum current. This results in a maximum magnetic flux density B of around 400 gauss at the bottom surface of the superconductor.

The 2 Tesla claim for the system in general does not seem to apply to the levitation coils. Even if one allows 850 gauss for each of 6 coils, right at their surfaces, the total field generated underneath the toroidal superconductor is only about 5100 gauss or 0.5 Tesla. And, this is only for static, not dynamic, magnetic fields. No estimate was made for eddy currents in the core, but intuitively it might be so high at the quoted frequency as to melt the device, or at least cause rapid boil-off of the cryogen due to high power dissipation. A dynamic analysis of this may be forthcoming from the author.

Estimates of the gravitoelectric and gravitomagnetic fields produced in such an arrangement are small, but large enough to be detected using a modern gravimeter. Of course, this estimate may very well be totally wrong, as the 10^{-9} coupling term in the Li/Torr paper may not apply to YBCO superconductors.

The author hopes that these calculations might enable others to reproduce Podkletnov's experiment more accurately. The author welcomes comments on the accuracy of this analysis.

References

- [1] Eugene Podkletnov, "Weak gravitation shielding properties of composite bulk YBCO superconductor below 70K under e.m. field." LANL database number [cond-mat/9701074](#), v. 3, 16 Sep 1997.
- [2] Radio Amateur's Handbook, 25th Edition, 1948, p. 555.
- [3] U.S. National Electrical Code, 1996.
- [4] Applied Electromagnetism, Second Edition, Liang Chi Shen and Jin Au Kong, PWS Publishers, 1987, p. 54, 57.
- [5] Private communication with Giovanni Modanese, 19 Sep 1997.
- [6] D.G. Torr and Ning Li, "Gravitoelectric-Electric Coupling via Superconductivity," Found. Phys. Let. v. 6, n. 4, 1993, p. 371-383.
- [7] R.L. Forward, "General Relativity for the Experimentalist," Proc. IRE, May 1961, p. 893. Symbols were changed from Forward's to the Li/Torr style.
- [8] Ning Li et al, "Static Test for a Gravitational Force Coupled to Type II YBCO Superconductors," Physica C 281, 1997, p. 260-267.

From a Relativistic Phenomenology of Anyons to a Model of Unification of Forces via the Spencer Theory of Lie Structures

Jacques L. Rubin*

May 6, 2002

Abstract

Starting from a relativistic phenomenology of anyons in crystals, we discuss the concept of relativistic interaction and the need to unify electromagnetism and gravitation within the Spencer cohomology of Lie equations. Then, from the sophisticated non-linear Spencer complex of the Poincaré and conformal Lie pseudogroups, we build up a non-linear relative complex assigned to a gauge sequence for electromagnetic and gravitational potentials and fields. Then, using a conformally equivariant Lagrangian density, we deduce, first, the two first steps of its corresponding Janet complex and second, the dual relative linear complex. We conclude by giving suggestions for higher unification with the weak and strong interactions and interpretations of the Lagrangian density as a thermodynamical function and quantum wave-function.

*Institut du Non-Linéaire de Nice, UMR 129 C.N.R.S.-Université de Nice-Sophia-Antipolis, 1361 route des Lucioles, 06560 Valbonne, France. E-mail: rubin@inln.cnrs.fr. Fax: 00/33/04-92-96-73-33

Contents

1. Introduction	3
2. Goals and problems	3
2.1. The physics of crystals and a relativistic phenomenology of anyons	3
2.2. Relativistic interaction in quantum mechanics	6
2.2.a. Polarization in quantum mechanics	6
2.2.b. Concerning the contradictions of an approach “à la Wigner” and the Einstein-Cartan unification	7
3. The Lie conformal pseudogroup associated to the unification model	8
3.1. The conformal finite Lie equations	9
3.2. The conformal Lie groupoid	13
3.3. The first non-linear Spencer differential operators	18
4. The potentials of interaction and the metric	22
4.1. The electromagnetic and gravitational potentials	23
4.2. The morphisms ϕ_1 , D'_1 and the metric of the “gauge space-time”	25
5. The fields of interaction	26
5.1. The second non-linear Spencer operator	26
5.2. The gravitational and electromagnetic fields	28
6. The dual linear Spencer complex and the Janet complex	32
6.1. The dual linear Spencer complex	32
6.2. The Janet complex of the Lagrangian density \mathfrak{L}	33
7. Conclusion	35
References	39

MSC: 35S05, 58G15, 58H15, 58G05, 53A30, 58H05, 53C10, 81T13

keywords: anyons, superconductivity, unification models, electromagnetism, gravitation, gauge theory, non-linear Spencer cohomology, conformally flat Riemannian manifolds, Lie groupoids, G-structures, Lie pseudogroups, deformations, differential complexes. Lie equations.

Short running title: From Anyons to Unification of Forces via the Spencer Theory

1. Introduction

In this paper we propose a model as well as suggestions for a unification of physical interactions. This is a model of electromagnetic and gravitational interactions well-founded on a phenomenological relativistic model of anyons in high- T_c superconductors (Rubin 1994). This unification has its roots, first in the Spencer cohomology of the conformal Lie pseudogroup which has abundantly been studied by J. Gasqui & H. Goldschmidt (Gasqui *et al.* 1984), and second, in the non-linear cohomology of Lie equations studied by H. Goldschmidt & D. Spencer (1976a, 1976b, 1978a, 1978b, 1981, see references therein). Meanwhile we only partially refer to some of its aspects to work out a relative non-linear cohomology explicitly associated with a model of unification. Such an approach was originally proposed by J.-F. Pommaret (1988, 1989, 1994) however, it appeared to us to be incomplete, indeed erroneous.

Thus, the purpose of the present paper is to discuss the Pommaret model and to suggest new developments based on the same assumption. Like Pommaret (1989), we think that the geometrical approach of the Maxwell theory has to be modified to be incorporated in a larger theory which explicitly includes gravitation through different equations describing the variations of potentials of gravitation. This result - or proposition - has not been obtained by Pommaret (Pommaret 1994, see conclusions page 456) who could not find any alternative descriptions and justifications neither for the Einstein theory, nor for the equations of fields of gravitation within the frame of the Spencer cohomology.

We conclude this paper a) by succinctly proposing a possible reinterpretation of the quantum wave-function as a classical thermodynamic function within the frame of the Misra-Prigogine-Courbage (MPC) ergodic theory of fields (Misra *et al.* 1979, Misra 1987), b) by suggesting ideas about a unification including the weak and strong interactions along a basic “à la Penrose” approach (Penrose *et al.* 1986).

Furthermore, this work is the result of informal reflexions about an increasing amount of contradictions and incoherences mainly concerning the concept of relativistic interaction, (that we find more and more serious) in the field of quantum physics as well as in classical physics. According to this observation, we first present our motives and a description of these contradictions in relation to F. Lurçat’s (1964), J.-M. Lévy-Leblond’s (1990) and J.-F. Pommaret’s arguments (1989) in order to justify a development via the Spencer cohomology of Lie pseudogroups (Goldschmidt *et al.* 1976a, 1976b, 1978a, 1978b, 1981).

2. Goals and problems

2.1. The physics of crystals and a relativistic phenomenology of anyons

Our initial motivation shall be seen as extremely far from the problems with unifications. Actually, we were more concerned in a simple minor model of a relativistic phenomenology of creation of anyons, accurate for certain crystals (Rubin 1994). At the

origin of this process of creation, we suggested the kinetico-magnetoelectrical effect as described by E. Asher (1973) and which has its roots in the former Minkowski works about the relations between tensors of polarization P and Faraday tensors F in a moving material of optical index $n \neq 1$. These relations are established by turning the following diagram into a commutative one:

$$\begin{array}{ccc} F' & \xrightarrow{\Lambda} & F \\ \downarrow \chi' & & \downarrow \chi \\ P' & \xrightarrow{\Lambda} & P \end{array}$$

where Λ is a Lorentz transformation, allowing us to shift from a frame R' to a frame R , and χ' and χ are respectively the tensors of susceptibility within those two frames, as well supposing P (or P') linearly depending on F (respectively F'). Resulting from this commutativity, the tensor χ linearly depends on χ' in general and also on a velocity 4-vector \tilde{u} associated to Λ (i.e. the relative velocity 4-vector between R and R'). In assimilating R' to the moving crystal frame and R to the laboratory frame, then to an applied electromagnetic field F fixed in R , corresponds in R' a field of polarization P which varies in relation to \tilde{u} . This is the so-called kinetico-magnetoelectrical effect.

Parallel to this phenomenon, A. Janner & E. Asher studied the concept of relativistic point symmetry in polarized crystals (Janner *et al.* 1969, 1978). Such a symmetry is defined, on the one hand, by a given discrete group G , sub-group of the so-called Shubnikov group $O(3)1'$ associated with the crystal, and on the other hand, as satisfying the following properties: to make this relativistic symmetry exist, there must be a $H(P)$ non-trivial group of Lorentz transformations depending on P , in which G is a normal sub-group, and that leaves the tensor of polarization P invariant. In other words, if $N(G)$ is the normalizer of G in the Lorentz group $O(1, 3)$, and $K(P)$ the sub-group of $O(1, 3)$ leaving P invariant, then $H(P)$ is the maximal sub-group such that:

$$\begin{cases} H(P) \subseteq K(P) \cap N(G) \\ H(P) \cap O(3)1' = G. \end{cases}$$

We can prove that $H(P)$ is about to exist only if a particular non-vanishing set V of velocity 4-vectors, invariant by action of G , is present and consequently compatible with a kinetico-magnetoelectrical effect (Asher 1973). Therefore, if there is an interaction between moving particles in the crystal and the polarization P , then the trajectories and P are obviously modified, and so is $H(P)$. In this process, only the group $N(G)$ is conserved so that the polarization and the trajectories are deducible during the time by the action of $N(G)$.

As we shall stipulate later on, the existence of an interaction will emerge due to a correlation between the position 3-vectors \vec{r} of the charge carriers and a particular 3-vector \vec{w} ($\notin V$ in general) associated with P ; \vec{w} becoming then a function of \vec{r} . In order to allow a cyclotron-type motion which is implicit within the theory of anyons, the group $N(G)$ must contain the group $SO(2)$ and the latter must non-trivially act on all the groups $H(P)$ associated to G . Then, only 12 groups G are compatible with such a description

(Rubin 1993). In fact, throughout this development, we implicitly use a principle of equivalence similar to the one formulated in general relativity: one cannot distinguish a cyclotron - type motion in a constant polarization field from a uniform rectilinear motion in a field of polarization varying in time by action of the normalizer $N(G)$. The time evolution of \vec{w} requiring the explicit knowledge of the gradient $\vec{\nabla}(\vec{w})$ of \vec{w} , it follows that \vec{w} and $\vec{\nabla}(\vec{w})$ are respectively the analogues (not the equivalents) of the tetrads and of the Killing vector fields. From an other point of view, the interaction is considered to allow the extension of an invariance with respect to $H(P)$ to an invariance with respect to $N(G)$. The lack of interaction is then what breaks down the symmetry!

This type of reasoning concerns in fact a large amount of physical phenomena such as the spin-orbit interaction for instance. In this context, the cyclotron-type motion of electrons in anyonic states would be similar to the Thomas or Larmor precessions (see also the Coriolis or Einstein-Bass effects). More precisely, taking up again a computation, analogous to the Thomas precession one (Bacry 1967) (i.e. considering as a constant the scalar product of two tangent vectors being two parallel transports along the trajectory (Dieudonné 1971)), concerning a charge carrier with the velocity 4-vector \tilde{u} in R , “polarized” by $\vec{w}(\vec{r})$ such as for example ($\tilde{v} = (0, \vec{v})_R$ constant and $\vec{v} \in V$):

$$\tilde{w} = (0, \vec{w})_R \equiv -P.\tilde{v} \text{ or } {}^*P.\tilde{v},$$

where P depends on \vec{r} , one can prove from $\tilde{w}.\tilde{u} = cst.$ that (t being the laboratory frame time and $(\tilde{r} = (t, \vec{r})_R)$:

$$\frac{d\tilde{u}}{dt} = (-e/m)F_{eff.}(\tilde{r}).\tilde{u}, \quad (1)$$

where m and e are respectively the mass and the electric charge of the carrier and $F_{eff.}(\tilde{r}) \equiv (\vec{E}_{eff.}(\tilde{r}), \vec{B}_{eff.}(\tilde{r}))$ is an effective Faraday tensor such that ($\gamma = (1 - \vec{w}^2)^{-\frac{1}{2}}$ and $\vec{j} = e\vec{u}$):

$$\begin{cases} \vec{B}_{eff.}(\tilde{r}) = (m/e^2) \left(\frac{\gamma}{1+\gamma} \right) \vec{w} \wedge [\vec{j} \cdot \vec{\nabla}] \vec{w} \\ \vec{E}_{eff.}(\tilde{r}) = \vec{0}. \end{cases}$$

Clearly, $F_{eff.}$ is an element of the Lie algebra of the group $SO(2)$ included in $N(G)$ and with $\vec{B}_{eff.} \in V$. Therefore this $\vec{B}_{eff.}$ magnetic field or \vec{v} (up to a constant) might be considered as the effective magnetic field of the flux-tube V generating the so-called Aharonov-Böhm effect at the origin of the statistical parameter in the anyons theory (Wilczek 1990). In this precession computation, from a more mathematical point of view, taking up the Lie groupoids theory, we think that perhaps we shift from a source “description” (at $t = 0$) to a target one (at any $t \neq 0$). It is definitely an equivalence principle analogous to the one occurring in general relativity, as shown by J.-F. Pommaret (1989). Let us add that in general $\text{div}(\vec{B}_{eff.}) \neq 0$ so that one gets a non-vanishing density of effective magnetic monopoles generated by the local variations (due to the interaction) of the polarization vector field \vec{w} in the crystal. Thus an anyon would be an effective magnetic monopole associated with a charge carrier, namely a dyon. Moreover, because this effective Faraday tensor is no more a closed two-from, a non-vanishing Chern-Simon

has to be taken into account in a Lagrangian description of anyons, from which non-vanishing spontaneous constant currents can occur.

2.2 Relativistic interaction in quantum mechanics

2.2.a. Polarization in quantum mechanics

The previous vector \vec{B}_{eff} . (or $\vec{v} \in V$) has the same status as the spin. Like the latter, it is defined by a torsor of order 2. From that time on, the transition from a classical description to a quantum one means that one must give an account of the interaction between a free particle (constituting a first sub-system) and a torsor field of order 2 (constituting a second sub-system). The problem seems to be solved and in particular the spin appears to be a “minor” complication of the wave-function defined on the Minkowski space-time, i.e. on a “non-polarized” space. This has to be taken as a postulate, an erroneous one according to part of F. Lurçat’s (1964) and J.-M. Lévy-Leblond’s (1990) arguments.

In fact, it is not even the case according to classical Galilean mechanics. Considering a body at a given time, one needs to know 6 parameters to describe it: 3 for the position and 3 others for the orientation. The latter are forgotten during the transition from the “extended body to the punctual particle” according to the quantum description. This can only be justified providing that the energy of rotation is negligible compared to the energy of translation. This is the indicator of an inadequacy of the principle of correspondence from classical mechanics to quantum mechanics. But the transition from quantum to classical mechanics is equally problematic: reaching the limit $\hbar \rightarrow 0$, the spin vanishes. Finally, we must say that this correspondance does not exist any more in chromodynamics.

Therefore, the fact that the wave-function only depends on the position should rather be considered as a postulate (moreover, the fact is inexistant in classical mechanics). For instance, if we measure the electric and magnetic fields at a point in space-time with a system of coordinates defined by a given Lorentz transformation, we can deduce that the Faraday tensor is a function of the 4 parameters of the position and 6 others defining Λ . It explicitly occurs within the kinetico-magnetoelectric effect.

Referring now to the wave-equations and to the methods usually accepted to determine them, we then use a principle of invariance. First, we identify the appropriate space-time symmetries, that is the group of relativity of the theory (for example the Poincaré group). Then, according to Wigner’s theory, we build the irreducible unitary representations (eventually projective) of the group to which correspond the elementary “kinetic” objects of the theory (i.e. the vector-valued wave-functions). Then again, we derive the infinitesimal generators of the group that we identify with physical and geometrical observables like the energy, the kinetic moment, etc...; the Lie algebra of the group and its “quantum” extensions defining the commutators. Finally, determining the invariants through the action of the group we obtain the other physical observables and their commutators.

Unfortunately, some ambiguities appear. For example, the generalization of the Dirac theory for spins higher than 1/2 gives different non-equivalent possible wave-equations.

Lastly, if we consider a theory of interacting fields, the particles associated with these fields can get out of their mass shell but not out of their spin shell! We then forget the spin again, which is impossible within a “(m,s)” theory. Therefore some of the aspects of the interaction and of the wave-equations - to be brief- can fully account for neither the free particles, nor the interacting particles!

In order to escape from these contradictions and to best describe the free particles, F. Lurçat proposed an approach “à la Wigner”. In this way, he postulated that first the scalar complex wave-function of a free particle was defined on the “Poincaré space” of the Poincaré Lie group. Second, this wave-function is an eigenfunction of the two Casimirs of the Poincaré group. Thus, the wave-function ϕ is a function first, of a 4-vector position \tilde{x} , element of the dual Lie algebra of the group of translations and second, of a second order torsor F (like the Faraday torsor which is an element of the dual Lie algebra of the Lorentz group). Also in fact, the Poincaré Lie group on the “Poincaré space” is the action of a so-called Lie groupoid on its associated Lie algebroid, defined on the Minkowski space-time. Finally, if we want to describe the interaction, making these modifications and requiring a gauge invariance, the involution of the infinitesimal generators “deformed” by the gauge fields is no longer satisfied, nor are the relativistic invariance and the correspondence principle.

This breaking of the invariance can be seen with the Dirac equation. More exactly, the Dirac equation is not *equivariant*, but only covariant, meaning that this equation is not invariant under any conformal changes of coordinates, but only under a change of frame, i.e. a change of basis of the tangent Minkowski space. In that case, for instance, one of the spectacular manifestations is Klein’s paradox of non-conservation of the current of probability during the scattering in a square potential (Itzykson *et al.* 1980). Still, the idea stays that ϕ is defined on a “larger” space-time than the Minkowski one, but for being a Kaluza-Klein type theory for example.

2.2.b. Concerning the contradictions of an approach “à la Wigner” and the Einstein-Cartan unification

Let us consider a complex wave-function ϕ depending on \tilde{x} and F on which the Poincaré group projectively acts. Because, then, of the phase arbitrariness, the ten infinitesimal generators of the group are defined with ten arbitrary gauge potentials, so-called Poincaré potentials. In order to keep the Lie algebra structure, i.e. the involution of the deformed generators, the fields associated with these potentials must satisfy some constraint equations. The electromagnetic field associated with the translations of the group has especially to be vanishing, which is first absolutely contradictory with ϕ as a function of F . Second, that would mean that one cannot describe any interaction with a field F without a Poincaré symmetry breaking. Nevertheless this symmetry is necessary to keep the relativistic equivariance.

Moreover, ϕ is independent of the gauge potentials although ϕ should be parametrized by these functions referring to the formal theory of system of partial differential equations (PDE). In fact, according to the Kähler-Cartan theorem, the analytical solutions of an involutive system of PDE (so formally integrable) depend on certain constants of integra-

tion (Dieudonné 1971), but, contrary to the ODE (ordinary differential equations), they depend on arbitrary C^∞ functions only constrained to verify Cauchy initial data (see also Shih 1986, 1987, 1991). These functions should be identified with the gauge potentials and with their fields. Still, we can say that the incoherences are not over!

Let us assume the presence of a relativistic interaction, then there consequently exists, if we refer to the equivalence principle, a proper frame in which ϕ is stationnary. If τ is the “proper” time then $\partial_\tau \phi = 0$. Within the “laboratory” frame, we will get a different equation, moreover, there won’t be any relativistic equivariance; a problem that is similarly encountered with either the Newton-Wigner position operator or the Dirac position operator. Indeed we will still be unable to write down this equation (!) for the reason that to shift to the Laboratory frame, one needs to know the classical motion of the particle, i.e. to know the equation that determines the evolution of the 4-vector speed \tilde{u} identified with the basic time type 4-vector \tilde{e}_0 of the proper frame.

Also at this point, the equation should be established in a certain system of coordinates and from \tilde{e}_0 only, one should obtain the other basic vectors of space type \tilde{e}_i ($i=1,2,3$) as well as their time evolutions. This equation should then be integrable in the Fröbenius sense. We think it is perhaps a matter of a generalisation of the Frénet moving frame method, as formulated by E. Cartan. Whereas we observe that by determining this moving frame from \tilde{e}_0 and $\dot{\tilde{e}}_0$, that the \tilde{e}_i ’s are defined from the \tilde{e}_μ ’s ($\mu=0,1,2,3$) and from the third order time derivatives of \tilde{e}_0 . Then, if $\dot{\tilde{e}}_0 \equiv F \cdot \tilde{e}_0$, we need to know the time derivatives of F up to the second order. Therefore, in the laboratory frame, ϕ should depend on the derivatives of F , contrary to the initial assumption, unless these derivatives are themselves functions of F and \tilde{x} . Condition which, first, is not the case in the Maxwell theory, second would set down some constraints on the moving frame and consequently a “partial” equivariance; unless one completes it with other fields - like those associated for example with the space-time curvature - which would be related to the derivatives of F . Let us remark also that this kind of discussion seems to be very similar to the one encountered in the demonstration of the Cauchy-Kowalewska theorem when starting from a pfaffian system to a “normal system” of PDE’s and considering the so-called “regular” tangent spaces and integrable manifolds (Dieudonné 1971).

On that subject, one can notice that the equation (1) can be rewritten in an orthonormal system of local coordinates ($\alpha, \beta, \gamma = 1, \dots, n$; the Γ ’s being the Christoffel symbols):

$$\dot{u}^\alpha + \Gamma_{\beta,\gamma}^\alpha u^\beta u^\gamma = 0.$$

We recognize the equation of the geodesics associated to a Riemannian connexion with torsion. This would suggest a unification in reference to the Einstein-Cartan theory. If we then keep on with the assumption that one has to add to the electromagnetic field a gravitational field and that the derivatives of the fields are functions of the fields themselves (as with the Bianchi identities according to the non-abelian theory for example), that means we make the assumption of the existence of a differential sequence. In electromagnetism, it is a matter of the de Rham sequence but gravitation does not interfere. The sequence integrating the latter - and being the purpose of this paper - might be a certain generalizing complex like the Spencer one, following then a method proposed by J.-F. Pommaret (1994) but largely modified.

3. The Lie conformal pseudogroup associated to the unification model

First of all, let us assume that the group of relativity is not the Poincaré group anymore but the conformal Lie group (we know from Bateman and Cunningham studies (1910) that it is the group of invariance of the Maxwell equations). In particular, this involves that no changes occur shifting from a given frame to a uniformly accelerated relative one. From a historical point of view, that happened to be the starting point of the Weyl theory which was finally in contradiction with experimental data and for various other reasons presented, for instance by J.-F. Pommaret in the framework of the Janet and Spencer complexes (Pommaret 1989). Then, starting from this mathematical framework, J.-F. Pommaret considered in trying this unification, the linear Spencer complex defined from the system of finite Lie equations associated to the Lie pseudogroup of conformal isometries. Unfortunately, the system of PDE proposed by J.-F. Pommaret is incomplete and its conditions of use are not really given. On the other hand, he claimed the Spencer complex of the conformal Lie pseudogroup would be the “unification complex” (Pommaret 1988), whereas we merely prove that it would rather be a relative complex deduced from an abelian extension (Godschmidt 1976a, 1976b, 1978a, 1978b, 1981), when reaching the conformal Lie pseudogroup starting from the Poincaré one. Before tackling these various complexes, we present and recall a few relations concerning the conformal Lie pseudogroup action on some tensors such as the metric and the Riemann and Weyl curvatures. Let us first call \mathcal{M} , the base space (or space-time), assumed to be of class C^∞ , of dimension $n \geq 4$, connected, paracompact, without boundaries, oriented and endowed with a metric 2-form ω , symmetric, of class C^2 on \mathcal{M} and non-degenerated. We also assume \mathcal{M} to be conformally flat.

3.1. The conformal finite Lie equations

These equations are deduced from the conformal action on the metric. Let us consider $\hat{f} \in \text{Diff}_{loc}^1(\mathcal{M})$ and any $\alpha \in C^0(\mathcal{M}, \mathbb{R})$, then if $\hat{f} \in \Gamma_{\widehat{G}}$ ($\Gamma_{\widehat{G}}$ being the pseudogroup of local conformal bidifferential maps on \mathcal{M}), \hat{f} is a solution of the following system of PDE (other PDE must be satisfied to completely define $\Gamma_{\widehat{G}}$):

$$\begin{cases} \hat{f}^* \omega = e^{2\alpha} \omega \\ \text{and with } \det(J(\hat{f})) \neq 0, \end{cases} \quad (2)$$

where $J(\hat{f})$ is the Jacobian of \hat{f} , and \hat{f}^* is the pull-back of \hat{f} . We denote $\tilde{\omega}$ the metric on \mathcal{M} such as:

$$\tilde{\omega} \stackrel{\text{def.}}{=} e^{2\alpha} \omega,$$

and we agree to put a tilde on each tensor or geometrical “object” relative to or deduced from this metric $\tilde{\omega}$. Let us notice that the latter depends on a fixed given element $\hat{f}^* \in \Gamma_{\widehat{G}}$. Also, in order to properly recall the last point, we shall sometimes use an alternative notation such as:

$$\tilde{\omega} \stackrel{def.}{\equiv} \hat{f}\omega.$$

This convention of notation will also be used on each geometrical object relative to this metric. Now, doing a first prolongation of the system (2), we deduce other second order PDE connecting the affine connexion 1-forms of Levi-Civita ∇ and $\widetilde{\nabla}$ respectively associated to ω and $\tilde{\omega}$. To obtain these PDE, we merely start from the following definition of $\widetilde{\nabla}$: let X, Y and Z be any vector fields in $C^1(T\mathcal{M})$, $\hat{f} \in \text{Diff}_{loc.}^2(\mathcal{M})$ and $\alpha \in C^1(\mathcal{M}, \mathbb{R})$, then by definition we have:

$$\begin{aligned} \tilde{\omega}(\widetilde{\nabla}_X Y, Z) = & \frac{1}{2}\{\tilde{\omega}([X, Y], Z) + \tilde{\omega}([Z, X], Y) + \tilde{\omega}([Z, Y], Y) \\ & + X.\tilde{\omega}(Y, Z) + Y.\tilde{\omega}(X, Z) - Z.\tilde{\omega}(X, Y)\}, \end{aligned} \quad (3)$$

from which we deduce with the relation (2) $\forall X, Y \in C^1(T\mathcal{M})$ and $\forall \hat{f} \in \text{Diff}_{loc.}^2(\mathcal{M})$,

$$\widetilde{\nabla}_X Y = \nabla_X Y + \mathbf{d}\alpha(X)Y + \mathbf{d}\alpha(Y)X - \omega(X, Y) * \mathbf{d}\alpha, \quad (4)$$

where \mathbf{d} is the exterior differential and $*\mathbf{d}\alpha$ is the dual vector field of the 1-form $\mathbf{d}\alpha$ with respect to the metric ω , i.e. such as $\forall X \in T\mathcal{M}$

$$\omega(X, *\mathbf{d}\alpha) = \mathbf{d}\alpha(X) = <\mathbf{d}\alpha|X>. \quad (5)$$

Let us also agree to denote in the sequel $\forall \tilde{x} \in \mathcal{M}$, $\forall \nu \in \wedge^r T^*\mathcal{M}$ and $\forall \xi_i \in T\mathcal{M}$ ($i = 1, \dots, r$):

$$\nu_{\tilde{x}}(\xi_{1,\tilde{x}}, \dots, \xi_{r,\tilde{x}}) = <\nu(\tilde{x})|\xi_{1,\tilde{x}} \otimes \dots \otimes \xi_{r,\tilde{x}}>.$$

Prolonging again and using the definition of the Riemann tensor ρ associated to ω , i.e. $\forall X, Y \in C^2(T\mathcal{M})$

$$\rho(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]},$$

one obtains the following relation $\forall X, Y, Z \in C^2(T\mathcal{M})$, $\forall \alpha \in C^2(\mathcal{M}, \mathbb{R})$ and $\forall \hat{f} \in \text{Diff}_{loc.}^3(\mathcal{M})$,

$$\begin{aligned} \tilde{\rho}(X, Y).Z = & \rho(X, Y).Z + \omega(X, Z)\nabla_Y(*\mathbf{d}\alpha) - \omega(Y, Z)\nabla_X(*\mathbf{d}\alpha) \\ & + \{\omega(\nabla_X(*\mathbf{d}\alpha), Z) + \omega(X, Z)\mathbf{d}\alpha(*\mathbf{d}\alpha)\}Y \\ & - \{\omega(\nabla_Y(*\mathbf{d}\alpha), Z) + \omega(Y, Z)\mathbf{d}\alpha(*\mathbf{d}\alpha)\}X \\ & + \{\mathbf{d}\alpha(X)\omega(Y, Z) - \mathbf{d}\alpha(Y)\omega(X, Z)\}*\mathbf{d}\alpha \\ & + \{\mathbf{d}\alpha(Y)X - \mathbf{d}\alpha(X)Y\}\mathbf{d}\alpha(Z) \end{aligned} \quad (6)$$

Assuming \mathcal{M} to be conformally flat, the Weyl tensor τ associated with ω vanishes. Hence, the Riemann tensor ρ can be rewritten $\forall X, Y, Z, U \in C^2(T\mathcal{M})$ as:

$$\begin{aligned} \omega(U, \rho(X, Y).Z) = & \frac{1}{(n-2)}\{\omega(X, U)\sigma(Y, Z) - \omega(Y, U)\sigma(X, Z) \\ & + \omega(Y, Z)\sigma(X, U) - \omega(X, Z)\sigma(Y, U)\}, \end{aligned} \quad (7)$$

where σ is the so-called Schouten tensor (Gasqui /it et al. 1984) $\forall X, Y \in C^2(T\mathcal{M})$,

$$\sigma(X, Y) = \rho_{ic}(X, Y) - \frac{\rho_s}{2(n-1)}\omega(X, Y), \quad (8)$$

where ρ_{ic} is the Ricci tensor and ρ_s is the Riemann scalar curvature. Thus, we might consider the relation (7) as the existence of a short exact sequence “symbolically” written as “ $0 \rightarrow \sigma \rightarrow \rho \rightarrow \tau \rightarrow 0$ ” and perhaps related to a sequence of cohomology spaces of symbols (Gasqui *et al.* 1984, Pommaret 1994). Consequently, the system of PDE (6) can be rewritten as a first order system of PDE concerning σ . To do this, we first define two suitable trace operators, used in the sequel to obtain the $\tilde{\rho}_{ic}$ and $\tilde{\rho}_s$ tensors and finally the $\tilde{\sigma}$ tensor. Let us denote Tr^1 the trace operator defined such that for any vector bundle E over \mathcal{M} we have:

$$Tr^1 : T\mathcal{M} \otimes T^*\mathcal{M} \otimes E \longrightarrow E,$$

with

$$Tr^1(X \otimes \alpha \otimes \mu) = \alpha(X)\mu$$

for any $X \in T\mathcal{M}$, $\alpha \in T^*\mathcal{M}$ and $\mu \in E$. Then, the second trace operator is the natural trace Tr_ω associated to ω and defined by:

$$Tr_\omega : \overset{2}{\otimes} T^*\mathcal{M} \longrightarrow \mathbb{R},$$

such that

$$Tr_\omega(u \otimes v) = v(*u).$$

Finally, with $Tr^1\tilde{\rho} = \tilde{\rho}_{ic}$ and $Tr_\omega\tilde{\rho}_{ic} = \tilde{\rho}_s$, we deduce first from the relations (6) and (8) $\forall \hat{f} \in Diff_{loc}^3(\mathcal{M})$, $\forall X, Y \in C^2(T\mathcal{M})$ and $\forall \alpha \in C^2(\mathcal{M}, \mathbb{R})$,

$$\begin{aligned} \tilde{\sigma}(X, Y) &= \sigma(X, Y) + (n-2)(\mathbf{d}\alpha(X)\mathbf{d}\alpha(Y) - \omega(\nabla_X(*\mathbf{d}\alpha), Y) \\ &\quad - \frac{1}{2}\omega(X, Y)\mathbf{d}\alpha(*\mathbf{d}\alpha)). \end{aligned} \quad (9)$$

In fact this expression can be symmetrized. Indeed, from the following property satisfied by ∇ :

$$\omega(\nabla_X(*\mathbf{d}\alpha), Y) + \omega(*\mathbf{d}\alpha, \nabla_X Y) = X \cdot \omega(*\mathbf{d}\alpha, Y),$$

and the definition of $*\mathbf{d}\alpha$, one obtains:

$$\omega(\nabla_X(*\mathbf{d}\alpha), Y) = X \cdot \mathbf{d}\alpha(Y) - \mathbf{d}\alpha(\nabla_X Y).$$

But, from the torsion free property of ∇ and from the relation

$$\mathbf{d}\alpha([X, Y]) = X \cdot \mathbf{d}\alpha(Y) - Y \cdot \mathbf{d}\alpha(X),$$

one deduces:

$$\omega(\nabla_X(*\mathbf{d}\alpha), Y) = Y \cdot \mathbf{d}\alpha(X) - \mathbf{d}\alpha(\nabla_Y X).$$

Now, $\forall \alpha \in C^2(\mathcal{M}, \mathbb{R})$, $\forall X, Y \in C^1(T\mathcal{M})$, defining $\mu \in C^0(S_2 T^*\mathcal{M})$ by:

$$\mu(X, Y) = \frac{1}{2} [X \cdot \mathbf{d}\alpha(Y) + Y \cdot \mathbf{d}\alpha(X)], \quad (10)$$

one has the relation:

$$\omega(\nabla_X(*\mathbf{d}\alpha), Y) = \mu(X, Y) - \frac{1}{2} \mathbf{d}\alpha(\nabla_X Y + \nabla_Y X).$$

Then, we can rewrite the first order PDE (9), $\forall \hat{f} \in \text{Diff}_{loc.}^3(\mathcal{M})$, $\forall X, Y \in C^2(T\mathcal{M})$ and $\forall \alpha \in C^2(\mathcal{M}, \mathbb{R})$ as:

$$\begin{aligned} \tilde{\sigma}(X, Y) &= \sigma(X, Y) + (n-2)(\mathbf{d}\alpha(X)\mathbf{d}\alpha(Y) \\ &\quad - \mu(X, Y) + \frac{1}{2}\mathbf{d}\alpha(\nabla_X Y + \nabla_Y X) \\ &\quad - \frac{1}{2}\omega(X, Y)\mathbf{d}\alpha(*\mathbf{d}\alpha)). \end{aligned} \quad (11)$$

It is worthy of note that if $\rho_s = c_0$ ($c_0 \in \mathbb{R}$), then the Schouten tensor σ satisfies the relation:

$$\sigma = c_0 \frac{(n-2)}{2} \omega. \quad (12)$$

and only in this case, the equation (11) must become the equation 2 and so it disappears. Then, considering the system (2), the system (11) reduces to a second order system of PDE such as $\forall \alpha \in C^2(\mathcal{M}, \mathbb{R})$ and $\forall X, Y \in C^1(T\mathcal{M})$ we have:

$$\begin{aligned} \mu(X, Y) &= \frac{1}{2} \{ [c_0(1 - e^{2\alpha}) - \mathbf{d}\alpha(*\mathbf{d}\alpha)] \omega(X, Y) + \mathbf{d}\alpha(\nabla_X Y + \nabla_Y X) \} \\ &\quad - \mathbf{d}\alpha(X)\mathbf{d}\alpha(Y). \end{aligned} \quad (13)$$

We also have the following PDE deduced from (4), $\forall X \in C^1(T\mathcal{M})$ and $\forall \hat{f} \in \text{Diff}_{loc.}^2(\mathcal{M})$:

$$Tr^1(\widetilde{\nabla}_X) = Tr^1(\nabla_X) + n \mathbf{d}\alpha(X). \quad (14)$$

Thus, we have a serie of PDE deduced from (2), in particular made of the systems of PDE (2), (11) and (14). But there are alternative versions of these PDE in which the function $\alpha \in C^2(\mathcal{M}, \mathbb{R})$ doesn't appear. These latters are the following: from the system (2), one deduces, $\forall \hat{f} \in \text{Diff}_{loc.}^1(\mathcal{M})$:

$$\begin{cases} \hat{f}^* \hat{\omega} = \hat{\omega} \\ \text{and with } \det(J(\hat{f})) \neq 0, \end{cases} \quad (15)$$

where $\hat{\omega} = \omega / |\det(\omega)|^{1/n}$. Then by prolongation, with $\widehat{\nabla}$ and $\hat{\rho}$ being respectively the connexion of Levi-Civita and the Riemann curvature tensor associated to $\hat{\omega}$, one obtains $\forall \hat{f} \in \text{Diff}_{loc.}^3(\mathcal{M})$:

$$\hat{f}\widehat{\nabla} = \widehat{\nabla} \quad (16)$$

and

$${}^{\hat{f}}\hat{\rho} = \hat{\rho}. \quad (17)$$

In the latter system (17) of PDE (first order), it has to be noted that $\hat{\rho} = \tau$, i.e. $\hat{\rho}$ is the Weyl curvature tensor associated to the metric ω . Furthermore, with the assumption that the conformal structure is flat, one has $\hat{\rho} = 0$ and $\hat{\sigma} = 0$. But in general, it is noteworthy to add that if $\tau = 0$ then obviously σ doesn't vanish. Then, the conformal Lie pseudogroup $\Gamma_{\widehat{G}}$ is the set of functions $\hat{f} \in \text{Diff}_{loc}^3(\mathcal{M})$ satisfying the following involutive system of PDE:

$$\begin{cases} \hat{f}^*\hat{\omega} = \hat{\omega} & \text{and with} \\ \hat{f}\hat{\nabla} = \hat{\nabla}, & \det(J(\hat{f})) \neq 0 \end{cases} \quad (18)$$

completed with a third system of PDE of order 3 defined $\forall X, Y \in C^2(T\mathcal{M})$ by:

$$\hat{f}\hat{\nabla}_X \hat{f}\hat{\nabla}_Y = \hat{\nabla}_X \hat{\nabla}_Y. \quad (19)$$

This system is formally integrable if and only if $\tau = 0$ (from the Weyl theorem) and involutive because the corresponding symbol \widehat{M}_3 vanishes. From a terminological point of view, one shall say that the system (18)-(19) is the “Lie form” of the system made of the PDE (2), (11) and (14) to which one adds the third order system deduced from the expression of $\hat{f}\hat{\nabla}_X \hat{f}\hat{\nabla}_Y$ with respect to $\nabla_X \nabla_Y$ and $\alpha \in C^3(\mathcal{M}, \mathbb{R})$. We shall call this latter system equivalent to the system (18)-(19), the “deformed” or “extended” system. It is remarkable that the deformed system brings out a supplementary system in comparison with its Lie form (18)-(19). It is about the system (11). If we consider the second order sub-system deduced from the system (18)-(19), we notice it is still formally integrable (again because of the Weyl theorem) but it is no longer involutive because the symbol \widehat{M}_2 of (18)-(19) is only 2-acyclic.

3.2. The conformal Lie groupoid

Before defining this groupoid, we need to recall some definitions concerning the sheafs of the k-jet fiber bundles (Kumpera *et al.* 1972). First of all, we denote $J_k(\mathcal{M})$ the affine fiber bundle of the k-jets of local $C^\infty(\mathcal{M}, \mathcal{M})$ functions on \mathcal{M} , $\theta_{\mathcal{M}}$ (or simply θ) the sheaf of local rings of germs of continuous functions on \mathcal{M} with values in \mathbb{R} , and J_k the affine fiber bundle of the k-jets of local functions in $C^\infty(\mathcal{M}, \mathbb{R})$. Then in what follows, we agree to underline all the names used for the sheafs of germs of local continuous sections associated with the various fiber bundles. Now, if ϵ is a sheaf of θ -modules on \mathcal{M} , we conventionally define the sheaf $\underline{J}_k(\epsilon)$ as:

$$\underline{J}_k(\epsilon) \equiv \underline{J}_k \otimes_{\theta} \epsilon.$$

Then, we have the injective sheafs map:

$$\begin{aligned} j_k : \quad & \underline{C^\infty(\mathcal{M})} \longrightarrow \underline{J_k(\mathcal{M})} \\ [m \rightarrow f(m)]_x & \longrightarrow [m \rightarrow j_k(f)(m)]_x \equiv j_k([f])_x, \end{aligned}$$

where $j_k(f)(m)$ is the set of germs at m of the derivatives of f up to order k , and $[]_x$ obviously being the equivalence classes of local sections at $x \in \mathcal{M}$. Let us also denote $\text{Aut}(\mathcal{M})$ the sheaf of germs of functions $f \in \text{Diff}_{loc}^\infty(\mathcal{M})$. The source map such as:

$$\begin{aligned} \alpha_k : \quad & J_k(\mathcal{M}) \longrightarrow \mathcal{M} \\ j_k(f)(x) & \longrightarrow x, \end{aligned}$$

and the target map:

$$\begin{aligned} \beta_k : \quad & J_k(\mathcal{M}) \longrightarrow \mathcal{M} \\ j_k(f)(x) & \longrightarrow f(x), \end{aligned}$$

are submersions on \mathcal{M} . One defines the composition on $J_k(\mathcal{M})$ by:

$$J_k(g)(y) \cdot J_k(f)(x) = J_k(g \circ f)(x),$$

with $y = f(x)$. The units in $\underline{J_k(\mathcal{M})}$ are the elements $j_k([id])_x$ and they can be identified with the points $x \in \mathcal{M}$. Then, let $\Pi_k(\mathcal{M})$ be the Lie groupoid of invertible elements of $J_k(\mathcal{M})$. The elements of $\Pi_k(\mathcal{M})$ are the k -jets of the functions $f \in \text{Diff}_{loc}^\infty$. $\underline{J_k(\mathcal{M})}$ (resp. $\underline{\Pi_k(\mathcal{M})}$) is also the sheaf of germs of local continuous sections f_k of α_k (resp. $\alpha_k/\Pi_k(\mathcal{M})$). The sheaf map j_k can be also restricted to the sheaf map:

$$j_k : \text{Aut}(\mathcal{M}) \longrightarrow \underline{\Pi_k(\mathcal{M})}.$$

An element $[f_k] \in \underline{\Pi_k(\mathcal{M})}$ shall be called “admissible” if $f = \beta_k \circ f_k \in \text{Aut}(\mathcal{M})$ (i.e. $\det([j_1(f)]) \neq 0$). The admissible elements are the germs of continuous sections f_k of $\alpha_k : \Pi_k(\mathcal{M}) \rightarrow \mathcal{M}$ such as $\beta_k \circ f_k \in \text{Diff}_{loc}^\infty(\mathcal{M})$. We denote $\Gamma_k(\mathcal{M})$ the sub-sheaf of admissible elements of $\underline{\Pi_k(\mathcal{M})}$. Then, we can define the sheaf epimorphism of groupoids:

$$j_k : \text{Aut}(\mathcal{M}) \longrightarrow \Gamma_k(\mathcal{M}).$$

Finally, we define the source map a_k and the target map b_k in $\underline{J_k(\mathcal{M})}$ by:

$$\begin{aligned} a_k : \quad & \underline{J_k(\mathcal{M})} \longrightarrow \mathcal{M} \\ [\sigma_k]_x & \longrightarrow x, \\ b_k : \quad & \underline{J_k(\mathcal{M})} \longrightarrow \mathcal{M} \\ [\sigma_k]_x & \longrightarrow \beta_k \circ \sigma_k(x), \end{aligned}$$

and the canonical projection Π_q^p ($p \geq q$) by:

$$\Pi_q^p : \quad \begin{aligned} & \underline{J_p(\mathcal{M})} \longrightarrow \underline{J_q(\mathcal{M})} \\ & [f_p]_x \longrightarrow [f_q]_x. \end{aligned}$$

Thus, at the sheafs level, the non-linear finite Lie groupoid $\widehat{\mathcal{R}}_3$ of the conformal pseudogroup $\Gamma_{\widehat{G}}$ is the set of germs of continuous sections in $\Gamma_3(\mathcal{M})$ satisfying the algebraic equations over each point $x \in \mathcal{M}$, obtained by substituting the germs $[\hat{f}_3]_x \in \Gamma_3(\mathcal{M})$ for the derivatives of \hat{f} up to order three in the system of PDE (18)-(19). In other terms, we substitute $[\hat{f}_3]$ for $j_3([\hat{f}])$. More precisely, one factorizes each differential operator of the system (18)-(19) with the operators j_k ($k = 1, 2, 3$). Then, one defines the morphisms:

$$\begin{cases} M(\hat{\omega}) : j_1(\mathcal{M}) \longrightarrow S_2 T^* \mathcal{M} \\ L(j_1(\hat{\omega})) : j_2(\mathcal{M}) \longrightarrow T\mathcal{M} \otimes T^* \mathcal{M} \otimes J_1^*(T\mathcal{M}) \\ K(j_2(\hat{\omega})) : j_3(\mathcal{M}) \longrightarrow T\mathcal{M} \otimes T^* \mathcal{M} \otimes J_1^*(T\mathcal{M}) \otimes J_2^*(T\mathcal{M}) \end{cases}$$

by the respective following relations:

$$\begin{cases} \hat{f}^* \hat{\omega} = M(\hat{\omega}) \circ j_1(\hat{f}) \\ \hat{f} \hat{\nabla} = L(j_1(\hat{\omega})) \circ j_2(\hat{f}) \\ \hat{f} \hat{\nabla} \hat{f} \hat{\nabla} = K(j_2(\hat{\omega})) \circ j_3(\hat{f}), \end{cases}$$

and consequently $\widehat{\mathcal{R}}_3$ can be rewritten as the system of PDE made of the two systems of PDE $\widehat{\mathcal{R}}_2$:

$$\begin{cases} [\hat{f}_1] \hat{\omega} \stackrel{\text{def.}}{=} M(\hat{\omega})([\hat{f}_1]) = \hat{\omega} \text{ with } \det([\hat{f}_1]) \neq 0 \text{ and } \det([j_1(\hat{f})]) \neq 0 \\ [\hat{f}_2] \widehat{\nabla} \stackrel{\text{def.}}{=} L(j_1(\hat{\omega}))([\hat{f}_2]) = \widehat{\nabla}, \end{cases} \quad (20)$$

completed with the third system:

$$[\hat{f}_3] (\widehat{\nabla} \widehat{\nabla}) \stackrel{\text{def.}}{=} K(j_2(\hat{\omega}))([\hat{f}_3]) = \widehat{\nabla} \widehat{\nabla}. \quad (21)$$

$\Gamma_{\widehat{G}}$ is then the set of germs $[\hat{f}] \in Aut(\mathcal{M})$ such as $j_3([\hat{f}]) \in \widehat{\mathcal{R}}_3$. Now since $\widehat{M}_3 = 0$, one also has the equivalence $\widehat{\mathcal{R}}_3 \simeq \widehat{\mathcal{R}}_2$ and in order to work out the sophisticated non-linear Spencer complex of $\widehat{\mathcal{R}}_3$, it is sufficient, as we shall see later, to obtain it for $\widehat{\mathcal{R}}_2$. That is because the exactness of this complex of length two, only needs the symbol of the corresponding Lie groupoid to be 2-acyclic. It is precisely the case for \widehat{M}_2 . Thus, the discussion in what follows will concern exclusively $\widehat{\mathcal{R}}_2$ defined by the “deformed” or “extended” system: $\forall X, Y \in C^2(T\mathcal{M})$ and $\forall [\alpha_2] = ([\alpha], [\beta], [\mu]) \in \underline{J}_2$,

$$\widehat{\mathcal{R}}_2 : \begin{cases} [\hat{f}_1] \hat{\omega} = e^{2[\alpha]} \omega \text{ with } \det([\hat{f}_1]) \neq 0 \text{ and } \det([j_1(\hat{f})]) \neq 0 \\ [\hat{f}_2] \nabla_X Y = \nabla_X Y + [\beta](X)Y + [\beta](Y)X - \omega(X, Y)_*[\beta] \\ [\hat{f}_1] \sigma(X, Y) = \sigma(X, Y) + (n-2) \left([\beta](X)[\beta](Y) - \frac{1}{2}\omega(X, Y)[\beta]_*[\beta] \right. \\ \left. - [\mu](X, Y) + \frac{1}{2}[\beta](\nabla_X Y + \nabla_Y X) \right). \end{cases} \quad (22)$$

Let us precise again that the equation (11) makes sense, and in the formula (22) the last equation must be considered. Indeed, in case of a conformally non-flat background metric ω , the Lie form of $\widehat{\mathcal{R}}_2$ given by the set of equations (15), (16) and (17), is equivalent

to the set of equations made of the two first equations in formula (22) together with the equation (6). Then setting $\hat{\rho} = \tau = 0$ (i.e. a conformally flat metric ω), doesn't change this equivalence, but in that case the equation (6) becomes equivalent to equation (11), and thus the "extended form" of $\widehat{\mathcal{R}}_2$ is the formula (22) and we have one equation more than in the Lie form case. The latter point is rather important to make the difference between the two forms. Obviously the groupoid $\mathcal{R}_2 \subset \widehat{\mathcal{R}}_2$ of the Poincaré pseudogroup corresponds to the case for which $[\alpha_2] = 0$. The symbol M_2 of \mathcal{R}_2 vanishes and so is involutive, and \mathcal{R}_2 is not formally integrable unless ρ_s is a constant. The present suggested model is associated to a particular split exact short sequence of groupoids (not of Lie groupoids because of \mathcal{R}_2):

$$1 \longrightarrow \mathcal{R}_2 \longrightarrow \widehat{\mathcal{R}}_2 \longrightarrow \mathcal{R}_2 \longrightarrow 1,$$

and in order to have a relative exact non-linear (even so fractional!) complex associated to \mathcal{R}_2 , the complex associated to \mathcal{R}_2 would also have to be exact. This is possible only if \mathcal{R}_2 is formally integrable and consequently involutive ($M_2 = 0$). Setting these conditions it follows that the relation (12) must be satisfied at the sheaf level. Then from relation (12) and (22), one deduces \mathcal{R}_2 is the set of elements $[\alpha_2] \in \underline{J}_2$ such as $\forall X, Y \in T\mathcal{M}$ and $\forall c_0 \in \mathbb{R}$:

$$\mathcal{R}_2 : \quad \left\{ \begin{array}{lcl} [\mu](X, Y) & = & \frac{1}{2} \left\{ \left[c_0 \left(1 - e^{2[\alpha]} \right) - [\beta](_*[\beta]) \right] \omega(X, Y) \right. \\ & & \left. + [\beta](\nabla_X Y + \nabla_Y X) \right\} - [\beta](X)[\beta](Y), \end{array} \right. \quad (23)$$

and only with these conditions does $\widehat{\mathcal{R}}_2$ reduce indeed to the system chosen by J.-F. Pommaret. Thus, $[\mu]$ is completely defined from $[\alpha]$ and $[\beta]$ so that the symbol M_2 of \mathcal{R}_2 obviously vanishes. Hence, \mathcal{R}_2 is involutive and one has the equivalence:

$$\mathcal{R}_2 \simeq \mathcal{R}_1 = \underline{J}_1,$$

deduced from the embedding of \underline{J}_1 in \underline{J}_2 defined by the system (23). Consequently, one has to work out the complex associated to \mathcal{R}_1 such that:

$$1 \longrightarrow \mathcal{R}_2 \longrightarrow \widehat{\mathcal{R}}_2 \xrightarrow{\phi_0} \mathcal{R}_1 \longrightarrow 1.$$

The sequences above are sequences of groupoids but not of Lie groupoids. That $\mathcal{R}_2 \equiv \underline{J}_1$ is a groupoid can be seen directly from the definition of ϕ_0 as we shall see in the sequel or first by considering locally, above each pair of open subsets $U \times V \subset \mathcal{M} \times \mathcal{M}$ the corresponding associated trivial local groupoids $U \times G_{U \times V} \times V \simeq \mathcal{R}_{2/U \times V}$ (and with analogous expressions for the other \mathcal{R} 's). Then we obtain a corresponding sequence of algebraic groups on a finite projective free module for the G 's. Since they are algebraic they are splittable and the sequence is split exact. Therefore we can find a splitting of groups (such as an Iwasawa decomposition for instance), by a good choice of back-connection. Then $\mathcal{R}_{1/U \times V}$ can be canonically injected in $\widehat{\mathcal{R}}_{2/U \times V}$ so that it acquires locally an algebraic group structure. Then, by gluing over all pairs of open subsets $U \times V$, we deduce the sequence of groupoids. Let us add from the definition of $\widehat{\mathcal{R}}_2$ that \mathcal{R}_1 is a natural bundle associated to $\widehat{\mathcal{R}}_2$. From a physical point of view, it is important to notice

that μ may be considered as an Abraham-Eötvos type tensor, leading to a first physical interpretation (up to a constant for units) of β as the acceleration 4-vector of gravity. On the other hand, completely in agreement with J.-F. Pommaret, α being associated with the dilatations it can be considered as a relative Thomson type temperature (again up to a constant for units):

$$\alpha = \ln(T_0/T), \quad (24)$$

where T_0 is a constant temperature of reference associated with the base space-time \mathcal{M} . Then, one can easily define the epimorphism ϕ_0 by the relations $\forall [\hat{f}_2] \in \widehat{\mathcal{R}}_2$:

$$\phi_0([\hat{f}_2]) = \begin{cases} [\alpha] &= \frac{1}{n} \ln |\det([\hat{f}_1])| \\ [\beta] &= \frac{1}{n} Tr_\omega ([\hat{f}_2] \nabla - \nabla) \end{cases}. \quad (25)$$

Thus, we get a first diagram:

$$\begin{array}{ccccc} & 1 & & 1 & \\ & \downarrow & & \downarrow & \\ 1 & \longrightarrow & \Gamma_G & \xrightarrow{j_2} & \mathcal{R}_2 \\ & \downarrow & & \downarrow & \\ 1 & \longrightarrow & \Gamma_{\widehat{G}} & \xrightarrow{j_2} & \widehat{\mathcal{R}}_2 \\ & \downarrow & & \downarrow \phi_0 & \\ 1 & \longrightarrow & \theta & \xrightarrow{j_1} & \underline{J}_1 \\ & & & & \downarrow \\ & & & & 0 \end{array} \quad (26)$$

Before presenting the various non-linear Spencer complexes we shall recall briefly certain definitions and results of this theory. The Spencer Theory presentation we give below is rather minimal since we think that it is impossible to describe it perfectly in few pages. It is especially a matter of indicating the notations chosen in the text and we do not claim to make a full and complete description. Moreover, this theory is presented and taken up historically with the “diagonal method” of Grothendieck (Kumpera *et al.* 1972), and the results and definitions we give don’t mention it. In this method, two copies of the base space \mathcal{M} are used. The first one \mathcal{M}_1 (the horizontal component) is attributed to the set of “points” on which the Taylor coefficients of particular Taylor series are defined, and the second \mathcal{M}_2 (the vertical component), to points on which these previous series are evaluated. The independence of this evaluation with respect to the points chosen in \mathcal{M}_1 allow to deduce the first Spencer differential operator (linear or non-linear) as an exterior differential operator on the horizontal component. Then the second Spencer differential

operator and a particular set of derivations are deduced from the equivariance of the first Spencer differential operator with respect to a particular groupoid action. Thus, the Spencer cohomology can be seen mainly as an equivariant cohomology on graded sheaves of the diagonal of $\mathcal{M}_1 \times \mathcal{M}_2$. Actually, we give the results and formulas after the vertical (diagonal isomorphism) projection on the vertical component of various diagonal graded sheaves defined on $\mathcal{M}_1 \times \mathcal{M}_2$ and following in parts a formulation given by J.-F. Pommaret especially when concerning the definition of the differential bracket. The other definitions are merely the vertical “translations” of the definitions given by Kumpera and Spencer applying the so-called “ ϵ ” isomorphism on the diagonal sheafs.

3.3. The first non-linear Spencer differential operators

First, we recall that $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} J_k(E)$, where E is a vector bundle, has a natural left $\underline{\Lambda T^*\mathcal{M}}$ -module structure.

Definition 1 a) The linear Spencer operator D is the unique differential operator (\mathbb{R} -linear sheaf map; $k, s \geq 0$)

$$D : \underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} J_{k+1}(E) \longrightarrow \underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} J_k(E),$$

satisfying the three following conditions:

1. $D \circ j_{k+1} = 0$,
2. $\forall \tau_{k+1} \in \underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} J_{k+1}(E)$, $D(\omega \wedge \tau_{k+1}) = \mathbf{d}\omega \wedge \tau_k + (-)^{d^0\omega} \omega \wedge D\tau_k$
where $\omega \in \underline{\Lambda T^*\mathcal{M}}$ is any homogeneous differential form and \mathbf{d} being the exterior differential,
3. D restricted to $J_{k+1}(E)$ satisfies:

$$\epsilon_1 \circ D = j_1 \circ \Pi_k^{k+1} - id_{J_{k+1}(E)},$$

where ϵ_1 is the monomorphism defined by the short exact sequence:

$$0 \longrightarrow T^*\mathcal{M} \otimes J_k(E) \xrightarrow{\epsilon_1} J_1(J_k(E)) \xrightarrow{\Pi_0^1} J_k(E) \longrightarrow 0.$$

b) The restriction of $(-D)$ to the symbol $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} S_k T^*\mathcal{M} \otimes_{\theta} E$ defines the $\underline{\Lambda T^*\mathcal{M}}$ -linear Spencer map $\underline{\delta}$ such that

$$\underline{\delta}(\omega \wedge \tau_k) = (-)^{d^0\omega} \omega \wedge \underline{\delta}(\tau_k),$$

with ω and τ_k as in a)-3..

■

Definition 2 We define the r -th Spencer sheaf ($r \geq 1$) of $J_k(E)$, the quotient sheaf

$$\mathcal{C}_k^r(E) = \underline{\Lambda^r T^* \mathcal{M} \otimes_{\theta} J_k(E)} / \zeta_k \circ \underline{\delta}(\underline{\Lambda^{r-1} T^* \mathcal{M} \otimes_{\theta} S_{k+1} T^* \mathcal{M} \otimes_{\theta} E}),$$

where ζ_k is the monomorphism defined by the short exact sequence:

$$0 \longrightarrow S_k T^* \mathcal{M} \otimes E \xrightarrow{\zeta_k} J_k(E) \xrightarrow{\Pi_{k-1}^k} J_{k-1}(E) \longrightarrow 0.$$

■

Let us add that $\mathcal{C}_k^r(E)$ has a module structure on the θ -algebra \underline{J}_k and we set $\mathcal{C}_k^r(E) = 0$ for $r > n$ and $\mathcal{C}_k^0(E) = \underline{J}_k(E)$.

Definition 3 The operator D can be factorized with a right θ -linear operator D' on $\mathcal{C}_k^r(E)$:

$$D' : \mathcal{C}_k^r(E) \longrightarrow \mathcal{C}_k^{r+1}(E).$$

■

Contrary to the operator D , there is no loss of order on k . More precisely, D' is such that the following diagram of split exact sequences is commutative:

$$\begin{array}{ccc}
0 & & 0 \\
\downarrow & & \downarrow \\
\underline{S_{k+1} T^* \mathcal{M} \otimes_{\theta} E} & \xrightarrow{(-\delta)} & \underline{T^* \mathcal{M} \otimes_{\theta} S_k T^* \mathcal{M} \otimes_{\theta} E} \\
\downarrow \zeta_{k+1} & & \downarrow id \otimes \zeta_k \\
\underline{J_{k+1}(E)} & \xrightarrow{D} & \underline{T^* \mathcal{M} \otimes_{\theta} J_k(E)} \\
\downarrow \Pi_k^{k+1} & & \downarrow \rho_k^{k+1} \\
\underline{J_k(E)} & \xrightarrow{D'} & \mathcal{C}_k^1(E) \\
\downarrow & & \downarrow \\
0 & & 0
\end{array}$$

It is to be noted that the sequences being split, then D' is built up from a connexion

$$c_{k+1}^k : \underline{J_k(E)} \longrightarrow \underline{J_{k+1}(E)},$$

such that by definition

$$\Pi_k^{k+1} \circ c_{k+1}^k = id_k.$$

But quotienting, then by definition, D' is independent of the choice of connexion c_{k+1}^k . Hence, whatever is c_{k+1}^k , one has

$$D' = \rho_k^{k+1} \circ D \circ c_{k+1}^k.$$

Finally, for $r \geq 1$, these definitions can be extended to the tangent bundle R_k of \mathcal{R}_k instead of $J_k(E)$, and to its corresponding symbol M_k . But in this case, from the definition of D' , 1) \mathcal{R}_{k+1} must be a fibered manifold, 2) to make a choice of connexion c_{k+1}^k we must have the epimorphism $\mathcal{R}_{k+1} \rightarrow \mathcal{R}_k \rightarrow 0$, and 3) the system \mathcal{R}_k must be formally transitive, i.e. we must have the epimorphism $\mathcal{R}_k \rightarrow \mathcal{M} \rightarrow 0$. Also we shall use the definitions and notations:

$$\mathcal{C}_k^r = \underline{\Lambda}^r T^*\mathcal{M} \otimes_{\theta} R_k / \zeta_k \circ \underline{\delta}(\underline{\Lambda}^{r-1} T^*\mathcal{M} \otimes_{\theta} M_{k+1}).$$

Definition 4 Let us define $B_k^r(\mathcal{M})$ and $B_k(\mathcal{M})$ such that

$$B_k^r(\mathcal{M}) = \underline{\Lambda}^r T^*\mathcal{M} \otimes_{\theta} J_k,$$

and

$$B_k(\mathcal{M}) = \bigoplus^r B_k^r(\mathcal{M}).$$

■

$B_k(\mathcal{M})$ has a natural structure of graded J_k -algebra defined by the exterior product of the J_k -valued forms, since we have the equivalence:

$$B_k(\mathcal{M}) \equiv \underline{\Lambda} T^*\mathcal{M} \otimes_{\theta} J_k.$$

$B_k(\mathcal{M})$ inherits also a natural structure of left graded $\Lambda T^*\mathcal{M}$ -module where the external operation on $\bigoplus^{r>0} B_k^r(\mathcal{M})$ is the exterior product $\omega \wedge \mu$ of $\omega \in \Lambda T^*\mathcal{M}$ and $\mu \in B_k(\mathcal{M})$ with respect to the pairing on $B_k^0(\mathcal{M})$:

$$([f], [g_k]) \in \theta \times \underline{J}_k \longrightarrow j_k([f]).[g_k] \in \underline{J}_k.$$

Let us denote by $Der_a B_k(\mathcal{M})$ the sheaf of germs of “admissible” graded derivations \mathcal{D} of $B_k(\mathcal{M})$, i.e. if \mathcal{D} is of degree p , one has:

$$\mathcal{D}(\underline{J}_k^0) \subset \underline{\Lambda}^p T^*\mathcal{M} \otimes_{\theta} \underline{J}_k^0,$$

$$\mathcal{D}(\underline{\Lambda}^r T^*\mathcal{M}) \subset \underline{\Lambda}^{r+p} T^*\mathcal{M},$$

where \underline{J}_k^0 is the kernel of the target map β_k defined on \underline{J}_k :

$$\beta_k : \underline{J}_k \longrightarrow \theta.$$

Obviously $Der_a B_k(\mathcal{M})$ is endowed with the bracket of derivations $[\quad , \quad]$, i.e. if \mathcal{D}_i is a derivation of degree p_i , then we have:

$$[\mathcal{D}_1, \mathcal{D}_2] = \mathcal{D}_1 \circ \mathcal{D}_2 - (-)^{p_1 p_2} \mathcal{D}_2 \circ \mathcal{D}_1.$$

Finally, we also have

$$Der_a B_k(\mathcal{M}) = \bigoplus^r Der_a^r B_k(\mathcal{M}),$$

where $Der_a^r B_k(\mathcal{M})$ is the module of admissible derivations of degree r on $B_k(\mathcal{M})$.

Definition 5 One defines \mathbf{D}_1 (the “twisting” of \mathbf{d}), the non-linear differential operator

$$\mathbf{D}_1 : \Gamma_{k+1}\mathcal{M} \longrightarrow Der_a^1 B_k(\mathcal{M})$$

such that:

$$\mathbf{D}_1[f_{k+1}] = \mathbf{d} - Ad[f_{k+1}] \circ \mathbf{d} \circ Ad[f_{k+1}^{-1}],$$

where $Ad[f_{k+1}]$ is the contravariant action at the sheaf level of $[f_{k+1}]$ on $\underline{\Lambda T^*\mathcal{M}} \otimes_\theta J_k$ corresponding to the action of the pull-back of $f \in Aut(\mathcal{M})$ on the tensors of $\Lambda T^*\mathcal{M} \otimes J_k$. ■

To this action on $\Lambda T^*\mathcal{M} \otimes J_k$ corresponds simultaneously an action of the “pull-back-push-forward of $f \in Aut(\mathcal{M})$ ” on the tensors of $\Lambda T^*\mathcal{M} \otimes J_k(\mathcal{M})$. Also, we deduce and define at the sheaf level, the extension of $Ad[f_{k+1}]$ on $\underline{\Lambda T^*\mathcal{M}} \otimes_\theta J_k(\mathcal{M})$.

Definition 6 Let

$$\mathbf{D}'_1 : \Gamma_k\mathcal{M} \longrightarrow \mathcal{C}_k^1(T\mathcal{M}),$$

be “the first non-linear Spencer operator” such that $\forall [f_k] \in \Gamma_k\mathcal{M}$:

$$\mathbf{D}'_1([f_k]) = \rho_k^{k+1} \circ [\mathbf{d} - Ad[f_{k+1}] \circ \mathbf{d} \circ Ad[f_{k+1}^{-1}]](id_k), \quad (27)$$

where $[f_{k+1}] = c_{k+1}^k([f_k])$ and $id_k \in \underline{J_k(\mathcal{M})}$ is the prolongation up to order k of $id_{\mathcal{M}} \equiv id_0$. ■

Sometimes this definition is given using the functor j_1 instead of \mathbf{d} . The result is that the derivation is made with respect to the source and not the target and difficulties appear when defining the brackets given further. Then Spencer defined the isomorphism ad of degree zero:

Definition 7 One defines the isomorphism ad , the operator

$$ad : \mathcal{C}_{k+1}^\bullet(T\mathcal{M}) \longrightarrow Der_a B_k(\mathcal{M})$$

such that $\forall v \in B_k(\mathcal{M})$ and $\forall u_{k+1} \in \mathcal{C}_{k+1}^r(T\mathcal{M})$:

$$1. ad(\mathcal{C}_{k+1}^\bullet(T\mathcal{M})) \stackrel{def.}{=} Der_\Sigma B_k(\mathcal{M}) = Der_{\Sigma,k}^\bullet \subseteq Der_a B_k(\mathcal{M}),$$

2. $\mathcal{L} : \Lambda T^*\mathcal{M} \otimes_{\theta} J_k(T\mathcal{M}) \longrightarrow \text{Der}_a B_k(\mathcal{M})$ being the Lie derivative of degree zero such that

$$\mathcal{L}(u_k)v = [i(u_k), \mathbf{d}]v \equiv u_k \bar{\wedge} \mathbf{d}v + (-)^r \mathbf{d}(u_k \bar{\wedge} v),$$

where $\bar{\wedge}$ is the extended Frölicher-Nijenhuis product and i the interior product:

$$i(u_k)v \equiv u_k \bar{\wedge} v,$$

3. $\text{ad}(u_{k+1})v = [\mathcal{L}(u_k) + (-)^{r+1} D'(u_{k+1}) \bar{\wedge}] v.$

■

Initially, ad is the θ -linear sheaf map corresponding to the $(k+1)$ -st order differential operator $\mathcal{L} \circ j_k$ where $\mathcal{L} : J_k(T) \longrightarrow \text{Der}_a B_k(\mathcal{M})$ is the Lie derivative of degree zero and of order 1. Then, one obtains the first step of the sophisticated non-linear Spencer complexes associated to the resolutions $\mathcal{C}_{k+1}^\bullet(T\mathcal{M})$ or $\text{Der}_{\Sigma,k}^\bullet$:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \text{Aut}(\mathcal{M}) & \xrightarrow{j_{k+1}} & \Gamma_{k+1}\mathcal{M} & \xrightarrow{\mathbf{D}'_1} & \mathcal{C}_{k+1}^1(T\mathcal{M}) \\ & & \parallel & & \parallel & & \downarrow \text{ad} \\ 1 & \longrightarrow & \text{Aut}(\mathcal{M}) & \xrightarrow{j_{k+1}} & \Gamma_{k+1}\mathcal{M} & \xrightarrow{\mathbf{D}_1} & \text{Der}_{\Sigma,k}^1 \end{array}$$

where the two horizontal sequences are split exact at $\Gamma_{k+1}\mathcal{M}$. From this, we can determine the induced fractional differential operator \mathbf{D}'_1 such that the following diagram is commutative ($M_3 = \widehat{M}_3 = 0$, $\mathcal{M}_2 = 0$):

$$\begin{array}{ccccccc} & & 1 & & 1 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 1 & \longrightarrow & \Gamma_G & \xrightarrow{j_2} & \mathcal{R}_2 \xrightarrow{\mathbf{D}'_1} \mathcal{C}_2^1 = \underline{T^*\mathcal{M}} \otimes_{\theta} R_2 \\ & & & & \downarrow & & \downarrow \\ & & 1 & \longrightarrow & \Gamma_{\widehat{G}} & \xrightarrow{j_2} & \widehat{\mathcal{R}}_2 \xrightarrow{\widehat{\mathbf{D}}'_1} \widehat{\mathcal{C}}_2^1 = \underline{T^*\mathcal{M}} \otimes_{\theta} \widehat{R}_2 & (28) \\ & & & & \downarrow & & \downarrow \\ & & 1 & \longrightarrow & \theta_{\mathcal{M}} & \xrightarrow{j_1} & \underline{J}_1 \xrightarrow{\mathbf{D}'_1} \mathcal{C}_1^1 = B_1^1(\mathcal{M}) \\ & & & & \downarrow & & \downarrow \\ & & & & 1 & & 0 \end{array}$$

Obviously, one will determine also ϕ_1 and in the sequel, one will call $B_1^1(\mathcal{M})$ the sheaf of electromagnetic and gravitational gauge potentials.

4. The potentials of interaction and the metric

4.1. The electromagnetic and gravitational potentials

In order to lighten the presentation of the results, first let us consider the following notations:

- 1) one will merely write f_k and $j_k(f)$ instead of respectively $[f_k]$ and $j_k([f])$,
- 2) one will denote $T^q f$ the restriction in $\underline{S_q T^* \mathcal{M} \otimes_\theta T \mathcal{M}}$ of $[f_p] \in \underline{J_p(\mathcal{M})}$ ($p \geq q \geq 1$),
- 3) one will denote $T(T^r f)$ the restriction in $\underline{T^* \mathcal{M} \otimes_\theta S_r T^* \mathcal{M} \otimes_\theta T \mathcal{M}}$ of $j_1([f_s]) \in \underline{J_1(J_s(\mathcal{M}))}$ ($s \geq r \geq 0$),
- 4) one sets analogous conventions concerning the presence of ordinary parenthesis in the notations for the differential of germs of the tangent maps $dT^q f$ and $d(T^q f)$ (let us note in these notations that d is not the exterior differential that one denotes by \mathbf{d} , but stands for differential maps).

Then, one has the following set of results $\forall \hat{f}_2 \in \widehat{\mathcal{R}}_2$ and $\forall X \in \mathcal{C}^1(T\mathcal{M})$:

$$\begin{aligned} Tr^1(j_1(\hat{f}_1)\nabla_X) &= \frac{1}{2} \sum_{i,j=1}^n \tilde{\omega}^{i,j} \left\{ \langle d(\omega) \circ \hat{f} | T(\hat{f}).X \rangle | T\hat{f}.e_i \otimes T\hat{f}.e_j \rangle \right. \\ &\quad \left. + 2 \langle \omega \circ \hat{f} | d(T\hat{f}) | X \rangle .e_i \otimes T\hat{f}.e_j \rangle \right\} \end{aligned} \quad (29)$$

$$= Tr^1(\nabla_X) + n \langle \mathbf{d}(\alpha) | X \rangle, \quad (30)$$

and

$$\begin{aligned} Tr^1(\hat{f}_2\nabla_X) &= \frac{1}{2} \sum_{i,j=1}^n \tilde{\omega}^{i,j} \left\{ \langle d(\omega) \circ \hat{f} | T\hat{f}.X \rangle | T\hat{f}.e_i \otimes T\hat{f}.e_j \rangle \right. \\ &\quad \left. + 2 \langle \omega \circ \hat{f} | d(T\hat{f}) | X \rangle .e_i \otimes T\hat{f}.e_j \rangle \right\} \end{aligned} \quad (31)$$

$$= Tr^1(\nabla_X) + n \langle \beta | X \rangle. \quad (32)$$

Now, let $\hat{\chi}^{(2)}$ be an element of $J_1(J_2(T\mathcal{M}))$, and its components $\hat{\chi}_q$ ($q = 0, 1, 2$), i.e. the restrictions of $\hat{\chi}^{(2)}$ to $\underline{T^* \mathcal{M} \otimes_\theta S_q T^* \mathcal{M} \otimes_\theta T \mathcal{M}}$ such that $\forall \hat{f}_3 \in \widehat{\mathcal{R}}_3$:

$$\hat{\chi}^{(2)} = \hat{f}_3^{-1} \circ j_1(\hat{f}_2) - id_3, \quad (33)$$

where by abuse of notations id_3 is the image of $id_3 \in J_3(T\mathcal{M})$ by the canonical injection $J_3(T\mathcal{M}) \longrightarrow J_1(J_2(T\mathcal{M}))$. In particularly, one has the relation (Pommaret 1994):

$$\begin{aligned} \hat{\chi}_0 &= \hat{f}_1^{-1} \circ j_1(\hat{f}) - id_1 \\ &\equiv \hat{A} - id_1. \end{aligned}$$

It follows that $\widehat{\mathbf{D}}'_1$ satisfies:

$$\epsilon_1 \circ \widehat{\mathbf{D}}'_1(\hat{f}_2) \equiv \hat{\tau}^{(2)} = \hat{\chi}^{(2)} \circ (\hat{B} \otimes id_2), \quad (34)$$

with $\hat{B} = \hat{A}^{-1}$ and $\hat{f}_3 = c_3^2(\hat{f}_2) \in \widehat{\mathcal{R}}_3$. In particular, $\hat{\tau}_0$ and $\hat{\tau}_1$ satisfy the relations:

$$\hat{\tau}_0 = id_{T\mathcal{M}} - T\hat{f} \circ T(\hat{f})^{-1} \equiv id_{T\mathcal{M}} - \hat{B} = \hat{\chi}_0 \cdot \hat{B} \in T\mathcal{M} \otimes T^*\mathcal{M},$$

and $\forall X, Y \in T\mathcal{M}$ (Pommaret 1994):

$$\langle dT\hat{f}|X\rangle \circ \hat{\tau}_0 \cdot Y + T\hat{f} \circ \langle \hat{\tau}_1|Y\rangle \cdot X = \langle d(T\hat{f}) - dT\hat{f}|\hat{B}.Y\rangle \cdot X,$$

where $\langle dT\hat{f}|X\rangle, \langle \hat{\tau}_1|Y\rangle \equiv \langle \hat{\chi}_1|\hat{B}.Y\rangle$ and $\langle d(T\hat{f}) - dT\hat{f}|\hat{B}.Y\rangle$, are considered as elements of $T\mathcal{M} \otimes T^*\mathcal{M}$. Then, if one substitutes

$$T\hat{f}^{-1} \circ T(\hat{f}).X = (\hat{\chi}_0 + id).X$$

for X in the relations (31)-(32), and also considering the Schwarz equalities:

$$\langle dT\hat{f}|X\rangle \cdot Y = \langle dT\hat{f}|Y\rangle \cdot X,$$

one obtains by subtracting the result from the relations (29)-(30) $\forall X \in T\mathcal{M}$:

$$\frac{1}{n} Tr^1 \left[\langle \hat{\chi}_1|X\rangle + \nabla_{\hat{\chi}_0.X} \right] = \langle \mathbf{d}(\alpha) - \beta|X\rangle - \langle \beta|\hat{\chi}_0.X\rangle. \quad (35)$$

From this latter relation, one can define the electromagnetic potential vector $\hat{\mathcal{A}} \in \underline{T^*\mathcal{M}}$ by $\forall X \in \underline{T\mathcal{M}}$:

$$\langle \hat{\mathcal{A}}|X\rangle = \frac{1}{n} Tr^1 \left[\langle \hat{\tau}_1|X\rangle + \nabla_{\hat{\tau}_0.X} \right] = \langle \mathbf{d}(\alpha)|\hat{B}.X\rangle - \langle \beta|X\rangle. \quad (36)$$

In an orthonormal system of coordinates, the latter definition becomes ($i, j, k = 1, \dots, n$):

$$\hat{\mathcal{A}}_i = \frac{1}{n} \left(\hat{\tau}_{k,i}^k + \hat{\tau}_{,i}^k \gamma_{j,k}^j \right) = \hat{B}_i^k \partial_k \alpha - \beta_k,$$

where γ is the Riemann-Christoffel 1-form associated to ω and thus satisfying:

$$Tr^1(\nabla_X) = Tr^1(\gamma(X)).$$

Prolonging the relation (35) (one does not prolong the relation (36) because $\hat{\tau}_2$ is not the first prolongation of $\hat{\tau}_1$, contrary to $\hat{\chi}_2$), one deduces and defines the mixed tensor potential of gravitation and electromagnetic $\hat{\mathcal{B}} \in \overset{2}{\otimes} \underline{T^*\mathcal{M}}$ such as $\forall X, Y \in T^*\mathcal{M}$:

$$\begin{aligned} \langle \hat{\mathcal{B}}|Y \otimes X\rangle &= \frac{1}{n} Tr^1 [i_Y \langle \hat{\tau}_2|X\rangle + \langle d(\gamma)|Y \otimes \hat{\tau}_0.X\rangle \\ &\quad + \gamma(\langle \hat{\tau}_1|X\rangle \cdot Y)] \\ &= \langle \mathbf{d}(\beta)|\hat{B}.X \otimes Y\rangle - \langle \beta|\langle \hat{\tau}_1|X\rangle \cdot Y\rangle \\ &\quad - \langle \mu|X \otimes Y\rangle \end{aligned} \quad (37)$$

where i_Y is the interior product by Y and $\hat{\tau}_2$ satisfies $\forall \hat{f}_3 \in \widehat{\mathcal{R}}_3$ and $\forall X, Y, Z \in T\mathcal{M}$:

$$\begin{aligned} << d(dT\hat{f}) - d^2T\hat{f} | \widehat{\mathcal{B}}.X > | Y \otimes Z > = \\ & < d^2T\hat{f} | Y \otimes Z > \circ \hat{\tau}_0 . X + < dT\hat{f} | Y > \circ < \hat{\tau}_1 | X > . Z \\ & + < dT\hat{f} | Z > \circ < \hat{\tau}_1 | X > . Y + T\hat{f} \circ << \hat{\tau}_2 | X > | Y \otimes Z > . \end{aligned}$$

Again, in an orthonormal system of coordinates $(i, j, k, h = 1, \dots, n)$:

$$\widehat{\mathcal{B}}_{j,i} = \frac{1}{n} \left(\hat{\tau}_{k,j,i}^k + \hat{\tau}_{j,i}^k \gamma_{h,k}^h + \hat{\tau}_{,i}^k (\partial_k \gamma_{h,j}^h) \right) = \widehat{B}_i^k \partial_k \beta_j - \mu_{ij} - \hat{\tau}_{j,i}^k \beta_k.$$

This definition for $\widehat{\mathcal{B}}$ hasn't been determined by J.-F. Pommaret (1994), he also gave a different definition for $\widehat{\mathcal{A}}$ rather associated to the relation (35). In conclusion to this chapter, one notices that \mathcal{D}'_1 (remaining to explicit) appears to be a Fröbenius-type operator and depends on $\hat{\tau}^{(1)}$ itself depending on $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{B}}$ as we shall see further. Lastly, ϕ_1 is quite defined by the relations (36) and (37), and if $n = 4$, one has 20 scalar gauge potentials. On the other hand, one can see as well that the definitions of $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{B}}$ can be deduced from the conformal Killing equations on $\underline{J}_1(T\mathcal{M})$, namely $\forall \xi_{(1)} \equiv (\xi_0, \xi_1) \in \underline{J}_1(T\mathcal{M})$ and $\forall \eta \in \theta$ then one has:

$$K_0(\xi^{(1)}) \stackrel{\text{def.}}{=} \frac{1}{n} Tr^1(\xi_1 + \gamma(\xi_0)) = \eta,$$

where K_0 is the conformal Killing operator. If K_1 is its first prolongation then setting $K^{(1)} \equiv (K_0, K_1)$ one obtains obviously from the equation above $\forall X \in \underline{T\mathcal{M}}$ and $\forall \hat{\tau}^{(2)} \in \widehat{\mathcal{C}}_2^1(T\mathcal{M})$:

$$\phi_1(\hat{\tau}^{(2)})(X) = K^{(1)}(< \hat{\tau}^{(2)} | X >).$$

4.2. The morphisms ϕ_1 , \mathcal{D}'_1 and the metric of the “gauge space-time”

From the preceding chapter, obviously one easily notices by definition that \mathcal{C}_2^1 is the kernel of ϕ_1 . On the contrary \mathcal{D}'_1 , depending on $\hat{\tau}^{(1)}$, is rather more tricky to determine. The sequences being split and ϕ_1 being θ -linear, one can deduce the important relation that $\hat{\tau}^{(2)}$ can be rewritten in the following form:

$$\hat{\tau}^{(2)} = \tau^{(2)} + < \widehat{\mathcal{A}} | \Omega^{(2)} > + < \widehat{\mathcal{B}} | \mathcal{K}^{(2)} >, \quad (38)$$

where $\tau^{(2)} \in \mathcal{C}_2^1$ and $\Omega^{(2)}$ and $\mathcal{K}^{(2)}$ are elements of $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} (\widehat{\mathcal{C}}_2^1 / \mathcal{C}_2^1)$, one calls the linear susceptibilities of the vacuum or the space-time associated respectively to $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{B}}$. Then, as a definition of \mathcal{D}'_1 , one obtains the following kind of relations:

$$\begin{aligned} < a_1 | \widehat{\mathcal{A}} > + < b_1 | \widehat{\mathcal{B}} > &= D'\alpha \equiv \mathbf{d}\alpha - \beta, \\ < a_2 | \widehat{\mathcal{A}} > + < b_2 | \widehat{\mathcal{B}} > &= D'\beta \equiv \mathbf{d}\beta - \mu, \end{aligned}$$

where a_1 is affine with respect to $\mathbf{d}\alpha$, b_1 linear with respect to $\mathbf{d}\alpha$, a_2 linear with respect to $\mathbf{d}\beta$ and β , and b_2 affine with respect to $\mathbf{d}\beta$ and β . Thus, \mathbb{D}'_1 is fractional with respect to $\mathbf{d}\alpha$, $\mathbf{d}\beta$, α and β and finally only depends on the susceptibilities of $\hat{\tau}^{(2)}$. Then, the exactness condition at \underline{J}_1 :

$$\mathbb{D}'_1 \circ j_1 = 0,$$

the relations (36)-(37) and the commutativity of the diagram (28) involve the necessary condition which must be satisfied by $\tau^{(2)}$:

$$\tau_0 = 0.$$

There remains the metric ν of the “gauge (or observable or measurable) space-time” defined in considering \hat{B} as a field of tetrads $\forall X, Y \in T\mathcal{M}$:

$$\langle \nu | X \otimes Y \rangle = \langle \omega | \hat{B}.X \otimes \hat{B}.Y \rangle.$$

Thus, one has the general relation between ν and ω :

$$\nu = \omega + \text{linear and quadratic terms in } \hat{\mathcal{A}} \text{ and } \hat{\mathcal{B}}.$$

Then from this metric ν , one can deduce the Riemann and Weyl curvature tensors of the gauge space-time. One has a non-metrical theory for the gravitation in the gauge space-time, since clearly ν doesn't appear as a gravitational potential. The space-time terminology we use is quite natural in the sense that one has simultaneously two types of space-time. The first one, which we call the “underlying” or “substrat” space-time, is endowed with the metric ω and is of constant scalar curvature. The other one, called the “gauge (observable or measurable) space-time”, endowed with the metric ν , is defined for any scalar curvature and by the gauge potentials $\hat{\mathcal{A}}$ and $\hat{\mathcal{B}}$. It can be considered as the underlying space-time deformed by the gauge potentials and the Weyl curvature does not necessarily vanish, contrary to Pommaret's assertions (Pommaret 1994, see page 456 and Pommaret 1989). Moreover, from a continuum mechanics of deformable bodies point of view, the metric ν can be interpreted as the tensor of deformation of the underlying space-time (Katanaev *et al.* 1992, Kleinert 1989).

5. The fields of interaction

5.1. The second non-linear Spencer operator

Before giving an explicit expression for this operator in the complex of \underline{J}_1 , one will briefly recall its definition, but before that one needs a few other definitions (Pommaret 1989, Kumpera *et al.* 1972).

Definition 8 Let the “algebraic bracket” $\{ , \}$ on $J_{k+1}(T\mathcal{M})$, be the \mathbb{R} -bilinear map ($\forall k \geq 0$)

$$\{ , \} : J_{k+1}(T\mathcal{M}) \times_{\mathcal{M}} J_{k+1}(T\mathcal{M}) \longrightarrow J_k(T\mathcal{M})$$

such as $\forall \xi_{k+1}, \eta_{k+1} \in J_{k+1}(T\mathcal{M})$ one has:

$$\{\xi_{k+1}, \eta_{k+1}\} = \{\xi_1, \eta_1\}_k,$$

where $\{\xi_1, \eta_1\} \in T\mathcal{M}$ is the usual Lie bracket defined on $J_1(T\mathcal{M})$ and $\{\xi_1, \eta_1\}_k$ its lift in $J_k(T\mathcal{M})$. ■

Definition 9 One calls “differential Lie bracket” on $\underline{J_k(T\mathcal{M})}$, the bracket $[\ , \]$ such that:

$$[\ , \] : \underline{J_k(T\mathcal{M})} \times_{\mathcal{M}} \underline{J_k(T\mathcal{M})} \longrightarrow \underline{J_k(T\mathcal{M})},$$

and $\forall \xi_k, \eta_k \in \underline{J_k(T\mathcal{M})}$ then

$$[\xi_k, \eta_k] = \{\xi_1, \eta_1\} + i_{\xi_0} D\eta_{k+1} - i_{\eta_0} D\xi_{k+1},$$

where i is the usual interior product and ξ_{k+1} and η_{k+1} are any lifts of ξ_k and η_k in $\underline{J_{k+1}(T\mathcal{M})}$. ■

Definition 10 For any decomposable elements

$$\alpha = u \otimes \xi_k \in \underline{\Lambda^r T^*\mathcal{M}} \oplus \underline{J_k(T\mathcal{M})},$$

$$\beta = v \otimes \eta_k \in \underline{\Lambda^s T^*\mathcal{M}} \oplus \underline{J_k(T\mathcal{M})},$$

and defining on $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} \underline{J_k(T\mathcal{M})}$ the interior product i by the relation: $\forall w \in \underline{\Lambda T^*\mathcal{M}}$,

$$i_{\alpha}w = u \wedge i_{\xi_k}w,$$

then with \mathbf{d} being the exterior derivative and

1) \mathcal{L} the Lie derivative on $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} \underline{J_k(T\mathcal{M})}$ such that:

$$\mathcal{L}(\alpha) = i_{\alpha} \circ \mathbf{d} + (-)^r \mathbf{d} \circ i_{\alpha},$$

2) $ad(\alpha) = \mathcal{L}(\alpha) + (-)^{r+1} i_{D\alpha}$,

one defines the “twisted” bracket on $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} \underline{J_k(T\mathcal{M})}$ (θ -bilinear), $[\ , \]$ by the relation:

$$[\alpha, \beta] = [ad(\alpha)v] \otimes \eta_k - (-)^{rs}[ad(\beta)u] \otimes \xi_k + (u \wedge v) \otimes [\xi_k, \eta_k] \in \underline{\Lambda^{r+s} T^*\mathcal{M}} \otimes_{\theta} \underline{J_k(T\mathcal{M})}. ■$$

This bracket defines a graded Lie algebra structure on $\underline{\Lambda T^*\mathcal{M}} \otimes_{\theta} \underline{J_k(T\mathcal{M})}$.

Definition 11 One calls 2nd non-linear Spencer operator of the resolution $\text{Der}_{\Sigma,k}^\bullet$, the differential operator \mathbf{D}_2 such that:

$$\mathbf{D}_2 : \text{Der}_{\Sigma,k}^1 \longrightarrow \text{Der}_{\Sigma,k}^2,$$

and $\forall u \in \text{Der}_{\Sigma,k}^1$

$$\mathbf{D}_2 u = [\mathbf{d}, u] - \frac{1}{2}[u, u],$$

■

Definition 12 To this operator \mathbf{D}_2 corresponds the 2nd non-linear Spencer operator \mathbf{D}'_2 of the resolution $\mathcal{C}_k^\bullet(T\mathcal{M})$ such that:

$$\mathbf{D}'_2 : \mathcal{C}_k^1(T\mathcal{M}) \longrightarrow \mathcal{C}_k^2(T\mathcal{M}),$$

and $\forall \tau \in \mathcal{C}_k^1(T\mathcal{M})$

$$\mathbf{D}'_2 \tau = D' \tau - \frac{1}{2}[\tau, \tau]',$$

where $[\ , \]'$ is the quotient twisted bracket.

■

Definition 13 With the latter definitions, the “sophisticated non-linear Spencer complex of $\Gamma_{k+1}(\mathcal{M})$ ”, is the truncated split exact differential sequence in the first row of the following commutative diagram of split exact sequences:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \text{Aut}(\mathcal{M}) & \xrightarrow{j_{k+1}} & \Gamma_{k+1}(\mathcal{M}) & \xrightarrow{\mathbf{D}'_1} & \mathcal{C}_{k+1}^1(T\mathcal{M}) & \xrightarrow{\mathbf{D}'_2} & \mathcal{C}_{k+1}^2(T\mathcal{M}) \\ & & \parallel & & \parallel & & \downarrow ad & & \downarrow ad \\ 1 & \longrightarrow & \text{Aut}(\mathcal{M}) & \xrightarrow{j_{k+1}} & \Gamma_{k+1}(\mathcal{M}) & \xrightarrow{\mathbf{D}_1} & \text{Der}_{\Sigma,k}^1 & \xrightarrow{\mathbf{D}_2} & \text{Der}_{\Sigma,k}^2 \end{array}$$

where ad is the isomorphism of graded Lie algebras given in the definition (10), i.e. $\forall \tau, \chi \in \mathcal{C}_k^\bullet(T\mathcal{M})$:

$$ad([\tau, \chi]'') = [ad(\tau), ad(\chi)].$$

■

These sequences can be restricted to $\widehat{\mathcal{R}}_{k+1}$ if $\widehat{\mathcal{R}}_{k+1}$ satisfies the same properties as those given in the sequel of definition (3) concerning D' , and moreover if it is 2-acyclic.

5.2. The gravitational and electromagnetic fields

On the one hand, one has the following commutative diagram:

$$\begin{array}{ccc}
\underline{J_1} & \xrightarrow{\mathbf{D}'_1} & \underline{T^*\mathcal{M}} \otimes_{\theta} \underline{J_1} \\
\uparrow \phi_0 & & \uparrow \phi_1 \\
\widehat{\mathcal{R}}_2 & \xrightarrow{\widehat{\mathbf{D}}'_1} & \widehat{\mathcal{C}}_2^1 \\
|| id & & \downarrow ad \\
\widehat{\mathcal{R}}_2 & \xrightarrow{\widehat{\mathbf{D}}_1} & Der_{\Sigma,a}^1(\widehat{B}_1)
\end{array}$$

where $\widehat{B}_1 = \Lambda T^*\mathcal{M} \otimes_{\theta} \underline{J}_1$, and on the other hand, one obtains the following relation deduced from the relations (36) and (37): $\forall \alpha_1 \in \underline{J}_1 \equiv \widehat{B}_1^0$,

$$\mathbf{D}'_1(\alpha_1) = D'(\alpha_1) - \widehat{\mathbf{D}}_1(\widehat{f}_2)(\alpha_1).$$

Also let us define $\widehat{\mathfrak{D}}_1 \equiv \widehat{\mathbf{D}}_1(\widehat{f}_2) \in Der_{\Sigma,a}^1(\widehat{B}_1)$ and \mathfrak{D} such that:

$$\mathfrak{D} = D' - \widehat{\mathfrak{D}}_1 \in Der_{\Sigma,a}^1(\widehat{B}_1).$$

It follows one can rewrite:

$$\mathfrak{D} \circ \mathfrak{D} = -\widehat{\mathbf{D}}_2(\widehat{\mathfrak{D}}_1),$$

where $\widehat{\mathbf{D}}_2$ is the 2nd non-linear Spencer operator of the sophisticated Spencer complex of $\widehat{\mathcal{R}}_2$:

$$1 \longrightarrow Aut(\mathcal{M}) \longrightarrow \widehat{\mathcal{R}}_2 \xrightarrow{\widehat{\mathbf{D}}_1} Der_{\Sigma,a}^1(\widehat{B}_1) \xrightarrow{\widehat{\mathbf{D}}_2} Der_{\Sigma,a}^2(\widehat{B}_1).$$

Then, from the definition of $Der_{\Sigma,k}^\bullet$, there exists $\widehat{\tau}^{(2)} \in \widehat{\mathcal{C}}_2^1$ such that $\widehat{\mathfrak{D}}_1 = ad(\widehat{\tau}^{(2)})$ and therefore one can write also:

$$\mathfrak{D} \circ \mathfrak{D} = -ad(\widehat{\mathbf{D}}_2'(\widehat{\tau}^{(2)})).$$

Hence $\forall \alpha_1 \in \underline{J}_1$ one deduces the relation:

$$\mathfrak{D} \circ \mathbf{D}'_1(\alpha_1) = -ad(\widehat{\mathbf{D}}_2'(\widehat{\tau}^{(2)}))(\alpha_1),$$

and also $\forall \widehat{\tau}^{(2)} \in \widehat{\mathcal{C}}_2^1$:

$$\begin{aligned}
\mathfrak{D} \circ \phi_1(\widehat{\tau}^{(2)}) &= D' \circ \phi_1(\widehat{\tau}^{(2)}) - \widehat{\mathfrak{D}}_1 \circ \phi_1(\widehat{\tau}^{(2)}) \\
&= D' \circ \phi_1(\widehat{\tau}^{(2)}) - ad(\widehat{\tau}^{(2)}) \circ \phi_1(\widehat{\tau}^{(2)}) \\
&= D' \circ \frac{1}{n} Tr^1(\widehat{\tau}^{(2)}) - ad(\widehat{\tau}^{(2)}) \circ \phi_1(\widehat{\tau}^{(2)}) \\
&= \frac{1}{n} Tr^1 \circ D'(\widehat{\tau}^{(2)}) - ad(\widehat{\tau}^{(2)}) \circ \phi_1(\widehat{\tau}^{(2)}) \\
&\stackrel{def. \phi_2}{=} \phi_2 \circ D'(\widehat{\tau}^{(2)}) - ad(\widehat{\tau}^{(2)}) \circ \phi_1(\widehat{\tau}^{(2)}),
\end{aligned}$$

which can be rewritten:

$$\mathfrak{D} \circ \phi_1(\hat{\tau}^{(2)}) = \phi_2(\widehat{\mathbf{D}}'_2(\hat{\tau}^{(2)})) + \frac{1}{2}\phi_2([\hat{\tau}^{(2)}, \hat{\tau}^{(2)}]) - ad(\hat{\tau}^{(2)}) \circ \phi_1(\hat{\tau}^{(2)}),$$

where ϕ_2 is the θ -linear morphism:

$$\phi_2 : \widehat{\mathcal{C}}_2^2 \longrightarrow Der_{\Sigma,1}^2 \equiv Der_{\Sigma,a}^2(\widehat{B}_1),$$

satisfying the relations: $\forall \hat{\sigma}^{(2)} \in \widehat{\mathcal{C}}_2^2$,

$$\begin{aligned} \phi_2(\hat{\sigma}^{(2)}) &\equiv \frac{1}{n} Tr^1(\hat{\sigma}^{(2)}) \\ \text{and } \phi_2 \circ D' &= D' \circ \phi_1. \end{aligned}$$

Then, defining $\underline{\mathbf{D}}'_2 \stackrel{\text{def.}}{=} \mathfrak{D} / \underline{T^*\mathcal{M} \otimes_\theta J_1}$, we deduce the theorem:

Theorem *The following diagram of differential sequences is commutative, i.e.* $\underline{\mathbf{D}}'_2 \circ \phi_1 = \phi_2 \circ \widehat{\mathbf{D}}'_2$:

$$\begin{array}{ccccccc} & 1 & & 1 & & 0 & 0 \\ & \downarrow & & \downarrow & & \downarrow & \downarrow \\ 1 & \longrightarrow & \Gamma_G & \xrightarrow{j_2} & \mathcal{R}_2 & \xrightarrow{\mathbf{D}'_1} & \mathcal{C}_2^1 & \xrightarrow{\mathbf{D}'_2} & \mathcal{C}_2^2 \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \Gamma_{\widehat{G}} & \xrightarrow{j_2} & \widehat{\mathcal{R}}_2 & \xrightarrow{\widehat{\mathbf{D}}'_1} & \widehat{\mathcal{C}}_2^1 & \xrightarrow{\widehat{\mathbf{D}}'_2} & \widehat{\mathcal{C}}_2^2 \\ & \downarrow & & \downarrow \phi_0 & & \downarrow \phi_1 & & \downarrow \phi_2 \\ 1 & \longrightarrow & \theta & \xrightarrow{j_1} & \underline{J_1} & \xrightarrow{\underline{\mathbf{D}}'_1} & \underline{T^*\mathcal{M} \otimes_\theta J_1} & \xrightarrow{\underline{\mathbf{D}}'_2} & \underline{\Lambda^2 T^*\mathcal{M} \otimes_\theta J_1} \\ & & & & \downarrow & & \downarrow & & \downarrow \\ & & & & 1 & & 0 & & 0 \end{array} \quad (39)$$

if the condition below is satisfied: $\forall \hat{\tau} \in \widehat{\mathcal{C}}_2^1$,

$$\phi_2([\hat{\tau}, \hat{\tau}]) = 2 ad(\hat{\tau}) \circ \phi_1(\hat{\tau}). \quad (40)$$

■

Finally, one notices from a diagram chasing in the diagram (39) that the sequence is exact at $\underline{T^*\mathcal{M} \otimes_\theta J_1}$ if and only if one restricts to $Im \underline{\mathbf{D}}'_1$, i.e. one has only the short split exact sequence:

$$1 \longrightarrow \theta \xrightarrow{j_1} \underline{J_1} \xrightarrow{\underline{\mathbf{D}}'_1} \underline{\mathbf{D}}'_1(\underline{J_1}) \xrightarrow{\underline{\mathbf{D}}'_2} 0. \quad (41)$$

This might be (?) an illustration of the fundamental theorem of Spencer on the deformations of Lie structures (Spencer 1962, 1965, Goldschmidt 1976a, 1976b, 1978a, 1978b, 1981) since \underline{J}_1 is endowed with a Lie group structure.

Let us denote $\hat{\mathcal{C}}$ and $\pi^{(2)}$ by:

$$\hat{\mathcal{C}} \equiv (\hat{\mathcal{A}}, \hat{\mathcal{B}}) \in \hat{B}_1^1 = \underline{T^*\mathcal{M}} \otimes_{\theta} \underline{J}_1$$

$$\pi^{(2)} \equiv (\Omega^{(2)}, \mathcal{K}^{(2)}) \in \hat{B}_1^1{}^* \otimes_{\theta} (\hat{\mathcal{C}}_2^1 / \mathcal{C}_2^1)$$

$$\text{and } <\hat{\mathcal{C}}|\pi^{(2)}> \stackrel{\text{def.}}{=} <\hat{\mathcal{A}}|\Omega^{(2)}> + <\hat{\mathcal{B}}|\mathcal{K}^{(2)}> = \hat{\tau} - \tau^{(2)}$$

with, as a consequence of the relations (36) and (37):

$$\phi_1(\pi^{(2)}) = 1d_{\hat{B}_1^1} \in \hat{B}_1^1{}^* \otimes_{\theta} \hat{B}_1^1 \quad \text{and} \quad \phi_1(\tau^{(2)}) = 0.$$

Then, from (40), one obtains the following equivalent relations:

$$\phi_2([\![\tau^{(2)}, \tau^{(2)}]\!]) = 0 \tag{42}$$

$$\phi_2([\![\tau^{(2)}, \pi^{(2)}]\!]) = ad(\tau^{(2)}) \tag{43}$$

$$\phi_2([\![\pi^{(2)}, \pi^{(2)}]\!]) = 2ad(\pi^{(2)}), \tag{44}$$

which are the defining constraints on the $\tau^{(2)}$ and the susceptibilities $\pi^{(2)}$.

In an orthonormal system of local coordinates, one can write $\mathcal{D}'_2(\hat{\mathcal{C}}) \equiv (\hat{\mathcal{G}}, \hat{\mathcal{H}}) = \phi_2(\hat{\sigma}^{(2)}) \in \Lambda^2 T^*\mathcal{M} \otimes_{\theta} \underline{J}_1 = \hat{B}_1^2$ in the form ($h, i, j, k, l, r, s = 1, \dots, n$):

$$\begin{aligned} \hat{\mathcal{G}}_{[ij]} &= \frac{1}{n} (\hat{\sigma}_{k,[ij]}^k + \hat{\sigma}_{,[ij]}^k \gamma_{hk}^h) \\ &\equiv \hat{B}_{[i}^k (\partial_{|k} \hat{\mathcal{A}}_{j]}) - \hat{\mathcal{F}}_{[ij]} - \hat{\tau}_{[j,i]}^k \hat{\mathcal{A}}_k \\ \hat{\mathcal{H}}_{j,[ki]} &= \frac{1}{n} (\hat{\sigma}_{hj,[ki]}^h + \hat{\sigma}_{j,[ki]}^l \gamma_{hl}^h + \hat{\sigma}_{,[ki]}^k (\partial_j \gamma_{hk}^h)) \\ &\equiv \hat{B}_{[k}^h (\partial_{|h} \hat{\mathcal{B}}_{j,|i]}) - \hat{\mathcal{E}}_{j[k,i]} - \hat{\tau}_{[i,k]}^r \hat{\mathcal{B}}_{j,r} - \hat{\tau}_{j,[k}^r \hat{\mathcal{B}}_{|r,i]} - \hat{\tau}_{j[i,k]}^r \hat{\mathcal{A}}_r \end{aligned}$$

where

1. $\hat{\mathcal{F}} \in \Lambda^2 T^*\mathcal{M}$ is the skew-symmetric part of $\hat{\mathcal{B}}$ one calls the electromagnetic field tensor (or the Faraday tensor),
2. $(\hat{\mathcal{E}}, \hat{\mathcal{B}}, \hat{\mathcal{A}}) \in \underline{T^*\mathcal{M}} \otimes_{\theta} \mathcal{R}_2$, i.e. $\forall X, Y, Z \in T\mathcal{M}, \forall c_0 \in \mathbb{R}$,

$$<\hat{\mathcal{E}}(X, Y)|Z> = \frac{1}{2} <\hat{\mathcal{B}}(\nabla_X Y + \nabla_Y X)|Z> - c_0 \omega(X, Y) <\hat{\mathcal{A}}|Z>,$$

3. $\hat{\tau}^{(2)}$ satisfies the relation (38).

The terminology one uses for $\widehat{\mathcal{F}}$, comes obviously from the previous relations analogous to those obtained in the Maxwell theory. Indeed, since $(\widehat{\mathcal{G}}, \widehat{\mathcal{H}}) = 0$ because of the exactness condition in the sequence (41), one deduces:

$$\widehat{\mathcal{F}}_{[i,j]} = \widehat{B}_{[i}^k (\partial_{|k|} \widehat{\mathcal{A}}_{j]}) + \widehat{\tau}_{[i,j]}^k \widehat{\mathcal{A}}_k,$$

and from the expression for $\widehat{\mathcal{H}}$:

$$\widehat{B}_{[k}^h (\partial_{|h} \widehat{\mathcal{F}}_{|j,i]}) + \widehat{\tau}_{[k,j]}^r \widehat{\mathcal{F}}_{i,r] = 0.$$

In the case of the weak fields limit, the latter become:

$$\begin{cases} \widehat{\mathcal{F}} = \mathbf{d}\widehat{\mathcal{A}} & \text{(Faraday tensor)} \\ \mathbf{d}\widehat{\mathcal{F}} = 0 & \text{(Bianchi identity).} \end{cases}$$

On the contrary, the symmetric part $\widehat{\mathcal{P}} \in S_2 T^* \mathcal{M}$ of $\widehat{\mathcal{B}}$, called “the gravitational field tensor”, satisfies a first order PDE depending on $(\widehat{\mathcal{A}}, \widehat{\mathcal{B}})$, and thus $\widehat{\mathcal{P}}$ varies even if there is only an electromagnetic field! One has to notice that the symmetric part $(\partial_i \widehat{\mathcal{A}}_j)$ is never taken into account in physics, in contradistinction to $\widehat{\mathcal{A}}$. As a result all its derivatives should be physical observables. Finally, one deduces from the exactness of the complex (41) that no current of magnetic charges can exist. In other words, the Bianchi identity must be satisfied. Hence, the lack of magnetic charges can be justified in the framework of the Spencer cohomology of conformal Lie structures but not in the de Rham cohomology framework.

6. The dual linear Spencer complex and the Janet complex

6.1. The dual linear Spencer complex

One shall refer in this chapter to the definitions given in references (Goldschmidt 1976a, 1976b, 1978a, 1978b, 1981, Gasqui *et al.* 1984, Pommaret 1995) relating to the dual linear Spencer complex. Mainly, in this chapter, one has to build up the dual operator ${}^*\mathbf{d}'_1$ of the infinitesimal operator \mathbf{d}'_1 of \mathbf{D}'_1 , and the morphisms L_{n-i} ($i = 0, 1$) in the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \theta_{\mathcal{M}} & \xrightarrow{j_1} & \underline{J}_1 & \xrightarrow{\mathbf{d}'_1} & \mathbf{d}'_1(\underline{J}_1) \subseteq \mathcal{C}_1^1 & \longrightarrow & 0 \\ & & & & \downarrow L_n & & \downarrow L_{n-1} & & \\ & & & & & & & & \\ & & \mathbb{A}_1^n & \xleftarrow{{}^*\mathbf{d}'_1} & \mathbb{A}_1^{n-1} & & & & 0 \end{array}$$

where the \mathbb{A}_1^{n-i} are the dualizing fiber bundles ($\mathcal{C}_1^0 \equiv \underline{J}_1$):

$$\mathbb{A}_1^{n-i} = \underline{\Lambda}^n T^* \mathcal{M} \otimes_{\theta} {}^*\mathcal{C}_1^i.$$

One uses a rather classical method to determine the adjoint operators on a connected and oriented compact without boundaries (see P. J. Olver for instance 1986). To this purpose, one gets before a conformally equivariant Lagrangian density \mathfrak{L} :

$$\mathfrak{L} : \underline{J}_1 \times_{\mathcal{M}} \mathcal{C}_1^1 \equiv \mathcal{C}_1^0 \times_{\mathcal{M}} \mathcal{C}_1^1 \longrightarrow \underline{\Lambda}^n T^* \mathcal{M},$$

and one defines the morphisms L_{n-i} and ${}^*\mathbf{d}'_1$ integrating by parts the infinitesimal variation of \mathfrak{L} , with \mathbf{d}'_1 defined by $(\alpha_1 \equiv (\alpha, \beta))$:

$$\left\{ \begin{array}{l} \widehat{\mathcal{A}} = \mathbf{d}\alpha - \beta \\ \widehat{\mathcal{B}} = \mathbf{d}\beta - \mu \end{array} \right\} \equiv \mathbf{d}'_1 \alpha_1$$

where $\forall X, Y \in \mathcal{C}^1(T\mathcal{M})$:

$$\mu(X, Y) = \frac{1}{2}\beta(\nabla_X Y + \nabla_Y X) - c_0 \alpha \omega(X, Y).$$

Let us define in \mathbf{A}_1^{n-1} , the images of $(\widehat{\mathcal{A}}, \widehat{\mathcal{B}})$ by the morphism L_{n-1} with the relations:

$$\left\{ \begin{array}{l} \widehat{\mathcal{J}} = (\partial \mathfrak{L} / \partial \widehat{\mathcal{A}}) \quad (\text{electric current}) \\ \widehat{\mathcal{N}} = (\partial \mathfrak{L} / \partial \widehat{\mathcal{B}}), \end{array} \right.$$

and in \mathbf{A}_1^n , the images of α_1 by the morphism L_n with the relations:

$$\left\{ \begin{array}{l} \widehat{\mathcal{S}} = (\partial \mathfrak{L} / \partial \alpha) \quad (\text{density of entropy}) \\ \widehat{\mathcal{Q}} = (\partial \mathfrak{L} / \partial \beta). \end{array} \right.$$

One obtains from the infinitesimal variation $\delta \mathfrak{L}$, the following defining relations of ${}^*\mathbf{d}'_1$:

$$\left\{ \begin{array}{l} \widehat{\mathcal{S}} = \text{div} \widehat{\mathcal{J}} - c_0 < \omega | \widehat{\mathcal{N}} > \\ \widehat{\mathcal{Q}} = \widehat{\mathcal{J}} + \text{div}_2 \widehat{\mathcal{N}} + < \zeta | \widehat{\mathcal{N}} >, \end{array} \right.$$

where div_2 and ζ are morphisms such as:

$$\begin{aligned} \text{div}_2 : \otimes^2 T\mathcal{M} &\longrightarrow T\mathcal{M} \\ u \otimes v &\longrightarrow v \text{div}(u) - u \text{div}(v) + [u, v] \end{aligned}$$

and

$$\begin{aligned} \zeta : \otimes^2 \mathcal{C}^1(T\mathcal{M}) &\longrightarrow \mathcal{C}^0(T\mathcal{M}) \\ u \otimes v &\longrightarrow \zeta(u \otimes v) = \gamma(u)v = \gamma(v)u \equiv < \xi | u \otimes v >. \end{aligned}$$

From these results, a further step would be to know what the relations between ${}^*\mathbf{d}'_1$ and the thermodynamical laws for the irreversible processes are, since the Lagrangian density \mathfrak{L} depends on temperature and variables of internal states. Hence, \mathfrak{L} might be identified with a Gibbs free enthalpy function, but moreover, setting certain conditions, the conformal equivariance laid down to \mathfrak{L} , might also allow us to consider \mathfrak{L} as a wavefunction, as we shall see in what follows.

6.2. The Janet complex of the Lagrangian density \mathfrak{L}

Again, one refers for the definitions to Pommaret's previous papers (Pommaret 1989, 1994) about the Janet complex and also to Gasqui-Goldschmidt's ones (Gasqui *et al.* 1988), but with an "un-named complex". The conformal equivariance of \mathfrak{L} will merely allow us to obtain explicitly the first Janet operator $\mathbf{D}_1 : \mathcal{F}_0 \longrightarrow \mathcal{F}_1$ where \mathcal{F}_0 stands for the line bundle over $\mathcal{C}_1^0 \times_{\mathcal{M}} \mathcal{C}_1^1$:

$$\begin{array}{ccc} & \mathcal{F}_0 & \\ & \downarrow \mathfrak{L} & \\ \mathcal{C}_1^0 \times_{\mathcal{M}} \mathcal{C}_1^1 & & \end{array}$$

From now on, one presents a dependent coordinates formulation choosing an orthonormal system of local coordinates. To make \mathbf{D}_1 , one must find the transformation rule of \mathfrak{L} by the action of the conformal Lie pseudogroup. Namely, if $\hat{f}_2 \in \widehat{\mathcal{R}}_2$ and $\hat{f}_2^{-1} = \hat{g}_2$, one has the transformation rules:

$$\left\{ \begin{array}{lcl} y & = & \hat{f}(x) \simeq x + \hat{\xi}(x) \\ \alpha' \circ \hat{f} & = & \alpha - \frac{1}{n} \ln |\det J(\hat{f}_1)| \\ \beta'_i \circ \hat{f} & = & \hat{g}_i^j \circ \hat{f} [\beta_j - \frac{1}{n} \hat{g}_l^k \circ \hat{f} \hat{f}_{kj}] \\ \hat{\mathcal{A}}'_i \circ \hat{f} & = & \hat{g}_i^j \circ \hat{f} \hat{\mathcal{A}}_j \\ \hat{\mathcal{B}}'_{i,j} \circ \hat{f} & = & \hat{g}_i^r \circ \hat{f} \hat{g}_j^s \circ \hat{f} \hat{\mathcal{B}}_{r,s} \end{array} \right.$$

and one must have:

$$\mathfrak{L}(\hat{f}, \alpha' \circ \hat{f}, \beta' \circ \hat{f}, \hat{\mathcal{A}}' \circ \hat{f}, \hat{\mathcal{B}}' \circ \hat{f}) \det J(\hat{f}_1) = \mathfrak{L}(x, \alpha, \beta, \hat{\mathcal{A}}, \hat{\mathcal{B}}). \quad (45)$$

One obtains the infinitesimal condition and the definition of \mathbf{D}_1 :

$$\mathbf{D}_1(\mathfrak{L}) \equiv \{v^\mu(\mathfrak{L}) + (\text{div } \xi^\mu) \mathfrak{L} \equiv \mathfrak{K}\} = 0,$$

whatever is v^μ ($\mu = 1, \dots, \dim \mathfrak{g}_c$), generator of the Lie algebra \mathfrak{g}_c of the conformal Lie group:

$$v^\mu = \hat{\xi}^{\mu,j} \partial_j + \phi^\mu(\alpha) \partial_\alpha + \phi^\mu(\beta_j) \partial_{\beta_j} + \phi^\mu(\hat{\mathcal{A}}_j) \partial_{\hat{\mathcal{A}}_j} + \phi^\mu(\hat{\mathcal{B}}_{k,l}) \partial_{\hat{\mathcal{B}}_{k,l}},$$

where

$$\left\{ \begin{array}{lcl} \phi^\mu(\alpha) & = & -\frac{1}{n}\hat{\xi}_k^{\mu,k} \\ \phi^\mu(\beta_j) & = & -[\hat{\xi}_j^{\mu,k}\beta_k + \hat{\xi}_{kj}^{\mu,k}] \\ \phi^\mu(\hat{\mathcal{A}}_j) & = & -\hat{\xi}_j^{\mu,k}\hat{\mathcal{A}}_k \\ \phi^\mu(\hat{\mathcal{B}}_{k,l}) & = & -[\hat{\xi}_k^{\mu,h}\hat{\mathcal{B}}_{h,l} + \hat{\xi}_l^{\mu,h}\hat{\mathcal{B}}_{k,h}] \end{array} \right.$$

and $\hat{\xi}_2^\mu \in \hat{R}_2$. Moreover, $\mathfrak{K} \equiv (\mathfrak{K}^\mu)$ is transformed like \mathfrak{L} and therefore the zero section of \mathcal{F}_1 is defined with $\dim \mathfrak{g}_c = \dim \mathcal{F}_1$ constants c^μ such that $\mathfrak{K}^\mu = c^\mu \mathfrak{L}$ meaning that (45) (or the Lagrangian density) is defined up to a multiplicative constant. The second Janet operator \mathbf{D}_2 such as $\mathbf{D}_2 \circ \mathbf{D}_1 = 0$ and $\mathbf{D}_2 : \mathcal{F}_1 \longrightarrow \mathcal{F}_2$, is defined from the involution of the generators of \mathfrak{g}_c :

$$[v^\mu, v^\nu] = c^{\mu\nu}_\lambda v^\lambda,$$

where the $c^{\mu\nu}_\lambda$ are the constants of structure of the \mathfrak{g}_c (see P. J. Olver 1986 for a precise definition of the bracket used above). Thus, one obtains:

$$\mathbf{D}_2(\mathfrak{K}) \equiv \left\{ v^\mu(\mathfrak{K}^\nu) - v^\nu(\mathfrak{K}^\mu) - c^{\mu\nu}_\lambda \mathfrak{K}^\lambda + (\text{div } \hat{\xi}^\mu) \mathfrak{K}^\nu - (\text{div } \hat{\xi}^\nu) \mathfrak{K}^\mu \right\}.$$

Leading down $\mathfrak{K}^\mu = c^\mu \mathfrak{L}$ in the latter expression, one deduces the following constraints on the constants $c^{\mu\nu}_\lambda$ of structure:

$$c^{\mu\nu}_\lambda c^\lambda = 0,$$

with $\mathbf{D}_1(\mathfrak{L})^\mu = c^\mu \mathfrak{L}$.

This latter equation gives in fact only one arbitrary constant among the c^μ 's and perhaps might be ascribed to a constant such as the Planck one.

As a particular case, if the metric ω and $\hat{\mathcal{A}}$ are constants, α and β vanish and \mathfrak{L} function of x and $\hat{\mathcal{A}}$ only, such that there exists η satifying the relation

$$\widehat{\mathcal{J}}^k = \eta^k \mathfrak{L},$$

then $\forall \mu, h$ one has $\hat{\xi}_k^{\mu,k} = \hat{\xi}_{kh}^{\mu,k} = 0$ and

$$[\hat{\xi}^{\mu,k} \partial_k - (\eta^k \hat{\xi}_k^{\mu,h}) \hat{\mathcal{A}}_h] \mathfrak{L} - c^\mu \mathfrak{L} = 0.$$

Thus, one obtains an analogous PDE to the Dirac equation but being equivariant and not only covariant like the Dirac equation. Nevertheless \mathfrak{L} being a real function, it can be interpreted as a wave-function only if one has a definition of a measure of probability. We propose two ways of doing this in the conclusion, and we make suggestions for taking the weak and strong interactions into account in the model.

7. Conclusion

The function \mathfrak{L} being real, let us suggest a notion of measure of probability. Two of them can be proposed. The first one is related to an “à la Misra-Prigogine-Courbage” (MPC) (Misra *et al.* 1979, Misra 1987) approach. With more details, the wave-function \mathfrak{S} of the model would be the function obtained when applying the integral action of the evolution operator to an initial Lagrangian density \mathfrak{L} . First, we thus obtain a wave interpretation. Second, within the MPC theory, the reduction of the wave-packet (or the collapse) would axiomatically be defined as the achievement of a K-partition of functions \mathfrak{S} on a compact (on which the initial Cauchy data would be defined) of a submanifold of the Minkowski space-time to determine a single function \mathfrak{S} . This K-partition of functions \mathfrak{S} would then be a set of new initial Lagrangian densities \mathfrak{L} before any time evolution. The physical measurement thus achieved would not lead us to obtain pure states and with the assumption (to be confirmed) that the conformally equivariant Lagrangian densities \mathfrak{L} would be Kolmogorov flows, it would then be possible, according to the MPC theory, to build up a non-commutative algebra of observables on the analogy of quantum mechanics. Moreover, the K-partition would be achieved on a set of Lagrangian densities of type (m,s) deduced by projection on the basis of the eigenstates of the Poincaré group during the process of measurement (from a certain point of view, the apparatus of measurement, being themselves of the type (m,s)). Thus, as a result of this projection, we shall see a particle state and initial Lagrangian densities of type (m,s) not depending on the fields of interaction anymore. The particle states might be interpreted as a split up of variables between the variable of position x and the fields variables (during specific physical measurements). In other words, one could say that the energy of the fields of interaction would be transferred by radiation to the apparatus of measurement. From a more general point of view, the process of “fragmentation” would be defined by the change from a tensorial Hilbert space to a decomposable Hilbert space (\mathcal{H}_i ; $i = 1, 2$: Hilbert spaces):

$$\mathcal{H}_1 \otimes \mathcal{H}_2 \longrightarrow \mathcal{H}_1 \oplus \mathcal{H}_2$$

Process that might be interpreted as well as a separation of phase in thermodynamics for the function \mathfrak{L} is similar to a Gibbs function. Under these conditions, the analytical developments of the initial Lagrangian densities define tensors of susceptibility directly associated with the “classical” states of the condensed matter, therefore with the classical notion of macroscopic particles. Thus, the macroscopicity would be associated with the “degree” of separability of the initial densities \mathfrak{L} . From our point of view, a second possibility used to define a measure of probability would be to refer to the Penrose approach (Penrose *et al.* 1986) on the complexification of the wave-function in the frame of the theory of twistors, with moreover, the notion of measure of probability defined as it usually is in the first quantization. This procedure is completely different from the one consisting in simply making an algebraic extension from the field of reals to the field of complexes, of the wave-functions satisfying PDE on the field of reals. Such an extension would lead to complex solutions whose real and imaginary parts would each separately be solutions. Consequently, we assume that a complexification presenting a real interest would give access to the determination of complex solutions which, on the contrary, would be nei-

ther their real parts, nor their imaginary ones, solutions for the given system of PDE. To achieve that, Penrose suggested decomposing a problem defined on the real Minkowski space \mathcal{M} into a set of physical subsystems defined on submanifolds of \mathcal{M} and having at least one (complex) spin structure, like the celestial spheres S^\pm are, or at fixed time, the sphere S^3 . The spin structures are associated with vector bundles like the tangent spaces, or on a rather similar manner (up to the target manifold), the 1-jet bundle $J_1(\epsilon)$ for example. In the present situation, we shall refer to the $\mathcal{A}'_{\underline{1}}(J_1) \subseteq \mathcal{C}^1_{\underline{1}}$ bundle over J_1 . We therefore must present the tensors $\hat{\mathcal{A}}$ and $\hat{\mathcal{B}}$ with spinors. In the more simple case with only one spin structure (on S^2 for example) and taking up again the Penrose indexed notation, one need at least 3 spinors to decompose the tensor $\hat{\mathcal{A}}$:

$$\hat{\mathcal{A}}_a = \alpha_A \cdot \bar{\alpha}_{A'} + \beta_A \cdot \bar{\beta}_{A'} - \gamma_A \cdot \bar{\gamma}_{A'}.$$

For only under these conditions, $\hat{\mathcal{A}}$ is of any norm, but with a spinor, $\hat{\mathcal{A}}$ must be of the type light and with two spinors, the norm of $\hat{\mathcal{A}}$ must be non-negative or non-positive. As far as I know this simple decomposition has never been presented yet (apart from the one with 2 spinors or 3 maximum and with a + mark). But it has indeed a few advantages. First, it is associated (Bars *et al.* 1990) with the representations $\{3\}$ and $\{\bar{3}\}$ of $SU(2, 1)$ isomorphic to the one of $SU(3)$. It is only about representations; the group $SU(2, 1)$ not being a dynamical gauge group of the model, like for example the Yang-Mills type, but only a group of internal classification and of invariance of the decomposition. The non-compactness of the group cannot appear determining to us within this context. But its finished unitary irreducible representations constructed from the two fundamental representations $\{3\}$ and $\{\bar{3}\}$ only do. Moreover, a symmetry breaking from $SU(2, 1)$ to $SU(2) \times U(1)$ or $SU(1, 1) \times U(1)$ is associated to a symmetry breaking of type T (for instance during the process of fragmentation) and $\hat{\mathcal{A}}$ is then of a non-negative or non-positive norm. All of that suggests a possible link with a theory of weak and strong interactions. Before that, formally deriving the spinors decomposing $\hat{\mathcal{A}}$, we obtain the following decomposition for $\hat{\mathcal{B}}$ (Penrose *et al.* 1986, see chapter 4.4.7.) using the Leibniz law for the spinors:

$$\hat{\mathcal{B}}_{a,b} = \alpha_A \cdot \bar{\Gamma}_{B'A',B} + \bar{\alpha}_{A'} \cdot \Gamma_{BA,B'} + \beta_A \cdot \bar{\Theta}_{B'A',B} + \bar{\beta}_{A'} \cdot \Theta_{BA,B'} + \gamma_A \cdot \bar{\Omega}_{B'A',B} + \bar{\gamma}_{A'} \cdot \Omega_{BA,B'},$$

where Γ , Θ and Ω are any mixed spinors of valence (2,1). They can decompose themselves into irreducible spinors with respect to $SL(2, \mathbb{C})$:

$$\Gamma_{AB,B'} = \Gamma_{(AB),B'} - \frac{1}{2} \epsilon_{[AB]} \Gamma_C{}^C_{,B'}$$

(with similar expressions for Θ and Ω). Let us notice that at a given fixed time t and with (α, β) fixed (i.e. during the measure), $SL(2, \mathbb{C})$ being associated with $O_1^+(1, 3)$, the irreducibility is thus, according to $SU(2)$, locally with respect to $O(3)$. Moreover, $\Gamma_{(AB),B'}$ is defined by 6 complex components, linearly independant and $\Gamma_C{}^C_{,B'}$ by 2 complex components. The interest of such a decomposition appears as soon as the assumption is made that the Lagrangian density \mathcal{L} is holomorphic in the various spinors on the

submanifolds endowed with spin structures. Then \mathfrak{L} satisfies complex PDE deduced from the real PDE defined on \mathcal{M} by lifting on the spinors bundle. But, the physical meaning of the holomorphy is that the physical system can precisely be confined on the submanifolds of M endowed with a spin structure. If we make the physical interpretation that to the spinors decomposing $\hat{\mathcal{A}}$ and $\hat{\mathcal{B}}$ are associated some fields of interactions or particles, it means that the latter can only be considered as “free” if they are precisely confined on those submanifolds; the density \mathfrak{L} being then a complex solution on the spinors bundle of the various lifted PDE of the model, from which neither the real part, nor the imaginary part are any solutions. On that subject, one cannot help thinking of the very controversial - and to my knowledge still un-confirmed - Larue *et al.* (1977) and Schaad *et al.* (1981) experiences on “free” quarks confined on bidimensional ($\simeq S^2$) layers of superconducting (!) nobium covering microballs of tungsten (see also Goldman 1995, 1996). Especially because the decomposition presented above could suggest that the simple spinors α , β and γ would be associated with some fields of fermion of mono-colored gluons of spin 1/2 (and not to bi-colored bosons!) and that the symmetric mixed spinors of valence (2,1) would be 3 quarks states of spin 1/2 determined by the 3 contracted spinors $\Gamma_{C,B'}^C$, $\Theta_{C,B'}^C$ and $\Omega_{C,B'}^C$ on which $SU(2,1)$ acts. At last, to finish as well as to come back to an application of the physics of the superconducting states (that some of the theoretical physicists like Mendelstam (1982) or t’Hooft (1978) used to explain the quarks confinement) the question is under which conditions the 4-vector current:

$$\widehat{\mathcal{J}} = (\partial \mathfrak{L} / \partial \hat{\mathcal{A}})$$

would be anomalous as well as the Faraday tensor $\widehat{\mathcal{F}}$. A possibility we suggest would be to consider a third metric λ deduced from ν like ν is from ω . The latter metric λ would appear in matter such as in crystals or amorphous materials for instance. Indeed in this case, the second metric ν (and its corresponding potentials and fields) would be associated to the “crystal field” and to a new kind of substrat space-time and new specific susceptibilities. But in the contrary to ω the Weyl curvature associated to ν is no longer necessarily vanishing as we pointed out in a previous chapter. Thus, the corresponding Lie equations for conformal transformations won’t be involutive. As a result, the corresponding Spencer sequences and the relative one won’t be exact any more as well. In particularly magnetic charges might occur and so anyons (Wilczek 1990). In the framework of symplectic cohomologies of the Lagrangian density \mathfrak{L} , it can be shown that the anomalies classification is associated to a second cohomology space knowledge as well as a third one (Cariñena *et al.* 1988). The question remaining to work out would be to know if one can obtain integer cohomologies from these latter in order to exhibit a kind of “quantum structure”. To conclude on a more philosophical note, the wave-function model we present here could be interpreted within the framework of a non-Lavoisian chemistry like G. Bachelard (1940) developped. In actual fact, the quantons would not be of “substance”, but as Bachelard named them, they would rather be “grains de réaction” and thus, from a certain point of view, like quantons of reactional synthesis or of separation of phase associated with the concept of fragmentation. We also think that this model can’t describe global evolution such as a big-bang model but only a local

evolution. In some ways as G. Deleuze *et al.* (1980) say about their striated space (like a substrat space-time) and smooth space concepts (see chapters: “12.1227. Traité de nomadologie: la machine de guerre” and “11.1837. De la ritournelle”), “one can only know the path by exploring it”. Also, the form concept for matter in space-time could be related to a boundary being the geometrical set of places onto which the “grains de réaction” occur.

Acknowledgements I wish to thank Florence Pittolo and Terence Blake who very kindly helped me in the translation of the paper and for their helpful comments and encouragements.

References

- Asher, E. 1973 In *Magnetoelectric Interaction Phenomena in Crystals*. Proceedings, Seattle, 69.
- Bachelard, G. 1940 *La philosophie du non*. Paris: Presse Universitaire de France.
- Bacry, H. 1967 *Leçons sur la théorie des groupes et les symétries des particules élémentaires*. Paris: Gordon and Breach.
- Bars, I. & Zhong-jian, T. 1990 *The unitary irreducible representations of $SU(2,1)$* . *J. Math. Phys.* (7)**31**, 1576-1586.
- Bateman, H. 1910 *Proc. Lond. math. Soc.* **8**, 223.
- Cariñena, J. F. & Ibort, L. A. 1988 *J. Math. Phys.* **29**, 541.
- Cunningham, E. 1910 *Proc. Lond. math. Soc.* **8**, 77.
- Deleuze, G. & Guattari, F. 1980 *Mille Plateaux*. Paris: Les éditions de minuit.
- Dieudonné, J. 1971 *Éléments d'analyse, Tome IV*. Paris: Gauthier-Villars Éditeur.
- Gasqui, J. & Goldschmidt, H. 1988 *Complexes of Differential Operators and Symmetric Spaces*. In *Deformation Theory of Algebras and Structures and Applications*. 797-827 (eds. M. Hazewinkel & M. Gerstanhaber) Dordrecht-Boston-London: Kluwer Academic Publishers.
- Gasqui, J. & Goldschmidt, H. 1984 *Deformations infinitesimales des structures conformes plates*. *Progress in Mathematics*, Vol. **52**, Boston-Basel-Stuttgart: Birkhaeuser.
- Goldman, V. J. 1996 *Surf. Sci.* **361**, 1.
- Goldman, V. J. & Su, B. 1995 *Science* **267**, 1010.

- Goldschmidt, H. & Spencer, D. 1976a *On the non-linear cohomology of Lie equations. Part I*, *Acta Math.* **136**, 103-170.
- Goldschmidt, H. & Spencer, D. 1976b *On the non-linear cohomology of Lie equations. Part II*, *Acta math.* **136**, 171-239.
- Goldschmidt, H. & Spencer, D. 1978a *On the non-linear cohomology of Lie equations. Part III*, *J. Differ. Geom.* **13**, 409-453.
- Goldschmidt, H. & Spencer, D. 1978b *On the non-linear cohomology of Lie equations. Part IV*, *J. Differ. Geom.* **13**, 455-526.
- Goldschmidt, H. 1981 *On the nonlinear cohomology of Lie equations. Part V*, *J. Differ. Geom.* **16**, 595-674.
- 't Hooft, G. 1978 *Nucl. Phys.* **B138**, 1.
- Itzykson, C. & Zuber, J.-B. 1980 *Quantum Field Theory*. McGraw-Hill.
- Janner, A. & Asher, E. 1969 *Phys. Lett.* (4)A**30**, 223.
- Janner, A. & Asher, E. 1978 *Physica* **48**, 425.
- Katanaev, M. O. & Volovich, I. V. 1992 *Ann. Phys.* **216**, 1-28.
- Kleinert, H. 1989 *Gauge fields in condensed matter*. Vol. **II**. World Scientific.
- Kumpera, A. & Spencer, D. C. 1972 *Lie Equations*, Vol. *I: General Theory. Annals of Mathematics Studies*, Vol. **73**.
- Larue, G. S. & Fairbank, W. M. & Hebard, A. F. 1977 *Evidence for the Existence of Fractional Charge on Matter**. *Phys. Rev. Lett.* (18)**38**, 1011-1014.
- Lévy-Leblond, J.-M. 1990 *ENIGMA OF THE SPINx*. In *Symposium on foundations of modern Physics*. 226-241. World Scientific.
- Lurçat, F. 1964 *Quantum Field Theory And The Dynamical Role Of Spin. Physics* (2)**1**, 95-106.
- Mandelstam, S. 1982 *Monopoles in Quantum Field Theory*. In *Proceedings of the Monopole Meeting*. Trieste, Italy, December 1981. World Scientific.
- Misra, B. & Prigogine, I. & Courbage, M. 1979 *From deterministic dynamics to probabilistic descriptions*. *Physica* **A98**, 1-26.
- Misra, B. 1987 *Fields as Kolmogorov Flows*. *J. Stat. Phys.* (5/6)**48**, 1295-1320.
- Olver, P. J. 1986 *Applications of Lie Groups to Differential Equations*. *Graduate texts in mathematics*, Vol. **107**. Springer.
- Penrose, R. & Rindler, W. 1986 *Spinors and Space-Time. Cambridge Monographs On Mathematical Physics*, Vol. **I & II**. Cambridge University Press.
- Pommaret, J.-F. 1994 *Partial Differential Equations and Group Theory, New Perspectives for Applications. Mathematics and Its Applications*, vol. **293**. Dordrecht-Boston-London: Kluwer Academic Publishers.

- Pommaret, J.-F. 1995 *Suites Differentielles et Calcul Variationnel*. *C. R. Acad. Sci. Paris* **320**, Série I, 207-212.
- Pommaret, J.-F. 1989 *Gauge Theory And General Relativity. Reports in Mathematical Physics* (3)**27**, 313-344.
- Pommaret, J.-F. 1988 *Thermodynamique et théorie des groupes*. *C. R. Acad. Sci. Paris* **307**, Série I, 839-842.
- Rubin, J. L. 1994 *Relativistic Crystalline Symmetry Breaking And Anyonic States in Magnetoelectric Superconductors*. *Ferroelectrics* **161**, 335.
- Rubin, J. L. 1993 *Broken Relativistic Symmetry Groups, Toroidal Moments and Superconductivity In Magnetoelectric Crystals*. *Il Nuovo Cimento D-Cond. Matt. At.* **15**, 59.
- Schaad, L. J. & Hess Jr., B. A. & Wikswo Jr., J. P. & Fairbank, W. M. 1981 *Quark chemistry*. *Phys. Rev. (4)A***23**, 1600-1607.
- Shih, W. 1986 *Une méthode élémentaire pour l'étude des équations aux dérivées partielles*. In *Diagrammes*, **16**. Paris.
- Shih, W. 1987 *C. R. Acad. Sci. Paris* **304**, Série I, 103-106, 187-190, 535-538.
- Shih, W. 1991 *Gazette. Société Mathématique de France*, 1.
- Spencer, D. C. 1962 *Deformation of Structures on Manifolds defined by transitive, continuous pseudogroups. Parts I & II*, *Annals of Math.* **76**, 306-445.
- Spencer, D. C. 1965 *Deformation of Structures on Manifolds defined by transitive, continuous pseudogroups. Part III*, *Annals of Math.* **81**, 289-450.
- Wilczek F. 1990 *Fractional Statistics and Anyons Superconductivity*. World Scientific Publishing Co. Pte. Ltd.

High Temperature Superconductivity

M. Brian Maple

Department of Physics and Institute for Pure and Applied Physical Sciences
University of California, San Diego, La Jolla, California 92093, USA

Abstract

The current status of basic research on the high temperature cuprate superconductors and prospects for technological applications of these materials is discussed. Recent developments concerning other novel superconductors are also briefly described.

Keywords:

Superconductivity – high- T_c , superconductors – ceramic, correlations – electronic, order parameter

Contact author:

M. Brian Maple, Department of Physics–0319, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093-0319, USA; FAX: (619) 534-1241; email: mbmaple@ucsd.edu

Introduction

The discovery of superconductivity at ~30 K in the La-Ba-Cu-O system by Bednorz and Müller in 1986 [1] ignited an explosion of interest in high temperature superconductivity. These initial developments rapidly evolved into an intense worldwide research effort that still persists after more than a decade, fueled by the fact that high temperature superconductivity constitutes an extremely important and challenging intellectual problem and has enormous potential for technological applications. During the past decade of research on this subject, significant progress has been made on both the fundamental science and technological applications fronts. For example, the symmetry of the superconducting order parameter and the identity the superconducting electron pairing mechanism appear to be on the threshold of being established, and prototype superconducting wires that have current carrying capacities in high magnetic fields that satisfy the requirements for applications are being developed. Prospects seem to be good for attaining a fundamental understanding of high temperature superconductors and realizing technological applications of these materials on a broad scale during the next decade.

The purpose of this paper is to provide a brief overview of the current status of the field of high temperature superconductivity. The emphasis is on experiment and recent developments on the high superconducting critical temperature (T_c) cuprates. Topics discussed include: (1) materials, (2) structure and charge carrier doping, (3) normal state properties, (4) symmetry of the superconducting order parameter, and (5) prospects for technological applications. At the end of the article, we describe recent progress involving other novel superconducting materials.

The immense scope of this subject dictated a very selective choice of the examples cited to illustrate the progress made in this field over the past decade. For comprehensive accounts of specific topics in high temperature superconductivity, the reader is referred to various review articles, such as those that appear in the series of volumes edited by

Ginsberg [2]. Because of space limitations, it was also not possible to discuss the fascinating subject of vortex phases and dynamics which has flourished since the discovery of the cuprate superconductors [3].

High T_c superconducting cuprates

The dramatic increases in T_c that have been observed since 1986 are illustrated in Fig. 1 where the maximum value of T_c is plotted vs. date. Prior to 1986, the A15 compound Nb_3Ge with $T_c \approx 23$ K held the record for the highest value of T_c [4]. The maximum value of T_c has increased steadily since 1986 to its present value of ~ 133 K for a compound in the Hg-Ba-Ca-Cu-O system [5,6]. When this compound ($\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_8$) is subjected to a high pressure, the T_c onset increases to ~ 164 K (more than half way to room temperature!) at pressures ~ 30 GPa [7,8]. While $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_8$ cannot be used in applications of superconductivity at such high pressures, this striking result suggests that values of T_c in the neighborhood of 160 K, or even higher, are attainable in cuprates at atmospheric pressure.

Values of T_c in excess of the boiling temperature of liquid nitrogen (77 K) immediately implicated high T_c cuprates as promising candidates for technological applications of superconductivity. Whereas liquid helium is currently employed to cool conventional superconducting materials such as Nb, NbTi, and Nb_3Sn into the superconducting state (Nb is employed in SQUIDs and NbTi and Nb_3Sn are used to make superconducting wires), the cuprate materials have the advantage that they can be cooled into the superconducting state using liquid nitrogen. Cuprates such as the $\text{LnBa}_2\text{Cu}_3\text{O}_{7-\delta}$ (Ln = lanthanide) compounds (T_c in the range 92-95 K) have enormous critical fields $\sim 10^2$ tesla [9,10] that are more than adequate for technological applications. Epitaxially grown thin films of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ on single crystal SrTiO_3 substrates have critical current densities $J_c \approx 10^6 \text{ A/cm}^2$ in zero field which decrease relatively slowly with magnetic field, making them suitable for technological applications [11]. Unfortunately, polycrystalline bulk materials have J_c 's that are disappointingly low, $\sim 10^3 - 10^4 \text{ A/cm}$, and are strongly

depressed by a magnetic field [12]. The situation can be improved substantially by subjecting $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ to a melt textured growth process which yields values of J_c of $\sim 10^5$ A/cm at 77 K that are not too strongly depressed by an applied magnetic field [13]. Fortunately, techniques have recently been devised which yield values of J_c in high fields for in-plane grain oriented thin films of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ on flexible substrates at 64 K (pumped liquid nitrogen temperatures) that exceed those of NbTi and Nb_3Sn at liquid helium temperatures [14]. These promising developments are briefly described near the end of this article.

The materials

Approximately 100 different cuprate materials, many of which are superconducting, have been discovered since 1986. Several of the more important high T_c cuprate superconductors are listed in Table 1, along with the maximum values of T_c observed in each class of materials. Included in the table are examples of abbreviated designations (nicknames) for specific cuprate materials which we will use throughout this article (e.g., $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ = YBCO, YBCO-123, Y-123). High quality polycrystalline, single crystal and thin film specimens of these materials have been prepared and investigated extensively to determine their fundamental normal and superconducting state properties. Presently, two of the leading candidates for technological applications of superconductivity are the $\text{LnBa}_2\text{Cu}_3\text{O}_{7-\delta}$ and $\text{Bi}_2\text{Sr}_2\text{Ca}_2\text{Cu}_3\text{O}_{10}$ materials.

Structure and charge carrier doping

The high T_c cuprate superconductors have layered perovskite-like crystal structures which consist of conducting CuO_2 planes separated by layers comprised of other elements A and oxygen, A_mO_n , and, in some cases, layers of Ln ions [15,16]. The mobile charge carriers, which can be electrons but are usually holes, are believed to reside primarily within the CuO_2 planes. The A_mO_n layers apparently function as charge reservoirs and control the doping of the CuO_2 planes with charge carriers. In several of the compounds

containing Ln layers, the Ln ions with partially-filled 4f electron shells and magnetic moments have been found to order antiferromagnetically at low temperature [9,12].

Many of the cuprates can be doped with charge carriers and rendered superconducting by substitution of appropriate elements into an antiferromagnetic insulating parent compound. For example, substitution of divalent Sr for trivalent La in the antiferromagnetic insulator La_2CuO_4 dopes the CuO_2 planes with mobile holes and produces superconductivity in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ with a maximum T_c of ~ 40 K at $x \approx 0.17$ [17]. Similarly, substitution of tetravalent Ce for trivalent Nd in the antiferromagnetic insulating compound Nd_2CuO_4 apparently dopes the CuO_2 planes with electrons, resulting in superconductivity in $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ with a maximum T_c of ~ 25 K at $x \approx 0.15$ for $y \approx 0.02$ [18,19]. The temperature T vs x phase diagrams for the $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ systems are shown in Fig. 2 [19], and the corresponding crystal structures of the La_2CuO_4 and Nd_2CuO_4 parent compounds are displayed in Fig. 3. The $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ systems have one CuO_2 plane per unit cell and are referred to as single CuO_2 layer compounds. Other superconducting cuprate systems have more than one CuO_2 plane per unit cell: $\text{LnBa}_2\text{Cu}_3\text{O}_{7-\delta}$ has two CuO_2 planes per unit cell (double CuO_2 layer compound), while $\text{Bi}_2\text{Sr}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_x$ has n CuO_2 layers per unit cell (n CuO_2 layer compound) and can be synthesized by conventional methods for $n = 1, 2, 3$.

Two features in Fig. 2 would appear to be relevant to cuprate superconductivity: (1) the apparent electron-hole symmetry may provide a constraint on viable theories of high T_c superconductivity in cuprates, and (2) the proximity of antiferromagnetism suggests that superconducting electron pairing in the cuprates may be mediated by antiferromagnetic spin fluctuations. An antiferromagnetic pairing mechanism is consistent with the occurrence of d-wave pairing with $d_{x^2-y^2}$ symmetry that is suggested by experiments on several hole-doped cuprates (discussed later in this article). A number of theoretical models (e.g., [20-22]) based on AFM spin fluctuations predict d-wave superconductivity with $d_{x^2-y^2}$

symmetry for the cuprates. Surprisingly, as discussed below, experiments on the electron-doped superconductor $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ suggest s-wave pairing similar to that of conventional superconductors where the pairing is mediated by phonons.

It is interesting that superconductivity with values of T_c in the neighborhood of 30 K have been found in two noncuprate materials: the cubic perovskite $\text{Ba}_{1-x}\text{K}_x\text{BiO}_3$ ($T_c \approx 30$ K) [23,24] and the fcc "buckeyball" compound Rb_3C_{60} ($T_c \approx 29$ K) [25,26]. Other features are consistent with non-phonon-mediated pairing in the hole-doped cuprates. The curve of T_c vs carrier concentration can be approximated by an inverted parabola with the maximum value of T_c occurring at an optimal dopant concentration x_0 [27]. (Note that the terminology "under-doped" refers to values of x smaller than the "optimally-doped" value x_0 , whereas "over-doped" refers values of x larger than x_0 .) The isotope effect on T_c for optimally-doped material is essentially zero (i.e., $T_c \propto M^{-\alpha}$ with $\alpha \approx 0$; M = ion mass) [28].

Normal state properties

It was realized at the outset that the normal state properties of the high T_c cuprate superconductors are unusual and appear to violate the Landau Fermi liquid paradigm [29-32]. Some researchers share the view that in so far as the normal state properties reflect the electronic structure that underlies high T_c superconductivity, it will be necessary to develop an understanding of the normal state before the superconducting state can be understood.

The anomalous normal state properties first identified in the high T_c cuprate superconductors include the electrical resistivity and Hall effect. The electrical resistivity $\rho_{ab}(T)$ in the ab-plane of many of the hole-doped cuprate superconductors near optimal doping has a linear temperature dependence between T_c and high temperatures ~ 1000 K, with an extrapolated residual resistivity $\rho_{ab}(0)$ that is very small; i.e., $\rho_{ab}(T) \approx \rho_{ab}(0) + cT$, with $\rho_{ab}(0) \approx 0$ and the value of c similar within different classes of cuprate materials [33]. The Hall coefficient R_H is inversely proportional to T and the cotangent of the Hall angle $\theta_H = R_H/\rho$ varies as T^2 ; i.e., $\cot(\theta_H) \equiv \sigma_{xx}/\sigma_{xy} = AT^2 + B$ [34]. The linear T -dependence

of $\rho(T)$ and the quadratic T-dependence of $\cot(\theta_H)$ have been attributed to longitudinal and transverse scattering rates τ_l^{-1} and τ_t^{-1} that vary as T and T^2 , respectively [35]. In the RVB model, the constant and T^2 terms in τ_t^{-1} and, in turn, $\cot(\theta_H)$, are ascribed to scattering of spinons by magnetic impurities and other spinons, respectively. Examples of the linear T-dependence of $\rho(T)$, inverse T-dependence of R_H , and quadratic T-dependence of $\cot(\theta_H)$ near optimal doping ($x \approx 0$) can be found in Figs. 4(a), 5(a), and 5(b) in which ρ_{ab} , R_H^{-1} , and $\cot(\theta_H)$ vs T data are displayed for the $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ system [36-38]. Experiments in which Ca^{2+} ions are counterdoped with Pr for Y in $YBa_2Cu_3O_{7-\delta}$ indicate that the Pr ions localize holes at a rate of ~one hole per substituted Pr ion [37]. Thus, as x is increased, the $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ system becomes more and more under-doped and T_c decreases, vanishing near the metal-insulator transition that occurs at $x_{mi} \approx 0.55$. Displayed in Fig. 4(b) is the T - x phase diagram for the $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ system which reveals the behavior of $T_c(x)$ as well as the Néel temperatures $T_N(x)$ for AFM ordering of Cu and Pr magnetic moments. It has been argued that the depression of T_c with x is primarily due to the decrease in the number of mobile holes with increasing Pr concentration, although magnetic pair breaking by Pr may also be involved [37,38]. In contrast, both $\rho_{ab}(T)$ and $\rho_c(T)$ of the optimally-doped electron-doped cuprate $Sm_{1.83}Ce_{0.17}CuO_{4-y}$ vary as T^2 , indicative of three dimensional Fermi liquid behavior [39].

The evolution of the normal ground state of the cuprates as a function of dopant concentration is particularly interesting. This is reflected in the temperature dependences of the ab-plane and c-axis electrical resistivities $\rho_{ab}(T)$ and $\rho_c(T)$ [40]. Both $\rho_{ab}(T)$ and $\rho_c(T)$ exhibit insulating behavior (i.e., $d\rho/dT < 0$) in the under-doped region, $\rho_{ab}(T)$ is metallic (i.e., $d\rho/dT > 0$) and $\rho_c(T)$ is insulating or metallic in the optimally-doped region, depending on the system, and $\rho_{ab}(T)$ and $\rho_c(T)$ are both metallic in the over-doped region. The linear T-dependence of $\rho_{ab}(T)$ and the insulating behavior of $\rho_c(T)$ suggest two dimensional non Fermi liquid behavior near the optimally-doped region, whereas the

metallic $\rho(T) \propto T^n$ with $n > 1$ reflects a tendency towards three dimensional Fermi liquid behavior in the over-doped region. Recent measurements in 61-tesla pulsed magnetic fields to quench the superconductivity have been particularly useful in elucidating the evolution of $\rho_{ab}(T)$ and $\rho_c(T)$ with dopant concentration in the $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ system [41]. Both $\rho_{ab}(T)$ and $\rho_c(T)$ were found to exhibit $-\ln T$ divergences in the under-doped region, indicative of a three dimensional non Fermi liquid [42]. As an example of the evolution of $\rho_{ab}(T)$ and $\rho_c(T)$ with doping, we again refer to the $\text{Y}_{1-x}\text{Pr}_x\text{Ba}_2\text{Cu}_3\text{O}_{7-\delta}$ system. Shown in Fig. 6 are $\rho_{ab}(T)$ and $\rho_c(T)$ data for $\text{Y}_{1-x}\text{Pr}_x\text{Ba}_2\text{Cu}_3\text{O}_{7-\delta}$ single crystals in the range of Pr concentrations $0 \leq x \leq 0.55$ [43]. The following features in the $\rho_{ab}(T)$ and $\rho_c(T)$ data in Fig. 6 are evident: a nonmonotonic evolution of $\rho_c(T)$ with x , the transformation of both $\rho_{ab}(T)$ and $\rho_c(T)$ from metallic to semiconducting with x , and the coexistence of metallic $\rho_{ab}(T)$ and semiconducting $\rho_c(T)$ for a certain range of doping. The nonmonotonic variation of $\rho_c(T)$ with x in Fig. 6(b) can be described with a phenomenological model [43] which assumes that the c-axis conductivity takes place via incoherent elastic tunneling between CuO_2 bilayers and CuO chain layers with a gap in the energy spectrum of the CuO chains (solid lines in Fig. 6(b)).

Perhaps the most remarkable aspect of the normal state is the pseudogap in the charge and spin excitation spectra of under-doped cuprates [44,45]. The pseudogap has been inferred from features in various transport, magnetic, and thermal measurements including $\rho_{ab}(T)$ [46,47,48], $R_H(T)$ [49], thermoelectric power $S(T)$ [50], NMR Knight shift $K(T)$ [51], NMR spin-lattice relaxation rate $1/T_1(T)$ [52,53,54], magnetic susceptibility $\chi(T)$ [49], neutron scattering [55], and specific heat $C(T)$ [56], as well as spectroscopic measurements such as infrared absorption [57,58,59] and angle resolved photoemission spectroscopy ARPES [60,61]. An example of the features in $\rho_{ab}(T)$ that are associated with the pseudogap can be seen in the $\rho_{ab}(T)$ data displayed in Figs. 4(a) and 6(a) for the $\text{Y}_{1-x}\text{Pr}_x\text{Ba}_2\text{Cu}_3\text{O}_{7-\delta}$ system. As the system becomes more under-doped with increasing x , $\rho_{ab}(T)$ deviates from linear behavior at higher temperature at a characteristic temperature T^*

which represents a crossover into the pseudogap state at $T < T^*$. The transport, thermal, magnetic, and infrared studies of the pseudogap have been carried out on several cuprate materials, including LSCO, YBCO-123, YBCO-124, and BSCCO-2212, while the ARPES investigations of the pseudogap have mainly focussed on BSCCO, although ARPES measurements have also been made on oxygen-deficient YBCO.

The ARPES measurements reveal several striking aspects of the pseudogap. The magnitude of the pseudogap has the same k -dependence in the ab-plane as the magnitude of the superconducting energy gap, with maxima in the directions of k_x and k_y and minima at 45° to these directions. In fact, the symmetry is consistent with $d_{x^2-y^2}$ symmetry inferred from Josephson tunneling measurements on hole-doped cuprates discussed in the next section. Furthermore, measurements of the temperature dependence of the pseudogap at the angles where it is a maximum show that the superconducting gap grows continuously out of the pseudogap and that the value of the sum of both gaps at low temperatures is constant, independent of the temperature T^* at which the pseudogap opens, or T_c . This is in marked contrast to the situation in conventional superconductors where the energy gap is proportional to T_c . These features are illustrated in Figs. 7(a) and (b) which show the k -dependence of the energy gap and the dependence of the maximum gap on temperature from the ARPES measurements of Ding et al. [60] on BSCCO-2212 samples with T_c 's of 87 K, 83 K, and 10 K. The pseudogap and the superconducting energy gap appear to be intimately related to one another, with the former the precursor of the latter. These results support the view that a unified theory of both the normal and superconducting states of the cuprates is imperative. The pseudogap and superconducting regions for the BSCCO-2212 derived from the ARPES measurements of Ding et al. [60] are shown in Fig. 8. Based upon investigations on LSCO, YBCO-123 and -124, BSCCO, and other systems, one can construct a generic T - x phase diagram which is shown schematically in Fig. 9. The phase diagram is very rich and contains insulating, antiferromagnetic, superconducting,

pseudogap, two dimensional (2D) non Fermi liquid like, and three dimensional (3D) Fermi liquid like regions.

A number of models and notions have been proposed to explain the part of the phase diagram delineated by the curves of T^* and T_c vs x (e.g., [62–67]). Generally, these models involve the local pairing of electrons (or holes) at the temperature T^* leading to a suppression of the low lying charge and spin excitations and the formation of the normal state pseudogap, followed by the onset of phase coherence at T_c that results in superconductivity. Since the phenomenon of superconductivity involves coherent pairing, a bell shaped curve of T_c vs x results as shown schematically in Fig. 10. For example, in resonating valence bond (RVB) models [30,62,63,64,67] which incorporate spin - charge separation into "spinons" with spin $s = 1/2$ and charge $q = 0$ and "holons" with $s = 0$ and $q = +e$, where e is the charge of the electron, the spinons become paired (spin pseudogap) at T^* and coherent pairing of holons (Bose-Einstein condensation) occurs at T_d , resulting in superconductivity.

Symmetry of the superconducting order parameter

During the last several years, a great deal of effort has been expended to determine the symmetry of the superconducting order parameter of the high T_c cuprate superconductors [68,69]. The pairing symmetry provides clues to the identity of the superconducting pairing mechanism which is essential for the development of the theory of high temperature superconductivity in the cuprates.

Shortly after the discovery of high T_c superconductivity in the cuprates, it was established from flux quantization, Andreev reflection, Josephson effect, and NMR Knight-shift measurements, that the superconductivity involves electrons that are paired in singlet spin states [70]. Possible orbital pairing states include s-wave, extended s-wave, and d-wave states. In the s-wave state, the energy gap $\Delta(\mathbf{k})$ is isotropic; i.e., $\Delta(\mathbf{k})$ is constant over the Fermi surface. This leads to "activated" behavior of the physical properties in the superconducting state for $T \ll \Delta$; e.g., the specific heat $C_e(T)$, ultrasonic

attenuation coefficient $\alpha_s(T)$, and NMR spin lattice relaxation rate $1/T_1(T)$ vary as $\exp(-\Delta/T)$. For the extended s-wave state, the energy gap $\Delta(T)$ is anisotropic; i.e., $\Delta(\mathbf{k})$ exhibits a variation over the Fermi surface which has the same symmetry as the rotational symmetry of the crystal. Similarly, for the d-wave case, the energy gap $\Delta(\mathbf{k})$ is anisotropic, but with a symmetry that is lower than the symmetry of the crystal. The d-wave state that is consistent with most of the experiments discussed below has $d_{x^2-y^2}$ symmetry, which can be expressed as $\Delta(\mathbf{k}) = \Delta_0[\cos(k_x a) - \cos(k_y a)]$. For both the extended s-wave and d-wave cases, $\Delta(\mathbf{k})$ vanishes at lines on the Fermi surface, resulting in a density of states $N_s(E)$ that is linear in energy E at low values of E . This leads to "power law" T^n ($n = \text{integer}$) behavior of the physical properties for $T \ll \Delta$; e.g., $C_e(T) \sim T^2$, the superconducting penetration depth $\lambda(T) \sim T$, and $1/T_1(T) \sim T^3$. The establishment of the symmetry of the superconducting order parameter requires the determination of both the magnitude and the phase of $\Delta(\mathbf{k})$. Shown in Fig. 11 is a schematic diagram of the variation of the energy gap over the Fermi surface and the density of states $N(E)$ vs E for the "s", extended "s", and "d" $_{x^2-y^2}$ states [68].

A number of different types of measurements have been performed on the high T_c cuprate superconductors which are sensitive to $|\Delta(\mathbf{k})|$. These include microwave penetration depth $\lambda(T)$ [71], microwave surface conductivity [72], NMR relaxation rate $1/T_1(T)$ [73], magnetic field dependence of the electronic specific heat $C_e(T)$ [74], thermal conductivity [75], ARPES [76], quasiparticle tunneling [77,78], and Raman scattering [79]. The results of these studies are generally consistent with a superconducting state with $d_{x^2-y^2}$ or extended s-wave symmetry for the hole-doped cuprates such as YBCO-123, 124, LSCCO, and BSCCO, and s-wave symmetry for the electron-doped superconductor NCCO [80,81,82]. However, the experiments discussed below indicate a dominant component of $d_{x^2-y^2}$ symmetry in the superconducting order parameter of hole-doped cuprates. This is a rather surprising result, considering the similarities in the structures of

the hole-doped and electron-doped superconductors and the fact that they are both derived from chemical substitution of antiferromagnetic insulating parent compounds.

Several different types of measurements which are sensitive to the phase of $\Delta(\mathbf{k})$ have been performed. These measurements, all of which involve the Josephson effect, include SQUID interferometry [83,84,85], single junction modulation [86,87], tricrystal ring magnetometry [88,89,90], c-axis Josephson tunneling [91,92,93,94,95], and grain boundary tunneling [96]. The SQUID interferometry, single junction modulation, and tricrystal ring magnetometry measurements were performed on YBCO, while tricrystal magnetometry experiments have also been carried out on TBCCO. These experiments indicate that the superconducting order parameter in the YBCO and TBCCO hole-doped materials has $d_{x^2-y^2}$ symmetry. However, c-axis Josephson tunneling studies on junctions consisting of a conventional superconductor (Pb) and twinned or untwinned single crystals of YBCO indicate that the superconducting order parameter of YBCO has a significant s-wave component [91,92]. Shown in Fig. 12 are Josephson critical current vs magnetic field B data for an untwinned YBCO/Pb tunnel junction at 4.2 K for $B \parallel a$ and $B \parallel b$ [92]. From the geometry of the junction, independent measurements of the penetration depth of Pb, and the period of the $I_c(B)$ Fraunhofer patterns, the YBCO penetration depths $\lambda_a = 1307 \text{ \AA}$ and $\lambda_b = 1942 \text{ \AA}$ were obtained. The large anisotropy ratio $(\lambda_a/\lambda_b)^2 \leq 2$ suggests significant pair condensation on the Cu-O chains in YBCO.

Recently, a new class of c-axis Josephson tunneling experiments in which a conventional superconductor (Pb) was deposited across a single twin boundary of a YBCO single crystal were performed by Kouznetsov et al. [97]. The Josephson critical current I_c was then measured as a function of magnitude and angle ϕ of magnetic field applied in the plane of the sample. For B aligned perpendicular to the twin boundary, a maximum in I_c as a function of B was observed at $B = 0$, similar to the case of the untwinned YBCO single crystal data shown in Fig. 12, whereas for B parallel to the twin boundary, a minimum in I_c was observed at $B = 0$. In all samples investigated, a clear experimental

signature of an order parameter phase shift across the twin boundary was observed. The results provide strong evidence for mixed d- and s-wave pairing in YBCO and are consistent with predominant d-wave pairing with $d_{x^2-y^2}$ symmetry and a sign reversal of the s-wave component across the twin boundary.

An experiment that provides evidence for a multicomponent superconducting order parameter in YBCO was recently reported by Srikanth et al [98]. Microwave measurements on YBCO single crystals prepared in BaZrO₃ (BZO) crucibles [99] yielded new features in the temperature dependence of the microwave surface resistance and penetration depth in the temperature range $0.6 \leq T/T_c \leq 1$ which the authors suggest constitute evidence for a multicomponent superconducting order parameter. These features were not observed in microwave measurements on YBCO single crystals prepared with yttria stabilized zirconia (YSZ) crucibles, suggesting that YBCO single crystals prepared in BZO crucibles are of higher quality than those prepared in YSZ crucibles [99].

Technological Applications of Superconductivity

Technological applications of superconductivity can be divided into two major areas: superconducting electronics and superconducting wires and tapes. While the widespread use of high T_c cuprate superconductors in technology has not yet been realized, steady and significant progress has been made towards this objective during the past decade. Recent developments indicate that high T_c cuprate superconductors will begin to have a significant impact on technology during the next 5 to 10 years. The applications in superconducting electronics that are likely to be realized on this time scale have been summarized in a recent article by Rowell [100]. In the order in which they are anticipated, these applications include: SQUIDS, NMR coils, wireless communications subsystems, MRI coils and NMR microscopes, and digital instruments. In the area of superconducting wires and tapes, applications that appear to be feasible within this same time period include: power transmission lines, motors and generators, transformers, current limiters, magnetic energy storage, magnetic separation, research magnet systems, and current leads.

An example of recent progress in the area of superconducting wires and tapes is the development of flexible superconducting ribbons consisting of deposits of YBCO on textured substrates which have critical current densities $J_c \sim 10^6 \text{ A/cm}^2$ in fields up to 8T at 64 K, a temperature that can be achieved by pumping on liquid nitrogen [14]. The performance of these prototype conductors in strong magnetic fields already surpasses that of NbTi and Nb₃Sn, which are currently used in commercial superconducting wires at liquid helium temperatures, in a comparable field range. The YBCO tapes are based on processes developed at two U. S. Department of Energy (DOE) National Laboratories, Los Alamos (LANL) and Oak Ridge (ORNL), under the DOE's National Superconductivity Program for Electric Power Applications. The essential step in both the LANL and ORNL processes is the preparation of a textured substrate, or "template," onto which a thick film of YBCO is deposited with well-aligned YBCO grains, that match the alignment of the underlying substrate. The alignment of the YBCO grains results in the high critical current density.

The LANL process, which is an extension of earlier work done at Fujikura Ltd. in Japan, uses a technique called "ion beam assisted deposition," or IBAD, to produce a preferentially oriented buffer layer on top of commercial nickel alloys, such as Hastelloy. The IBAD method uses four beams of charged particles to grow YSZ crystals with only one particular orientation on top of a ceramic oxide-buffered Hastelloy tape. Because of the excellent lattice match between YBCO and YSZ, the YBCO grains are "in-plane aligned" like the grains of the underlying YSZ layer.

The ORNL process involves fabricating long lengths of biaxially-textured metal (e.g., nickel) strips. Oxide buffer layers are then deposited on top of the metal substrate in order to transfer the alignment to the superconducting layer, while avoiding chemical degradation. The ORNL substrate technology is referred to as "RABiTS," or rolling-assisted, biaxially-textured substrates. In both the LANL and ORNL processes, pulsed

laser deposition (PLD) is used to deposit the superconducting YBCO layer and some of the buffer layers.

Shown in Fig. 13 is the magnetic field dependence of the critical current density for the IBAD and RABiTS based YBCO coated samples recently produced at LANL and ORNL. These coated samples operated in the liquid nitrogen temperature range clearly outperform the metallic superconductors (NbTi, Nb₃Sn) at 4.2 K. Furthermore, even in the worst field direction ($H \parallel c$), and for temperatures below 65 K, the short sample YBCO coated conductors operated in a 8-tesla background field have at least an order of magnitude higher J_c than pre-commercial BSCCO-2223 wire with no applied field.

The high T_c YBCO coated tapes have been shown to be extremely flexible and to retain the high current carrying capacity, so that they appear to be suitable for wound magnet and coil applications. A significant challenge that remains is the development of efficient, continuous commercially viable processes for fabricating long lengths of these high current carrying in-plane aligned YBCO coated conductors with uniform properties.

Other superconducting materials

Although the high T_c cuprates have been the focus of research on superconducting materials during the past decade, a number of other noteworthy novel superconducting materials have been discovered during this period. A few examples are briefly described below.

Rare earth transition metal borocarbides

Superconductivity was discovered in a series of compounds with the formula RNi_2B_2C with a maximum T_c of 16.5 K for $R = Lu$ [101,102]. These materials have attracted a great deal of interest because they display both superconductivity and magnetic order and effects associated with the interplay of these two phenomena, similar to the RRh_4B_4 and RMo_6X_8 ($X = S, Se$) ternary superconductors that were studied extensively during the '70's and early '80's [103]. Investigations of superconducting and magnetic order and their interplay have been greatly facilitated by the availability of large single

crystals of these materials [104]. Recently, values of T_c that rival the $T_c = 23$ K value of the intermetallic compound Nb_3Ge , the high T_c record holder for an intermetallic compound, have been found in mixed phase materials of the composition $\text{YPd}_5\text{B}_3\text{C}$ ($T_c = 23$ K) [105] and $\text{ThPd}_3\text{B}_3\text{C}$ ($T_c = 21$ K, $H_{c2}(0) \approx 17$ T) [106].

Sr_2RuO_4

The superconducting compound Sr_2RuO_4 has the same structure as the $\text{La}_{2-x}\text{M}_x\text{CuO}_4$ ($\text{M} = \text{Ba}, \text{Sr}, \text{Ca}; \text{Na}$) high T_c cuprate superconductors [107]. While the T_c of Sr_2RuO_4 is only ~ 1 K, this compound is of considerable interest because it is the only layered perovskite superconductor without Cu. The anisotropy of the superconducting properties of Sr_2RuO_4 is very large ($\gamma = \xi_{ab}/\xi_c \approx 26$). Although this anisotropy is larger than that of $\text{La}_{2-x}\text{M}_x\text{CuO}_4$, the in-plane and c-axis resistivities of Sr_2RuO_4 vary as T^2 at low temperature, indicative of Fermi liquid behavior. It has been suggested that Sr_2RuO_4 may exhibit p-wave superconductivity [108].

Alkali-metal doped C₆₀

Exploration of the physical properties of materials based on the molecule C_{60} revealed superconductivity with relatively high values of T_c in metal-doped C_{60} compounds [109]. For example, the fcc compounds K_3C_{60} and Rb_3C_{60} have T_c 's of 18 K and 29 K, respectively [25,26].

LiV_2O_4

The metallic transition metal oxide LiV_2O_4 , which has the fcc normal-spinel structure, has been found to exhibit a crossover with decreasing temperature from localized moment to heavy Fermi liquid behavior [110], similar to that which has been observed in strongly correlated f-electron materials [111]. At 1 K, the electronic specific heat coefficient $\gamma \approx 0.42$ J/mol K², is exceptionally large for a transition metal compound. No superconducting or magnetic order was detected in this compound to temperatures as low as ~ 0.01 K. This behavior can be contrasted with that of the isostructural compound

LiTi_2O_4 which displays nearly T-independent Pauli paramagnetism and superconductivity with $T_c = 13.7$ K [112].

Quantum Spin Ladder materials

Quantum spin ladder materials have attracted much recent interest [113,114]. These materials consist of ladders made of AFM chains of $S = 1/2$ spins coupled by inter-chain AFM bonds. Examples of 2-leg ladder materials are SrCu_2O_3 and $\text{LaCuO}_{2.5}$; an example of a 3-leg ladder material is $\text{Sr}_2\text{Cu}_2\text{O}_5$. Superconductivity has apparently been discovered in the ladder material $\text{Sr}_{0.4}\text{Ca}_{13.6}\text{Cu}_{24}\text{O}_{41.84}$ under pressure with $T_c \approx 12$ K at 3 GPa [115]. Part of the interest in quantum spin ladder materials stems from the fact that they are simple model systems for theories of superconductivity based on magnetic pairing mechanisms.

Concluding remarks

During the past decade, remarkable progress in the areas of basic research and technological applications has been made on the high T_c cuprate superconductors. The availability of high quality polycrystalline and single crystal bulk and thin film materials has made it possible to make reliable measurements of the physical properties of these materials and to optimize superconducting properties (e.g., J_c) that are important for technological applications. These investigations have provided important information regarding the anomalous normal state properties, the symmetry of the superconducting order parameter, and vortex phases and dynamics in the cuprates. The next decade of research on the high T_c cuprate superconductors as well as other novel superconducting materials promises to yield significant advances toward the development of a theory of high temperature superconductivity as well as the realization of technological applications of these materials on a broad scale. It is possible that significantly higher values of T_c will be found in new cuprate compounds or other classes of materials. Nature may even have some more surprises in store for us, as it did in 1986!

Acknowledgments

Assistance in preparing the Plenary Lecture on which this paper is based was kindly provided by D. Basov, D. K. Christen, D. L. Cox, M. C. de Andrade, R. Chau, N. R. Dilley, R. C. Dynes, A. S. Katz, M. P. Maley, N. J. McLaughlin, and F. Weals. This research was supported by the U. S. Department of Energy under Grant No. DE-FG03-86ER-45320.

References

- [1] J. G. Bednorz and K. A. Müller, *Z. Phys. B* 64 (1986) 189.
- [2] *Physical Properties of High Temperature Superconductors I - V*, D. M. Ginsberg, ed. (World Scientific, Singapore, 1989 - 1996).
- [3] See, for example, G. W. Crabtree and D. R. Nelson, *Physics Today*, April (1997) 38.
- [4] J. R. Gavalier, *Appl. Phys. Lett.* 23 (1973) 480.
- [5] S. N. Putilin, E. V. Antipov, O. Chmaissem, and M. Marezio, *Nature* 362 (1993) 226.
- [6] A. Schilling, M. Cantoni, J. D. Guo, and H. R. Ott, *Nature* 363 (1993) 56.
- [7] C. W. Chu, L. Gao, F. Chen, Z. J. Huang, R. L. Meng, and Y. Y. Xue, *Nature*, 365 (1993) 323.
- [8] M. Nuñez-Regueiro, J.-L. Tholence, E. V. Antipov, J.-J. Capponi, and M. Marezio, *Science* 262 (1993) 97.
- [9] M. B. Maple, Y. Dalichaouch, J. M. Ferreira, R. R. Hake, B. W. Lee, J. J. Neumeier, M. S. Torikachvili, K. N. Yang, H. Zhou, R. P. Guertin, and M. V. Kuric, *Physica B* 148 (1987) 155.
- [10] T. P. Orlando, K. A. Delin, S. Foner, E. J. McNiff, Jr., J. M. Tarascon, L. H. Greene, W. R. McKinnon, and G. W. Hull, *Phys. Rev. B* 36 (1987) 2394.
- [11] P. Chaudhari, R. H. Koch, R. B. Laibowitz, T. R. McGuire, and R. J. Gambino, *Phys. Rev. Lett.* 58 (1987) 2684.
- [12] For a review, see J. T. Markert, Y. Dalichaouch, and M. B. Maple, in *Physical Properties of High Temperature Superconductors I*, D. M. Ginsberg, ed. (World Scientific, Singapore, 1989), Ch. 6.
- [13] S. Jin, T. H. Tiefel, R. C. Sherwood, M. E. Davis, R. B. van Dover, G. W. Kammlott, R. A. Fastnacht, and H. D. Keith, *Appl. Phys. Lett.* 52 (1988) 2074.
- [14] R. Hawsey and D. Peterson, *Superconductor Industry*, Fall, 1996; and references cited therein.
- [15] A. W. Sleight, *Physics Today*, June (1991) 24.
- [16] J. D. Jorgensen, *Physics Today*, June (1991) 34.
- [17] R. J. Cava, R. B. van Dover, B. Batlogg, and E. A. Rietman, *Phys. Rev. Lett.* 58 (1987) 408.
- [18] Y. Tokura, H. Takagi, and S. Uchida, *Nature* 337 (1989) 345.
- [19] M. B. Maple, *MRS Bulletin* XV, No. 6 (1990) 60.
- [20] D. J. Scalapino, *Physics Reports* 250 (1995) 329; and references cited therein.
- [21] D. Pines, *Physica B* 199 & 200 (1994) 300; and references cited therein.

- [22] M. T. Béal-Monod and K. Maki, to be published.
- [23] L. F. Mattheis, E. M. Gyorgy, and D. W. Johnson, Jr., *Phys. Rev. B* 37 (1988) 3745.
- [24] R. J. Cava, B. Batlogg, J. J. Krajewski, R. Farrow, L. W. Rupp, Jr., A. E. White, K. Short, W. F. Peck, and T. Kometani, *Nature* 332 (1988) 814.
- [25] M. J. Rosseinsky, A. P. Ramirez, S. H. Glarum, D. W. Murphy, R. C. Haddon, A. F. Hebard, T. T. M. Palstra, A. R. Kortan, S. M. Zahurak, and A. V. Makhija, *Phys. Rev. Lett.* 66 (1991) 2830.
- [26] K. Holczer, O. Klein, S.-M. Huang, R. B. Kaner, K.-J. Fu, R. L. Whetten, and F. Diederich, *Science* 252 (1991) 1154.
- [27] S. Uchida, *Jpn. J. Appl. Phys.* 32 (1993) 3784 .
- [28] J. P. Franck, in *Physical Properties of High Temperature Superconductors IV*, D. M. Ginsberg, ed. (World Scientific, Singapore, 1994), p. 189.
- [29] See, for example, B. G. Levi, *Physics Today*, March (1990) 20; and references therein.
- [30] P. W. Anderson, *Science* 235 (1987) 1196.
- [31] R. B. Laughlin, *Science* 244 (1988) 525.
- [32] C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams, and A. E. Ruckenstein, *Phys. Rev. Lett.* 63 (1989) 1996.
- [33] See, for example, Y. Iye, in *Physical Properties of High Temperature Superconductors III*, D. M. Ginsberg, ed. (World Scientific, Singapore, 1992), Ch. 4.
- [34] T. R. Chien, Z. Z. Wang, and N. P. Ong, *Phys. Rev. Lett.* 67 (1991) 2088.
- [35] P. W. Anderson, *Phys. Rev. Lett.* 67 (1991) 2092.
- [36] M. B. Maple, C. C. Almasan, C. L. Seaman, S. H. Han, K. Yoshiara, M. Buchgeister, L. M. Paulius, B. W. Lee, D. A. Gajewski, R. F. Jardim, C. R. Fincher, Jr., G. B. Blanchet, and R. P. Guertin, *J. Superconductivity* 7 (1994) 97.
- [37] J. J. Neumeier, T. Bjornholm, M. B. Maple, and I. K. Schuller, *Phys. Rev. Lett.* 63 (1989) 2516.
- [38] J. J. Neumeier and M. B. Maple, *Physica C* 191 (1992) 158.
- [39] Y. Dalichaouch, C. L. Seaman, C. C. Almasan, M. C. de Andrade, H. Iwasaki, P. K. Tsai, and M. B. Maple, *Physica B* 171 (1991) 308.
- [40] See, for example, S. L. Cooper, and K. E. Gray, in *Physical Properties of High Temperature Superconductors IV*, D. M. Ginsberg, ed. (World Scientific, Singapore, 1994), Ch. 3.

- [41] G. S. Boebinger, Y. Ando, A. Passner, T. Kimura, M. Okuya, J. Shimoyama, K. Kishio, K. Tamasaku, N. Ichikawa, and S. Uchida, *Phys. Rev. Lett.* 77 (1996) 5417.
- [42] Y. Ando, G. S. Boebinger, A. Passner, T. Kimura, and K. Kishio, *Phys. Rev. Lett.* 75 (1995) 4662.
- [43] C. N. Jiang, A. R. Baldwin, G. A. Levin, T. Stein, C. C. Almasan, D. A. Gajewski, S. H. Han, and M. B. Maple, *Phys. Rev. B* 55 (1997) R3390.
- [44] See, for example, B. G. Levi, *Physics Today*, January (1996) 19; and references therein.
- [45] K. Levin, J. H. Kim, J. P. Lu, and Q. Si, *Physica C* 175 (1991) 449.
- [46] B. Bucher, P. Steiner, J. Karpinski, E. Kaldis, and P. Wachter, *Phys. Rev. Lett.* 70 (1993) 2012.
- [47] B. Batlogg, H. Y. Hwang, H. Takagi, R. J. Cava, H. L. Rao, and J. Kwo, *Physica* 235-240C (1994) 130; and references therein.
- [48] T. Ito, K. Takenaka, and S. Uchida, *Phys. Rev. Lett.* 70 (1993) 3995.
- [49] H. Y. Hwang, B. Batlogg, H. Takagi, H. L. Kao, J. Kwo, R. J. Cava, J. J. Krajewski, and W. F. Peck, Jr., *Phys. Rev. Lett.* 72 (1994) 2636.
- [50] J. L. Tallon, J. R. Cooper, P. deSilva, G. V. M. Williams, and J. W. Loram, *Phys. Rev. Lett.* 75 (1995) 4114.
- [51] W. W. Warren, Jr., R. E. Walstedt, G. F. Brennert, R. J. Cava, R. Tycko, R. F. Bell, and G. Dabbagh, *Phys. Rev. Lett.* 62 (1989) 1193.
- [52] M. Takigawa, A. P. Reyes, P. C. Hammel, J. D. Thompson, R. H. Heffner, Z. Fisk, and K. C. Ott, *Phys. Rev. B* 43 (1991) 247.
- [53] H. Alloul, A. Mahajan, H. Casalta, and O. Klein, *Phys. Rev. Lett.* 70 (1993) 1171.
- [54] T. Imai, T. Shimizu, H. Yasuoka, Y. Ueda, and K. Kosuge, *J. Phys. Soc. Japan* 57 (1988) 2280.
- [55] J. Rossat-Mignod, L. P. Regnault, C. Vettier, P. Bourges, P. Burlet, J. Bossy, J. Y. Henry, and G. Lapertot, *Physica* 185-189 C (1991) 86.
- [56] J. W. Loram, K. A. Mirza, J. R. Cooper, and W. Y. Liang, *Phys. Rev. Lett.* 71 (1993) 1740.
- [57] C. C. Homes, T. Timusk, R. Liang, D. A. Bonn, and W. N. Hardy, *Phys. Rev. Lett.* 71 (1993) 1645.
- [58] D. N. Basov, T. Timusk, B. Dabrowski, and J. D. Jorgensen, *Phys. Rev. B* 50 (1994) 3511.

- [59] A. V. Puchkov, P. Fournier, D. N. Basov, T. Timusk, A. Kapitulnik, and N. N. Kolesnikov, *Phys. Rev. Lett.* 77 (1996) 3212.
- [60] H. Ding, T. Yokoya, J. C. Campuzano, T. Takahashi, M. Randeria, M. R. Norman, T. Mochiku, K. Kadowaki, and J. Giapintzakis, *Nature* 382 (1996) 51.
- [61] A. G. Loeser, Z.-X. Shen, D. S. Dessau, D. S. Marshall, C. H. Park, P. Fournier, and A. Kapitulnik, *Science* 273 (1996) 325.
- [62] Y. Suzumura, Y. Hasegawa, and H. Fukuyama, *J. Phys. Soc. Japan* 57 (1988) 2768; H. Fukuyama, *Prog. Theoret. Phys. Suppl.* 108 (1992) 287.
- [63] N. Nagaosa and P. A. Lee, *Phys. Rev. B* 45 (1992) 966.
- [64] M. Randeria, N. Trivedi, A. Moreo, and R. T. Scalettar, *Phys. Rev. Lett.* 69 (1992) 2001.
- [65] V. J. Emery and S. A. Kivelson, *Nature* 374 (1995) 434.
- [66] Y. J. Uemura, in *Physics of the 10th Anniversary HTS Workshop on Physics, Materials, and Applications*, B. Batlogg, C. W. Chu, W. K. Chu, D. U. Gubser, and K. A. Müller, eds. (World Scientific, Singapore, 1996), p. 68.
- [67] S.-C. Zhang, *Science* 275 (1997) 1089.
- [68] See, for example, D. L. Cox and M. B. Maple, *Physics Today*, February (1995) 32.
- [69] See, for example, B. G. Levi, *Physics Today*, May (1993) 17; January (1996) 19; and references therein.
- [70] B. Batlogg, in *High Temperature Superconductivity; Proc. Los Alamos Symp. 1989*, K. S. Bedell, D. Coffey, D. E. Meltzer, D. Pines, and J. R. Schrieffer, eds. (Addison-Wesley, Redwood City, 1990), p. 37.
- [71] W. N. Hardy, D. A. Bonn, D. C. Morgan, R. Liang, and K. Zhang, *Phys. Rev. Lett.* 70 (1993) 3999.
- [72] D. A. Bonn, R. Liang, T. M. Riseman, D. J. Baar, D. C. Morgan, K. Zhang, P. Dosanjh, T. L. Duty, A. MacFarlane, G. D. Morris, J. H. Brewer, W. N. Hardy, C. Kallin, and A. J. Berlinsky, *Phys. Rev. B* 47 (1993) 11314.
- [73] J. A. Martindale, S. E. Barrett, K. E. O'Hara, C. P. Schlichter, W. C. Lee, and D. M. Ginsberg, *Phys. Rev. B* 47 (1993) 9155.
- [74] K. A. Moler, D. J. Baar, J. S. Urbach, R. Liang, W. N. Hardy, and A. Kapitulnik, *Phys. Rev. Lett.* 73 (1994) 2744.
- [75] H. Aubin, K. Behnia, M. Ribault, R. Gagnon, and L. Taillefer, *Phys. Rev. Lett.* 78 (1997) 2624.

- [76] Z.-X. Shen, D. S. Dessau, B. O. Wells, D. M. King, W. E. Spicer, A. J. Arko, D. Marshall, L. W. Lombardi, A. Kapitulnik, P. Dickenson, S. Doniach, J. DiCarlo, T. Loeser, and C. H. Park, *Phys. Rev. Lett.* 70 (1993) 1553.
- [77] D. Mandrus, J. Hartge, C. Kendziora, L. Mihaly, and L. Forro, *Europhys. Lett.* 22 (1990) 460.
- [78] D. Coffey and L. Coffey, *Phys. Rev. Lett.* 70 (1993) 1529.
- [79] T. P. Deveraux, D. Einzel, B. Stadlober, R. Hackl, D. H. Leach, and J. J. Neumeier, *Phys. Rev. Lett.* 72 (1994) 396.
- [80] D.-H. Wu, J. Mao, S. N. Mao, J. L. Peng, X. X. Xi, R. L. Greene, and S. M. Anlage, *Phys. Rev. Lett.* 70 (1993) 85.
- [81] S. M. Anlage, D.-H. Wu, J. Mao, S. N. Mao, X. X. Xi, T. Venkatesan, J. L. Peng, and R. L. Greene, *Phys. Rev. B* 50 (1994) 523.
- [82] B. Stadlober, G. Krug, R. Nemetschek, R. Hackl, J. L. Cobb, and J. T. Markert, *Phys. Rev. Lett.* 74 (1995) 4911.
- [83] D. A. Wollman, D. J. Van Harlingen, W. C. Lee, D. M. Ginsberg, and A. J. Leggett, *Phys. Rev. Lett.* 71 (1993) 2134.
- [84] D. A. Brauner and H. R. Ott, *Phys. Rev. B* 50 (1994) 6530.
- [85] A. Mathai, Y. Gin, R. C. Black, A. Amar, and F. C. Wellstood, *Phys. Rev. Lett.* 74 (1995) 4523.
- [86] D. A. Wollman, D. J. Van Harlingen, J. Giapintzakis, and D. M. Ginsberg, *Phys. Rev. Lett.* 74 (1995) 797.
- [87] J. H. Miller, Q. Y. Ying, Z. G. Zou, N. Q. Fan, J. H. Xu, M. F. Davis, and J. C. Wolfe, *Phys. Rev. Lett.* 74 (1995) 2347.
- [88] C. C. Tsuei, J. R. Kirtley, C. C. Chi, L.-S. Yu-Jahnes, A. Gupta, T. Shaw, J. Z. Sun, and M. B. Ketchen, *Phys. Rev. Lett.* 72 (1994) 593.
- [89] J. R. Kirtley, C. C. Tsuei, J. Z. Sun, C. C. Chi, L. S. Yu-Jahnes, A. Gupta, M. Rupp, and M. B. Ketchen, *Nature* 373 (1995) 225.
- [90] C. C. Tsuei, J. R. Kirtley, M. Rupp, J. Z. Sun, A. Gupta, M. B. Ketchen, C. A. Wang, Z. F. Ren, J. H. Wang, and M. Bhushan, *Science* 271 (1996) 329.
- [91] A. G. Sun, D. A. Gajewski, M. B. Maple, and R. C. Dynes, *Phys. Rev. Lett.* 72 (1994) 2267.
- [92] A. G. Sun, S. H. Han, A. S. Katz, D. A. Gajewski, M. B. Maple, and R. C. Dynes, *Phys. Rev. B* 52 (1995) R15731.
- [93] J. Lesueur, M. Aprili, A. Goulian, T. J. Horton, and L. Dumoulin, *Phys. Rev. B* 55 (1997) 3308.

- [94] A. G. Sun, A. Truscott, A. S. Katz, R. C. Dynes, B. W. Veal, and C. Gu, *Phys. Rev. B* 54 (1996) 6734.
- [95] R. Kleiner, A. S. Katz, A. G. Sun, R. Summer, D. A. Gajewski, S. H. Han, S. I. Woods, E. Dantsker, B. Chen, K. Char, M. B. Maple, R. C. Dynes, and J. Clarke, *Phys. Rev. Lett.* 76 (1996) 2161.
- [96] P. Chaudhari and S.-Y. Lin, *Phys. Rev. Lett.* 72 (1994) 1084.
- [97] K. A. Kouznetsov, A. G. Sun, B. Chen, A. S. Katz, S. R. Bahcall, J. Clarke, R. C. Dynes, D. A. Gajewski, S. H. Han, M. B. Maple, J. Giapintzakis, J.-T. Kim, and D. M. Ginsberg, to appear in *Phys. Rev. Lett.*, 1997.
- [98] H. Srikanth, B. A. Willemse, T. Jacobs, S. Sridhar, A. Erb, E. Walker, and R. Flükiger, *Phys. Rev. B* 55 (1997) R14733.
- [99] A. Erb, E. Walker, and R. Flükiger, *Physica C* 245 (1995) 245.
- [100] J. M. Rowell, *Solid State Commun.* 102 (1997) 269.
- [101] R. Nagarajan, C. Mazumdar, Z. Hossain, S. K. Dhar, K. V. Gopalakrishnan, L. C. Gupta, C. Godart, B. D. Padalia, and R. Vijayaraghavan, *Phys. Rev. Lett.* 72 (1994) 274.
- [102] R. J. Cava, H. Takagi, B. Batlogg, H. W. Zandbergen, J. J. Krajewski, W. F. Peck, Jr., R. B. van Dover, R. J. Felder, K. Mizuhashi, J. O. Lee, H. Eisaki, and S. Uchida, *Nature* 367 (1994) 252.
- [103] For a review, see *Superconductivity in Ternary Compounds II; Superconductivity and Magnetism*, M. B. Maple and Ø. Fischer, eds. (Springer, Berlin, Heidelberg, New York, 1982).
- [104] B. K. Cho, P. C. Canfield, and D. C. Johnston, *Phys. Rev. B* 52 (1995) R3844.
- [105] R. J. Cava, H. Takagi, B. Batlogg, H. W. Zandbergen, J. J. Krajewski, W. F. Peck, Jr., R. B. van Dover, R. J. Felder, T. Siegrist, K. Mizuhashi, J. O. Lee, H. Eisaki, S. A. Carter, and S. Uchida, *Nature* 367 (1994) 146.
- [106] J. L. Sarrao, M. C. de Andrade, J. Herrmann, S. H. Han, Z. Fisk, M. B. Maple, and R. J. Cava, *Physica C* 229 (1994) 65.
- [107] Y. Maeno, H. Hashimoto, K. Yoshida, S. Nishizaki, T. Fujita, J. G. Bednorz, and P. Lichtenberg, *Nature* 372 (1994) 532.
- [108] T. M. Rice and M. Sigrist, *J. Phys., Condens. Matter* 7 (1995) 643.
- [109] A. F. Hebard, *Physics Today*, November (1992) 26; and references therein.
- [110] S. Kondo, D. C. Johnston, C. A. Swenson, F. Borsa, A. V. Mahajan, L. L. Miller, T. Gu, A. I. Goldman, M. B. Maple, D. A. Gajewski, E. J. Freeman, N. R. Dilley, R. P. Dickey, J. Merrin, K. Kojima, G. M. Luke, Y. J. Uemura, O. Chmaissem, and J. D. Jorgensen, *Phys. Rev. Lett.* 78 (1997) 3729.

- [111] M. B. Maple, M. C. de Andrade, J. Herrmann, R. P. Dickey, N. R. Dilley, and S. Han, *J. Alloys and Compounds* 250 (1997) 585.
- [112] D. C. Johnston, *J. Low Temp. Phys.* 25 (1976) 145.
- [113] M. Takano, *Physica C* 263 (1996) 468.
- [114] S. Maekawa, *Science* 273 (1996) 1515.
- [115] M. Uehara, T. Nagata, J. Akimitsu, H. Takahashi, N. Môri, and K. Kinoshita, *J. Phys. Soc. Japan* 65 (1996) 2764.

Table caption

Table 1.

- (a) Some important classes of cuprate superconductors and the maximum value of T_c observed in each class.
- (b) Examples of the abbreviated designations (nicknames) used to denote cuprate materials.

Figure captions

Fig. 1.

Maximum superconducting critical temperature T_c vs date.

Fig. 2.

Temperature-dopant concentration (T - x) phase diagram delineating the regions of superconductivity and antiferromagnetic ordering of the Cu^{2+} ions for the hole-doped $La_{2-x}Sr_xCuO_4$ and electron-doped $Nd_{2-x}Ce_xCuO_{4-y}$ systems. AFM = antiferromagnetic phase, SG = spin-glass phase, and SC = superconducting phase. After Reference [19].

Fig. 3.

Crystal structures of La_2CuO_4 (T-phase) and Ln_2CuO_4 ($Ln = Pr, Nd, Sm, Eu, Gd$; T'-phase) parent compounds. From Reference [19].

Fig. 4.

- (a) In-plane electrical resistivity ρ_{ab} vs temperature T curves for $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ ($0 \leq x \leq 0.51$) single crystals. From Reference [36].
- (b) Temperature T vs Pr concentration x phase diagram for the $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ system, delineating metallic, superconducting, insulating, and antiferromagnetically ordered regions. From References [36,37].

Fig. 5.

- (a) Temperature dependence of the Hall carrier number $n_H = V/eR_H$ vs temperature T for $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ single crystals with different x values. From Reference [36].
- (b) Cotangent of the Hall angle $\cot(\theta_H)$ vs T^2 for $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ single crystals with different x values. Inset: Slope A and intercept B vs T . From Reference [36].

Fig. 6.

- (a) In-plane resistivity $\rho_{ab}(T)$ and (b) out-of-plane resistivity $\rho_c(T)$ for $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ single crystals. The solid lines in (b) are fits to the data using a model described in Reference [43]. Inset to Fig. 6(a): The configuration of leads used in the measurements. Inset to Fig. 6(b): Anisotropy ρ_c/ρ_{ab} vs Pr concentration x at different temperatures. The solid lines are guides to the eye. From Reference [43].

Fig. 7.

Momentum and temperature dependence of the energy gap estimated from leading edge shifts of ARPES spectra for BSCCO-2212. (a) k -dependence of the gap in the $T_c = 87$ K, 83 K, and 10 K samples, measured at 14 K. The inset shows the Brillouin zone with a large Fermi surface (FS) closing the (π, π) point, with the occupied region shaded. (b) Temperature dependence of the maximum gap in a near-optimal $T_c = 87$ K sample (circles), and underdoped $T_c = 83$ K (squares) and $T_c = 10$ K (triangles) samples. From Reference [60].

Fig. 8.

Schematic phase diagram of BSCCO-2212 as a function of doping. The filled symbols represent the T_c 's determined from magnetic susceptibility measurements. The open symbols are the T^* 's at which the pseudogap closes derived from the data shown in Fig. 7. From Reference [60].

Fig. 9.

Generic temperature T - dopant concentration x phase diagram for cuprates (schematic). The solid lines labelled T_N and T_c delineate the antiferromagnetic (AFM) and superconducting regions, respectively. The "hatched" line denoted T^* represents the crossover into the pseudogap state.

Fig. 10.

Schematic temperature T - dopant concentration x phase diagram for the cuprates. The dashed line labelled T^* represents the temperature below which some type of local pairing occurs leading to a suppression of low energy excitations and the formation of the pseudogap. The solid line labelled T_ϕ denotes the temperature below which phase coherence develops, resulting in superconductivity. The dark solid line labelled T_c delineates the superconducting region.

Fig. 11.

Fermi surface gap functions and densities of states of a superconductor with tetragonal symmetry for various pairing symmetries. The gap functions in the $k_z = 0$ plane (top) are represented by the light solid lines; distance from the Fermi surface (dark solid lines) gives the amplitude, a positive value being outside the Fermi surface, a negative value inside. The corresponding density of states for one-quasiparticle excitations $N(E)$ is shown below each gap function, with N_0 the normal state value. Gap node surfaces are represented by

the dashed lines. Left: The classic s-wave case, where the gap function is constant, with value Δ . This gives rise to a square-root singularity in $N(E)$ at energy $E = \Delta$. Middle: The extended s-wave case derives from pairs situated on nearest-neighbor square lattice sites in real space, with an approximate k-space form of $\cos(k_x a) + \cos(k_y a)$. For the Fermi surface shown here, the gap function has lines of nodes running out of the page. Right: A d-wave function of $x^2 - y^2$ symmetry. The extended s-wave and d-wave functions shown here each have a linear density of states up to order Δ , which measures the maximum gap amplitude about the Fermi surface. From Reference [68].

Fig. 12.

Josephson critical current I_c vs magnetic field B for c-axis Josephson tunneling between an untwinned $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ single crystal and a Pb counterelectrode for $B \parallel a$ (upper curve) and $B \parallel b$ (lower curve). The upper curve is offset by 0.5 mA along the y-axis. From Reference [92].

Fig. 13.

Magnetic field dependence of the critical current density for a range of short sample YBCO conductors produced using either IBAD or RABIT substrates. These data are compared with typical values obtained for NbTi and Nb_3Sn wires at 4.2 K and for BSCCO, oxide-powder-in-tube wires at 77K. From Reference [14].

Table 1. (a) Some important classes of cuprate superconductors and the maximum value of T_c observed in each class. (b) Examples of the abbreviated names (nicknames) used to denote cuprate materials.

(a)

Material	Max. T_c (K)
• $\text{La}_{2-x}\text{M}_x\text{CuO}_4$; M = Ba, Sr, Ca, Na	~40
• $\text{Ln}_{2-x}\text{M}_x\text{CuO}_{4-y}$; Ln = Pr, Nd, Sm, Eu; M = Ce, Th	~25
• $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$	92
• $\text{LnBa}_2\text{Cu}_3\text{O}_{7-\delta}$ Ce, Tb do not form phase Pr forms phase; neither metallic nor SC'ing	~95
• $\text{RBa}_2\text{Cu}_4\text{O}_8$	~80
• $\text{Bi}_2\text{Sr}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$ ($n = 1,2,3,4$)	($n = 3$) 110
• $\text{TlBa}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+3}$ ($n = 1,2,3,4$)	($n = 4$) 122
• $\text{Tl}_2\text{Ba}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$ ($n = 1,2,3,4$)	($n = 3$) 122
• $\text{HgBa}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+2}$ ($n = 1,2,3,4$)	($n = 3$) 133

(b)

Material	Nickname	T_c (K)
• $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$	YBCO; YBCO-123; Y-123	92
• $\text{Bi}_2\text{Sr}_2\text{Ca}_2\text{Cu}_3\text{O}_{10}$	BSCCO; BSCCO-2223; Bi-2223	110
• $\text{Tl}_2\text{Ba}_2\text{Ca}_2\text{Cu}_3\text{O}_{10}$	TBCCO; TBCCO-2223; Tl-2223	122
• $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_8$	HBCCO; HBCCO-1223; Hg-1223	133
• $\text{La}_{1.85}\text{Sr}_{0.15}\text{CuO}_4$	LSCO	40
• $\text{Nd}_{1.85}\text{Ce}_{0.15}\text{CuO}_{4-y}$	NCCO	25




Fig. 1

ELECTRON-HOLE SYMMETRY (QUALITATIVE)

Metallic ←———— Insulating —————→ Metallic




Fig. 2




Fig. 3





Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13

A possibility of emission of high frequency gravitational radiation from d-wave to s-wave type superconductor junctions.

Giorgio Fontana

Copyright April 1998

*Faculty of Science, University of Trento,
38050 Povo, TN, Italy.*

Recent measurements on a class of high-T_c superconductors (HTSC) have shown that Cooper-pairs binding may be associated to a d-wave, while in another class, d and s waves may coexist.

When d-wave Cooper-pairs are injected in a superconductor that can sustain s-wave binding, d-wave pairs decay to s-wave pairs and energy is irradiated by means of gravitons.

We show that in s-wave to d-wave type superconductor (SDS) junctions in an equilibrium condition no net gravitational wave energy is emitted, on the other hand under non equilibrium conditions a net gravitational wave energy is emitted by the junction.

Experiments which show a gravitational interaction between inhomogeneous high-T_c superconductors, under non equilibrium conditions, and test objects may be understood by accepting a possibility of emission of gravitational radiation from SDS junctions.

Introduction. In spite of the fact that quantum gravity is a theory currently under development (1), there are few elementary concepts that are involved in the application of some basic ideas related to the quantization of the gravitational field. For instance, it has been demonstrated (2) that gravitational waves, because of their quadrupolar nature, are carried by particles of spin 2, named gravitons. Moreover it has been suggested that atomic transitions for which the quantum number L changes by + or - 2, and for which the total quantum number J changes by 0 or + or - 2 are quadrupolar transitions and are permitted for the emission of gravitons, while the emission of photons is forbidden (2) (3).

Atomic transitions from orbitals 3d to 1s, 3d to 2s and 3d to 3s are possible candidates for transitions which may be applicable for the generation of gravitational radiation by atoms of a suitable material. The material could be pumped by photons and let decay gravitationally. The efficiency of the process depend on the half-lives of metastable states and the possibility of

obtaining a population inversion, as we have in LASERs.

While looking for a natural source of high frequency gravitational radiation it has been computed that the energy of gravitons involved in some possible stellar processes are very high, 14.4 keV for ⁵⁷Fe in the sun and 16.1 MeV from supernovae (2). Halpern's work (2) (3) also defines the gravitational counterpart of the laser, called a "gaser", for which we could prefer the name GWASER, for Gravitational Wave Amplification by Stimulated Emission of Radiation.

In spite of the fact that there is a lack of substantial theoretical and experimental work in this field, we could extend the above mentioned considerations to different quantum systems, looking for a possible candidate to efficient high frequency gravitational wave generators and detectors. Our attention has been focused on superconductors, where the carriers of electric current are bound in atom-like structures composed of two electrons, the Cooper-pairs. Cooper-pairs obey to quantum rules, both individually and collectively. With superconductors we are in the very lucky situation where the collective behaviour may show the individual state of each Cooper pair, being them collectively at the ground state for the specific superconducting material.

Very recently it has been experimentally observed the existence of two different symmetries of the order parameter in high temperature superconductor, a symmetry with an s-wave component and a symmetry with a d-wave component (5), therefore we know that Cooper-pairs are in s-orbitals and d-orbitals respectively. We can also predict that when Cooper-pairs move under non equilibrium conditions, i.e. under the effect of magnetic fields, from a superconductor where the symmetry of the order parameter is of type d to a superconductor where this symmetry is of type s, the Cooper-pairs must decay loosing energy by the emission of gravitons. This working principle is similar to the photon generation mechanism in LEDs and semiconductor injection LASERs, giving the gravitational counterparts that we may call GWEDs and injection GWASERs.

Pairing states in HTSC. The symmetry of the order parameter D (k) in superconductors has been found to have an s-wave symmetry in conventional low T_c superconductors and, after an intense research both theoretical and experimental (4) (5) (6) (7), it has been found that YBCO HTSC support both an s-wave symmetry and a d-wave symmetry depending on the probe direction. The s-wave component is due to the distortion by the orthorombic symmetry and it has been found that different HTCS which do not have the orthorombic distortion are nearly pure d-

wave (7). A background discussion on the symmetry of pairing states in both conventional and high Tc superconductors can be found in (4). We recall that electron pairing in low Tc superconductors is attributed to phonons, while in HTSC a strong magnetic coupling is believed to cause electron pairing, giving an inevitable *d*-wave symmetry and the high transition temperatures observed.

Emission of gravitational waves from HTSC.

The efforts required for the study of the nature of pairing states in HTSC has led to many experiments where Josephson junctions have been prepared using conventional superconductors injecting a supercurrent in HTSC through an insulating barrier. While the obtained interference patterns corroborated the cited theory, we may ask about the angular momentum quantum number of Cooper-pairs while traversing a *s*-wave to *d*-wave superconductors (SDS) junction. Experiments show that the supercurrent is not destroyed, it follows that each Cooper-pair should emit or absorb a particle carrying a spin of + or - 2, which may be identified with graviton and antigraviton respectively. In a fully superconductive circuit comprising the junction with no current, the number of Cooper-pairs that traverse the junction averages to zero, with no overall emission; on the other hand if a current flows in the superconductor, the distance between *s*-wave to *d*-wave junction and *d*-wave to *s*-wave junction along the current flow can become large enough to create two independent sources of gravitons and antigravitons which are not subjected to a local annihilation, therefore a net emission of gravitational wave could be possible, energy is subtracted from the current loop (flux vortex) which collapses and new loops are created by external fields. It is not known theoretically how much gravitational wave energy could be released by the process, as a complete theory of HTSC has not been fully developed and quantum gravity has only speculative foundations (1), nevertheless we may estimate the order of magnitude of the predicted phenomenon using some well known quantum relations.

We may estimate that the coupling energy that is released by a single decay could be a large fraction of

$$1) |T_{cp} - T_{cs}| k_B$$

where T_{cp} is the critical temperature of the *p*-wave superconductor and T_{cs} is the critical temperature of the *s*-wave superconductor and k_B the Boltzmann constant.

If we make the hypothesis that this fraction is a factor of one, we may write:

$$2) k_B |T_{cp} - T_{cs}| = hf$$

obtaining graviton frequencies of the order of hundreds of GHz.

The maximum power emitted by the process could be found with the hypothesis of currents slightly below the critical currents of most superconductors, and of about 10 kA/cm^2 , obtaining a power density of :

$$3) k_B |T_{cp} - T_{cs}| (10 \text{kA}/3.2 \times 10^{-19}) \text{ W/cm}^2$$

which is of the order of ten W/cm^2 .

We observe that the conditions required for the emission of gravitational radiation from a superconductor may have occurred incidentally during a sequence of experiments (8) (9) where a dual layered YBCO superconducting disk had been subjected to strong e-m fields. The disk consisted in two layers, the upper part had the orthorombic phase with the *c*-axis parallel to the surface of the disk and a T_c of 94.2 K, the lower part had a markedly different structure, which was 40 % tetragonal, and T_c equal to 60.5 K. It is not known which was the pairing state of Cooper pairs in the lower part, nevertheless it is supposed to be pure *d*-wave because of the absence of the orthorombic phase. We also observe that the orthorombic layer, because of its orientation supports both *d*-wave and *s*-wave for a supercurrent traversing the junction between the two layers. It follows that a current induced by the complex coil arrangement and traversing the partially SDS junction may have generated a beam of gravitational radiation.

The observed effects were attributed to a possible gravitational shielding; this hypothesis is not corroborated by any currently accepted theory or other supporting experiments, nor the observed region of "interaction", which has exactly the shape of the superconducting disk or toroid for distances of many disk diameters from the device surface, is compatible with a static field modification. On the contrary it is compatible with a beam of millimetric or submillimetric gravitational radiation generated by the decay of Cooper-pairs. The emission is orthogonal to the plane of the junction where small domains should have emitted coherent and thus beamed gravitational radiation, these domains are associated to current fluxes that locally align Cooper-pairs and locally determine the phase of the gravitational wave radiation; we also recall that HTSC are type 2 superconductors which allow the partial penetration of magnetic field by flux vortices.

It has been already shown that gravitational radiation may transfer energy and momentum (10), because of that, the repulsive effect observed in (8) (9) could be a straightforward

consequence of this property of gravitational radiation.

Conclusions. A mechanism for the efficient generation of high frequency gravitational radiation using high T_c superconductors has been proposed and discussed. The mechanism could explain a controversial experiment and clarify its essential features and construction details.

Additional experimental work and theoretical research will be required for a detailed quantitative analysis of the interaction between superconductors and gravitational radiation, nevertheless the possible existence of such a phenomenon will certainly promote the research in this field.

References.

1. Prima Facie Questions in Quantum Gravity (lecture), C. J. Isham, gr-qc/9310031 22 Oct 1993 preprint.
2. On the Gravitational Radiation of Microscopic Systems, L. Halpern, B. Laurent, IL NUOVO CIMENTO Vol. XXXIII, N. 3, 728, 1964.
3. On the stimulated photon-graviton conversion by an electromagnetic field, L. Halpern, B. Jouvet, Annales H. Poincare, Vol. VIII, NA1, p 25, 1968.
4. Phase-sensitive tests of the symmetry of the pairing state in high-temperature superconductors-Evidence for d symmetry, D.J. Harlingen, Reviews of Modern Physics, Vol. 67 No. 2, 515, 1995.
5. c-axis Josephson Tunneling between YBa₂Cu₃O_{7-x} and Pb: Direct Evidence for Mixed Order Parameter Symmetry in a High T_c Superconductor, K.A. Kouznetsov et al., PHYSICAL REVIEW LETTERS, 3050, 20 Oct. 1997.
6. Unusual paramagnetic phenomena in granular high-temperature superconductors-A consequence of d-wave pairing ?, M. Sigrist and T.M. Rice, , Reviews of Modern Physics, Vol. 67 No. 2, 503, 1995.
7. Angle-resolved photoemission spectroscopy study of the superconducting gap anisotropy in Bi₂Sr₂CaCu₂O_{8+x} , H. Ding et al. , PHYSICAL REVIEW B Vol. 54, N. 14, 54, 1 Oct. 1996-II
8. A possibility of gravitational force shielding by bulk YBa₂Cu₃O_{7-x} superconductor., E. Podkletnov and R. Nieminen, PHYSICA C, 203 (1992) 441-444.
9. Weak gravitational shielding properties of composite bulk YBa₂Cu₃O_{7-x} superconductor below 70 K under e.m. field., MSU-chem 95 – cond-mat/9701074v2 5 Feb 1997 preprint.
10. The gravitational wave rocket, W. B. Bonnor and M. S. Piper, Class. Quantum Grav. 14 (1997) 2895-2904.

Gravitation shielding properties of composite bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$

superconductor below 70 K under electromagnetic field.

E.E. Podkletnov, A.D. Levit

Report MSU-95 chem.

Abstract

High-temperature $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ ceramic superconductor with composite structure has revealed shielding properties against gravitational force in the state of levitation at temperatures below 70 K. Bulk ceramic superconductor with a shape of a toroidal disk was prepared using a conventional ceramic technology in combination with melt-texture growth.

Two solenoids were placed around the disk in order to initiate the current inside it and also to provide the rotation around its central axis. The samples placed over the rotating disk demonstrated a weight loss of 0.3-0.5%. When the rotation speed was slowly reduced by electromagnetic field in the solenoids, the shielding effect became considerably higher and reached 1.9-2.1% at maximum.

1. INTRODUCTION

The behavior of high-temperature ceramic superconductors under high-frequency magnetic field and their shielding properties are of great interest for practical application. Crystal structure seems to be the key factor determining all physical properties of bulk superconductors, and the interaction of the structure with the external and internal electro-magnetic fields might result in quite unusual effects. Despite a large number of studies [1-3] the nature of these interactions still remains unresolved. Our recent experimental work [4] clearly indicated that under certain conditions single-phase bulk dense $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ revealed a moderate shielding effect against gravitation force.

In order to obtain more information about this unusual phenomenon, a new installation was created which allowed operation with magnetic fields up to 2 T and frequencies up to 10^8 Hz at temperatures from 40 to 70 K. A new experimental technique was applied to modify the structure of the ceramic superconductor. All these efforts yielded a much bigger value in shielding effect (up to 2%) and provided good hopes for industrial application.

2. EXPERIMENTAL

The shielding superconducting element was made of the dense, bulk, almost single-phase $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ and had a shape of a toroidal disk with the outer diameter of 275 mm, the inner diameter of 20 mm, and the thickness of 10 mm. The preparation of the 123-compound consisted of mixing the initial oxides followed by calcining the powder at 930 C in air,

grinding, pressing the disk at 120 MPa, and sintering it in oxygen at 930 C for 12 h with slow cooling down to room temperature. After that the disk was kept in the furnace at 600 C, and the upper surface was quickly heated to 1200 C using planar high-frequency inductor as shown in Figure 1, with the following cooling down. During last heating the gap between the sample and the inductor was chosen precisely to provide only the heating of the upper 2 mm-thick layer of the disk, though high heat conductivity of the material caused the heating of some part of the material below this layer. Finally the disk was slowly cooled down to room temperature in a flow of oxygen and treated mechanically in order to obtain good balance during rotation.

Two solenoids were placed around the superconductor as shown in Figure 2. The gaps between these solenoids and the superconductor were big enough so that it could easily move about 20 mm in each direction. The toroidal disk was placed inside a cryostat equipped with a set of three coils which could keep it levitating when it reached the superconducting state. High-frequency electric current was first connected to a couple of two main solenoids around the toroidal disk and thus the current was initiated inside the ceramics. Then the system was cooled by liquid helium vapors to the temperature of 65-70 K and the disk became superconducting. After that the current was sent to the coils below the disk and the superconductor raised up (about 15 mm) because of the Meissner effect.

At this moment small current was sent to the main solenoids and the disk began rotating with increasing speed. The rotation speed was increased up to 5000 rpm. At this moment the measurements of weight for various subjects were taken. Finally the rotation speed was slowly reduced by changing the current in the main solenoids. The speed of rotation was controlled with the help of a laser beam reflected by a small piece of plastic light-reflecting foil attached to the disk.

The measurements of weight were taken constantly during this period too. The frequency of the electromagnetic field was varied from 10^3 to 10^8 Hz. Various materials were used for the measurements of weight over the levitating disk, including metals, glass, plastic, wood and so on. All these samples were placed hanging over the cryostat on a thread connected to sensible balance. The distance from the samples to the cryostat varied from 25 to 1500 mm. The weight of the samples was typically from 10 to 50 grains. Every precaution was taken to avoid all possible disturbances or the induced magnetic field and air flow.

The phase and crystal structure of the superconductor were studied using X-ray diffraction analysis (XRD) and scanning electron microscope (SEM) equipped with energy dispersive spectral (EDS) analyzer. The samples were cut layer by layer from the bulk ceramic disk.

The transition temperature T_c was determined from the resistive transition in a temperature variable cryostat, under zero magnetic field, using an AC current and sputtered golden contacts. The critical current density was measured for the samples cut from the top and from the bottom of the superconducting disk. Je

measurements were carried out at 75 K using an AC current, four-probe method and direct transport measurements.

3. RESULTS

The levitating superconducting ceramic disk revealed clearly visible shielding effect against the gravitational force even without rotation. The values of the weight loss for various samples were within the range of 0.05-0.07%. As soon as the main solenoids were switched on and the disk began rotating in the vapors of liquid helium, the shielding effect increased, and at the speed of 5000 rpm, the air over the cryostat began to raise slowly up to the ceiling. The particles of dust or smoke made the effect clearly visible. The boundaries of the flow could be seen clearly and they corresponded exactly to the shape of the toroid.

The weight of various samples decreased no matter what material they were made of. The samples of the same material but with different masses lost the same fraction of their weight. The loss of weight depended on the shape and the position of the sample. The maximum loss of weight could be reached if the sample was oriented with the flat surface parallel to the surface of the disk, so that the projection of the sample had the maximum area. The best measurement gave the value of the weight loss of 0.5%, while typical values were from 0.3 to 0.5%. The areas close to the inner edge of the toroid (5-7mm from the edge) gave lower values of shielding, from 0.1 to 0.25% only.

During the time when the rotation speed was decreased from 5000 to 3500 rpm using the solenoids as braking tools, the shielding effect reached maximum values and the weight loss of the samples was from 1.9 to 2-1% depending on the position of the sample from the outer edge of the disk. These peak values were measured during 25-30 seconds as the speed decreased rather quickly. Because of considerable vibration of the disk at the rotation speed close to 3000-3300 rpm further braking was done very quickly in order to avoid unbalanced rotation, and the measurements of weight could not be carried out.

The maximum shielding properties were observed for the maximum current inside the superconducting disk. According to preliminary measurements the upper layer of the disk was able of carrying over 15 thousand A/cm².

The maximum loss of weight of the samples was observed only at high frequencies of the magnetic field in the interval from 3.2 to 3.8 Mhz, but further detailed investigations are needed as the deviations of weight were sometimes rather large.

In order to prove that the weight loss occurred due to the difference of atmospheric pressure over the cryostat and below it but not as a result of some kind of high frequency field, the air pressure was measured with high precision using mercury barometer. The difference of pressures was clearly observed and was equal to 8mm for the maximum shielding effect.

The analysis of the cross-section of the ceramic YBa₂Cu₃O_{7-x} disk revealed the existence of two different zones of crystal structure. The upper part of

the disk (6-7 mm of thickness) had an orthorhombic structure typical for the quench and melt growth process [5,6] and consisted mainly of single-phase orthorhombic 123-compound. The material was dense and had pure and hardly visible grain boundaries. The grains had the size of less than 2 um and were oriented (75%) with c-axis parallel to the surface of the disk. The transition temperature for the material as defined by direct measurements was 94.2 K.

The lower part of the disk which had close contact with water-cooled base during high-frequency heat treatment had a structure with randomly-oriented grains, and the average grain size from 5 to 15 um. The porosity of this zone varied from 5 to 9%. The transition temperature T_c was equal to 65.5 K, and the material contained about 40% of the tetragonal phase. Crystal lattice parameters for these two layers as calculated from XRD are listed below, the values are given in mm.

$$\begin{aligned} \text{upper layer: } & a=0.381; b=0.387; c=1.165; \\ \text{lower layer: } & a=0.384; b=0.388; c=1.170; \\ & a=0.387; c=1.183. \end{aligned}$$

The first (upper) layer was quite homogenous with even distribution of elements in the volume of all the samples. EDS analysis showed the presence of small inclusions of Y₂BaCuO₅ in the lower layer.

4. DISCUSSION

The interaction of a superconducting ceramic body with gravitational field (or with gravitational waves) seems to be a complicated problem and cannot be explained by one law or one physical phenomenon. Common theory of field has little to say about the nature of gravitational field and its interaction with electric, magnetic and nuclear forces. Also an overwhelming explanation of the mechanism of high-temperature superconductivity has not yet been found. Still all these facts do not make the observed phenomenon less interesting.

In our previous work [4] the loss of weight of the sample over the levitating superconductor was extremely small and varied from 0.05 to 0.3%. At that time it was difficult to exclude entirely the influence of radio-frequency field because the sample was separated from the disk and the magnets by a thin plastic film. Now the superconductor was situated in a stainless steel cryostat and the influence of other factors causing the loss of weight should be negligible. The modification of the crystal structure of a superconductor allowed to obtain a composite body with a dense and oriented upper layer and a porous lower layer with random orientation of the grains. The upper layer is able of carrying high J_c current under considerable magnetic field while the lower layer cannot conduct high currents and is not resistant to the external magnetic field. The lower part of the disk with wide intergrain boundaries is also a source of a great number of Josephson junctions and is responsible for the direct and reverse, primary and secondary Josephson effect. The presence of tetragonal non-superconducting phase allows certain interaction with the external magnetic field.

A combination of two different crystal structures with different behaviour under magnetic field, creates a composite ceramic body with new properties. According to Faraday the replacement of a normal conductor in a magnetic field causes electric current inside it. Usually the magnetic field does not penetrate a superconductor more than a penetration depth, thus the interaction with the field is extremely small. But in the described experiment the superconductor also carries the high frequency electric current modified by Josephson effect. It is possible to admit that some interaction between the composite ceramic body and the external magnetic field takes place. This interaction depends on the coherence length, the flux pinning, the field frequency and the field force, the penetration depth and the parameters of the crystal lattice. These characteristics are interrelated in a complex way. As analysed in [7] pinning centres with different origins may exist inside the superconducting disk, and fluxes will be trapped at some of them. Fluxes trapped at weak centres will begin to move first and those trapped at strong ones will not move until the Lorentz force exceeds the pinning force. The overall current will be composed of the superposition of the flux motions with different speeds. Generally speaking, the quantized fluxes move as a bundle locally formed in a flux lattice by the magnetic interaction between them.

The temperature is also of great importance as it determines the state of energy inside a superconductor. The shielding effect was observed only below 70 K, while the ceramic disk became superconducting already at 94 K.

A specific state of energy of the superconducting body, caused by the conditions of the experiment, seems to change the behaviour of the whole atomic structure in such a way that the interaction with gravitational field (or waves) becomes different. Then, in order to keep a stable level of energy and a stable atomic and electronic structure, the superconductor absorbs some part of the energy from the gravitational field and slightly decreases it.

Figure 3 gives a schematic diagram of the shielding effect under usual atmospheric pressure (10000 kg/m^2). A thick horizontal line is a superconducting shield which reduces the gravitational force. Every object placed over the superconducting disk will loose only about 2% of its weight, which is not enough to make it levitating. Meanwhile the air over the object will also loose 2% of the weight. The bigger the surface area of the superconductor, the bigger becomes the lifting force. A 2% shielding creates a lifting force of about 200 Kg/m^2 because of the difference of atmospheric pressures under the shield and over it. The shield creates a kind of a vertical tunnel in the air with the reduced gravitational field. The observed effect also works in various gases and liquid media.

Various approaches to the problem of shielding properties of high T_c materials are possible and some of them can be based on the mechanism of superconductivity. All elementary particles are divided by their physical properties into two classes: bosons and fermions. It means that the behaviour of the particles is described by two different quantum statistics: Bose-Einstein or Fermi-Dirac. The electrons

of conductivity interact with each other according to Coulomb law and this interaction is strong enough; the binding energy of electrons in metals is comparable with their own kinetic energy, but at low temperatures electrons form sooner liquid than gas. Under vacuum equally charged electrons always undergo repulsion. Inside crystal ionic lattice the energy of the bond of free electrons can be negative and they can attract one another. Electrons with different impulses and spins are able to form pairs, producing one particle with zero spin and zero impulse.

The particles with zero spin obey the statistics of Bose-Einstein. These particles are called bosons and they "tend" to exist in equal quantum state. A good example of bosons properties is Bose-Einstein condensation: at low enough temperature, when kinetic energy of particles is small, they exist at the lowest energy level, revealing new properties. The condensate which is formed from Cooper pairs has no internal friction and demonstrates superfluidity. But superfluidity of a charged liquid is nothing more but superconductivity.

According to BCS theory if we regard weak bond, electrons of conductivity and phonons in the crystal lattice interact from time to time, particles collide but they preserve their individual positions and properties. If we deal with strong bond the interaction takes place all the time and free electrons and phonons exist no longer giving birth to a certain mixture called electron-phonon liquid. This liquid has specific properties and the behaviour of this electron-phonon mixture under various conditions is not yet studied. It is possible to admit that this liquid has some properties similar to those which are typical for magnetic liquids, especially if we take into consideration that magnetic hysteresis is characteristic for high T_c compounds. Also the experimental equipment described above has much in common with magneto-hydro-dynamic (MHD) generator. According to Faraday any conductor moving in the magnetic field produces electric current. The ceramic high T_c disc has a composite structure as it consists of a normal conductor and a superconductor both having good electrical contact. So we can admit that there is a certain interaction of the whole ceramic body with the external magnetic field.

According to the experiment ceramic superconductor kept at the temperature below 70 K does not reveal any unusual shielding if it has no contact with the alternative magnetic field. The shielding effect is extremely small when the sample is levitating over solenoid but has no current inside and is not rotating and has no rotation moment of its own [4]. Only composite structure allows some penetration of the external field inside the body. High-frequency electromagnetic field provides small wave length and facilitates the interaction with solid body. Various disturbances of the electro-magnetic picture inside the superconductor caused by Josephson junctions do not play any important role, though they should be studied in detail.

There are no grounds to claim that the rotation momentum of the disk interacts with gravitation force, but it seems that fast rotation is favourable for the stabilisation of the shielding effect. The biggest shielding was obtained when the rotation speed was

decreased by magnetic field in the solenoids. The most reasonable explanation at present is that the rotating disk when slowing down is loosing some part of its energy (and of the electron-phonon liquid as well) and in order to compensate [for] the loss of energy, absorbs some part of the energy of gravitation field. Superfluidity of liquid helium at low temperatures is based on the fact that the system tends to loose some energy in order to get a stable energy state. In the observed phenomenon the high T_c system tends to gain some additional energy also in order to keep a stable energy level and stable electronic structure. The first attempt to a theoretical explanation of the effect has been done by G.Modanese [8]. Further investigations that are in progress, will help to prove or change or to complete the present understanding of the observed phenomenon.

CONCLUSION

A superconducting ceramic levitating disk of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ with composite structure demonstrated stable and clearly visible shielding effect against gravitational force under 70 K and high-frequency electro-magnetic field. A combination of the high-frequency current inside the rotating toroidal disk and the high-frequency external magnetic field created a specific state of energy inside the material. This resulted in the ability of the superconductor to absorb or to modify the energy of the gravitational force and yielded 1.9-2.1% weight loss of the samples.

Various samples made of metals, plastic, ceramic, wood etc. were situated over the disk and their weight was measured with high precision. All the samples showed partial loss of weight no matter what material they were made of. In order to obtain the maximum loss of weight the samples should be oriented with the flat surface parallel to the surface of the disk. The maximum shielding effect (2.1%) was obtained when the speed of the rotation and the corresponding centrifugal force was slightly decreased by the magnetic field.

It was found that gravitation shielding effect depended on the temperature, the speed of rotation, the frequency and the force of the magnetic field. At present it seems a bit early to discuss the mechanisms and to give the detailed analysis of the observed phenomenon as farther investigation is necessary. It is not wise to regard any possibility of anti-gravitation but it seems reasonable to admit that under certain conditions the interaction of solid bodies with electromagnetic and nuclear force fields can demonstrate unusual properties, resulting in specific interaction of the solid substance with gravitational field. The experimentally obtained value of 2% shielding effect might be interesting for scientific and industrial application but no doubt that optimisation of the operation parameters might sufficiently improve the results.

ACKNOWLEDGEMENT

The authors are grateful to the Institute for High Temperatures of the Russian Academy of Sciences for the help in preparation of the unique superconducting ceramic discs and for the possibility to use their technological equipment.

REFERENCES

- (1) Ruse A B, Johansen T H, Bratsberg H, and Yang Z J, *Appl.Phys.Lett* 61, (1988) 1419.
- (2) Brand E H, *Am.J.Phys* 58, (1990) 43.
- (3) Lofland S, Huang M X, and Bhagat S M, *Physica C* 203, (1992) 271.
- (4) Podkletnov E and Nieminen R, *Physica C* 203, (1992) 441.
- (5) Murakami M, Morita M, Doi K and Miyamoto K, *Jpn.J.Appl.Phys.* 28, (1989) 1189.
- (6) Murakami M, Morita M and Koyama N, *Jpn.J.Appl.Phys.* 28, (1989) 1125.
- (7) Takizawa T, Kanbara K, Morita M and Hashimoto M, *Jpn.J.Appl.Phys.* 32, (1993) 774.
- (8) Giovanni Modanese, electronic preprint MPI-PhT/95-44, hep-th/9505094

[Search Results](#) [New Search](#)

3 records selected from **Compendex** for (((anti-gravity) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or e-mail results.

3. **Accession Number:** 6840449

Title: Unidentified flying objects with unusual propulsive techniques: Generation of antigravity by the absorption of gravitational vortices

Author(s): [Holwerda, Martin](#);

First author affiliation: New Energy Foundation ,3318 CG Dordrecht ,Netherlands

Serial Title: Journal of New Energy

Abbreviated Serial Title: J New Energy

Volume: v 6

Issue number: n 1

Issue date: Summer , 2001

ISSN: 1086-8259

CODEN: JNENFI

Publication year: 2001

Pages: p 169-178

Abstract: Propulsion and anti-gravity effect of an unidentified flying object (UFO) was discussed. A space liquid model establishing a relation of electrical, gravitational and magnetic phenomena to a concrete flow-pattern of space-liquid was used to explain the studied effects. The pair-formation process and electron-positron annihilation revealed the presence of electrons in the propulsive mechanism of the spacecraft. Gravity was found to be caused due to an excess of anticlockwise spiral-shaped space currents.

Abstract type: (Edited abstract)

Main heading: Spacecraft propulsion

Controlled terms: Gravitational effects; Astrophysics; Positrons; Electrons; Mathematical models

Uncontrolled terms: Unidentified flying objects (UFO); Anti-gravity effects; Positron annihilation

Treatment type: Theoretical;Experimental;

Classification codes: 656.1; 931.5; 657.2; 931.3; 921

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

4. **Accession Number:** 6840444

Title: The mechanical anti-gravity setups of AG-1, AG-2, AG-3

Author(s): [Selin, Alexey](#);

Serial Title: Journal of New Energy

Abbreviated Serial Title: J New Energy

Volume: v 6

Issue number: n 1

Issue date: Summer , 2001

ISSN: 1086-8259

CODEN: JNENFI

Publication year: 2001

Pages: p 128-132

Abstract: Two basic experiments confirm the existence of an energetic aether. One is the bending of light rays from remote stars as reported by A. Eddington in 1919. The second is the observation of the violet frequency shift using the Mossbauer effect as reported by R.V. Pound and G. Rebka in 1961. This paper is a discussion of these observations and the later observations of mechanical means to demonstrate anti-gravity are discussed.

Main heading: Gravitational effects

Controlled terms: Gravitation; Astrophysics; Weightlessness; Hydroelectric generators

Uncontrolled terms: Energetic aether; Mossbauer effects

Treatment type: Theoretical;Experimental;

Classification codes: 931.5; 657.2; 656.1; 611.1; 705.2

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

8. **Accession Number:** 4

Title: Anti-gravity implies infinite free energy

Author(s): [Bass, Robert](#);

Serial Title: Journal of New Energy

Abbreviated Serial Title: J New Energy

Volume: v 1

Issue number: n 4

Issue date: Winter, 1996

ISSN: 1086-8259

Publication year: 1996

Pages: p 76-78

Abstract: If the empirical Podkletnov Effect (Journal of New Energy, vol 2, no 2, ref 4, p 136) is confirmed to be physically valid as described, then it will be simple to produce unlimited free energy from some hitherto untapped renewable source. Indeed, the device described loc.cit. [cf. also Business Week, Sept. 30, 1996, p 42] produces a constant reduction in the measured weight of an arbitrary test mass when run in steady-state at constant power. But this means that the total energy input grows linearly with time, whereas the total energy output stored in a vertically-arrayed flywheel (driven by the torque producible from asymmetric weight-reduction and consequent unbalanced unidirectional torque generated as each successive portion of the flywheel's rim becomes tangent to the Podkletnov beam) grows quadratically with time. The announced result follows at once.

Abstract type: (Author abstract)

Number of references: 4 Refs.

Main heading: Free energy

Controlled terms: Rotating machinery; Flywheels; Calculations; Torque; Energy utilization; Acceleration; Energy storage

Uncontrolled terms: Angular momentum; Rotational kinetic energy

Treatment type: Theoretical;

Classification codes: 641.1; 601.1; 921.6; 931.1; 525.3

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

Search Results

New Search

Results Manager

[Search Results](#) [New Search](#)

7 records selected from **Compendex** for (((podkletnov) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or e-mail results.

1. **Accession Number:** 2001025492810

Title: New theoretical approach of the gravitational shielding: Podkletnov's effect

Author(s): [Buzea, C.Gh.](#); [Agop, M.](#);

First author affiliation: Inst of Technical Physics ,Iasi ,Rom

Serial Title: Physica C: Superconductivity and its Applications

Abbreviated Serial Title: Phys C Supercond Appl

Volume: v341-348 (I)

ISSN: 0921-4534

CODEN: PHYCE6

Conference Name: International Conference on Materials and Mechanisms of Superconductivity

High Temperature Superconductors VI

Conference date: Feb 20-Feb 25 2000

Conference location: Houston, TX, USA

Sponsor: Air Force Office of Scientific Research; Argone National Laboratory; DANKA; ISSO

Conference code: 57787

Publisher: Elsevier Science Publishers B.V., Amsterdam, Netherlands

Publication year: 2000

Pages: p 307-308

Abstract: We show an electromagnetic field can produce a gravitational Meissner effect and, as a result, a gravitational shielding (Podkletnov's effect).

Abstract type: (Author abstract)

Number of references: 7 Refs.

Main heading: Electromagnetic field effects

Controlled terms: Gravitational effects; Maxwell equations; Current density; Electric charge; Light velocity; Permittivity; Magnetic permeability; Superconducting materials

Uncontrolled terms: Gravitational shielding; Podkletnov effect; Gravitational Meissner effect; London equation; Mass density

Treatment type: Theoretical;

Classification codes: 701.2; 701.1; 931.5; 921.2; 741.1; 708.1

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

2. **Accession Number:** 4

Title: Anti-gravity implies infinite free energy

Author(s): [Bass, Robert](#);

Serial Title: Journal of New Energy

Abbreviated Serial Title: J New Energy

Volume: v 1

Issue number: n 4

Issue date: Winter, 1996

ISSN: 1086-8259

Publication year: 1996

Pages: p 76-78

Abstract: If the empirical Podkletnov Effect (Journal of New Energy, vol 2, no 2, ref 4, p 136) is confirmed to be physically valid as described, then it will be simple to produce unlimited free energy from some hitherto untapped renewable source. Indeed, the device described loc.cit. [cf. also Business Week, Sept. 30, 1996, p 42] produces a constant reduction in the measured weight of an arbitrary test mass when run in steady-state at constant power. But this means that the total energy input grows linearly with time, whereas the total energy output stored in a vertically-arrayed flywheel (driven by the torque producible from asymmetric weight-reduction and consequent unbalanced unidirectional torque generated as each successive portion of the flywheel's rim becomes tangent to the Podkletnov beam) grows quadratically with time. The announced result follows at once.

Abstract type: (Author abstract)

Number of references: 4 Refs.

Main heading: Free energy

Controlled terms: Rotating machinery; Flywheels; Calculations; Torque; Energy utilization; Acceleration; Energy storage

Uncontrolled terms: Angular momentum; Rotational kinetic energy

Treatment type: Theoretical;

Classification codes: 641.1; 601.1; 921.6; 931.1; 525.3

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

3. **Accession Number:** 1

Title: Alternative explanation of 'gravitational screening' experiments

Author(s): [de Podesta, Michael](#); [Bull, Martyn](#);

First author affiliation: Birkbeck Coll ,London ,Engl

Serial Title: Physica C: Superconductivity

Abbreviated Serial Title: Phys C Supercond

Volume: v253

Issue number: n1-2

Issue date: Oct 20, 1995

ISSN: 0921-4534

CODEN: PHYCE6

Publisher: Elsevier Science B.V., Amsterdam, Netherlands

Publication year: 1995

Pages: p 199-200

Abstract: We show that the phenomena tentatively ascribed by Podkletnov and Nieminen [Physica C 203 (1992) 441] to gravitational screening may be easily understood in terms of a buoyancy correction to the authors' weighing procedure.

Abstract type: (Author abstract)

Number of references: 2 Refs.

Main heading: Oxide superconductors

Controlled terms: Gravitation; Weighing; Silica; Electric currents; Buoyancy; Cryostats; Calculations; Density (specific gravity); Electric fields

Uncontrolled terms: Gravitational screening; Buoyancy correction; Weight reduction

Treatment type: Experimental;

Classification codes: 708.3.1; 931.5; 943.3; 804.2; 701.1; 931.2

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

4.

Accession Number: 1924832

Title: Electroconducting ceramics on the base of cerium, indium oxides and the variation of its properties under the action of different gaseous mediums

Author(s): [Akopov, F.A.](#); [Barykin, B.M.](#); [Novov, Yu.D.](#); [Pakhomov, E.P.](#); [Podkletnov, E.E.](#);

First author affiliation: Inst Vysokikh Temperatur RAN ,Moscow ,Russia

Serial Title: Ogneupory

Abbreviated Serial Title: Ogneupory

Issue number: n 2

Issue date: Feb, 1994

ISSN:0369-7290

CODEN: OGNPA2

Publication year: 1994

Pages: p 10-13

Abstract: High temperature electroconductivity, porosity, compressive and tensile strengthes, thermal stability of ceramics with varying ratio of cerium to indium oxides was studied. Effect of gaseous products of propane combustion on the electroconductivity was examined. N-type conductivity under a such conditions was higher due to the reduction of indium oxide. It was shown that the components of mixed ceramics did not interact.

Number of references: 4 Refs.

Main heading:Ceramic materials

Controlled terms: Refractory materials; Cerium compounds; Indium compounds; High temperature properties; Electric conductivity of solids; Porosity; Thermodynamic stability; Strength of materials; Combustion; Reduction

Uncontrolled terms: Electroconducting ceramics; Propane combustion; N-type conductivity; Mixed ceramics

Treatment type: Experimental;Applications;

Classification codes: 812.1; 812.2; 802.2; 801.4.1

Language: Russian.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Order Document from Linda Hall Library](#)

5.

Accession Number: 1502273

Title:Possibility of gravitational force shielding by bulk $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ superconductor

Author(s): [Podkletnov, E.](#); [Niemenen, R.](#);

First author affiliation: Tampere Univ of Technology ,Finl

Serial Title: Physica C: Superconductivity

Abbreviated Serial Title: Phys C Supercond

Volume:v 203

Issue number: n 3-4

Issue date: Dec 10, 1992

ISSN:0921-4534

CODEN: PHYCE6

Publication year: 1992

Pages: p 441-444

Abstract: Shielding properties of single-phase dense bulk superconducting ceramics of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ against the gravitational force were studied at temperatures below 77 K. A small non-conducting and non-magnetic sample weighing 5.48 g was placed over a levitating superconducting disk and the loss of weight was measured with high precision using an electro-optical balance system. The sample was found to lose from 0.05 to 0.3% of its weight, depending on the rotation speed of the superconducting disk. Partial loss of weight might be the result of a certain state of energy which exists inside the crystal structure of the superconductor at low temperatures. The unusual state of energy might have changed a regular interaction between electromagnetic, nuclear and gravitational forces inside a solid body and is responsible for the gravity shielding effect.

Abstract type: (Author abstract)

Number of references: 8 Refs

Main heading:High temperature superconductors

Controlled terms: Yttrium compounds; Copper oxides; Ceramic materials; Crystal structure; Electromagnetic shielding; Gravitational effects; Thermal effects; Superconductivity; Magnetic levitation

Uncontrolled terms: Yttrium barium copper oxides; Gravitational force shielding; Levitating superconducting disks; Disk rotation speed; Weight loss; Energy state; Electrooptical balance system

Treatment type: Applications;Experimental;

Classification codes: 708.3.1; 804.2; 812.1; 933.1.1; 701.2; 701.1

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

6. **Accession Number:** 1171584

Title:Morphology and microstructure of $\text{YBa}_2\text{Cu}_3\text{O}_7$ thin films on SrTiO_3 single crystal by RF magnetron sputtering

Author(s): [Podkletnov, E.E.](#); [Jaervinen, R.J.O.](#); [Vuorinen, P.T.](#); [Maentylae, T.A.](#);

First author affiliation: Tampere Univ of Technology ,Finl

Serial Title: Journal of Crystal Growth

Abbreviated Serial Title: J Cryst Growth

Volume:v 114

Issue number: n 1-2

Issue date: Oct, 1991

ISSN:0022-0248

CODEN: JCRGAE

Publication year: 1991

Pages: p 198-202

Abstract: Superconducting thin films of 123 compound were deposited 'in situ' on (100) single crystals of SrTiO_3 by RF magnetron sputtering. Oriented films with $T_c(R = 0)$ of 85 K and $J_c = 10^5 \text{ A/cm}^2$ (at 78 K in zero field) were obtained. In order to provide constant substrate temperature during the deposition and thus to obtain perfect crystal structure, the transparent substrate was coated from the back side by a non-transparent layer of 123 compound before the main 'in situ' sputtering. To reduce the resputtering effect the substrates were placed aside from the projection of the target on the substrate table. Film properties such as crystallinity, surface morphology and composition were analysed by X-ray diffraction (XRD), scanning electron microscopy (SEM) and X-ray energy dispersive spectroscopy (EDS).

Abstract type: (Author abstract)

Number of references: 12 Refs

Main heading:High Temperature Superconductors

Subheading: Thin Films

Controlled terms: Crystals--Growing; Yttrium Compounds; X-Rays--Diffraction; Microscopic Examination--Scanning Electron Microscopy; Ceramic Materials

Uncontrolled terms: Oxide Superconductors; Yttrium Barium Copper Oxides; RF Magnetron Sputtering

Treatment type: Experimental;

Classification codes: 708; 932; 741; 531; 812; 547

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

7. **Accession Number:** 1042102
Title:Cerium dioxide refractory materials
Author(s): [Akopov, F.A.](#); [Romanov, A.I.](#); [Podkletnov, E.E.](#); [Valjano, G.E.](#);
Monograph Title: Proceedings of the Unified International Technical Conference on
Conference Name: Proceedings of the Unified International Technical Conference on
Refractories - UNITECR '89
Conference date: Nov 1-4 1989
Conference location: Anaheim, CA, USA
Conference code: 14079
Publisher: Publ by American Ceramic Soc, Westerville, OH, USA
Publication year: 1989
Pages: p 1071
Document type: Conference Articles
Database: Compendex
Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Order Document from Linda Hall Library](#)

[Search Results](#)

[New Search](#)

Results Manager

Select Results Format

Citation

Abstract / Links

Detailed Record / Links

[New Display](#)

[Print](#)

[E-mail](#)

[Save/Download](#)

[Clear Selected Records](#)

[Privacy Policy](#) | [Contact Us](#) | [Feedback](#)

Copyright © 2000 - 2001 by Engineering Information Inc., Hoboken, New Jersey, U.S.A.

[Search Results](#) [New Search](#)

4 records selected from **Compendex** for (((gravitational shielding) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or e-mail results.

1. **Accession Number:** 2001025492810

Title: New theoretical approach of the gravitational shielding: Podkletnov's effect

Author(s): [Buzea, C.Gh.](#); [Agop, M.](#);

First author affiliation: Inst of Technical Physics ,Iasi ,Rom

Serial Title: Physica C: Superconductivity and its Applications

Abbreviated Serial Title: Phys C Supercond Appl

Volume: v341-348 (I)

ISSN: 0921-4534

CODEN: PHYCE6

Conference Name: International Conference on Materials and Mechanisms of Superconductivity

High Temperature Superconductors VI

Conference date: Feb 20-Feb 25 2000

Conference location: Houston, TX, USA

Sponsor: Air Force Office of Scientific Research; Argone National Laboratory; DANKA; ISSO

Conference code: 57787

Publisher: Elsevier Science Publishers B.V., Amsterdam, Netherlands

Publication year: 2000

Pages: p 307-308

Abstract: We show an electromagnetic field can produce a gravitational Meissner effect and, as a result, a gravitational shielding (Podkletnov's effect).

Abstract type: (Author abstract)

Number of references: 7 Refs.

Main heading: Electromagnetic field effects

Controlled terms: Gravitational effects; Maxwell equations; Current density; Electric charge; Light velocity; Permittivity; Magnetic permeability; Superconducting materials

Uncontrolled terms: Gravitational shielding; Podkletnov effect; Gravitational Meissner effect; London equation; Mass density

Treatment type: Theoretical;

Classification codes: 701.2; 701.1; 931.5; 921.2; 741.1; 708.1

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

2. **Accession Number:** 7

Title: Local gravitoelectromagnetic effects on a superconductor

Author(s): [Agop, M.](#); [Buzea, C.Gh.](#); [Nica, P.](#);

First author affiliation: Technical Univ 'Gh. Asachi', Iasi ,Rom

Serial Title: Physica C: Superconductivity and its Applications

Abbreviated Serial Title: Phys C Supercond Appl

Volume: v339

Issue number: n2

Issue date: Oct, 2000

ISSN: 0921-4534

CODEN: PHYCE6

Publisher: Elsevier Science Publishers B.V., Amsterdam, Netherlands

Publication year: 2000

Pages: p 120-128

Abstract: Maxwell's and London's generalized equations, generalized Meissner effect and gravitational shielding in an electromagnetic field are obtained. In such a context, we show that a neutral particle beam in an infinitely thin superconducting cylinder placed in the Earth's gravitoelectric field is focused, and that the penetration depth increases with the pulsation of the electromagnetic field.

Abstract type: (Author abstract)

Number of references: 25 Refs.

Main heading: Superconducting materials

Controlled terms: Gravitational effects; Electromagnetic dispersion; Maxwell equations; Electromagnetic shielding

Uncontrolled terms: Maxwell-London equations; Meissner effects; Gravitoelectromagnetic effects

Treatment type: Experimental;

Classification codes: 708.3; 931.5; 711.1; 701.1; 701.2; 921.2

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

3. **Accession Number:** 3443589

Title:Theoretical analysis of a reported weak-gravitational-shielding effect

Author(s): [Modanese, G.](#);

Serial Title: Europhysics Letters

Abbreviated Serial Title: Europhys Lett

Volume:v 35

Issue number: n 6

CODEN: EULEEJ

Publication year: 1996

Pages: p 413

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Order Document from Linda Hall Library](#)

4. **Accession Number:** 2

Title:Does a superconductor shield gravity?

Author(s): [Unnikrishnan, C.S.](#);

First author affiliation: Tata Inst of Fundamental Research ,Bombay ,India

Serial Title: Physica C: Superconductivity

Abbreviated Serial Title: Phys C Supercond

Volume:v266

Issue number: n1-2

Issue date: Jul 20, 1996

ISSN:0921-4534

CODEN: PHYCE6

Publisher: Elsevier Science B.V., Amsterdam, Netherlands

Publication year: 1996

Pages: p 133-137

Abstract: Recently there were some experimental observations which were interpreted as due to

a shielding of the gravitational interaction by a superconducting disc in a static configuration as well as when set in rotation. We examine the experiments in detail and point out some difficulties which should be eliminated before reliable results can be claimed. The data from these experiments provide an internal check on the correctness of the hypothesis and we argue that the observed results are inconsistent with the hypothesis of shielding and therefore they are not due to shielding of the Earth's gravity. Our preliminary experiments in the static case do not show any evidence for the reported shielding.

Abstract type: (Author abstract)

Number of references: 4 Refs.

Main heading: Superconducting materials

Controlled terms: Superconductivity; Magnetic levitation; Gravitational effects

Uncontrolled terms: Meissner effect; Gravitational shielding; Superconductor shield gravity

Treatment type: Experimental;

Classification codes: 708.3; 701.1; 701.2; 931.5

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

[Search Results](#) [New Search](#)

Results Manager

Select Results Format

Citation

Abstract / Links

Detailed Record / Links

[New Display](#)

[Print](#)

[E-mail](#)

[Save/Download](#)

[Clear Selected Records](#)

[Privacy Policy](#) | [Contact Us](#) | [Feedback](#)

Copyright © 2000 - 2001 by Engineering Information Inc., Hoboken, New Jersey, U.S.A.



[Take your sex life places it's never been – like Vancouver. For travel, click here.](#)

ONE MILLION UNIQUE USERS | REACH YOUR TARGET AUDIENCE | WE MAKE IT EASY!

ADVERTISE

on sites that dominate this region

A Service of The Seattle Times Company

[seattletimes.com](#)

[NWclassifieds](#)

[NWsource](#)

[seattletimes.com](#)
The Seattle Times

WEB ARCHIVE

:: Home delivery
:: Contact us
:: Search archive

[HOME](#)
[Site index](#)

[Web archive](#)

[Help](#)
[Other searches](#)

**Bellevue | Burlington
Lynnwood | Renton**

(866) 204-4444
Home of Service
Loaners for life
soundford.com

Health & Science: Friday, February 27, 1998

Why universe expands? Nature adores a vacuum

by K.C. Cole
Los Angeles Times

In what may be a major insight into the nature of space itself, astronomers have discovered evidence that the universe is expanding rapidly under pressure of an anti-gravity-like force first proposed by Albert Einstein.

While Einstein later called his proposal the worst blunder of his life, his idea has resurfaced over the past decade as a way to solve a host of cosmological conundrums, including suggestions that the universe is younger than some of its stars.

"This is the first believable direct evidence that there might be" such a repulsive force, said astrophysicist Rocky Kolb of Fermi National Accelerator Laboratory in Illinois. "It may have been a blunder for (Einstein) to think it was a blunder."

The discovery was made by an international group of astronomers who looked at exploding stars, or supernovas, that appear to be hurtling away from each other at surprisingly rapid rates.

Astronomers have long known that the universe has continued to expand ever since its explosive birth in the Big Bang. However, it was thought that the mutual gravitational attraction of all the matter and energy in the universe would slow the expansion down. The inward pull of gravity would put a brake on the explosive outward expansion.

These supernovas appeared to be so far away, however, that they imply that the expansion of the universe is actually speeding up, rather than slowing down. It is as if the supernovas were actually picking up speed from some outward pushing force.

"Our observations show that the universe is expanding faster today than yesterday," said University of California, Berkeley, astronomer Adam Riess, one of the scientists who presented the findings at a meeting at UCLA last week. After trying to rule out other explanations, he said, his team proposed the reason might be Einstein's long-abandoned repulsive force.

Anti-gravity it's not, however. Gravity is the mutual attraction of matter and energy. Anti-gravity would imply that matter and energy would somehow mutually repel. If gravity causes two planets to attract each other, anti-gravity would cause them to repel each other.

The repulsive force, however, is the property of empty space itself.

"There's energy in the vacuum of space, and that energy wants to stretch that vacuum," he said. The repulsive force doesn't cause the planets to push away from each other. Rather, it causes the space between them to expand.

"If it's true, it's the craziest thing you could think of," said physicist Lawrence Krauss, of Case Western Reserve University in Cleveland.

Krauss is pleased because he's been proposing for years that Einstein's cosmological constant - the repulsive force - would solve vexing problems. For example, it would resolve the possible contradiction between the age of the universe and the oldest stars.

Security Camera

have FUN!
feel SAFE!

Tiny,
Wireless,
Goes
Anywhere!

Go now!

FREE SHIPPING!

The problem would be solved by a universe that is expanding more rapidly today than yesterday, because such a universe would be older than a steadily expanding universe.

Copyright © 2002 The Seattle Times Company

[Search web archive](#)

Advertising



SeattleFSBO.com

[Buy or Sell Homes](#)
[Commission FREE!](#)



Find the perfect dining places in the Northwest.

NWSOURCE
We find it. You do it.

[seattletimes.com home](#)

[Local news](#) | [Sports](#) | [Business & technology](#) | [Education](#) | [Investigation & special projects](#)

[Nation & world](#) | [Personal technology](#) | [Obituaries](#) | [Editorials & opinion](#) | [Columnists](#) | [Arts & entertainment](#)

[Northwest Life](#) | [Health & science](#) | [Travel](#) | [Northwest Weekend](#) | [Pacific Northwest magazine](#)

[Home delivery](#) | [Contact us](#) | [Search archive](#) | [Site index](#)
[NWclassifieds](#) | [NWSOURCE](#) | [Advertising info](#)

[Copyright © 2002 The Seattle Times Company](#)

↑ [Back to top](#)

[PG Home](#)[PG News](#)[Sports](#)[Lifestyle](#)[A&E](#)[Photo Journal](#)[AP Wire](#)[Business](#)[Classifieds](#)[Web Extras](#)[Weather](#)[Health & Science](#)[Search](#)[PG Store](#)[PG Delivery](#)[post-gazette.com](#)**Headlines
by E-mail**[Nation & World](#)[Sports](#)[Local News](#)[Health & Science](#)[Neighborhoods](#)[Lifestyle](#)[Business & Technology](#)[Opinion](#)

Pitt physicist offers spin on a universal theory

Beyond the rim of mainstream physics, a professor offers his ideas on matter and the nature of the universe

Monday, April 06, 1998

By Byron Spice, Science Editor, Post-Gazette

Astrophysicists are abuzz this spring over observations of distant exploding stars that suggest the expansion of the universe is picking up speed.

It's a new twist on an old question: Will the expansion of the universe that began in the Big Bang 15 billion years ago continue forever, eventually stop under the tug of gravity, or even reverse itself in a Big Crunch?

Evidence that the expansion actually is accelerating, first announced in January by an international team of astronomers, is still preliminary and hotly debated. But it raises the intriguing possibility that some strange force may be at work - some sort of anti-gravity, or what Albert Einstein called the cosmological constant.

Physicist Ernest Sternglass, on the other hand, offers his own spin - literally - to the question. The universe, he suggests, is not only expanding but rotating. If the expansion is accelerating, it's due to nothing more than plain old centrifugal force.

"It's just inertial force," said Sternglass, 74, an emeritus professor of radiological physics at the University of Pittsburgh, "the same kind of force that keeps the moon from falling down." It's the same force that makes his grandchildren hold on tight as they twirl on a park merry-go-round.

The rotating universe will continue to expand until the force of gravity balances the inertial forces, resulting in a stable cosmos.

This view, which Sternglass explains in his book, "Before the Big Bang,"



**Physicist Ernest Sternglass (Bill Wade
Post-Gazette)**

published late last year by Four Walls Eight Windows, is consistent with the predictions of Immanuel Kant, the 18th-century German philosopher. But it is well outside the mainstream of contemporary physics.

Likewise, Sternglass' theory that the negatively charged electron and its oppositely charged twin, the positron, are the fundamental building blocks of the universe flies in the face of the standard model of particle physics.

"I'm very skeptical," said Allen Janis, a theoretical physicist at Pitt who has ruminated and argued with his friend Sternglass for four decades. "He's a very imaginative person ... but I'm not convinced ... There is a coherent, well-verified body of knowledge and this just doesn't fit with that at all."

Sternlass is accustomed to life outside the mainstream. It was Einstein, he says, who encouraged him to pursue his physics theories outside of academia while earning a living in applied science. Though his work at Westinghouse Research Laboratories led to development of the video camera tube that captured Neil Armstrong's first steps on the moon and to ultraviolet-light detectors used in orbiting telescopes, his outspoken concerns about the dangers of radioactive fallout and nuclear power plant emissions made him a controversial figure in the 1960s and '70s.

He says he is satisfied with his accomplishments and has no regrets about the choices he has made in his life and career. And he retains the enthusiasm of a young boy.

"When I look in the mirror," he said, "I cannot believe I'm as old as I am."

A meeting with Einstein

It was a decidedly fresh-faced Sternlass who uneasily rang the doorbell at Einstein's house in Princeton, N.J., one cool spring day in 1947.

Newly hired by the Naval Ordnance Laboratory, Sternlass had been asked to explore the possibility of developing a night vision system. He had some ideas for building a television camera that could detect infrared light and he had written to Einstein in hopes of discussing them with the great physicist.

Dressed in a baggy gray gym suit, Einstein showed Sternlass to the back porch. What Sternlass expected to be a brief discussion turned into a five-hour conversation. Einstein and Sternlass, who had immigrated from Berlin with his physician parents in 1938, chatted in German about the nature of light particles called photons, about Einstein's dislike for quantum mechanics and his frustration with trying to develop a unified theory that would explain all of the forces of the universe.

"I found him to be very depressed and very sad," Sternlass recalled. Einstein was upset that he had made little progress on his unified field theory, his wife had died,

and his son suffered from mental illness. "Here was this man so widely admired and respected ... yet he felt a failure."

What most confounded Sternglass was Einstein's career advice. Sternglass was considering going back to school for an advanced degree in physics. Einstein advised him to pursue physics on his own and, perhaps most importantly, always have "a cobbler's job."

"Don't do what I have done," Sternglass, in his book, quotes Einstein as saying. He made a mistake, he told Sternglass, when he left his job as a patent examiner for a position with the University of Berlin. He had no real duties at the university, Einstein explained: "Nothing to do except wake up and solve the problems of the universe every morning. Nobody can do that."

Sternglass decided to return to Cornell University for graduate studies, eventually earning his doctorate, but he followed Einstein's advice in seeking a "cobbler's job." In 1952, he moved to Pittsburgh to join the Westinghouse lab. There, he could continue his studies in electron scattering and a phenomenon known as secondary electron emission, which interested Westinghouse because of its potential use in building electronic X-ray image intensifiers.

In 1967, he joined the Pitt Medical School's radiology department, where he investigated electronic techniques that would permit lower doses of diagnostic X-rays and applied computer techniques to enhance diagnostic images.

Earning and learning

While these jobs put food on the table, Sternglass pursued his interest in physics on the side. As Einstein had predicted, this arrangement allowed him to pursue far-out ideas and make his mistakes in private.

In 1963, Sternglass' work gained him a brief moment of fame. At a meeting of the American Physical Society in New York City, he outlined his theory on the nature of matter. He argued that all matter is composed of electrons and positrons.

Later that year, however, Murray Gell-Mann and George Zweig of the California Institute of Technology developed a theory of quarks, fractionally charged subatomic particles that comprise the protons and neutrons of atomic nuclei. Sternglass' theory was soon forgotten.

He would gain much more fame -- or infamy -- for his opposition to nuclear power. He argued that even low levels of nuclear fallout and radioactive emissions from nuclear power plants could damage children, raising infant mortality rates and depressing Scholastic Aptitude Test scores.

The vast majority of scientists rejected these notions. Neither Westinghouse nor Pitt ever tried to muzzle him, but Sternglass knows that his stands isolated him within the scientific community.

"Many of my former friends were very upset," he said.

His own interest in electrons and positrons was revived in 1974 when physicists at Brookhaven National Laboratory and the Stanford Linear Accelerator Center discovered a particle they called the J/Psi meson. It was both an unusually dense and unusually long-lived particle and it got Sternglass wondering about the nature of the "primeval atom."

The primeval atom was a concept developed by Georges Lamaitre, a Belgian priest and astrophysicist, in the 1920s. He envisioned the universe being created by the breakup of this primeval atom, an idea that presaged the Big Bang theory.

Sternglass saw this primeval atom as an electron-positron pair. This pair of particles rapidly orbited each other and contained all of the mass of the universe.

In a series of steps beginning billions of years before the Big Bang, this electron-positron pair went through a series of splits, creating thousands of electron-positron "seed pairs" from which galaxies later would emerge.

By contrast, according to the standard model of particle physics, the universe was condensed into an infinitely dense point before the Big Bang.

Sternglass said his model and the standard model don't differ markedly in terms of how elementary particles were formed in the minutes following the Big Bang.

Astrophysicists have long worried that the universe behaves as if it has much more mass than humans can see. Sternglass suggests some of this "missing mass" may be in the form of seed pairs that did not expand immediately after the Big Bang and remain sprinkled through the universe.

A universe in ether

In addition to suggesting that the universe is rotating, Sternglass resurrects the 3,000-year-old notion of the universe existing within an "ether," a liquid-like medium. Scientists once considered ether essential. For instance, scientists who observed light behaving like a wave reasoned that light would have to move through some sort of medium if it was to make waves.

But Einstein's photoelectric theory dispensed with the need for ether. He showed that light behaved as a particle, a photon. His theory of relativity also dispensed with ether, by showing that the speed of light need not be measured relative to an ether.

In Sternglass' view, matter exists as circular vortices -- something like a smoke ring -- moving through an ether. Just as air is capable of uprooting trees and tossing cars when whipped into a tornado, these vortices transform energy into mass, he said.

No one has ever been able to prove that ether exists, Sternglass acknowledged, but

that doesn't necessarily mean it doesn't.

"There are fashions in science just as there are fashions in art," he added. Ether is clearly out of fashion.

Fashions do exist, "but that's a double-edged sword," said Adolf Grunbaum, the Andrew Mellon Professor of philosophy of science at Pitt.

"The mere fact of the existence of fashions in science does not show, in a given case, that an unorthodox point of view is necessarily soundly supported by the evidence."

If most physicists aren't ready to embrace Sternglass' view, they might have trouble dismissing it, admitted Pitt's Janis. "As far as specific predictions he makes, there is nothing at the moment that could convincingly demonstrate that his ideas are wrong."

One important piece of evidence might come once a new particle accelerator, called the Large Hadron Collider, is completed at the European Laboratory for Particle Physics, or CERN, near Geneva, Switzerland. Scientists say the machine might produce a particle called the Higgs; if this particle is proven to exist, scientists believe it will confirm much of the standard model. Sternglass' theory predicts no such particle exists.

In the meantime, Sten Odenwald, an astronomer who works under contract with Goddard Space Flight Center in Greenbelt, Md., and runs the popular Astronomy Cafe Web site, said evidence from the Cosmic Background Explorer satellite in the early 1990s suggests the universe is not rotating, or at least not rotating much.

And even the recent evidence of possible acceleration of the universe's expansion may be questionable, Odenwald said. Those findings come from observing just a handful of exploding stars called supernovae. More observations could alter those findings dramatically.

Sternglass is realistic about his chances of convincing other physicists.

"Let's face it: When I present these things at astronomical meetings, it's a 10-minute presentation and people don't know what to make of it," he said.

"Nobody may believe me until after I'm dead," but that's OK, he added. "You have to take the long view."



There are seven deadly sins. One for each night out. For nightlife, click here.

ONE MILLION UNIQUE USERS | REACH YOUR TARGET AUDIENCE | WE MAKE IT EASY!

ADVERTISE

on sites that dominate this region

A Service of The Seattle Times Company

seattletimes.com

[NWclassifieds](#)

[NWsource](#)

seattletimes.com
The Seattle Times

WEB ARCHIVE

:: Home delivery
:: Contact us
:: Search archive

[HOME](#)
[Site index](#)

[Web archive](#)

[Help](#)
[Other searches](#)



(866) 204-4444
Home of Service
Loaners for life
www.soundford.com

Nation & World: Friday, January 23, 1998

Warped idea? NASA grasps for the stars

by Robert S. Boyd
Knight Ridder Newspapers

WASHINGTON - Not content with exploring our dinky little solar system, NASA is studying ways to travel across trillions of miles of empty space to the stars - first by robots, ultimately perhaps by humans.

NASA Administrator Daniel Goldin says he hopes the space agency will be able to launch an interstellar spaceship within the next 25 years and to reach an alien planet around a nearby star by the end of the 21st century, when today's children's children will be alive.

"We're going to want to send spacecraft toward distant worlds," Goldin told the American Astronomical Society here. "It sounds a little far out, but we've come a long way already. We can get there with a robotic explorer in a few generations."

As a preliminary step, Goldin has directed NASA's Jet Propulsion Laboratory in Pasadena, Calif., which managed the spectacular Mars Pathfinder landing last summer, to develop plans for an "interstellar precursor mission" - perhaps using exotic technology such as sails powered by light, or anti-gravity, warp drives or wormholes.

Goldin said NASA is not expecting short-term practical benefits from interstellar travel, but rather sees it as a logical continuation of humanity's age-old itch to explore. Furthermore, he believes researchers should push beyond the frontiers of existing technology.

Even enthusiasts agree that star travel won't be easy. Landing on the moon or Mars is a stroll on the beach compared with the difficulties of reaching even the nearest star, Alpha Centauri, 24 trillion miles away.

The basic problem is distance. If the sun were the size of a tennis ball, Pluto would be 100 yards away, but 600 lonesome miles would lie between Earth and Alpha Centauri. The star's light, traveling at 186,281 miles per second, takes more than four years to reach the Earth.

The current distance record for a spacecraft - 6.3 billion miles - is held by NASA's Pioneer 10, which was launched 26 years ago and is coasting away from the sun at 27,000 miles per hour. At that rate, it would take 100,000 years to get to Alpha Centauri.

So, fundamental breakthroughs in physics, chemistry and biology would be required. Today's chemical-fueled rockets don't have enough oomph.

Last year, NASA sponsored a workshop in Cleveland on exotic space-propulsion systems; 84 government, university and industrial scientists and engineers plotted how to break the universal speed limit.

"Imagine it," said Catherine Asaro, a physicist in Greenbelt, Md., who attended. "NASA invites you to an invitation-only conference, sits you down at a table with some of the best minds in physics, and says, 'OK, figure out how to go faster than the speed of light!'"

According to Einstein's theory of relativity, an object cannot exceed the speed of light.

Security Camera

have FUN!
feel SAFE!

Tiny,
Wireless,
Goes
Anywhere!

Go now!

FREE SHIPPING!

But science is constantly doing things once thought impossible, such as cloning an adult sheep, creating anti-matter or recovering the DNA of a Neanderthal man.

"One should never say 'never' in science," Floyd Bloom, editor of the respected journal Science, wrote last month. "Barriers that are acceptable by today's logic could eventually yield to persistent research."

One technique discussed at the Cleveland conference is to somehow warp, or bend, the fabric of space-time to shorten the distance between Earth and a star. A rough analogy would be folding a piece of paper to connect two points that used to be separated. Such a shortcut through space is known as a wormhole.

Background and Related Info.

- [NASA's Web site](#)

Copyright © 2002 The Seattle Times Company

[Search web archive](#)

Advertising

SeattleAreaHomes.com



[Search For Homes. Free Professional Assistance](#)

FOR RENT AFFORDABLE HOUSING FOR STUDENTS.



NWclassifieds

[seattletimes.com home](#)

[Local news](#) | [Sports](#) | [Business & technology](#) | [Education](#) | [Investigation & special projects](#)

[Nation & world](#) | [Personal technology](#) | [Obituaries](#) | [Editorials & opinion](#) | [Columnists](#) | [Arts & entertainment](#)

[Northwest Life](#) | [Health & science](#) | [Travel](#) | [Northwest Weekend](#) | [Pacific Northwest magazine](#)

[Home delivery](#) | [Contact us](#) | [Search archive](#) | [Site index](#)
[NWclassifieds](#) | [NWsource](#) | [Advertising info](#)

[Copyright © 2002 The Seattle Times Company](#)

↑ [Back to top](#)



ELSEVIER

A New Theoretical Approach of the Gravitational Shielding: Podkletnov's Effect

C. Gh. Buzea^a and M. Agop^b^aInstitute of Technical Physics, D. Mangeron 47, 6600-Iasi, Romania^bTechnical University "Gh. Asachi", Physics Department, 6600-Iasi, Romania

We show an electromagnetic field can produce a gravitational Meissner effect and, as a result, a gravitational shielding (Podkletnov's effect).

1. INTRODUCTION

Gravitational Meissner effect can not be found in usual matter. Consequently, gravitational shielding does not occur here. However, particular space-time structures like wormholes [1] or cosmic strings [2], which violate the null energy condition (NEC theorem), might reveal such peculiar features [3].

In this paper we show that the presence of an electromagnetic field can cause a gravitational Meissner effect and, as a result, a gravitational shielding.

2. GENERALIZED MAXWELL AND LONDON EQUATIONS

It is well known that Einstein's field equations in the weak field approximation and at low speeds reduce to Maxwell's type equations [4]. Then, by superposition, the generalized Maxwell equations (of the electromagnetic and gravitational field) become :

$$\begin{aligned} \nabla \cdot \mathbf{g} &= \left(-4\pi G + \frac{1}{\epsilon_0 m^2} \frac{q^2}{m^2} \right) \rho \\ \nabla \cdot \mathbf{B} &= 0 \\ \nabla \times \mathbf{g} &= -\frac{\partial \mathbf{B}}{\partial t} \\ \nabla \times \mathbf{B} &= \left(-\frac{4\pi G}{c^2} + \mu_0 \frac{q^2}{m^2} \right) \mathbf{j} + c^{-2} \frac{\partial \mathbf{g}}{\partial t} \end{aligned} \quad (1)$$

where the following substitutions have been used :

$$\begin{aligned} \mathbf{g} &= \mathbf{g}_g + \frac{q}{m} \mathbf{g}_e, \quad \mathbf{B} = \mathbf{B}_g + \frac{q}{m} \mathbf{B}_e, \\ \rho_e &= \frac{q}{m} \rho, \quad \mathbf{j}_e = \frac{q}{m} \mathbf{j} \end{aligned} \quad (2)$$

In relations (1) and (2) (\mathbf{g}_e , \mathbf{B}_e) are the electromagnetic field components, (\mathbf{g}_g , \mathbf{B}_g) the gravitational field components, \mathbf{j}_e the electric field current density, \mathbf{j} the mass current density, ρ_e the electric charge density, ρ the mass density, q the

electric charge of a particle, m its mass, G is Newton's constant, c the speed of light, ϵ_0 the electric permittivity and μ_0 the magnetic permeability.

If the wave function $\psi = |\psi| e^{i\phi}$ is minimally coupled to the generalized vector potential $\mathbf{A} = \mathbf{A}_g + (q/m) \mathbf{A}_e$, the mass current density \mathbf{j} is

$$\begin{aligned} \mathbf{j} &= m \mathbf{j}_p = \frac{\hbar}{2i} \left(\psi^* \bar{\nabla} \psi - \psi \bar{\nabla} \psi^* \right) = \\ &= \frac{\hbar}{2i} \left(\psi^* \nabla \psi - \psi \nabla \psi^* \right) - \hbar g \mathbf{A} |\psi|^2 \end{aligned} \quad (3)$$

where \mathbf{j}_p is the probability current density, $\bar{\nabla} = \nabla - ig \mathbf{A}$ the covariant derivative, \mathbf{A}_g the gravitoelectromagnetic vector potential, \mathbf{A}_e the electromagnetic vector potential, g a coupling constant, q and m the charge and the mass of the superconducting pair, respectively. For $n = |\psi|^2 = \text{constant}$ and $g = m/\hbar$, the curl of relation (3) leads to the second London equation

$$\mathbf{B} = -\frac{1}{\rho} \nabla \times \mathbf{j}, \quad (4)$$

The first generalized London equation results from the third eq. (1) and eq. (4) using the standard procedure [5]

$$\mathbf{g} = \frac{1}{\rho} \frac{\partial \mathbf{j}}{\partial t} \quad (5)$$

Under these circumstances "matter" satisfying generalized London equations (4) and (5) defines a generalized superconductor. The electrono-positronic vacuum, type I and II superconductors are only few examples of generalized superconductors.

3. GENERALIZED MEISSNER EFFECT AND GRAVITATIONAL SHIELDING

Let us consider the last eq. (1) where we neglect the electromagnetic $\partial \mathbf{g}_e$ and gravitational $\partial \mathbf{g}_g$ displacement currents. Using the substitutions

$$\lambda_e = (m \mu_0^{-1} q^{-2} n^{-1})^{1/2} \text{ and } \lambda_g = (\mu_0^{-1} m^{-1} n^{-1})^{1/2} \text{ with}$$




Figure 1 Change of the lines of the gravitational field \mathbf{g} close to a cylinder coated with high temperature superconductors.

$\mu_0 g = 4\pi G c^2$, the curl of this equation with (4) becomes

$$\Delta \mathbf{B} - (\lambda_e^{-2} - \lambda_g^{-2}) \mathbf{B} = 0 \quad (6)$$

Equation (6) argues that for $\lambda_g > \lambda_e$, i.e. $q/m > (\mu_0 g / \mu_0 e)^{1/2} = 4.25 \cdot 10^{-11} \text{ C/m}$, a generalized Meissner effect might occur in generalized superconductors.

In this context the parameter $\lambda = \lambda_g \lambda_e (\lambda_g^2 - \lambda_e^2)^{-1/2}$ corresponds to the generalized penetration depth. It is easy to test such an effect may occur in conventional superconductors. Consequently, for $m = 9.1 \cdot 10^{-31} \text{ kg}$, $q = 1.6 \cdot 10^{-19} \text{ C}$, $n = 10^{28} \text{ m}^{-3}$, $\lambda_e \approx 10^{-8} \text{ m}$ and $\lambda_g \approx 10^{13} \text{ m}$ and (6) becomes

$$\Delta \mathbf{B} - \lambda_e^{-2} \mathbf{B} \approx 0 \quad (7)$$

For $\lambda_g < \lambda_e$, i.e. $q/m < (\mu_0 g / \mu_0 e)^{1/2} = 4.25 \cdot 10^{-11} \text{ C/m}$, eq. (6) indicates that in a generalized superconductor, the generalized Meissner effect is not present, in other words, the field \mathbf{B} induces in superconductor spatial oscillations with the wavelength $\lambda' = \lambda_g \lambda_e (\lambda_e^2 - \lambda_g^2)^{-1/2}$.

Now, following the same method used in [5] one finds for the component \mathbf{g} the equation

$$\Delta \mathbf{g} - (\lambda_e^{-2} - \lambda_g^{-2}) \mathbf{g} = 0 \quad (8)$$

Therefore, for $\lambda_g > \lambda_e$ the lines of the field \mathbf{g} are expelled from the superconductor, in other words, having in view its structure, it is possible to shield the gravitational field by means of an electromagnetic field. Such a shielding effect is present in Podkletnov's effect : a sample body placed over a spinning superconducting disk looses its weight [6].

In such a context, a superconducting cylinder placed in the gravitational field of the Earth will act

as a particle focusing device. Indeed, any particle moving along this cylinder with the speed v is subjected to a force (see figure 1)

$$\mathbf{g}_g = g_0 \exp\left(-\frac{R-x}{\lambda_e}\right), \quad (9)$$

solution of eq. (8).

Then for $(R-x)/\lambda_e \ll 1$ the particle's equation of motion becomes, on a first approximation

$$\ddot{y} + \omega^2 y = 0, \quad y = R - x, \quad \omega^2 = g_0 \lambda_e^{-1} \quad (10)$$

Under these circumstances, taking into account results from refs. [7] the focusing length is

$$f = 2\pi v \omega^{-1} \quad (11)$$

E.g., for a beam of ultracold neutrons with speed $v \sim 10^2 \text{ m/s}$ moving inside a cylinder coated with HTS, the focusing length calculated with (11) is $f \sim 1 \text{ cm}$.

4. CONCLUSIONS

The main conclusions of the present work are as follows :

- assuming the superposition principle as valid we get Maxwell type equations for the generalized field (electromagnetic plus linear gravitational);
- we get the generalized London equations;
- we establish the conditions necessary for the generalized Meissner effect to occur;
- we show that if an electromagnetic field is present, the gravitational shielding is possible to appear; in our opinion, this way Podkletnov's effect may be seen and understood [6];
- we propose a new experiment of focusing a beam of ultracold neutrons by a superconducting cylinder with its generatrix aligned parallel to the lines of the Earth's gravitational field.

REFERENCES

1. M. Visser, Phys. Rev. D 54 (1996) 5103; Phys. Rev. D 56 (1997) 936.
2. J. R. Gott, Phys. Rev. Lett. 66 (1991) 1126.
3. M. Agop and C. Gh. Buzea, Austr. Jour. Phys. (2000) in press.
4. H. Peng, Gen. Rel. Grav. 15 (1983) 725.
5. M. Agop, N. Rezlescu and G. Kalogirou, Nonlinear Phenomena in Materials Science, Plaisio Publishing House, Athenna, Greece, (1999).
6. E. Podkletnov, R. Nieminen, Physica C 203 (1992) 441.
7. C. Gh. Buzea, M. Agop and N. Rezlescu, J.J.A.P. 38 10 (1999) 5863.

Alternative explanation of “gravitational screening” experiments

Michael de Podesta *, Martyn Bull

Physics Department, Birkbeck College, Malet Street, London, WC1E 7HX, UK

Received 21 June 1995

Abstract

We show that the phenomena tentatively ascribed by Podkletnov and Nieminen [Physica C 203 (1992) 441] to gravitational screening may be easily understood in terms of a buoyancy correction to the authors’ weighing procedure.

Podkletnov and Nieminen [1] report that they have observed phenomena consistent with gravitational screening by a bulk superconductor. Their experiments consist of weighing a sample of SiO_2 above a liquid helium cryostat containing samples of YBCO superconductor. They report apparent reductions in sample weight from that observed at room temperature varying between 0.05% with no AC current in the superconductor to around 0.5% with AC currents flowing in the superconductor. Our assumptions concerning their apparatus are illustrated in Fig. 1 of their paper, reproduced here.

However, the authors have made no correction for the buoyancy of their SiO_2 samples in the air in which they are weighed. A simple calculation shows that the downward force which is counteracted by the weighing apparatus is given by

$$F = \rho_{\text{Sample}} V g - \rho_{\text{Gas}} V g \quad (1)$$

Assuming a density of 2100 kg m^{-3} for the SiO_2 and that the atmosphere within the cryostat is air, we have calculated the ratio of the density of air within the cryostat to the density of the SiO_2 sample. By assuming that the density of air is similar to that of

dry nitrogen [2], we predict an apparent weight loss of a similar magnitude to that observed (Fig. 2).

What this amounts to is an explanation of the phenomenon observed in terms of the *temperature* of the gas around the sample. In the steady state (no rotating fields) the sample is reported to show weight reduction of 0.05%, which indicates a sample temperature of the order of 150 K. When AC fields are in place, convection within the cryostat may reduce this temperature significantly, thus increasing the density of the gas. The convection would not need to exert a force on the sample directly but would merely lower the temperature of the air surrounding the sample. The convection could be driven by slight




Fig. 1. Schematic of the experimental apparatus described by the authors.

* Corresponding author.




Fig. 2. Prediction of the percentage reduction in apparent sample weight of an SiO_2 sample compared with its apparent weight at 290 K. The data assumes that the sample is weighed in dry nitrogen gas at atmospheric pressure, but the results are likely to be similar for dry air.

heating of the cryostat walls. This suggestion is directly falsifiable by measurement of the time variation of the temperature in the region of the sample. Note that the calculations reported in Fig. 2 are based on the density of dry nitrogen.

While the matter of a buoyancy correction in a weighing experiment is not of itself particularly important, the interpretation of the results that the authors have suggested is of great significance. We thus feel that it is important to ascertain that there are no alternative explanations for these observations before accepting the possibility of gravitational shielding.

References

- [1] E. Podkletnov and R. Nieminen, Physica C 203 (1992) 441.
- [2] R.T. Jacobsen and R.B. Stewart, J. Phys. Chem. Ref. Data. 2 (1973) 802.



ELSEVIER

Physica C 339 (2000) 120–128

PHYSICA C

www.elsevier.nl/locate/physc

Local gravitoelectromagnetic effects on a superconductor

M. Agop ^{a,*}, C. Gh. Buzea ^b, P. Nica ^a

^a Department of Physics, Technical University 'Gh. Asachi', Iasi, Romania

^b Superconductivity Research Laboratory, Institute of Technical Physics, Iasi, Romania

Received 16 February 2000; received in revised form 17 April 2000; accepted 1 May 2000

Abstract

Maxwell's and London's generalized equations, generalized Meissner effect and gravitational shielding in an electromagnetic field are obtained. In such a context, we show that a neutral particle beam in an infinitely thin superconducting cylinder placed in the Earth's gravitoelectric field is focused, and that the penetration depth increases with the pulsation of the electromagnetic field. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: Maxwell's and London's generalized equations; Generalized Meissner effect; Gravitational shielding in an electromagnetic field; Dispersion relations for superconductors

1. Introduction

Several experiments have been proposed to detect local gravitoelectromagnetic effect [1], but the practical difficulties are high. One of the reasons for studying the extreme sensitivity of superconducting devices may enable one to test gravitational effects. De Witt [2] and Anandan [3] studied gravitomagnetic effects on a superconducting circular ring. From the flux quantization condition, they calculated the magnetic field inside a superconducting ring, which is induced by an applied gravitomagnetic field. Papini [4] studied gravitomagnetic effects on a Josephson interferometer and designed an experiment to test the Machian principle. Cerdonio [5] discussed two kinds of detectors for the local gravitomagnetic effect of the Earth,

one of them based on the London momentum phenomenon. Recently, Podkletnov and Nieminen experimentally showed that a superconducting disk under electromagnetic field behaves as a gravitational shield [6].

In this paper, we propose an approach of studying gravitoelectromagnetic effects on superconductors. Assuming the gravitational and electromagnetic field superposition principle as valid, we obtain the generalized Maxwell's equation, and from here, on minimal coupling basis, the generalized London's equations. In such a context, we study the generalized Meissner effect and then, the gravitational shielding in an electromagnetic field. We build the solution of differential equations corresponding to the gravitational shielding effect for an infinitely thin superconducting cylinder placed in the Earth's gravitoelectric field. From this solution, we show that a neutral particle beam has cnoidal oscillation modes. By degeneration, we obtain harmonic, soliton, soliton packet modes. Since for the harmonic mode the beam is focused,

* Corresponding author. Address: St. Moara de Foc, no. 10, Bl. 406, sc. B, et. 8, ap. 35, Iasi, 6600, Romania.

E-mail address: pnica@ch.tuiasi.ro (M. Agop).

we suggest an experimental test. Using a dispersion relation, we show that the penetration depth increases with the pulsation of the electromagnetic field.

2. Generalized Maxwell equations

When four-dimensional space–time is split into space plus time ($3 + 1$), the electromagnetic field $F^{\alpha\beta}$ breaks up into two parts, the electric field \mathbf{g}_e and the magnetic field \mathbf{B}_e . These fields verify Maxwell's equations:

$$\nabla \cdot \mathbf{g}_e = \frac{\rho_e}{\epsilon_0}, \quad (1)$$

$$\nabla \cdot \mathbf{B}_e = 0, \quad (2)$$

$$\nabla \times \mathbf{g}_e = -\frac{\partial \mathbf{B}_e}{\partial t}, \quad (3)$$

$$\nabla \times \mathbf{B}_e = \mu_0 \mathbf{j}_e + c^{-2} \frac{\partial \mathbf{g}_e}{\partial t}, \quad (4)$$

where \mathbf{j}_e is the electric current density vector $\mathbf{j}_e = \rho_e \mathbf{v}$, ρ_e the electric charge density, μ_0 the vacuum magnetic permeability and c the speed of light in vacuum. Similarly, the general relativistic gravitational field ($g^{\alpha\beta}$) breaks into three parts [7]: (i) an electric like part, g^{00} , whose gradient for weak gravity is Newtonian acceleration \mathbf{g}_g ; (ii) a magnetic part, g^{0i} , whose curl for weak gravity is the gravitomagnetic field \mathbf{B}_g ; (iii) a spatial metric, g^{ij} , whose curvature tensor is the ‘curvature of space’ [8].

Although in astrophysics some theories of quasar and galactic nuclei rely on the gravitomagnetic field of a supermassive black hole for energy storage, power generation, jet formation and jet alignment, experiments have only probed g^{00} and g^{ij} with less than 0.1% accuracy. The gravitomagnetic potential g^{0i} is so weak in the Solar system that it has never been detected. According to the magnetic-type gravitation theory [9], in a two-body system, say, the moon–earth system, the body moving on the orbit produces a gravitomagnetic field \mathbf{B}_g at the center of the orbit. \mathbf{B}_g makes the rotation axis of the central body

precess. An experiment, which seems sufficiently feasible, is the gravitomagnetic-induced precession, relative to the star, of a superconducting gyroscope in an earth-orbiting satellite [10].

The relativistic weak-field approximation is constructed beginning from the axiom of relativity that the local coordinates, covering some patch on the space–time manifold which forms an open neighborhood about the origin, in terms of which the metric is Minkowskian up to first order in the coordinate over this patch. This construction may be carried out everywhere on the manifold [11].

In the weak gravitational field, we assume the metric in the form [12]

$$g_{\mu\nu} \approx \eta_{\mu\nu} + h_{\mu\nu} = \eta_{\mu\nu} + \bar{h}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \bar{h},$$

where $\eta_{\mu\nu} = (+, -1, -1, -1)$ is the usual Minkowski metric, $h_{\mu\nu}$ is to be treated as a small perturbation to $\eta_{\mu\nu}$, and

$$h_{\mu\nu} = \bar{h}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \bar{h}, \quad \bar{h} = h_\alpha^\alpha.$$

The contravariant form of the metric to first order is

$$g^{\mu\nu} \approx \eta^{\mu\nu} - h^{\mu\nu}, \quad h^{\mu\nu} = \eta^{\mu\alpha} \eta^{\nu\beta} h_{\alpha\beta},$$

and the determinant of the contravariant metric

$$g = \det(g_{\mu\nu}) \approx -1 - h, \quad (-g)^{1/2} = 1 + \frac{h}{2},$$

$$h = h_\alpha^\alpha.$$

Since all the covariant derivatives reduce to partial derivatives,

$$\nabla_\mu h_{\alpha\beta} \approx \partial_\mu h_{\alpha\beta},$$

the Riemann tensor, $R_{\mu\nu\lambda}^\sigma$, and the Ricci tensor $R_{\mu\nu}$ become

$$R_{\mu\nu}^\sigma = \partial_\nu \Gamma_{\mu\lambda}^\sigma - \dots$$

$$\approx \frac{1}{2} (\partial^\sigma \partial_\lambda h_{\mu\nu} + \partial_\mu \partial_\nu h_\lambda^\sigma - \partial_\mu \partial_\lambda h_\nu^\sigma - \partial_\nu \partial_\lambda h_{\mu\nu}),$$

and

$$R_{\mu\nu} = R_{\mu\lambda\nu}^\lambda \approx \frac{1}{2} (\partial^\lambda \partial_\lambda h_{\mu\nu} + \partial_\mu \partial_\nu h - \partial_\mu \partial_\lambda h_\nu^\lambda - \partial_\nu \partial_\lambda h_{\mu\nu}),$$

respectively.

We introduce the tensor

$$G^{\mu\nu\lambda} = \frac{1}{2} (\bar{h}^{\mu[\nu,\lambda]} + \eta^{\mu[\nu} \bar{h}_{\lambda]}^{\lambda\alpha}) \quad (5)$$

so that with the gauge condition,

$$\begin{aligned}\partial_\mu \left[g^{\mu\nu} (-g)^{1/2} \right] &= 0, \\ \partial_\mu h^{\mu\nu} \equiv \partial_\mu h_{,\mu}^{\mu\nu} &\cong \frac{1}{2} \partial^\nu h, \quad \bar{h}_{,\mu}^{\mu\nu} = 0,\end{aligned}\tag{6}$$

Einstein's equations reduce to

$$G_{,\lambda}^{\mu\nu\lambda} = -\frac{8\pi G}{c^4} T^{\mu\nu},\tag{7}$$

where G is Newton's constant and $T^{\mu\nu}$ is the energy-momentum tensor that is the source of the gravitational field. From Eqs. (5) and (6), we get

$$\begin{aligned}G^{\mu\nu\lambda} &= -G^{\mu\lambda\nu}, \quad G^{\mu\nu\lambda} + G^{\nu\lambda\mu} + G^{\lambda\mu\nu} = 0, \\ G^{\mu\nu\lambda} + G^{\nu\lambda\mu} + G^{\lambda\mu\nu} &= 0.\end{aligned}\tag{8}$$

As in the case of usual Maxwell's equations, since we do not require a completely covariant formulation, it is convenient to break the tensors into scalar, vector and tensor parts according to the transformation properties under pure rotations.

We introduce the vectors

$$\begin{aligned}\mathbf{g}_g &= (g_g^1, g_g^2, g_g^3), \quad g_g^i = G^{00i} \quad (i = 1, 2, 3), \\ \mathbf{A}_g &= (A_g^1, A_g^2, A_g^3), \quad A_g^i = \frac{1}{4} h^{0i} \\ \mathbf{B}_g &= (B_g^1, B_g^2, B_g^3), \quad B_g^1 = G^{023}, \quad B_g^2 = G^{031}, \\ B_g^3 &= G^{012}.\end{aligned}\tag{9}$$

Substituting Eq. (9) into Eq. (5), we find

$$G^{0ij} = A_g^{i,j} - A_g^{j,i}, \quad \mathbf{B}_g = \nabla \times \mathbf{A}_g.\tag{10}$$

Using Eqs. (9) and (10), from Eqs. (7) and (8), we get the Maxwell-type gravitational equations for the weak gravitational field:

$$\nabla \cdot \mathbf{g}_g = -4\pi G\rho,\tag{11}$$

$$\nabla \cdot \mathbf{B}_g = 0,\tag{12}$$

$$\nabla \times \mathbf{g}_g = -\frac{\partial \mathbf{B}_g}{\partial t},\tag{13}$$

$$\nabla \times \mathbf{B}_g = -\frac{4\pi G}{c^2} \mathbf{j} + \frac{1}{c^2} \frac{\partial \mathbf{g}_g}{\partial t},\tag{14}$$

where \mathbf{j} is the mass current density vector, $\mathbf{j} = \rho \mathbf{v}$ and ρ , the mass density. This form of the decom-

posed equations is valid in any coordinate system. They are covariant under transformations of the spatial coordinates $\bar{x}^i = \bar{x}^i(x^1, x^2, x^3)$, but it should be stressed that, in general, once a particular coordinates system is chosen, the equations cannot be taken simply over another system of coordinates by an arbitrary coordinate transformation of the type $\bar{x}^x = \bar{x}^x(x^0, x^1, x^2, x^3)$. It is necessary to return to the original term of the decomposed equations and substitute the quantities appropriate to the new system of coordinates.

Quantitatively, the linearizing approximation is extremely well justified on the surface of any member of our Solar system. Even on the surface of a white dwarf, the deviation of g_{00} from unity does not exceed a value of the order of 10^{-5} . Therefore, we can postulate, ab initio, the Maxwell-type gravitational equations for linear gravitation.

Now, if in a space-time manifold, Eqs. (1)–(4) as well as Eqs. (11)–(14) are valid, based on the superposition principle, we can introduce the generalized fields:

$$\mathbf{g} = \mathbf{g}_g + \frac{q}{m} \mathbf{g}_e, \quad \mathbf{B} = \mathbf{B}_g + \frac{q}{m} \mathbf{B}_e.\tag{15}$$

These fields verify the generalized Maxwell's equations

$$\nabla \cdot \mathbf{g} = \left(-4\pi G + \frac{1}{\epsilon_0 m^2} q^2 \right) \rho,\tag{16}$$

$$\nabla \cdot \mathbf{B} = 0,\tag{17}$$

$$\nabla \times \mathbf{g} = -\frac{\partial \mathbf{B}}{\partial t},\tag{18}$$

$$\nabla \times \mathbf{B} = \left(-\frac{4\pi G}{c^2} + \mu_0 \frac{q^2}{m^2} \right) \mathbf{j} + \frac{1}{c^2} \frac{\partial \mathbf{g}}{\partial t},\tag{19}$$

where we accounted for the following relations:

$$\rho_e = \frac{q}{m} \rho, \quad j_e = \frac{q}{m} j.\tag{20}$$

As an example, Eq. (16) is obtained adding Eq. (1) multiplied by q/m to Eq. (11) and taking into account the first relation in Eqs. (15) and (20), respectively.

3. Generalized London equations

If the wave function $\psi = |\psi|e^{i\varphi}$ is minimally coupled to the generalized vector potential

$$\mathbf{A} = \mathbf{A}_g + \frac{q}{m}\mathbf{A}_e, \quad (21)$$

the mass current density \mathbf{j} is

$$\begin{aligned} \mathbf{j} &= m\mathbf{j}_p = \frac{\hbar}{2i} (\psi^* \bar{\nabla} \psi - \psi \bar{\nabla} \psi^*) \\ &= \frac{\hbar}{2i} (\psi^* \nabla \psi - \psi \nabla \psi^*) - \hbar g \mathbf{A} |\psi|^2, \end{aligned} \quad (22)$$

where \mathbf{j}_p is the probability current density, $\bar{\nabla} = \nabla - ig\mathbf{A}$ the covariant derivative, \mathbf{A}_g the gravito-electromagnetic vector potential, \mathbf{A}_e the electromagnetic vector potential, g a coupling constant, q and m the effective charge and the effective mass of the superconducting pair, respectively.

For $n = |\psi|^2 = \text{constant}$, relation (22) becomes

$$\mathbf{j} = \hbar |\psi|^2 (\nabla \varphi - g\mathbf{A}), \quad (23)$$

or with $g = m/\hbar$ and taking the curl,

$$\mathbf{B} = -\frac{1}{\rho} \nabla \times \mathbf{j}, \quad (24)$$

where the following notations have been used:

$$\begin{aligned} \mathbf{B} &= \nabla \times \mathbf{A}, & \mathbf{B}_g &= \nabla \times \mathbf{A}_g, \\ \mathbf{B}_e &= \nabla \times \mathbf{A}_e, & \rho &= m|\psi|^2, \end{aligned} \quad (25)$$

We refer to Eq. (24) as the second generalized London equation. This relation embodies some particular interesting cases. They read as follows:

(i) For $\mathbf{B}_g = 0$ and

$$\lambda_e = \left(\frac{m}{\mu_{0e} q^2 n} \right)^{1/2}, \quad (26)$$

Eq. (24) reduces to the second London equation [13]

$$\mathbf{B}_e = -\mu_{0e} \lambda_e^2 \nabla \times \mathbf{j}_e, \quad (27)$$

where λ_e is the electromagnetic penetration depth.

(ii) For $\mathbf{B}_e = 0$ and

$$\lambda_g = \left(\frac{1}{\mu_{0g} nm} \right)^{1/2}, \quad \mu_{0g} = \frac{4\pi G}{c^2}, \quad (28)$$

Eq. (24) reduces to the second gravitational London equation

$$\mathbf{B}_g = -\mu_{0g} \lambda_g^2 \nabla \times \mathbf{j}, \quad (29)$$

where μ_{0g} defines the “gravitational permeability of vacuum” and λ_g the “gravitational penetration depth” [14].

Let us write Eq. (18) in the form

$$\nabla \times \left(\mathbf{g} + \frac{\partial \mathbf{A}}{\partial t} \right) = 0. \quad (30)$$

Integrating results in

$$\mathbf{g} + \frac{\partial \mathbf{A}}{\partial t} = 0. \quad (31)$$

Since from Eq. (24), one gets

$$\mathbf{A} = -\frac{1}{\rho} \mathbf{j}, \quad (32)$$

substituting the time derivative from this relation in Eq. (31), the first generalized London equation results

$$\mathbf{g} = \frac{1}{\rho} \frac{\partial \mathbf{j}}{\partial t}. \quad (33)$$

This relation embodies some particular interesting cases. They read as follows:

(i) For $\mathbf{g}_g = 0$ and using substitutions (26), relation (33) reduces to the first London equation [13]

$$\mathbf{g}_e = \mu_{0e} \lambda_e^2 \frac{\partial \mathbf{j}_e}{\partial t}. \quad (34)$$

(ii) For $\mathbf{g}_e = 0$ and using substitutions (28), relation (33) reduces to the first gravitational London equation [14]

$$\mathbf{g}_g = \mu_{0g} \lambda_g^2 \frac{\partial \mathbf{j}}{\partial t}. \quad (35)$$

Under these circumstances, “matter” satisfying generalized London equations (24) and (33) defines a generalized superconductor. The electrono-positronic vacuum, type I and II superconductors

are only few examples of generalized superconductors.

4. Generalized Meissner effect

Let us consider Eq. (19) wherein we neglect the electromagnetic $\partial_t \mathbf{g}_e$ and gravitational $\partial_t \mathbf{g}_g$ displacement currents. With substitutions (26) and (28) and applying the curl it becomes

$$\Delta \mathbf{B} - \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) \mathbf{B} = 0, \quad (36)$$

where we took into account Eq. (17). Eq. (36) argues that for $\lambda_g > \lambda_e$, i.e. $q/m > (\mu_{0g}/\mu_{0e})^{1/2}$, a generalized Meissner effect might occur in generalized superconductors. In this context the parameter $\lambda = \lambda_g \lambda_e (\lambda_g^2 - \lambda_e^2)^{-1/2}$ corresponds to the generalized penetration depth.

It is easy to test that such an effect may occur in conventional superconductors. Consequently, for $\lambda_e \approx 10^{-8}$ m [13] and $\lambda_g \approx 10^{13}$ m [14] and Eq. (36) becomes

$$\Delta \mathbf{B} - \frac{1}{\lambda_e^2} \mathbf{B} \approx 0. \quad (37)$$

For $\lambda_g < \lambda_e$, i.e. $q/m < (\mu_{0g}/\mu_{0e})^{1/2}$, Eq. (36) indicates that in a generalized superconductor, the generalized Meissner effect is not present, in other words, the field \mathbf{B} induces, in superconductor, spatial oscillations with the wavelength $\lambda' = \lambda_g \lambda_e (\lambda_e^2 - \lambda_g^2)^{-1/2}$.

Eq. (36) embodies some interesting cases. Thus,

(i) For $\mathbf{B}_g = 0$, Eq. (36) becomes Eq. [13]

$$\Delta \mathbf{B}_e - \frac{1}{\lambda_e^2} \mathbf{B}_e = 0 \quad (38)$$

namely, the usual Meissner effect;

(ii) For $\mathbf{B}_e = 0$, Eq. (36) becomes

$$\Delta \mathbf{B}_g + \frac{1}{\lambda_g^2} \mathbf{B}_g = 0, \quad (39)$$

namely, the gravitational Meissner effect is absent [15]. These results are different compared with those obtained by Peng in Ref. [16]. Thus, from Ref. [16] and with notations (26) and (28) we obtain

$$\Delta \mathbf{B} - \left(\frac{1}{\lambda_e^2} + \frac{1}{\lambda_g^2} \right) \mathbf{B} = 0.$$

From here, for the gravitational field, it gives the gravitational Meissner effect:

$$\Delta \mathbf{B} - \frac{1}{\lambda_g^2} \mathbf{B} = 0.$$

This relation is not in agreement with Refs. [15,17,18]. In the references, mentioned we showed that the gravitational Meissner effect is not present in a pure gravitomagnetic field. The absence of the effect was interpreted as a self-structuring of the physical space-time in a boundless crystal named “world crystal”.

5. Generalized shielding

Substituting Eq. (24) into Eq. (19) and taking into account notations (26) and (28), it gives

$$\Delta \mathbf{j} - \nabla(\nabla \cdot \mathbf{j}) - \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) \mathbf{j} = 0. \quad (40)$$

Since the conservation law

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} = 0 \quad (41)$$

with $\rho = \text{constant}$ implies

$$\nabla \cdot \mathbf{j} = 0, \quad (42)$$

Eq. (40) becomes

$$\Delta \mathbf{j} - \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) \mathbf{j} = 0. \quad (43)$$

From here, differentiating with t and taking into account Eq. (31), we get

$$\Delta \mathbf{g} - \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) \mathbf{g} = 0. \quad (44)$$

Therefore, for $\lambda_g > \lambda_e$, the lines of the field \mathbf{g} are expelled from the superconductor; in other words, having its structure in view, it is possible to shield

the gravitational field by means of an electromagnetic field. Podkletnov and Nieminen experimentally verified this effect in Ref. [6].

6. Oscillation modes in thin cylindrical superconductors

For a superconductor in the Earth's gravitoelectric field, Eq. (44) takes the form

$$\Delta g_g - \frac{1}{\lambda_e^2} g_g = 0. \quad (45)$$

In the case of an infinitely thin superconductor cylinder of radius R (Fig. 1), the solution of Eq. (45) (see the method from Ref. [19]), i.e.

$$g_g(x) = g_0 e^{-(R-x)/\lambda_e}, \quad x \leq R, \quad (46)$$

where g_0 is the gravitoelectric field on the cylinder's surface, (from Eq. (46) $x = R$ corresponds to the cylinder's surface), indicates that on a particle moving with speed v , acts the specific repulsive force:

$$\frac{d^2x}{dt^2} = -g_g(x). \quad (47)$$

Consequently, a beam placed near the superconductor cylinder axis is focused. Let us consider the dimensionless differential equation (47) assuming $y = R - x/\lambda_e \ll 1$, i.e.,




Fig. 1. Thin superconducting cylinder in a gravitoelectric field.

$$\frac{d^2y}{d\tau^2} = \frac{g_0}{\lambda_e \omega^2} \left(\frac{y^2}{2} - y + 1 \right), \quad \tau = \omega t. \quad (48)$$

Integrating twice, results in

$$\left(\frac{dy}{d\tau} \right)^2 = \frac{g_0}{3\lambda_e \omega^2} P(y), \quad (49)$$

and, respectively,

$$\int \frac{du}{[(1-u^2)(1-k^2u^2)]} = \gamma \int d\tau, \quad (50)$$

where we used the substitutions,

$$\begin{aligned} y &= e_1 + (e_2 - e_1)u^2, & A &= \frac{e_2 - e_1}{2}, \\ k^2 &= \frac{e_2 - e_1}{e_3 - e_1}, & \gamma^2 &= \frac{g_0(e_3 - e_1)}{12\lambda_e \omega^2}, \end{aligned} \quad (51)$$

$$\begin{aligned} P(y) &= y^3 - 3y^2 + 6y + 6C \\ &= (y - e_1)(y - e_2)(y - e_3), \\ C &= \text{const.} \end{aligned} \quad (52)$$

with $e_1 < e_2 < e_3$, the real roots of the polynomial $P(y)$. Finally, using the method given in Ref. [20], the solution becomes

$$y = e_2 - 2Acn^2[\gamma(\tau - \tau_0); k], \quad (53)$$

where cn is Jacobi's elliptic function with modulus k [20]. Thus, the oscillation modes of a particle beam, which travels parallel and near the cylinder axis, are cnoidal (Fig. 2) and have the following particularities:

(i) The period

$$T = 2 \left(\frac{6\lambda_e}{g_0 A} \right)^{1/2} K(k)k. \quad (54)$$

This result is obtained by integrating Eq. (39) with substitution $u = \sin \varphi$ in the limits 0 and $\pi/2$ and taking into account that $cn^2(\theta + 2K) = cn^2(\theta)$, i.e.,

$$\begin{aligned} 2K(k) &= 2 \int_0^{\pi/2} \frac{d\varphi}{(1 - k^2 \sin^2 \varphi)^{1/2}} \\ &= \frac{1}{k} \left(\frac{g_0 A}{6\lambda_e} \right)^{1/2} \int_0^T dt. \end{aligned} \quad (55)$$




Fig. 2. Cnoidal oscillation modes.

(ii) The mean elongation

$$\bar{y} = -\frac{2A}{k^2} \frac{E(k)}{K(k)} + e_3, \quad (56)$$

where

$$E(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \varphi)^{1/2} d\varphi. \quad (57)$$

This result is obtained using the averaging relation

$$\bar{y} = \frac{1}{T} \int_0^T y(t) dt. \quad (58)$$

Under these conditions, the focusing distance for a particle beam of speed v has the form

$$f = \frac{\lambda}{4} = \frac{1}{2} \left(\frac{6\lambda_e}{g_0 A} \right)^{1/2} v K(k) k. \quad (59)$$

By specifying the modulus k , we distinguish the following degenerations:

(i) For $k = 0$, Eq. (53) reduces to the harmonic mode

$$y = e_1 - A \cos \omega t \quad (60)$$

relative to $\bar{y} = e_1$ with pulsation

$$\omega = \left(\frac{g_0}{\lambda_e} \right)^{1/2}. \quad (61)$$

Then, the focusing distance becomes

$$f = \frac{v\pi}{2} \left(\frac{\lambda_e}{g_0} \right)^{1/2}. \quad (62)$$

(ii) For $k \rightarrow 0$, Eq. (53) reduces to the harmonic modes packet

$$y \rightarrow e_1 - A \left[\cos \omega t + \frac{1}{8} k^2 \cos 2\omega t + O(k^4) \right] \quad (63)$$

relative to $\bar{y} \rightarrow e_1$, with pulsation

$$\frac{2g_0 A}{3\lambda_e \omega^2} \rightarrow k^2 \left(1 + \frac{k^2}{2} \right) + O(k^6). \quad (64)$$

Then, the focusing distance becomes

$$f \rightarrow \frac{v\pi}{2} \left(\frac{3\lambda_e}{2g_0 A} \right)^{1/2} k \left(1 + \frac{k^2}{4} \right) + O(k^5). \quad (65)$$

(iii) For $k = 1$, Eq. (53) reduces to the ‘solitonic’ mode

$$y = e_3 - \frac{a}{\operatorname{ch}^2 \left[\left(\frac{g_0 a}{12\lambda_e} \right)^{1/2} t \right]} \quad (66)$$

relatively to $\bar{y} = e_3$, with amplitude $a = e_2 - e_1$ and ‘width’ $D = (12\lambda_e/g_0 a)^{1/2}$. Since the pulsation is infinite, see relation (54) for $K(1) = \infty$, the focusing distance is infinite.

(iv) For $k \rightarrow 1$, Eq. (53) reduces to a ‘solitonic’ modes packet relative to $\bar{y} = e_3$, with pulsation

$$\omega \rightarrow \frac{2\pi}{\left(\frac{6\lambda_e}{g_0 A} \right)^{1/2} \ln \left(\frac{16}{1-k^2} \right)}. \quad (67)$$

Then, the focusing distance becomes

$$f \rightarrow \frac{v\pi}{4} \left(\frac{6\lambda_e}{g_0 A} \right)^{1/2} \ln \left(\frac{16}{1-k^2} \right). \quad (68)$$

Evaluating the focusing distance in the Earth’s gravitoelectric field, corresponding to the harmonic mode, we find

$$f \approx 5 \times 10^{-5} v(s). \quad (69)$$

Since such cylindrical superconductors (supertrons) are already built [21,25], we suggest the use

of thermal neutrons (10^3 m/s) to be focused in a 5 cm cylinder, according to Eq. (69).

7. The dispersion relations: the dependence of the penetration depth on the pulsation of the electromagnetic field

Let us suppose that the current density \mathbf{j} is determined by the contribution of the normal and superconducting electrons, i.e.,

$$\mathbf{j} = \mathbf{j}_N + \mathbf{j}_S, \quad (70)$$

where

$$\mathbf{j}_N = -\sigma_g \mathbf{g}_g \mathbf{j}_{Ne} = -\sigma_e \mathbf{g}_e, \quad (71)$$

and \mathbf{j}_S satisfy London equations (24) and (33). In Eq. (71), σ_g is the gravitational conductivity (for details see Refs. [22,23]), and σ_e is the electrical conductivity.

By applying the curl to Eq. (19), and taking into account Eqs. (15)–(18) and (20), (24), (26), (28) and restriction

$$\mu_0 g = \mu_0 e \sigma_e, \quad (72)$$

we obtain the propagation equation

$$\mathbf{B} = \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) \mathbf{B} + \mu_0 g \sigma_g \frac{\partial \mathbf{B}}{\partial t}. \quad (73)$$

From here, using $\mathbf{B} = \mathbf{B}_0 e^{i(kr-\omega t)}$, we find the dispersion equation

$$k^2 = \mu_0 g \sigma_g \omega i - \left(\frac{1}{\lambda_e^2} - \frac{1}{\lambda_g^2} \right) + \frac{\omega^2}{c^2}. \quad (74)$$

This relation embodies some interesting particular cases. They read as follows:

(i) For $B_g = 0$, Eq. (74) reduces to the dispersion relation of the electromagnetic waves in superconductor two-fluid model [24], i.e.,

$$k^2 = \mu_0 e \sigma_e \omega i - \frac{1}{\lambda_e^2} + \frac{\omega^2}{c^2}. \quad (75)$$

(ii) For $B_e = 0$, Eq. (74) reduces to a dispersion relation corresponding to a gravitational perturbation, i.e.,

$$k^2 = \mu_0 g \sigma_g \omega i + \frac{1}{\lambda_g^2} + \frac{\omega^2}{c^2}. \quad (76)$$

Taking into account Eq. (74) and Ref. [23], the refraction index has the form

$$N^2 = \frac{k^2(\omega)}{k^2(0)} = 1 - \left(\frac{\omega \lambda}{c} \right)^2 - \mu_0 g \sigma_g \omega^2 i, \quad (77)$$

where

$$\lambda^2 = \lambda_e^2 \left(1 - \frac{\lambda_e^2}{\lambda_g^2} \right)^{-1}. \quad (78)$$

In relation (77), N^2 is complex; the real part, i.e.,

$$\text{Re} N^2 = 1 - \left(\frac{\omega \lambda}{c} \right)^2 \quad (79)$$

corresponds to the transparency, and the imaginary part

$$\text{Im} N^2 = -\mu_0 g \sigma_g \omega \lambda^2 \quad (80)$$

to the absorption. Since for a superconductor $\lambda \approx \lambda_e$, the functional dependence $\lambda_e = \lambda_e(\omega)$ is obtained by extending relation (26) in the form

$$\lambda_e(\omega) = \left(\frac{\varepsilon_0 |N|^2 mc^2}{q^2 n} \right)^{1/2} = \lambda_e(0) |N| \quad (81)$$

with

$$\lambda_e(0) = \left(\frac{\varepsilon_0 mc^2}{q^2 n} \right)^{1/2}. \quad (82)$$

This results in

$$\begin{aligned} \lambda_e(\omega) = \lambda_e(0) & \left[\left(1 - \left(\frac{\omega \lambda_e(0)}{c} \right)^2 \right)^2 \right. \\ & \left. + \left(\mu_0 e \sigma_e \omega \lambda_e^2(0) \right)^2 \right], \end{aligned} \quad (83)$$

which shows that the penetration depth increases with increasing pulsations of the electromagnetic field. This result is in agreement with Podkletnov's experiment [6].

8. Conclusions

The main conclusions of the present work are as follows:

1. assuming that the superposition principle is valid, we get Maxwell type equations for the generalized field (electromagnetic plus linear gravitational);
2. we get the generalized London equations;
3. we establish the conditions necessary for the generalized Meissner effect to occur;
4. the gravitational shielding in an electromagnetic field is analyzed;
5. we build the solution of differential equation corresponding to the gravitational shielding effect for an infinitely thin superconductor cylinder placed in the Earth's gravitoelectric field;
6. using this solution, we show that the oscillation modes of a neutral particle beam is cnoidal;
7. by the degeneration of cnoidal modes, we get harmonic, soliton, and solitons packet modes;
8. for the harmonic mode, we calculate the focusing length; in these conditions, we suggest an experimental test;
9. using a dispersion relation, we show that the penetration depth increases with the pulsation of the electromagnetic field.

Acknowledgements

We wish to thank Prof. S.W. Hawking, E. Podkletnov, and H. Matsuzawa for professional observations. We also thank the referees for their comments.

References

- [1] H. Peng, C. Qin, Phys. Lett. 103A (1984) 197.
- [2] B.S. De Witt, Phys. Rev. Lett. 16 (1966) 1092.
- [3] J. Anandan, in: M. Greensberger (Ed.), *New Techniques and Ideas in Quantum Measurement Theory*, New York Academy of Sciences, New York, 1986.
- [4] G. Papini, Phys. Lett. 24A (1967) 32.
- [5] M. Cerdonio, Gen. Rel. Grav. 20 (1988) 83.
- [6] E. Podkletnov, R. Nieminen, Physica C 203 (1992) 441.
- [7] V.B. Braginsky, A.G. Polnarev, K.S. Thorne, Phys. Rev. Lett. 53 (1984) 863.
- [8] P. Meystre, M.O. Scully, *Quantum optics, Experimental Gravitation and Measurement Theory*, Plenum Press, New York, 1983.
- [9] H. Peng, Gen. Rel. Grav. 15 (1983) 725.
- [10] B. Bertotti, *Experimental gravitation*, Academic Press, New York, 1984.
- [11] M.S. Madsen, Class. Quant. Grav. 87 (1990) 4.
- [12] C. Ciubotariu, I. Gottlieb, I. Simaciu, V. Griga, C.I. Ciubotariu, S. Croitoru, Tensor N.S. 52 (1993) 138.
- [13] G. Burns, *High Temperature Superconductivity – an Introduction*, Academic Press, San Diego, 1992.
- [14] M. Agop, N. Rezlescu, G. Kalogirou, *Nonlinear Phenomena in Materials Science*, Graphics Art Publishing House, Athenna, Greece, 1999.
- [15] C. Ciubotariu, M. Agop, Gen. Rel. Grav. 28 (1996) 405.
- [16] H. Peng, Gen. Rel. Grav. 22 (1990) 609.
- [17] M. Agop, B. Ciobanu, C.Gh. Buzea, N. Rezlescu, T. Horgos, C. Stan, Il Nuovo Cimento 113B (1998) 17.
- [18] M. Agop, C.Gh. Buzea, V. Griga, C. Ciubotariu, C. Buzea, C. Stan, D. Tatomir, Austr. J. Phys. 49 (1996) 1063.
- [19] C.P. Poole, H.A. Farach, R.J. Creswick, *Superconductivity*, Academic Press, New York, 1995.
- [20] F. Bowman, *Introduction to Elliptic Function with Applications*, English University Press, London, 1955.
- [21] C. Buzea, M. Agop, N. Rezlescu, Jpn. J. Appl. Phys. 38 (1999) 5863.
- [22] C. Ciubotariu, Phys. Lett. A 158 (1991) 27.
- [23] I. Oprea, M. Agop, Studia Geoph. Et Geod. 42 (1998) 431.
- [24] R.D. Deusch, *Introduction in Low Temperature Physics*, Academic Press, Bucharest, 1970.
- [25] H. Matsuzawa, J. Appl. Phys. 74 (1993) 1111.

Does a superconductor shield gravity?

C.S. Unnikrishnan¹

Gravitation Experiments Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay, 400 005, India

Received 22 March 1996

Abstract

Recently there were some experimental observations which were interpreted as due to a shielding of the gravitational interaction by a superconducting disc in a static configuration as well as when set in rotation. We examine the experiments in detail and point out some difficulties which should be eliminated before reliable results can be claimed. The data from these experiments provide an internal check on the correctness of the hypothesis and we argue that the observed results are inconsistent with the hypothesis of shielding and therefore they are not due to shielding of the Earth's gravity. Our preliminary experiments in the static case do not show any evidence for the reported shielding.

Keywords: Superconductivity; Meissner effect; Levitation; Gravitational shielding

Recently there were experimental observations by Podkletnov et al. of an apparent reduction in weight of a sample hung above a rotating superconducting disc, for certain ranges in the rotation rate [1,2]. These results were interpreted as due to the partial shielding of the gravitational field by the rotating superconducting disc. Shielding of gravity is an important issue and so far there is no experimental evidence for such an effect, and therefore it is important to carefully study the results and data in Refs. [1] and [2]. Of course, there is no theoretical framework at present which can give rise to such a shielding and any such effect can only be due to some effect which is beyond the general theory of relativity, possibly from quantum effects and so on. The problem even with such a view is that the measured effect is very large, amounting to more than 1% shielding of the local gravitational field. If the coupling which gives rise to such a shielding depends on fundamental constants like G , c , and \hbar , then such

a large shielding is certainly very unnatural for situations involving small gravitational fields and small velocities. In fact, since one is looking at the Newtonian field (the electric part of the full gravitational interaction, with the Newtonian potential ϕ given by $1 + 2\phi = g_{00}$, where g_{00} is the time-time component of the metric), the expectation would be an enhancement of the weight since the superconductor is in rotation and the rotational energy should contribute to the effective mass and effective gravity. But this effect also is very small, being of the order of $GI\omega^2m/c^2r$, where I is the moment of inertia of the disc and r the typical separation between the disc and test mass m . Such effects amount to corrections to the weight of less than 10^{-23} !

A fundamental difference between electromagnetism and gravitation is the fact that in gravity there is only one type of charge as far as we know and therefore many phenomena in electromagnetism which involve two types of charges are absent in gravitation. That is why there is no shielding of the Newtonian

¹ E-mail: unni@tifrvax.tifr.res.in.

gravitational field. With Einstein's theory the situation regarding shielding is not different; in fact, in general relativity typically the potential is enhanced due to other interactions since they also contribute to gravity with the same attractive sign. But there is the possibility that some part of the potential is diminished in the presence of an object like a superconductor where there are macroscopic quantum effects. An analogue of the Meissner effect could be thought of in which the gravitomagnetic potential (from the g_{0i} part of the metric) is shielded by the superconducting electrons. There is a difficulty, however, since the corresponding penetration depths are very large and therefore it is equivalent to no shielding at all in any imaginable situation. In any case such a shielding or enhancement, if any, cannot affect the weight of a test mass since the gravitomagnetic effects are velocity dependent and do not exert a force on any static object.

The reasons for starting with a pessimistic view on the results presented in Refs. [1] and [2] are three-fold:

- (1) the theoretical framework of the phenomenon of classical gravity is now well tested and general relativity has emerged as more or less the correct theory describing all known gravitational phenomena.
- (2) In a recent experiment measuring the levitation force on magnets near a superconductor in static situations we have not seen any anomalous weight reduction even though our sensitivity was about 50 times higher than what was required to see the effect seen by Podkletnov in the static situation, and
- (3) new results obtained by Podkletnov [2] in an improved experimental situation are internally inconsistent as we will argue in this paper. All these point to the need for better experiments with several control experiments to compare with.

Before we discuss these results, it is appropriate to highlight the essential details of the experiment, as described in Refs. [1] and [2]. A large superconducting toroidal disc (YBCO, cooled in He vapour around 60 K) was levitated by the Meissner effect in an oscillating magnetic field and then rotated with rotating magnetic fields. The weight of a SiO_2 sample (and several other samples in the later experiments) hung from a balance was measured as a function of the rotation rate. They report fluctuating weight readings, including both increase and decrease for some ranges of rotation rates and there are reproducible values of

the reduction in weight for a certain range of rotation rate. Also, there is some weight reduction noticed when the disc is not rotating at all. The reduction is around 0.05% for the static situation and it is 0.1% when the disc is rotating. The fluctuating values are also very large, a few percent.

In a more recent version of the experiment [2], the rotating superconducting disc was housed in a cryostat to avoid air currents during rotation. In all these experiments the weight reduction of the sample amounting to even 2% was seen. Also, it was observed that the shielding and weight reduction was independent of the distance above the rotating disc. We will argue later that this fact is inconsistent with any natural model of shielding of the Earth's gravity.

Before we discuss the internal inconsistency of Podkletnov's data, we would like to make some comments about the possibilities suggested by Podkletnov for the shielding of gravity. Any mechanism which invokes additional magnetic lines inside the superconductor affecting the gravitational interaction itself will not work since the energy stored in flux lines will be proportional to B^2 , and it is positive. This can only enhance the gravitational effect, though such enhancements are minuscule. Classically, the gravitational constant itself cannot be modified by the stored magnetic field, whichever way it is distributed between the grain boundaries of the superconductor. So, even if the effect is genuine it has to come from a new interaction and there are some attempts to invoke such a mechanism [3]. A more probable cause would be unidentified electromagnetic interaction (this could be from pinned flux lines rotating with the disc, for example) or more mundanely, but more probably, air currents and drags due to temperature gradients and rotation affecting the sample on the balance. There is good reason to believe that this might be a factor in at least some of the experiments reported in Refs. [1] and [2], since it was seen that the shielding is better when the sample is kept such that its projected area above the rotating disc is maximum, very characteristic of levitation by air currents.

Now we will elaborate on an important internal consistency check available in Podkletnov's data, which rules out shielding of Earth's gravity by the rotating superconducting disk. Podkletnov has done experiments in which the rotating disc was enclosed in a cryostat to eliminate air currents affecting the measurements.




Fig. 1. Diagram showing the decrease of volume and mass of the Earth which can be shielded in principle by the disc (D), as a function of the distance of the sample (S) above the disc. See text for details.

But, this does not eliminate convection currents outside the cryostat and in fact the experimenters state that an annular cylindrical flow of air is clearly visible above the cryostat (the disc used also is annular). Fortunately, Podkletnov has measured the weight of the sample above the disc for various distances from the disc and one of these measurements is done in a room above the room in which the disc was rotating. As mentioned earlier the shielding was found to be independent of the distance of the sample above the rotating disc. The observation of weight reduction when the sample is kept in a separate room above the room where the disc was rotating helps in quenching doubts about the role of air currents in the observed weight reduction. But the observed distance independence contradicts the shielding hypothesis and it provides the crucial test of the hypothesis of shielding of the earth's gravity. It is known from Newton's time that the Earth's gravity (or gravity of any spherical

body) when added as a vectorial interaction acts as if the whole mass is concentrated at the centre. This of course does not mean that the source is concentrated at the centre (even if the centre is hollow, this result remains true). For any arbitrary mass distribution the correct total gravitational field is given by the vectorial integral of the field from elementary masses distributed throughout the large mass. Therefore, in any hypothesis of shielding what is important is the mass of the Earth as seen by the sample being weighed. If the vector connecting the sample and a mass element passes through the disc (like the arrow numbered "1" in Fig. 1), there is a possibility of shielding of gravity of that particular mass element, but if the disc does not intersect the vector (arrow numbered "2") then the sample will be pulled by gravity of that mass element. Referring to Fig. 1, it is clear that only the gravitational force from the region inside the intersection of the cone with the spherical earth would be shielded in the configuration shown and that the rest of the Earth's gravity will still pull down on the sample. The figure also shows the fact that the shielded region diminishes in volume as the distance (h) between the sample and the disc is increased. The region which could be shielded has the approximate shape of a truncated cone and it is easy to estimate that the reduction in shielding as the height of the sample above the disc is increased. Since the radius of the earth (R) is very large compared to the diameter (d) of the disc and the height h , the mass which is effectively shielded can be approximated as that of the cone and is given by $M_c = (2\pi\rho_c/3)(d^2R^3)/h^2$, where ρ_c is the average density of the conical region. The mass of the earth is given by $M = (4\pi\rho R^3)/3$. The ratio of the masses is $M_c/M = \rho_c d^2/(2\rho h^2)$. To find the ratio of the gravitational contributions from the whole earth and the conical region, we need to include the angular factors and integrate the contributions from elementary masses in these regions with the density as a function of radius. For $d \approx 20$ cm, and $h \geq 100$ cm, the volume of the conical region ($d^2/2h^2$) is less than 1% of the volume of the earth and then the angle factors are not important for an approximate estimate. (Podkletnov has taken data up to a distance of 300 cm, and the disc is actually toroidal with an outer diameter of 24.5 cm and an inner diameter of 8 cm. We approximate the effective shielding diameter to around 20 cm and estimate the effects for a circular disc.)

Even after allowing for the fact that ρ_c is somewhat larger than (but less than a factor of 2) ρ , we can see that the shielding should reduce significantly for distances much larger than the diameter of the disc, since the reduction is quadratic in d/h . For $d/h \approx 5$, the shielding should be only about 5% of its value at $d/h \approx 1$, and for $d/h \approx 10$, this should have reduced to about 1%. For the largest value of $d/h \approx 15$ in the experiment, the shielding would diminish to 0.05% of its value near the disc and this distance dependence is not observed in Podkletnov's experiments. On the contrary the shielding was seen to be accurately distance independent. The observed shielding is around 1%-2% near the disc as well as 3 m above the disc whereas it should have diminished to less than 0.01% at these heights. Similar results have been obtained by Modanese employing a Monte Carlo simulation [4]. This crucial consistency check clearly points to some spurious reason for the data observed by Podkletnov and rules out shielding of the earth's gravity as a possible explanation.

In addition to this consistency check, which is the most significant point, we want to point out some additional facts which supports the suspicion that the observed effect is not due to shielding of gravity. We conducted two simple experiments to check for shielding in the following cases:

(1) Measurement of the weight of a sample hung above a superconducting YBCO disc in the superconducting state, immersed in liquid nitrogen, without rotation or magnetic field.

(2) Measurement of the weight of the sample over the disc when there are static magnetic fields below the disc, generating supercurrents, but with the disc nonrotating.

The sensitivity of these measurements (using a Metler AE240 balance) was such that a weight change of 0.001% could be measurable. In the first measurement we did see a weight reduction amounting to 0.05% or so, but the same weight reduction was observed even when the superconducting disc was removed, but the liquid nitrogen dewar remained in place! The observed weight reduction in this case was verified to be due to the additional buoyancy in the cold nitrogen atmosphere and partly due to air currents from evaporating nitrogen.

In the second measurement with magnets directly underneath the superconducting disc, the results were

the same as in the first set and this suggests that the weight reduction observed by Podkletnov in the static situation is spurious (the magnetic field was about 20 times the lower critical field and considerable flux penetration was expected). Of course, we have not reproduced exactly the same experimental condition as in Podkletnov's experiment, but we certainly have reproduced some of the physical ingredients which could have given such an effect, like supercurrents. Also, Podkletnov's experiments were done at a lower temperature. It might still be worthwhile to do an experiment replicating Podkletnov's experiment, but the real value of such an experiment would be in finding out whether it is buoyancy-related effects or electromagnetic effects that are responsible for the observed data.

It is possible to improve the experiments described in Refs. [1] and [2] on several fronts. The overall sensitivity itself can be improved by a factor of 5 to 10 using better commercial balances and this will allow a more careful measurement to be made on the weight changes as a function of height from the disc as well as with position in the horizontal plane over the disc. To see whether free rotation without time-varying electromagnetic fields can lead to weight changes, the disc can be rotated with a gas jet and the levitation itself can possibly be done on static magnetic fields. It is also important to get the functional dependence of the weight change on the rate of rotation. A more convincing data set could be obtained by keeping the sample on the balance in a chamber isolated from the disc, to avoid air currents, and repeatedly rotating and stopping the disc to see whether there are any reproducible periodic weight changes.

In summary, we have analysed the data from Podkletnov's experiments on the shielding of gravitational field by a superconductor and concluded that the data fail to pass a crucial consistency test. This is important since it is not easy to repeat Podkletnov's experiments exactly and the claimed effect touches fundamentally important issues. Our critique of the claim is more or less independent of the model assumed for possible shielding and thus the shielding of gravity by rotating or static superconductors at the level of 0.05% is ruled out. We have also conducted some preliminary experiments in the static situation and there is no evidence for any shielding.

Acknowledgements

I thank V.P.S. Awana for drawing my attention to a paper by G. Modanese and for lending YBCO discs. I am thankful to G. Modanese and E. Podkletnov for several clarifications.

References

- [1] E. Podkletnov and R. Nieminen, Physica C 203 (1992) 441.
- [2] E.E. Podkletnov and A.D. Levit, Report MSU-95 chem, (1995).
- [3] G. Modanese, preprint, hep-th 9505094 (1995).
- [4] G. Modanese, preprint, supr-con/9601001 (1996);
G. Modanese, private communication.

Quick Search:

within

[\(?\) Search tips](#)

26 of 27

[results list](#)[◀ previous](#)[next ▶](#)**Technol Rep Tohoku Univ**

Volume 43, Issue 2, 1978, Pages 443-450

ISSN: 0040-8816, Coden: TRTUA9

Ei Compendex © 1998 Elsevier Engineering Information, Inc. All Rights Reserved

GENERATION OF ANTI-GRAVITY, OWING TO COUNTERCLOCKWISE TURNING OF GYROSCOPE

Hayasaka, Hideo

Abstract

The weight reduction owing to the counterclockwise turning of a gyroscope discovered by Kozyrev, is considered as the perturbation to the clockwise turning. The perturbation is due to the singularity of time concerning a mirror transformation. The singularity of time corresponds to the singularity of space expressing a mass point in gravitational field and leads to the generation of **anti-gravity**. The condition of no weight change in the clockwise turning leads to the reduction of the gravitational acceleration in the counterclockwise turning; that is, the counterclockwise turning of a gyroscope creates positive potential energy and then generates the **anti-gravity**. [17 Refs]

This Document

- Abstract-Compendex

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

Index Terms: GRAVITATION
Technol Rep Tohoku Univ

Volume 43, Issue 2, 1978, Pages 443-450

ISSN: 0040-8816, Coden: TRTUA9

This Document

- Abstract-Compendex

Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

26 of 27

[results list](#)[◀ previous](#)[next ▶](#)

[Register](#) or Login:

Password:

[Home](#)[Publications](#)[Search](#)[My Alerts](#)[My Profile](#)[Help](#)

Quick Search:

within

[\(?\) Search tips](#)

25 of 27

[results list](#)[◀ previous](#)[next ▶](#)**Technology Reports of the Tohoku University**

Volume 44, Issue 2, 1979, Pages 515-535

ISSN: 0040-8816, Coden: TRTUA9

Ei Compendex © 1998 Elsevier Engineering Information, Inc. All Rights Reserved

RELATIVISTIC MECHANISM OF LOCAL VANISHING OF THE ATTRACTIVE FIELD OF THE EARTH BY VIRTUE OF THE LEFT TURNING MOTION OF A MATTER

This Document[► Abstract-Compendex](#)**Actions**

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

Hayasaka, Hideo

Abstract

The phenomenon of weight reduction in the left turning of a gyroscope discovered by Kozyrev, is explained from the spontaneous symmetry breaking of Christoffel's affinity for left and right turnings. The spontaneous symmetry breakdown is due to the appearance of the latent singularity in the action integral concerning the moment of gravity generated by turning motion with the flow of time. The left turning of gyroscope adds positive potential energy to the gyroscope by absorbing energy from the flow of time and then generates **anti-gravity.** [50 Refs]

Index Terms: GRAVITATIONAL EFFECTS; GYROSCOPES**Technology Reports of the Tohoku University**

Volume 44, Issue 2, 1979, Pages 515-535

ISSN: 0040-8816, Coden: TRTUA9

This Document[► Abstract-Compendex](#)**Actions**

- [Cited By](#)
- [Save as Citation Alert](#)
- [Export Citation](#)

25 of 27

[results list](#)[◀ previous](#)[next ▶](#)[Home](#)[Publications](#)[Search](#)[My Alerts](#)[My Profile](#)[Help](#)

We are a **LIVE** online global community!

Join now for a

FREE membership!

Search news

Free newsletter

NewScientist.com

PPA Interactive Magazine of the Year

HOME | NEWS | HOT TOPICS | THE LAST WORD | OPINION | WEBLINKS | PRINT EDITION | SUBSCRIBE | ARCHIVE | JOBS & CAREERS

NEWS

ALL LATEST NEWS

CONFERENCE NEWS

SEARCH NEWS

TOP NEWS STORIES

[Triple-action drug battles botulism](#)

[Red tide microbe's killer appetite revealed](#)

[Middle-eastern farmers 'civilised' Europe](#)

[Giant galaxy means model success](#)

[vCJD risk from blood transfusion "appreciable"](#)

['Clean' human stem cells grown](#)

[All latest news](#)

TOP SCIENCE JOBS

The World's No.1 Science & Technology News Service

Gravity's quantum leaps detected

19:00 16 January 02

Gravity's subtle influence in the quantum world has been directly observed for the first time.

On tiny scales, nature makes particles behave according to curiously rigid rules. For instance, negatively charged electrons trapped around a positive nucleus under the pull of the electromagnetic force cannot have any energy they want -they have to fall into a set of distinct energy levels.

In the same way, the pull of gravity should make particles fall into discrete energy levels. But because gravity is extremely weak on small scales, the effect has been impossible to spot. "To be able to measure it, you need to suppress interference from all the other fields," says Valery Nesvizhevsky of the Laue-Langevin Institute in Grenoble, France.

Now Nesvizhevsky and his colleagues have achieved the feat using a beam of neutrons. Neutrons were ideal because they're neutral, so they don't feel the electromagnetic force and can ignore its quantum rules.

"The experiment is a real tour-de-force in making the first demonstration of a quantum effect due to gravity," says Thomas Bowles of Los Alamos National Laboratory in New Mexico. "Although everyone feels the pull of gravity on their bodies, the pull of gravity on a neutron is smaller by more than a trillion trillion times."

Bowles says the accuracy of the experiment is stunning. "Imagine putting a 747 aeroplane on a scale and weighing it. Then take one grain of sand and add it to the scale. The fractional change you would observe on the scale readings would still be a billion times larger than the fractional change that Nesvizhevsky was able to observe in their experiment."

Bouncing neutrons

Nesvizhevsky's team took a beam of ultracold neutrons with tiny energies, moving from left to right at less than eight metres per second. Under the force of gravity, the neutrons fell down onto a reflecting mirror and bounced off it before arriving at a detector.

The team could limit the energies of the neutrons arriving at the

SUBSCRIBE

Subscribe to New Scientist print edition and enjoy

4 FREE ISSUES

& UNLIMITED ACCESS to our 10 year **Archive**



ONLINE CONFERENCE REPORTS

[AIDS 2002 - Barcelona](#)

MORE ON THIS STORY

Related Stories

[Flowing liquid revealed as quantum wave](#)

4 July 2001

[Quantum dots to barcode DNA](#)

2 June 2001

[Delicate quantum states can be shielded from destructive noise](#)

4 January 2001

For more related stories
search the print edition [Archive](#)

Weblinks

[Laue-Langevin Institute](#)

[Nature](#)

ELSEWHERE TODAY

All the best science stories from the web

[Mouse genome mapped](#)

Independent, UK

[Intriguing clues in anthrax killer case](#)

Newsweek

[UK legionnaires' outbreak flattens out](#)

BBC

Best live jobs in the hottest areas of sciences[Biotechnology](#)[Environmental Science](#)[Computer Science](#)[Biochemistry](#)[Courses](#)Or [search all](#) of our
jobs and
opportunities

detector by placing an absorbing material at different heights above the mirror. The material mopped up all the neutrons that bounced too high.

Forgetting quantum mechanics, you would expect neutrons with any energy to arrive at the detector. But no neutrons appeared unless the neutron-mop was at least 15 micrometres above the mirror. This means the neutrons have to have a certain, minimum energy (equal to 1.41×10^{-12} electronvolts) in the Earth's gravitational field.

Step up

There were also hints that neutron transmission took little leaps at different, higher energies, corresponding to higher quantum levels. However, the team has still to confirm this.

Nesvizhevsky says the technology is exciting because it could test some other key ideas in physics - for instance, whether or not the neutron carries some minuscule amount of electric charge. "If it's there, it's very, very small," says Nesvizhevsky.

It could also put on trial the equivalence principle, a famous concept of Einstein's. It says that all particles, regardless of their mass or composition, should fall with the same acceleration in a uniform gravitational field.

Journal reference: *Nature* (vol 415, p 297)

Hazel Muir

This story is from NewScientist.com's news service - for more exclusive news and expert analysis every week [subscribe](#) to New Scientist print edition.

[Print this article](#)[Send to a friend](#)

- ▶ For what's in New Scientist magazine this week see [Print Edition](#)
- ▶ Search the [Archive](#) for more stories like this, originally published in the Print Edition
- ▶ [Subscribe](#) to New Scientist Print Edition
- ▶ [Correspondence](#) about this story should be directed to onlinenews@newscientist.com
- ▶ [Sign up for our free newsletter](#)

[GPS sparks boundary wars on Rhode Island](#)[MSNBC](#)[Is this the iceberg that sank the Titanic?](#)[Discovery Channel](#)[Spam and the email arms race](#)[CNN](#)**NewScientist Archive**


We are a **LIVE** online global community!

Join now for a

FREE membership!

Search news

Free newsletter

NewScientist.com

PPA Interactive Magazine of the Year

HOME | NEWS | HOT TOPICS | THE LAST WORD | OPINION | WEBLINKS | PRINT EDITION | SUBSCRIBE | ARCHIVE | JOBS & CAREERS

NEWS

ALL LATEST NEWS

CONFERENCE NEWS

SEARCH NEWS

TOP NEWS STORIES

[Triple-action drug battles botulism](#)

[Red tide microbe's killer appetite revealed](#)

[Middle-eastern farmers 'civilised' Europe](#)

[Giant galaxy means model success](#)

[vCJD risk from blood transfusion "appreciable"](#)

['Clean' human stem cells grown](#)

[All latest news](#)

TOP SCIENCE JOBS

The World's No.1 Science & Technology News Service

Anti-gravity research on the rise

12:31 30 July 02

NewScientist.com news service

Researchers around the world are opening their minds to the possibility that the phenomenon of anti-gravity is not just science fiction.

Most respected physicists still scoff at the idea that experimental equipment can reduce gravity, but several groups have been working on it independently and are coming to the same conclusion: it might just be true.

On Monday, reports re-emerged that Boeing, the American aircraft manufacturer, is interested in exploring the possibility of building an anti-gravity device. The news, first revealed in **New Scientist** magazine in January 2002, centres around Russian scientist, Evgeny Podkletnov. In 1992 he claimed to be the first person to witness the reduction of gravity above a spinning superconducting disc.

Podkletnov, a specialist in superconductors, says he stumbled on the effect whilst performing a routine test on a large superconductor in his laboratory at the Tampere University of Technology, Finland.

High speed spin

Podkletnov met **New Scientist** in late 2001 to outline the experiment, in which a large yttrium-barium-copper-oxide (YBCO) superconducting disc was suspended in nitrogen vapour and cooled to around —233 °C. The disk was levitated in a magnetic field and finally spun at speeds of up to 5000 revolutions per minute by means of an alternating electric current.

He claimed that objects placed above the disc lost around one per cent of their weight. But so far no one has managed to successfully repeat his experiment.

However, several high-profile organisations have taken an active interest in his work. NASA has paid Superconductive Components of Columbus, Ohio, \$600,000 to reproduce the apparatus Podkletnov used in his experiment. There have been delays, but NASA's Ron Koczor told **New Scientist**: "We expect to be ready to test the device in late September 2002."

SUBSCRIBE

Subscribe to New Scientist print edition and enjoy

4 FREE ISSUES

& UNLIMITED ACCESS to our 10 year **Archive**



ONLINE CONFERENCE REPORTS

[AIDS 2002 - Barcelona](#)

MORE ON THIS STORY

Related Stories

[Gravity's quantum leaps detected](#)
16 January 2002

For more related stories
search the print edition [Archive](#)

Weblinks

[Breakthrough Propulsion Physics, NASA](#)

[Greenglow, BAe Systems](#)

[Boeing](#)

[Raymond Chiao, UCB](#)

[The Podkletnov Effect](#)

ELSEWHERE TODAY

All the best science stories from the web

[Mouse genome mapped](#)
Independent, UK

[Intriguing clues in anthrax killer case](#)
Newsweek

[UK legionnaires' outbreak flattens out](#)
BBC

Best live jobs in the hottest areas of sciences[Biotechnology](#)[Environmental Science](#)[Computer Science](#)[Biochemistry](#)[Courses](#)

Or [search all](#) of our
jobs and
opportunities

British defence contractor, BAe Systems, is also interested in the work and set up Project Greenglow to explore the subject. Other groups in Japan, France and Canada are also rumoured to be working on the subject though they have so far kept their identities secret.

Predicted effect

The most intriguing aspect of the affair is that Ning Li, then at the University of Alabama, Huntsville, says she independently predicted Podkletnov's observation in 1989.

Li's theory predicts that if a time-varying magnetic field were applied to a superconductor, charged and deformed lattice ions within the superconductor could absorb enormous amounts of energy. This would cause the lattice ions to spin rapidly about their equilibrium positions and create a minuscule gravitational field.

Li claimed that if these charged, rotating, lattice ions were aligned with each other by a strong magnetic field, the resulting change in local gravity would be measurable.

Early in 2002, Raymond Chiao, a respected physicist at the University of California at Berkley, put forward his own theory relating gravity and superconductors. He predicted that bombarding a superconductor with electromagnetic waves would produce gravitational radiation and is now attempting to prove his theory by experimentation.

David Cohen

This story is from NewScientist.com's news service - for more exclusive news and expert analysis every week [subscribe](#) to New Scientist print edition.

[Print this article](#)



[Send to a friend](#)



- ▶ For what's in New Scientist magazine this week see [Print Edition](#)
- ▶ Search the [Archive](#) for more stories like this, originally published in the Print Edition
- ▶ [Subscribe](#) to New Scientist Print Edition
- ▶ [Correspondence](#) about this story should be directed to onlinenews@newscientist.com
- ▶ [Sign up for our free newsletter](#)

[GPS sparks boundary wars on Rhode Island](#)

MSNBC

[Is this the iceberg that sank the Titanic?](#)

Discovery Channel

[Spam and the email arms race](#)

CNN

NewScientistArchive

FREE
2 SUBSCRIBERS

New Scientist---Antigravity

*See updates at bottom

Date: Fri, 11 Jan 2002

Evgeny Podkletnov has convinced NASA to spend \$600,000 on a machine he claims will shield matter from the Earth's pull. Has the agency flipped? David Cohen tracked down the man who wants to turn the laws of physics on their head

Going up

SEATED in a near-empty restaurant in a backstreet of Tampere in Finland, Evgeny Podkletnov certainly doesn't look crazy—even when he holds up the superconducting disc he says he used to reduce the effects of gravity. The Russian emigre's claim caused such a storm he was thrown out of his job at Tampere University of Technology five years ago. He now works as a researcher in superconducting materials at the nearby University of Tampere, but he's not about to give up his quest to be taken seriously.

Podkletnov claims others have repeated the experiments with great success, and for the moment at least, influential scientists around the world are giving him the benefit of the doubt. Researchers at Toronto University in Canada, at CNRS—France's national research agency—and even an employee of Boeing in the US all want to repeat his experiment, Podkletnov says. And perhaps most significantly of all, NASA is ready to give the idea a shot. This month, after a two-year wait, Ron Koczor and his team at NASA's Marshall Space Flight Center in Huntsville, Alabama, will take delivery of a machine that Koczor believes could shield matter from gravity.

Koczor persuaded NASA to pay Superconductive Components (SCI) of Columbus, Ohio, \$600,000 to build a copy of Podkletnov's apparatus. If SCI's replica works, it could change our way of interacting with a fundamental force of nature. And that, Koczor says, would change everything. Wave goodbye to rockets and the internal combustion engine. Say hello to energy-saving gravity-powered spaceships, aeroplanes, cars and elevators—and a whole new branch of theoretical physics.

Koczor is aware of what the critics will say, but he believes there are hints that it might work and he is determined to keep an open mind. This kind of investigation lies within the Marshall Center's remit to seek out new and exotic forms of propulsion and the potential payoff is huge, he says. "It's worth a little bit of effort to pursue it to its end." But that "little bit of effort" is, essentially,

a gamble on Podkletnov's claims. In 1992 he published a paper describing how he had stumbled across a "gravity shielding" effect while running a routine test on one of his superconductors. The details were sketchy. But the basics are these: make a superconducting disc 145 millimetres in diameter and 6 millimetres thick, according to a special chemical recipe that Podkletnov did not make public. Cool the disc to below -233°C, then levitate it using a magnetic field. Finally, apply an electric current alternating at around 100 kilohertz to coils surrounding the disc. The current makes the disc rotate in the constantly changing magnetic field, something like an electric motor (see Graphic). So far, there's nothing extraordinary here.

But Podkletnov claimed that when the disc was spinning at more than 5000 revolutions per minute, objects placed above it lost around 1 percent of their weight. Increasing the spin speed, he claimed, reduced their weight still further. In subsequent experiments, he claims to have seen weight reductions of up to 2 percent.

Podkletnov concluded that this apparatus somehow reduced the strength of the Earth's pull on any object placed above it and called it a "gravity shielding" device. Stick a more powerful version of this apparatus on the bottom of a spacecraft and rocket propulsion would be history: just the slightest nudge would be needed for lift-off into space. Terrestrial transport would be revolutionized too, together with a large chunk of theoretical physics.

At the time, the paper was greeted without fanfare. It would probably have been forgotten, but for the fact that Podkletnov continued his experiments and, in 1996, produced another paper. Physica D reviewed and accepted it, but its contents were leaked to the press before publication. "The world's first anti-gravity device", as The Sunday Telegraph in Britain called it, was rubbished by scientists around the globe, who loudly proclaimed that it broke the known laws of physics. In the academic scuffle that ensued, Podkletnov was dismissed from his post. After withdrawing the paper (to protect his co-author's career, he says) he disappeared—for a while, at least. I caught up with him in Tampere at the end of last year and found him still adamant that his superconducting disc can shield matter from gravity.

Podkletnov has the air of a persecuted man. While talking about his work his mood shifts constantly between suspicion, seriousness and wild excitement—there are echoes of the cold fusion debate here. But his frustration is clear. "I am a professional scientist and have published more than 30 papers and hold many patents," he

says. "Some people say 'Podkletnov is a fool,' but there are too many other people in the world who have seen this and they all can't be wrong." His English is almost perfect with only a faint Russian accent. He peppers his conversation with references to private conversations with eminent scientists who would come right out and support him, were they not so scared of losing their credibility.

His confidence-and Koczor's-stems from the fact that he is not alone in suggesting a way to modify gravity. In 1995, Koczor and his team were approached by Ning Li, a researcher at the University of Alabama in Huntsville. Li had never met or even heard of Podkletnov, yet she was developing a theory, based on the idea of converting electromagnetic fields into gravitational fields, that came very close to explaining Podkletnov's experiment. She claimed her theory pointed to the possibility of producing a "gravito-magnetic effect" by spinning it super cooled superconducting disc: the angular momentum of fast-spinning ions in the superconductor would produce a gravitational field, she said. By 1995 Li felt she had reached the point where she could approach NASA to fund an experimental test of her ideas.

"We were interested in her theories," says Koczor. "But we thought her experiment was undoable." Then, in a literature search, Koczor and Li found Podkletnov's 1992 paper in the journal *Physica C*. "We were intrigued. It was essentially the same experiment, only simpler," Koczor says. "*Physica C* is not a trivial journal. If [the experiment] got in there then it must have got through sufficient scientific vetting to take to a higher level, so we decided we'd try it ourselves."

For the following two years, Koczor and Li tried to duplicate Podkletnov's experiment. They bought some small superconducting discs, levitated them, put high-frequency electromagnetic fields into them and did a few experiments to measure the gravitational effects. "We tried to see if there was one or other of these factors that could be isolated and identified as responsible for the Podkletnov effect," explains Koczor.

Their experiments were unsuccessful. In 1997 Koczor's team reported their lack of findings in *Physica C*, saying that for their 10-centimetre discs the measurable effect on gravitational pull was a mere two millionths of 1 per cent-small enough to have been background noise in the measuring equipment. But they were not disheartened.

"Podkletnov told us we wouldn't see any effect unless we repeated his experiment faithfully," Koczor says. "We never did the full Podkletnov experiment-we were still learning to work with

these superconductors." And so the team focused on producing a 30-centimetre yttrium-barium-copper-oxide (YBCO) superconducting disc like that used by Podkletnov. But they still didn't have his recipe. Eventually, in 1999 Koczor gave up and commissioned SCI to build a replica of Podkletnov's apparatus. At the same time, Li set up an independent laboratory to pursue the research. SCI contracted Podkletnov as a consultant on its project, asking him to advise on some technical aspects of building the superconductor. "Podkletnov has been as helpful as he could be to get our mission fulfilled," says J.R. Gaines, vice president and general manager of SCI. And so this month-a year behind schedule-Gaines will hand over the finished apparatus.

High hopes

NASA is not the only bona fide organization that has been taking Podkletnov seriously. When British military and aerospace company BAE Systems learned that Clive Woods, a superconductor researcher from Sheffield University, was trying to replicate the experiment it decided it too would hedge its bets and help fund his attempts. "We know we're out on a limb," says Ron Evans, director of Project Greenglow-BAE Systems' research program into alternative forms of propulsion. "But even though we got negative advice from several professors, it seemed to me that for a small amount of money it's worth the gamble. Experts have been wrong before and that's the only thing that makes it worth doing."

Evans is giving Woods an undisclosed sum to reproduce Podkletnov's experiment. So far Woods, too, has been unsuccessful. Like Koczor, Woods believes this could be because he has not managed to reproduce all the conditions Podkletnov says are necessary-the specifications are extremely demanding. &That does not mean there is no effect to be observed, Woods says.

Meanwhile, Podkletnov has been quietly continuing his research. "I am not a rich man," he says. "But I have some funds from other projects and I put everything I have into gravity research. This is my life's dream, my hobby, my goal."

He has made good progress, he says. With the help of friends in a laboratory that once belonged to Moscow's Institute for High Temperatures, he claims to have built an "impulse gravity generator". He says its pulse-produced by a spinning superconductor with a strong electrical charge-is capable of knocking over a book placed on end more than a kilometre away.

The pulse has the same properties as a gravitational field, says Podkletnov. It is

unaffected by an inch-thick steel plate fixed in the beam path, and the force it exerts is changed only with the target's mass, not its constituent material nor its chemical or electromagnetic properties. As he talks about it, he suddenly becomes animated. He thinks it could one day be used to nudge satellites into the correct orbit, and even knock incoming missiles off course. "This is a very powerful device, and I am now in the process of arranging a future project on the gravity generator with serious European firms," he says, almost in a whisper. But, he adds, he cannot divulge which firms-he has signed confidentiality agreements. Although Podkletnov is happy to discuss his work, he says no one can come and look at the gravity pulse experiment. It requires extremely high voltages, and the required generating equipment is, unfortunately, in a restricted area of Moscow State University's campus. So he refused my request to watch the gravity generator in action.

Evans, too, has suggested that an independent observer might visit Podkletnov's Moscow laboratory. Again, Podkletnov refused. "He told me that he once hosted some Japanese visitors to his lab, but they tried to bribe his technicians for the secrets on how the experiment worked," Evans says. &As a result he decided not to bring any other visitors."

Podkletnov, who says he is in the process of patenting his work, is also scared someone might steal his intellectual property rights to the experiment. But Robin Tucker, a theoretical physicist at Lancaster University who is also investigating possible ways to control matter with gravity, thinks Podkletnov's secretive behaviour is odd, to say the least. &Any normal physicist who found this kind of effect would be shouting about it from the tops of the trees and asking people to come and verify it," he says. "It would mean a Nobel Prize if you'd actually discovered some kind of gravity focusing."

Podkletnov's refusal to open up to scrutiny leaves the scientific world lacking any independent, verifiable observations of gravity modification. He gave me an untraceable e-mail address for a Takashi Nakamura, who he claimed was a senior physics professor employed at Toshiba Electronics in Japan. Nakamura responded to my e-mail question, saying that he had managed to reproduce Podkletnov's experiments with even better results. "With all my respect to Evgeny-san, our ceramics is better and we got 8.79% of the weight reduction," he wrote. "Our programme of research has already shown much better efficiency."

However, when I asked for references documenting these results, Nakamura terminated the correspondence.

Quantum suspects

Podkletnov's only current collaboration is with Giovanni Modanese, an Italian physicist who is trying to build a theoretical explanation for Podkletnov's results. But because physicists have such a poor understanding of the mechanisms behind both gravity and high-temperature superconductivity, his explanations are necessarily vague. He suggests that quantum processes within the superconducting material are interacting with quantum processes in the gravitational field. But, he admits, he can't go far with the work because there are too many unknowns. Again, Tucker is skeptical even of this attempt to formalize what's going on. "I think the correlation between the experiment and the theoretical description is very tenuous," he says.

So, frustrating as it is, that's as much as we know at the moment. I contacted several of Ning Li's ex-colleagues, but all said they did not know of her current whereabouts or the state of her research. NASA and BAE Systems still don't know whether they have been sent up a blind alley by Podkletnov's enthusiasm. But Koczor believes he'll soon have the answer. "Running the experiment will take six months at most," he says. If it fails to confirm Podkletnov's experiment, that will be the end of the matter. But if the experiment succeeds, and they can modify gravity, then who knows what could be possible? In the end, pigs really might fly.

*Update From Eugene Podkletnov
<epodkletnov@hotmail.com> July 02, 2002*

It might be possible that the wobble in the superconductor experiments is favorable for the antigravitational effect but I don't think that it is the main reason for the effect. In my experiments the key element is creating a super high density of electrons or Cooper pairs that move in strong magnetic field. Actually, antigravitation is caused by the polarization of vacuum or polarization of sub-atomic particles that constitute the vacuum. There exist several methods of causing this polarization:

1. High speed rotation (80 thousand rpm) of metal crystal lattice.
2. Rotation of magnetic fields that coincide in phase with electric field.
3. High voltage static electric field (the lifter).
4. Resonance frequencies using ELF waves and De Aquino configuration.
5. Topological effect, using grids with special geometry
6. Acoustical vibration at resonance frequencies.

Comment from editor: Podkletnov mentions "antigravitation is caused by the polarization of vacuum." Is gravity a vacuum polarization effect? Perhaps yes? See Polarizable-Vacuum (PV) representation of general relativity by H.E. Puthoff

New Scientist Update 30 July 02,

Anti-gravity research on the rise. Researchers around the world are opening their minds to the possibility that the phenomenon of anti-gravity is not just science fiction. And related to this please see Reverse Engineering at Area- 51? Were Einstein and others involved in the initial program?

Additional References

Static Test for a Gravitational Force Coupled to

TYPE II YBCO Superconductors,

Gravitational Modification---Super Conductors,
varying magnetic field,

Possible quantum gravity effects in a charged
Bose condensate under variable e.m. field,

Impulse Gravity Generator Based on Charged
 $\text{YBa}_2\text{Cu}_3\text{O}_{\{7-y\}}$ Superconductor with
Composite Crystal Structure and,

Role of a "Local" Cosmological Constant in
Euclidean Quantum Gravity Also, related to this
please see Stiff challenge to space-time,
Einstein's theory of relativity tells us that gravity
bends space. Now a physicist at the University of
Portsmouth, UK, has worked out that magnetic
fields may smooth bent space.

Copyrighted© by Robert M. Collins, Editor,2002

Polarizable-Vacuum (PV) representation of general relativity

H. E. Puthoff

Institute for Advanced Studies at Austin

4030 W. Braker Lane, Suite 300, Austin, Texas 78759

puthoff@aol.com

ABSTRACT

Standard pedagogy treats topics in general relativity (GR) in terms of tensor formulations in curved space-time. Although mathematically straightforward, the curved space-time approach can seem abstruse to beginning students due to the degree of mathematical sophistication required. As a heuristic tool to provide insight into what is meant by a curved metric, we present a polarizable-vacuum (PV) representation of GR derived from a model by Dicke and related to the "*THeμ*" formalism used in comparative studies of gravitational theories.

I. INTRODUCTION

Textbook presentations treat General Relativity (GR) in terms of tensor formulations in curved space-time. Such an approach captures in a concise and elegant way the interaction between masses, and their consequent motion. "Matter tells space how to curve, and space tells matter how to move [1]."

Although conceptually straightforward, the curved space-time approach can seem rather abstract to beginning students, and often lacking in intuitive appeal. During the course of development of GR over the years, however, alternative approaches have emerged that provide convenient methodologies for investigating metric changes in less abstract formalisms, and which yield heuristic insight into what is meant by a curved metric.

One approach that has a long history in GR studies, and that does have intuitive appeal, is what can be called the polarizable-vacuum (PV) representation of GR. Introduced by Wilson [2] and developed further by Dicke [3], the PV approach treats metric changes in terms of equivalent changes in the permittivity and permeability constants of the vacuum, ϵ_0 and μ_0 , essentially along the lines of the so-called "*THeμ*" methodology used in comparative studies of gravitational theories [4-6].

In brief, Maxwell's equations in curved space are treated in the isomorphism of a polarizable medium of variable refractive index in flat space [7]; the bending of a light ray near a massive body is modeled as due to an induced spatial variation in the refractive index of the vacuum near the body; the GR reduction in the velocity of light in a gravitational potential as compared to a flat-space reference frame at infinity is represented by an effective increase in the refractive index of the vacuum, and so forth. As elaborated in Refs. [3-7], PV modeling can be

carried out in a self-consistent way so as to reproduce to appropriate order both the equations of GR, and the match to the classical experimental tests of those equations. In what follows we shall continually cross-reference PV-derived results to those obtained by conventional GR techniques to confirm that the PV approach as applied does not introduce spurious results.

II. THE POLARIZABLE VACUUM

At this point one approach would be to introduce an appropriate Lagrangian for the PV formulation, and then derive equations of motion. Recognizing, however, that the PV approach is likely unfamiliar to most readers, we defer such formal derivation to Section IV. In its place we simply present and apply here certain key results of the more formal derivation to problems of interest, such as the three classical tests of GR. In this way we build up for the reader an intuitive picture of what is meant by a polarizable-vacuum representation of curved space-time.

Under non-curved-space conditions the electric flux vector \mathbf{D} in a linear, homogeneous medium can be written

$$\mathbf{D} = \epsilon \mathbf{E} = \epsilon_0 \mathbf{E} + \mathbf{P} \quad (1)$$

where ϵ and ϵ_0 are the permittivities of the medium and of the vacuum, respectively, and the polarization \mathbf{P} corresponds to the induced dipole moment per unit volume in the medium. Writing \mathbf{P} in terms of the polarizability per unit volume, α_V , as $\mathbf{P} = \alpha_V \mathbf{E}$, we obtain the symmetrical form

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \alpha_V \mathbf{E} . \quad (2)$$

The above expression leads naturally to the interpretation of ϵ_0 as the polarizability per unit volume of the vacuum, treated as a medium in its own right. That this interpretation is correct is explicitly corroborated in detail by the picture of the vacuum as developed in quantum field theory. There it is shown that the vacuum acts as a polarizable medium by virtue of induced dipole moments resulting from the excitation of virtual electron-positron pairs [8].

With regard to representing curved-space conditions, the basic postulate for the PV approach is that the polarizability of the vacuum in the vicinity of, say, a mass differs from its asymptotic far-field value by virtue of vacuum polarization effects induced by the presence of the body. That is, we postulate for the vacuum itself

$$\mathbf{D} = \epsilon \mathbf{E} = K \epsilon_0 \mathbf{E} , \quad (3)$$

where K is the (altered) dielectric constant of the vacuum (typically a function of position) due to (GR-induced) vacuum polarizability changes under consideration. *Throughout the rest of our study the vacuum dielectric constant K constitutes the key variable of interest.*

We next examine quantitatively the effects of a polarizable vacuum on the various measurement processes that form the basis of the PV approach to general relativity.

A. Velocity of Light in a Vacuum of Variable Polarizability

We begin our discussion of the polarizable-vacuum model by examining constraints imposed by observation. An appropriate starting point is the expression for the fine structure constant,

$$\alpha = \frac{e^2}{4\pi \epsilon_0 \hbar c} , \quad \text{where} \quad c = \frac{1}{\sqrt{\mu_0 \epsilon_0}} . \quad (4)$$

By the conservation of charge for elementary particles, e can be taken as a constant. Similarly, by conservation of angular momentum for a circularly polarized photon propagating through the vacuum (even with variable polarizability), \hbar can be taken as a constant. Given that ϵ_0 and c can be expected with a variable vacuum polarizability to change to $\epsilon(K) = K\epsilon_0$ and $c'(K) = 1/\sqrt{\mu(K)\epsilon(K)}$, the fine structure constant takes the form

$$\alpha = \frac{e^2}{4\pi \epsilon_0 \hbar c} \sqrt{\frac{\mu(K)/\mu_0}{K}} \quad (5)$$

which is potentially a function of K .

Studies that consider the possibility of the variability of fundamental constants under varying cosmological conditions, however, require that the fine structure constant remain constant in order to satisfy the constraints of observational data [9-11]. Under this constraint we obtain from Eq. (5) $\mu(K) = K\mu_0$; thus the permittivity and permeability constants of the vacuum must change together with vacuum polarizability as

$$\epsilon_0 \rightarrow \epsilon = K\epsilon_0 , \quad \mu_0 \rightarrow \mu = K\mu_0 . \quad (6)$$

This transformation, which maintains constant the ratio $\sqrt{\mu/\epsilon} = \sqrt{\mu_0/\epsilon_0}$ (the impedance of free space) is just what is required to maintain electric-to-magnetic energy ratios constant during adiabatic movement of atoms from one point to another of differing vacuum polarizability [3]. Detailed analysis shows that it is also a necessary condition in the $T\mu\epsilon\mu$ formalism for an electromagnetic test body to fall in a gravitational field with a composition-independent acceleration (WEP, or weak equivalence principle, verified by Eötvös-type experiments) [4-6]. Finally, this condition must be satisfied by any metric theory of gravity, which constitutes the class of viable gravity theories.

The above arguments therefore lead us to our first major conclusion concerning the effects of a variable vacuum polarizability; namely, the permittivity and permeability constants, ϵ and μ , change linearly with the vacuum dielectric constant K as given in Eq. (6), and the velocity of light therefore changes inversely with K in accordance with

$$c (= 1/\sqrt{\mu_0 \epsilon_0}) \longrightarrow c/K (= 1/\sqrt{\mu \epsilon}) . \quad (7)$$

Thus, the dielectric constant of the vacuum plays the role of a variable refractive index under conditions in which vacuum polarizability is assumed to change in response to GR-type influences.

B. Energy in a Vacuum of Variable Polarizability

As will be made explicit in the detailed derivations later, the PV treatment of GR effects is based on the use of equations that hold in special relativity, but with the modification that the velocity of light c in the Lorentz transformations and elsewhere is replaced by the velocity of light in a medium of variable refractive index, c/K , as given in the previous section. Expressions such as $E = mc^2$ are still valid, but now take into account that $c \rightarrow c/K$, and E and m may be functions of K , and so forth.

With regard to expressions for energy, Dicke has shown by application of a limited principle of equivalence that the energy of a system whose value is E_O in flat space ($K = 1$) takes on the value

$$E = \frac{E_O}{\sqrt{K}} \quad (8)$$

in a region where $K \neq 1$ [3]. This is due to the fact that the self-energy of a system changes in response to changes in the local vacuum polarizability. This is analogous to the change in the stored energy of a charged air capacitor during transport to a region of differing dielectric constant.

The energy relationship given by Eq. (8) also implies, via $E_O = m_Oc^2$, which becomes $E(K) = m(K)(c/K)^2$, a corollary change in mass

$$m = m_O K^{3/2} , \quad (9)$$

again a consequence of the change in self-energy.

C. Rod and Clock (Metric) Changes in a Vacuum of Variable Polarizability

Another consequence of the change in energy as a function of vacuum polarizability is a change in associated frequency processes which, by the quantum condition $E = \hbar \omega$ and Eq. (8), takes the form

$$\omega = \frac{\omega_O}{\sqrt{K}} . \quad (10)$$

This, as we shall see, is responsible for the red shift in light emitted from an atom located in a gravitational potential.

From the reciprocal of Eq. (10) we find that time intervals marked by such processes are related by

$$\Delta t = \Delta t_o \sqrt{K} . \quad (11)$$

Therefore, in a gravitational potential (where it will be shown that $K > 1$) the time interval between clock ticks is increased (that is, the clock runs slower) relative to a reference clock at infinity.

With regard to effects on measuring rods, we note that, for example, the radius of the ground-state Bohr orbit of a hydrogen atom

$$\Delta r_o = \frac{\hbar}{m_o c} \frac{1}{\alpha} \quad (12)$$

becomes (with $c \rightarrow c/K$, $m_o \rightarrow m$, and α constant as discussed earlier)

$$\Delta r = \frac{\Delta r_o}{\sqrt{K}} . \quad (13)$$

Other measures of length such as the classical electron radius or the Compton wavelength of a particle lead to the relationship Eq. (13) as well, so this relationship is general. This dependence of fundamental length measures on the variable K indicates that the dimensions of material objects adjust in accordance with local changes in vacuum polarizability - thus there is no such thing as a perfectly rigid rod. From the standpoint of the PV approach this is the genesis of the variable metric that is of such significance in GR studies.

We are now in a position to define precisely what is meant by the label "curved space." In the vicinity of, say, a planet or star, where $K > 1$, if one were to take a ruler and measure along a radius vector R to some circular orbit, and then measure the circumference C of that orbit, one would obtain $C < 2\pi R$ (as for a concave curved surface). This is a consequence of the ruler being relatively shorter during the radial measuring process (see Eq. (13)) when closer to the body where K is relatively greater, as compared to its length during the circumferential measuring process when further from the body. Such an influence on the measuring process due to induced polarizability changes in the vacuum near the body leads to the GR concept that the presence of the body "influences the metric," and correctly so.

Of special interest is the measurement of the velocity of light with "natural" (i.e., physical) rods and clocks in a gravitational potential which have become "distorted" in accordance with Eqns. (11) and (12). Let the end points of a measurement be a "true" distance ΔX apart, and let the time for propagation of a light signal from one end point to the other be a "true" time interval ΔT . Since the measurement is assumed to take place in a gravitational potential with a propagation velocity c/K , ΔX and ΔT are related by

$$\frac{\Delta X}{\Delta T} = \frac{c}{K} . \quad (14)$$

However, a rod whose length is Δr_O at infinity will have shortened in the gravitational potential in accordance with Eq. (13), and therefore the *measured* length ΔX_m will be given not by ΔX but by $\Delta X_m = \Delta X \sqrt{K}$. Similarly, a clock whose rate is determined by Eq. (11), and therefore runs slower in the gravitational potential, will yield for a *measured* propagation time $\Delta T_m = \Delta T / \sqrt{K}$. As a result the *measured* velocity of light, v_m , is given by

$$v_m = \frac{\Delta X_m}{\Delta T_m} = \frac{\Delta X \sqrt{K}}{\Delta T / \sqrt{K}} = \frac{c}{K} \bullet K = c . \quad (15)$$

We thus arrive at the interesting (and significant) result that the measured velocity of light obtained by the use of physical rods and clocks renormalizes from its "true" value c/K to the value c . The PV formalism therefore maintains the universal constancy of the locally *measured* velocity of light.

D. The Metric Tensor

At this point we can make a crossover connection to the standard metric tensor concept that characterizes the conventional GR formulation. In flat space a (4-dimensional) infinitesimal interval is given by the expression

$$ds^2 = c^2 dt_O^2 - (dx_O^2 + dy_O^2 + dz_O^2) . \quad (16)$$

If rods were rigid and clocks non-varying in their performance in regions of differing vacuum polarizability, then the above expression would hold universally. However, a dx_O -length measuring rod placed in a region where $K > 1$, for example, shrinks according to Eq. (13) to $dx = dx_O / \sqrt{K}$. Therefore, the infinitesimal length which would measure dx_O were the rod to remain rigid is now expressed in terms of the dx -length rod as $dx_O = \sqrt{K} dx$. With a similar argument based on Eq. (11) holding for clock rate, Eq. (16) can be written

$$ds^2 = \frac{1}{K} c^2 dt^2 - K(dx^2 + dy^2 + dz^2) . \quad (17)$$

Therefore, the infinitesimal interval takes on the form

$$ds^2 = g_{ij} dx^i dx^j , \quad (18)$$

where g_{ij} in the above expression defines the metric tensor, and

$$dx^0 = c dt, \quad g_{00} = 1/K, \quad g_{11} = g_{22} = g_{33} = -K, \quad g_{ij} = 0 \text{ for } i \neq j . \quad (19)$$

The metric tensor in this form defines an *isotropic coordinate system*, familiar in GR studies.

III. CLASSICAL EXPERIMENTAL TESTS OF GENERAL RELATIVITY IN THE PV MODEL

In the previous sections we have established the concept of the polarizable vacuum and the effects of polarizability changes on metric (rods and clocks) behavior. In particular, we found that metric changes can be specified in terms of a single parameter K , the dielectric constant of the vacuum. This is the basis of the PV approach to GR.

In this section we shall explore, with the aid of expressions to be derived in detail in Section IV, specifically how K changes in the presence of mass, and the effects generated thereby. The effects of major interest at this point comprise the three classical tests of GR; namely, the gravitational redshift, the bending of light and the advance of the perihelion of Mercury. These examples constitute a good testbed for demonstrating the techniques of the PV alternative to the conventional GR curved-space approach.

For the spherically symmetric mass distribution of a star or planet it will be shown later from the basic postulates of the PV approach that the appropriate PV expression for the vacuum dielectric constant K is given by the exponential form

$$K = e^{2GM/rc^2}, \quad (20)$$

where G is the gravitational constant, M is the mass, and r is the distance from the origin located at the center of the mass M . For comparison with expressions derived by conventional GR techniques, it is sufficient for our purposes to restrict consideration to a weak-field approximation based on expansion of the exponential to second order, viz.,

$$K \approx 1 + \frac{2GM}{rc^2} + \frac{1}{2} \left(\frac{2GM}{rc^2} \right)^2. \quad (21)$$

A. Gravitational Redshift

In a gravitation-free part of space, photon emission from an excited atom takes place at some frequency ω_0 , uninfluenced by vacuum polarizability changes. That same emission process taking place in a gravitational field, however, will, according to Eq. (10), have its emission frequency altered (redshifted) to $\omega = \omega_0 / \sqrt{K}$. With the first-order correction to $K = 1$ given by the first two terms in Eq. (21), emission by an atom located on the surface of a body of mass M and radius R will therefore experience a redshift by an amount

$$\frac{\Delta\omega}{\omega_0} = \frac{\omega - \omega_0}{\omega_0} \approx -\frac{GM}{Rc^2}, \quad (22)$$

where we take $GM/Rc^2 \ll 1$. Once emitted, the frequency of the photon remains constant during its propagation to a relatively gravitation-free part of space where its frequency can then be compared against that of local emission, and the spectral shift given by Eq. (22) observed.

Measurement of the redshift of the sodium D_I line emitted on the surface of the sun and received on earth has verified Eq. (22) to a precision of 5% [12].

Experiments carried out on the surface of the earth involving the comparison of photon frequencies at different heights have improved the accuracy of verification still further to a precision of 1% [13-14]. In this case, with the two ends of the experiment separated by a vertical height h , the first-order frequency shift is calculated with the aid of Eqns. (10) and (21) as

$$\frac{\Delta\omega}{\omega} \approx \frac{GM}{R^2c^2} h, \quad (23)$$

where M and R are the mass and radius of the earth. This experiment, carried out at Harvard University, required a measurement accuracy of $\Delta\omega/\omega \sim 10^{-15}$ for a height $h = 22.5$ meters. It was accomplished by the use of Mössbauer-effect measurement of the difference between γ -ray emission and absorption frequencies at the two ends of the experiment.

B. Bending of Light Rays

We consider now the case of a light ray passing in close proximity to a massive body, say, the sun. Given that the velocity of light in a gravitational potential is given by c/K , and given that K varies as shown in Eq. (21), the light ray in essence passes through a medium of variable refractive index.

Since K rises in value near the body, there is a corresponding reduction in the velocity of light in that region. Therefore, for a light ray grazing the body, that part of the wavefront closest to the body will, by virtue of its reduced velocity, cause the wavefront to "wheel about" the body, so to speak. As a result the light ray is deflected toward the body. This deflection, in GR terms, yields a measure of "local curvature," while in the PV approach it is interpreted as a measure of the spatially dependent vacuum polarizability.

Quantitatively, the magnitude of the deflection can be calculated as follows, where we use the sun as an example. Since the deflection is small (less than two seconds of arc), small-angle approximations can be used throughout.

As a light ray grazes the sun, the wavefront tilt, and hence the deflection angle, is found by calculating the accumulated difference in wavefront advancement across the ray (transverse to the direction of propagation) due to the slight difference in phase velocity. Since the velocity of light is given by $v_L = c/K$, to first order Eq. (21) yields

$$v_L = c/K \approx c \left(1 + \frac{2GM}{rc^2} \right) \approx c \left(1 - \frac{2GM}{rc^2} \right). \quad (24)$$

With the geometry of a grazing ray as shown in Fig. 1(a), Eq. (24) can be written

$$v_L \approx c \left[1 - \frac{2GM}{c^2} \frac{1}{\sqrt{(R+\delta)^2 + z^2}} \right] . \quad (25)$$

Therefore, the differential velocity across the ray is (for $\delta \ll R$)

$$\Delta v_L \approx \frac{2GM}{c} \frac{R\delta}{(R^2 + z^2)^{3/2}} . \quad (26)$$

Reference to Fig. 1(b) indicates that, as the ray travels a distance $dz \approx c dt$, the differential velocity across the ray results in an accumulated wavefront difference Δz given by

$$\Delta z = \Delta v_L dt \approx \frac{2GM}{c^2} \frac{R\delta}{(R^2 + z^2)^{3/2}} dz \quad (27)$$

and an accumulated tilt angle

$$d\alpha \approx \tan(d\alpha) = \frac{\Delta z}{\delta} \approx \frac{2GM}{c^2} \frac{R}{(R^2 + z^2)^{3/2}} dz . \quad (28)$$

Integrating over the entire ray path from $z = -\infty$ to $z = +\infty$ we obtain

$$\alpha \approx \frac{4GM}{Rc^2} . \quad (29)$$

For the sun this gives 1.75 seconds of arc, a value first verified within experimental error by measurements of grazing starlight images made during total eclipses of the sun in 1919. Further confirmation was obtained in the form of deflection by the planet Jupiter of waves from a strong celestial radio source by an amount of 300 microseconds of arc [15].

C. Advance of the Perihelion of Mercury

The third classical test of GR theory involves a prediction of a slight precession (rotation) of the elliptical orbit of a planet or satellite with each orbital revolution. For even the closest planet to the sun, Mercury, however, the advance of the perihelion amounts to only 43 seconds of arc per century, and is therefore an extremely small effect.

Analysis of the perihelion advance in the PV approach exhibits quite explicitly how changes in vacuum properties lead to GR effects. We begin with the standard Lagrangian for a *free* particle [16],

$$L = -mc^2 \sqrt{1 - \left(\frac{v}{c}\right)^2} , \quad (30)$$

only we note that the mass $m(K)$, given by Eq. (9), and the velocity of light $c(K)$, given by Eq. (7), are now functions of the vacuum dielectric constant K ,

$$L = -m(K)c(K)^2 \sqrt{1 - \left(\frac{v}{c/K}\right)^2} = -m_0 K^{3/2} \left(\frac{c}{K}\right)^2 \sqrt{1 - \frac{\left(\dot{r}^2 + r^2 \dot{\theta}^2\right)}{(c/K)^2}} . \quad (31)$$

With K given by Eq. (20), the above takes the form

$$L = -m_0 c^2 e^{-GM/rc^2} \sqrt{1 - \frac{\left(\dot{r}^2 + r^2 \dot{\theta}^2\right) e^{4GM/rc^2}}{c^2}} . \quad (32)$$

Substitution into Lagrange's general equations of motion

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) - \frac{\partial L}{\partial q} = 0 \quad (33)$$

then leads to the specific equations of motion

$$\frac{d}{dt} \left(m r^2 \dot{\theta} \right) = 0 , \quad (34a)$$

$$\frac{d}{dt} \left(m \dot{r} \right) = -\frac{GMm}{r^2} e^{-4GM/rc^2} + m r \dot{\theta}^2 - \frac{2GMm}{r^2 c^2} \left(\dot{r}^2 + r^2 \dot{\theta}^2 \right) , \quad (34b)$$

where $m = m_0 K^{3/2} = m_0 \times \exp(3GM/rc^2)$ and, once derivatives are taken, we assume $\sqrt{1 - (v/c(K))^2} \approx 1$ for nonrelativistic motion.

The solution to Eq. (34a) yields an expression for the angular momentum ℓ ,

$$m r^2 \dot{\theta} = \ell . \quad (35)$$

Substitution into Eq. (34b), followed by elimination of the variable t by means of $d/dt = \dot{\theta} d/d\theta$, then leads to an equation for the planetary trajectory in the (r, θ) plane,

$$\frac{d}{d\theta} \left(\frac{1}{r^2} \frac{dr}{d\theta} \right) = -\frac{GMm_0^2}{\ell^2} e^{2GM/rc^2} + \frac{1}{r} - \frac{2GM}{c^2} \left[\frac{1}{r^2} + \frac{1}{r^4} \left(\frac{dr}{d\theta} \right)^2 \right] . \quad (36)$$

With a change of variable to $u = 1/r$, we obtain, finally,

$$\frac{d^2u}{d\theta^2} + u = \frac{GMm_o^2}{\ell^2} e^{2GMu/c^2} + \frac{2GM}{c^2} \left[u^2 + \left(\frac{du}{d\theta} \right)^2 \right]. \quad (37)$$

Taking $\exp(2GM/rc^2) \approx 1 + 2GM/rc^2$, that is, $\exp(2GMu/c^2) \approx 1 + 2GMu/c^2$, we assume a solution of the form

$$u = A(1 + e \cos \gamma \theta), \quad (38)$$

where here e is the eccentricity of the orbit ($e = 0.206$ for Mercury). This leads to good approximation to an expression for γ of the form

$$\gamma = \sqrt{1 - 6 \left(\frac{GMm_o}{\ell c} \right)^2} \approx 1 - 3 \left(\frac{GMm_o}{\ell c} \right)^2. \quad (39)$$

For one revolution of the planet in its solar orbit, the second term in Eq. (39) represents a correction to the nonprecessing Newtonian orbit. In particular, it corresponds to a perihelion advance per orbit of

$$\Delta\theta = 2\pi \times 3 \left(\frac{GMm_o}{\ell c} \right)^2 = 6\pi \left(\frac{GMm_o}{\ell c} \right)^2, \quad (40)$$

which for the planet Mercury accumulates to the observed value of 43 seconds of arc per century.

From the standpoint of conventional GR theory the predicted and observed perihelion advance constitutes remarkable corroboration of the curved-space concept. From the standpoint of the PV approach such a result can be interpreted as corroboration of the fact that the dielectric constant of the vacuum in the vicinity of mass is modified by the presence of that mass. The epistemological link between the two approaches can be seen in the fact that in neither is gravity treated as a force *per se*. Rather, in both cases the test body is treated as a *free* particle (note the use of a free-particle Lagrangian in the PV approach) whose trajectory is determined "by the metric properties of space itself."

D. Comparison of the Metric Tensors in the GR and PV Approaches

Having shown by specific calculation that the PV approach to the three classical tests of GR reproduces the traditional GR results, we now establish the underlying generality of such correspondence.

In standard textbook treatments of the three classical tests of GR one begins with the Schwarzschild metric, which in isotropic coordinates is written [17]

$$ds^2 = g_{ij} dx^i dx^j$$

$$= \left(\frac{1 - \frac{GM}{2rc^2}}{1 + \frac{GM}{2rc^2}} \right)^2 c^2 dt^2 - \left(1 + \frac{GM}{2rc^2} \right)^4 (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2). \quad (41)$$

Expanding the metric tensor for small departures from flatness as a Maclaurin series in (GM/rc^2) , we obtain

$$g_{00} = \left(\frac{1 - \frac{GM}{2rc^2}}{1 + \frac{GM}{2rc^2}} \right)^2 = 1 - 2 \left(\frac{GM}{rc^2} \right) + 2 \left(\frac{GM}{rc^2} \right)^2 - \dots \quad (42)$$

$$g_{11} = g_{22} = g_{33} = - \left(1 + \frac{GM}{2rc^2} \right)^4 = - \left[1 + 2 \left(\frac{GM}{rc^2} \right) + \dots \right]. \quad (43)$$

Similarly, in the PV approach one begins with the exponential metric defined by Eqns. (17) - (20),

$$ds^2 = g_{ij} dx^i dx^j$$

$$= e^{-2GM/rc^2} c^2 dt^2 - e^{2GM/rc^2} (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2). \quad (44)$$

This, when expanded to the same order as the Schwarzschild metric tensor above for small departures from unity vacuum dielectric constant, yields

$$g_{00} = e^{-2GM/rc^2} = 1 - 2 \left(\frac{GM}{rc^2} \right) + 2 \left(\frac{GM}{rc^2} \right)^2 - \dots, \quad (45)$$

$$g_{11} = g_{22} = g_{33} = -e^{2GM/rc^2} = - \left[1 + 2 \left(\frac{GM}{rc^2} \right) + \dots \right]. \quad (46)$$

Comparison of Eqns. (45) - (46) with Eqns. (42) - (43) reveals that, to the order of expansion shown, the two metric tensors are identical. Since the three classical tests of GR do not require terms beyond these explicitly displayed, the agreement between theory and experiment is accounted for equally in both the conventional GR and in the alternative PV formalisms [18].

IV. COUPLED MATTER-FIELD EQUATIONS

In the preceding section we have seen that the three classical tests of GR theory, redshift, bending of light rays, and the advance of the perihelion of Mercury, are accounted for in the PV formalism on the basis of a variable vacuum dielectric constant, K . To carry that out we stated without proof that the appropriate mathematical form for the variation in K induced by the presence of mass is an exponential form, which, under the weak-field conditions prevailing in the solar system, can be treated in terms of the series expansion

$$K = e^{2GM/rc^2} = 1 + 2\left(\frac{GM}{rc^2}\right) + \dots . \quad (47)$$

In this section we show how the exponential form is derived from first principles, and, in the process, establish the general approach to the derivation of field equations as well as the equations for particle motion. The approach consists of following standard Lagrangian techniques as outlined, for example, in Ref. 16, but with the proviso that in our case the dielectric constant of the vacuum is treated as a variable function of time and space.

A. Lagrangian Approach

As stated earlier, the standard Lagrangian for a free particle is given by the form Eq. (30)

$$L^p = -mc^2 \sqrt{1 - \left(\frac{v}{c}\right)^2} , \quad (48)$$

which, in the presence of a variable vacuum dielectric constant K is modified to Eq. (31) with the aid of Eqns. (7) and (9) to read

$$L^p = -\frac{m_0 c^2}{\sqrt{K}} \sqrt{1 - \left(\frac{v}{c/K}\right)^2} . \quad (49)$$

This implies a Lagrangian *density* for the particle of

$$L_d^p = -\frac{m_0 c^2}{\sqrt{K}} \sqrt{1 - \left(\frac{v}{c/K}\right)^2} \delta^3(\mathbf{r} - \bar{\mathbf{r}}) , \quad (50)$$

where $\delta^3(\mathbf{r} - \bar{\mathbf{r}})$ is the three-dimensional delta function that serves to locate the (point) particle at $\mathbf{r} = \bar{\mathbf{r}}$.

Following standard procedure, the particle Lagrangian density can be extended to the case of interaction with electromagnetic fields by the addition of terms involving the scalar and vector potentials (Φ, \mathbf{A}) ,

$$L_d^p = - \left(\frac{m_o c^2}{\sqrt{K}} \sqrt{1 - \left(\frac{v}{c/K} \right)^2} + q\Phi - q\mathbf{A} \bullet \mathbf{v} \right) \delta^3(\mathbf{r} - \bar{\mathbf{r}}), \quad (51)$$

where (Φ, \mathbf{A}) are related to the electromagnetic field vectors (\mathbf{E}, \mathbf{B}) by

$$\mathbf{E} = -\nabla\Phi - \frac{\partial\mathbf{A}}{\partial t}, \quad \mathbf{B} = \nabla \times \mathbf{A}. \quad (52)$$

The Lagrangian density for the electromagnetic fields themselves, as in the case of the particle Lagrangian, is given by the standard expression (see, e.g., Ref. 16), except that again K is treated as a variable,

$$L_d^{em} = -\frac{1}{2} \left(\frac{B^2}{K\mu_o} - K\epsilon_o E^2 \right). \quad (53)$$

We now need a Lagrangian density for the dielectric constant variable K , which, being treated as a scalar variable, must take on the standard Lorentz-invariant form for propagational disturbances of a scalar,

$$L_d^K = -\lambda f(K) \left[(\nabla K)^2 - \frac{1}{(c/K)^2} \left(\frac{\partial K}{\partial t} \right)^2 \right], \quad (54)$$

where $f(K)$ is an arbitrary function of K . As indicated by Dicke in the second citation of Ref. 3, a correct match to experiment requires that we take $\lambda = c^4/32\pi G$ and $f(K) = 1/K^2$; thus,

$$L_d^K = -\frac{\lambda}{K^2} \left[(\nabla K)^2 - \frac{1}{(c/K)^2} \left(\frac{\partial K}{\partial t} \right)^2 \right]. \quad (55)$$

We can now write down the total Lagrangian density for matter-field interactions in a vacuum of variable dielectric constant,

$$L_d = - \left(\frac{m_o c^2}{\sqrt{K}} \sqrt{1 - \left(\frac{v}{c/K} \right)^2} + q\Phi - q\mathbf{A} \bullet \mathbf{v} \right) \delta^3(\mathbf{r} - \bar{\mathbf{r}}) - \frac{1}{2} \left(\frac{B^2}{K\mu_o} - K\epsilon_o E^2 \right) - \frac{\lambda}{K^2} \left[(\nabla K)^2 - \frac{1}{(c/K)^2} \left(\frac{\partial K}{\partial t} \right)^2 \right]. \quad (56)$$

B. General Matter-Field Equations

Variation of the Lagrangian density $\delta \int L_d dx dy dz dt$ with regard to the particle variables as per standard action principle techniques (Ref. 16) leads to our first important equation, the equation for particle motion in a variable dielectric vacuum,

$$\frac{d}{dt} \left[\frac{(m_o K^{3/2}) \mathbf{v}}{\sqrt{1 - \left(\frac{v}{c/K} \right)^2}} \right] = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}) - \nabla \left(\frac{m_o c^2}{\sqrt{K}} \sqrt{1 - \left(\frac{v}{c/K} \right)^2} \right) \quad (57)$$

or

$$\frac{d}{dt} \left[\frac{(m_o K^{3/2}) \mathbf{v}}{\sqrt{1 - \left(\frac{v}{c/K} \right)^2}} \right] = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}) + \frac{(m_o c^2 / \sqrt{K})}{\sqrt{1 - \left(\frac{v}{c/K} \right)^2}} \left(\frac{1 + \left(\frac{v}{c/K} \right)^2}{2} \right) \frac{\nabla K}{K}. \quad (58)$$

We see that accompanying the usual Lorentz force is an additional dielectric force proportional to the gradient of the vacuum dielectric constant. This term is equally effective with regard to both charged and neutral particles and accounts for the familiar gravitational potential, whether Newtonian in form or, as in the last section, taken to higher order to account for GR effects. One note of passing interest is the fact that as $m_o \rightarrow 0$, but $v \rightarrow c/K$, the deflection of a zero-mass particle (e.g., a photon) in a gravitational field is twice that of a slow-moving particle ($v \rightarrow 0$), an important result in GR dynamics.

Variation of the Lagrangian density with regard to the K variable leads to the second equation of interest, an equation for the generation of GR vacuum polarization effects due to the presence of matter and fields. (In the final expression we use $\nabla^2 K = (1/2K)(\nabla K)^2 + 2\sqrt{K}\nabla^2\sqrt{K}$ to obtain a form convenient for the following discussion.)

$$\begin{aligned} \nabla^2 \sqrt{K} - \frac{1}{(c/K)^2} \frac{\partial^2 \sqrt{K}}{\partial t^2} &= -\frac{\sqrt{K}}{4\lambda} \left\{ \frac{(m_o c^2 / \sqrt{K})}{\sqrt{1 - \left(\frac{v}{c/K} \right)^2}} \left[\frac{1 + \left(\frac{v}{c/K} \right)^2}{2} \right] \delta^3(\mathbf{r} - \bar{\mathbf{r}}) \right. \\ &\quad \left. + \frac{1}{2} \left(\frac{B^2}{K \mu_o} + K \epsilon_o E^2 \right) - \frac{\lambda}{K^2} \left[(\nabla K)^2 + \frac{1}{(c/K)^2} \left(\frac{\partial K}{\partial t} \right)^2 \right] \right\}. \end{aligned} \quad (59)$$

Thus we see that changes in the vacuum dielectric constant K are driven by mass density (first term), EM energy density (second term), and the vacuum polarization energy density itself (third term). The fact that the latter term corresponds to the energy density of the K variable can be seen by the following argument. We start with the Lagrangian density Eq. (55), define the momentum density as usual by $\pi = \partial L_d^K / \partial(\partial K / \partial t)$, and form the Hamiltonian energy density by standard techniques to obtain

$$H_d^K = \pi \left(\frac{\partial K}{\partial t} \right) - L_d^K = \frac{\lambda}{K^2} \left[(\nabla K)^2 + \frac{1}{(c/K)^2} \left(\frac{\partial K}{\partial t} \right)^2 \right]. \quad (60)$$

Again, in passing, of interest is the fact that the energy densities of the EM fields and the K variable enter into Eq. (59) with opposite signs. Therefore, EM field effects can counteract gravitational field effects. This will become more apparent when we examine the so-called "electrogravitic repulsion forces" associated with the Reissner-Nordstrøm solution to Eq. (59).

Eqns. (58) and (59), together with Maxwell's equations for propagation in a medium with variable dielectric constant, thus constitute the master equations to be used in discussing matter-field interactions in a vacuum of variable dielectric constant as required in the PV formulation of GR.

V. STATIC FIELD SOLUTIONS

We bring our tutorial on the PV approach to a close by demonstrating application of the field Eq. (59) to two static field cases with spherical symmetry: derivation of the expression introduced earlier for the gravitational field alone, and derivation of the corresponding expression for charged masses.

A. Static Fields (Gravitational)

In space surrounding an uncharged spherical mass distribution the static solution ($\partial K / \partial t = 0$) to Eq. (59) is found by solving

$$\nabla^2 \sqrt{K} = \frac{1}{4K^{3/2}} (\nabla K)^2 = \frac{1}{\sqrt{K}} (\nabla \sqrt{K})^2 \quad (61a)$$

or

$$\frac{d^2 \sqrt{K}}{dr^2} + \frac{2}{r} \frac{d\sqrt{K}}{dr} = \frac{1}{\sqrt{K}} \left(\frac{d\sqrt{K}}{dr} \right)^2, \quad (61b)$$

where we have used $(\nabla K)^2 = 4K(\nabla \sqrt{K})^2$.

The solution that satisfies the Newtonian limit is given by

$$\sqrt{K} = e^{GM/r c^2} \quad (62a)$$

or

$$K = e^{2GM/r c^2} = 1 + 2\left(\frac{GM}{rc^2}\right) + \dots , \quad (62b)$$

which can be verified by substitution into the equation for particle motion, Eq. (58). We have thus derived from first principles the exponential form of the variable dielectric constant in the vicinity of a mass as used in earlier sections. As indicated in Section III-D, this solution reproduces to appropriate order the standard GR Schwarzschild metric predictions as they apply to the weak-field conditions prevailing in the solar system.

B. Static Fields (Gravitational Plus Electrical)

We now take up the important case of a static metric generated by a mass M with charge Q . Assuming spherical symmetry, we obtain the electric field appropriate to a charged mass imbedded in a variable-dielectric-constant medium,

$$\int \mathcal{D} \bullet d\vec{a} = K\epsilon_0 E 4\pi r^2 = Q \quad (63a)$$

or

$$E = \frac{Q}{4\pi K\epsilon_0 r^2} . \quad (63b)$$

Substitution into Eq. (59) yields

$$\nabla^2 \sqrt{K} = \frac{1}{\sqrt{K}} \left[(\nabla \sqrt{K})^2 - \frac{b^2}{r^4} \right] \quad (64a)$$

or

$$\frac{d^2 \sqrt{K}}{dr^2} + \frac{2}{r} \frac{d\sqrt{K}}{dr} = \frac{1}{\sqrt{K}} \left[\left(\frac{d\sqrt{K}}{dr} \right)^2 - \frac{b^2}{r^4} \right] , \quad (64b)$$

where $b^2 = Q^2 G / 4\pi \epsilon_0 c^4$.

The solution to Eq. (64) as a function of charge (represented by b) and mass (represented by $a = GM/c^2$) is given below. Substitution into Eq. (58) verifies that as $r \rightarrow \infty$ this expression asymptotically approaches the standard flat-space equations for particle motion about a body of charge Q and mass M .

$$\sqrt{K} = \cosh\left(\frac{\sqrt{a^2 - b^2}}{r}\right) + \frac{a}{\sqrt{a^2 - b^2}} \sinh\left(\frac{\sqrt{a^2 - b^2}}{r}\right) , \quad a^2 > b^2 . \quad (65)$$

(For $b^2 > a^2$ the solution is trigonometric.)

Now let us compare these results for the weak-field case with the comparable Reissner-Nordström solution of the conventional GR approach, which in isotropic coordinates is written [19]

$$\begin{aligned}
ds^2 &= g_{ij} dx^i dx^j \\
&= \left[\frac{1 - \left(\frac{a}{2r} \right)^2 + \left(\frac{b}{2r} \right)^2}{\left(1 + \frac{a}{2r} \right)^2 - \left(\frac{b}{2r} \right)^2} \right]^2 c^2 dt^2 \\
&\quad - \left[\left(1 + \frac{a}{2r} \right)^2 - \left(\frac{b}{2r} \right)^2 \right]^2 (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2). \tag{66}
\end{aligned}$$

Expanding the metric tensor for small departures from flatness as a Maclaurin series in $(a/r) = (GM/rc^2)$, we obtain

$$g_{00} = \left[\frac{1 - \left(\frac{a}{2r} \right)^2 + \left(\frac{b}{2r} \right)^2}{\left(1 + \frac{a}{2r} \right)^2 - \left(\frac{b}{2r} \right)^2} \right]^2 = 1 - 2 \left(\frac{a}{r} \right) + \left(2 + \frac{b^2}{a^2} \right) \left(\frac{a}{r} \right)^2 - \dots, \tag{67}$$

$$g_{11} = g_{22} = g_{33} = - \left[\left(1 + \frac{a}{2r} \right)^2 - \left(\frac{b}{2r} \right)^2 \right]^2 = - \left[1 + 2 \left(\frac{a}{r} \right) + \dots \right]. \tag{68}$$

The comparable expansion in the PV approach [see Eqns. (18) - (19)],

$$\begin{aligned}
ds^2 &= g_{ij} dx^i dx^j \\
&= \frac{1}{K} c^2 dt^2 - K (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2), \tag{69}
\end{aligned}$$

when expanded by the use of Eq. (65) to the same order as the Reissner-Nordström metric tensor above, yields

$$g_{00} = 1 - 2 \left(\frac{a}{r} \right) + \left(2 + \frac{b^2}{a^2} \right) \left(\frac{a}{r} \right)^2 - \dots, \tag{70}$$

$$g_{11} = g_{22} = g_{33} = - \left[1 + 2 \left(\frac{a}{r} \right) + \dots \right]. \tag{71}$$

Comparison of Eqns. (70) - (71) with Eqns. (67) - (68) reveals that, to the order of expansion shown, the two metric tensors are identical.

We note from Eq. (70) that the sign of the contribution to the metric due to charge (represented by b) is counter to that of mass (represented by a). In the literature this effect is sometimes discussed in terms of an "electrogravitic repulsion" force [20].

VI. DISCUSSION

In overview, we have shown that a convenient methodology for investigating general relativistic (GR) effects in a non-abstract formalism is provided by the so-called polarizable-vacuum (PV) representation of GR. The PV approach treats metric perturbation in terms of a vacuum dielectric function K that tracks changes in the effective permittivity and permeability constants of the vacuum, a *metric engineering* approach, so to speak [21]. The structure of the approach is along the lines of the *THEμ* formalism used in comparative studies of gravitational theories. A summary of the important relationships derived here is presented in the accompanying Table.

The PV-derived matter-field Eqns. (58)-(59) are in principle applicable to a wide variety of problems. This short exposition, covering but the Schwarzschild and Reissner-Nordstrøm metrics and the three experimental tests of GR, is therefore clearly not exhaustive. Consideration was confined to cases of spherical symmetry [22], and important topics such as gravitational radiation and frame-dragging effects were not addressed. Therefore, further exploration and extension of the PV approach to specific problems of interest is encouraged, again with cross-referencing of PV-derived results to those obtained by conventional GR techniques to ensure that the PV approach does not generate spurious results.

With regard to the epistemology underlying the polarizable-vacuum (PV) approach as compared with the standard GR approach, one rather unconventional viewpoint is that expressed by Atkinson who carried out a study comparing the two [23]. "It is possible, on the one hand, to postulate that the velocity of light is a universal constant, to define 'natural' clocks and measuring rods as the standards by which space and time are to be judged, and then to discover from measurement that space-time, and space itself, are 'really' non-Euclidean; alternatively, one can *define* space as Euclidean and time as the same everywhere, and discover (from exactly the same measurements) how the velocity of light, and natural clocks, rods, and particle inertias 'really' behave in the neighborhood of large masses. There is just as much (or as little) content for the word 'really' in the one approach as in the other; provided that each is self-consistent, the ultimate appeal is only to convenience and fruitfulness, and even 'convenience' may be largely a matter of personal taste..."

On the other hand, as we have discussed in Section II, from the standpoint of what is actually measured with physical rods and clocks, the conventional approach captures such measurements in a concise, mathematically self-consistent formalism (the tensor approach), despite its somewhat formidable appearance to a neophyte. Therefore, the standard approach is more closely aligned with the positivist viewpoint that underlies modern scientific thought. Nonetheless, the PV model, with its intuitive, physical appeal, can be useful in bridging the gap between flat-space Newtonian physics and the curved-spacetime formalisms of general relativity.

ACKNOWLEDGEMENTS

I wish to express my appreciation to G. W. Church, Jr., for encouragement and useful suggestions in the development of this effort. I also wish to thank M. Ibison and E. Davis for stimulating discussions of the concepts presented herein.

Table
Metric Effects in the Polarizable Vacuum (PV) Representation of GR
(For reference frame at infinity, $K = 1$)

<i>Variable</i>	<i>Determining Equation</i>	$K \geq 1$ <i>(typical mass distribution, M)</i>
vel. of light $v_L(K)$	$v_L = c/K$ Eq. (7)	vel. of light $< c$
mass $m(K)$	$m = m_0 K^{3/2}$ Eq. (9)	effective mass increases
frequency $\omega(K)$	$\omega = \omega_0 / \sqrt{K}$ Eq. (10)	redshift toward lower frequencies
time interval $\Delta t(K)$	$\Delta t = \Delta t_0 \sqrt{K}$ Eq. (11)	clocks run slower
energy $E(K)$	$E = E_0 / \sqrt{K}$ Eq. (8)	lower energy states
length dim. $L(K)$	$L = L_0 / \sqrt{K}$ Eq. (13)	objects (spaces) shrink
dielectric-vacuum "grav." forces $F_K(K)$	$F_K(K) \propto \nabla K$ Eq. (58)	attractive grav. force

REFERENCES

1. C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973), p. 5.
2. H. A. Wilson, "An electromagnetic theory of gravitation," Phys. Rev. **17**, 54-59 (1921).
3. R. H. Dicke, "Gravitation without a principle of equivalence," Rev. Mod. Phys. **29**, 363-376 (1957). See also R. H. Dicke, "Mach's principle and equivalence," in *Proc. of the Intern'l School of Physics "Enrico Fermi" Course XX, Evidence for Gravitational Theories*, ed. C. Møller (Academic Press, New York, 1961), pp.1-49.
4. A. P. Lightman and D. L. Lee, "Restricted proof that the weak equivalence principle implies the Einstein equivalence principle," Phys. Rev. D **8**, 364-376 (1973).
5. C. M. Will, "Gravitational red-shift measurements as tests of nonmetric theories of gravity," Phys. Rev. D **10**, 2330-2337 (1974).
6. M. P. Haugan and C. M. Will, "Principles of equivalence, Eötvös experiments, and gravitational red-shift experiments: The free fall of electromagnetic systems to post--post-Coulombian order," Phys. Rev. D **15**, 2711-2720 (1977).
7. A. M. Volkov, A. A. Izmest'ev, and G. V. Skrotskii, "The propagation of electromagnetic waves in a Riemannian space," Sov. Phys. JETP **32**, 686-689 (1971). See also F. de Felice, "On the gravitational field acting as an optical medium," Gen. Rel. Grav. **2**, 347-357 (1971).
8. W. Heitler, *The Quantum Theory of Radiation*, 3rd ed. (Oxford University Press, London, 1954), p. 113.
9. R. A. Alpher, "Large numbers, cosmology, and Gamow," Am. Sci. **61**, 52-58 (Jan.-Feb. 1973).
10. E. R. Harrison, "The cosmic numbers," Phys. Today **25**, 30-34 (Dec. 1972).
11. J. K. Webb et al., "Search for time variation of the fine structure constant," Phys. Rev. Lett. **82**, 884-887 (1999).
12. J. W. Brault, "Gravitational red shift of solar lines," Bull. Amer. Phys. Soc. **8**, 28 (1963).
13. R. V. Pound and G. A. Rebka, "Apparent weight of photons," Phys. Rev. Lett. **4**, 337-341 (1960).
14. R. V. Pound and J. L. Snider, "Effect of gravity on nuclear resonance," Phys. Rev. Lett. **13**, 539-540 (1965).
15. R. N. Treuhaft and S. T. Lowe, "A measurement of planetary relativistic deflection," Astron. Jour. **102**, 1879-1888 (1991).

16. H. Goldstein, *Classical Mechanics*, (Addison-Wesley, Reading MA, 1957), pp. 206-207.
17. Ref. 1, p. 840.
18. For increasingly larger departures from flatness, however, the two approaches, although initially following similar trends, begin to diverge with regard to specific magnitudes of effects. In the PV approach the solutions for the metric tensor are exponential in form; in the conventional GR approach somewhat more complex. This discrepancy has shown up previously in other general curved-space approaches to GR as well. Of special interest is the so-called Einstein-Yilmaz form, in which Einstein's equations are modified by inclusion of the stress-energy tensor of the gravitational field itself on the R.H.S. in addition to the usual matter/field stress-energy. (For a review of the Einstein-Yilmaz formalism, see Y. Mizobuchi, "New theory of space-time and gravitation - Yilmaz's approach," Hadronic Jour. **8**, 193-219 (1985)). The Yilmaz modification yields exponential solutions in the form derived here by means of the PV approach. The Einstein-Yilmaz equations satisfy the standard experimental tests of GR, as well as addressing a number of mathematical issues of concern to general relativists, and are thus under study as a potentially viable modification to the original Einstein form. (See C. O. Alley, "The Yilmaz theory of gravity and its compatibility with quantum theory," in *Fundamental Problems in Quantum Theory: A Conference Held in Honor of Professor John A. Wheeler*, ed. D. M. Greenberger and A. Zeilinger (Vol. 755 of the *Annals of the New York Academy of Sciences*, New York, 1995), pp. 464-475.) Should further work in the development of general relativity lead to acceptance of the Einstein-Yilmaz form as the canonical form, then the PV-derived results would be applicable under strong-field conditions as well.
19. Ref. 1, p. 841.
- 20 S. M. Mahajan, A. Qadir and P. M. Valanju, "Reintroducing the concept of <>force<> into relativity theory," Il Nuovo Cim. **65B**, 404-417 (1981).
21. H. E. Puthoff, "SETI, the velocity-of-light limitation, and the Alcubierre warp drive: an integrating overview," Physics Essays **9**, 156-158 (1996).
22. However, it is known that the PV-related *THEμ* approach is sufficiently general that results obtained for spherically symmetric gravitational fields can be generalized to hold for nonsymmetric conditions as well.
23. R. d'E. Atkinson, "General relativity in Euclidean terms," Proc. Roy. Soc. **272**, 60-78 (1962).




Fig. 1(a)




Fig. 1(b)

Fig. 1. Geometry of bending of light rays

“Anti-Gravity Personae”



Yevgeny Podkletnov at the Tampere Lake, Finland




Podkletnov at the Tampere University of Technology



Yevgeny Podkletnov



Dr. Giovanni Modanese



Whitt Brantley - Chief of the Advanced Concepts Office at NASA MSFC



Ron Koczor - Chief Engineer and Project Manager for DeltaG, NASA MSFC



Tony Robertson, Staff Engineer for DeltaG, NASA MSFC




John Schnurer



Dr. Ning Li and two members of the NASA MSFC team, Neil Tyson and Jerald Oakly



Dr. Noever with large cryostat, NASA



NASA team - from left: Tony Robertson, Neil Tyson, Whitt Brantley, David Noever, Jerald Oakly, Ronald Koczor, and Ning Li

Disclaimer

The author of this document makes no representation about the suitability of the information presented in this document for any purpose. All information is presented "As Is" without warranty of any kind. In no event shall the author of this document or his representatives be liable for any direct or indirect or consequential damages resulting from use or lack of use of this document or the information contained in this document, in any form what so ever.

This document contains information entered by third parties as articles and graphics for their own use while interacting with other readers. The author of this document will not be responsible or liable for any of the language used in such entries nor does the author of this document make any representations about the truthfulness of said entries.

Copyright

All information presented on this site, in whole or in part, may not under any circumstances be copied, reverse engineered, resold, redistributed or used on an Internet site for purposes of any kind without prior written permission from respective copyright holders. Any rights not expressly granted herein are reserved. This copyright notice is intended to protect the rights of copyright holders other than myself. For accepted use of material prepared by Venik, please consult the Copyright Fair Use Clause below. Please keep in mind that, depending on your geographical location, neither Russian nor the US copyright regulations may apply, so you may need to consult your local laws and regulations before using or

distributing this document. This document resides on a Web server in Russia and is protected under the Intellectual Property and Associated Rights Law of the Russian Federation, Section II: *Intellectual Property Rights*, Article 19: *Use of a work without the consent of the author and without monetary compensation*, Paragraph 2.

Copyright Fair Use Clause

US Copyright Law

Material in this document has been (a) created by the document author, Venik, or (b) assembled from a variety of Internet web sites and been placed here for ease and rapidity of local access, or, (c) for small amounts of text, derived from printed material.

A good faith effort has been made to comply with US copyright law. This does not mean that none of the material is copyright, but that the "fair use" clause of US Copyright Law and the Section II, Article 19 of the Intellectual Property and Associated Rights Law of the Russian Federation has been adhered to. The "fair use" clause, as explicated at the University of Pennsylvania web site (<http://www.upenn.edu/>), is given below. In particular, any copyright material used here is (a) not used for commercial gain and used exclusively for educational purposes; and (b) used in limited amounts in comparison to the published source. The latter condition is not required under Section II, Article 19, Paragraph 2 of the Intellectual Property and Associated Rights Law of the Russian Federation.

If there are any objections that material placed here does not conform to the "fair use" provisions outlined, contact Venik (venik4@comcast.net) and material will be removed immediately pending resolution of

the issue. As for material prepared by Venik, permission is given to use freely in electronic form, and in print at educational institutions, as long as source and author are indicated.

"107: Limitations on exclusive rights: Fair Use Notwithstanding the provisions of section 106, the fair use of a copyrighted work, including such use by reproduction in copies or phone records or by any other means specified by that section, for purposes such as criticism, comment, news reporting, teaching (including multiple copies for classroom use), scholarship, or research, is not an infringement of copyright. In determining whether the use made of a work in any particular case is a fair use, the factors to be considered shall include:

1. The purpose and character of the use, including whether such use is of a commercial nature or is for nonprofit educational purposes;
 2. The nature of the copyrighted work;
 3. The amount and substantiality of the portion used in relation to the copyrighted work as a whole; and
 4. The effect of the use upon the potential market for or value of the copyrighted work.
- (added pub. I 94-553, Title I, 101, Oct 19, 1976, 90 Stat 2546)"

Intellectual Property and Associated Rights Law of the Russian Federation

The following is an English translation of the excerpt from the Intellectual Property and Associated Rights Law of the Russian Federation, Section II: Intellectual Property Rights, dealing with acceptable use of published works without the consent of the author and without monetary compensation.

Article 19. Use of a work without the consent of the author and without monetary compensation

Without the consent of the author and without monetary

compensation, but with mandatory indication of the name of the author whose work is being used and the name of the source the following is permitted:

1) quoting of the original and in translation in scientific, research, polemic, analytical, and informational purposes from legally published materials in the amount justified by the purpose of quoting, including reproduction of excerpts from newspapers and magazine articles in the form of press reviews;

2) use of legally published materials and excerpts from them as illustrations in publications, sound and video recordings of educational nature in the amount justified by the established goal;

3) reproduction in newspapers, broadcasting via radio or cable for the purpose of public awareness of legally published in newspapers or magazines information dealing with current economic, political, social and religious subject matters or broadcast information of similar nature in cases when such reproduction, broadcasting via radio or cable were not explicitly prohibited by the author;

4) reproduction in newspapers, broadcasting via radio or cable for the purpose of public awareness of public political speeches, addresses, reports and other similar materials in the amount justified by informational purposes. At the same time the author retains the right to publish such materials as collected works;

5) reproduction or transmission in reviews of current events by means of radio or cable broadcasting for the purpose of public awareness of materials that were seen or heard in the course of such events in the amount justified by informational purposes. At the same time the author retains the right to publish such materials as collected works;

6) reproduction of legally published materials on non-profit basis by means of raised dot fonts or using other special means for blind, except for works specifically created for such methods of publication.

The following is the excerpt from the Intellectual Property and Associated Rights Law of the Russian Federation, Section II: Intellectual Property Rights, dealing with acceptable use of published works without the consent of the author and without monetary compensation.

Статья 19. Использование произведения без согласия автора и без выплаты авторского вознаграждения

Допускается без согласия автора и без выплаты авторского вознаграждения, но с обязательным указанием имени автора, произведение которого используется, и источника заимствования:

1) цитирование в оригинале и в переводе в научных, исследовательских, полемических, критических и информационных целях из правомерно обнародованных произведений в объеме, оправданном целью цитирования, включая воспроизведение отрывков из газетных и журнальных статей в форме обзоров печати;

2) использование правомерно обнародованных произведений и отрывков из них в качестве иллюстраций в изданиях, в радио- и телепередачах, звуко- и видеозаписях учебного характера в объеме, оправданном поставленной целью;

3) воспроизведение в газетах, передача в эфир или сообщение по кабелю для всеобщего сведения правомерно

опубликованных в газетах или журналах статей по текущим экономическим, политическим, социальным и религиозным вопросам или переданных в эфир произведений такого же характера в случаях, когда такие воспроизведение, передача в эфир или сообщение по кабелю не были специально запрещены автором;

4) воспроизведение в газетах, передача в эфир или сообщение по кабелю для всеобщего сведения публично произнесенных политических речей, обращений, докладов и других аналогичных произведений в объеме, оправданном информационной целью. При этом за автором сохраняется право на опубликование таких произведений в сборниках;

5) воспроизведение или сообщение для всеобщего сведения в обзорах текущих событий средствами фотографии, путем передачи в эфир или сообщения для всеобщего сведения по кабелю произведений, которые становятся увиденными или услышанными в ходе таких событий, в объеме, оправданном информационной целью. При этом за автором сохраняется право на опубликование таких произведений в сборниках;

6) воспроизведение правомерно обнародованных произведений без извлечения прибыли рельефно - точечным шрифтом или другими специальными способами для слепых, кроме произведений, специально созданных для таких способов воспроизведения.

[Search Results](#) [New Search](#)

1 record selected from **Compendex** for (((London equations for superconductors) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or email results.

1. **Accession Number:** 9

Title: Modelling of thermomagnetoelectric effects in type-II superconductors in the mixed state

Author(s): Zhou, Shu-Ang;

First author affiliation: Ericsson Telecom AB ,Stockholm ,Sweden

Serial Title: Physica C: Superconductivity

Abbreviated Serial Title: Phys C Supercond

Volume: v 228

Issue number: n 1-2

Issue date: Jul 10, 1994

ISSN: 0921-4534

CODEN: PHYCE6

Publisher: Publ by Elsevier Science Publishers B.V., Amsterdam, Neth

Publication year: 1994

Pages: p 122-128

Abstract: This article presents a phenomenological theory for studying thermomagnetoelectric effects in type-II superconductors in the mixed state. In the theoretical model, the anisotropic effect, the vortex dynamic effect, the flux-flow Hall effect and the thermomagnetic effects in the type-II superconductor in the mixed state are all taken into account self-consistently. It is shown that, at the London approximation, the classical first and second London equations for superconductors in the Meissner state are modified for the superconductors in the mixed state due to vortex dynamic effects, which are, in general, nonlinear. At the linear approximation, effective constitutive equations and field equations are derived to describe the thermomagnetoelectric behavior of anisotropic type-II superconductors in the mixed state. It is shown that the derived field equations are, in general, coupled due to the vortex dynamic effects. A possible solution of a plane thermomagnetic wave in the type-II superconductor in the mixed state is also studied illustratively.

Abstract type: (Author abstract)

Number of references: 27 Refs.

Main heading: Superconducting materials

Controlled terms: Mathematical models; Anisotropy; Hall effect; Thermal effects; Electric field effects; Approximation theory; Equations of state; Nonlinear equations

Uncontrolled terms: Thermomagnetoelectric effects; Phenomenological theory; London approximation; Vortex dynamics effects

Treatment type: Theoretical;

Classification codes: 708.3; 921.6; 931.2; 701.2; 641.1; 701.1

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

[Search Results](#)

[New Search](#)

Results Manager

Select Results Format

Citation

Abstract / Links

Detailed Record / Links

[New Display](#)

[Print](#)

[E-mail](#)

[Save/Download](#)

[Clear Selected Records](#)

[Privacy Policy](#) | [Contact Us](#) | [Feedback](#)

Copyright © 2000 - 2001 by Engineering Information Inc., Hoboken, New Jersey, U.S.A.

[Search Results](#) [New Search](#)

1 record selected from **Compendex** for (((Gravitoelectric) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or email results.

1. **Accession Number:** 7

Title: Local gravitoelectromagnetic effects on a superconductor

Author(s): [Agop, M.](#); [Buzea, C.Gh.](#); [Nica, P.](#);

First author affiliation: Technical Univ `Gh. Asachi' ,Iasi ,Rom

Serial Title: Physica C: Superconductivity and its Applications

Abbreviated Serial Title: Phys C Supercond Appl

Volume: v339

Issue number: n2

Issue date: Oct, 2000

ISSN: 0921-4534

CODEN: PHYCE6

Publisher: Elsevier Science Publishers B.V., Amsterdam, Netherlands

Publication year: 2000

Pages: p 120-128

Abstract: Maxwell's and London's generalized equations, generalized Meissner effect and gravitational shielding in an electromagnetic field are obtained. In such a context, we show that a neutral particle beam in an infinitely thin superconducting cylinder placed in the Earth's gravitoelectric field is focused, and that the penetration depth increases with the pulsation of the electromagnetic field.

Abstract type: (Author abstract)

Number of references: 25 Refs.

Main heading: Superconducting materials

Controlled terms: Gravitational effects; Electromagnetic dispersion; Maxwell equations; Electromagnetic shielding

Uncontrolled terms: Maxwell-London equations; Meissner effects; Gravitoelectromagnetic effects

Treatment type: Experimental;

Classification codes: 708.3; 931.5; 711.1; 701.1; 701.2; 921.2

Language: English.

Document type: Journal Articles

Database: Compendex

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

Search Results

New Search

Results Manager

Select Results Format

Citation

Abstract / Links

Detailed Record / Links

[New Display](#)

[Print](#)

[E-mail](#)

[Save/Download](#)

[Clear Selected Records](#)

[Privacy Policy](#) | [Contact Us](#) | [Feedback](#)

Copyright © 2000 - 2001 by Engineering Information Inc., Hoboken, New Jersey, U.S.A.

[Search Results](#) [New Search](#)

1 record selected from **Compendex** for (((Gravitoelectric) WITHIN ALL))

Use the [Results Manager](#) at the bottom of this page to save, print or email results.

2. **Accession Number:** 4133022

Title: Some physical implications of the gravitoelectromagnetic field in fractal space-time theory

Author(s): [Agop, M.](#); [Griga, V.](#); [Buzea, C.Gh.](#); [Petreus, I.](#); [Buzea, C.](#); [Marin, C.](#); [Rezlescu, N.](#);

First author affiliation: Technical 'Gh. Asachi' Univ ,Iasi ,Rom

Serial Title: Australian Journal of Physics

Abbreviated Serial Title: Aust J Phys

Volume: v51

Issue number: n1

ISSN: 0004-9506

CODEN: AUJPAS

Publication year: 1998

Pages: p 9-19

Abstract: It is shown that in terms of the fractal space-time theory the gravitoelectric potential is responsible for the quantization of the planetary and binary galaxy motions. On a cosmic scale a homogeneous gravitomagnetic field allows not only an ordering of the Universe, but a 'global' redshift quantization of galaxies as well.

Language: English.

Document type: Journal Articles

Database: Compendex

Compilation and Indexing Terms, ©2002, Elsevier Engineering Information, Inc.

Full Text Options

[Access Electronic Full-Text](#) | [Order Document from Linda Hall Library](#)

[Search Results](#) [New Search](#)[Results Manager](#)

Дата 20020805 Дата загрузки 20020805 Время загрузки 1727 Источник Журнал Итоги, Москва Заголовок **АНТИГРАВИТАЦИОННЫЙ ДВИГАТЕЛЬ** Автор Светлана Сухова Тема США Выпуск 31 Текст

"Отменить" закон земного притяжения ученые мечтали давно, вот только воплотить фантазию в жизнь никому не удавалось. Дошло до того, что антигравитация стала считаться абсурдной идеей, чем-то сродни вечному двигателю. Однако на днях представители американской компании Boeing заявили, что корпорация заинтересовалась разработками российского ученого Евгения Подклетнова в области антигравитации и намерена спонсировать дальнейшие изыскания в рамках программы Phantom Works. Американская компания далеко не первопроходец в этой области: до нее реализацией аналогичной программы Project Greenglow занималось оборонное подразделение британской компании BAE Systems. Но ближе всех к реальному воплощению антигравитационной мысли подошло НАСА. Американское аэрокосмическое агентство в 1997 году попыталось повторить эксперимент Подклетнова, но безуспешно. Однако на этом исследования не закончились, в лабораториях НАСА строят новую копию антигравитационного аппарата.

Еще в 1992 году Подклетнов опубликовал в специализированном журнале *Physica C* статью, в которой утверждал, что ему удалось "оградить" область пространства от сил гравитации с помощью охлажденного намагниченного кольца из сверхпроводящего материала. Пропуская через конструкцию импульсный электроток, Подклетнов добился, чтобы вес объекта, размещенного над вращающимся кольцом, уменьшился на 2 процента. Он также установил, что чем быстрее вращение, тем сильнее сокращается сила гравитации.

Понятно, почему исследованиями Подклетнова заинтересовались представители именно аэрокосмической отрасли: любой летательный аппарат, помещенный над таким гигантским кольцом, оторвется от Земли без ракетоносителя. С самолетом сложнее - понадобится нечто вроде сверхпроводящих намагниченных трасс. В общем, если опыты Подклетнова удастся повторить, то в ближайшем будущем ученым предстоит решить еще две задачи: как управлять объектом в антигравитационном поле и, что еще важнее, как остановить его движение.

Дата 20020801 Дата загрузки 20020731 Время загрузки 1800 Источник Комсомольская правда, Москва Заголовок **АМЕРИКАНЦЫ СОБИРАЮТСЯ ПРЕОДОЛЕТЬ ГРАВИТАЦИЮ** Выпуск 137 Текст

И сделают они это с помощью нашего ученого

Американская аэрокосмическая компания "Boeing" заинтересовалась открытием российского ученого Евгения Подклетнова, которое может помочь преодолеть силу гравитации.

На объекте компании в Сиэтле, который занимается наиболее сложными и перспективными проектами, научные результаты работы Подклетнова всерьез изучаются для их практического применения, сообщает Би-би-си.

Еще в 1992 году Подклетнов и его финский коллега Вуоринен провели в университете города Тампера опыты, результаты которых стали сенсацией. Ученые, помещая предметы в центр сверхпроводящего керамического диска, вращающегося с высокой скоростью, обнаружили, что их вес уменьшается на 2 процента!

Однако в те времена к пока не объяснимому открытию отнеслись скептически, и авторы даже не смогли сразу опубликовать статью о нем в научных журналах.

Теперь, должно быть, в Сиэтле "доведут до ума" наработки россиянина, и американские объекты, выводимые в космос, смогут сильно терять в весе, что сделает их запуск значительно дешевле.

Евгений УМЕРЕНКОВ. (Наш соб. корр.). Нью-Йорк.

Дата 20020731 Дата загрузки 20020730 Время загрузки 1727 Источник Труд, Москва Заголовок **ЗЕМНОЕ ПРИТЯЖЕНИЕ БУДЕТ ПОБЕЖДЕНО?. ОТКРЫТИЕ**

РОССИЙСКОГО УЧЕНОГО ЗАИНТЕРЕСОВАЛО "БОИНГ" Автор Сергей БАЗАВЛУК. Тема МИР СЕГОДНЯ Выпуск 131 Текст

В условиях строжайшей секретности крупнейшая в мире авиастроительная компания "Боинг" занимается разработкой так называемого антигравитационного эффекта. Ее специалисты решили воспользоваться разработками российского ученого десятилетней давности. О начале работы по преодолению силы земного притяжения сообщил глава сверхсекретной лаборатории "Боинга" Джордж Мильнер. "Фантом уоркс", так называется этот тщательно охраняемый исследовательский центр в Сиэтле, занимается инновациями в области самолетостроения.

Специалисты, работающие в лаборатории "Боинга", решили воспользоваться наработками российского ученого Евгения Подклетнова, десять лет назад создавшего в технологическом университете финского города Тампер антигравитационное устройство. По заявлению конструктора, изобретение способно ослаблять воздействие земного притяжения на различные объекты. Устройство, в частности, можно использовать и в самолетостроении. При этом будет значительно сокращен расход энергии, затрачиваемой на полет крылатой машины.

Суть открытия заключается в том, что объекты, расположенные над керамическим сверхпроводниковым диском, вращающимся под воздействием мощных электромагнитов, теряют вес. Уменьшение силы гравитации, правда, было небольшим - всего около 2% - но последствия открытия могут быть весьма заметными.

В 1992 году об этом как-то вяло написали некоторые газеты, но идея преодоления земного притяжения была замечена американцами. Исследованием такой возможности занялась НАСА. Вместе с началом работы космического агентства по воссозданию антигравитации всяческое упоминание об открытии российского ученого и вовсе пропало из СМИ.

В данный момент, как стало известно из авторитетного журнала "Джейнс дефенс уикли", над открытием работают как минимум три серьезные организации. Это "Боинг", который после десяти лет изучения материалов эксперимента Евгения Подклетного пришел к выводу, что данный эффект вполне возможен. Это британская группа специалистов по высоким технологиям компании "BAE Systems", проводящая аналогичные исследования. И, конечно, это НАСА.

Несмотря на столь авторитетный сонм ученых, работающих над этой проблемой, у открытия есть и противники. Некоторые специалисты с подозрением относятся к возможности победить силу земного притяжения. Они, в частности, считают, что эксперимент российского ученого проводился некорректно и отрицательной гравитации не существует. Тем не менее работа продолжается. Ведь в случае победы над земным притяжением удастся решить многие проблемы, одна из которых - участившиеся авиакатастрофы и десятки невинно погибших людей.

Дата 20020731 Дата загрузки 20020730 Время загрузки 2345 Источник Ведомости, Москва Заголовок **Boeing готовит революцию Автор Ник Кул Выпуск 133 Текст**

Тема антигравитации, находившаяся далеко за пределами интересов инженеров аэрокосмической области, возможно, вскоре станет чем-то более привычным и респектабельным. На этой неделе стало известно, что корпорация Boeing, крупнейший в мире производитель самолетов, ведет исследования, результаты которых могут перевернуть наши представления о перемещении в пространстве.

В руки журналистов отраслевого издания JaneXs Defence Weekly попал внутренний документ Boeing, связанный с исследовательским проектом GRASP ("Исследования гравитации для усовершенствованного перемещения в пространстве"). "Если управление гравитацией окажется возможным, это произведет революцию во всей аэрокосмической индустрии", - говорится в документе. Это слишком скромное утверждение. Ведь если бы энергию гравитации можно было как-то использовать, это означало бы революцию во всем

устройстве мира - достаточно сказать, что весь транспорт можно будет перевести на двигатели нового типа, использующие вместо горючего энергию гравитационного поля.

Документ по проекту GRASP ссылается на работу русского ученого Евгения Подклетнова, который, как он утверждает, разработал систему, ослабляющую гравитационное поле. В 1992 г. Подклетнов, работавший в Технологическом университете г. Тампере в Финляндии, направил в британский физический журнал статью с описанием действия системы, основанной на сверхпроводимости - способности некоторых веществ утрачивать электрическое сопротивление при температурах, близких к абсолютному нулю. По утверждению ученого, объекты, помещенные над быстро врачающимся "сверхпроводником", теряют 2% своего веса.

Физики подвергли Подклетнова остракизму, но работой ученого заинтересовались в NASA. У этой организации уже были материалы Нин Ли, исследовательницы из Университета Алабамы, которая якобы научилась создавать поле, сходное с гравитационным и позволяющее притягивать или отталкивать объекты, используя опять-таки врачающийся сверхпроводящий диск.

В середине 1990-х исследователи из Центра космических полетов NASA в Алабаме безуспешно пытались воспроизвести эксперименты Подклетнова. Однако они сочли, что для получения положительных результатов им не хватает уникальной технологии изготовления сверхпроводящих дисков, примененной русским ученым.

В 1999 г. NASA заплатила \$600 000 компании Superconductive Components of Columbus за изготовление дисков, подобных подклетновским. Сам Подклетнов был нанят в качестве консультанта. Эксперимент был отложен, но Рон Кошор, руководитель проекта NASA, рассчитывает осуществить его до конца этого года. Тем временем Подклетнов, работающий уже в Москве, занят развитием своей теории. В прошлом году он опубликовал детальное описание работы "импульсного гравитационного генератора", якобы способного отталкивать любые объекты. Генератор использует мощный электрический разряд и сверхпроводящий излучатель, с помощью которых, по словам Подклетнова, создается гравитационный импульс, сохраняющийся в течение очень короткого времени и распространяющийся с огромной скоростью (практически мгновенно). Импульс "проходит сквозь различные объекты без видимых потерь энергии". Этот луч оказывает на все встречающиеся объекты отталкивающее воздействие, пропорциональное их массе. Утверждается, что луч воздействует таким образом на объекты, удаленные более чем на километр. Именно эта работа привлекла внимание Boeing. В документе по GRASP также описан луч, способный проникать сквозь любые среды и невосприимчивый к каким-либо электромагнитным полям. (FT, 29. 07. 2002)

Дата 20020730 Дата загрузки 20020729 Время загрузки 2130 Источник Известия, Москва Заголовок "**Боинг**" интересуется "хозяином" гравитации Автор Сергей ЛЕСКОВ Выпуск 132 Текст

Крупнейшая американская авиастроительная компания Boeing приступила к изучению экспериментов российского ученого Евгения Подклетнова, которые уже 10 лет являются предметом бурных научных дискуссий. Евгений Подклетнов утверждает, что создал антигравитационное устройство, которое уменьшает силу тяжести. Проект под кодовым названием "Захват" запущен в секретном отделе Phantom Works, занимающемся наиболее сложными разработками. Глава отдела Джордж Милнер заявил, что разработка внушает ему доверие.

В 1988 году Подклетнова для работы над диссертацией пригласили в университет финского города Тампере. За плечами у него Химико-технологический институт имени Менделеева и 15 лет работы в Институте высоких температур РАН. В Тампере Подклетнов привез 30-санитметровый диск из расплава оксидов меди, иттрия и бария, который является его "ноу-хай". Если этот диск охладить до температуры сверхпроводимости и раскрутить электромагнитами, то, как установил Подклетнов, все предметы, помещенные над диском

(будь они из железа, бумаги, стекла и проч.), становятся легче на 2%. Это переворачивает науку: согласно общей теории относительности создать экран от гравитации невозможно.

Подклетнов, будучи ученым с серьезной школой, понимал, что рискует оказаться в стане лженауки. Но эксперименты раз за разом повторяли результат. Он в полной мере познал все злоключения, которые испытывают непризнанные гении. Под давлением официальной науки (а она существует и на Западе) перед ним закрывались двери журналов, от статей отказывались соавторы, а из университета изобретателю пришлось уволиться.

Как часто бывает, спорными исследованиями заинтересовался располагающий большими деньгами военно-промышленный комплекс, на эпизодические гранты которого ученый теперь и существует. Компания Superconductive Systems получила от НАСА 70 тысяч долларов на опыт с использованием 8 спутников. Эффект не наблюдался, но, как заявил побывавший в Центре космических исследований автор изобретения, диск был изготовлен неправильно, а скорость его вращения оказалась слишком мала. В ряде университетов США и Западной Европы получено подтверждение его результатов. Недавно подобные опыты начали в военном подразделении английской компании BAE Systems.

- Бред, - академик Виталий Гинзбург категоричен. - Взаимодействие электромагнитного и гравитационного полей возможно, но оно очень слабое. Хотя я точно не знаю, что он измерял и в чем ошибся, уверен, все это полная чепуха. Типичная лженаука.

- На Западе выделяют гранты на телепатию, но из этого не следует, что она существует, - сказал заместитель директора Института физических проблем РАН Генрих Богомолов. - В лучших лабораториях эти эффекты не наблюдаются, а Подклетнов их получает "на коленке". Что-то подобное может происходить в ядрах звезд, но не на Земле.

Наши попытки найти изобретателя успехом не увенчались. Из университета Тампере пришел ответ, что Евгений Подклетнов в штате давно не числится, а посещение лабораторий ему запрещено...

Дата 20020729 Дата загрузки 20020730 Время загрузки 0002 Источник Lenta.Ru, Москва Заголовок **КОРПОРАЦИЯ Boeing И РУССКИЙ ФИЗИК СОВМЕСТНО ПРЕОДОЛЕВАЮТ ГРАВИТАЦИЮ** Тема В МИРЕ Примечания <http://lenta.ru/world/2002/07/29/experiment/> Текст

Корпорация [<http://www.boeing.com/>] Boeing заинтересовалась исследованиями российского физика Евгения Подклетнова в области антигравитации, сообщает BBC News со ссылкой на [<http://jdw.janes.com/>] Jane's Defence Weekly.

В 1992 году, в ходе [<http://ufolog.nm.ru/ag2.htm>] эксперимента, проведенного в Технологическом университете Тампере в Финляндии, Подклетнов смог добиться потери веса предмета помещенного над высокотемпературным сверхпроводящим керамическим диском, вращающимся в электромагнитном поле. Общее сокращение силы тяжести при этом было сравнительно мало и достигало лишь 2-х процентов.

Подклетнов и его коллеги считают, что нашли способ "заслониться" от земного притяжения, но традиционная наука отнеслась к этим исследованиям с некоторым недоверием.

Со своей стороны, специалисты корпорации Boeing весьма заинтересовались результатами Подклетнова. Его разработками занялись в секретных лабораториях Phantom Works в Сиэтле. Гипотезу Подклетного пытаются применить в рамках новой разработки корпорации Boeing, получившей название Project Grasp.

Напомним, что исследования Подклетнова заинтересовали не только Boeing. Над ними также работают английская корпорация BAe Systems в рамках военных исследований Project Greenglow и корпорация NASA. Впрочем, специалисты NASA сообщают, что не могут добиться того же эффекта.

ССЫЛКИ ПО ТЕМЕ

[<http://news.bbc.co.uk/1/hi/sci/tech/2157975.stm>] Boeing tries to defy gravity - BBC News, 29.07.2002

"Корпорация Boeing и русский физик совместно преодолевают гравитацию", 30.07.2002
<http://lenta.ru/world/2002/07/29/experiment/>
(C) 2002 Lenta.Ru/В МИРЕ

Дата 20020418 Дата загрузки 20020513 Время загрузки 1257 Источник Алфавит, Москва Заголовок **ВЕЧНОСТЬ НА ПРОДАЖУ** Автор Леонид Попов Выпуск 16 (178) Текст

Официальная наука то и дело натыкается на природные явления, не стыкующиеся с господствующими представлениями физики. Каким образом форель умудряется почти неподвижно стоять в горном потоке, бегущем со скоростью десятки метров в секунду? Нет однозначного объяснения скорости, которую развивают в воде дельфины. А майский жук с точки зрения классической аэrodинамики вообще не мог бы подняться в воздух. Возможно, мы подошли к той черте, когда эксперимент, опыт, действующий прибор будет двигать прогресс человечества, не дожидаясь, пока подтянется теоретический арьергард.

Февраль, ВВЦ, выставка "Двойные технологии". Народ клубится вокруг малопримечательного железного бака с изогнутыми трубами. Кто-то записывает координаты автора - Юрия Потапова, кто-то растерянно пожимает плечами. Особо грамотные, ползая на коленках, заглядывают под установку в поисках обмана - скрытого источника тепла, замаскированного провода. Увы! У всех на глазах происходит явное научное безобразие: холодная вода, закачиваемая электронасосом в систему трубок "теплогенератора Потапова", пройдя по ним, почему-то нагревается до 70 градусов. Просто от механического движения. То есть дополнительная энергия (на 1 кВт потребляемого из сети электричества установка выдаёт около 2 кВт тепла) появляется "ниоткуда". Словно не зная, что это категорически запрещено вторым законом термодинамики.

Особую пикантность происходившему придаёт то обстоятельство, что один из организаторов выставки - РАН, уже объявившая установку Потапова очередным "вечным двигателем", то есть шарлатанством.

Вердикт вынесен известным академиком Эдуардом Кругляковым, уже несколько лет возглавляющим академическую "комиссию по борьбе с лжен наукой" и издавшим на тему "лжеучёных" несколько разоблачительных книжек.

Юрий Семёнович Потапов, доктор технических наук из Молдавии, ныне академик РАЕН, с академиком Кругляковым и не думал спорить. Человек практичный, обладающий к тому же кое-какими инженерными связями, он просто... наладил серийный выпуск еретической установки. Сначала в Молдавии на фирме "Юсмар", потом и в России - на оборонном заводе им. В.А. Дегтярёва.

Естественно, перед этим странный теплогенератор прошёл придирчивую экспертизу: в небезызвестном РКК "Энергия" в г. Королёве, самарском ЦСКБ, киевском НПО "Холод". А под занавес и в Национальном ядерном центре в Лос-Аламосе. За своё изобретение Потапов получил в США премию в области инноваций - "Факел Бирмингема", его генератор запатентован в 42 странах мира. В России он давно обогревает частные коттеджи, цеха и склады от Калининграда до Владивостока.

Так что же всё-таки придумал этот "псевдоизобретатель", как ему удалось "обшарлатанить" такое количество людей?

Сам Юрий Семёнович поясняет, что лишь довёл до ума опыты, начатые задолго до него. Ещё в 20-х годах XX века французский инженер Ж. Ранке изобрёл вихревую трубу. Насос закачивает в неё воздух через наклонный канал так, чтобы он закручивался вдоль оси трубы. Воздух расслаивается на два потока: один - горячее исходного, второй - холоднее. Внятного объяснения этому эффекту нет до сих пор. Несколько лет назад Потапов задал себе

простой вопрос, никому почему-то ранее в голову не приходивший: что если в трубу Ранке запустить воду? Оказалось, водный поток тоже разделяется на разные температурные составляющие.

Но вот загадка - даже "холодная" часть имела температуру выше, чем вода на входе. Закон сохранения энергии трещал по швам!

Юрий Семёнович чисто опытным путём пришёл к хитрой системе решёток, конусов и завихрителей, стабильно "нагревающих" проходящую через них воду до 50-70 градусов за счёт "чего-то". Так появился "теплогенератор Потапова", невозможный с точки зрения традиционной физики.

К слову, сам изобретатель уверен, что закон сохранения остался незыблёмым. Просто им найден новый источник энергии... в самой воде.

Вскоре подоспели и теоретики. Академик РАН Леонид Фоминский подвёл под работающий де-факто аппарат следующую теорию. В книге, выпущенной в соавторстве с Потаповым, Фоминский доказывает (кстати, основываясь на теории относительности Эйнштейна), что во всяком теле при его раскручивании часть массы должна превращаться в энергию. Это первая составляющая "излишков" тепла в установке. Далее в ход идут пресловутые "торсионные поля", категорически не признаваемые РАН. Эти самые поля вращения, по мнению Фоминского, стимулируют... холодный ядерный синтез в воде.

По словам исследователей, в одном из опытов под нагревательной установкой установили дозиметр, и тот показал превышение гамма-излучения над естественным фоном в полтора раза.

Час от часу не легче! Холодный термояд уже многоократно открывали и закрывали по всему миру. Была, помнится, "буря в стакане воды" американских учёных Понса и Флейшмана. А теперь вот... "буря в батарее отопления".

При всей внешней анекдотичности ситуации у "теплогенератора Потапова" есть существенное отличие и от спорных лабораторных установок учёных, и от проектов мечтателей-Кулибиных. Он работает! Так или иначе, но вода прямо через стенки греет воздух в помещении. В последних опытных образцах тепловой энергии на выходе получается в десять раз больше, чем берёт насос из электросети. При такой разнице, усмехается Потапов, трудновато будет списать результат на ошибку измерений...

Среди благодарных покупателей бытового "термоядерного реактора" оказались даже... монахи из Свято-Вознесенского Ново-Нямецкого монастыря в с. Кицканы (Молдавия). Для них-то чудо - понятие привычное. Правда, в случае с отоплением и верующие, и атеисты, и учёные, и неучи - все поступают до безобразия прозаично: считают деньги, затраченные на электричество и топливо до и после приобретения установки. И когда выводят баланс, посыпают подальше разоблачительный пафос академиков.

Так что же это: случай массового помешательства или слепота официальной науки, упорно не желающей замечать открытие века?

Блаженный Августин метко заметил: "Чудо - не то, что противоречит законам природы, а то, что противоречит нашему знанию этих законов".

Эдуарду Павловичу Круглякову, конечно, можно посочувствовать. В России-матушке сегодня действительно перебор с патологически активными оккультистами, равно как и с циничными шарлатанами, прикрывающимися научообразной лексикой. Хватает и сумасшедших. А ещё есть - как бы поточнее выразиться - фантазёры. Допустим, результаты экспериментов удивительные, а теоретическая база под них подводится невнятная, неудачная. С такими тоже бороться? Но без них наука совсем закиснет в своих "правильных" тупиках!

Понятно, США и Канада, финансирующие наших изобретателей-«еретиков», для нас не указ. Недавно канадский бизнесмен, глава фирмы "Агримекс Интернэшнл" Василиос Кацабанис подписал контракт на сумму 168 млн долларов с волгоградским НПЦ "ГРУС". На основе открытия академика РАН Валериана Соболева эта фирма наладила производство мини-электростанций, вырабатывающих ток из... окружающего пространства. На вердикт

уважаемых академиков РАН по поводу соболевского открытия - "обман или ошибка" - канадский воротила внимания не обратил. Ещё раньше НАСА профинансировало опыты изгоя большой российской науки Евгения Подклетнова по экранированию гравитации. А ныне, по слухам, Национальное агентство по аэрокосмическим исследованиям занимается этим уже самостоятельно и за закрытыми дверями... Комиссии по лженауке на них нет!

Леонид Попов

Справка "Алфавита"

С теорией, объясняющей "лишнюю" энергию в установках Потапова, вы можете ознакомиться в книге "Энергия вращения"; Ю.С. Потапов, Л.П. Фоминский, С.Ю. Потапов, Кишинёв, 2001.

Противоположную точку зрения вы найдёте в публикациях председателя комиссии РАН по борьбе с лженаукой академика Эдуарда Круглякова - "Что же с нами происходит?" и "Учёные с большой дороги" - www-psb.ad-sbras.nsc.ru/krugl1w.htm

Критика деятельности Э. Круглякова в статье "Борьба с "лженаукой" как способ подавления научного инакомыслия?" На www.membrana.ru

Современные вечные двигатели, проекты получения энергии из вакуума, ссылки на патенты - www.adventure.dv.ru/project/index.htm

Дата 20020211 Дата загрузки 20020221 Время загрузки 0020 Источник Utro.Ru, Москва
Заголовок **ПОЛЕТЫ НЕ ВО СНЕ, А НАЯВУ** Автор Вадим САМОКАТОВ Тема
АНАЛИТИКА:ЖИЗНЬ Выпуск 388 (710) Примечания
<http://www.utro.ru/articles/2002021102065461204.shtml> Текст

Однажды я собственными глазами видел, как человек взлетел. Точнее - оторвался от пола и на несколько секунд повис в воздухе. Произошло это в середине девяностых годов на выступлении одного довольно популярного в то время колдуна. Был ли это трюк, какие проделывает на своих концертах Дэвид Копперфильд, или колдун действительно умел летать, я не знаю (склоняюсь все-таки к мысли, что я стал свидетелем фокуса), но увиденное произвело на меня сильнейшее впечатление. Зато стоявший рядом со мной во время демонстрации "чуда" человек профессорского вида, похоже, ничуть не удивился. Дернув себя за бородку и подмигнув мне, он отчетливым шепотом пояснил: "Это, молодой человек, левитация. Такое я уже видел". К сожалению, мне не удалось побеседовать с бородачом и расспросить его ни о сути самого явления, ни о том, при каких обстоятельствах и где он смог его наблюдать. Но с тех пор этот феномен, который Большая британская энциклопедия определяет как "сверхъестественную способность становиться легким по собственному желанию и подниматься в воздух без применения механизмов", очень меня занимает. Увы, никогда больше "вживую" летающих людей я не видел, только по телевизору в передачах, посвященных индийским йогам и буддийским монахам. Читать же о левитации приходилось много, благо литературы, в том числе фантастической, мистической и псевдонаучной и даже серьезной научной, посвященной этому удивительному явлению, предостаточно.

Описания случаев левитации встречаются с незапамятных времен. Самое удивительное - в старинных книгах способность к левитации отмечается не только у колдунов, ведьм и одержимых дьяволом, но и у праведников. Только если колдунов и ведьм за "полеты" сжигали на кострах, у праведников они считались признаком святости. Если верить историческим источникам, во время молитвенного экстаза в воздух поднимались многие религиозные деятели. Среди русских "левитантов" можно назвать Серафима Саровского, архиепископа Новгорода и Пскова Иоанна. Кроме того, московские летописи рассказывают о Василии Блаженном, который не раз на глазах у толпы переносился через Москву-реку. Способностью подниматься в воздух обладали и католические святые. Есть документы, описывающие, как один испанский паломник умудрился перелететь через широкую реку, а епископ Валенсии и вовсе парил над землей 12 часов. В 1550 году в Верте, Брабант, несколько монахов "поднялись в воздух, карабкались по деревьям как кошки", и их якобы поддерживали невидимые пальцы. А свидетелями полетов монахини-кармелитки Терезы

однажды стали 230 католических священников. О своем даре Тереза рассказала в автобиографии, датированной 1565 годом: "Вознесение приходит как удар, неожиданный и резкий, и, прежде чем ты можешь собраться с мыслями или прийти в себя, тебе кажется, будто облако уносит тебя в небеса или могучий орел на своих крыльях... Я вполне осознавала себя, чтобы видеть, что нахожусь в воздухе... Должна сказать, что, когда вознесение оканчивалось, я ощущала необыкновенную легкость во всем теле, словно я совсем невесомая". Умел летать и Иосиф Деза (1603-1663), прозванный Купертинским. Этот человек с детства отличался необычайной набожностью и всячески истязал себя, чтобы испытать состояние религиозного экстаза. А после того как был принят в орден францисканцев, стал действительно впадать в экстаз и в этом состоянии... взмывал в воздух. Более ста случаев левитации Иосифа наблюдали тогдашние ученые, оставившие на сей счет официальные свидетельства. Всего же, по сведениям из церковных книг, количество святых и истинно верующих, демонстрировавших феномен левитации, приближается к трем сотням. В Новом Завете есть прямое указание на то, что не один Христос мог ходить по воде. В Евангелии от Матфея сказано: "Петр вышел из лодки и пошел по воде, чтобы подойти к Христу. Но, видя сильный ветер, испугался и начал утопать, закричал: "Господи! Спаси меня". Иисус тотчас простер руку, поддержал его и говорит: "Маловерный! Зачем ты усомнился?.."».

Левитация была известна и на Востоке. Но там этому искусству обучались специально. В индийских Ведах содержится даже практическое руководство по левитации, которое описывает, как привести себя в такое состояние, чтобы оторваться от земли. Левитация в древности практиковалась также в Тибете. Буддистские тексты повествуют о том, что когда в 527 году нашей эры в тибетский монастырь Шаолинь пришел индийский основоположник дзэн-буддизма Бодхидхарма, он научил монахов управлять энергией тела - непременному условию для полетов. Пользовались левитацией и сам Будда, и его наставник маг Саммат, которые могли часами оставаться висящими в воздухе. Говорят, что в Индии и в Тибете искусство левитации сохранилось до наших дней. Например, британская путешественница Александра Давид-Неель наблюдала, как на высокогорном плато Чанг-Танга один из буддистских монахов, сидя неподвижно с подогнутыми под себя ногами, пролетал десятки метров, касаясь земли и вновь взмывал в воздух. А в 1986 году в Вашингтоне даже проходило соревнование "летающих йогов", в котором принимало участие около 20 человек. В программу входили полеты в высоту (был достигнут рекорд около 60 см), полеты в длину (рекорд - 1,8 м) и полеты на скорость на дистанции 25 м с препятствием высотой 20 см.

Но парить в воздухе способны не только буддисты, йоги и христианские святые. В литературе описано несколько случаев, когда левитировали и вполне обычное (на первый взгляд) люди. В XIX веке российский журнал "Ребус", специализирующийся на всяких необычных явлениях, опубликовал материал о случае, произошедшем с восьмилетним Н.Юрловым. 19 июля 1837 года мальчик спал в своей детской, на втором этаже. Ночью он был разбужен грозой и к ужасу своему увидел, что недалеко от его кровати на балконе, держась за ручку стеклянной двери, стоит высокий лысый человек, с седой бородой, в длинной синей рубашке. От ужаса мальчик скатился вниз по лестнице со второго этажа дома, выбежал из него и, ничего не соображая, стремительно побежал в сторону реки Свияги, протекавшей у села Спешневки Сенгилеевского уезда, Симбирской губернии. Прислуга спохватилась быстро, и мальчика нашли уже через десять минут, но на другом берегу реки и - совершенно сухого! Бежавшие за мальчиком люди, Василий Кондаков и Федор Плотников, садовник Николай Ермаков и кузнец Архипов, переплы whole вплавь Свиягу, клятвенно удостоверили, что они ясно видели, как ребенок быстро по воздуху переносился через реку, почти наравне с водою.

А чего стоит случай с ярмарочной гадалкой Марысей Лозинкой, который, если верить газете "Жиче Варшавы", случился польском городке Гошковице! В воскресный день на людной ярмарочной площади Лозинская сидела за столиком и занималась своим обычным делом, то есть гадала на картах всем желающим. Вдруг пани Марыся пожаловалась на

сильнейшую головную боль и, вскрикнув: "Как больно!", в тот же момент начала медленно подниматься над землей. При этом ее тело оставалось совершенно неподвижным. Все вокруг с изумлением смотрели на женщину, неведомым образом зависшую в воздухе на высоте около трех метров. Продолжалось это две минуты. Затем Марыся стала медленно снижаться. Едва ее ноги коснулись земли, как она тут же упала ничком. Свидетелем полета гадалки были сотни людей.

О знаменитом медиуме прошлого века Дэниеле Дугласе Юме рассказывали, что он однажды, поднявшись на воздух и облетев комнату, в горизонтальном положении вылетел в окошко и влетел в другое, произведя эту процедуру на высоте 25 футов над землей. Потом, когда его спрашивали об этом, он рассказывал, что ничего не помнит.

Левитация до недавнего времени напрочь отвергалась наукой. Хотя ученым уже давно было известно, что необычные психические состояния часто связаны с изменением веса человека. Так, например, профессор Санкт-Петербургской Военно-медицинской академии Ковалевский взвешивал пациентов во время приступов эpileпсии и после. Результаты этих взвешиваний показали, что падение веса эpileптика при самой легкой форме болезни - головокружении - составляет от 2 до 9 фунтов. А при эpileптических судорогах - до 12 фунтов. В случае же глубокой психической болезни, которой часто сопровождается эpileпсия, и при длительном припадке эти потери достигают четверти веса. Современными исследованиями установлено, что в особо тяжелых случаях потери составляют до 33-35% веса больного; при этом такая резкая потеря веса вовсе не оказывается на его внешнем виде. Через некоторое время после приступа вес восстанавливается. Резкие потери веса случаются и при других психических заболеваниях, например, истерии.

Отношение к феномену левитации у современных ученых начало изменяться после ряда опытов, проведенных российским физиком Евгением Подклетновым в лаборатории технологического университета финского города Тампера. Специальный диск охлаждался физиком до минус 167оС и помещался в электромагнитном поле, заставляющем его вращаться. При достижении 3000 об/мин предметы, помещенные под вращающимся диском, начали терять свою массу. Доклад о результатах эксперимента Подклетнов опубликовал в ряде научных журналов, после чего одно за другим начали появляться сообщения специалистов о наблюдаемых ими антигравитационных эффектах. Джон Шнурер из Энтиосского колледжа штата Огайо сообщил, что если над магнитом поместить сверхпроводник, он зависает в воздухе. Вообще-то, это явление уже было известно как "эффект Мейснера". Любопытно вот что: измерения показывают, что над сверхпроводящей системой возникает зона, где предметы теряют до 5% веса.

А голландские ученые умудрились подвесить в воздухе живую лягушку. Земноводное помещалось над сверхпроводящей катушкой: мощное поле воздействовало прямо на клетки тела, создавая в них магнитный момент, противоположный магнитному полю Земли. Таким образом, создавались силы, нейтрализующие тяготение. Голландцы утверждают, что их лягушка - первое живое существо, поднявшееся в воздух с помощью так называемого "молекулярного магнетизма". Исследователи заявили, что с помощью своей аппаратуры они смогут поднимать в воздух любой объект, в том числе и человека. Пока же, после эксперимента с лягушкой, они подвесили в воздухе бутерброд.

Кстати, в одном из российских университетов на кафедре психологии был проведен очень любопытный эксперимент. Во время сеанса гипноза подопытному внущили, что он находится в космическом корабле, который выходит на орбиту. Как сообщали потом участники эксперимента, если они и рассчитывали зафиксировать изменения показателей, то незначительные. И потому, когда приборы показали полное исчезновение веса, ученые просто не поверили своим глазам. Но этот невероятный факт был зафиксирован приборами. По свидетельству профессоров В.М.Инюшина и А.П.Дуброва, вес испытуемого, пусть и на незначительное время, исчез полностью.

После подобных экспериментов рассказы о летающих монахах и людях, ходящих по воде аки посуху, уже не выглядят столь фантастично. И как знать, может, тот колдун, на сеансе которого мне довелось присутствовать, вовсе не был шарлатаном.

(С) 1999-2002 Ежедневная электронная газета "УТРО"

Дата 20011206 Дата загрузки 20011205 Время загрузки 1917 Источник Московская правда Заголовок **МИНЗДРАВ НЕ ПРЕДУПРЕЖДАЛ** Автор Виола ЕГИКОВА. Выпуск 225 Текст

Наука: вести отовсюду

Как человек некурящий, но регулярно общающийся с заядлыми любителями подымить, позволю себе перефразировать известное выражение: обидеть курильщика может каждый. Вот и я добавлю с подачи ИнформНауки, предложившей обратиться за любопытной информацией на кафедру аналитической химии Московского университета.

Химики МГУ решили сравнить, сколько канцерогенных веществ поглощается человеком при выкуривании дорогих и дешевых сигарет. Исследование привело ученых к неожиданным выводам. Оказалось, большую или меньшую опасность представляет отнюдь не класс сигарет и не сорт табака. А еще, как выяснилось, не столь уж и важны указания на пачке о количестве никотина и смол, содержащихся в одной сигарете. Вовсе не эти вещества обещают печальную перспективу схлопотать рак горла или легких, о чем намекает в своем предупреждении Минздрав. Канцерогенными свойствами обладают так называемые ПАУ - полигарматические углеводороды, образующиеся в сигарете при горении.

На химфаке МГУ исследовали сигареты различного класса - от простейшей "Примы" до самых дорогих иномарок, дым которых собирали на специальные фильтры. Толщину последних подбирали так, чтобы обеспечить полную сорбцию дыма, затем фильтры промывали небольшими порциями органического растворителя, а полученный экстракт анализировали. Что же оказалось? Что цена сигарет ровным счетом не имеет никакого значения: в дыму как дорогих, так и самых дешевых сигарет ПАУ остается практически неизменным, на уровне миллионной доли грамма. И только в сигаретах с угольным фильтром содержание ПАУ снижается в 10 - 15 раз.

Кстати, цена сигарет вовсе не означает наличие именно такого фильтра. Бывает все как раз наоборот. Так что выбор дорогих сигарет, утверждают химики, чаще всего - лишь дань рекламе, сознательное или подсознательное стремление поддержать определенный имидж, но не более того.

Коли так, обратимся к подсознательным возможностям человека. Интересную публикацию на эту тему поместил журнал Nature, рассказав об экспериментах группы американских исследователей под руководством Такео Ватанабе из Бостонского университета. Добровольные помощники ученых выполняли задание, не подозревая, что они на самом деле тренируются для выполнения совершенно другой задачи. И когда она была поставлена, участники эксперимента справились с нею гораздо быстрее и успешнее, нежели другая группа, которая ранее не была задействована в исследованиях.

Суть проведенного эксперимента, на первый взгляд, довольно проста: его участникам предлагали называть буквы, периодически появлявшиеся на экране. Задание состояло именно в этом, и никто не подозревал, что главной целью эксперимента были перемещающиеся на том же экране точки. Они двигались хаотично, и лишь одна из двадцати выдерживала определенное направление. Экран показывали испытуемым по часу в день в течение месяца, после чего им показали экран уже без букв, но с точками. На этот раз в строго определенном направлении двигалась уже каждая десятая, что заметить все равно было довольно трудно. Но группа, до этого наблюдавшая экран, справилась с задачей гораздо лучше новичков.

Эти опыты, полагают психологи, подтверждают, что подсознательно воспринятая информация помогает обучению и стимулирует человека к принятию заданного решения.

Именно так воздействует на нас, например, реклама. Что же касается использования подсознательного восприятия в процессе обучения, здесь, как считает профессор Ватанабе, еще много неизученных возможностей. Так, регулярное прослушивание речи на незнакомом языке не приведет, по его мнению, к пониманию значения слов, но поможет улучшить произношение и слуховое восприятие.

А еще на наши языковые возможности влияет музыка. Вывод сделан уже не психологами, а специалистами из Института неврологии Макса Планка (Германия), о работе которых также рассказывает Nature. Исследования, проведенные учеными этого научного центра, показали, что умение различать построение слов в предложении, равно как и музыкальные фразы, контролирует один и тот же участок головного мозга. Именно он помогает не только понимать язык, но и отличать гармонию звуков от какофонии. Называется этот отдел мозга центром Брука. Собственно, его роль в нашем языковом общении - не новость, а вот то, что существует прямая связь между восприятием грамматических правил и музыки, это, по мнению германских ученых, важное открытие, которое, не исключено, поможет лечить людей с речевыми расстройствами.

Об открытии судить специалистам. Слово это довольно часто используется, когда упоминают о научных исследованиях, хотя, строго говоря, открытием как таковым можно называть далеко не любой научный результат. Открытий в науке не так уж и много. В России, например, за последнее десятилетие (не самое благоприятное для ученых) было зарегистрировано 172 научных открытия. Больше всего их в медицине и биологии. В физике в наши дни они чрезвычайно редки. Тем приятнее сообщить об открытии, зарегистрированном недавно учеными Томского политехнического университета.

Связано оно с физикой твердого тела. Сибирские ученые установили, что под воздействием малых доз ионизирующего излучения происходит упорядочение структуры кристаллов. Как считают специалисты, открытие этого физического явления найдет широкое практическое применение. Оно позволит, например, значительно повысить надежность и продлить срок службы ядерных реакторов, а также установок для добычи нефти и газа.

Будет ли сказано новое слово в физике и по поводу гравитации? Возможность преодоления гравитационных сил когда-то категорически исключалась наукой, да и сегодня многие ученые считают разговоры на эту тему полнейшей ересью. Но далеко не все! Год назад НАСА даже выделило солидную сумму на поощрение работ по поиску возможностей преодоления гравитации.

Всерьез на эту тему заговорили после экспериментов российского ученого Евгения Подклетнова, который, работая со сверхпроводниками в начале 1990-х в Технологическом университете Тампере (Финляндия), обнаружил явление частичной потери веса. Затем стало известно об экспериментах американского физика из Эндиоического колледжа (штат Огайо) Джона Шнурера... А недавно на международном семинаре в Москве о любопытных научных результатах рассказал сотрудник Института общей физики РАН Дмитрий Ципенюк. В ходе экспериментов российским физикам удалось зафиксировать отклонения, вызванные изменением гравитационного поля...

Ученый был крайне осторожен в своих оценках, считая необходимым продолжение экспериментов. Так что о преодолении гравитации - пока ни слова. Но уж если кто-то получит такой результат, вот уж будет открытие так открытие! (Наш дом - Москва 36)

Дата 20001013 Дата загрузки 20001013 Время загрузки 1346 Источник Эпиграф, Новосибирск Заголовок компьютера+ page Тема компьютера+ page Выпуск 40 (290) Текст

Вне власти притяжения

NASA планирует к лету будущего года заполучить в свои руки технологию, способную вывести эту организацию в абсолютные лидеры среди космических агентств мира. Речь идет о... гравитационном щите, или попросту - антиграве.

В прошлом году NASA вложило 600 тыс. долларов в разработку компанией Superconductive Components Inc. прототипа устройства, способного снижать гравитацию.

Компания пытается повторить эксперимент 1992 года, когда российский ученый Евгений Подклетнов, работающий в Финляндии, заявил об успешном испытании хитрой конструкции, основой которой был левитирующий в магнитном поле быстро вращающийся диск из суперпроводящей керамики. По утверждению автора, объекты, помещенные над диском, теряли до 2% своего веса. В настоящее время Подклетнов занят в Superconductive в качестве консультанта; однако, до сих пор не ясно, удастся ли NASA повторить его эксперимент; по крайней мере, до сих пор это получилось, по словам того же Подклетнова, только у нескольких групп исследователей, в точности следовавших его указаниям.

Первое применение, придуманное NASA для разрабатываемой технологии, таково: антигравитационный щит должен будет снизить гравитацию на месте запуска космических кораблей, существенно уменьшив тем самым накладные расходы на преодоление притяжения Земли - топливо и пр.

Большинство физиков считает, что антигравитация невозможна, поскольку она нарушает фундаментальные законы природы. Однако, природа исследуемых явлений слишком необычайна, а потому, надежда на успешный исход остается.

Не как все

Интернациональная группа астрономов, в которую вошли представители Испании, Германии и США, опубликовала в журнале "Science" отчет об исследовании, которое может в корне изменить представления науки о формировании планетных систем.

До настоящего времени считалось, что планеты возникают из вращающихся вокруг молодых звезд газопылевых облаков. Однако восемнадцать объектов, обнаруженных учеными по отражаемому ими свету в туманности Ориона, судя по всему, имеют совершенно иное прошлое. Странствующие объекты (каждый из которых - газовый гигант, близкий по параметрам к Юпитеру, но больший его по массе в 8-15 раз) принадлежат к звездной системе, родившейся только несколько миллионов лет назад (планеты возле нашего Солнца родились, когда ему было уже несколько миллиардов лет). Объяснить их существование с позиций современной астрофизики весьма затруднительно. Если бы это были старые, потухшие, но еще относительно горячие звезды, то их существование было бы понятным, однако, это именно холодные планеты (что доказано спектральным анализом). Теперь науке предстоит разработка теории, способной убедительно описать появление и эволюцию таких планет.

Как это будет по-русски...

С 10 октября в России начала продаваться локализованная версия ОС Windows ME, последней ОС, построенной на ядре Win9x.

Оригинальная англоязычная версия системы вышла почти за месяц до этого, 14 сентября. Тогда одновременно появилось 15 версий Windows ME на разных языках, сейчас вышла русская редакция, а к концу года ожидается, что Windows ME "заговорит" на 28 языках.

В русскоязычной версии полностью русифицирован интерфейс, переведена справочная система. Кроме русского языка Windows ME имеетстроенную поддержку украинского и белорусского языков. В остальном же операционная система практически идентична своему англоязычному собрату за исключением того, что в русской версии системы отсутствует пакет WebTV, в наших условиях бесполезный.

Кроме того, уже в следующем году в России будет запущена служба интерактивной техподдержки пользователей продуктов Microsoft. Windows ME - это первый программный продукт Microsoft, в который встроены функции, способные работать с этой службой. Для облегчения диагностирования проблем ОС может передавать данные о компьютере пользователя в службу поддержки. Справедливости ради надо заметить, что эта функция, которая далеко не всем придется по вкусу, может быть отключена.

Могуч и призрачен

Солнечные паруса, применить которые для путешествия в пределах ближайших к нам планет планирует NASA, внезапно обрели новое свойство - они стали прозрачны!

Ученые предложили заменить парус на... магнитный пузырь - "сплетенный" вокруг летательного аппарата кокон магнитных полей, наподобие тех, что есть у Земли и многих других планет Солнечной системы. Магнитосфера Земли защищает нас от потоков космических частиц, искривляя траектории и гася их энергию над полюсами планеты (вспомните явление северного сияния). Но она действует еще и как огромный парус, улавливающий потоки солнечного ветра и толкающий Землю с орбиты по направлению от Солнца. Наша планета слишком массивна, чтобы сдвинуться с места под воздействием этих усилий, однако, представьте себе небольшой летательный аппарат. Генерируемая им магнитосфера диаметром в 15 км, находясь на расстоянии в одну астрономическую единицу от Солнца, способна за три месяца разогнать 200-килограммовый космический кораблик до скорости в 80 км/с. Однако, как создать магнитное поле таких размеров?

Исследователи из NASA предложили использовать вместе с обычным мощным электромагнитом поток плазмы. Высокотемпературный ионизированный газ (получаемый нагревом в своеобразной микроволновой печи гелий-argonовой смеси), впрыскиваемый в магнитное поле близ катушки электромагнита, дает удивительный эффект: размеры поля увеличиваются во много раз. Диаметр 15 км уже может быть достигнут, правда, только по словам авторов, поскольку развернуться до таких пределов на Земле не позволяют размеры лаборатории. Для питания установки необходим килограмм газовой смеси и 1 киловатт электроэнергии в день. А мощность солнечного ветра, который будет захвачен полученной магнитосферой, равна 600 киловаттам. Интересно, что природа уже давно использует похожий плазменный "надув": магнитосфера Юпитера своими огромными размерами обязана именно плазме, вдуваемой в поля планеты-гиганта действующими вулканами его спутника Ио.

Два интересных побочных свойства нового космического движка: защищенность экипажа корабля, путешествующего с его применением, от космической радиации (потоки излучения просто не смогут пробиться сквозь магнитный заслон); и второй - по мере удаления от Солнца размер магнитного паруса будет увеличиваться, поскольку давление солнечного ветра, который его "сжимает", падает. Таким образом, разработчики надеются, что двигатель этот можно будет эффективно использовать в пределах всей Солнечной системы и даже за ней - на больших удалениях от звезды.

Хорошего - помногу

Компания Texas Instruments объявила о выпуске чипа, способного совершить революцию на рынке переносных проигрывателей цифрового звука.

Речь идет о микросхеме DA250, которая содержит все необходимое для построения на ее основе цифровых проигрывателей. Это аппаратные кодер/декодер (с перепрограммируемой флэш-памятью для поддержки различных форматов звука, не только MP3), поддержка нескольких коммуникационных интерфейсов (в том числе USB) и пр. Но самое главное преимущество перед аналогами - режим управления энергопотреблением. Специалисты TI встроили в свое детище способность отключения от питания неиспользуемых вычислительных блоков прямо "на ходу". Кроме того, энергопотребление удалось существенно снизить также благодаря интеграции всех функций на едином кристалле. Результатом стал увеличенный до 70 часов ресурс работы от двух батареек типа АА.

Поставки чипа намечены на начало следующего года. Планы по построению на его основе плейеров уже есть у Samsung и еще почти 60 более мелких разработчиков.

Дата 20001002 Дата загрузки 20001005 Время загрузки 2231 Источник Журнал Огонек, Москва Заголовок **ЛЕГЧЕ ЛЕГКОГО** Тема ПАНОРАМА Выпуск 36 Текст

Услышав, что в NASA полным ходом идут работы по программе "Прорыв в области физики движения", а проще говоря, по антигравитации, специалисты из BAe Systems (бывшая British Aerospace) решили: "А что мы, хуже?" и заявили, что они тоже занимаются разработкой принципов антигравитационных двигателей. А ведь еще в 1996 году они все

вместе объявляли шарлатаном российского физика Евгения Подклетнова, заявившего, что он нашел способ экранирования сил притяжения. Вот вам и научная этика. Теперь будут выдавать научно обоснованную "нуль-транспортировку" за свое открытие.

Дата 20000801 Дата загрузки 20000731 Время загрузки 2005 Источник Комсомольская правда, Москва Заголовок **ЛЮДИ БУДУТ ЛЕТАТЬ, КАК ПТИЦЫ!** Автор СВЕТЛАНА ХАБЛИЦКАЯ, АННА ЕРОШОВА, АЛЕКСАНДР МИЛКУС Выпуск 138 Текст

Российский ученый Евгений Подклетнов доказывает: антигравитация возможна

Ах, почему люди не летают, как птицы? - мечтала Катерина в хрестоматийной Грозе . А если всерьез, без литературной лирики разобраться - почему? Что мешает нам воспарить под облака? Да все просто: сила тяготения - объяснил даже тот, у кого в школе по физике была круглая двойка . Просто? Но ведь до сих пор механизм тяготения, припечатывающего всех нас к Земле, или, по-другому, гравитация, для человечества - загадка!

- Происхождение силы тяжести неизвестно. Никто из ученых не изучал гравитацию на субатомном уровне, - рассказывает физик из Института Макса Планка в Германии Джованни Моданес. - Нам просто не хватает знаний для того, чтобы сделать какие-то определенные выводы. Значит, и создание прибора для преодоления гравитации возможно. Так все-таки есть шанс взлететь у Катерины?

Диск полегчал

...Лет сто назад об этом решались писать разве что романтики-фантасты. А сегодня в США уже второй раз проходит серьезная научная конференция по антигравитационным устройствам. Главная тема для обсуждения - попытка исследовательского центра в Хансвилле создать технологию, позволяющую преодолеть земное тяготение. Обосновывая свои выкладки, американцы ссылаются на недавнюю работу русского ученого Евгения Подклетнова.

Ожидается, что антигравитационный эффект прежде всего может наблюдаться в сверхпроводниках. Первая серьезная работа по этому поводу была опубликована еще в 1976 году. Американские физики обратили внимание - при работе с такими материалами происходят гравитационные нарушения. Но продолжения исследований на эту тему авторы в научных журналах не помещали. То ли засомневались в своих выводах, то ли работы засекретили...

Потому и произвела фурор статья нашего Евгения Подклетнова, опубликованная в 1996-м в солидном научном издании. Ученый рассказывал, что диск из сверхпроводящей керамики, вращающийся над мощным электромагнитом, становится легче на 0,3 процента. Немного? Но ведь лиха беда начало!

- Евгений окончил Московский химико-технологический университет имени Менделеева, потом пришел в наш институт, - говорит сотрудник Института теоретической и прикладной электродинамики РАН доктор физико-математических наук Александр Львович Рахманов. - Он всегда был отличником. Его избирали секретарем комсомольской организации института. Специализировался Евгений на сверхпроводящих пленках. Несколько лет назад уехал в Финляндию. Занимается исследованиями в университете города Тампере. Но диссертация его никакого отношения к антигравитации не имела. Сначала исчез сигаретный дым

Эффект антигравитации Подклетнов обнаружил случайно во время обычных испытаний сверхпроводящего диска из высокотемпературной керамики. Три электрические катушки создавали магнитное поле, в котором этот диск вращался со скоростью пять тысяч оборотов в минуту. И вдруг экспериментатор заметил, что дым от сигареты, тянувшийся струйкой через комнату, почему-то над прибором исчезал. Выяснилось: в этом месте возникла область с пониженным давлением атмосферы.

Стали проверять почему. И вот тогда обнаружилось, что быстро вращающийся диск уменьшает гравитационное поле и предметы, подвешенные над экспериментальной установкой, теряют в весе.

С самим ученым нам связаться не удалось. Говорят, теперь в Москве он бывает нечасто, а о результатах своих новых исследований предпочитает не распространяться. Особенно журналистам. Тем более что открытие его, если оно действительно будет подтверждено другими специалистами, грозит в корне изменить серьезные фундаментальные научные теории, и для серьезных заявлений нужны новые эксперименты, которыми Подклетнов сейчас и занимается.

Представляете, какие перспективы открываются, если российскому ученому удастся увеличить полученный эффект и заставить пластинку пусть пока не парить - просто приподняться над землей? Оседлав антигравитацию человечеству, придется пересмотреть все нынешние транспортные системы. Самолетам уже не понадобятся крылья - можно будет обойтись антигравитационным экраном. Не нужно будет вкладывать сумасшедшие деньги в автомобильные дороги: поднялся в небо - и лети себе. Да и ракеты, заполненные тоннами ядовитого топлива, не станут больше буравить небо! В США вкладывают деньги в сумасшедшие проекты

Поэтому-то, несмотря на то что опыты по антигравитации не вписываются в современные правильные теории, НАСА и не отвергает их с порога. Не так давно здесь создана особая лаборатория, призванная проверять самые безумные идеи, если только они каким-то боком имеют отношение к полетам в космос. Бюджету этой лаборатории позавидовало бы любое российское научное учреждение - 685 тысяч долларов. Определено пять направлений исследований. Под номером четыре - антигравитация.

- Уже несколько лет мы пытаемся добиться антигравитации, но наши устройства оказываются не настолько удачными, как это бы хотелось, - признает один из участников доктор Джон Брэнтли. - Подклетнов мог бы помочь нам, но он категорически отказывается говорить о некоторых аспектах своей работы.

Мнение эксперта КЛ Бориса ВЕЙЦМАНА (Центр космических исследований имени Годдарда, НАСА):

- Не думаю, что эффект Подклетнова - шарлатанство. Видно, что ученый умеет работать и с низкими температурами, и со сверхпроводниками. Но я бы не торопился с утверждением о том, что средство для достижения антигравитации найдено. Ученые в Англии и Индии, занимавшиеся работами Евгения, считают: результаты, полученные русским специалистом, могут быть объяснены неучтенными в эксперименте эффектами. В общем, то, что предмет несколько похудел, - скорее всего, ошибка экспериментатора.

И все-таки, уверены оптимисты, рано или поздно людям удастся справиться с гравитацией.

- Если уже сегодня можно создать экраны для электромагнитных полей, то и антигравитационный экран не нарушает никаких законов физики, - уверен профессор физики из университета штата Мэриленд Ричард

МакНикол.

В общем, поводов для уныния пока нет. Нужно просто ждать. Каждому открытию - свое время. Тело подают на тарелочке

В Японии уже лет пять демонстрируют такой аттракцион. Упитанного человека ставят на тарелку, отлитую из сверхпроводящей керамики. Под ним располагают электромагнит и поливают его жидким азотом (чтобы тот охладился). Магнит замерзает, и человек приподнимается в воздухе. Ничего фантастического, увы, тут нет. Сверхпроводящую пластину выталкивает магнитное поле. На этом эффекте построены железные дороги на сверхпроводящем подвесе. Да, движутся по таким рельсам поезда быстрее, так как отсутствует трение. Но, увы, для поддержания сверхпроводимости нужно расходовать гораздо больше энергии, чем для преодоления трения.

КСТАТИ И все-таки воспарим...

Бритиш эйрспейс - одна из дочерних фирм гиганта английской авиакосмической индустрии концерна BAE Systems - объявила о начале финансирования разработки устройства, способного преодолевать земное притяжение. Научный проект получил название

Гринглоу , что в переводе на русский означает Зеленый отблеск . Мы считаем наш проект началом необыкновенного приключения, - говорят организаторы проекта, - к которому приглашаем присоединиться академиков, членов правительства и финансистов со всего света

- По моему мнению, высокие посты в Бритиш аэроспейс занимают люди, которые не знают основных законов фундаментальной физики. Шанс, что этот проект будет успешным, равен нулю, - так саркастично прокомментировал это событие Боб Парк из Американского физического общества. Почему бутерброд падает маслом вниз?

Красивым подтверждением того, что не все тела притягиваются Землей с одинаковой силой, могла бы стать судьба злосчастного бутерброда, всегда падающего маслом вниз. Увы, этот прискорбный момент объясняется физиками гораздо проще. Если один предмет соединить с другим, более тяжелым, то во время полета он перевернется тяжелым предметом вниз, просто потому, что тот весит больше. Тело занимает такое положение, чтобы его центр тяжести находился как можно ниже. Поэтому, если трясти ведро с картошкой, крупные картофелины оказываются вверху, а мелкие плотно укладываются на дно. Масло тяжелее хлеба, вот его и тянет к земле сильнее.

Светлана ХАБЛИЦКАЯ, Анна ЕРОШОВА, Александр МИЛКУС.

Дата 20000711 Дата загрузки 20000825 Время загрузки 2303 Источник Тульские известия Заголовок **А ВДРУГ ПОЛУЧИТСЯ?** Тема С МИРУ ПО СТРОЧКЕ Выпуск 154 Текст

Одна из дочерних фирм гиганта авиакосмической индустрии "Бритиш аэроспейс" приняла решение финансировать проект создания устройства, способного преодолевать силу земного притяжения. В основе революционной идеи лежат, как стало известно в Лондоне, изыскания российского физика Евгения Подклетнова, который в данный момент работает в Финляндии.

Российский ученый утверждает, что ему удалось разработать антигравитационное устройство, состоящее из врачающегося диска и электромагнитов. По словам самого Е. Подклетнова, путем экспериментов он смог снизить силу тяжести более чем на два процента. Если подобные данные подтвердятся, это потенциально способно вызвать настоящий переворот в представлениях о законах физики, а также революцию в истории перемещения человека в земном и околоземном пространстве.

Современная наука достаточно скептически относится к самой возможности создания антигравитационного аппарата. По мнению ряда специалистов, в конечном счете проект может быть признан бесплодной тратой средств и времени на фантастические разработки.

Дата 20000621 Дата загрузки 20000620 Время загрузки 1802 Источник Независимая газета, Москва Заголовок **Антигравитация все-таки существует?** Автор Леонид Ширшов Выпуск 6 (32) Текст

В рамках физических программ некоторых английских и американских научных центров ведутся эксперименты по экранированию притяжения Земли

В "Ленте" научно-технических новостей Би-би-си от 27 марта 2000 г. появилась информация, посвященная принятию решения о начале работ по созданию приборов, экранирующих притяжение Земли.

Военное крыло группы высоких технологий ВАЕ, прежде известной как Британское Аэрокосмическое объединение, подтвердило, что начата программа исследования антигравитации. Высказывается надежда, что проект "Гринглоу" объединит ученых, работающих над созданием двигателевых установок для транспортных средств будущего.

В рамках физических программ НАСА также ведутся эксперименты по экранированию гравитации. Как сообщается, они были инициированы исследованиями российского ученого Евгения Подклетного, заметившего, что врачающийся диск из высокотемпературной сверхпроводящей керамики (ВТСП-керамика) уменьшает гравитационное поле.

Американское Национальное аэрокосмическое агентство (НАСА) выделило грант фирме Superconductive Components на разработку больших сверхпроводящих дисков со специальной микроструктурой для антигравитационной машины.

В 1996 г. Евгений Подклетнов подготовил для научного журнала статью, в которой утверждал, что вращающийся ВТСП-диск уменьшает гравитацию. В неопубликованной работе по экранированию гравитации сверхпроводником он сообщал о потере 2% от веса груза. Однако большинство ученых были уверены, что эксперименты по антигравитации были проведены не вполне корректно и противоречат тому, что мы знаем из законов природы, и поэтому такие результаты невозможны.

Комментируя нынешнее решение англичан, Боб Парк из Американского физического общества сказал: "Я нахожу достаточно характерным, что они пошли на это. Можно только заключить, что на высшем уровне этой организации имеются люди, не обладающие достаточной подготовкой в области фундаментальной физики".

Однако такая точка зрения вряд ли остановит работы по антигравитационным исследованиям. Так, по запросу с ключевыми словами "Anti-Gravity" системы информационного поиска в Интернете дают ссылку на добрую тысячу документов

(см., например, описание истории злоключений Евгения Подклетнова)

Неожиданное явление было обнаружено в ходе стандартных испытаний с использованием сверхпроводящего диска из высокотемпературной керамики, которые проводились в 1992 году на базе технологического университета г. Тампера (Финляндия). Три электрические катушки создавали магнитное поле, в котором вращался ВТСП-диск, скорость которого достигала 5 тыс. оборотов в минуту.

Прибор размещался в криостате диаметром 11 дюймов (около 30 см) с жидким азотом. В процессе экспериментов исследователи наблюдали странную картину: дым от сигарет тянулся струйкой через комнату и исчезал над криостатом. Над ним была обнаружена область с пониженным давлением атмосферы. Для объяснения поведения дыма не было видимых причин, поскольку воздушных потоков в комнате не было.

Экспериментаторы предположили, что быстровращающийся диск из сверхпроводящего материала уменьшает гравитационное поле, и стали проверять этот эффект, подвешивая разные объекты выше криостата. При этом обнаружилось, что вес предметов уменьшался на два процента, что является существенным значением для такого рода измерений. Размещение двух криостатов один над другим, показало, что уменьшение веса грузов составило уже четыре процента.

"Мы предполагали, что возможна ошибка, но предприняли все меры предосторожности", - отметил Подклетнов, отправляя подготовленную публикацию об этом эффекте в престижный "Журнал прикладной физики", издаваемый английским институтом (The Journal of Physics-D: Applied Physics, a publication of England's Institute Physics). Группа авторов выдвинула гипотезу о том, что вращающийся диск сверхпроводника в какой-то мере блокировал действие гравитации. Несмотря на необычный характер сообщения, посвященного антигравитации, его рассмотрели независимые эксперты, которые рекомендовали материал к печати. Публикация планировалась в октябре 1996 года.

Но немногим ранее сенсационные новости стали достоянием широкой прессы. "Санди-телеграф" поместил 1 сентября 1996 г. статью "Антигравитационные приборы снимают ограничения". В ней предсказывалось большое будущее технологических использований антигравитации и сообщалось, что НАСА собирается финансировать эти исследования.

Широкий резонанс в мире, вызванный обсуждением антигравитационной сенсации, вызвал ряд непредсказуемых последствий. Представитель финского технологического университета в Тампере выступил с заявлением о том, что в стенах этого учреждения вопросами антигравитации не занимались. Возможные соавторы статьи отрицали свое участие в подготовке публикации, и 9 сентября 1996 г. Евгений Подклетнов предложил редакции журнала снять подготовленный текст.

Вопрос с испытанием "машины антигравитации" остался открытым до сих пор. Возможно, произошло непреднамеренное искажение результатов эксперимента, как случилось при наблюдении "холодного синтеза ядер" - другой нашумевшей и развенчанной сенсации. Достаточно простая конструкция установки побудила многих исследователей повторить эксперименты по наблюдению экранирующего воздействия вращающегося сверхпроводящего диска, но про результаты успешного функционирования "машины антигравитации" до сих пор не сообщалось.

Интересным представляется тот факт, что за 7 лет до экспериментов Подклетнова, теоретик Нинг Ли, работавший в Центре космических полетов имени Маршалла, опубликовал три статьи в серьезных журналах, где предсказывал, что сверхпроводник, вращающийся в сильном магнитном поле, может нарушать гравитационные силы поблизости от диска. Как сообщается в статье "NASA's Antigravity Machine", Нинг Ли, работающий в настоящее время научным сотрудником научно-исследовательского факультета университета штата Алабама, оказывает помощь центру Маршалла в создании сверхпроводящего диска для прибора антигравитации.

Планируется воспроизвести установку Подклетнова, усовершенствовав конструкцию. Антигравитационное устройство

НАСА включает композиционный диск диаметром 20 дюймов, который будет вращаться в жидким гелии и охлаждаться до сверхнизких температур. Главные трудности представляет изготовление большого диска из сверхпроводящей керамики из-за высокой хрупкости материала.

В заключение можно добавить, что 23-25 июня в штате Невада состоится

2-я международная конференция по антигравитационным исследованиям.

(с) НГ Наука N 6 (32) 21.06.2000

Дата 20000511 Дата загрузки 20000510 Время загрузки 2002 Источник Труд, Москва Заголовок "**ЗЕЛЕНЫЙ ОТБЛЕСК" РУССКОГО ФИЗИКА**" Автор Игорь ВАДИМОВ. Тема НА ГРАНИ ФАНТАСТИКИ Выпуск 84 Текст

Еще недавно об антигравитации писали только фантасты. И вот на днях научный мир был потрясен сенсационным известием - солидная организация (одна из дочерних фирм гиганта авиакосмической индустрии "Бритиш аэроспейс") сообщила, что будет финансировать проект создания устройства, способного преодолевать силу земного притяжения. В основе революционной идеи лежат, как стало известно в Лондоне, изыскания российского физика Евгения Подклетнова, который в данный момент работает в Финляндии. Российский ученый утверждает, что ему уже удалось разработать антигравитационное устройство, состоящее из вращающегося диска и электромагнитов. По словам самого Е.Подклетнова, путем экспериментов он смог снизить силу тяжести пока только на два процента. Но если эти данные подтвердятся, даже такой незначительный на первый взгляд антигравитационный эффект способен вызвать революцию в истории перемещения человека в земном и околоземном пространствах.

Компания "Бритиш аэроспейс систем" подтвердила сообщения о своем намерении финансировать изыскания в этой области. Проекту уже присвоено рабочее название "Гринглоу" - "Зеленый отблеск".

Дата 20000426 Дата загрузки 20000428 Время загрузки 1629 Источник Алфавит, Москва Заголовок **АНТИГРАВИЦАПА** Автор Андрей Самохин Выпуск 17 (75) Текст

Человек успешно делает научную карьеру, идет по восходящей ученых степеней. И вдруг: вспышка, озарение (или "обольщение") - и он уже одержим некой идеей, корпит по ночам, конструируя перпетуум-мобиле или "летающую тарелку". Не признанный официальной наукой, досадуя на косность специальных журналов и медвежьи услуги желтой прессы, укоряемый семьей и друзьями, он будет заниматься этим, пока не поймет, в чем ошибка, или... не докажет своего. По обочинам его пути маячат тюрьма, сумма и желтый дом,

а на горизонте призывающе мерцают звезды нобелевки. Чаще всего такие люди становятся предметом насмешек, иногда - горьких сожалений современников... И снисходительного сочувствия потомков, реже - их уважения.

Роман Герберта Уэллса "Путешествие на Луну" (1899) повествует о том, как ученый Кавор изобрел некую пасту "каворит", экранирующую земное притяжение. Обмазав ею круглый аппарат, он тотчас отправился с друзьями прогуляться по Селене. Великий английский фантаст как живой стоит перед глазами, когда читаешь о "трудах и днях" российского ученого Евгения Подклетнова. Ведь он претендует в наши дни на открытие... реального "каворита" в виде сверхпроводящего материала "YBCO".

Прочитав в подборке информации из агентств, приходящей в редакцию, о том, что дочерняя фирма "Бритиш Аэроспейс" начала финансировать "создание устройства, преодолевающего гравитацию на основе изысканий Е. Подклетнова", мы поначалу сочли это первоапрельской уткой. Однако, полазив по Сети, нашли массу интересных подробностей. Оказывается, в прошлом году НАСА истратило 600 тыс. долл. на исследование эффекта, обнаруженного нашим соотечественником, а количество статей о Подклетнове в западной прессе исчисляется уже десятками...

Евгений Подклетнов, выпускник знаменитого Менделеевского института в Москве, проработал 15 лет в Институте высоких температур АН СССР, а в 1988 г. был приглашен в Финляндию в университет г. Тампере по совместному проекту синтеза сверхпроводников. Он ехал работать над докторской, а с собой вез сверхпроводящий диск, который изготовил в Москве из расплава оксидов меди, иттрия и бария, - совсем как в романе Уэллса. В диаметре диск достигал почти 30 сантиметров. Огромный по тем временам. Да и сейчас американцы, по их собственному признанию, отстают от России в изготовлении таких дисков.

Там, в Тампере, и произошло "озарение". Конечно, случайно. В одном из штатных опытов диск охладили до температуры сверхпроводимости с помощью жидкого гелия и раскрутили электромагнитами. Далее, со слов самого ученого, озвученных журналистом американского журнала "Вайдед" Чарлзом Платтом, "кто-то в лаборатории курил трубку, и мы заметили, что дым, проходя над сверхпроводящим диском, поднимается вверх. Тогда мы поместили на диск оказавшийся под рукой намагниченный шарик, соединив его с весами. Показания весов нас удивили. Вместо металлического шарика взяли неметаллический и немагнитный материал - кремний. Весы вели себя по-прежнему очень странно. Оказалось, что любой предмет, помещенный над диском, чуть-чуть терял в весе, а если диск вращали, этот эффект еще увеличивался".

Подобрав определенную скорость вращения, экспериментатор добился максимального уменьшения веса предметов - на 2%. Ученый подчеркнул, что ошибки быть не может, что эксперименты он вел не один, с ним было шесть или семь опытных специалистов. Взвешиваемые предметы помещали в вакуум, экранировали листом металла - вес все равно уменьшался. Мало того, ртутный барометр, помещенный над диском, показал уменьшение атмосферного давления на 4 мм, и это уменьшение веса воздуха регистрировалось и на втором этаже, точно над экспериментальной установкой!

Открывшиеся перспективы так увлекли Подклетнова, что он потерял покой. Остался в Финляндии и лихорадочно продолжал опыты. В 1992 г. ученому удалось опубликовать предварительное сообщение о новом эффекте, после чего он с финским соавтором написал более подробную статью. Рукопись отвергли более десятка научных журналов, пока ее не принял солидный английский "Journal of Physics-D". Каким-то путем сотруднику желтоватой английской газеты удалось добыть гранки готовой к публикации статьи, и 1 сентября 1996 г. на видном месте в газете появился сенсационный материал, начинавшийся так: "Ученые из Финляндии вскоре опубликуют подробности устройства первого в мире генератора антигравитации". (Сам Подклетнов никогда не употреблял слово "антигравитация", он говорил лишь о возможности экранировать силу тяжести.) Явственно запахло Бэтменом из детских комиксов. Вскоре в мастихой "Санди телеграф" вышла разгромная заметка о сомнительных опытах "горе-ученого". Руководство университета в Тампере отреагировало

примерно так, как реагировали когда-то у нас на статьи в "Правде". Коллегу Подклетнова вынудили отказаться от соавторства, а его самого выгнали из университета. Новые друзья и знакомые отвернулись от русского ученого. В 1997 г. он сделал попытку вернуться в Москву, но нашел в своем родном институте "пепелище".

Оказавшись снова в Тампере, он устроился материаловедом в одну из местных фирм и продолжает опыты в свободное от работы время. И по-прежнему полагает, что сделал одно из крупнейших открытий XX в. и что не позднее чем через 10 лет оно найдет практическое применение. Евгений живет анахоретом, прячется от журналистов и ходит иногда около университета, куда его непускают...

Как вы уже догадались, это вовсе не финал. Большинство физиков скептически (мягко говоря) восприняли результаты опытов Подклетнова, основываясь на положениях общей теории относительности, согласно которой гравитация вообще не сила, а искривление четырехмерного пространства-времени. И вдруг, несмотря ни на что, НАСА неожиданно самостоятельно начало эксперименты по проверке эффекта Подклетнова в Центре космических полетов имени Маршалла в Хантсвилле (штат Алабама)**.

Результаты оказались малоутешительные - вес над сверхпроводящим диском уменьшался не на 2%, как у Подклетнова, а в миллион раз меньше. Американцы были обескуражены...и пригласили Подклетнова к себе. Он приехал в Хантсвилл в конце 1998 года. Верный своей привычке, избегал корреспондентов и не давал интервью. Тем не менее сведения о его визите просочились в печать. Причиной неудачи, по словам ученого, было то, что американцы не вращали диск. Кроме того, они работали с маленьким диском. Поэтому для точного воспроизведения опытов Подклетнова было решено сделать диск в два раза больше. Возникли сложности с его изготовлением. НАСА объявило конкурс среди профильных фирм. Создание "такого же диска, как в России" было оценено в 750 тыс. долларов. Наверное, приобретение диска у нас или у Подклетнова обошлось бы дешевле. Но американцы решили сделать его сами: на карту поставлен престиж страны!

За это время появилось немало сообщений о якобы повторенных и даже превзойденных результатах опытов русского ученого: группой студентов в Шеффилде (Англия), учеными из Торонто. Активным сторонником Подклетнова выступает итальянский физик Джованни Моданезе из университета г. Тренто. Выяснилось, что о волшебном антигравитационном экране Герберта Уэллса задолго до Подклетнова думали физики-теоретики из университета Алабамы - американец Дуглас Торр и китаянка Нинь Ли. По их настоянию в НАСА в 1990 г. была даже организована лаборатория для опытов с гравитацией. Само Национальное Агентство космических исследований отказывается пока публично комментировать окончательные результаты "эффекта Подклетнова", призывая нетерпеливых журналистов "немного подождать". А компания "Бритиш Аэроспейс", как уже было сказано выше, решила вовсе не ждать, а ковать, "пока горячо"...

Несмотря на поднятый "антигравитационный" шум и гам, крупные мировые физики продолжают по этому поводу хранить презрительное олимпийское спокойствие. Как и российские ученые. В адрес Подклетнова от них можно услышать лишь сакраментальное "типичная лженаука" и совет не писать об этом. На растерянный вопрос: "А как же НАСА и "Бритиш Аэроспейс"?" - следует ответ: "У этих организаций столько денег на исследования, что они могут позволить себе проверять весьма сомнительные идеи". А один собеседник, подмигнув, высказал небанальную мысль: "Женя все правильно делает - его не зря столько готовили".

Видя недоумение в моих глазах, собеседник пояснил: "В прошлую "холодную войну" американцы много раз через организацию "дезы" увлекали нас на тупиковый путь в исследованиях. Чтобы мы деньги на ветер бросали. Вот пускай теперь антигравицапу поконструируют по нашей наводке!"

Следуя его логике, янки из НАСА вместе с гордыми британцами - полные идиоты. Добро, если так. Но как бы нам (пользуясь лексикой из того же популярного фильма) не пришлось в конце концов приседать перед ними с растерянным возгласом "Ку!".

Андрей Самохин

В подготовке статьи использованы материалы из Интернета: аналитической службы ПЕРсТ и Александра Шермана.

* Американский журнал "Вайред" посвящен в основном проблемам компьютеров и Интернета, но уделяет немало внимания открытиям и изобретениям и в других областях.

**Эксперимент готовили со всей тщательностью: приборы, измеряющие гравитацию с точностью до восьмого знака, 8 космических спутников, учитывающих слабые изменения гравитационного поля Земли, вызванные приливами и отливами... Гравитацию измеряли в вакууме, в огромном металлическом ящике, экранированном от электромагнитного поля.

Дата 20000404 Дата загрузки 20000405 Время загрузки 0031 Источник Vesti.Ru, Москва
Заголовок **МИФ О ЖИРАФЕ** Автор Никита Максимов Тема Наука Примечания
<http://vesti.ru/nauka/2000/04/04/giraffe/> Текст

Не ожидал столь бурной реакции от читателей по поводу [<http://vesti.ru/nauka/2000/03/31/women/>] первоапрельской статьи. Борис Баанов вспомнил о статье, посвященной жирафе, в которой доказывалась невозможность ее существования. Степан Пачиков приводит более удачную редакцию анекдота про тещу:

Мальчик: Папа, почему бабушка бегает по огороду так странно?

Отец: Кому - бабушка, а кому - теща. Подай лучше, сынок, еще патронов.

Но основной же поток писем связан с просьбой рассказать об истории с жирафой. Я поставлен в очень неловкое положение - с одной стороны, уже не первое апреля, а с другой - дожидаться следующего дня смеха глупо. Выручили меня, как всегда, информационные агентства. Появилось [<http://www.gazeta.ru/podkletnov.shtml>] сразу два [http://www.nauka.newmail.ru/news_0204.html] сообщения о так называемом "антигравитационном эффекте Евгения Подклетнова". История с уменьшением на несколько процентов сил гравитации началась еще года четыре назад. Даже Wired сподобился на очень большую [<http://www.wired.com/wired/archive/6.03/antigravity.html>] статью о русском физике-гении. Но тогда все окончилось ничем - серьезного научного обсуждения работа Подклетнова не выдержала. На сей раз британская компания BAe Systems решила начать исследовательские работы по применению антигравитационного эффекта Подклетнова. Удивительные все-таки эти люди, англичане...

Вернусь, однако, к жирафе. В журнале "Знание - сила" была опубликована такая статья:

Читателя, знакомого с методами настоящей научной дискуссии, не надо убеждать, что жирафы не существуют. Ему достаточно четко и ясно сказать: "Ее нет и не было!" Подобный метод доказательств носит название "принцип Лавуазье" - в честь знаменитого химика. Как известно, он отрицал существование метеоритов на том основании, что "камни не могут падать с неба, потому что на небе нет камней". Среди последователей Лавуазье был и Симон Ньюкомб, который отверг идею создания летательных аппаратов тяжелее воздуха потому, что аппараты тяжелее воздуха летать не могут. Независимо от великого французского ученого так называемый принцип Лавуазье был сформулирован также в России: "Этого не может быть, потому что не может быть никогда".

Первое упоминание о "жирафе" - мифологическом звере с неестественно длинной шеей - мы находим у арабского путешественника Аль Кварима. Вслед за Аль Кваримом многие путешественники уверяли, что они видели - а некоторые даже фотографировали - это "длинношеене". Совсем недавно журнал "Планета", рискуя ввести в заблуждение излишне доверчивого читателя, несмотря на протесты всемирно известного научного деятеля Андре Парено, опубликовал статью, где легенда о жирафе трактуется как непреложная истина. Памятая о "принципе Лавуазье", мы ограничимся утверждением, что жирафа не существует, и основное внимание уделим тому, каким образом могла возникнуть "Легенда о жирафе". Общеизвестно, что в Африке, которой приписывается сомнительная честь быть родиной этого гипотетического животного, довольно часто наблюдаются явления, известные под названием фата моргана, или мираж, когда наблюдатель видит отдаленные предметы в

местах, где на самом деле они отсутствуют. При этом сам предмет или часть его могут предстать перед наблюдателем в деформированном виде - вытянутыми в горизонтальном и вертикальном направлении, как в кривом зеркале. Естественно предположить, что поводом для возникновения версии о существовании жирафы послужил мираж, из-за которого обычное, хорошо изученное животное, например, лошадь, могло представиться путешественнику в искаженном виде - с непомерно длинной шеей.

Действие фата морганы могло быть усугублено фактором чисто психологическим. Исследователю, заблудившемуся в пустыне и мучимому жаждой, в условиях действия высокой температуры и заторможенного сознания могло начать казаться, что его шея растет и вытягивается до размеров, позволяющих ему дотянуться до ближайшего оазиса. В этих условиях немудрено, что фантазия путешественника на основании увиденных им миражей могла создать образ животного со столь же длинной шеей.

Польский исследователь Гржибовский в легенде о жирафе видит пробуждение биологической памяти человека о вымерших доисторических животных, тоже отличавшихся исключительно длинной шеей.

Следует упомянуть также о работе английского биоматика Дж. Гедвика, который, давая дальнейшее развитие мифу о жирафе, доказывает, что понятие "жирафы" могло быть введено физиками, когда они пробовали применять методы статистики к сравнительной патоанатомии парнокопытных. Тогда понятие "жирафы" - лишь некая абстракция, которая совершенно не предполагает существования материального носителя ее свойств.

Таким образом, мы видим, как применение современных методов позволяет объяснить возникновение легенды о жирафе с позиций науки, не прибегая к заведомо не выдерживающей критики гипотезе о реальном существовании жирафы. Применение этих методов может оказаться плодотворным при подходе к рассмотрению и других вопросов во всех областях знания.

В заключение хотелось бы отметить достопримечательный факт, ярко демонстрирующий единство здравого смысла и новейших достижений современной науки. Когда простому американскому фермеру показали рисунок, изображающий так называемую жирафу, он воскликнул: "Не может быть!" Разве это не служит наглядным подтверждением народной мудрости и близости к народу современной науки?

После публикации этой статьи появились два письма на эту тему:

"УВАЖАЕМАЯ РЕДАКЦИЯ! Мы постоянные подписчики журнала "Знание - сила". Очень любим замечательные материалы, сопровождаемые иллюстрациями. Убедительно просим вас разъяснить нам следующее: В журнале № 5 за май месяц 1967 года на странице 28 ("Веселая наука") помещено "ЖИРАФА? НЕТ, МИФ!"

Автор статьи не указан. Надо полагать, что это статья редакционная. В статье говорится о том, что жирафы не существует, что ее нет и не было! Среди учителей нашего города эта статья вызвала целое смятение. Нашлись такие, что "лично видели" это животное в Африке. Другие утверждают, что в английском зоопарке эти жирафы спокойно проживают и т.д. Просим вас разъяснить нам этот вопрос. Ждем ответа по адресу:

Пенсионеры ПОВЕЛИЦА, РУДЕНКО

УВАЖАЕМЫЙ ТОВ. РЕДАКТОР! В номере 5 Вашего журнала опубликована статья "Жирафа? Нет, миф!" Считаю необходимым отметить своевременность и актуальность затронутой в указанной статье темы - разоблачения антинаучного мифа о жирафе.

Животное, обладающее столь длинной шеей, не имело бы никаких шансов выжить в процессе длительной эволюции. В самом деле, известно, что наиболее уязвимым местом травоядного является шея, и следовательно, животное, обладающее длинной шеей, имеет во столько раз меньше шансов выжить в процессе борьбы за существование, во сколько раз его шея длиннее обычной средней нормы для животных такой величины.

С уважением У. ЧЕНОВ.

А спустя два года в журнале появился вот такой текст:

ОТ ПРЕЗИДИУМА Академии Веселых Наук:

Мы празднуем своеобразный юбилей. Ровно два года назад появилась публикация, положившая начало нашей Академии (см. "Знание - сила", № 5, 1967, стр. 231). Некоторые считали, что проблемы жирафологии не найдут горячего отклика. Но маловеры и на этот раз были посрамлены. Письма со всех концов страны стекались в редакцию. Звонил телефон. Приходили телеграммы. Некоторые читатели являлись лично. И у всех на устах был один вопрос: "Да или нет?"

Мы - тогда еще не академики и даже не член-корреспонденты - без устали вскрывали конверты с вырезками из "Огонька", "Вокруг света", "Науки и жизни", на которых красовались длинношеие животные из зоопарков Копенгагена, Токио и даже города Дадли. Как же так, недоумевали авторы писем, жирафы - нет, а фотографируют! Один только - "Огонек" трижды (№№ 19, 33 в 1967 году) порадовал своих читателей видом жираф и жирафов. Назревала дискуссия, и мы не ушли от нее. Дав немного утихнуть нездоровому ажиотажу, в № 6 за прошлый год (см. стр. 47) опубликовали письмо бывших учителей, ныне пенсионеров, из Новомосковска Днепропетровской области и ответ на него У. Ченова. Таким образом, были представлены обе противоборствующие точки зрения: "жирафа есть" и "жирафы нет".

Костер полемики вспыхнул. Наши новомосковские корреспонденты ответили немедленно: "Мы ожидаем более полного разъяснения на страницах вашего журнала и со стороны У. Ченова, потому что наш ответ детям должен быть точным - существует ли такое животное, или же это миф". И хотя сам У. Ченов по понятным причинам от дальнейших выступлений воздержался, недостатка в защитниках его - как, впрочем, и противоположной - точки зрения не было. Но обработав получаемую корреспонденцию с помощью ЭВМ и математико-статистических методов, мы увидели, что идейных противников в процентном отношении не так уж и много. Основная масса жирафологов оказалась просто дезориентированной. Вот только некоторые отрывки из писем, демонстрирующие всю меру отчаяния их авторов.

"Дорогая редакция! Я являюсь постоянным подписчиком Вашего журнала и с интересом его читаю, находя много нового и интересного. Письмо У. Ченова, а также заметка в № 5 за 1967 г. и все прочитанное мною в других журналах ввели меня в полное заблуждение. Например, журнал "Наука и жизнь", № 6 за 1968 год на стр. 125 не только не отвергает существования жирафы, но даже помещает фотографию с сидящим верхом человеком. Еще более точное описание жирафы дает "Календарь школьника" за 1969 год, выпущенный "Полигиздатом". На листке от 11 марта написано, что "нескладные" шеи у жирафов отлично складываются. А на листке 20 июля под тем же заголовком пояснено, что спят жирафы - лишь 20 минут в сутки. Объясните, пожалуйста, чему верить? С уважением, В.В., агроном, п/о Светлобово, Усть-Уфимского р-на Иркутской области".

"Уважаемая редакция! Помещенная на страницах журнала (№ 5, 1967 г.) статья "Жирафа? Нет, миф!" вызывает явное недоумение. Допустим, что неизвестный автор статьи наконец "открыл" миру глаза, но становится совершенно непонятным, чем же вызван огромный поток информации, предложенный нам другими источниками, повествующими о жизни жираф. Так, в журнале "Наука и жизнь" (№ 1, 1968 г. на стр. 61) помещена удивительно поэтическая фотография двух жираф, поистине упоенных счастьем своего существования. В комментариях к фотографии отмечено, что жирафы прописаны в Экваториальной Африке, обитают в степных районах с редкой растительностью.

Далее в журнале № 8 за 1967 год на стр. 138 иллюстрирован поединок жирафа, в № 11 за 1966 год на стр. 45 все того же журнала "Наука и жизнь" изображена группа жираф, среди которых не только пятнистые, но и белая. Это уже совсем не в пользу неизвестного автора. И еще, в книге "В мире занимательных фактов" (издательство "Казахстан", Алма-Ата, 1965 год) на стр. 283 утверждается, что жирафа может не пить шесть месяцев и есть акации с острыми колючками. В этой же книге на стр. 299 говорится, что жирафа, будучи близоруким и длинношеим животным, нередко задевает за провода, срывает их и, бросившись от испуга, запутывает десятки метров проводов.

Подчеркивается также, что жирафы обитают в Южной Африке. Приведенные факты вселяют сомнение в том, что жираф не существует. Ведь иначе откуда взяться фотографиям и прочим данным? Убедительно просим журнал "Знание - сила" осветить этот вопрос как можно конкретнее. Братья Т., Анатолий и Михаил, г. Кишинев".

"Уважаемая редакция! В журнале "Вокруг света" № 8 напечатана фотография обитательницы Токийского зоопарка жирафы Томи с детенышем. Как же объяснить статью, напечатанную в Вашем журнале? Так как достать Ваш журнал в киоске бывает очень трудно, а я боюсь пропустить Ваш ответ, очень Вас прошу ответить по адресу: г. Казань..."

Иногда в письмах содержались конструктивные предложения:

"Уважаемая редакция! Пастухи нашего колхоза не согласны с утверждением читателя У. Ченова. Они говорят, что "длинная шея животного - замечательное приспособление, выработавшееся в процессе борьбы за существование. Подножный корм весь выеден, можно и листочками деревьев питаться. Если бы вывели породу овец с длинными шеями, хотя бы в 2-3 м, была бы благодать! Овцы очень любят листики осины и березы, даже к зиме запасают много веников из осин и березы, чтобы скормливать овцам. А как бы увеличилась кормовая база!" Правы колхозники. Д. А., Вологодская обл., Кадуйский район, с. Пречистое".

Иногда - корректный вопрос.

"Я лично сам не могу быть уверенным, что жирафы есть. Так как нет веских доказательств. Поэтому я и написал это письмо, чтобы узнать истину. Ю. Л., г. Шемонаиха".

Но порой - резкий окрик:

"Простите за беспокойство, но меня тревожит подрыв авторитета вашего журнала среди многих его читателей. Знание - сила! Казалось бы, эти два слова должны говорить сами за себя. Но почему же тогда в нем помещаются такие несоответствующие правде (если не сказать больше) статьи, как вызвавшая смятение статья "Жирафа? Нет, миф!", напечатанная в № 5 за 1967г., о разъяснении которой Вас уже просили пенсионеры из г. Новомосковска т.т. Повелица и Руденко. Не хочется думать, что журнал в будущем будет ставить на карту свою солидность и ради заполнения пустых мест печатать подобные вымыслы. А. А., г. Таллин".

Но лишь когда пришло полное неверия в жизнь письмо: "Где же правда? Кому верить? Кто кого вводит в заблуждение? Ю.Д., г. Ленинград" редакция призналась в мистификации.

Но шутка этим не закончилась... В 1976 году, когда журналу исполнялось 50 лет, редакция решила выпустить юбилейный номер с жирафом, прогуливающейся по 2-му Волконскому переулку в Москве. Однако цензура "зарезала" номер, не разрешив его отпечатать. Неудивительно, что после всех этих злоключений жирафа стала негласным символом журнала "Знание - сила" и вручается в уменьшенном размере всем отличившимся...

А вы говорите, антигравитация...

За сим разрешите откланяться. ОБЗОР ПРЕССЫ:

Газета "Наука в Сибири" весело [<http://www-sbras.nsc.ru/HBC/2000/n13/f32.html>] вспоминает 1-е апреля.

журнал "Огонек": [<http://www.ropnet.ru/ogonyok/win/200011/11-42-43.html>] Тело не мыть - долго не жить!

"Общая газета": [<http://www.og.ru/archieve/10/mat/na1.shtml>] Судьба Нобелей в России

С директором Главного ботанического сада РАН Львом Андреевым в "Московских новостях" [<http://www.mn.ru/2000/12/202.html>] беседует писатель и журналист Владимир Губарев.

[<http://science.ng.ru/current/>] "НГ-наука":

Мультимедийный вариант наскальной живописи
В пространстве науки и образования
Жизнь и смерть радиоактивных отходов
Молекулярная информация: миф или реальность?
Несколько замечаний по поводу "миссии российской науки"
Бистабильный Каспий
Разведчик озоновых дыр
АЭС учится плавать
На пути к искусственному интеллекту
Куда податься тугодуму?

Газета "Сегодня": Японские мистики
[http://www.segodnya.ru/w3s.nsf/Contents/2000_73_life_col_apostolova4.html] демонстрируют возможности человека.

Газета "Известия":
Российские биологи [<http://www.izvestia.ru/izvestia/article/213089>] добились сенсационного результата, к которому ученые всего мира стремились несколько десятилетий.

[http://www.accessnet.ru/vivovoco/VV/JOURNAL/NATURE/03_00/GINZBURG.HTM]
Ответы на вопросы журнала Physics World Б.Л. Гинзбурга в журнале "Природа"

ССЫЛКИ ПО ТЕМЕ НОВОСТИ

Швейцарские физики впервые непосредственно [<http://focus.aps.org/v5/st12.html>] измерили величину силового взаимодействия между индивидуальными атомами. Сотрудники Базельского университета с помощью силового сканирующего микроскопа определили силу притяжения между двумя атомами кремния, разделенными расстоянием в несколько атомных диаметров.

В Институте катализа СО РАН завершена работа над новым проектом - созданием теплого окна, не пропускающего холодный воздух и идеально сохраняющего тепло в доме. Создан сверхновый материал - аэрогель диоксида кремния, с помощью которого стало возможным полностью устраниТЬ утечку тепла. Самый лучший на сегодняшний день теплоизолятор имеет гелеобразное состояние, но при этом бесцветен, у него нет запаха, его удельный вес максимально приближен к весу воздуха. [<mailto:>] По сообщению [<mailto:agency@online.nsk.su>] Агентства общественных связей "БЮРО ПРОПАГАНДЫ"

"Миф о жирафе", 05.04.2000
<http://vesti.ru/nauka/2000/04/04/giraffe/>
(C) 2000 Vesti.Ru/Наука

Дата 20000404 Дата загрузки 20000406 Время загрузки 0301 Источник Симбирский курьер, Ульяновск Заголовок **В ПОПЫТКЕ ИЗМЕНИТЬ СИЛУ ТЯЖЕСТИ НА ЗЕМЛЕ**
Выпуск 50 (1627) Текст

Известная британская военно-промышленная компания "BAE Systems" заявила о том, что будет разрабатывать устройство, уменьшающее силу притяжения. Возможно, в скором будущем человечество сможет долететь до Луны на одном литре горючего.

Идея создания такого устройства впервые была выдвинута российским физиком Евгением Подклетновым. Подклетнов, выпускник Менделеевского института в Москве,

проработал 15 лет в Институте высоких температур АН СССР, а в 1988 году был приглашен в университет Тампера (Финляндия). Там, изучая свойства сверхпроводников, он вращал диск из сверхпроводящей керамики над мощным электромагнитом. Случайно он заметил, что вес предметов, помещенных над диском, уменьшается на 2%.

В 1992 году Подклетнов опубликовал предварительное сообщение о новом эффекте. Однако в профессиональной среде его открытие подвергли резкой критике, а с работы за занятия посторонними делами уволили. Сейчас российский физик живет в Тампера и работает в одной из местных фирм, продолжая свои опыты. На днях "BAE Systems" объявила о своем решении начать разработки по созданию гравитационного экрана, уменьшающего силу тяжести.

Однако в России по-прежнему скептически относятся к изобретению Подклетнова, считая это бредом. По мнению доктора наук Бориса Шустова, если в одном месте установить гравитационный экран и уменьшить тем самым силу тяжести, то в другом месте в соответствии с законом сохранения энергии гравитация обязательно увеличится. Кроме того, над таким гравитационным экраном будет большое скопление воздуха, в результате чего "Земля может лишиться атмосферного слоя". "Но с другой стороны, - иронизирует Шустов, - космонавты смогут долететь до Луны, потратив всего 1 кг горючего. Да и передвигать тяжести, например целые платформы, будет под силу даже ребенку".

"Симбирский КУРЬЕР" N 50(1627) 4 апреля 2000г., г.Ульяновск Тел.: (8422) 41-12-23, 41-14-00, 41-13-98, факс 41-12-11 E-mail: skos@nova-s.ul.ru, <http://ck.ul.ru>

Дата 20000401 Дата загрузки 20000331 Время загрузки 1617 Источник Якутия, Якутск
Заголовок **РОЖДАЕТСЯ УСТРОЙСТВО, УМЕНЬШАЮЩЕЕ ГРАВИТАЦИЮ** Автор
ОЛЕГ ЕМЕЛЬЯНОВ Тема С МИРУ ПО ФАЙЛУ Выпуск 59 Текст

Британская компания оборонного комплекса BAe Systems подтвердила намерение финансировать разработку устройства ученых Института физико-технических проблем Севера, которое позволит изменять силу тяжести.

Идея такого устройства впервые была предложена российским физиком Евгением Подклетновым. В ходе проведенных экспериментов удавалось снизить силу тяжести на 2% и более, что может иметь огромные практические последствия.

BAe Systems уже присвоила программе исследования антигравитации название Проект Гринглоу.

Сама идея возможности изменения одного из основополагающих физических явлений вызывает скептицизм у многих ученых, которые склонны относить эти исследования к области научной фантастики. Именно поэтому Российская академия наук отказалась в финансировании проекта, и якутским ученым пришлось обратиться к британским коллегам.

Дежурный по Интернету ОЛЕГ ЕМЕЛЬЯНОВ.

Дата 19991008 Дата загрузки 19991012 Время загрузки 1822 Источник Рыбак Приморья, Владивосток Заголовок **ПО ВОДЕ, АКИ ПОСУХУ** Выпуск 41 Текст
ПОЛЕТЫ НАЯВУ

Как известно - мир полон чудес. Имеются и в наши дни удивительные и непознанные для всех нас с вами тайны и загадки. Возможно их полное отрицание или безоговорочное признание. И, наконец, попытка понять и объяснить их. Для науки, конечно, важнее последнее.

Сегодня мы остановимся на хождении человека по водам, словно посуху, т.е. о левитации, - сверхъестественной его способности становиться легким по собственному желанию. Феномен же левитации, по утверждению различных исторических источников, наблюдался неоднократно. Так, во времена молитвенного экстаза в воздух поднимались многие святые и религиозные деятели.

Если же верить преданиям, то не один Христос мог ходить по воде. В Евангелии от Матфея можно прочесть: "Петр вышел из лодки и пошел по воде, чтобы подойти к Христу.

Но, видя сильный ветер, испугался и начал утопать, закричал: "Господи! Спаси меня". Иисус тотчас простер руку, поддержал его и говорит: "Маловерный! Зачем ты усомнился?.."

В приведенном примере необходимо обратить внимание на силу веры, а если точнее, на психологический фактор, который, по наблюдениям, в исключительных обстоятельствах приводит к резкому уменьшению массы тела (к левитации), когда человек вдруг становится легче воды и не тонет в ней. Из отечественных "левитантов" называют Серафима Саровского, архиепископа Новгорода и Пскова Иоанна... Московские же летописцы рассказывают о Василии Блаженном, который не раз на глазах у изумленной толпы переносился через Москву-реку.

По ряду исторических источников католические святые ничем не уступали нашим. Так, один испанский поломник умудрился перелететь через широкую реку, а епископ Валенсии парил над землей порядка 12 часов. Всего же, как утверждают церковные книги, количество людей, которые на глазах у верующих передвигались по воздуху и ходили по воде, приближается к трем сотням. А сколько ведь во времена инквизиции за то же самое горело на костре?

В XIX веке российский журнал "Ребус", который специализировался на различных таинственных явлениях, опубликовал материал о случае, произошедшем с Н.Юрловым, когда тому было 8 лет. 19 июля 1837 года мальчик спал. Ночью он был разбужен грозой и к ужасу своему увидел, что недалеко от его кровати на балконе, держась за ручку стеклянной двери, стоит высокий лысый человек, с седой бородой, в длинной синей рубашке... От ужаса мальчик скатился вниз по лестнице со второго этажа дома, выбежал из него и, ничего не соображая, стремительно побежал в сторону реки Свияги, протекавшей у села Спешневки Сенгилеевского уезда, Симбирской губернии. Через некоторое время мальчика нашли на другом берегу реки совершенно сухим. Домашние видели, как он переносился через реку, едва касаясь воды...

О подобных явлениях рассказывалось в 1900 году в сборнике "Из области таинственного". Так, например, в нем шла речь об известном итальянском церковном деятеле Бенидикте Нурсийском и его ученике Маурусе, который, увидев, как на воде тонет мальчик, бросился к нему на помощь. "Только вернувшись на землю, он пришел в себя и посмотрел назад. Когда же он увидел, что пробежал по воде, то страшно испугался происшедшему..."

Одним из известных левитантов прошлого века считают англичанина Даниэля Дугласа Юма. Очевидцы описали выше ста случаев его полетов по воздуху. Случаи левитации случались и с солдатами в годы Великой Отечественной войны. Так, одна из бывших участниц этой войны рассказала:

- Когда наши войска форсировали Днепр, я была радиисткой. Мы переправлялись через реку под ураганным огнем. На меня напал неописуемый страх. Плыла вдвоем с солдатом на какой-то лодочки. Вдруг бах - и мы оказались в воде. Я плавать не умею, ухватилась за ватник солдата, но он оттолкнул меня, и я стала тонуть. Помню, как какая-то сила вытолкнула меня на поверхность, и вода подо мною вдруг стала плотной, как земля. Вскоре я, сама не знаю как, очутилась на том берегу и стала благодарить Господа за чудесное спасение.

То, что фронтовичка восприняла как чудо, для жрецов некоторых племен Индии является делом обыденным. Они умеют приводить свою психику в такое состояние, которое вызывает левитацию, то есть резкую потерю массы своего тела. Долго жившая в Индии писательница Л.Шапошникова рассказывает: когда она плыла на пароходе по реке, то увидела, как бородатый старик в чалме и набедренной повязке перешел на противоположный берег по воде, не замочив ноги.

Любое явление требует научного объяснения. До недавнего времени официальная наука относилась к сообщениям о феноменах левитации со значительным скепсисом. Но это положение начало изменяться после серии опытов, проведенных российским физиком Евгением Подклетновым в лаборатории технологического университета финского города

Тампера. Специальный диск охлаждался физиком до -167оС и помещался в электромагнитном поле, заставляющем его вращаться. При достижении 3000 лб/м предметы, помещенные под вращающимся диском, начали терять свою массу. Сообщение в печати об этом эксперименте послужило своеобразным детонатором, после которого и другие ученые стали проводить свои опыты и сообщать о наблюдаемых ими антигравитационных эффектах. Любопытные результаты по данному вопросу были получены Джоном Шнурером из Энтиосского колледжа, что в штате Огайо (США) и рядом голландских ученых и особо доктором Андре Гейном и его помощниками из Университета города Наннели.

Однако в специальной литературе по проблемам левитации встречаются и такие утверждения: она порой достигает столь высокой степени, что человек становится массой с пушинку и приобретает возможность не только ходить по воде, но даже летать в пространстве без каких-либо технических приспособлений. Это одно из самых загадочных, схожих с мистикой проявлений биоэнергетики человека. Так, немецкая газета "Вельт ам зонитаг" несколько лет тому назад на своих страницах рассказывала о том, что англичанин Роберт Стюарт, путешествуя по Монголии, посетил буддийский монастырь, расположенный в пустынной местности в Южно-Гобийском аймане и во время богослужения он наблюдал загадочное явление. Монахи зажгли множество ароматных курительных палочек, затем стали читать молитвы. Вдруг Стюарт увидел, как один из послушников начал медленно отрываться от земли. Поднявшись на высоту примерно трех метров он остановился, а затем начал медленно опускаться на землю.

Необъясненным пока остается феномен левитирования людей, овладевших учением о трансцендентальной медитации, созданным в начале 50-х годов Махариши Махен Ионом (в мире - Махариши). Внешне это выглядит как подскоки в воздух в положении "поза лотоса".

В американском городе Вашингтоне в 1986 году проходило соревнование "летающих йогов", в котором принимало участие около 20 человек. В программу этих соревнований входили полеты в высоту (был достигнут рекорд около 60 см), полеты в длину (рекорд - 1.8 м) и полеты на скорость на дистанции 25 м с препятствием высотой 20 см. Парапсихологи сопоставляют прыжки "летающих йогов" с проявлением феномена левитации. Физики же уверяют, что секрет кроется совершенно в другом.

Какая же из этих гипотез ближе к истине, нам думается, рассудит время.

Дата 19990401 Дата загрузки 19990429 Время загрузки 1630 Источник Наше время, Рига, Латвия Заголовок "**ЗАБУДЬТЕ ОБ ЭЙНШТЕЙНЕ! Я ЕГО ОТМЕНИЛ**" Выпуск 013 Текст

Года два назад один из лидеров Государственной думы, известный своими невоздержанными высказываниями, выехал за пределы отчизны и там на весь мир заявил, что Россия обладает неким фантастическим сверхоружием. Цивилизованный мир, у которого и от "обычных"-то ядерных ракет слабеющей России голова трещит, ужаснулся.

Лидером тем был Жириновский, а под сверхоружием он имел в виду гравитационные технологии, о которых прочел в журнале известного политического оракула Кургиняна, славного своей любовью к мифам. Тем не менее зерно истины в публикации было. Имя зерна - Поляков.

Поляков - личность в определенных кругах легендарная. И иностранцы кино о нем снимали. И в военных кругах его знают как непризнанного гения и исследователя неожженых научных троп. После окончания Кишиневского университета Поляков с дипломом, в котором гордо красовалось - "физик широкого профиля", попал по распределению во Фрязино, в "Исток", один из самых знаменитых наших "ящиков". Из "Истока" вытекла вся отечественная электроника, первые масс-спектрографы, лазеры и полупроводники. Все электронные предприятия отечества имели началом "Исток". Вот какое это славное было предприятие! Именно там ковал оборонный щит родины Поляков, там в свободное от ковки время и увлекся проблемами гравитации. Должно же быть у человека приличное хобби! Не марки же физику широкого профиля собирать!

Сейчас Поляков давно уже на пенсии, но продолжает исправно ходить на родное предприятие, благо не гонят. Там он садится на стул в лаборатории и начинает ждать чуда. Телефонных звонков. Он, конечно, видит, что все кругом перевернулось, но в глубине души не верит, что его новая физика никому так уж и не нужна.

...Сразу видно, что Фрязино - город "почтовых ящиков". Нечищенные дороги, разбитый асфальт, какие-то черные люди, неизвестно зачем и куда бредущие по мрачным улицам, перекошенные оранжевые автобусы, по самую крышу заляпанные серой грязью. К подъезду Полякова я проехать не смог: не на тракторе приехал. Бросил машину, дошел пешком.

Квартира пенсионера Полякова под стать городу. В ней пахло кошками и вообще видно было, что живут тут холостяки - отец с сыном.

- Вы извините, у нас тут немного не убрано. Два года назад умерла моя жена, и мы с сыном еще никак в себя не придем.

Конечно, их мир рухнул. И даже не два года назад. Гораздо раньше.

...Началось все с неутоленной душевной жажды. Не может русский человек со своей загадочной душой просто работать.

Решение хоть и творческих, но сугубо прикладных задач не может успокоить русскую душу. И тут два выхода есть у души - либо водку пить, либо заниматься проблемами мироздания. Гуманитарии в ситуации душевной неуспокоенности всегда смысл жизни ищут, в религию ударяются, а люди образованные... Например, проблемы гравитации решают.

Мало чего знает современная физика о гравитации. Меньше, чем отдельно взятый Поляков, который сидит передо мной на полуразваленном стуле и курит самые дешевые сигареты без фильтра. Интересно курит: отрывается от газеты, покрывающей стол, часть текста, сворачивает из обрывка мундштук, засовывает в него сигарету и курит. Так экономнее получается: сигарету можно до упора выкурить, до последней табачинки - и губы не обжигает. Зато коричневеет от сигаретного жара конец газетного мундштука. Ну да ничего, газеты на столе еще много, можно скрутить сколько угодно таких мундштуков.

За квартиру Спартак Михайлович уже полтора года не платит. И за свет. И за газ. И за телефон. Поэтому домашний телефон ему давно отключили. Вся стариковская пенсия уходит на еду и исследования. Да вот еще недавно взятку дал электрику, чтобы свет за неуплату ему не отрубил. Обидно: непроизводительные расходы. Гараж свой Поляков продал еще года три назад и деньги тоже вбухал в научную работу.

А началось-то все с пустяка. Задумался как-то молодой "физик широкого профиля" о том, что такое квант света. Видимо, это оттого произошло, что недоучили его. Обычно дипломированные специалисты подобными вопросами не задаются, им еще на студенческой скамье все правильно объясняют. Никто, в общем-то, не знает, что такое "поле", "заряд", "квант", но глупых вопросов тем не менее не задают. Потому что уже к старшим курсам студент привыкает к терминам и считает эту свою привычку знанием.

А Поляков привычку за знание считать отказывался. У него вообще о знании особые представления:

- Знания - это когда вы можете сделать вещь из того, что валяется под ногами и никому не нужно. Вот, например, физики пользуются своими понятиями, но не знают их. Если я могу пользоваться телевизором, это не значит, что я знаю, как он устроен. А вот когда я его сам смогу собрать, тогда да, это знание...

Короче говоря, для начала построил Поляков новую модель фотона, состоящего из 137 длин волн, с нее плавно на электродинамику перешел, а потом на гравитацию. Так хорошо все шло! Ранее плохообъяснимые факты, типа магнитострикции или оптического намагничивания, легко и просто объяснялись в рамках теории Полякова, каковая, кстати, вошла в полное противоречие с теорией относительности. У Эйнштейна скорость света - предельная величина, а у Полякова скорость распространения гравитационной волны на десятки порядков превышает скорость света... Ах, лучше бы я не упоминал при Полякове имени великого физика!

- Да х.. с ним, с Эйнштейном! Забудьте о нем! - взорвался до того тишины Спартак Михайлович. - Наврал он все, и об этом сейчас только ленивый не говорит! Вот профессор Логунов, бывший ректор МГУ, всегда против теории Эйнштейна выступал. А я так вообще опытно доказал, что скорость гравитационной волны в 500 миллионов раз превышает скорость света. По изменению импульса на своей установке доказал! Вот они, результаты...

От установки так просто не отмахнешься. А она стоит себе на его родном предприятии, гудит, работает...

Понимаете, из поляковской физики следует не только, что гравитация быстрее скорости света. Из нее вытекает еще и крепкая взаимосвязь вращения и гравитации. Вращающееся тело, по Полякову, излучает гравитационные волны. Из этого много чего вытекает. Во-первых, принципиально новые виды связи. Во-вторых, новые виды двигателей и источники энергии. В-третьих, новые виды оружия. Короче, очередной переворот в физике, технологии, жизни на Земле. Сравнимый с вхождением в нашу жизнь электричества.

Здесь необходимо небольшое отвлечение, друзья мои... На протяжении последних 10-15 лет я с любопытством наблюдаю, как многие любители науки и серьезные ученые увлеклись связью вращения с тяготением. Попадаются среди этих увлеченных откровенные шизофреники, недоучившиеся инженеры, но есть и люди с нормальным физическим образованием и даже работающие в академических институтах. Что-то в этом есть, черт возьми! Какая-то надинтуиция коллективного бессознательного. Какой-то научный сюр, предвосхищение мысли, парящей в воздухе. Полет шмеля за секунду до пробуждения. Это должно было чем-то кончиться. Может быть, и кончилось практической установкой Полякова? И началось в качестве новой технологической эры?

Он сам ее строил, установку эту. Еще когда работал в своем "Истоке". Сам добывал приборы, сам вытачивал, сам фрезеровал, сам детали и материалы доставал.

- У меня руки тогда были - мог доски мозолями шлифовать.

Зато все у него получилось: удельная тяга, развиваемая вращением тяготеющей массы, - 2,5 кг на киловатт. Если Поляков не ошибся, это практически первый в истории гравитационный двигатель. Сам себя он пока еще в воздух не поднимает, потому что весит 40 кг, но для первой опытной установки это совсем неплохо, если учесть, что тяга современных вертолетных двигателей - до 8 кг/кВт.

Штукой, разумеется, заинтересовались военные, но к тому времени денег у армии уже не было не только на перспективные разработки, но и на зарплату. И Поляков, приезжая в Министерство обороны к заинтересовавшимся военным, привозил с собой молотый кофе - угождал оголодавших защитников. Кофе у Полякова в ту пору было много: сын работал на небольшой кофемольной фабричке, и дома кофею этого было просто завались - запах стоял по всем углам. Военные радовались и угождались поляковским кофе. С тех пор прошло еще года два.

И вот сегодня Поляков сидит передо мной, вставляет очередную сигарету в очередной газетный мундштук и мечтает:

- Я бы согласился отдать разработку в обмен на самосвал с деньгами. Продаю свой памятник - берите! Только дайте мне закончить исследования. Здесь всем хватит. Стадию честолюбия и жажды славы я уже давно прошел, старый стал. Всю жизнь ведь на это дело, считай, положил. Мне главное теперь - чтоб не пропало все.

Жизнь у изобретателей-фанатиков не сладкая, смею заверить. Один из знакомых Полякова - Коровяков, который тоже занимался схожими проблемами и на пару хороших мыслей Полякова натолкнул, на увлечении своем все потерял - благополучие, семью, работу. Полякову тут больше повезло. Его жена очень поддерживала. Любила, наверное. Была она учительницей английского и помогала мужу деньгами на исследования. Поляков показал мне ее портреты - на кухне, и в комнате, и в другой комнате. Хорошее лицо, доброе.

...Я ехал домой на машине с двигателем внутреннего сгорания и думал, что, может быть, еще доживу до времен, когда под капотом моего летающего лимузина разместится

антигравитационный двигатель. И слово "поляков" будет писаться с маленькой буквы, как сейчас пишут "дизель".

А.НИКОНОВ

Кстати: NASA выделило более полумиллиона долларов на проверку эффекта Подклетнова - очередного не признанного официальной наукой физика, который вроде бы открыл связь сверхпроводимости с антигравитацией. Охлажденный жидким гелием диск из сверхпроводящих материалов после раскручивания его магнитным полем начинал вдруг экранировать земное тяготение - у Подклетнова получалось, что вес тела, расположенного над вращающимся диском, уменьшается на 2. Пока что в лабораториях NASA эффект Подклетнова воспроизвести не удалось, но американцы очень на это надеются.

Дата 19990326 Дата загрузки 19990325 Время загрузки 2300 Источник Россия, Москва
Заголовок **И ГРАВИТАЦИЮ ПРЕОДОЛЕЕМ СКОРО** Выпуск 18 Текст

Идея победы человека над гравитацией и создания летательных аппаратов, неподвластных всемирному тяготению, давно стала почти обязательным атрибутом произведений в жанре научной фантастики. Однако эта идея мало - помалу начинает перекочевывать со страниц романов и киносценариев в программы работ научно-исследовательских центров. Национальное управление по аeronавтике и исследованию космического пространства (НАСА) - ведомство, не склонное заниматься беспочвенными фантазиями, - выделило 600 тыс. долларов на работы, связанные с поиском средств защиты от гравитационного воздействия. Деньги пойдут на изготовление сверхпроводящего диска диаметром в 31 сантиметр и на продолжение экспериментов по экранированию от гравитации. В случае успешного завершения проекта устройство сможет создать некий антигравитационный экран, который защитит летательный аппарат от поля тяготения. Исходной причиной для проведения этих работ стали эксперименты российского ученого, специалиста в области материаловедения Евгения Подклетнова, который, работая в начале 90-х годов в Технологическом университете Тампере (Финляндия) со сверхпроводниками, обнаружил явление частичной потери веса.

Дата 19990322 Дата загрузки 19990323 Время загрузки 2200 Источник Журнал Огонек, Москва
Заголовок **"ЗАБУДЬТЕ ОБ ЭЙНШТЕЙНЕ! Я ЕГО ОТМЕНИЛ"** Выпуск 12 Текст
СПАРТАК ПОЛЯКОВ, ГЕНИЙ ИЗ ФРЯЗИНА

Года два назад один из лидеров Государственной думы, известный своими невоздержанными высказываниями, выехал за пределы отчизны и там на весь мир заявил, что Россия обладает неким фантастическим сверхоружием. Цивилизованный мир, у которого и от "обычных" - то ядерных ракет слабеющей России голова трещит, ужаснулся.

Лидером тем был Жириновский, а под сверхоружием он имел в виду гравитационные технологии, о которых прочел в журнале известного политического оракула Кургиняна, славного своей любовью к мифам. Тем не менее зерно истины в публикации было. Имя зерна - Поляков.

Поляков - личность в определенных кругах легендарная. И иностранцы кино о нем снимали. И в военных кругах его знают как непризнанного гения и исследователя неожженых научных троп. После окончания Кишиневского университета Поляков с дипломом, в котором гордо красовалось - "физик широкого профиля", попал по распределению во Фрязино, в "Исток", один из самых знаменитых наших "ящиков". Из "Истока" вытекла вся отечественная электроника, первые масс-спектрографы, лазеры и полупроводники. Все электронные предприятия отечества имели началом "Исток". Вот какое это славное было предприятие! Именно там ковал оборонный щит родины Поляков, там в свободное от ковки время и увлекся проблемами гравитации. Должно же быть у человека приличное хобби! Не марки же физику широкого профиля собирать!

Сейчас Поляков давно уже на пенсии, но продолжает исправно ходить на родное предприятие, благо не гонят. Там он садится на стул в лаборатории и начинает ждать чуда.

Телефонных звонков. Он, конечно, видит, что все вокруг перевернулось, но в глубине души не верит, что его новая физика никому так уж и не нужна.

... Сразу видно, что Фрязино - город "почтовых ящиков". Нечищенные дороги, разбитый асфальт, какие-то черные люди, неизвестно зачем и куда бредущие по мрачным улицам, перекошенные оранжевые автобусы, по самую крышу заляпанные серой грязью. К подъезду Полякова я проехать не смог: не на тракторе приехал. Бросил машину, дошел пешком.

Квартира пенсионера Полякова под стать городу. В ней пахло кошками и вообще видно было, что живут тут холостяки - отец с сыном. Но и давние следы женской руки тоже можно было разглядеть за слоем грязи, огрызков и окурков. Когда-то здесь можно было жить, когда-то это все было даже красиво и уютно, пока не облетело, не оборвалось, не повисло беспомощно, не замечаемое хозяйственным глазом. Впрочем, нет, замечаемое...

- Вы извините, у нас тут немного не убрано. Два года назад умерла моя жена, и мы с сыном еще никак в себя не приDEM.

Конечно, их мир рухнул. И даже не два года назад. Гораздо раньше.

- Значит, вы из "Огонька"? - старик пожевал губами. - Журнал ваш закуплен Израилем на корню!

- Не знаю, не продавал... На кухне Spartak Mихайлович мне, как гостю, предложил неполоманный стул и заварил крепкого чаю прямо в бокале с отбитым краешком, накрыв бокал пепельницей - чтобы заварилось покруче. Отпить из него я не рискнул.

... Началось все с неутоленной душевной жажды. Не может русский человек со своей загадочной душой просто работать. Даже творчески. Скучно ему делать какие-то электронные лампы для ракет, пусть даже по самому сумасшедшему, почти невыполнимому заказу обронки. Зачем это нужно, чтобы электронная лампа для ядерной ракеты выдерживала перегрузки в 10 000 g, если ракета ускоряется максимум на 10 g? Тысячекратный запас прочности получается! И на чем испытывать такие лампы прикажете, перегрузки создавать? Для этого же нужно отдельные сверхцентрифуги строить, а это особые подшипники, особые смазки, отдельная отрасль промышленности... Так одна сверхпрочная лампочка лавиной тянет за собой целый шлейф проблем, целую экономику. А ракета же не из одних только ламп состоит.

Так вот и надорвалась советская экономика.

- Зато ракеты наши дубовые были! - вспоминает молодость Spartak Mихайлович. - По электронике мы от Запада лет на 15 отставали, по точности наши ракеты им всегда в два раза уступали, зато по прочности - все как у Собакевича было. Только прямое попадание в такую ракету могло сбить ее с курса. А на взрывы рядом она просто не реагировала...

Еще бы ей реагировать - с тысячекратным-то запасом прочности! Не зря Поляков лампу изобретал.

Но, повторимся, решение хоть и творческих, но сугубо прикладных задач не может успокоить русскую душу. И тут два выхода есть у души - либо водку пить, либо заниматься проблемами мироздания. Гуманитарии в ситуации душевной неуспокоенности всегда смысл жизни ищут, в религию ударяются, а люди образованные... Например, проблемы гравитации решают.

Мало чего знает современная физика о гравитации. Меньше, чем отдельно взятый Поляков, который сидит передо мной на полуразваленном стуле и курит самые дешевые сигареты без фильтра. Интересно курит: отрывает от газеты, покрывающей стол, часть текста, сворачивает из обрывка мундштук, засовывает в него сигарету и курит. Так экономнее получается: сигарету можно до упора выкурить, до последней табачинки - и губы не обжигает. Зато коричневеет от сигаретного жара конец газетного мундштука. Ну да ничего, газеты на столе еще много, можно скрутить сколько угодно таких мундштуков.

За квартиру Spartak Mихайлович уже полтора года не платит. И за свет. И за газ. И за телефон. Поэтому домашний телефон ему давно отключили. Вся старицкая пенсия уходит на еду и исследования. Да вот еще недавно взялку дал электрику, чтобы свет за неуплату ему

не отрубил. Обидно: непроизводительные расходы. Гараж свой Поляков продал еще года три назад и деньги тоже вбухал в научную работу.

А началось-то все с пустяка. Задумался как-то молодой "физик широкого профиля" о том, что такое квант света. Видимо, это оттого произошло, что недоучили его. Обычно дипломированные специалисты подобными вопросами не задаются, им еще на студенческой скамье все правильно объясняют. Никто, в общем-то, не знает, что такое "поле", "заряд", "квант", но глупых вопросов тем не менее не задают. Потому что уже к старшим курсам студент привыкает к терминам и считает эту свою привычку знанием.

А Поляков привычку за знание считать отказывался. У него вообще о знании особые представления:

- Знания - это когда вы можете сделать вещь из того, что валяется под ногами и никому не нужно. Вот, например, физики пользуются своими понятиями, но не знают их. Если я могу пользоваться телевизором, это не значит, что я знаю, как он устроен. А вот когда я его сам смогу собрать, тогда да, это знание...

Короче говоря, для начала построил Поляков новую модель фотона, состоящего из 137 длин волн, с нее плавно на электродинамику перешел, а потом на гравитацию. Так хорошо все шло! Ранее плохообъяснимые факты, типа магнитострикции или оптического намагничивания, легко и просто объяснялись в рамках теории Полякова, каковая, кстати, вошла в полное противоречие с теорией относительности. У Эйнштейна скорость света - предельная величина, а у Полякова скорость распространения гравитационной волны на десятки порядков превышает скорость света... Ах, лучше бы я не упоминал при Полякове имени великого физика!

- Да х.. с ним, с Эйнштейном! Забудьте о нем! - взорвался до того тишайший Спартак Михайлович. - Наврал он все, и об этом сейчас только ленивый не говорит! Вот профессор Логунов, бывший ректор МГУ, всегда против теории Эйнштейна выступал. А я так вообще опытно доказал, что скорость гравитационной волны в 500 миллионов раз превышает скорость света. По изменению импульса на своей установке доказал! Вот они, результаты...

Поляков горячится. Тыкает в свою книжку, которую издал за свой счет в начале 90-х. Когда книжка вышла, весь тираж хотели скупить некие коммерсанты. Но Поляков не продал!

- Они хотели под нож пустить весь тираж, эти представители Израиля! - негодует Спартак Михайлович.

- Зачем? - не понимаю я.

- А зачем они Союз развалили?..

На этот убийственный аргумент ответить нечего. Действительно, от людей, разваливших Советский Союз, можно ожидать чего угодно...

Ну да, не любит Поляков евреев. Такой странный эффект, эту загадочную aberrацию сознания я и раньше наблюдал у многих бывших работников Оборонпрома и военных. Наверное, это хорошая тема для психологической диссертации - "Влияние оборонного сознания на адекватность мировосприятия и развитие ксенофобии". Они, может, даже и не виноваты в своем антисемитизме, просто так сложилось...

Но ведь главное в Полякове не его неприличные взгляды на деяния семитских народов, а его опытная установка. Та, на которой он скорость гравитационной волны определил. В конце концов книжкой, на которую покушались проклятые сионисты, можно было бы ипренебречь, как и всеми поляковскими теориями (мало ли теорий люди выдумывают!), но вот от установки так просто не отмахнешься. А она стоит себе на его родном предприятии, гудит, работает. Понимаете, из поляковской физики следует не только, что гравитация быстрее скорости света. Из нее вытекает еще и крепкая взаимосвязь вращения и гравитации. Вращающееся тело, по Полякову, излучает гравитационные волны. Из этого многое чего вытекает. Во-первых, принципиально новые виды связи. Во-вторых, новые виды двигателей и источники энергии. В-третьих, новые виды оружия. Короче, очередной переворот в физике, технологии, жизни на Земле. Сравнимый с вхождением в нашу жизнь электричества.

Здесь необходимо небольшое отвлечение, друзья мои... На протяжении последних 10 - 15 лет я с любопытством наблюдаю, как многие любители науки и серьезные ученые увлеклись связью вращения с тяготением. Попадаются среди этих увлеченных откровенные шизофреники, недоучившиеся инженеры, но есть и люди с нормальным физическим образованием и даже работающие в академических институтах. Что-то в этом есть, черт возьми! Какая-то надинтуиция коллективного бессознательного. Какой-то научный сюр, предвосхищение мысли, парящей в воздухе. Полет шмеля за секунду до пробуждения. Это должно было чем-то кончиться. Может быть, и кончилось практической установкой Полякова? И началось в качестве новой технологической эры?

Он сам ее строил, установку эту. Еще когда работал в своем "Истоке". Сам добывал приборы, сам вытачивал, сам фрезеровал, сам детали и материалы доставал.

- У меня руки тогда были - мог доски мозолями шлифовать.

Зато все у него получилось: удельная тяга, развиваемая вращением тяготеющей массы, - 2,5 кг на киловатт. Если Поляков не ошибся, это практически первый в истории гравитационный двигатель. Сам себя он пока еще в воздух не поднимает, потому что весит 40 кг, но для первой опытной установки это совсем неплохо, если учесть, что тяга современных вертолетных двигателей - до 8 кг/кВт.

Штукой, разумеется, заинтересовались военные, но к тому времени денег у армии уже не было не только на перспективные разработки, но и на зарплату. И Поляков, приезжая в Минобороны к заинтересовавшимся военным, привозил с собой молотый кофе - угождал оголодавших защитников. Кофе у Полякова о ту пору было много: сын работал на небольшой кофемольной фабричке, и дома кофею этого было просто завались - запах стоял по всем углам. Военные радовались и угождались поляковским кофе. С тех пор прошло еще года два.

И вот сегодня Поляков сидит передо мной, вставляет очередную сигарету в очередной газетный мундштук и мечтает:

- Я бы согласился отдать разработку в обмен на самосвал с деньгами. Продаю свой памятник - берите! Только дайте мне закончить исследования. Здесь всем хватит. Стадию честолюбия и жажды славы я уже давно прошел, старый стал. Всю жизнь ведь на это дело, считай, положил. Мне главное теперь - чтоб не пропало все.

Жизнь у изобретателей-фанатиков не сладкая, смею заверить. Один из знакомых Полякова - Коровяков, который тоже занимался схожими проблемами и на пару хороших мыслей Полякова натолкнул, на увлечении своем все потерял - благополучие, семью, работу. Полякову тут больше повезло. Его жена очень поддерживала. Любила, наверное. Была она учительницей английского и помогала мужу деньгами на исследования. Поляков показал мне ее портреты - на кухне, и в комнате, и в другой комнате. Хорошее лицо, доброе.

... Я ехал домой на машине с двигателем внутреннего сгорания и думал, что, может быть, еще доживу до времен, когда под капотом моего летающего лимузина разместится антигравитационный двигатель. И слово "поляков" будет писаться с маленькой буквы, как сейчас пишут "дизель".

Александр НИКОНОВ КСТАТИ: НАСА выделило более полумиллиона долларов на проверку эффекта Подклетнова - очередного не признанного официальной наукой физика, который вроде бы открыл связь сверхпроводимости с антигравитацией. Охлажденный жидким гелием диск из сверхпроводящих материалов после раскручивания его магнитным полем начинал вдруг экранировать земное тяготение - у Подклетнова получалось, что вес тела, расположенного над вращающимся диском, уменьшается на 2 %. Пока что в лабораториях НАСА эффект Подклетнова воспроизвести не удалось, но американцы очень на это надеются.

Дата 19990312 Дата загрузки 19990312 Время загрузки 1100 Источник Известия, Москва Заголовок **ВЛАСТЬ ВСЕМИРНОГО ТЯГОТЕНИЯ** Автор Александр МАРГОЛИН Выпуск 43 Текст

Национальное аэрокосмическое агентство США (НАСА) выделило 600 тысяч долларов на создание защитного экрана от силы земного притяжения. В основу работ положены идеи российского физика Евгения Подклетнова.

Идея - та же, что в фантастическом романе Герберта Уэллса "Первые люди на Луне". Его герой, ученый-отшельник, изобрел вещество по имени "кейворит", непроницаемое для земного тяготения. Любые предметы, помещенные над кейворитом, лишались веса и свободно летели вверх. В мире, оборудованном таким экраном, появятся самолеты без крыльев, а морские танкеры пойдут на слом. Космические перевозки гигантских грузов станут пустячным делом. Кто владеет всемирным тяготением, тот владеет миром.

В авторитетнейшем научном журнале "Нью сайентист", далеком, кстати, от фантастики, сообщается, что НАСА намерено проверить спорные опыты русского ученого Евгения Подклетнова, который заявил, что открыл вещество, уменьшающее силу тяготения. Специалисты весьма скептически отнеслись к фантастическому заявлению. "Осторожно!" - говорил себе и я. Так недалеко и до зеленых человечков...

Прежде всего, кто такой Подклетнов? Обращаюсь с электронным запросом в НАСА. Почтовый робот подтверждает существование Евгения Подклетнова. Его идея в настоящее время действительно проверяется в одном из отделений НАСА, в Центре космических полетов имени Маршалла в Хантсвилле, штат Алабама.

Тампере, Финляндия

В 1988 году Евгения Подклетнова для работы над докторской диссертацией пригласили в Финляндию, в университет Тампере. За плечами у него было 15 лет работы в Институте высоких температур РАН. С собой он привез диск из сверхпроводника, который изготовил из расплава оксидов меди, иттрия и бария - совсем как в романе Уэллса. В диаметре диск достигал почти 30 сантиметров. Американцы, по их признанию, отстают от нас в изготовлении таких дисков.

По словам Подклетнова, открытие было сделано случайно, когда этот самый диск охладили до температуры сверхпроводимости с помощью жидкого гелия и раскрутили электромагнитами. В этот момент в лаборатории кто-то закурил, и все увидели, как облако табачного дыма втягивается в невидимую колонну над диском и столбом медленно поднимается к потолку. Примерно так же рассказывал о действии кейворита Герберт Уэллс сто лет назад.

Открывшиеся перспективы так ошеломили и увлекли Подклетнова, что он потерял покой. Остался в Финляндии и лихорадочно продолжал опыты. Оказалось, что все предметы, помещенные над диском, будь они из железа, бумаги, меди, глины, стекла и т. д., уменьшили свой вес на два процента. Это вовсе не маленький эффект: столбик из ста дисков уменьшил вес почти в десять раз. Вместе с финским соавтором Подклетнов послал статью об открытии в британский журнал "Джорнал о физикс". С этого момента начались злоключения.

Английские газетчики пронюхали о сенсационной, опровергающей законы физики статье. В сентябре 1995 года в "Санди телеграф" вышла разгромная заметка о сомнительных опытах. Руководство университета в Тампере отреагировало примерно так, как реагировали когда-то у нас на статьи в "Правде". Вместо того чтобы проверить результаты, работу запретили. Коллегу вынудили отказаться от соавторства. Подклетнова под угрозой увольнения заставили забрать статью из редакции, а затем все равно выгнали из университета. Знакомые отвернулись от него. Так он оказался изобретателем-одиночкой, а эффект уменьшения веса назвали его именем.

С тех пор он исчез из поля зрения широкой публики, превратившись в ученого-отшельника. Прячется от журналистов, избегая новых подвохов. Не дает интервью. По слухам, живет в Тампере, ходит иногда вокруг университета, куда его не пускают. И молчит.

Последователи

Морозы, грипп, кризис волнуют понятно почему. Но почему нас волнуют идеи? Иные сжигают людей, как пожар солому. И мы лишь разводим руками: так устроено природой. Поколения фанатиков ломали голову над теоремой Ферма. Неужели так будет и с эффектом

Подклетнова? Неведомой силой приковал он внимание людей в разных странах. Появились общества последователей Подклетнова. В Интернете имеются сотни страниц с его именем.

Энтузиасты собирают материалы об эффекте, выпускают просветительскую литературу о гравитации и ее колоссальных возможностях. Как правило, это те, кто любит в свободное от работы время паять, мастерить, возиться с микросхемами, ставить самодельные опыты. Например, Джон Шнурер из штата Огайо. Он заявил, что воспроизвел эффект Подклетнова и достиг нового рекорда - уменьшил силу гравитации на 5 процентов. Правда, очевидцы говорят, что в их присутствии опыт не получается. Весы качаются под действием пузырьков кипящего жидкого азота, как лапша в супе, и при желании можно увидеть какой угодно результат. Любители на то и любители, что необъективно относятся к объекту своего увлечения.

Но среди сторонников Подклетнова есть и профессионалы. Самый активный - итальянский физик Джованни Моданезе из Института Макса Планка в Мюнхене. Правда, в штате сотрудников института его фамилию найти не удалось, хотя там действительно есть группа физиков, работающих над теорией гравитации.

Обращаюсь в Лос-Аламосскую лабораторию, собравшую самую большую библиотеку по физическим наукам. Да, статьи Моданезе о квантовой теории эффекта Подклетнова опубликованы в солидных физических журналах. Последняя статья - 1999 года. В настоящее время он работает в университете города Тренто (Италия). Вопреки общепринятой науке Моданезе верит в реальность волшебного экрана. Он считает, что сверхпроводник способен поглощать гравитацию и тем самым освобождать от ее власти.

Еще раньше о волшебном экране начали подумывать физики-теоретики из университета Алабамы в Хантсвилле - Дуглас Торр и Нинь Ли. Их первая статья на эту тему датирована 1990 годом. По их настоянию в НАСА после сообщения об эффекте Подклетнова была немедленно организована научная группа для опытов с гравитацией. Потом Дуглас Торр переехал в другой штат и начал самостоятельные исследования. Он увлекся совсем странными вещами и договорился до того, что будто бы изобрел "генератор гравитации", способный послать силовой луч в любом заданном направлении. Но его заявления мало кто воспринимает всерьез. Что касается доктора физики Нинь Ли, то она продолжает работу с НАСА.

Хантсвилл, США

Эксперимент в НАСА по проверке эффекта Подклетнова готовили с тщательностью, на какую способны физики высшего класса. Достаточно сказать, что приборы способны измерить гравитацию с точностью до восьмого знака. В опытах задействованы 8 космических спутников, которые учитывают слабые изменения гравитационного поля Земли, вызванные приливами и отливами. Гравитацию измеряли в вакууме, в огромном металлическом ящике, экранированном от электромагнитного поля.

Оказалось, что вес над сверхпроводящим диском если и уменьшается, то мало. Не на два процента, как у Подклетнова, а в миллион раз меньше. Американцы были обескуражены. И пригласили Подклетнова в НАСА.

Он приехал в Центр космических полетов в Хантсвилле в конце прошлого года. Верный привычке, избегал корреспондентов и не давал интервью. Тем не менее сведения о его визите просочились в печать. Причиной неудачи, как он предположил, было то, что экспериментаторы не врашивали диск. А с неподвижным диском и у него не было эффекта. Кроме того, работали с маленьким диском. Поэтому было решено сделать диск в два раза больше.

Представитель компании по сверхпроводникам заявил, однако, что сложность изготовления дисков неизменно возрастает с увеличением их размеров. Самые большие диски, которые удается делать в Америке, имеют диаметр около 15 сантиметров. В феврале этого года НАСА предоставило компании 70 тысяч долларов с одним условием: если удастся сделать такой же диск, как в России, величина вознаграждения возрастет до 750 тысяч

долларов. Наверное, приобретение диска у нас или у Подклетнова обошлось бы дешевле. Но в США решили сделать его сами. На карту поставлены большие деньги и престиж.

Москва, Россия

В России, как и в Америке, гравитация издавна привлекает внимание ученых. Физик Г. Вилковыский считает, что "Ее Величество Квантовая Гравитация порабощает каждого, кто взглянул на нее хотя бы раз". В Москве проходят международные семинары по квантовой гравитации, съезжаются научные светила со всего мира. Однако в России мнение физиков об эффекте Подклетнова явно скептическое.

- Не надо об этом писать, - убеждали меня. - Это типичная лженеука. К счастью, в России Подклетнова не знают. И незачем говорить о нем народу. У нас и без него хватает колдунов и лжеученых.

Выходит, народу не нужно знать, о чем думают и спорят физики? Но за границей тоже хватает безумцев и лжеученых. Тем не менее там не боятся давать информацию. Ибо именно замалчивание кажущихся лжесенсаций работ помогает расцвету лженеуки. Под покровом секретности, к слову, легче получить деньги на безграмотные проекты.

Я спросил об эффекте Подклетнова академика Виталия Гинзбурга, одного из создателей теории сверхпроводимости, известной под названием теории Гинзбурга-Ландау. Он отнесся к эффекту с большим сомнением. "Экранов от гравитации создать нельзя, таков вывод из общей теории относительности, - сказал он. - Одна из важнейших проблем современной физики и космологии - создание квантовой теории гравитации".

Эффект Подклетнова согласился прокомментировать известный специалист по сверхпроводникам профессор Максим Каган из Института физических проблем им. П. Капицы РАН:

- Опыты Подклетнова выглядят довольно странно. На двух недавних международных конференциях по сверхпроводимости в Бостоне и Дрездене, где я принимал участие, его опыты не обсуждались. Специалистам он широко не известен. Уравнения Эйнштейна, в принципе, допускают взаимодействие электромагнитных и гравитационных полей. Но для того, чтобы такое взаимодействие стало заметным, нужна колossalная электромагнитная энергия, сравнимая с эйнштейновской энергией покоя. Нужны электрические токи на очень много порядков выше тех, что достижимы в современных лабораториях. Поэтому нет реальных возможностей изменять в эксперименте гравитационное взаимодействие.

- А как же НАСА? - спросил я.

- У НАСА есть большие деньги. Они проверяют многие идеи. Проверяют даже идеи сомнительные, но привлекательные для широкой аудитории. Большинство других организаций, включая даже такие научные фирмы, как "Белл" и "ИБМ", не может себе этого позволить. Мы изучаем реальные свойства сверхпроводников. Они очень интересны сами по себе. За последние 10 лет резко повышена температура сверхпроводящего перехода. Созданы сверхпроводники с критической температурой около минус 110 градусов Цельсия. В принципе возможно создание еще более высокотемпературных сверхпроводников. В результате может произойти настоящая революция в энергетике и в вычислительной технике. Появятся очень мощные компьютеры и сверхпроводящие магниты. В США уже сейчас испытывается первая короткая линия электропередач на базе высокотемпературных сверхпроводников (ВТСП). В Японии строят первый поезд на сверхпроводящей магнитной подушке. До широкого применения ВТСП-систем в промышленности требуется еще лет 10-15. Однако существенный прогресс в теоретическом понимании природы сверхпроводимости и в экспериментальном изготовлении образцов высокого качества наблюдается уже сейчас.

Эйнштейн, Сахаров. Кто следующий?

Недавно появилось первое упоминание о Подклетнове в российской прессе. Его опыты уже пытаются использовать для обоснования каких-то паранормальных явлений, психофизической энергии и прочей дребедени. Между тем эксперимент в НАСА еще не закончен, а то, что сделано, не подтверждает нашумевший эффект.

На пороге XXI века физики творят чудеса. Получены атомы антиводорода, сделан кристалл из электронов, скорость света замедлена до нескольких метров в секунду. Неудивительно, что случаются и фальстарты.

Альберт Эйнштейн последние 30 лет жизни работал над общей теорией поля. Эстафету подхватил другой гений - Андрей Сахаров. Кто знает, какими были бы наши знания, если бы внезапная смерть не оборвала его работу 10 лет назад? По свидетельству знаменитого физика Джона Уилера, именно работы Сахарова заложили основы современной теории гравитации.

Может быть, где-то под Архангельском или в Дели растет новый гений. Может быть, он заинтересуется гравитацией. Его жизнь будет нелегкой. Ему придется долго учиться, сдать много экзаменов и тяжело работать, чтобы принять огромное богатство знаний. Придется изучить и эффект Подклетнова. Возможно, ему удастся сделать то, над чем бился этот ученый. И человек наконец овладеет тайнами гравитации...

Дата 19981005 Дата загрузки 19990515 Время загрузки 2030 Источник Аномалия, Санкт-Петербург Заголовок **Обнаружены следы суперцивилизации!** Автор Михаил Герштейн Тема Вселенские новости Выпуск 19 Текст

Подводные пирамиды возрастом старше египетских найдены водолазами у японских берегов. Рядом с небольшим островком Йонагуни (неподалеку от Окинавы) на глубине 25 метров находится загадочное сооружение, больше всего напоминающее ступенчатую пирамиду.

Высота каждой ступени - около метра. Все сооружение возвышается над морским дном на 30 метров. Основание пирамиды не квадратное, а прямоугольное; длина наибольшей стороны - около 200 метров.

Ученые предполагают, что этой пирамиде более 10000 лет, то есть она вдвое старше знаменитых египетских собратьев!

Возможно, под водой она оказалась после окончания ледникового периода, когда поднялся уровень Мирового океана...

Эти данные мы почерпнули из бюллетеня "Рубежи науки" ("Science Frontiers", July/Aug. 1998). Ученые сообщают, что к пирамиде ведет что-то вроде дороги. Правда, это не убедило скептиков, которые полагают, что под водой находится всего лишь причудливым образом размытая скала.

Кто был прав, покажут дальнейшие исследования.

NASA строит антигравитрон по нашим "чертежам"

Американский журнал "Популярная механика" ("Popular Mechanics") сообщил о предстоящем испытании антигравитационной машины. Специалисты NASA надеются, что это произойдет в самое ближайшее время, поскольку машина собрана уже на 90 процентов.

Ее принцип основан на идеях нашего соотечественника Евгения Подклетнова, который демонстрировал опыты по подавлению гравитации в университете Тампере (Финляндия).

В машине NASA будут использованы диски из сверхпроводящих материалов, погруженные в смесь жидкого гелия и азота при температуре -400° по Фаренгейту, которые будут разогнаны до нескольких тысяч оборотов в минуту.

Любой предмет, помещенный над этой машиной, потеряет в весе; если его притяжение к Земле будет полностью компенсировано силой антигравитации, предмет зависнет в воздухе.

Если сила антигравитации будет превышать силу притяжения, предмет полетит вверх - пока обе силы не уравняются.

Подклетнову уже удавалось подвешивать в воздухе маленькие вещи; новый антигравитрон, скорее всего, будет способен на большее.

"Близкий контакт" произошел в Австралии

Колин Норрис, главный редактор австралийского журнала "Australian International UFO-Flying Saucer Research", что переводится как "Международный журнал по исследованиям НЛО/летающих тарелок в Австралии" (Aug. 1998), прислал в редакцию

последний номер с важным сообщением. 3 июня 1998 года в Австралии произошел случай, который уже стали называть "контактом десятилетия". Частный летчик Марк Мускат поднялся в воздух, чтобы показать группе туристов виды ночной Аделаиды с воздуха...

Он заметил в воздухе шесть тусклых оранжевых шаров. НЛО неслись на его самолет со скоростью 200 километров в час!

"Я прошел прямо посреди их строя, - вспоминал впоследствии Марк. - Шары были размером с большой автомобиль; они бесшумно пронеслись метрах в двадцати по обе стороны моего самолета. В этот миг я думал только о том, как бы они в меня не врезались".

Как пишет журнал все это видели не только потрясенные туристы, сидевшие за спиной Марка, но и десятки людей, находившихся на земле. Им казалось, что самолет вот-вот столкнется с одним из шаров, но НЛО в самый последний момент отвернулся в сторону.

Авиадиспетчеры так и не заметили ничего странного на своих экранах. Однако вскоре после "контакта" их радары засекли какую-то загадочную "цель", - возможно, корабль-матку, который ожидал возвращения "шаров".

Спасибо, коллега!

Космический "линкор" над Северным морем

В том же австралийском журнале мы откопали весеннюю "новость". 26 апреля с. г. радарная станция дальнего обнаружения Министерства обороны Великобритании засекла над Северным морем НЛО "размером с линкор". Гигантский треугольный объект летел по причудливой зигзагообразной траектории со скоростью 27000 километров в час!

За чудовищным "треугольником" следили и Военно-воздушные силы Дании. Датчане выслали на перехват два истребителя F-16, однако они не смогли догнать НЛО. Объект разогнался до скорости 38000 км/ч и улетел в сторону Атлантики.

"Судя по его маневрам, он определенно находился под чьим-то управлением," - заявил представитель Министерства обороны Великобритании.

По данным австралийской газеты "Advertiser", радарные записи "космического линкора" будут предоставлены ученым и военным на очередной конференции по освоению космоса в военных целях. Представители Министерства обороны признались, что за последние два года не раз засекали НЛО, но они предпочитают все данные по другим случаям держать в секрете, чтобы не раскрывать слишком много информации о возможностях радарной станции.

Найдено свидетельство крушения НЛО... в 1865 году

Канадский охотник наблюдал взрыв светящегося тела и нашел его обломки.

На эти сведения натолкнулся канадский исследователь Ритчи Бенедикт, просматривая старинные газеты. В одной из них он прочитал совершенно фантастическую статью.

"Мистер Джеймс Ламли, пожилой охотник из Скалистых Гор... рассказал нам весьма удивительную историю, каковая, если является правдивой, вызовет величайшее возбуждение в научном мире", - писала газета "British Daily Whig" от 15 ноября 1865 года.

Ламли видел падение "ярко светящегося тела", которое взорвалось в воздухе. Тряслась земля, пронеслась ударная волна. В воздухе пахло "странным сернистым запахом".

На следующее утро он наткнулся на гигантскую просеку, пропаханную через лес. "Огромные деревья были вырваны с корнем и изломаны у самой земли, вершины холмов начисто срезаны, земля взрыта во многих местах, - пересказывали его слова журналисты. - Повсюду виднелись следы страшного разрушения".

Идя по столь четкому следу, Джеймс наткнулся на гигантский "камень", врезавшийся в склон горы. Он был "разделен на отсеки", а в нескольких местах "покрыт иероглифами"!

"М-р Ламли также обнаружил осколки вещества, напоминающего стекло; там и сям виднелись темные пятна, словно оставленные жидкостью. Он был уверен, что иероглифы являются работой рук человеческих и что этот камень является частью гигантского тела, которое было использовано для каких-то целей разумными существами".

Падение "камня" видел не только Джеймс, но и другие очевидцы.

В конце статьи говорилось: "Быть может, в будущем некие Колумбы с Меркурия или Урана приземлятся на нашей планете в своих метеорных экипажах и овладеют ею... загнав человеческую расу в состояние самого жалкого рабства". Точно такие же мысли, судя по фильму "День независимости", не покидают человечество и в наши дни...

Заметку об этом мы прочитали в последнем номере журнала "Fate".

Дата 19980911 Дата загрузки 20001016 Время загрузки 1904 Источник Кузбасс, Кемерово Заголовок **ЛЯГУШКИ ЛЕТАЮТ, НО ПОКА НИЗЕНЬКО-НИЗЕНЬКО** Автор Игорь ЛЕЛЯНОВ Тема В МИРЕ ЗАГАДОЧНОГО Выпуск 169 Текст

Левитация, о которой так долго говорили только чудаки, теперь изучается в научных лабораториях

Наш рассказ о загадочной силе, которая якобы способна поднимать в воздух людей и предметы, мы начнем с истории, которая недавно случилась в польском городке Гошковице. Там, если верить газете "Жиче Варшавы", десятки людей стали свидетелями невероятного происшествия, когда местная гадалка Марыся Лозинская неведомо как взлетела в небо...

Дело происходило в воскресный день на людной ярмарочной площади. Марыся сидела за столиком и привычно раскладывала карты. Рядом находилось несколько "клиенток", желающих узнать у гадалки свою судьбу. Одна из них, Аннели Возгольская, так описывает дальнейшие события:

- Пани Марыся пожаловалась вдруг на сильнейшую головную боль и сказала, что вынуждена прервать гадание. Она и впрямь выглядела неважко - лицо побледнело и даже приобрело зеленоватый оттенок. Гадалка даже воскликнула: "Как больно!" - и в тот же момент начала медленно подниматься над землей. При этом ее тело оставалось совершенно неподвижным.

Дружное "ах!" пронеслось над ярмаркой. Все вокруг с изумлением смотрели на женщину, неведомым образом зависшую в воздухе на высоте около трех метров. Это продолжалось минуты две. Затем пани Марыся стала медленно снижаться. Едва ее ноги коснулись земли, как она тут же упала ничком.

А вот что рассказал журналистам врач А. Добжинский, который первым осматривал бесчувственную Марысю Лозинскую: "Глаза ее были закрыты. Пульс прощупывался, хотя и очень слабо. Постепенно на щеках появился румянец. Женщина спросила: "Почему я здесь лежу?". Оказалось, что она ничего не помнит..."

Следует сказать, что, с точки зрения исследователей так называемых аномальных явлений, собирающих в своих архивах подобные факты, случай с Марысей Лозинской далеко не самый выдающийся. Если верить историческим источникам, во время молитвенного экстаза в воздух поднимались многие религиозные деятели. Из наших отечественных "левитантов" можно назвать Серафима Саровского, архиепископа Новгорода и Пскова Иоанна... Московские летописи рассказывают о Василии Блаженном, который не раз на глазах у толпы переносился через Москву-реку. Католические святые (по тем же историческим источникам) ничем не уступали нашим. Один испанский паломник тоже умудрился перелететь через широкую реку, а епископ Валенсии однажды и вовсе парил над землей целых 12 часов. Всего же, как утверждают церковные книги, количество людей, демонстрировавших на глазах верующих феномен левитации, приближается к трем сотням. А число ведьм, которых во времена инквизиции за то же самое сожгли на костре, и вовсе не поддается учету.

Русский журнал "Ребус" XIX века, специализировавшийся на всяких таинственных историях, сообщал, например, такой случай, описанный неким Н. Юрловым. В возрасте 8 лет, 18 июля 1837 года, он спал в своей детской, на втором этаже в селе Спешневке (Сенгилеевский уезд, Симбирская губерния). Ночью была гроза. Разбуженный вспышками молний и громом, мальчик сел в постели.

"И вдруг, когда молния озарила комнату, к ужасу моему, увидал, что в двух шагах от моей кровати на балконе, держась за ручку стеклянной двери, стоит высокий лысый старик, с седою бородою, в длинной синей рубашке..."

Ужас мальчика был неописуем. Он скатился по лестнице вниз, выбежал из дверей в сад и, ничего не соображая, бросился в сторону реки Свияги. Прислуга спохватилась быстро, и мальчика нашли уже через 10 минут, но, если верить рассказчику, на другом берегу реки и - совершенно сухого!

По словам Юрлова, "...бежавшие за мною люди, Василий Кондаков и Федор Плотников, а в особенности садовник Николай Ермаков и кузнец Архипов, переплы whole="v"ые вплавь Свиягу, клятвенно удостоверили, что они ясно видели, как я быстро по воздуху переносился через реку, почти наравне с водою..."

Еще несколько подобных случаев приведено в большом сборнике "Из области таинственного" (1900 год). Так, к примеру, там рассказывается об известном итальянском церковном деятеле Бенедикте Нурсийском. Среди его учеников был некто Маурес. Увидев как-то тонущего мальчика, Маурес (если верить рассказам) стремглав бросился к несчастному на помощь, выхватил его из воды и выбежал обратно на берег.

"Только вернувшись на землю, он пришел в себя и посмотрел назад. Когда же он увидел, что побежал по воде, то страшно испугался происшедшего..."

Сегодня, как утверждают парапсихологи, феномен левитации успешно демонстрируют так называемые "летающие йоги" - люди, овладевшие учением о трансцендентальной медитации, которое было создано в начале 50-х годов Махариши Махеш Йогом (в мире - Махариши). Внешне полеты "летающих йогов" выглядят как подскоки в воздух в положении "поза лотоса". В 1986 году в городе Вашингтоне проходило публичное соревнование "летающих йогов", в котором приняло участие около 20 человек. В соревнование входили следующие дисциплины: высота полета (рекорд около 60 сантиметров), длина перелета (рекорд 1,8 метра) и полеты на скорость на дистанции 25 метров с препятствиями высотой около 20 сантиметров.

Словом, рассказов о левитации и левитантах хватает. Проблема в другом - можно ли верить таким историям?

До недавнего времени при упоминании слов "левитация" и "антигравитация" ученые дружно восклицали: "Чушь собачья!" Сегодня такого единодушия уже не наблюдается. Смятение в академические умы внес российский физик Евгений Подклетнов, работающий сейчас в технологическом университете финского города Тампере.

Специальный диск охлаждался исследователем до температуры минус 167 градусов по Цельсию и помещался в электромагнитное поле, заставляющее его вращаться. При достижении трех тысяч оборотов в минуту предметы, помещенные над вращающимся диском, начинали терять вес.

Доклад Подклетнова, опубликованный в научных журналах, сработал наподобие своеобразного детонатора. Одно за другим посыпались сообщения специалистов о наблюдавших ими антигравитационных эффектах.

Любопытные результаты получены Джоном Шнурером из Энтиоцского колледжа (штат Огайо). Суть его опытов заключается в следующем: если над магнитом поместить сверхпроводник, он зависает в воздухе. Впрочем, это уже известное явление, получившее название "эффект Мейснера". Непонятное, как и в опытах Подклетнова, начинается, когда над парящим без опоры сверхпроводником помещается какой-либо объект. Точные измерения показали, что над сверхпроводящей системой возникает зона, где предметы теряют до 5 процентов веса. По мнению Шнурера, это реальный шаг к созданию антигравитационных установок.

Еще более поразительные эксперименты проводились голландскими учеными. Они умудрились подвесить в воздухе живую лягушку. Земноводное помещалось над сверхпроводящей катушкой. Мощное поле воздействовало прямо на клетки тела, создавая в

них магнитный момент, противоположный магнитному полю Земли. Таким образом создавались силы, нейтрализующие тяготение.

Доктор Андре Гейн и его коллеги из университета в городе Наннели уверены, что вскоре с помощью своей аппаратуры они смогут поднимать в воздух любой объект, в том числе и человека. По крайней мере, сразу же после эксперимента с лягушкой эти ученые успешно подвесили в воздухе бутерброд, оставшийся у них от завтрака...

Голландцы нахально утверждают, что их лягушка - первое живое существо, поднявшееся в воздух без посредства преобразования механической энергии (то есть без прыжка, броска, крыльев, пропеллера, реактивной струи или взрывной волны), а иным способом, получившим название "молекулярный магнетизм". Но так ли это? В свете успешных экспериментов доктора Гейна уже совсем иначе выглядят церковные истории о левитирующих святых и ведьмах. Ведь лабораторные опыты только повторяют условия, возникающие порой в естественной природной среде.

И теперь на вопросы о левитации бывшие скептики вынуждены вместо ответа смущенно пожимать плечами. После "летающего голландского бутерброда" было бы глупостью голословно отрицать возможность такого явления. Значит, нужны специальные исследования, призванные подтвердить (или опровергнуть) легенды о том, что человек без всякой аппаратуры (вольно или невольно) способен так изменить работу клеток своего тела, что они начинают отталкиваться от магнитного поля планеты...

И такие исследования уже ведутся. Из самых серьезных организаций, заинтересовавшихся практическими возможностями подобных эффектов, можно назвать американский Центр космических полетов Джорджа Маршалла и НАСА. Руководитель отдела перспективных концепций Национального управления по аeronавтике США Уитт Брэнтли признает, что работы Подклетнова, Шнурера и других произвели в современной науке эффект разорвавшейся бомбы и в случае подтверждения откроют перед человечеством грандиозные перспективы.

Игорь ЛЕЛЯНОВ, сотрудник пресс-центра объединения "Феномен".

Дата 19980514 Дата загрузки 19980521 Время загрузки 1130 Источник Куранты, Москва Заголовок **Летающие лягушки, диски и бутерброды** Автор Игорь Кириллов Выпуск 10 Текст

Смятение в академические умы внес российский физик Евгений Подклетнов, работающий сейчас в Технологическом университете финского города Тампere.

Специальный диск охлаждался исследователем до температуры минус 167 градусов по Цельсию и помещался в электромагнитное поле, заставляющее его вращаться. При достижении трех тысяч оборотов в минуту предметы, помещенные над вращающимся диском, начинали терять вес.

Доклад Подклетнова, опубликованный в научных журналах, сработал наподобие своеобразного детонатора. Одно за другим посыпались сообщения специалистов о наблюдаемых ими антигравитационных эффектах.

Любопытные результаты получены Джоном Шнурером из Энтиохского колледжа (штат Огайо). Суть его опытов заключается в следующем: если над магнитом поместить сверхпроводник, он зависает в воздухе. Впрочем, это уже известное явление, получившее название "эффект Мейснера". Непонятное, как и в опытах Подклетнова, начинается, когда над парящим без опоры сверхпроводником помещается какой-либо объект. Точные измерения показали, что над сверхпроводящей системой возникает зона, где предметы теряют до 5 процентов веса. По мнению Шнурера, это реальный шаг к созданию антигравитационных установок.

Еще более поразительные эксперименты проводились голландскими учеными. Они умудрились подвесить в воздухе живую лягушку. Земноводное помещалось над сверхпроводящей катушкой. Мощное поле воздействовало прямо на клетки тела, создавая в

них магнитный момент, противоположный магнитному полю Земли. Таким образом создавались силы, нейтрализующие тяготение.

Доктор Андре Гейн и его коллеги из университета в городе Наннели уверены, что вскоре с помощью своей аппаратуры они смогут поднимать в воздух любой объект, в том числе и человека. По крайней мере сразу же после эксперимента с лягушкой эти ученые успешно подвесили в воздухе бутерброд, оставшийся у них от завтрака...

И теперь на вопросы о левитации бывшие скептики вынуждены вместо ответа смущенно пожимать плечами. После "летающего голландского бутерброда" было бы глупостью голословно отрицать возможность такого явления. Значит, нужны специальные исследования, призванные подтвердить (или опровергнуть) легенды о том, что человек без всякой аппаратуры (вольно или невольно) способен так изменить работу клеток своего тела, что они начинают отталкиваться от магнитного поля планеты... И такие исследования уже ведутся. Руководитель отдела перспективных концепций Национального управления по аэронавтике США Уитт Брэнтли признает, что работы Подклетнова, Шнурера и других произвели в современной науке эффект разорвавшейся бомбы и в случае подтверждения откроют перед человечеством грандиозные перспективы.

Дата 19970911 Время 0507 Дата загрузки 19971021 Время загрузки 1100 Источник Куйбышевский железнодорожник, Самара Заголовок **ЧЕЛОВЕК МЕДЛЕННО ПОДНИМАЛСЯ НАД ЗЕМЛЕЙ...** Автор Игорь Царев Выпуск 97 Текст

Наш рассказ о загадочной силе, которая способна поднимать в воздух людей и предметы, мы начнем с истории, которая недавно случилась в польском городке Гошковице. Там, если верить газете "Жиче Варшавы", десятки людей стали свидетелями невероятного происшествия, когда местная гадалка Марыся Лозинская неведомо как взлетела в небо...

Дело происходило в воскресный день на людной ярмарочной площади. Марыся сидела за столиком и привычно раскладывала карты. Рядом находилось несколько "клиенток", желающих узнать у гадалки свою судьбу. Одна из них (Аннели Возгольская) так описывает дальнейшие события:

- Пани Марыся пожаловалась вдруг на сильнейшую головную боль и сказала, что вынуждена прервать гадание. Она и впрямь выглядела неважко - лицо побледнело и даже приобрело зеленоватый оттенок. Гадалка даже воскликнула: "Как больно!" - и в тот же момент начала медленно подниматься над землей. При том ее тело оставалось совершенно неподвижным.

Дружное "ах!" пронеслось над ярмаркой. Все вокруг с изумлением смотрели на женщину, неведомым образом зависшую в воздухе на высоте около трех метров. Это продолжалось минуты две, затем пани Марыся стала медленно снижаться. Едва ее ноги коснулись земли, как она тут же упала ничком.

А вот что рассказал журналистам врач А. Добжинский, который первым осматривал бесчувственную Марысю Лозинскую: "Глаза ее были закрыты. Пульс прощупывался, хотя и очень слабо. Постепенно на щеках появился румянец. Женщина спросила: "Почему я здесь лежу?" Оказалось, что она ничего не помнит..."

Следует сказать, что с точки зрения исследователей так называемых аномальных явлений, собирающих в своих архивах подобные факты, случай с Марысей Лозинской далеко не самый выдающийся. Если верить историческим источникам, во время молитвенного экстаза в воздух поднимались многие религиозные деятели. Из наших отечественных "левитантов" можно назвать Серафима Саровского, архиепископа Новгорода и Пскова Иоанна... А число ведьм, которых во времена инквизиции за то же самое сожгли на костре, и вовсе не поддается учету. Проблема в другом - можно ли верить таким историям?

До недавнего времени при упоминании слов "левитация" и антигравитация" ученые дружно восклицали: "Чушь собачья!" Сегодня такого единодушия уже не наблюдается. Смятение в академические умы внес российский физик Евгений Подклетнов, работающий сейчас в Технологическом университете финского города Тампере.

Специальный диск охлаждался исследователем до температуры минус 167 градусов по Цельсию и помещался в электромагнитное поле, заставляющем его вращаться. При достижении трех тысяч оборотов в минуту предметы, помещенные над вращающимся диском, начинали терять вес.

Доклад Подклетнова, опубликованный в научных журналах, сработал наподобие своеобразного детонатора. Одно за другим посыпались сообщения специалистов о наблюдаемых ими антигравитационных эффектах.

Любопытные результаты получены Джоном Шнурером из Энтиохского колледжа (штат Огайо). Точные измерения показали, что над сверхпроводящей системой возникает зона, где предметы теряют до 5 процентов веса. По мнению Шнурера, это реальный шаг к созданию антигравитационных установок.

Еще более поразительные эксперименты проводились голландскими учеными. Они умудрились подвесить в воздухе живую лягушку. Земноводное помещалось над сверхпроводящей катушкой. Мощное поле воздействовало прямо на клетки тела, создавая в них магнитный момент, противоположный магнитному полю Земли. Таким образом создавались силы, нейтрализующие тяготение.

Доктор Андре Гейн и его коллеги из университета в городе Наннели уверены, что вскоре с помощью своей аппаратуры смогут поднимать в воздух любой объект, в том числе и человека. По крайней мере, сразу же после эксперимента с лягушкой эти ученые успешно подвесили в воздухе бутерброд, оставшийся у них от завтрака.

Голландцы нахально утверждают, что их лягушка - первое живое существо, поднявшееся в воздух без посредства преобразования механической энергии (то есть без прыжка, броска, крыльев, пропеллера, реактивной струи или взрывной волны), а иным способом, получившим название "молекулярный магнетизм". Но так ли это? В свете успешных экспериментов доктора Гейна уже совсем иначе выглядят церковные истории о левитирующих святых и ведьмах.

И теперь на вопросы о левитации бывшие скептики вынуждены вместо ответа смущенно пожимать плечами. После "летающего голландского бутерброда" было бы глупостью голословно отрицать возможность такого явления. Значит, нужны специальные исследования, призванные подтвердить (или опровергнуть) легенды о том, что человек без всякой аппаратуры (вольно или невольно) способен так изменить работу клеток своего тела, что они начинают отталкиваться от магнитного поля планеты...

И такие исследования уже ведутся. Из самых серьезных организаций, заинтересовавшихся практическими возможностями подобных эффектов, можно назвать американский Центр космических полетов Джорджа Маршалла (США).

Ведутся похожие исследования и в России. В информационный банк комиссии "Феномен", куда мы собираем самые невероятные идеи, открытия и технологии, поступила видеокассета, демонстрирующая волчок, свободно висящий примерно в десяти сантиметрах от подставки. При этом эффект достигается без использования сверхпроводящих систем и вообще без подвода какой-либо энергии. Как утверждают авторы "летающего волчка", при наличии необходимых материалов они могут изготовить такое устройство практически любых размеров. Эксперты "Феномена" сейчас внимательно изучают эту любопытную работу.

Дата 19970304 Время 0633 Дата загрузки 19970516 Время загрузки 2104 Источник Ставропольская правда Заголовок **ОНИ К НАМ НА ТАРЕЛКАХ , А МЫ - НА ДИСКАХ** Выпуск 41 Текст

Российский ученый Евгений Подклетнов, как сообщается в последнем номере журнала Business Week, охладил сверхпроводимый диск примерно до минус 167 градусов по Цельсию, а затем поместил его в электромагнитное поле, которое заставило этот диск, вращаться. Приблизительно при трех тысячах оборотов в минуту все, что помещалось сверху над диском, утратило два процента массы.

Успех эксперимента Е. Подклетнова в Технологическом университете Тампере (Финляндия) вызвал бурю в научном мире. Ведь это означает открытие антигравитации!"

Сторонники сулили ученому Нобелевскую премию, скептики - кинематографическую премию Оскар по категории спецэффекты.

Руководитель отдела прогрессивных концепций НАСА Уитт Брэнтли заявил, что намерен повторить эксперимент Е. Подклетнова. Если антигравитация действительно существует, считает Брэнтли, то НАСА должно выяснить это первым.

Сделать сверхпроводящий диск размером 27,5 см в диаметре мы еще никогда не пробовали, - признался Рональд Кокзор, главный инженер Центра космических полетов имени Джорджа Маршалла. На это, судя по всему, у американских ученых уйдет не меньше двух месяцев.

Business Week считает, что, если создание антигравитационных приборов станет возможно, автомобили и самолеты смогут экономить невероятное количество топлива. А вот если использовать сразу 100 сверхпроводящих дисков, подобных диску Подклетнова, то с их помощью можно запустить космический корабль.

Дата 19961126 Дата загрузки 19970520 Время загрузки 2017 Источник Новгородские ведомости Заголовок "ОСКАР" ИЛИ НОБЕЛЕВКА? Выпуск 77 Текст

("Труд").

Настоящий переполох среди американских ученых вызвало открытие их российского коллеги Евгения Подклетнова в области антигравитации.

Подклетнов, сообщает в последнем номере журнал "Бизнес уик", экспериментируя с материалами при экстремальных температурах в Технологическом университете Тампере (Финляндия), получил эффект частичной потери веса (в пределах 2 процентов). Российский ученый охладил специальный диск примерно до минус 167 градусов по Цельсию, а затем поместил его в электромагнитное поле, которое заставило его вращаться. Приблизительно при 3 тыс. оборотов в минуту все, что помещалось сверху над диском, как утверждает Подклетнов, теряло ориентировочно 2 процента веса.

Впрочем, отмечает еженедельник, появились утверждения, что речь может идти о мошенничестве. Однако, как бы там ни было, "мы намерены продолжать наши усилия по дублированию такого опыта", заявил Уитт Брэнтли, руководитель отдела прогрессивных концепций в Национальном управлении по аeronавтике и исследованию космического пространства (НАСА).

"Антигравитационный прибор, - подчеркивает "Бизнес уик", - может найти огромное практическое применение. Если его вмонтировать в автомашины, самолеты или железнодорожные вагоны, то можно получить значительную экономию топлива. Это также позволит при конструировании машин не особенно заботиться об их размерах и весе. А если использовать сразу 100 суперпроводящих дисков, каждый из которых обеспечивает 2-процентную потерю веса, то из совокупного воздействия можно запустить космический корабль".

Что касается российского ученого, то, отмечает журнал, "если результаты эксперимента Подклетнова подтвердятся, то он наверняка получит Нобелевскую премию или... высшую американскую награду в области кинематографа - "Оскара" по категории "спецэффекты".

Дата 19961102 Дата загрузки 19970520 Время загрузки 2018 Источник Коммерсантъ-daily, Москва Заголовок Загадки российской антигравитации Автор ИВАН ЛЕНСКИЙ Тема Наука и Образование, Приложение/Последняя Полоса Выпуск 188 Страница 17 Примечания Отдел Выпускающий Редактор Текст

Аппарат Подклетнова

Загадки российской антигравитации

Одной из любимых тем авторов научно-фантастических романов были и остаются антигравитационные машины, с помощью которых можно преодолевать силу тяжести и совершать всевозможные подвиги в состоянии левитации. С недавнего времени научный мир США и Западной Европы взбудоражен информацией о том, что российскому физику Евгению Подклетнову удалось создать такую машину.

По данным германской газеты *Frankfurter Allgemeine* и британского журнала *New Scientist*, Подклетнов отправил рукопись, в которой описал разработанный им аппарат, в редакцию именитого американского физического журнала *Journal of Physics*. Эксперты журнала представили рукопись Подклетнова на экспертизу трем специалистам. Ни один из них не смог обнаружить в ней логических или фактических ошибок.

Антигравитационная машина Подклетнова представляет собой сосуд цилиндрической формы диаметром 27,5 см, в котором "подвешено" вращающееся под воздействием электромагнитного поля со скоростью 3 тыс. оборотов в минуту кольцо из керамического сверхпроводника. По словам Подклетнова, при температуре -200 по Цельсию этот прибор уменьшает вес любого объекта, лежащего на нем, на 2%. Из какого материала выполнен находящийся над аппаратом объект, значения не имеет. Более того: при вращении одного кольца вокруг другого вес расположенного сверху объекта снижается уже на 4%.

После предварительной проверки американцы приняли решение опубликовать рукопись Подклетнова. Однако этого все же не произошло: вскоре после того как по всему научному миру разлетелась информация о грандиозном открытии российского физика, спикер университета в городе Тампере (Финляндия) обнародовал опровержение данных Подклетнова. При чем тут финны? Подклетнов сообщил американцам, что свое открытие совершил в сотрудничестве с финским физиком Петри Вуориненом. А спикер университета утверждает, что в Финляндии ни о каких антигравитационных исследованиях ничего не известно. Одновременно спикер подтвердил, что Подклетнов действительно работал в Тампере в начале девяностых годов. Что касается господина Вуоринена, то он категорически опровергает, что когда-либо занимался проблемами антигравитации.

В ответ на опровержение из Финляндии Подклетнов объявил, что свои эксперименты произвел в 1992 году. И если "теперешний" Вуоринен отрицает, что с ним работал, значит, это был какой-то его однофамилец. На это финны заявили, что другого Вуоринена у них нет и никогда не было. Нетрудно догадаться, что российско-финская коллизия привела американцев в немалое замешательство. К большому облегчению редакции *Journal of Physics*, Подклетнов вскоре отозвал свою рукопись.

Так совершено научное открытие или нет? Без экспериментального исследования аппарата Подклетнова американцы отвечать на этот вопрос отказываются. Однако прежние работы Подклетнова дают основания говорить о нем как о серьезном ученом. Сам он утверждает, что пока не намерен предоставлять никакой дополнительной информации о своем открытии "по соображениям финансового характера".

Как бы то ни было, активные работы по проверке экспериментов и выводов Подклетнова начались во многих научных центрах мира - в лаборатории американского космического агентства NASA в городе Хантсвилле (штат Алабама), Институте имени Макса Планка в Мюнхене (ФРГ) и др.

ИВАН Ъ-ЛЕНСКИЙ



Wells Fargo

Search the Web:

powered by MSN Search

Qwest

[Print](#) | [E-mail](#) | [Discuss](#) | [Bookmark](#) | [E-mail Services](#) | [Help](#)

Slate

brave new world

Feeling Antigravity's Pull

Can NASA stop the apple from falling on Newton's head?

By Adam Rogers

Posted Friday, October 18, 2002, at 8:30 AM PT

"Don't call it antigravity research," Ron Koczor pleads. He's a physicist at NASA's Marshall Space Flight Center in Huntsville, Ala., and he's talking about a project he's been working on for almost a decade. "Call it 'gravity modification.' 'Gravity anomalies.' Anything but antigravity. That's a red flag."

When people find out that the National Aeronautics and Space Administration has researchers working on sci-fi stuff like antigravity—or rather, "gravity modification"—the red flags do indeed start waving. Reputable scientists like Koczor earn polite disdain from colleagues (or worse, from funders of research). But truth's truth: NASA has been studying the manipulation of gravity for at least 10 years, as have nongovernment researchers.

NASA began its work after a Russian physicist named Evgeny Podkletnov published an article in the peer-reviewed journal *Physica C* in 1992. Podkletnov claimed that a device built around a [superconductor](#) and a magnet could shield an object from gravity. The trick, he said, was to make a superconducting disc about a foot in diameter, chill it, levitate it over magnets—a nifty property of superconductors is that they repel magnetic fields—and set it revolving like a compact disc. Podkletnov said an object placed above that contraption lost 0.3 percent of its weight. The object itself didn't change. Rather, gravity's effect on it lessened.

If that effect could be harnessed and strengthened, the aerospace industry would be upended. Vessels bound for space wouldn't have to ride atop massive, barely controlled explosions. All the energy human beings expend moving things around, from cargo to cars, could be reduced or eliminated. And post-Einsteinian physics would have to be rewritten to explain what the hell was going on. Podkletnov called the effect "gravitational force shielding," and even in the absence of a good theory to explain the phenomenon, other researchers took notice. "Because his experiment and results were published in a peer-reviewed, scientific journal, that gave it a level of credibility," Koczor says.

After Podkletnov published his article, it took NASA until 1999 to figure out how to make a large, thin superconducting disc. Ceramic high-temperature superconductors are brittle as cheap china, and the discs kept shattering. Once they solved that problem, NASA paid Columbus, Ohio-based SCI Engineered Materials \$650,000 to build the entire apparatus. But Podkletnov had called for a disc with two layers, one superconducting and one not, and SCI didn't solve that engineering challenge until last year. Then they hit another roadblock. The disc wouldn't spin. SCI engineers stuck a rotor through the

Also in Slate

Sports

The Anaheim Angels

The Anaheim Angels aren't the worst franchise in baseball history, but they are the most pathetic ... [More](#)

Arts & Life

Video Killed the TV Star

In *Auto Focus*, Paul Schrader seems almost relieved to have as his protagonist a befuddled pipsqueak ... [More](#)

News & Politics

A



How Smart Are Our Smart Bombs?

In the debate over going to war with Iraq, hawks and doves tend to share one tenet—that the ... [More](#)

Confidential Pharmacy

- ✓ ONLINE PHARMACY
- ✓ FREE ONLINE CONSULTATION
- ✓ NEXT DAY SHIPPING
- ✓ LOWEST PRICES ON THE INTERNET
- ✓ SECURE ONLINE ORDERING

CLICK HERE

disc's center to turn it mechanically, but Podkletnov specified 5,000 revolutions per minute. SCI's device barely pulls 30 rpm.

Why not just ask Podkletnov how to build the thing? SCI brought him over to consult a couple of years ago, to little avail. "His excuse basically was that he was a ceramics physicist, not an electrical or mechanical engineer, and other people built the device for him," Koczor says. "Draw your own conclusions. All I know is, if I were a principal investigator on something like this, I would know the size and thread-depth of every screw in the damn thing. But you know, the Europeans and the Russians, they're different. They're much more, 'this is your job and this is my job.' So it's plausible that he didn't know the details." It might not matter. SCI's contract is ending, and Koczor's budget to explore "way-out physics" is spent. He hasn't got the money to actually test the device even if it did meet Podkletnov's specs.

But researchers outside NASA are working on the problem, too. This summer Nick Cook, a writer for *Jane's Defence Weekly*, reported that aerospace giant Boeing was pursuing antigravity research. Boeing denied it. "We are aware of Podkletnov's work on 'anti-gravity' devices and would be interested in seeing further development work being done," said a company statement. "However, Boeing is not funding any activities in this area at this time." Note Boeing's use of the Clintonian present tense. They never contacted *Jane's* to ask for a correction, Cook says. Meanwhile, British aerospace company BAE Systems says it's keeping an eye on the research, and that it had once funded its own antigravity project, Greenglow.

Unfortunately, Cook strains his own credibility somewhat. A couple of weeks after his *Jane's* piece appeared, Cook's book on antigravity research, *The Hunt for Zero Point*, came out. In it, he claims that the Nazis built an antigravity device during World War II. Its absence from present-day science, Cook says, implies a vast "black" world of secret antigravity aircraft that might explain the UFOs people see over Area 51. He's a careful investigative reporter, but once you start talking about UFOs and Nazi antigravity you're not far from hidden tunnels under the White House full of lizard-men disguised as Freemasons.

Even without Nazis, there are plenty of reasons to doubt Podkletnov. My e-mails to the account listed on his recent articles (not peer-reviewed) went unanswered. Even more problematic, I can't find the institution he lists as his affiliation in Moscow. "Eugene always expressed his worries that others could copy his work, although as far as I know he never applied for a patent," Giovanni Modanese, a collaborator of Podkletnov's at the University of Bolzano in Italy, wrote in an e-mail (using a Western version of Podkletnov's first name). "Nonetheless, at the scientific level if one wants a confirmation by others and a successful replication, one must give all the necessary elements." Well, yeah. Modanese says that the current version of the device, now called an "impulse gravity generator," is simpler and could be built "by a big-science team of people expert in superconductivity." A Boeing spokesperson didn't respond to follow-up questions. So, either there's nothing going on here, or it's an X-File.

And the science? Ten years is a long time to go without replication. Combine that with Podkletnov's cagey behavior and it's enough to make even sci-fi geeks like me lose hope. But like the core of any good conspiracy, antigravity research has the ring of plausibility. One of the outstanding problems in physics and cosmology today involves the existence of so-called dark matter and dark energy. They're by far the main constituents of matter in the universe, and nobody knows what they're made of—researchers have only inferred their existence from gravitational effects. Coming up with a new theory of how gravity works might explain that, though it'd be a scientific revolution on a par with relativity. "Changing gravity is in the cards," says Paul Scheckter, an astronomer at MIT. "But so far no one's been able to do

better than Einstein." Still, Einstein worked in a lowly patent office. Ron Koczor works for NASA.

Advertisement



[Leatherman Pulse with Leather](#) -

Leatherman Pulse with Leather Belt Sheath

[Amazon.com](#)

New Super Saver Shipping!

Adam Rogers, a reporter for *Newsweek*, is a Knight Science Journalism fellow at the Massachusetts Institute of Technology.

Slate

More brave new world

[NASA's mysterious antigravity research.](#)

posted October 18, 2002

Adam Rogers

[eBabe](#)

posted October 28, 1999

William Saletan

[Frankenstein's Minister](#)

posted December 10, 1998

William Saletan

Search for more [Brave New World](#) in our archive.



What did you think of this article?

[Join the Fray, our reader discussion forum](#)

[POST A MESSAGE](#) [READ MESSAGES](#)

[MSN - More Useful Everyday](#)

[MSN Home](#) | [My MSN](#) | [Hotmail](#) | [Search](#) | [Shopping](#) | [Money](#) | [People & Chat](#)

Â©2002 Microsoft Corporation. All rights reserved. [Terms of Use](#) [Advertise](#) [TRUSTe Approved](#) [Privacy Statement](#) [GetNetWise](#)



The Following is an essay by Paul A. LaViolette, Ph.D.

It is also an excerpt from the book "Electrogravitics Systems: Reports on a New Propulsion Methodology" by Thomas Valone, M.A., P.E.

The U.S. Antigravity Squadron

by Paul A. LaViolette, Ph.D.

Abstract

Electrogravitic (antigravity) technology, under development in U.S.

Air Force black R&D programs since late 1954, may now have been put to practical use in the B-2 Advanced Technology Bomber to provide an exotic auxiliary mode of propulsion.

This inference is based on the recent disclosure that the B-2 charges both its wing leading edge and jet exhaust stream to a high voltage.

Positive ions emitted from its wing leading edge would produce a positively charged parabolic ion sheath ahead of the craft while negative ions injected into its exhaust stream would set up a trailing negative space charge with a potential difference in excess of 15 million volts.

According to electrogravitic research carried out by physicist T. Townsend Brown, such a differential space charge would set up an artificial gravity field that would induce a reactionless force on the aircraft in the direction of the positive pole.

An electrogravitic drive of this sort could allow the B-2 to function with over-unity propulsion efficiency when cruising at supersonic velocities.

For many years rumors circulated that the U.S. was secretly developing a highly advanced, radar-evading aircraft.

Rumor turned to reality in November of 1988, when the Air Force unveiled the B-2 Advanced Technology Bomber.

Although military spokesmen provided the news media with some information about the craft's outward design, and low radar and infrared profile, there was much they were silent about.

However, several years later, some key secrets about the B-2 were leaked to the press.


On March 9, 1992, "Aviation Week and Space Technology" magazine made a surprising disclosure that the B-2 electrostatically charges its exhaust stream and the leading edges of its wing-like body.(1)

Those familiar with the electrogravitics research of American physicist T. Townsend Brown will quickly realize that this is tantamount to stating that the B-2 is able to function as an antigravity aircraft. "Aviation Week" obtained their information about the B-2 from a small group of renegade west coast scientists and engineers who were formerly associated with black research projects.

In making these disclosures, these scientists broke a code of silence that rivals the Mafia's.

They took the risk because they felt that it was important for economic reasons that efforts be made to declassify certain black technologies for commercial use.

Two of these individuals said that their civil rights had been blatantly abused (in the name of security) either to keep them quiet or to prevent them from leaving the tightly controlled black R&D community. Several months after "Aviation Week" published the article, black world security personnel went into high gear.



That sector of the black R&D community received VERY STRONG warnings and, as a result, the group of scientists subsequently broke off contact with the magazine.

Clearly, the overseers of black R&D programs were substantially concerned about the information leaks that had come out in that article. To completely understand the significance of what was said about the B-2, one must first become familiar with Brown's work.

Beginning in the mid 1920's, Townsend Brown discovered that it is possible to create an artificial gravity field by charging an electrical capacitor to a high-voltage.(2)

He specially built a capacitor which utilized a heavy, high charge-accumulating (high K-factor) dielectric material between its plates and found that when charged with between 70,000 to 300,000 volts, it would move in the direction of its positive pole.

When oriented with its positive side up, it would proceed to lose about 1 percent of its weight. (3, 4)

He attributed this motion to an electrostatically-induced gravity field acting between the capacitor's oppositely charged plates.

By 1958, he had succeeded in developing a 15 inch diameter model saucer that could lift over 110% of its weight!(5)

Brown's experiments had launched a new field of investigation which came to be known as electrogravitics, the technology of controlling gravity through the use of high-voltage electric charge. As early as 1952, an Air Force major general witnessed a demonstration in which Brown flew a pair of 18 inch disc airfoils suspended from opposite ends of a rotatable arm.

When electrified with 50,000 volts, they circuited at a speed of 12 miles per hour.(6) About a year later, he flew a set of 3 foot diameter saucers for some Air Force officials and representatives from a number of major aircraft companies.

When energized with 150,000 volts, the discs sped around the 50 foot diameter course so fast that the subject was immediately classified.

"Interavia" magazine later reported that the discs would attain speeds of several hundred miles per hour when charged with several hundred thousand volts.(7) Brown's discs were charged with a high positive voltage, on a wire, running along their leading edge and a high negative voltage, on a wire, running along their trailing edge.

As the wires ionized the air around them, a dense cloud of positive ions would form ahead of the craft and corresponding cloud of negative ions would form behind the craft.


Brown's research indicated that, like the charged plates of his capacitors, these ion clouds induced a gravitational force directed in the minus to plus direction.

As the disc moved forward in the response to its self generated gravity field, it would carry with it its positive and negative ion clouds and their associated electrogravity gradient.

Consequently, the discs would ride their advancing gravity wave much like surfers ride an ocean wave. Dr. Mason Rose, one of Townsend's colleagues, described the discs, principle of operation as follows:(8) The saucers made by Brown have no propellers, no jets, no moving parts at all.

They create a modification of the gravitational field around themselves, which is analogous to putting them on the incline of a hill.

They act like a surfboard on a wave...The electrogravitational saucer creates its own "hill", which is a local distortion of the



gravitational field, then it takes this "hill" with it in any chosen direction and at any rate. The occupants of one of [Brown's] saucers would feel no stress at all no matter how sharp the turn or how great the acceleration.

This is because the ship and its occupants and the load are all responding equally to the wave like distortion of the local gravitation field. Although skeptics at first thought that the discs were propelled by more mundane effects such as the pressure of negative ions striking the positive electrode, Brown later carried out vacuum chamber tests which proved that a force was present even in the absence of such ion thrust.

He did not offer a theory to explain this nonconventional electrogravitic phenomenon; except to say that it was predicted neither by general relativity nor by modern theories of electromagnetism.

However, recent advances in theoretical physics provide a rather straightforward explanation of the principle. According to the novel physics of subquantum kinetics, gravity potential can adopt two polarities, instead of one. (9-13)

Not only can a gravity field exist in the form of a matter attracting gravity potential well, as standard physics teaches, but it can also exist in the form of a matter repelling gravity potential hill.

Moreover, it predicts that these gravity polarities should be directly matched with electrical polarity; positively charged particles such as protons generating gravity wells and negatively charged particles such as electrons generating gravity hills.

(Thus contrary to conventional theory, the electron produces a MATTER-REPELLING gravity field.

Electrical neutral matter remains gravitationally attractive because of the proton's G-well marginally dominates the


electron's G-hill.) Consequently, subquantum kinetics predicts that the negative ion cloud behind Brown's disc should form a matter repelling gravity hill while the positive ion cloud ahead of the disc should form a matter attracting gravity well.

As increasing voltage is applied to the disc, the gravity potential hill and well become increasing prominent and the gravity potential gradient between them increasing steep.

In Rose's terminology, the craft would find itself on the incline of a gravitational "hill".

Since gravity force is known to increase in accordance with the steepness of such a gravity potential slope, increased voltage would induce an increasingly strong gravity force on the disc and would act in the direction of the positive ion cloud.

The disc would behave as if it was being tugged by a very strong gravitational field emanating from an invisible planet sized mass positioned beyond its positive pole.



Emerging Possibilities for Space Propulsion Breakthroughs

Originally published in the Interstellar Propulsion Society Newsletter, Vol. I, No. 1, July 1, 1995.

Marc G. Millis
Space Propulsion Technology Division
NASA
Lewis Research Center
Cleveland, Ohio

The ideal interstellar propulsion system would be one that could get you to other stars as quickly and comfortably as envisioned in science fiction. Before this can become a reality, two scientific breakthroughs are needed: discovery of a means to exceed light speed, and discovery of a means to manipulate the coupling between mass and spacetime. This article explains why these breakthroughs are needed and introduces the emerging possibilities that may eventually lead to these breakthroughs. It should be noted that either of these breakthroughs by itself would have revolutionary consequences which would be of enormous value.


The need to exceed light speed: Simply put, the universe is big. The fastest thing known is light, yet it takes over four years for light to reach our nearest neighboring star. When NASA's Voyager spacecraft left our solar system it was traveling around 37-thousand mph. At that rate it couldn't reach the nearest star until after 80-thousand years. If we want to cruise to other stars within comfortable time spans (say, less than a term in Congress), we have to figure out a way to go faster than light.

The need to manipulate mass and spacetime coupling: This need is less obvious than the light speed issue. The problem is fuel, or more specifically, rocket propellant. Unlike a car that has the road to push against, or an airplane that has the air to push against, rockets don't have roads or air in space. Rockets have to carry along all the mass that they'll need to push against. To circumvent this problem, we need to find a way to interact with spacetime

itself to induce propulsive forces without using propellant. This implies that we'll need to find a way to alter a vehicle's inertia, its gravitational field, or its connectivity to the structure of spacetime itself.

Just how limited are rockets for interstellar travel? Although rockets are reasonable for journeys into orbit or to the moon, they become unreasonable for interstellar travel. If you want to deliver a modest size payload, say a full Shuttle cargo (20,000 kg), and you are patient enough to wait 900 years for it to just fly by the nearest star, here's how much propellant you'll need: If you use a rocket like on the Shuttle ($I_{sp} \sim 500s$), there isn't enough mass in the universe to get you there. If you use a nuclear fission rocket ($I_{sp} \sim 5,000s$) you need about a billion super-tankers of propellant. If you use a nuclear fusion rocket ($I_{sp} \sim 10,000s$) you only need about a thousand super-tankers. And if you assume that you'll have a super-duper Ion or Antimatter rocket ($I_{sp} \sim 50,000s$), well now you only need about ten railway tankers. It gets even worse if you want to get there sooner. (Based on mass fractions from ref 1, p. 52) There are other ideas, like using laser pushed light-sails that don't need propellant, but these have limitations. The biggest limitation is their dependence on the laser that remains near Earth. To make an unplanned course change they need to radio back for the laser to track their new course and wait for it to do so. At interstellar distances this is prohibitive. At one light-year from Earth, for example, it would take two years for the command to be sent and the new pointing received.

Is there hope?: Science continues to advance. In addition to the continuing refinements of general relativity and other attempts to better understand mass, space, and time, there have been some recently published theories which provide new perspectives, theories which have been reported in various news articles (Refs 2-6). Each of these theories has some relevance to propulsion and presents new avenues from which to start searching for the breakthrough physics. These recent theories are summarized next.



A theory about "warp drive": Using the formalism of general relativity, it has been shown that faster than light travel may be possible (ref 7). All you need to do is contract spacetime in front of your ship and expand spacetime behind your ship. This "warped" space and the region within it would propel itself "with an arbitrarily large speed" (ref 7). Observers outside this "warp" would see it move faster than the speed of light. Observers inside this "warp" would feel no acceleration as they zip along at warp speed.

So what's the catch? First, to expand spacetime behind the ship you'll need matter having a negative energy density like negative mass, and lots of it too. It is unknown in physics whether negative mass or negative energy densities can exist. Classical physics tends toward a "no," while quantum physics leans to a "maybe, yes." Second, you'll need equal amounts of positive energy density matter, positive mass, to contract spacetime in front of the ship. Third, you'll need a way to control this effect to turn it on and off at will. And lastly, there is the debate about whether this whole "warp" would indeed move faster than the speed of light. To address this speeding issue, the theory draws on the "inflationary universe" perspective. The idea goes something like this: Even though light-speed is a limit within spacetime, the rate at which spacetime itself can expand or contract is an open issue. Back during the early moments of the Big Bang, spacetime expands faster than the speed of light. So if spacetime can expand faster than the speed of light during the Big Bang, why not for our warp drive? Just prior to the publication of the above theory, there was a workshop held at JPL to examine the possibilities for faster-than-light travel (ref 8). Wormholes, tachyons, and alternate dimensions were just some of the topics examined. The conclusions from this informal two-day workshop are as follows: (1) Faster-than-light travel is beyond our current horizons. Not only is the physics inadequately developed, but this physics is not oriented toward space propulsion or toward laboratory scale experiments.

(2) Causality violations (where effect precedes cause) are unavoidable if faster-than-light travel is possible, but it is uncertain whether causality violations are themselves physically prohibited.


(3) A few experimental approaches are feasible to address the science associated with faster-than-light travel, including: (a) Search for evidence of wormholes using astronomical observations: look for a group of co-moving stars or for the visual distortions indicative of a negative mass hole entrance.

(b) Measure the velocity of light inside a Casimir cavity (between closely spaced conductive plates) to search for evidence of negative space energy. This pertains to wormholes, tachyons, and the negative energy density issue.

(c) Resolve the rest mass issue of the Neutrino, determining whether the unconfirmed experimental evidence of imaginary mass is genuine.

(d) Study cosmic rays above the atmosphere, using scattering targets of known composition to look for characteristic evidence of tachyons and more general particle physics events.

New ways to think of inertia and gravity: As mentioned earlier, the ideal interstellar drive would have the ability to manipulate the connection between mass and spacetime. One approach is to look for ways to use electromagnetism, a phenomenon for which we are technologically proficient, to control inertial or gravitational forces. It is known that gravity and electromagnetism are coupled phenomena. In the formalism of general relativity this coupling is described in terms of how mass warps the spacetime against which electromagnetism is measured. In simple terms this has the consequence that gravity appears to bend light, red-shift light and slow time. These observations and the general relativistic formalism that describes them have been confirmed (ref 9, 10). Although gravity's



affects on electromagnetism have been confirmed, the possibility of the reverse, of using electromagnetism to affect gravity, is unknown.

New perspectives on the connection between gravity and electromagnetism have just emerged. A theory published in February 1994 (ref 11) suggests that inertia is nothing but an electromagnetic illusion. This theory builds on an earlier work (ref 12) that asserts that gravity is nothing other than an electromagnetic side-effect. Both of these works rely on the perspective that all matter is fundamentally made up of electrically charged particles, and they rely on the existence of Zero Point Energy.


Zero Point Energy (ZPE) is the term used to describe the random electromagnetic oscillations that are left in a vacuum after all other energy has been removed (ref 13). This can be explained in terms of quantum theory, where there exists energy even in the absolute lowest state of a harmonic oscillator. The lowest state of an electromagnetic oscillation is equal to one-half the Planck constant times the frequency. If all the energy for all the possible frequencies is summed up, the result is an enormous energy density, ranging from 10³⁶ to 10⁷⁰ Joules/m³. In simplistic terms there is enough energy in a cubic centimeter of the empty vacuum to boil away Earth's oceans. First predicted in 1948, ZPE has been linked to a number of experimental observations. Examples include the Casimir effect (ref 14), Van der Waal forces (ref 15), the Lamb-Rutherford Shift (ref 10, p. 427), explanations of the Planck blackbody radiation spectrum (ref 16), the stability of the ground state of the hydrogen atom from radiative collapse (ref 17), and the effect of cavities to inhibit or enhance the spontaneous emission from excited atoms (ref 18).

Regarding the inertia and gravity theories mentioned earlier, they take the perspective that all matter is fundamentally constructed of electrically charged particles and that these particles are constantly interacting with this ZPE background. From this perspective the

property of inertia, the resistance to change of a particle's velocity, is described as a high-frequency electromagnetic drag against the Zero Point Fluctuations. Gravity, the attraction between masses, is described as Van der Waals forces between oscillating dipoles, where these dipoles are the charged particles that have been set into oscillation by the ZPE background.

It should be noted that these theories were not written in the context of propulsion and do not yet provide direct clues for how to electromagnetically manipulate inertia or gravity. Also, these theories are still too new to have either been confirmed or discounted. Despite these uncertainties, typical of any fledgling theory, these theories do provide new approaches to search for breakthrough propulsion physics. Their utility and correctness remains to be determined.

Another viewpoint on gravity and spacetime: As mentioned earlier, the ideal interstellar drive must not use propellant. Instead the ideal drive would have to use some means to push against spacetime itself. One of the major objections to this notion is the issue of conservation of momentum (ref 19). In order to satisfy conservation of momentum, something must act as a reaction mass. For rockets it is the expelled propellant; for aircraft it is the air. If one considers propelling against spacetime itself, then one must entertain the possibility that the fields of spacetime have an energy or momentum that can serve as a reaction mass. Although existing physics does not provide this perspective, a recent theory has emerged that might. A news article published in December 94 (ref 6) introduced a theory (ref 20) that is challenging Einstein's general theory of relativity. The theory is generating a bit of controversy because it claims that the Einstein field equations need a slight correction. Without this correction it is claimed that the Einstein equations can only predict the behavior of simple one-body problems (where only one gravitating mass exists whose affect on an inconsequential test particle is described). For two-body or n-body problems, this new theory shows that the



Einstein equations are inadequate. The required correction is that another term must be added to the matter tensor, specifically a term for the stress-energy tensor of the gravitational field itself. This suggests that gravitational fields have an energy and momentum of their own. This may be a foundation to address the issue of a reaction mass for the ideal space drive.

Like the previously mentioned theories, it is uncertain whether this theory is correct or not, but it is certain that this theory adds yet another research path to search for breakthrough propulsion.


But wait, there's more: Another avenue to explore pushing against space is to examine the contents of the vacuum that may be indicative of a reaction mass. In addition to the items mentioned above, consider the following phenomena: Cosmic Background Radiation (ref 21), Virtual Pair Production (ref 22), and Dark Matter (ref 23). Whether any of these may constitute a reaction mass or may be evidence for a reaction mass is uncertain.

In addition to these recent events, there have been occasional surveys by the Air Force and others to examine science that may be applicable to propulsion technology (refs 24-29). The options identified by these studies include assessments of the technological status of many popular ideas, such as light-sails, nuclear rockets, and antimatter rockets, plus they include mention of more speculative work. Many of the more speculative ideas, from alternative theories of gravity and electromagnetism through unconfirmed anomalous effects, would be relatively simple to test. Very few of these possibilities have been rigorously investigated.


As you can see, there are a number of dangling loose ends in physics that may prove to be fruitful paths to the goal of creating the breakthroughs for practical interstellar travel. Pick your favorite idea and let us know what you discover.

REFERENCES Note: An annotated bibliogoraphy is available.

1. Mallove, E.F., and Matloff, G.L., *The Starflight Handbook*, Wiley Science Editions, John Wiley and Sons, Inc., New York (1989).
2. Gonick, L., "Science Classics (Warp-and-woof drive)", In *Discover*, p. 44-54, (DEC 1994).
3. Szipir, M, "Spacetime Hypersurfing?", In *American Scientist*, Vol. 82, p. 422-423, (September-October 1994)
4. Clarke, A. C., "Space Drive: A Fantasy That Could Become Reality", In *Ad Astra*, p. 38, (Nov-Dec 1994)
5. Matthews, R., "Inertia: Does Empty Space Put Up the Resistance?", In *Science*, Vol. 263, p. 612-613, (4 Feb 1994).
6. Peterson, I., "A New Gravity?, Challenging Einstein's General Theory of Relativity", In *Science News*, Vol. 146, p. 376-378, (Dec 3, 1994).
7. Alcubierre, M., "The warp drive: hyper-fast travel within general relativity", In *Classical and Quantum Gravity*, Vol 11, p. L73- L77, (1994).
8. Bennett, G., "Warp Speed, Fact or Fiction?", In *Final Frontier*, p. 35-39, (September-October 1994).
9. Pool,R., "Closing in On Einstein's Special Theory of Relativity", In *Science*, Vol. 250, p. 1207-1208, (Nov. 9, 1990).
10. Misner,C.W., Thorne,K.S., and Wheeler,J.A., *Gravitation*, W.H.Freeman and Co., New York (1973).
11. Haisch, B., Rueda, A., and Puthoff, H.E., "Inertia as a Zero-Point Field Lorentz Force", In *Physical Review A*, Vol. 49, No. 2, p. 678- 694, (FEB 1994)
12. Puthoff,H.E., "Gravity as a zero-point-fluctuation force", In *Physical Review*



- A, Vol. 39, N. 5, (A89-33278), p. 2333-2342, (Mar 1, 1989).
13. Boyer,T.H., "The Classical Vacuum", In Scientific American, p. 70-78, (Aug 1985).
14. Spaarnay, M. J., Measurements of Attractive Forces between Flat Plates, In Physica, Vol. 24, p. 751-764, (1958).
15. Casimir,H.B.G., and Polder, D., "The Influence of Retardation on the London-van der Waals Forces", In Physical Review, Vol.73, N.4, p.360-372, (February 15, 1948).
16. Boyer,T.M., "Random Electrodynamics: The Theory of Classical Electromagnetic Zero-Point Radiation" In Physical Review D, Vol.11, N.4, p.790-808, (Feb 15, 1975).
17. Puthoff,H.E., "Ground State of Hydrogen as a Zero-Point-Fluctuation-Determined State", In Physical Review D, Vol.35, No.10, p.3266-3269, (15 May 1987).
18. Haroche,S. and Kleppner,D., "Cavity Quantum Electrodynamics" In Physics Today, p.24-30, (January, 1989).
19. Millis, M. G., "Exploring the Notion of Space Coupling Propulsion", In Vision 21: Space Travel for the Next Millennium, Symposium Proceedings, Apr 1990, NASA-CP-10059, p. 307-316, (1990).
20. Yilmaz, H., "Toward a Field Theory of Gravitation", In Il Nuovo Cimento, Vol. 107B, no. 8, p. 941-960, (Aug 1992).
21. Muller, R. A., "The Cosmic Background Radiation and the new Aether Drift", In Scientific American, Vol. 238, N. 5, p. 64-74, (May 1978).
22. Kaufmann, W. J. III, Black Holes and Warped Spacetime, pp 206-208, W. H. Freeman and Co., San Francisco, (1979).
23. Krauss, L. M., "Dark Matter in the Universe", In Scientific American, p. 58-68, (Dec 1986).
24. Mead, F. B. ,Jr., et al, Advanced Propulsion Concepts - Project Outgrowth, AFRPL-TR-72-31, (JUN 1972).
25. Forward, R. L., "Feasibility of Interstellar Travel: A Review", In Acta Astronautica, Vol. 14, p 243-252, (1986) 26. Mead, F. B. ,Jr., "Exotic Concepts for Future Propulsion and Space Travel", In Advanced Propulsion Concepts, 1989 JPM Specialist Session, (JANNAF) Chemical Propulsion Information Agency, CPIA Publication 528, p.93-99, (May 24, 1989).
27. Cravens,D.L., Electric Propulsion Study, AL-TR-89-040, Final Report on Contract FO4611-88-C-0014, Air Force Astronautics Lab (AFSC), (Aug 1990).
28. Evans, R.A., British Aerospace Ltd. Co (BAe) University Round Table on Gravitational Research, Report on Meeting held at the NOVOTEL Conf.
- Ctr., Preston , on March 26-27, 1990, FBS 007, (Nov 1990).
29. Forward, R. L., 21st Century Space Propulsion Study, AL-TR-90-030, Final Report on Contract FO4611-87-C-0029, Air Force Astronautics Lab (AFSC), (Oct 1990). --AND-- Forward,R.L., 21st Century Space Propulsion Study (Addendum), PL-TR-91-3022, Final (Addendum), OLAC Phillips Lab, formally known as Air Force Astronautics Lab (AFSC), (June 1991).



Annotated Bibliography

(source: <http://www.grc.nasa.gov/WWW/PAO/html/warp/bibliog.htm>)

Note: Annotations are provided as an aid to the reader and are the personal opinions of the author. No endorsements of the works or ideas presented are implied.

Categories:

Overviews, Fact & Fiction Introductions to Emerging Physics Detailed Technical Papers
Overviews, Fact & Fiction:

1. Mallove, E.F., and Matloff, G.L., The Starflight Handbook, Wiley Science Editions, John Wiley & Sons, Inc., New York (1989).

Book; (274 pages) An overview of interstellar travel, the problems and proposed solutions. It contains technical details sufficient to lead researchers to more substantive material, yet is easy enough to read for general interest. The book constrains itself mostly to existing science and projected technology.

2. Mauldin, J.H., Prospects for Interstellar Travel, American Astronautical Society by Univelt Inc. San Diego CA, (1992).

Book; (370 pages) An overview of interstellar travel, the problems, proposed solutions and social issues. It contains sufficient details to lead researchers to more substantive material. It also touches on longer-term and more speculative ideas, including space warps, Zero Point Energy and Higgs fields.

3. Emme, E.D., (ed), Science Fiction and Space Futures Past & Present, American Astronautical Society Historical Series, Vol 5, (1982).

Book; Explores the connection between Science Fiction and advances in astronautics. Conclusions: Science fiction is not technically accurate at predicting the future, but has been

instrumental in inspiring famous pioneers. Examples are cited.

4 Hirsch, D. & Zimmerman, H., Starlog Photo Guidebook to Spaceships (Enlarged Edition), Starlog Press, O'Quinn Studios, NY NY, 1980, 2nd printing, ISBN 0-931064-23-6, (1982) Book; (98 pages); Chronicles the cinematic and television art of science fiction space vehicles from their beginning through the publication date. Photos of the vehicles and brief descriptions of the movies or series from which they came are provided.

5. Herbert, N., Faster Than Light, Superluminal Loopholes in Physics, Plume Books, New York (1988,89).

Book; Presents the variety of evidence and speculation toward faster-than-light possibilities.

6. Hujasak, John T, & Hujasak Edward, Interstellar Propulsion Society A non-profit organization for professional and public membership that provides a means for scientists and engineers, worldwide, to join in collaborative efforts to accelerate scientific and engineering advancements in space propulsion, leading to manned missions to other star systems at fractional light speeds, relativistic velocities and beyond. Its World Wide Web Internet address is:

<http://www.digimark.net/ips/>.


7. Kurtis,B., producer, "The Science of Star Trek,"; a video episode of "THE NEW EXPLORERS"; series, Kurtis Productions, Ltd. and WTTW Chicago, aired January 1995.

One hour video comparing the science fiction of the popular series "Star Trek"; to the science fact of today.

Introductions to Emerging Physics:

8. Bennett, G., "Warp Speed, Fact or Fiction?", In Final Frontier, p. 35-39,

- (September-October 1994) Article (5 page); Discusses the issues and unanswered questions of Faster-Than-Light phenomena; causality issues, time travel implications, tachyon theory requiring imaginary mass, and wormhole theory. It mentions the conclusions of the May 1994 workshop at NASA JPL on this topic.
9. Szpir, M, "Spacetime Hypersurfing?", In American Scientist, Vol. 82, p. 422-423, (September-October 1994) Article (2 page); Explanation of the 1994 Alcubierre "warp drive"; article, including a copy of the figure of the warped spacetime from Alcubierre's paper. Discusses the necessity for the existence of negative energy density and the relation of that issue to the Casimir effect.
10. Gonick, L., "Science Classics (Warp-and-woof drive)", In Discover, p. 44- 54, (DEC 1994). Comic-Strip (2 page); Playful and succinct explanation of the 1994 Alcubierre "warp drive"; article.
11. Peterson, I., "A New Gravity?, Challenging Einstein's General Theory of Relativity", In Science News, Vol. 146, p. 376-378, (Dec 3, 1994). Article; Introduces the theory of Yilmaz and Alley that suggest that corrections are required to the Einstein field equations in order for them to correctly predict real behavior. The article also introduces experiments planned by NASA Goddard to test the theory by measuring the light propagation velocities over two different paths.
12. Matthews, R., "Inertia: Does Empty Space Put Up the Resistance?", In Science, Vol. 263, p. 612-613, (4 Feb 1994). Article (2 pages); An introduction to the theory of Haisch, et al. (Phys Rev A, Feb 94), where inertia is described as a high-frequency electromagnetic drag relative to the Zero Point Fluctuations of the vacuum. An important note is that this article mentions an experiment that would have a bearing on this theory: Exposing high-energy electrons to a terawatt beam from a neodymium-YAG laser, an experiment planned by K.McDonald et. al. @ Stanford Linear Accelerator Center, for late 1994.
13. Clarke, A. C., "Space Drive: A Fantasy That Could Become Reality", In Ad Astra, p. 38, (Nov-Dec 1994) Article (1 page); Arthur C. Clarke suggests that the time has come to revisit the idea of the "mythical space drive"; based on the recent theories of Haisch, Rueda and Puthoff (Phys Rev A, Feb 94).
14. Von Baeyer,H.C., "Vacuum Matters", In Discover, Vol. 13, No. 3, p. 108- 112, (March 1992). Article: Description of vacuum properties, including a historical perspective. Describes Zero Point Energy, Virtual Pairs, and the Casimir effect. Introduces recent experiments by D.Kleppner of MIT and N.Lawandy and J.Martorell of Brown University where a Casimir cavity is used to inhibit the spontaneous emission of exited atoms.
15. Boyer,T.H., "The Classical Vacuum", In Scientific American, p. 70-78, (Aug 1985). Article (10 pg): Summary of the contents of the vacuum: Thermal radiation; Zero Point Energy; Casimir Effect; and the ZPE as viewed during constant and accelerated motion.
16. Muller,R.A., "The Cosmic Background Radiation and the new Aether Drift", In Scientific American, Vol. 238, N. 5, p. 64-74, (May 1978). Article; Introduction to the CBR. "The three-degree cosmic background radiation provides an all-pervasive radiation 'aether' ..." (it also mentions Mach's Principle) ... against which we can measure our velocity.



(Earth/Sun are moving about 300km around our galaxy which itself is moving 600 km/sec relative to the universe. The universe is rotating less than 10-9 arc-sec/century) There is some discussion of the expansion of the universe and the slight non-isotropic nature of the CBR. Most of the article deals with the difficulties and methods of measuring the CBR.

17. Krauss, L. M., "Dark Matter in the Universe", In Scientific American, p. 58-68, (Dec 1986). Article; Describes evidence for and the theories leading to the conclusion that there must be more matter in the universe than is visible. Estimates range that only 2%-30% of the mass is visible.

The missing mass is called "dark matter."; Unconfirmed candidates for this missing mass are the neutrino and the following forms of "cold dark matter:"; axions, photinos, and cosmic strings. There is no mention of ZPE or virtual pairs as candidate explanations. The evidence: observed rotation rate of spiral galaxies is too high; Hubble's constant and the assumption of a "flat" universe requires more mass than is observed; Viral Theorem and observations of the relative velocities of stellar masses for stable systems requires more mass than is observed, and nucleosynthesis which predicts proportions between light elements and heavier elements requires more mass than is observed. The assumption that the universe is flat is based on the isotropy of the CBR and theories that predict that a non-flat universe would be obvious.

18. Hough,J., (Dept. of Natural Philosophy, Univ. of Glasgow, Glasgow G12 8QQ, UK), "New Detectors for Gravity Waves", In Nature, Vol. 316, p. 576-577, (15 AUG 1985).

Article (2 pg); Clear explanation of a variety of gravitational wave detectors, who is doing what, and speculates on sources of gravitational radiation.


19. Pool,R., "Closing in On Einstein's Special Theory of Relativity", In Science, Vol. 250, p. 1207-1208, (Nov. 9, 1990).

Article; Summarizes a variety of recent positive tests of relativity theory.

Detailed Technical Papers:

20. Alcubierre, M., (Dept. of Physics and Astronomy, Univ. of Wales, College of Cardiff CF1 3YB, UK), "The warp drive: hyper-fast travel within general relativity", In Classical and Quantum Gravity, Vol 11, p. L73-L77, (1994).

Letter to the editor (5 pg); Quoting from the abstract: "It is shown how, within the framework of general relativity and without the introduction of wormholes, it is possible to modify space-time in a way that allows a spaceship to travel with an arbitrarily large speed. By a purely local expansion of space-time behind the spaceship and an opposite contraction in front of it, motion faster than the speed of light as seen by observers outside the disturbed region is possible. The resulting distortion is reminiscent of the "warp drive"; of science fiction. However, just as happens with wormholes, exotic matter will be needed in order to generate a distortion of space-time like the one discussed here."; It is unknown in physics whether such "exotic matter"; having a negative mass or negative energy density can exist. Classical physics tends toward a "no," while quantum physics leans to a "maybe, yes."; Equal amounts of positive energy density matter will also be required. There is presently no known way to induce or control such effects. It is also uncertain whether this whole "warp"; would indeed move faster than the speed of light. Even though light-speed is a limit within spacetime, the rate at which spacetime itself can expand or contract is an open issue. Back during the early moments of the Big Bang, spacetime is assumed to have been able to expand faster than the speed of light. This is known



as the "inflationary universe" perspective. 21. Haisch, B. (Lockheed Palo Alto CA 94304), Rueda, A. (Dept. of E.E.

California State Univ.

Long Beach CA 90840) & Puthoff, H.E. (Inst. for Advanced Studies at Austin TX 78759); "Inertia as a Zero-Point Field Lorentz Force"; In Physical Review A, Vol. 49, No. 2, p. 678-694, (FEB 1994) Article: Asserts that inertia, the resistance to a change in velocity, is a high-frequency electromagnetic drag relative to the Zero Point Fluctuations of the vacuum (there is also a lower frequency drag effect called the Davies-Unruh effect, 1976). Inertial mass is calculated and compared to the gravitational mass calculated from Puthoff's 1989 paper that states that gravity is a consequence of ZPF interactions. They are off by a factor of two which is discussed. Unfortunately, this perspective does yet not provide clues for how to experimentally manipulate inertia or gravity using electromagnetics. The paper offers no examples of calculating mass from this perspective, such as calculating the mass of a neutron. Theoretically, the possible weak links of the theory are the reliance on "parton"; perspectives (assuming that all matter is fundamentally constructed of charged particles) and the Abraham-Lorentz radiation damping equation.

22. Puthoff,H.E., (Institute for Advanced Studies at Austin, Austin TX 78746), "Gravity as a zero-point-fluctuation force", In Physical Review A, Vol. 39, N. 5, (A89- 33278), p. 2333-2342, (Mar 1, 1989).

Article; Describes gravity as a side-effect of the Zero Point Fluctuation effects on charged matter. It assumes that all matter is fundamentally constructed of charged particles. Specifically it equates gravity to Van der Waals forces resulting from the dipoles induced by ZPE oscillations. Newton's $G = \frac{c^5}{(h\bar{\nu})} \times (\text{cut-off frequency } \sim 1042 \text{ Hz})$ Unfortunately, this perspective does not yet provide clues for how to

experimentally manipulate gravity using electromagnetics.

23.Yilmaz, H. (Hamamatsu Photonics, K. K. Hamamatsu City, 435 Japan, Electro-Optics Technology Center, Tufts Univ. Medford Mass, 02155 USA), "Toward a Field Theory of Gravitation", In Il Nuovo Cimento, Vol. 107B, no. 8, p. 941-960, (1991?).

Article; Suggests that corrections are required to the Einstein field equations in order to correctly predict behavior for systems beyond the simple one-body problem (the case of a single gravitating body affecting an insignificant test particle). The required correction is to add a term for the gravitational field energy itself to the matter tensor. This implies that the gravitational field has a mass-energy equivalence itself.

24.Misner,C.W., Thorne,K.S., and Wheeler,J.A., Gravitation, W.H.Freeman & Co., New York (1973).

Book; (1279 pages) This is a primary text book on General Relativity and related physics.

25.Physics Through the 1990's: Gravitation, Cosmology, and Cosmic Ray Physics, PB 86-241486, National Research Council, Washington DC, for the Department of Defense, (ISBN 0-309-03579-1), (1986).

Report (187 pages): Easy to understand compilation of the "State Of the Art"; of experimental work on the subject and recommends for future work, including candidate space experiments.

26.Forward,R.L., (Forward Unlimited, Malibu CA 90265-7783), "Space Warps: A Review of One Form of Propulsionless Transport", AIAA-89- 2332, 25th Joint Propulsion Conference, Monterey CA, (1989).

Paper; Examines a variety of ideas for creating worm holes, space warps or other propulsive methods based on theories from

General Relativity. None of the methods can be practically accomplished.

27.Winterberg,F., (Desert Research Institute, Univ of Nevada System, Reno, Nevada), "On Negative Mass Propulsion", AIF Paper 89-668, 40th Congress of the International Astronautical Federation, Malaga, Spain, (Oct, 1989)..

Article; A propulsion scheme is described consisting of negative and normal mass side-by-side. Because of the attraction / repulsion properties of the respective masses, both masses accelerate continuously in the direction pointing from the negative to the positive mass. Suggests that there is experimental evidence for the existence of negative mass. It also discusses an older interpretation of relativity by Lorentz and Poincar, that relies on an absolute reference frame; ether. This treatment is consistent with experimental observations including the Michelson-Morley experiment. The null result of the Michelson-Morley experiment is explained that the apparatus is equally affected as the light, canceling any observed effect. It also discussed the role of ZPE and mentions Zitterbewegung in the context of experimental evidence for the existence of negative mass.

28.Forward,R.L., (Forward Unlimited, Malibu CA 90265-7783), "Negative Matter Propulsion", AIAA-88-3168 preprint, 24th Joint Propulsion Conference, Boston, Mass., (1988). {Note: a shorter version that omits the gravitational coupling details exists as: Forward,R.L., "Negative Matter Propulsion", In AIAA Journal of Propulsion, Vol. 6 No. 1, p. 28- 37, (Jan-Feb 1990)} Paper (19 pg); Shows that the negative mass propulsion scheme does NOT violate conservation of momentum even though no reaction mass is expelled. Three cases are analyzed classically; where the masses are gravitationally coupled, coupled with an ideal spring, and coupled by electrostatic forces. Each of these couplings is analyzed assuming the magnitudes of the negative and positive masses are equal, and again assuming they are unequal. For all situations,


energy and momentum are found to be conserved. It is important to note that in the case of gravitational coupling, the analysis is simplified by setting the initial velocities such that their momentums sum to zero. This zero sum continues throughout the different resulting motions of each mass. For the case of spring-coupled unequal masses, oscillations occur and energy is conserved using the potential energy of the spring.

29 Millis,M.G., (NASA, Lewis Research Center), "Exploring the Notion of Space Coupling Propulsion", In Vision 21: Space Travel for the Next Millennium, Symposium Proceedings, Apr 1990, NASA-CP-10059, p. 307- 316, (1990).

Paper; Identifies the 3 major objections to the notion of propelling a vehicle by coupling to the properties of space: (1) violates conservation of momentum, (2) no reactive media in space, (3) no grand unified theories to link gravity with other controllable phenomena. Introduces a variety of avenues to possibly satisfy these objections, including introducing the notion of coupling to distant masses via the fundamental structure of the intervening space and the associated need for alternative treatments of space physics.

30.Bennett, G.L. & Knowles, H.B., (NASA HQ & Physics consulting, El SoBrante, CA 94803), "Boundary Conditions on Faster-Than-Light Transport Systems", AIAA paper 93-1995, presented at the 29th Joint Propulsion Conference, Monterey CA, (1993).

Paper (16 pg); Outlines the prerequisite physics to enable Faster-Than-Light transport: The theory of FTL must equally describe sub-light physics, must not violate causality, must match physics at quantum scales, and must satisfy conservation of energy. The paper discusses Tachyons and Wormholes concluding that it is "very unlikely that interactive tachyons exist";, and the we don't know enough about Wormholes to render judgement. 31. Forward,R.L., (Hughes Research Lab, Malibu CA 90265) "Feasibility of Interstellar Travel: A Review",



In Acta Astronautica, Vol. 14, p 243-252, (1986) Article: Comments on a variety of scientifically plausible propulsion schemes. States that Centauri cluster is reachable within 50 yrs given a velocity of 10% lightspeed, and 25 stars are within reach given 30% lightspeed. Various propulsion schemes are compared, but the comparisons were not on equal footing:

Nuclear Electric Propulsion: 10,000 year trip
Nuclear (Fission) Pulse Propulsion: 130 year trip.

Antimatter Propulsion: 50 year mission requires 9 kg of anti-hydrogen for 1 ton payload.

Nuclear Fusion: To be determined.

Interstellar Ramjet: To be determined, particularly the issue of fusing protons without having to slow them down.

Beamed Power: 20 year trip for a crewed vehicle, but needs lasers that are about 10,000 times more powerful than all the power used on the Earth today.



Distribution A:

Approved for public release; distribution is unlimited.

Document created: 25 March 02
Aerospace Power Journal - Spring 2002

Operation LUSTY

The US Army Air Forces' Exploitation of the Luftwaffe's Secret Aeronautical Technology, 1944–45

Dik Alan Daso, PhD

Editorial Abstract: The bewildering pace of development in aerospace-power technology immediately following World War II was no accident. The author's account of the highly successful efforts of Army Air Corps leaders to exploit German technology at the end of the war is a story that still has lessons for us today.


IN New World Vistas, the US Air Force's science and technology (S&T) study of 1995, Dr. Gene McCall wrote about the relationship of technology to the Air Force after almost 50 years as an independent service: "It was clear in 1945 that the technology gains of the first half of the twentieth century should be consolidated to create a superior, technology- and capability-based Air Force which could respond to threats not yet imagined. The world which emerged from the destruction of World War II could not have been predicted in 1945, but the emphasis on technology and capability rather than on assumptions about future geopolitical scenarios served us well as we entered the Cold War."¹

Technology is fundamental to the culture of the US Air Force. For the most part, this technology culture appeared at the same time as the air service itself, due to the

nature of heavier-than-air flight. For nearly a century, technological progress has occurred in starts and fits as well as leaps and bounds, exploding geometrically as it accompanied the visionary efforts of key individuals and programs. In conducting analyses of technological efficiencies in anticipation of tomorrow's complex threat environment, one would do well to consider the past successes of some of these key players in technological development. In particular, a seminal turning point occurred on the heels of World War II as part of a plan to exploit German scientific advancements. The plan was called Operation LUSTY (for Luftwaffe secret technology).

Technological change during World War II proceeded at a frightening pace. Developments in aircraft design, propulsion, weapons, and electronics contributed vitally to the outcome of events in the global conflict. At the heart of these developments were scientists, largely civilians, who worked to produce military equipment that would turn the tide of the war. Among them was the youthful Hungarian aerodynamicist Dr. Theodore von Kármán. Since his arrival in the United States from Europe, having obtained Guggenheim funding and hoping to avoid rising nationalism and Nazism, he had become acquainted with several Army air officers, among them a young major named Henry "Hap" Arnold, who would later command the US Army Air Forces (AAF) throughout World War II.

Since their first meeting at the California Institute of Technology (Caltech) in the early 1930s, Arnold had witnessed the professor's skilled use of mathematical equations to solve complex aerodynamic problems. After inheriting command of the Army Air Corps in 1938 and driven by a



near- obsessive belief in the efficacy of scientific approaches to Air Corps problems, Arnold called civilian scientists to a meeting at the National Academy of Sciences building in Washington, D.C., in 1939. Among the visitors was a team from Caltech, including Kármán. At that meeting, Arnold doled out scientific projects, such as finding a solution to high-altitude windshield icing and developing aircraft radios and jet-assisted takeoff (although the term jet was a misnomer). Kármán assigned the difficult rocket project to his most senior students at Caltech, the “suicide club.” From that small project grew what is today the Jet Propulsion Laboratory near Pasadena, California. More importantly, Arnold’s trust in Kármán grew as the Caltech program continued to tackle the most difficult projects without hesitation. Arnold did not tolerate a “no-can-do” attitude.


By war’s end, General Arnold had decided that the AAF was in a position to capitalize on World War II’s many technological developments. He also realized that the United States and its Allies by no means led the world in military aeronautical development. He used his influence with Kármán, convincing him to head a task force of scientists who would evaluate captured German aeronautical data and laboratories for the AAF. As the Allies advanced into Europe during the spring of 1945, Kármán’s team, close on the heels of the advancing wave, scoured German laboratories. For the AAF, Operation LUSTY began during a supersecret meeting between General Arnold and Dr. von Kármán on the runway at LaGuardia Airport, New York.²

Only after D day and the realization of several key elements in wartime operations

did Arnold believe that Allied victory in Europe was a foregone conclusion. The air war had become a deadly routine. At that point, it was merely a numbers game—Allied air strength versus dwindling Axis air capability.

The Normandy invasion occurred under the umbrella of air supremacy. The P-51 had operated successfully with drop tanks for several months with encouraging results. Additionally, B-29 production had increased to acceptable levels. For the operational needs of combat, this long-range, heavy bomber became Arnold’s Pacific trump card. He had devoted a great deal of personal effort to ensure its development, despite severe engine problems initially. Only after assuring himself that these production and procurement programs were succeeding did the general set his sights on developing S&T for the AAF.³

General Arnold and Dr. von Kármán stayed in “continual conference” after the LaGuardia encounter. Kármán recalled that he was “more impressed than ever with Arnold’s vision,”⁴ and Arnold insisted that Kármán examine everything and let his “imagination run wild.”⁵ This challenge fitted perfectly into Kármán’s philosophy, including the belief that imagination was a vital part of the invention process.⁶ To ensure the excellence of this crucial task, Arnold imposed no completion deadline (a luxury he later rescinded) and insisted that Kármán’s group travel to many foreign countries, assess their aeronautics programs, and then fashion a bold final report—a viable forecast for maintaining future American air supremacy.⁷ Arnold’s establishment of the forecasting group itself was totally secret—almost “cloak and dagger.”⁸ To accomplish his mission, Kármán officially became an AAF



consultant on scientific matters on 23 October 1944.⁹


Kármán's first, unofficial AAF report was organizational in nature, naming as his deputy Dr. Hugh L. Dryden, long-time head of the National Bureau of Standards. November 1944 saw endless conferences and establishment of "relations with the various agencies in the labyrinth of military and scientific aviation."¹⁰ Arnold drafted official, written instructions on 7 November, solidifying the LaGuardia Agreement, a four-page letter that set the boundaries for the report of Kármán's group. They were not very restrictive: "Except perhaps to review current techniques and research trends, I am asking you and your associates to divorce yourselves from the present war in order to investigate all the possibilities and desirabilities for postwar and future war's development as respects the AAF. Upon completion of your studies, please then give me a report or guide for recommended future AAF research and development [R&D] programs."¹¹ Initially, Kármán's group was called the AAF Consulting Board for Future Research, but apparently AAFCBFR proved too long an acronym, even for the Army. Redesignated the Scientific Advisory Group (SAG) on 1 December 1944, it reported directly to General Arnold.¹²

Germany's last, desperate attempt to end the war at the Bulge occurred as the scientists gathered, anticipating their chance to exploit the work that German scientists had done over the last five to seven years. In January 1945, Kármán's handpicked, scientific team of "thirty-one giant brains" congregated in Washington to begin the monumental task Arnold had given them. Initially, Kármán met internal resistance to a few of his choices for the

group—for example, Sir William Hawthorne, an Englishman. Col Frederick E. "Fritz" Glantzberg, Kármán's military assistant, voiced his objection to having any "foreigners" in the group. Kármán reminded the colonel that Arnold wanted the best people, regardless of their origins. Glantzberg relented, conceding that "the British were, after all, our Allies." Kármán also insisted upon adding a naval officer, William Bollay (a former Caltech student). When the colonel insisted that the professor had gone too far, Kármán responded with the simple question, "But Colonel, the Navy are surely our Allies too?" After considering this for a moment, Glantzberg finally agreed that they were: "Not as close as the British, but a damn sight closer than the Russians."¹³ For administrative reasons, neither of these men served in the group until 1949, but Arnold wanted the best and did not care how Kármán carried out that order. Arnold envisioned and enacted the concept of "jointness" long before the term was formalized in the Pentagon almost half a century later with the enactment of the Goldwater-Nichols Department of Defense Reorganization Act.

A five-star general since December 1944, Arnold insisted that the group throw conservative thinking to the wind. Kármán then reminded the scientists in his quiet, broken English that they had to deliver on their promises. Unsurprisingly, the younger members of the team found working in the SAG the "equivalent of a semester of grad school each day."¹⁴ In mid-January 1945, Arnold suffered a severe heart attack and retreated to Florida to recuperate. Fortunately, he had already given Kármán his marching orders.¹⁵

SAG meetings held during the first weeks in February, March, and April



accomplished the basic research and finalized the general format for the report. Kármán emphasized that these spring meetings had a threefold purpose: (1) the SAG would search for ways to secure “scientific insight in a standing Air Force”; (2) it would ensure the continued interest of American scientists in the future of the Air Force; and (3) the group would educate the American public in the necessity of maintaining a strong Air Force.¹⁶ These objectives may have seemed remarkably vague, but specifics in design and engineering were not really part of the SAG’s overall task. Actually, this sweeping view predated America’s entry into the war. In the Pasadena Star News of 24 February 1941, reporters quoted Kármán as saying, “So rapid has been the development of military aircraft during the present war, it is impossible to forecast what performance limits will be obtained by warplanes before the war ends.”¹⁷ For reasons such as this, a broad approach to technology forecasting remained uppermost in Kármán’s mind, and he convinced Arnold of the same.

As mentioned above, in late April 1945, SAG members departed for Europe to inspect liberated enemy laboratories. Operation LUSTY, a name that the cosmopolitan, unmarried young professor sardonically suggested was “unlikely but pleasant,” fulfilled Arnold’s insistence that the SAG investigate the most advanced S&T aeronautical information available worldwide.¹⁸ LUSTY was the code name for a much larger operational, exploitation expedition of European technologies initiated by the US Army, of which the SAG represented only one small part. Arnold’s instructions to Gen Carl “Tooey” Spaatz, commander of US Strategic Air Forces in Europe, were crystal clear: “May I ask . . . in view of the importance of this

project that you give it your personal attention.”¹⁹ Already alerted to Arnold’s belief in science, Spaatz did just that. In September 1944, while traveling to the second wartime Quebec Conference, Arnold had informed Spaatz of his belief in the “value and the importance of these long-haired scientists.”²⁰ Already, Arnold had secretly established the SAG as proof of this commitment. Spaatz’s immediate cooperation was vital to the success of the SAG’s contribution to Operation LUSTY.

After the scientists arrived in Paris on 1 May 1945, one member of the team, H. Guyford Stever, observed the critical nature of timing during the Allied advance. He recalled that, although local looting often presented a problem, the advancing Russians were the real concern. More significantly, Stever mentioned that “until this von Kármán mission, we [scientists] had to piece the enemy’s facts together. Now we had the advantage of actually talking to the German scientists and engineers, seeing their laboratories, and hearing them describe their total programs.”²¹ Dr. Dryden echoed Stever’s conclusion: “I think we found out more about what had been going on in the war in a few days [sic] conversations with some of these key German leaders, than all the running around and digging for drawings and models . . . could bring.”²² Only after Kármán arrived did the group discover the scope of Germany’s S&T efforts.

To preserve that scientific picture, the American teams boxed up everything they could and immediately shipped it to Wright Field, Ohio, the AAF’s center for aeronautical R&D. First on the scene at one location, Navy exploitation teams quickly boxed up the hardware and technical data in large crates and labeled them “US Navy.” Two days later, Army



teams made it to the same location, whereupon they crated the Navy boxes in larger crates and relabeled them “US Army.”²³ For these reasons—some good, some ridiculous—immediate access to targets became crucial. Spaatz provided the transportation capability to meet these requirements.²⁴ His personal involvement in the early days of the SAG’s visit to Europe helped strengthen his own understanding of its capabilities during his tour as the first chief of staff of the independent Air Force.

Among the discoveries in Germany during the “scientists’ invasion” were rocket-propelled fighter planes, radio-controlled bombs, guided antiaircraft missiles, and practically every type of fighter aircraft in the Luftwaffe inventory. The most surprising ones included a jet-powered helicopter built by Doblhoff, swept-back wings hung in high-speed wind tunnels, hidden assembly locations for V-1 and V-2 “vengeance” weapons, and plans for V-3 (intercontinental) rockets capable of reaching targets on the east coast of the United States; many of these rockets were deep below ground in hidden caves. Under the watchful eye of American scientists, including Kármán, German technicians launched several of the V-2 rockets from test sites during the summer and fall months.


Perhaps of even greater significance were thousands of linear feet of data and documents that accompanied these projects—the teams shipped more than 100,000 tons to a London clearinghouse that spring. Upon close examination, many of these German studies confirmed the path that American science had already taken. Some, the jet-powered helicopter for instance (the fourth modification of the

original autorotator design), came as a total surprise.²⁵

After six weeks of traveling throughout the devastated European countryside, the professor met Arnold—now recovered from his January heart attack—in Paris on 13 July 1945 to discuss the team’s initial findings. General Arnold, who was traveling to join President Truman at Potsdam, Germany, and did not have much time, asked the professor to prepare a report that summarized the SAG’s discoveries. Kármán submitted Where We Stand on 22 August, satisfying that request.

This summary of the exploitation of German S&T that Kármán’s men had unearthed began by listing a set of eight aspects of aerial warfare which, Kármán believed, had become “fundamental realities”:

1. Aircraft- manned or pilotless- will move with speeds far beyond the velocity of sound.
2. Due to improvements in aerodynamics, propulsion, and electronic control, unmanned devices will transport means of destruction to targets at distances up to several thousand miles.
3. Small amounts of explosive material will cause destruction over areas of several square miles.
4. Defense against present-day aircraft will be perfected by target-seeking missiles.
5. Only aircraft or missiles moving at extreme speeds will be able to



- penetrate enemy territory protected by such defenses.
6. A perfect communication system between fighter command and each individual aircraft will be established.
 7. Location and observation of targets; takeoff, navigation, and landing of aircraft; and communication will be independent of visibility and weather.
 8. Fully equipped airborne task forces will be enabled to strike at far-distant points and will be supplied by air.²⁶

Additionally, the report sought to explain why Germany was more advanced in some areas but lagged in others. The title itself reflected Kármán's evaluation of US posture in regard to foreign scientific developments.

For example, the report did not attribute German achievements in aeronautics to superior scientists but to "very substantial support enjoyed by their research institutions in obtaining expensive research equipment such as large supersonic wind tunnels many years before such equipment was planned in this country."²⁷ These tunnels supported development in the field of transonic and supersonic wing design to the point of "practical application," whereas advanced design ideas were only at the discussion stage in America, spearheaded by Kármán and others after the Volta Conference of 1935.

Kármán's summary added a warning: "We cannot hope to secure air superiority in any future conflict without entering the supersonic speed range." Additionally, the

report stated that "V-2 development was successful not so much because of striking scientific developments [but] because of an early start, military support, and boldness of execution."²⁸ An early start, unlimited funding, and bold execution of German scientific plans became a recurring theme throughout the report.

However, the United States held substantial leads over the Axis in some areas, such as radar development:

It must be realized that radar is not a facility of attachment which will occasionally be used under bad conditions. Rather, the Air Force of the future will be operated so that radar is the primary facility, and visual methods will only occasionally be used. . . . Hence, in an all-weather Air Force, radar must be the universally used tool for bombing, gunfire, navigation, landing, and control. The whole structure of the Air Force, the planning of its operations, its training program, and its organization must be based on this premise. The development and perfection of radar and the techniques for using it effectively are as important as the development of the jet-propelled plane (emphasis in original).²⁹

Today, this realization appears the most prescient of all those made during a period when the AAF's primary doctrine (in Europe certainly)—that of precision, strategic, daylight bombing—was based largely on the ability to acquire the intended target visually.³⁰ Kármán also pointed out that the Germans had failed to keep stride with the rest of the world because "most of the development took place in industrial laboratories . . . but the very brilliant group of German physicists in universities were never called in to participate. Consequently, while



engineering design was good, imaginative new thinking was lacking.” Kármán could detect the absence of imagination and individual brilliance—whether in his students or in notable scientists. Further, he predicted that “the ability to achieve Air Force operations under all conditions of darkness and weather contributes more than any other single factor to increasing the military effectiveness of the air forces. Hence, any research program designed to overcome the limitations to flight at night and in bad weather will pay big dividends.” Aware of the rapid improvements in radar technology, the professor suggested that the Air Force “be alert in swiftly utilizing any new developments.”³¹


By emphasizing radar, Kármán also indirectly assured that the Massachusetts Institute of Technology (MIT) would share in future military research projects. During the war, the MIT Radiation Laboratory led the development of American radar. Generally, just as Caltech held the reins of AAF aeronautical science, so did MIT direct AAF radar programs. In fact, the addition of Dr. Edward Bowles to Arnold’s staff in 1943 linked radar and electronic programs to the AAF, much as Kármán’s association had linked aeronautics in earlier years. The rivalry that developed between these schools was friendlier than Caltech’s rivalry with the National Advisory Committee for Aeronautics (NACA). Both schools held particular expertise in different areas of technological development, and, for the most part, each respected the other’s accomplishments.³²

After publication of this initial report, Kármán began the arduous task of compiling the SAG’s detailed work. Suddenly, the deliberate pace normally associated with scientific research was

replaced by a great sense of urgency to complete the project. Fearing radical budget cuts at war’s end, Arnold cabled Kármán, still in Europe, wondering if the report might be finalized by 15 December 1945. To accommodate the general’s request, Kármán canceled an inspection of Japanese aeronautical laboratories (which he had helped to establish at Kobe in 1927) and sent a few of his team members to the Orient instead. From October through December, work proceeded at a frenetic pace. After many sleepless nights, Kármán had the draft version of the final report, *Toward New Horizons*, delivered to Arnold’s desk on 15 December 1945.³³

Kármán’s summary volume, *Science: The Key to Air Supremacy*, introduced the classified 12-volume report.³⁴ In essence, this volume amplified the tenets of the August report with a few significant additions. It addressed the problems associated with “research and development from the point of view of the technical requirements which the Air Force must meet in order to carry out its task, securing the safety of the nation.” The third chapter elaborated upon correcting the organizational and administrative problems addressed in *Where We Stand*. Most notable of these was a plea for government authority to “foster,” not “dictate,” basic research.³⁵ This long-range, extremely detailed study was the first of its kind in American military history. Along with *Where We Stand*, it would serve as the blueprint for building the Air Force during the next two decades.

General Arnold was so interested in the possibilities of future airpower development that, based upon Kármán’s preliminary report, he offered his personal perceptions of the SAG’s importance to General Spaatz. Arnold reminded Spaatz,




his successor, that the AAF had no great scientists in its ranks. Military R&D labs had stagnated during the war, largely due to increased production requirements and personnel shortages. The AAF had required civilian help during the war to solve aircraft power-plant and structural-design problems. Only with civilian assistance had the service been able to realize its S&T potential. Arnold reminded Spaatz that “these men did things that the average Army officer could never have accomplished. We must not lose these contacts.”³⁶ Today, through organizations such as the Scientific Advisory Board (SAB), the Air Force continues this tradition through a variety of contacts in civilian industry and academia.

Spaatz took Arnold’s advice to heart and established the SAB as a permanent group; it met for the first time on 17 June 1946. It was not, however, attached to the commanding general, as Kármán had suggested, but was relegated to Gen Curtis E. LeMay, deputy chief of the Air Staff for R&D.³⁷ Nevertheless, the SAB survived postwar cuts by providing scientific advice to higher levels of Air Force leadership. The imperfection of the new system was eventually repaired.

The Arnold/Kármán team, although it existed officially only from November 1944 to February 1946, created the plan that has since evolved into the S&T infrastructure of today’s Air Force. Gen Bernard Schriever, father of the Air Force’s missile program, once said of Arnold, “There’s no question, his greatness was that he created the infrastructure. He visualized the kind of infrastructure that the Air Force needed to really get into the technology age.”³⁸

In addition to the SAG, Arnold established Project RAND and the Office of Scientific Liaison and funded dozens of guided-missile programs before postwar demobilization and inevitable budget cuts slowed the procedure. Kármán and the SAG assumed the strenuous task of traveling the world in search of the most advanced technologies, constantly mindful of how these advances might be applied to American airpower. In the end, Arnold’s recognition of the need for such a study and Kármán’s unique ability to apply scientific findings to the practical, technological needs of the Air Force — helped along by lifelong associates at opportune times and places—produced a report that had great potential for long-term success.

Such success, however, carried no guarantees. Initially, funds were scarce, and leadership was in constant flux. The reorganization of the National Military Establishment into the Department of Defense only added to the quagmire. Somehow, by the nature of his association with both officers and scientists, Kármán prevented the newly formed SAB from stalling. He nurtured its structure and function in the face of misunderstandings, opposing agendas, and, at times, lack of interest until the board was capable of independent growth. By 1954, a decade after the process began, Kármán’s vision-guided by his own perseverance—proved directly responsible for sustaining the SAB. Yet, Arnold’s ghost was never far away. During this period, the general’s lifelong associations with officers, industrialists, and scientists continually surfaced. Such people as General Schriever, Lt Gen Laurence Craigie, Lt Gen Benjamin Chidlaw, Larry Bell, Donald Douglas, and Dr. Bowles were all



vital to the eventual institutionalization of Toward New Horizons.

Only after Kármán was certain that the SAB would thrive did he resign his chair. Similarly, just as Arnold's influence continued to be felt, so did Kármán's. Former students, colleagues, and friends who had been educated by or employed with the professor carried his broad-based, practical- applications approach to problem solving as part of their own methodology—in several cases, into the mid-1990s. Arnold's drive and Kármán's method, embodied in the institutionalized SAB, had become the Air Force's foundation for S&T matters.


Airpower institutions have evolved erratically since World War I. In Ideas and Weapons, I. B. Holley concluded that “the postwar [World War I] Air Service made use of only a relatively small portion of the experience of the war regarding the problem of weapons.”³⁹ One lesson learned, however, was that quality was preferable to quantity as far as weapons were concerned. Arnold had internalized that lesson. Unfortunately, administrative organizations that might have assured high-quality weapons development during those years had been neglected. Another lesson learned—perhaps the most significant one—concerned unity of command. According to Holley, “the available evidence shows that after the war the Air Service learned the importance of organization for decision and established channels of command for unified, decisive, and authoritative action in contrast to the dispersed, ill-defined, and overlapping channels that existed during the war.”⁴⁰ This very development allowed Arnold to act as a stopgap, ensuring that the lessons of the Great War had not fallen on totally deaf ears. Arnold acted as the AAF's

tangible link between the lessons of World War I and the institutionalization of S&T that became reality after World War II. Ordering the blueprint that became the S&T cornerstone of American airpower is one of Arnold's legacies—creating it is one of Kármán's.

As hundreds of American military soldiers and CIA operatives rummage through caves in war-torn Afghanistan in search of enemy documents and leaders, we are reminded that we have accomplished similar missions before. The intelligence retrieved during Operation LUSTY was part of the process by which the US Air Force entered its technological infancy. Over the past 60 years, developing air and space technology has created the means to improve the human condition or to bring devastating destruction to unsuspecting enemies anywhere in the world. Lines have become blurred between military and civilian aviation and space technology, just as it is difficult to distinguish among those who utilize these technologies for their own purposes.

Today, it remains clear that technology developments made during the twentieth century should be consolidated to create a superior technology- and capability-based twenty-first-century Air Force that will be able to respond to threats not yet imagined. What is unclear is the kind of world that is emerging from the rubble of the World Trade Center and Pentagon. The emphasis placed upon technology and capability that has been the hallmark of the Air Force over the past half century must be molded more and more by future geopolitical scenarios.

As much as ever before in Air Force history, national aerospace power faces difficult challenges in its quest to achieve



desired effects against complex, ruthless, and elusive enemies both at home and on foreign soil. Fortunately, because of the efforts of people like Arnold and Kármán, the Air Force now has many technological tools that contribute to achieving national-security objectives and attaining global stability. Today's perceived technological superiority is forcing the Air Force and the nation to reassess how best to apply these technologies during the uncertainties already developing during this new millennium.

Notes

1. Gene H. McCall and John A. Corder, *New World Vistas: Air and Space Power for the 21st Century: Summary Volume* (Washington, D.C.: Scientific Advisory Board, 1995), iii. For an examination of the technological culture of the Air Force, see Carl H. Builder, *The Icarus Syndrome: The Role of Air Power Theory in the Evolution and Fate of the U.S. Air Force* (New Brunswick, N.J.: Transaction Publishers, 1994). See also my article "New World Vistas: Looking toward the Future, Learning from the Past," *Aerospace Power Journal* 13, no. 4 (Winter 1999): 67–76. The author wishes to thank the USAF Historical Research Agency and Air University Press, both at Maxwell AFB, Ala., for their support early in the development of this article.
2. The LaGuardia meeting holds a quiet, yet legendary, place in the history of the Air Force's S&T development. In his later years, Kármán recalled the meeting but not the details. Arnold's plane arrived, jostled by the rough winds of a passing cold front, and Kármán, recovering from recent abdominal surgery, was transported by Army staff car to the end of the runway, where the general joined him after deplaning. Arnold dismissed the military driver and then, in total secrecy, discussed his plans for Kármán and his desires for the exploitation project. Arnold spoke of his concerns about the future of American airpower, and he wondered how jet propulsion, radar, rockets, and other "gadgets" might affect that future. "What do you wish me to do?" Kármán asked with a thick Hungarian accent. "I want you to come to the Pentagon and gather a group of scientists who will work out a blueprint for air research for the next 20, 30, perhaps 50 years," Arnold replied. After promising to give all of the orders on Kármán's behalf (the professor insisted on that caveat), Arnold hopped back in his plane, the deal done. Kármán, flattered and excited, was impressed that General Arnold had the vision to look beyond the war, seeking the help of university scientists. The timing of Arnold's request was not accidental. See Theodore von Kármán with Lee Edson, *The Wind and Beyond: Theodore von Kármán, Pioneer in Aviation and Pathfinder in Space* (Boston: Little, Brown, 1967).
3. Wesley Frank Craven and James Lea Cate, eds., *The Army Air Forces in World War II*, vol. 6, *Men and Planes* (1955; new imprint, Washington, D.C.: Office of Air Force History, 1983), 218–19.
4. Kármán, *The Wind and Beyond*, 267–68; and Clark Millikan to William Knudsen, letter, 3 October 1944.
5. Gen H. H. Arnold to Gen Carl Spaatz, letter, 6 December 1945; and Arnold to Gen Ira C. Eaker, letter, 22 May 1945, both located in the Hap Arnold Murray Green Collection, USAF Academy Library, Special Collections (hereinafter MGC). See also Craven and Cate, vol. 6, 234; and Thomas A. Sturm, *USAF Scientific Advisory Board: Its First Twenty Years, 1944–1964* (Washington, D.C.: USAF Historical Division Liaison Office, 1967), 37.
6. "Disney Folder," Kármán Papers, California Institute of Technology, no. 59.2.
7. Henry H. Arnold, *Global Mission* (New York: Harper and Brothers, 1949), 532–33, reinforced by a cable sent to Spaatz toward the end of the war, 15 April 1945, in MGC.
8. On 25 October, in a reply to a letter from Lt Gen George Kenney concerning future



- planning, Arnold detailed more than 30 specific actions pertaining to aircraft production and design, but he did not mention the Kármán project, already under way. Arnold added only a brief clue in a postscript: "There is still more that is being prepared now but will not be actuated until the Post-War Period." In a speech to the Aeronautical Research Laboratory of the National Advisory Committee for Aeronautics (NACA) on 9 November 1944, Arnold cryptically told the gathering of scientists and engineers that when the AAF got stuck in a development problem or when it looked toward the future of aeronautics, "normally we go to the NACA and ask you people to do that work for us." But Arnold would not go to the NACA this time. Just as he had secretly given the Whittle jet-engine-development problem to the Bell/ General Electric team in 1941, he now gave the critical task of forecasting the requirements for obtaining future air supremacy to Kármán and his scientists.
9. Scientific Advisory Board (SAB) Office, 1944–1945 file, Pentagon, Washington, D.C.
 10. Kármán's first report for the Scientific Advisory Group (SAG), 23 November 1944, in MGC; Kármán to Clark Millikan, letter, 4 November 1944, Kármán Papers, California Institute of Technology, no. 73.6; and Michael H. Gorn, *The Universal Man: Theodore von Kármán's Life in Aeronautics* (Washington, D.C.: Smithsonian Institution Press, 1992), 99.
 11. Arnold to Kármán, letter, subject: Instructions for Forecasting Group, 7 November 1944, SAB Office, 1944–1945 file, Pentagon, Washington, D.C.
 12. SAB Office, Pentagon, Washington, D.C.; Kármán's first report for the SAG, 23 November 1944, MGC, ref. L/C box 79; Gorn, 99–100; H. H. Arnold Papers, Headquarters Office Instruction 20-76, MGC, L/C box 40; and Arnold to Kármán, letter, 7 November 1944, SAB Office, Pentagon, Washington, D.C.
 13. Kármán, *The Wind and Beyond*, 269–70; and William Rees Sears, Stories from a Twentieth Century Life (Stanford, Calif.: Parabolic Press, 1993), 219.
 14. Kármán, interviewed by Shirley Thomas, cassette tape, University of Indiana Library, 1960; T. F. Walkowicz, "Von Kármán's Singular Contributions to Aerospace Power," *Air Force Magazine*, May 1981, 60–61; and Gorn, 47.
 15. Arnold to Lois Snowden, letter, 22 February 1945, MGC. The general described his condition to Lois, his daughter, in mechanical terms: "Apparently one of my cylinders blew a gasket and I had to get down here to have an overhaul job done. . . . While I was here they checked my lubrication, ignition, and gasoline system and they said they were working alright."
 16. Sturm, 5.
 17. "No Way to Predict Future of Warplane Performance," Pasadena Star News, 24 February 1941, Kármán Papers, California Institute of Technology, no. 157.2.
 18. Kármán, transcript of oral history interview, n.d., US Air Force Academy Library, Colorado Springs, Colo.; Chester Hasert, National Academy of Sciences, Washington, D.C., interviewed by author, 10 November 1994; and Operation LUSTY folder, Air Force Historical Research Agency, Maxwell AFB, Ala.
 19. Lt Gen Barney Giles (for Arnold) to Spaatz, letter, 19 April 1945, Kármán Papers, California Institute of Technology, no. 90.2; Kármán, *The Wind and Beyond*, 272; Gorn, 103–5; and Kármán, oral history interview.
 20. In a reply to an earlier letter praising radar developments, Arnold wrote Spaatz on 12 September 1944, affirming his trust in scientists, MGC, roll 12.
 21. H. Guyford Stever, Washington, D.C., interviewed by author, 18 May 1995.
 22. Hugh L. Dryden, Columbia University Oral History Review, 24.



23. Dr. Homerjoe Stewart, Pasadena, Calif., interviewed by author, 21 July 1995.
24. Stever interview. Dr. Stever was working with the British radiation laboratory as part of the MIT exchange team when LUSTY operations began. He was attached to Kármán's group in place of Dr. L. DuBridge, who was unavailable. Stever is a former chairman of the SAB from 1962 to 1964 and a former presidential science advisor.
25. Ibid.; Dr. Richard P. Hallion, interview for New World Vistas, videotape, Office of Air Force History, Bolling AFB, Washington, D.C.; summary of memo from Kármán to Arnold, 30 July 1945, which documented the group's travels to that point, MGC, roll 12; and "History of Operation Lusty, 6 June 1944–1 February 1945," US Air Force Historical Research Agency, Maxwell AFB, Ala., file 570.650A.
26. Theodore von Kármán, "Where We Stand: First Report to General of the Army H. H. Arnold on Long Range Research Problems of the Air Forces with a Review of German Plans and Developments, 22 August 1945," typed manuscript (Wright-Patterson AFB, Ohio: Air Force Materiel Command History Office, August 1945), 1–2. Sturm's USAF Scientific Advisory Board: Its First Twenty Years includes the evolution and decline of the group through 1964. Alan Gropman has nicely summarized the report itself in "Air Force Planning and the Technology Development Planning Process in the Post-World War II Air Force- The First Decade (1945–1955)," in Military Planning in the Twentieth Century: The Proceedings of the Eleventh Military History Symposium, USAFA, 10–12 October 1984 (Washington, D.C.: Office of Air Force History, 1986), 154–230. Kármán's reports may be found in the appendices of my book Architects of American Air Supremacy: Gen. Hap Arnold and Dr. Theodore von Kármán (Maxwell AFB, Ala.: Air University Press, 1997), the only publication in which the Kármán reports are published together and in their entirety.
27. Kármán, Where We Stand, 5.
28. Ibid., 8, 12, 21.
29. Ibid., 75–76.
30. Maj Gen Haywood Hansell Jr., to author, letter, 4 October 1979. Although the AAF did accomplish a limited number of area-bombing missions in Europe, these were supplemental to precision attacks in almost every case.
31. Daso, Architects of American Air Supremacy, 283–85.
32. Jack H. Nunn, "MIT: A University's Contribution to National Defense," Military Affairs, October 1979, 120–25.
33. Kármán, The Wind and Beyond, 290; and Gorn, 113–14.
34. Theodore von Kármán, Science: The Key to Air Supremacy, summary vol. to Toward New Horizons: A Report to General of the Army H. H. Arnold, Submitted on Behalf of the A.A.A. Scientific Advisory Group (Wright Field, Dayton, Ohio: Air Materiel Command Publications Branch, Intelligence, T-2, 15 December 1945), 1.3.
35. Kármán, Toward New Horizons, commemorative version (Wright-Patterson AFB, Ohio: Headquarters Air Force Systems Command History Office, 1992), 69–84. Although future attempts were made to repeat the forecast, none made such a monumental impact on the structure or the vision of the US Air Force. Originals of the Kármán report are located in both the Arnold Papers and Spaatz Papers in the Library of Congress as well as at the Air Force Materiel Command Archives at Wright-Patterson AFB, Ohio.
36. Arnold to Spaatz, letter, 6 December 1945, MGC.
37. Sturm, 14–15.
38. Bernard A. Schriever, Washington, D.C., interviewed by author, 10 November 1994.



39. I. B. Holley Jr., Ideas and Weapons: Exploitation of the Aerial Weapon by the United States during World War I: A Study in the Relationship of Technological Advance, Military Doctrine, and the Development of Weapons (1953; new imprint, Washington, D.C.: Office of Air Force History, 1983), 176. The following discussion is based on Dr. Holley's summary and conclusions.


40. Ibid., 177.

Contributor

Dik Alan Daso (USAFA; MA, PhD, University of South Carolina) is curator of modern military aircraft at the Smithsonian Institution, National Air and Space Museum, Washington, D.C. During his career in the Air Force, his assignments included T-38 pilot instructor training at Randolph AFB, Texas; RF-4C instructor pilot; F-15 pilot; and chief of Air Force doctrine at Headquarters Air Force, the Pentagon. Daso is the author of Architects of American Air Supremacy: Gen. Hap Arnold and Dr. Theodore von Kármán (Maxwell AFB, Ala.: Air University Press, 1997) and Hap Arnold and the Evolution of American Airpower (Washington, D.C.: Smithsonian Institution Press, 2000), winner of the American Institute of Aeronautics and Astronautics Manuscript Award in 2001.

Disclaimer

The conclusions and opinions expressed in this document are those of the author cultivated in the freedom of expression, academic environment of Air University. They do not reflect the official position of the U.S. Government, Department of Defense, the United States Air Force or the Air University.



RS Electrogravitic References

(source: http://www.padrak.com/ine/RS_REF1)

"This file contains an electrogravitics reference list, copied ad hoc from various other files and sources, with commentary by yours truly.

Prepared by: Robert Stirnimann
robert@skylink.net

This Update: March 1, 1996

[Email address updated: Dec. 23, 1997].

Danger Will Robinson! Some of the following information is serious, and some is nonsense. Some of the things that might at first seem to be nonsense, are not. And some things referenced below, which come from serious credentialed scientists, are in fact nonsense. Whatever the case, it's been included. Good luck sorting it out.

Understanding gravity is a matter of time.

----- Internet Sites

Elektromagnum web site by David Jonsson:
<http://nucleus.ibg.uu.se/elektromagnum>

KeelyNet:
<http://www protree.com/KeelyNet/>

Los Alamos National Lab Physics E-Print Archive: <http://xxx.lanl.gov/>

Center for Gravitational Physics and Geometry: <http://vishnu.nirvana.phys.psu.edu/>

Bill Beaty's Weird Science, Anomalous Physics, Free-Energy, Tesla Society:
<http://www.eskimo.com/~billb/>

The Institute For New Energy, Patrick Bailey, homepage : <http://www.padrak.com/ine/>

Digital Equipment Corp's Alta Vista web search engine.

If you can't find it with this, it ain't out there yet. <http://www.altavista.digital.com/>

Elsevier Science. Search or browse the table of contents of more than 900 science and technology journals. Data since early 1995.
<http://www.elsevier.nl/cas/estoc/>

Norman Redington's website, The Net Advance of Physics, recent preprints and papers describing new developments in physics: <http://pobox.com/~redingtn>

Embry-Riddle Aeronautical University's Aerospace Virtual Library:
http://macwww.db.erau.edu/www_virtual_lib/aerospace.html

Jack R. Hunt Memorial Library (aerospace):
<http://amelia.db.erau.edu/>

American Institute of Aeronautics & Astronautics (AIAA) home page: <http://www-leland.stanford.edu/group/aiaa/national>

NASA Langley Research Center Library:
<http://blearg.larc.nasa.gov/library/larc-lib.html>

NASA Scientific and Technical Information:
<http://www.sti.nasa.gov/STI-homepage.html>

University of Alabama at Huntsville. Dr Ning Li and Dr Douglas Torr. Microgravity research consultants to NASA's Marshall Space Center.
<http://isl-garnet.uah.edu/RR93/uahmatsci.html>


The Microgravity Research Experiments (MICREX) Data Base
<http://samson2.msfc.nasa.gov/fame/exp/kaw-sl3.html>

Interstellar Propulsion Society:
<http://www.digimark.net/ips/>

National Science Foundation World Wide Web Server.
 Find out where your science tax dollars are going. <http://stis.nsf.gov/>

Nexus magazine web page:
<http://www.peg.apc.org/~nexus/>

Home page of New Scientist magazine:



<http://www.newscientist.com/pstourist/index.html>

The Farce of Physics:

<http://www.germany.eu.net/books/farce>

The World Wide Web Virtual Library:
Sumeria/Technology

<http://lablinks.com/sumeria/tech.html>

The Society for the Advancement of
Autodynamics website:

<http://www.webcom.com/~saa>

Popular Mechanics' Tech Update Article
Archive: <http://popularmechanics.com/cgi-bin/wais.pl>

Fortean web site:

[http://www.clas.ufl.edu/anthro/fortpages.html.](http://www.clas.ufl.edu/anthro/fortpages.html)

Homepage of Apeiron Magazine:

<http://montreal.aei.ca:80/~apeiron/>

Borderland Sciences Research Foundation ftp
site: <ftp://northcoast.com/pub/bsrf>

Homepage of the International Society of
Unified Science, for advancing the Reciprocal
System Theory of Dewey B. Larson:
<http://infox.eunet.cz/interpres/sr/isus/index.html>

Frank Lofaro's homepage, including alternative
science links, and two articles by Whittaker
written in 1903 and 1904 about scalar field
theory and free energy:

<http://www.unlv.edu/~ftlofaro/>

Homepage of the Oppositely Charged Twin
Monopole (OCTM) theory of matter, "Gravity
is a Push", US patent number 5,377,936:
<http://www.epicom.com/gravitypush>

Dr Eujin Jeong's Dipole Theory of Gravity
homepage: <http://www.realtime.net/~ejeong/>

Levesque's (laurient@ee.umanitoba.ca) web
site: <http://www.ee.umanitoba.ca/~laurient>

UFOs and the New Physics:

<http://www.hia.com/hia/pcr/ufo.html>

There is a fairly large body of evidence which
supports the idea of a strong relationship, and
possibly an equivalent fundamental source, for
electromagnetism and gravitation. Many
references to this effect are contained in this
resource list. But for now, let's forget about the
experimental evidence and theoretical ideas
which are presented here, and begin with first
principles.

What if our knowledge of physics had evolved
differently?

What if no one had ever given a thought to any
theory of gravitation, before we discovered the
principles and theories of electromagnetics and
the two nuclear forces. We might have
developed some fairly good theories which
unify the "three" forces. We would know that
clumps of matter are held together primarily by
electromagnetic forces. And we would find
experimentally that if we separate some of
these clumps of matter, a small force continues
to exist which tries to bring them back together.
Would it seem rational to speculate that this
force is something entirely new and
completely different from electromagnetics?
Would it not be a great foolishness to invent
something new and call it gravity and claim
that it has no relationship with the known
forces, and then write elaborate mathematical
theories which describe it solely as geometry?
Or, would it be more rational to see it as what
it probably is -- a manifestation of the
electromagnetic forces which we already know
to hold matter together?

Could it be that electric charge is a
fundamental thing, and inertial mass is merely
a shadow of something primal, and what we
know as a gravitational field is merely the net
result of other primary fields? Geometrize it if
you find it useful to do so, but please recognize
that defining gravity as geometry lends no
information to the understanding of its cause.

Of all the forces we know, there is none stronger than a paradigm. -- Robert Stirnimann

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9512027 From: kldalton@cs.clemson.edu Date: Tue, 12 Dec 1995 11:30:30 -0500

Author(s): Kenneth Dalton

Journal-ref: Hadronic J. 17 (1994) 483-501 Hypothesis: The electromagnetic field is the source of gravitation. This treatment of gravitation is consistent with the quantum theory of matter, which holds that electric charge (or 'generalized charge') is the most fundamental attribute of matter. Experimental predictions of the theory include: (1) any massive body generates a time-dependent gravitational field; (2) there is a linear correlation between the gravitational red-shift of a stellar source and the energy of cosmic rays emitted by that source, given by $\Delta \nu / \nu_0 = \text{energy (eV)} / 10^{27}$; (3) the maximum energy of cosmic rays is 10^{27} eV; (4) this limit is associated with an infinitely red-shifted stellar object, an "electrostatic black-hole," at the potential $c^2/G^{1/2} = 10^{27}$ volts. Finally, the theory predicts that the gravitational potential near any charged elementary particle is many orders of magnitude greater than the Newtonian value.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9601066 From: Kenichi Horie Date: Sat, 13 Jan 1996 14:41:29 +0900

Geometric Interpretation of Electromagnetism in a Gravitational Theory with Torsion and Spinorial Matter

Author(s): Kenichi Horie (KEK Japan)

Comments: Ph.D. thesis, 98 pages, LaTeX file, ca 276kB Possible geometric frameworks for a unified theory of gravity and electromagnetism are investigated: General relativity is enlarged by allowing for an arbitrary complex linear connection and by constructing an extended spinor derivative based on the complex connection. Thereby the spacetime torsion not only is coupled to the spin of fermions and causes a four-fermion contact interaction, but the non-metric vector-part of torsion is also

related to the electromagnetic potential. However, this long-standing relation is shown to be valid only in a special U(1) gauge, and it is a formal consequence of the underlying extended geometry.

Salem, Kenneth G.

The new gravity : a new force, a new mass, a new acceleration : unifying gravity with light / Kenneth G. Salem. 1st ed. Johnstown, PA : Salem Books, c1994. xiii, 181 p. : ill. ; 22 cm. LC CALL NUMBER: QC794.6.G7 S26 1994 SUBJECTS: Unified field theories. Gravitation. Electromagnetic interactions. ISBN: 0962539813

Green, James A.

Gravitation & the electroform model : from general relativity to unified field theory / by James A. Green. 7th ed. [Wichita, Kan.] : Greenwood Research, c1994. 33 p. : ill. ; 24 cm.

LC CALL NUMBER: QC178 .G68 1994 SUBJECTS: Gravitation. Unified field theories. Astrophysics. "Wichita State University Physics Graduate Seminar, Dec.1993 and Dec. 1994"

Another very interesting research on anti-gravity is done (and still going on) by the Japanese prof. Shinichi SEIKE. He published his findings in the book " The Principles of Ultra Relativity ". For his highly mathematical (no nonsense) book write to:

Shinichi SEIKE
G Research Institute
Box 33
UWAJIMA/Ehime (798)
JAPAN

Patents for anti-gravity devices and systems have been issued to Brown, Hooper, Wallace, and others.

US Patents Awarded to Townsend Brown --

300,311 T.T.Brown	Nov.	15,	1928	A
Method of and an Apparatus or Machine for Producing Force or Motion				
1,974,483 T.T.Brown	Sept.	25,	1934	
Electrostatic Motor				
2,949,550 T.T.Brown	Aug.	16,	1960	
Electrokinetic Apparatus				
3,022,430 T.T.Brown	Feb.	20,	1962	
Electrokinetic Generator				
3,187,206 T.T.Brown	June	1,	1965	
Electrokinetic Apparatus				
3,296,491 T.T.Brown	Jan.	3,	1967	Method and Apparatus for Produc- ing Ions and Electrically-Charged Aerosols
3,518,462 T.T.Brown	June	30,	1970	Fluid Flow Control System

Dr. late William J. Hooper, BA, MA, PhD in Physics was affiliated with the University of California at Berkley, and was Professor Emeritus, when he died in 1971. His works are documented and he gained two U.S. patents for his "ALL-

ELECTRIC MOTIONAL FIELD GENERATOR". He claimed use of the "Motional Electric Field" to produce gravity and anti-gravity for use in SPACECRAFT and AIRCRAFT. Indeed, in U.S. patent #3,610,971 you can see a Flying Saucer diagram is used as an example in Figure 7.

-- James Hartman, CaluNET Future Science Administrator

US Patent #3,610,971. "All Electric Motional Electric Field Generator", Awarded to William Hooper, April 1969

US Patent # 3,656,013. "Apparatus for Generating Motional Electric Field", Awarded to William Hooper, April 1972

Hooper, W. J. (1974). New Horizons in Electric, Magnetic and Gravitational Field Theory, Electrodynamics Gravity, Inc. 1969

Frances G. Gibson, "THE ALL-ELECTRIC FIELD GENERATOR AND ITS POTENTIAL", Electrodynamics Gravity, Inc., 1983

"Electric Propulsion Study", Dr. Dennis Cravens, SAIC Corp, prepared for USAF Astronautics Lab at Edwards AFB, August 1990 -- Section 3.7 Non-Inductive Coils


Several authors have suggested that $v \times B$ term in the Lorentz expression should be called into question. Several unverified experimental results have ever been made. An experiment is suggested to test one or several of these theoretical views. This is an area where the experimental procedure is workable and the outcome could have direct results in the area of inertia forces.

During the late 60's William J. Hooper put forth an interesting theory involving the $v \times B$ terms dynamic electrical circuits. There was and is uncertainty as to the exact physical understanding of the Biot-Savart-Lorentz law and Ampere's law involving the set of reaction forces. Peter Graneau has studied these expressions. Hoopers view was that there are three different types of electric fields due to the distribution of electric field, and two due to induction.

At the heart of the issue is the connection of the magnetic field and its source in the charged particles. EM theory is presently consistent with the idea that spinning magnetic dipoles create effects indistinguishable from charged particles. There has been no critical experiment which can disprove whether a magnetic flux rotates with its source. If it does co-move with its source then it is logical to assume that a motional electric field in a fixed reference frame of the current induces a magnetic field. This concept is likewise consistent with a field-free interpretation such as Ampere's original laws.

(with 4 pages more about Hooper's theories)

FREE FALL OF ELEMENTARY PARTICLES: ON MOVING BODIES AND THEIR ELECTROMAGNETIC FORCES, by Nils Rognerud 1994 (nils@ccnet.com) (available at the elektromagnum web site) This paper is a review of the problem of the



observable action of gravitational forces on charged particles. The author discusses the induced electric fields and the sometimes overlooked unique physical properties. He analyzes several experiments, showing the reality of the induced electric fields. The current interpretation, based on the idea of only one electric field, with certain characteristics, is compared with alternative approaches.

The Hooper Coil:

The author has tested a setup by pulsing strong currents, opposite and equal, through multiple parallel conductors. The configuration of the conductors in this type of experiment will cancel the B-fields, while still producing an Em field, in accordance with Eq. 4.2. This is similar to an experiment by Hooper (W. J. Hooper), who successfully predicted and measured the motional electric field - all in zero resultant B-field.

Interestingly, all of the above experiments can influence an electron with a zero B-field, in the region of the electron. This has some profound implications - one of which is that the motional electric force field is immune to electrostatic or magnetic shielding. Experimentally, it can be confirmed that the motional electric field is immune to shielding and follows the boundary conditions of the magnetic (not electric) field. The only way to shield a motional electric field is to use a magnetic shield around the source of the magnetic flux - containing it at the source. These effects are not startling if one remembers that the motional electric field is a magnetic effect and that a magnetic field has a different boundary condition than the electric field.

US Patent #3626605 -- "Method and Apparatus for Generating a Secondary Gravitational Force Field"

Awarded to Henry Wm Wallace of Ardmore PA Dec 14, 1971

US Patent #3626606 -- "Method and Apparatus for Generating a Dynamic Force Field"

Awarded to Henry Wm Wallace of Ardmore PA Dec 14, 1971

US Patent #3823570 -- "Heat Pump" (based on technology similar to the above two inventions)

Awarded to Henry Wm Wallace of Freeport NY July 16, 1973

Gravity is a PUSH!

United States Patent Number 5,377,936
NET KINETIC ENERGY DIFFERENTIAL
GUIDANCE AND PROPULSION SYSTEM
FOR SATELLITES AND SPACE VEHICLES

In the early 1960s, Erwin Saxl conducted a series of experiments which seemed to illustrate a non-zero coupling between EM and gravitational fields. He claimed to see a change in the period of a torque pendulum when its electric potential was raised.


US Patent # 3357253 -- "Device and Method for Measuring Gravitational and Other Forces", awarded to E.J. Saxl, December 1967

"An Electrically Charged Torque Pendulum", by E.J. Saxl, Nature 203, Page 136, July 11 1963.

US patent number #5,076,971.

Barker places radioactive elements inside the sphere of a Van de Graaff generator, runs it at a negative potential for several minutes/hours/days -- and finds that the rate of radioactive decay is extremely enhanced -- with some relationship to the magnitude of the negative potential.

The principal investigator undertook a series of experiments to test the "Barker effect" and the "Keller Catalytic Process" in changing the rate of radioactive decay of heavy elements (elements heavier than lead, such as radium, thorium, or uranium, all of which are radioactive). Barker claims that subjecting radioactive materials to high electrostatic potentials (50,000 volts to 500,000 volts) can increase or decrease the rate of radioactive decay, with short exposures of the high voltage



capable of inducing erratic decay rates which slowly return to normal over a period of weeks. Keller claims that subjecting radioactive materials to the high heat and fusing reaction of a chemical process (Keller Catalytic Process) can eliminate the radioactivity completely.

-- Michael Mandeville
<http://www.aa.net/~mwm/dexmrad1.html>

Carr, Otis (1959). "Amusement Device," (i.e. A Flying Saucer), US Patent No. 2,912,244.

Otis Carr's work involved counter-rotating charged discs that supposedly produced thrust when they reached a certain speed in relation to the earth's rotational speed and became activated by free energy from space. Maybe he did have something." -- James E. Cox

Carr's work is similar in some respects to Hooper's inventions. In both cases, an anti-gravitational effect is reported to result from equal and opposite electric currents. Furthermore, one of Hooper's embodiments, the pancake coil, has an uncanny resemblance to the gravitational shielding experiments which were recently conducted in Tampere Finland (1992 and 1995). Except that in the Tampere experiments, the equal and opposite current is generated in a superconductor disk by way of the Meissner effect. Will we soon begin to recognize value of the discoveries that Carr made nearly 40 years ago, and Hooper made over 25 years ago? -- Robert Stirniman

EXPERIMENTAL RESULTS OF HOOPER'S GRAVITY-ELECTROMAGNETIC COUPLING CONCEPT National Aeronautics and Space Administration. Lewis Research Center, Cleveland, OH. MILLIS, MARC G. WILLIAMSON, GARY SCOTT JUN. 1995 12 PAGES Presented at the 31st Joint Propulsion Conference and Exhibit, San Diego CA, 10-12 Jul. 1995; sponsored by AIAA, ASME, SAE, and ASEE NASA-TM-106963 E-9719 NAS 1.15:106963 AIAA PAPER 95-2601 Avail: CASI HC A03/MF A01 Experiments were conducted to test assertions

from Patent 3,610,971, by W.J. Hooper that self-cancelling electromagnetic coils can reduce the weight of objects placed underneath. No weight changes were observed within the detectability of the instrumentation. More careful examination of the patent and other reports from Hooper led to the conclusion that Hooper may have misinterpreted thermal effects as his 'Motional Field' effects. There is a possibility that the claimed effects are below the detection thresholds of the instrumentation used for these tests. CASI Accession Number: N95-28893

I have two problems with the methodology used by the NASA scientists in the above experiment.

First -- The amount of ampere-turns used in the NASA experiment was substantially lower than the amount used by Hooper. Hooper found that his effect increased in proportion to the square of the current. If you were motivated to verify that the Hooper effect exists, would you not try to conduct the experiment with MORE current, rather than less? Second -- NASA conducted it's tests by energizing the coils and making measurements in an immediate on-off mode, rather than letting things run for a while as Hooper did. NASA's reason for doing this was to avoid errors due to thermal effects. This makes sense. But what does not make sense is that if you are trying to verify an original experiment and you make changes, you have an obligation to also conduct the experiment in it's original mode. To do otherwise is bad science. But what could be wrong with testing things in an immediate on-off mode? Well, it can be seen in other experiments that a gravitational effect sometimes results from macroscopic spin alignment of the quantum angular momentum of a large number of microscopic particles. It has been demonstrated in other experiments that it takes time for these particles to come into alignment. For example in the inventions of Henry Wallace it sometimes took minutes for the "kinemassic" gravito-magnetic field to fully manifest itself. The reason that it takes time for particles to come into alignment, could be much the same reason that it takes

time to permanently magnetize a magnet. Wallace found that the "kinemassic" effect occurs with elemental materials which have a component of unpaired spin in the atomic nucleus. This includes all common isotopes of copper, which of course is the material used in Hooper's coils.

Incidently, NASA essentially has an economic monopoly in the lucrative market for microgravity materials research. -- Robert Stirnimann

The Hooper effect can be readily demonstrated in the "Two Moving

Magnets Experiment". In this experiment, magnetic flux is provided by equal strength opposite pole magnets, moving uniformly in opposite directions. The induced motional electric field that is generated in a conductor, is found to be twice that which would result from a single magnet, while remarkably, the sum of the magnetic B field is zero. This experiment is easy to setup and verify in any electronics laboratory with a pair of magnets, a wire, and a voltmeter. In fact, you may wrap the conductor, in electrostatic or magnetic shielding, and find the same result.

-- Nils Rognerud

Oleg Jefimenko, "Causality, Electromagnetic Induction, and Gravitation", Electret Scientific, Star City, (1992)

Oleg Jefimenko, "Force Exerted on a Stationary Charge by a Moving Electric Current or a Moving Magnet", American Journal of Physics, Vol 61, pages 218-222 (1993)

Apparently, there are some VERY interesting clues to the nature of the universe that are related to the phenomenon of SPIN. It might get very interesting if someone were to make a project of assembling in one place all the information that has been observed, alleged, suspected, or speculated about concerning unexpected effects related to spin, along with

all the traditional Newtonian results, stir, add some seasoning, and see what comes out. For example, in quantum mechanics, if you want to measure the spin axis of an electron, you do an experiment in which you ASSUME an axis, make a measurement of the correlation (the dot product) of that axis with the actual axis of spin for that electron, and theory says you can determine at least how close your guess was. It was a major surprise for the first experimenters with this to find that the guess was always right: whatever spin axis you assume turns out to be correct, exactly dead accurate. You must be a VERY good guesser. Out of this experimental result came the concept of "isospin". Which in itself is kind of weird in that objects with zero radius can still exhibit spinx. But I find the idea that the spin is wherever you guess it might be to be even weirder and to need a better model that predicts this result. -- John Sangster

Paper: gr-qc/9311036

From: jaegukim@cc.kangwon.ac.kr

Date: Tue, 30 Nov 93 13:47:52 +0900

Gravitational Field of a Moving Spinning Point Particle, by Jaegu Kim, 7 pages,

The gravitational and electromagnetic fields of a moving charged spinning point particle are obtained in the Lorentz covariant form by transforming the Kerr-Newman solution in Boyer-Lindquist coordinates to the one in the coordinate system which resembles the isotropic coordinates and then covariantizing it. It is shown that the general relativistic proper time at the location of the particle is the same as the special relativistic one and the gravitational and electromagnetic self forces vanish.

Jaegu Kim, "Gravitational Field of a Moving Point Particle", Journal of the Korean Physical Society, Vol 27 No 5, Oct 94, Pages 484-492

Jaegu Kim, "Gravitational Field of a Moving Spinning Point Particle", Journal of the Korean Physical Society, Vol 27 No 5, Oct 94, Pages 479-483

In the above papers, Dr. Kim derives solutions for the Einstein-Maxwell equations for: a charged massless point particle, a point particle having mass but no charge, a point particle having mass and charge, a massless point particle with charge and spin, and finally -- a point particle having charge, mass, and spin. He determines that there is a region of space around a charged spinning mass in which the gravitational force is negative.

The ability to generate a negative gravity effect may come as no surprise to experimenters who have worked with Bose-Einstein condensates, superfluids, or superconductor material in which the angular momentum of quantum level particles can become aligned along a "macroscopic" spin axis. And it is probably also not a surprise to those who have looked at devices such as the inventions of Henry Wallace, in which a macroscopic body is mechanically spun at high speed in order to cause a "kinemassic" gravito-magnetic field due to spin alignment of the nucleus of elemental materials having an odd number of nucleons (un-paired spin).

Paper: GR-QC/9504023

Date: Mon, 17 Apr 1995 10:43:50 +0900

Title: Pure spin-connection formulation of gravity and classification of energy-momentum tensors

Author: Mathias PILLIN Report-no: YITP/U-95-12

It is shown how the different irreducibility classes of the energy-momentum tensor allow for a pure spin-connection formulation. Ambiguities in this formulation especially concerning the need for constraints are clarified.

From: R.Bursill@sheffield.ac.uk (R Bursill)
Subject: Hi Tc SC and gravitational shielding
Date: Fri, 6 Oct 1995 03:14:41 GMT

Is anyone familiar with the experiments in Tampere Finland, by Podkletnov et al on weak gravitational shielding from a Meissner


levitating, rotating disk of high-T_c superconducting material? The paper is: E. Podkletnov and R. Nieminen, Physica C 203 (1992) 441. E. Podkletnov and A. D. Levit have another paper now, a Tampere University of Technology report, January 1995 (Finland), the experiment having being repeated (I assume no one believed it the first time?).

In the 1st experiment a 5 g sample of silicon dioxide was found to loose around 0.05 % of its weight when placed at a distance of 15 mm from the SC disk. The SC disk had diameter 145 mm and thickness 6 mm. Under rotation of the disk the effect increased up to 0.3 %. In the 2nd experiment samples of different composition and weight (10-50 g) were placed at distances of 25 mm to 1.5 m from the disk. The mass loss went as high as around 2 %. I found out about this through a theoretical preprint by Giovanni Modanese, a Von Humboldt Fellow from the Max Plank institute. The preprint no. is MPI-PhT/95-44, May 1995. A colleague got it from hep-th@babbage.sissa.it, paper 9505094. Modanese thinks that it is something to do with the bose condensate from the SC interacting with the gravitational field. He uses some non-perturbative quantum theory on the Regge lattice to attempt to understand the effect. Must be a little bit like explaining cold fusion with the standard tools - couldn't be done. We all know what happened to cold fusion but at the time a professor from my department said in a public lecture that the product of the believability and the potential importance if true was of order 1.

- Robert Bursill

E. Podkletnov and R. Nieminen, "A Possibility of Gravitational Force Shielding by Bulk YBa₂Cu₃O_{7-x} Superconductor", Physica C 203 (1992) pp 441-444.

E. Podkletnov and A.D. Levi, "Gravitational Shielding Properties of Composite Bulk YBa₂Cu₃O_{7-x} Superconductor Below 70 C Under Electro-Magnetic Field", Tampere University of Technology report MSU-95 chem, January 1995.



HEP-TH/9505094

Theoretical analysis of a reported weak gravitational shielding effect Author: G. Modanese (Max-Planck-Institut, Munich) Report-no: MPI-PhT/95-44 May 1995 Under special conditions (Meissner-effect levitation and rapid rotation) a disk of high-Tc superconducting material has recently been found to produce a weak shielding of the gravitational field. We show that this phenomenon has no explanation in the standard gravity theories, except possibly in the non-perturbative quantum theory on the Regge lattice. More data, and independent repetitions of the experiment are however necessary.

ABSTRACT SUPR-CON/9601001

From: Modanese Giovanni Date: Wed, 17 Jan 1996 21:54:45 +0100 (MET) Updating the analysis of Tampere's weak gravitational shielding experiment Author: Giovanni Modanese

Report-no: UTF-367/96

The most recent data about the weak gravitational shielding produced in Tampere by Podkletnov and coworkers through a levitating and rotating HTC superconducting disk show a very weak dependence of the shielding value ($\sim 1\%$) on the height above the disk. We show that whilst this behaviour is incompatible with an intuitive vectorial picture of the shielding, it is consistently explained by our theoretical model. The expulsive force observed at the border of the shielded zone is due to energy conservation.

NASA is conducting experiments similar to the anti-gravity shielding experiments done in Tampere Finland. A scientist named Ning Li at the University of Alabama Huntsville, is reported to be consulting with NASA. She has written some interesting articles about the relationship between superconductors and gravtiation. Here are references to some of her published articles, and a few related items:

AUTHOR(s): Li, Ning and Torr, D.G.

TITLE(s) Effects of a Gravitomagnetic Field on pure superconductors
In: Phys. Rev. D, JAN 15 1993 v 43 n 2 Page 457

AUTHOR(s): Torr, Douglas G. Li, Ning
TITLE(s): Gravitoelectric-Electric Coupling via Superconductivity.
In: Foundations of physics letters.
AUG 01 1993 v 6 n 4 Page 371

AUTHOR(s): Li, Ning and Torr, D.G.
TITLE(s): Gravitational effects on the magnetic attenuation of superconductors.
In: Physical review. b, condensed matter. SEP 01 1992 v 46 n 9 Page 5489

AUTHOR(s): Peng, Huei
TITLE(s): A New Approach to Studying Local Gravitomagnetic Effects on a Superconductor.
In: General relativity and gravitation.
JUN 01 1990 v 22 n 6 Page 609

AUTHOR(s): Mashhoon, Bahram Paik, Jung Ho Will, Clifford M.
TITLE(s): Detection of the gravitomagnetic field using an orbiting superconducting gravity gradiometer. Theoretical principles. In: Physical review. D, Particles and fields. MAY 15 1989 v 39 n 10 Page 2825

I haven't had the opportunity to read the articles by Drs. Li and Torr, but I am told that in one of her articles, Dr Li provides the following interesting comment --

" a detectable gravitomagnetic field, and in the presence of a time-dependent applied magnetic vector potential field, a detectable gravitoelectric field could be produced"

There is also some information about Dr Ning Li at: <http://isl-garnet.uah.edu/RR93/uahmatsci.html>

Dr Li is with the Applied Materials Lab at the University of Alabama at Huntsville. She



works closely with Dr Douglas Torr. One of their primary interests is development and production of exotic materials in a microgravity environment -- a peculiar coincidence, or maybe not, with the writing of physical theories about how to produce anti-gravity in the laboratory.

Here's an unusual article from the website. -----

Can gravity be 'made' in the laboratory?

A theory that might lead to the creation of measurable manmade gravitational fields has been developed by physicists at UAH.

If the theoretical work is borne out in the laboratory, it will prove that physicist Albert Einstein was correct in predicting that moving matter generates two kinds of gravitational fields: gravito-magnetic and gravito-electric. The 'artificial' gravitational field would be generated inside a container made of a superconducting material, said Dr. Douglas Torr, a research professor of physics and director of UAH's Optical Aeronomy Laboratory. "I think we can at the very least generate a microscopic field ..." If Einstein was right, the amount of gravito-magnetic energy produced by an object is proportional to its mass and its movement, explained Dr. Ning Li, a research scientist in UAH's Center for Space Plasma and Aeronomics Research. To create the artificial gravitational fields, Torr and Li propose placing a superconducting container in a magnetic field to align ions that are spinning or rotating in tiny circles inside the superconducting material. Their theory predicts the existence of ionic spin or rotation in a superconductor in a magnetic field.

There are persistent rumors among UFO-buffs that NASA already has an operating microgravity chamber, located in Houston TX and/or Huntsville AL. One person, Robert Oechsler, reports that he has personally been inside NASA's antigrav chamber. But, that's another story. For more info, see the books "Alien Contact" and "Alien Update" by Timothy Good.

Paper: hep-th/9412243

From: Vu.Ho@sci.monash.edu.au

Date: Sat, 31 Dec 1994 17:06:38 +1100


Title: Gravity as a coupling of two

electromagnetic fields Author: Vu B Ho

A discussion on a possibility to represent gravity as a coupling of two equal and opposite electrogmanetic fields. Classically the existence of equal and opposite electromagnetic fields can be ignored altogether. However, the problem can be viewed differently if we want to take into account possible quantum effects. We know that in quantum mechanics the potentials themselves may be significant and they may determine the dynamics of a particle in a region where the fields vanish. (Aharonov and Bohm 1959, Peshkin and Tonomura 1983)

AN EXPERIMENT TO TEST THE GRAVITATIONAL AHARONOV-BOHM EFFECT Ho, Vu B. Morgan, Michael J. Monash University, Clayton, Victoria, Australia 1994 8 PAGES, Australian Journal of Physics (ISSN 0004-9506) vol. 47, no. 3 1994 p. 245-252 HTN-95-92507 The gravitational Aharonov-Bohm (AB) effect is examined in the weak-field approximation to general relativity. In analogy with the electromagnetic AB effect, we find that a gravitoelectromagnetic 4-vector potential gives rise to interference effects. A matter wave interferometry experiment, based on a modification of the gravity-induced quantum interference experiment of Colella, Overhauser and Werner (COW), is proposed to explicitly test the gravitoelectric version of the AB effect in a uniform gravitational field. CASI Accession Number: A95-87327

I recommend you get a copy of Aharonov and Bohm's classic paper "Significance of Electromagnetic Potentials in the Quantum Theory" published in The Physical Review in 1959. One of the important things that Aharonov and Bohm did was to demonstrate that the electromagnetic potentials are richer in properties than the Maxwell fields. The field is



an artificial mathematical construct from which emerges the whole idea of a continuum. When you can wean yourself of this intellectual crutch you will be ready to do real physics. Both GR and QM are addicted to the same falsehood.

-- Charles Cagle

In the Aharonov-Bohm effect it has been determined theoretically and experimentally that there is a measurable effect on a charged particle due to the electromagnetic vector potential. Which of course would be no surprise, except that the effect occurs even in areas of space where the value of the classical electromagnetic fields vanish. A quantum phase shift, detectable via particle interferometry, is found to occur due to the magnetic vector potential A . The effect on a charged particle occurs in regions which are completely shielded from classical electromagnetic fields.

A dual of the Aharonov-Bohm effect is the Aharonov-Casher effect, where it is shown that measurable effects of spin-precession of a particle's magnetic moment can occur due to the electric potential, even in areas of space where the classical electrical field is completely absent.

Prior to the revolutionary paper by Aharonov and Bohm in 1959, the importance of the electromagnetic potential and related interferometry effects, was suggested in articles by Edmund Whittaker in 1903 and 1904. And, what is now known as the Aharonov-Bohm effect, was explicitly identified in an earlier paper on electron optics by Ehrenberg and Siday in 1949.

E.T. Whittaker, "On the partial differential equations of mathematical physics," *Mathematische Annalen*, Vol 57, 1903, pages 333-355. In this paper Whittaker demonstrates that all scalar EM potentials have an internal, organized, bidirectional EM plane-wave structure. Thus there exists an electromagnetics that is totally internal to the scalar EM potential. Since vacuum/spacetime

is scalar potential, then this internal EM is in fact "internal" to the local potentialized vacuum/ spacetime.

-- Tom Bearden


E.T. Whittaker, "On an expression of the electromagnetic field due to electrons by means of two scalar potential functions," *Proceedings of the London Mathematical Society*, Series 2, Vol 1, 1904, pages 367-372. In this paper Whittaker shows that all of classical electromagnetics can be replaced by scalar potential interferometry. This ignored paper anticipated the Aharonov-Bohm (AB) effect by 55 years, and drastically extended it as well. Indeed, it prescribes a macroscopic AB effect that is distance-independent, providing a direct and engineerable mechanism for action-at-a-distance. It also provides a testable hidden-variable theory that predicts drastically new and novel effects. -- Tom Bearden

W. Ehrenberg and R. W. Siday, Proc. Phys. Soc. London, B62, 8 (1949) Ten years earlier than Aharonov and Bohm, Ehrenberg and Siday formulated the science of electron optics by defining the electron refractive-index as a function of electromagnetic potential. Near the end of their paper, they discuss "a curious effect", which is exactly the AB effect. On the two sides of a magnetic flux, the vector potential has different values. This means a different refractive index for two geometrically equivalent paths. This difference in refractive index would cause an observable phase shift.

-- Jun Liu

Y. Aharonov and D. Bohm, "Significance of Electromagnetic Potentials in the Quantum Theory," *Physical Review*, Second Series, Vol 115 no 3, pages 485-491 (1959)

Effects of potentials on charged particles exist even in the region where all the fields (and therefore the forces on the particles) vanish, contrary to classical electrodynamics. The quantum effects are due to the phenomenon of interference. These effects occur in spite of Faraday shielding. The Lorentz force does not appear anywhere in the fundamental quantum



theory, but appears only as an approximation that holds in the classical limit. In QM, the fundamental physical entities are the potentials, while the fields are derived from them by differentiation.

Herman Erlichson, "Aharonov-Bohm Effect and Quantum Effects on Charged Particles in Field-Free Regions," American Journal of Physics, Vol 38 No 2, Pages 162-173 (1970).

M. Danos, "Bohm-Aharonov effect. The quantum mechanics of the electrical transformer," American Journal of Physics, Vol 50 No 1, pgs 64-66 (1982).

Bertram Schwarzschild, "Currents in normal-metal rings exhibit Aharonov-Bohm Effect," Physics Today, Vol 39 No 1, pages 17-20 (Jan 1986)

S. Olariu and I. Iovitzu Popescu, "The quantum effects of electromagnetic fluxes," Reviews of Modern Physics, Vol 57 No2, April 1985.

Yoseph Imry and Richard Webb, "Quantum Interference and the Aharonov- Bohm Effect", Scientific American, April 1989, pages 56-62

E. Merzbacher, "Single Valuedness of Wave Functions", American Journal of Physics, Vol 30 No 4, pages 237-247 (April 1962)

Yoseph Imry, "The Physics of Mesoscopic Systems", Directions in Condensed Matter Physics, World Scientific Publishing (1986)

Richard Webb and Sean Washburn, "Quantum Interference Fluctuations in Disordered Metals", Physics Today, Vol 41 No 12 pages 46-53, Dec 1989

"STAR WARS NOW! The Bohm-Aharonov Effect, Scalar Interferometry, and Soviet Weaponization" By T. E. Bearden, Tesla Book Company

Peshkin M. and Lipkin H.J. "Topology, Locality, and Aharonov-Bohm Effect with

Neutrons" Physical review letters APR 10 1995 v 74 n 15

Yakir Aharonov and Ady Stern, "Origin of the geometric forces accompanying Berry's geometric potentials", Physical Review letters. DEC 21 1992 v 69 n 25 Page 3593


Yakir Aharonov, Jeeva Anandan, and Sandu Popescu, "Superpositions of time evolutions of a quantum system and a quantum time-translation machine." Physical review letters. JUN 18 1990 v 64 n 25 Page 2965

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9506038 From: "Jun Liu" Date: Sun, 25 Jun 1995 03:25:05 -0400

Potential Effect: Aharonov-Bohm Effect of Simply Connected Region Author: Jun Liu
Comments: Prediction of a new effect. Numerical estimate given for experimental verification. The referees disagree with each other on the existence of this effect.

We study a generalization of Aharonov-Bohm effect, the potential effect. The discussion is focused on field-free effects in simply connected region, which obviously can not have any local field-flux. Among the published discussions about this kind of effects, it is generally agreed that this kind of effect does not exist due to gauge invariance. However, there are also opinions that this effect is a trivial variation of Aharonov-Bohm effect and therefore there is no need to check its existence. To my knowledge, it has neverbeen tested. My first goal here is to supply enough theoretical reason to motivate the experimental test of this effect. I start with an intuitive derivation, then I introduce a wave-front theory as a theoretical consideration. Logically, the existence of potential effect implies the existence of the AB effect, but not vice versa. The purpose of this paper is to provide a physical connection in the opposite direction.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9510004 From: "Jun Liu" Date: Thu, 5 Oct 1995 04:30:27 -0400



The Real Significance of the Electromagnetic Potentials Author(s): J'un L'iu

The importance of the potential is revealed in a newly discovered effect of the potential. This paper explores the same issue introduced in quant-ph/9506038 from several different aspects including electron optics and relativity. Some people fail to recognize this effect due to a wrong application of gauge invariance.

In the above two papers, Dr Liu proposes a theory of the electromagnetic potential which is a radical extension of the well known Aharonov-Bohm effect. In the second paper he is barely able to contain his frustration about repeated publication rejections over the last four years from leading physics journals. He provides a theoretical foundation for his potential theory, as well as some relatively straight forward suggestions for experiments which might confirm the theory. But there is an enormous problem. Liu's theory violates the concept of invariance of physical parameters under an electromagnetic gauge transformation. Electromagnetic gauge invariance is a cornerstone in the foundation of quantum theory and QED, and it is also part and parcel linked with the dogma of light speed invariance. In other words, heresy.

The AB effect is invariant under an electromagnetic gauge transformation. While a phase-shift occurs in the AB effect, it can be identified only over a closed path and is impossible to identify with any specific "local" region of space. Furthermore, in the AB effect, there is no interaction relating to a transfer of energy or momentum. Maintaining the idea of gauge invariance is a little harder to do in the Aharonov-Casher effect, but it can be accomplished by "gauging away" the physical effects of magnetic spin precession by using a combination of factors from the classical Maxwell fields along with the electromagnetic potential. It has the look of an elaborate parlor trick, but so does most of QED.


Liu's theory predicts that the electromagnetic potential acts like a kind of "refractive index" to wave propagation, and is similar in some

respects to what was predicted in the earlier paper on electron optics by Ehrenberg and Siday in 1949. The result is that in some circumstances an electromagnetic potential causes a change in wavelength, and in other circumstances causes a change in phase (AB effect). An effect on wavelength would be manifested as a change in the envelope of the interference pattern, rather than merely a shift in the pattern. In Liu's theory an exchange of energy and momentum becomes possible. His theory is relatively easy to test and verify, but oddly or not, no one has yet done so. Maybe because we already "know" it can't be true?

One interesting prediction of Liu's theory is that electromagnetic potential will result in time dilation. He doesn't appear to be aware that there is already experimental evidence that this occurs. See references to inventions and experiments by people such as Saxl, Barker, and Keller, which demonstrate time dilation in an electric potential. Time dilation can be viewed equivalently as a shift in wavelength. Liu wishes for someone to conduct an experiment to test for a change in wavelength by using a quantum interferometer. A fine idea. But what about those experimenters who have already measured this effect with a clock? Also see a variety of references here to theories and experiments which relate the scalar electric potential to the gravitational field, and time dilation is a well known, and experimentally verified, prediction of general relativity.

The Aharonov-Bohm effect has sparked a revolution in physical thought. There are a variety of new ideas and experiments, such as verification of Liu's theory, which could soon begin to fan it to a flame. When the flame becomes sufficiently illuminating, watch the political scientists begin to scramble for a comfortable seat nearer the fire. -- Robert Stirnimann

Over the last five years, there have been over 300 papers published about various aspects of Aharonov-Bohm and Aharonov-Casher



effects, and quantum interferometry. The subject relates to nearly all aspects of modern physics. Here are selected examples:

AUTHOR(s): Semon, Mark D.
TITLE(s): The Aharonov-Bohm Effect: Still a Thought-Provoking Experiment.
In: Foundations of physics.
JUL 01 1988 v 18 n 7 Page 731

AUTHOR(s): Furuya, Kazuhito
TITLE(s): Transient Response of the Aharonov-Bohm Effect.
In: Japanese journal of applied physics. part 1, FEB 01 1989 v 28 n 2 Page 303

AUTHOR(s): Chetouani, L. Guechi, L. Hammann, T.F.
TITLE(s): Exact path integral solution of the coulomb plus Aharonov-Bohm potential.
In: Journal of mathematical physics.
MAR 01 1989 v 30 n 3 Page 655

AUTHOR(s): Lee, Patrick A.
TITLE(s): Gauge field, Aharonov-Bohm Flux, and high-Tc superconductivity.
In: Physical review letters.
AUG 07 1989 v 63 n 6 Page 680

AUTHOR(s): Bezerra, V.B.
TITLE(s): Gravitational analogs of the Aharonov-Bohm effect.
In: Journal of mathematical physics.
DEC 01 1989 v 30 n 12 Page 2895

AUTHOR(s): Reznik, B. Aharonov, Y.
TITLE(s): Question of the nonlocality of the Aharonov-Casher effect.
In: Physical review. D, Particles and fields.
DEC 15 1989 v 40 n 12 Page 4178

AUTHOR(s): Stovicek, P.
TITLE(s): The Green function for the two-solenoid Aharonov-Bohm effect.
In: Physics letters: [part A]
NOV 27 1989 v 142 n 1 Page 5

AUTHOR(s): Ellis, J.R.
TITLE(s): Dirac magnetic monopole and the Aharonov-Bohm solenoid in the Poincare gauge.

In: Journal of physics A: Mathematical and general. JAN 07 1990 v 23 n 1 Page 65

AUTHOR(s): Gerber, A. Deutscher, G.
TITLE(s): AC-to-DC conversion and Aharonov-Bohm effect in percolating superconducting films.
In: Physical review letters.
MAR 26 1990 v 64 n 13 Page 1585

AUTHOR(s): Hagen, C.R.
TITLE(s): Exact equivalence of spin-1/2 Aharonov-Bohm and Aharonov-Casher effects.
In: Physical review letters.
MAY 14 1990 v 64 n 20 Page 2347


AUTHOR(s): Afanase'ev, G.N.
TITLE(s): Old and new problems in the theory of the Aharonov-Bohm effect.
In: Soviet journal of particles and nuclei. JAN 01 1990 v 21 n 1 Page 74

AUTHOR(s): Silverman, M.P.
TITLE(s): Two-solenoid Aharonov-Bohm experiment with correlated particles.
In: Physics letters: [part A]
AUG 13 1990 v 148 n 3/4 Page 154

AUTHOR(s): Gornicki, Pawel
TITLE(s): Aharonov-Bohm Effect Vacuum Polarization.
In: Annals of physics.
SEP 01 1990 v 202 n 2 Page 271

AUTHOR(s): Gal'tsov, D.V.
Voropaev, S.A.
TITLE(s): Bremsstrahlung polarization in the Aharonov-Bohm effect.
In: Moscow University physics bulletin.
1990 v 45 n 1 Page 8

AUTHOR(s): Padmanabhan, T.
TITLE(s): Vacuum polarization around an Aharonov-Bohm solenoid.



- In: Pramana.
MAR 01 1991 v 36 n 3 Page 253
- AUTHOR(s): Hagen, C.R.
TITLE(s): Spin dependence of the Aharonov-Bohm Effect.
In: International journal of modern physics A. JUL 30 1991 v 6 n 18 Page 3119
- AUTHOR(s): Dupuis, Nicolas Montambaux, Gilles
TITLE(s): Aharonov-Bohm flux and statistics of energy levels in metals.
In: Physical review B: Condensed matter. JUN 15 1991 v 43 n 18 Page 14390
- AUTHOR(s): Ortiz, M.E.
TITLE(s): Gravitational anyons, Chern-Simons-Witten gravity and the gravitational Aharonov-Bohm effect.
In: Nuclear physics. b. SEP 30 1991 v 363 n 1 Page 185
- AUTHOR(s): Bezerra, V.B.
TITLE(s): Gravitational Aharonov-Bohm effect in a locally flat spacetime.
In: Classical and quantum gravity. OCT 01 1991 v 8 n 10 Page 1939
- AUTHOR(s): Sitenko, Y.A.
TITLE(s): The Aharonov-Bohm effect and the inducing of vacuum charge by a singular magnetic string.
In: Nuclear physics. b. MAR 23 1992 v 372 n 3 Page 622
- AUTHOR(s): March-Russell, John Preskill, John Wilczek, Frank
TITLE(s): Internal frame dragging and a global analog of the Aharonov-Bohm effect.
In: Physical review letters. APR 27 1992 v 68 n 17 Page 2567
- AUTHOR(s): Krive, I.V. Rozhavsky, A.S.
TITLE(s): Non-Traditional Aharonov-Bohm Effects in Condensed Matter.
In: International journal of modern physics. B. MAY 10 1992 v 6 n 9 Page 1255
- AUTHOR(s): Krive, I. V. Zvyagin, A. A.
TITLE(s): Aharonov-casher effect in half-integer spin antiferromagnets.
In: Modern physics letters. B, Condensed matter ph JUN 20 1992 v 6 n 14 Page 871
- AUTHOR(s): Zubkov, M.A. Polikarpov, M.I.
TITLE(s): Aharonov-Bohm effect in lattice field theory.
In: JETP letters. APR 25 1993 v 57 n 8 Page 461
- AUTHOR(s): Duru, I. H.
TITLE(s): Casimir Force Between Two Aharonov-Bohm Solenoids.
In: Foundations of physics. MAY 01 1993 v 23 n 5 Page 809
- AUTHOR(s): Takai, Daisuke Ohta, Kuniichi
TITLE(s): Aharonov-Bohm effect in the presence of magnetic flux and electrostatic potential.
In: Physical review. b, condensed matter. JUL 15 1993 v 48 n 3 Page 1537
- AUTHOR(s): Allman, B.E. Cimmino, A. Klein, A.G.
TITLE(s): Observation of the scalar Aharonov-Bohm effect by neutron interferometry.
In: Physical review. A. SEP 01 1993 v 48 n 3 Page 1799
- AUTHOR(s): Jensen, Bjorn Kucera, Jaromir
TITLE(s): On a gravitational Aharonov-Bohm effect.
In: Journal of mathematical physics. NOV 01 1993 v 34 n 11 Page 4975
- AUTHOR(s): Maeda, J. Shizuya, K.
TITLE(s): Aharonov-Bohm and Aharonov-Casher effects and electromagnetic angular momentum.
In: Zeitschrift fur Physik C; particles and fields. 1993 v 60 n 2 Page 265
- AUTHOR(s): Afanasiev, G.N.



- TITLE(s): Toroidal solenoids in an electromagnetic field and toroidal Aharonov-Casher effect.
In: *Physica scripta*. OCT 01 1993 v 48 n 4 Page 385
- AUTHOR(s): Moreau, William Ross, Dennis K.
TITLE(s): Complementary electric Aharonov-Bohm effect.
In: *Physical review. A, Atomic, molecular, and opt* JUN 01 1994 v 49 n 6 Page 4348
- AUTHOR(s): Ho, Vu B. Morgan, Michael J.
TITLE(s): An Experiment to Test the Gravitational Aharonov-Bohm Effect.
In: *Australian journal of physics*. 1994 v 47 n 3 Page: 245
- AUTHOR(s): Zeiske, K. Zinner, G. Helmcke, J.
TITLE(s): Atom interferometry in a static electric field: Measurement of the Aharonov-Casher phase.
In: *Applied physics. b, lasers and optics*. FEB 01 1995 v 60 n 2/3 Page: 205
- AUTHOR(s): Sazonov, S.N.
TITLE(s): On Aharonov-Bohm Effect in Multiconnected Superconductor.
In: *Acta physica Polonica, A.* DEC 01 1994 v 86 n 6 Page 987
- AUTHOR(s): Reznik, B.
TITLE(s): Gravitational analogue of the Aharonov-Casher effect.
In: *Physical review d: particles, fields, gravitat* MAR 15 1995 v 51 n 6 Page 3108
- AUTHOR(s): Oh, Sangchul Ryu, Chang-Mo
TITLE(s): Persistent spin currents induced by the Aharonov-Casher effect in mesoscopic rings.
In: *Physical review B: Condensed matter*. MAY 15 1995 v 51 n 19 Page 13441
- AUTHOR(s): Leadbeater, M. Lambert, C.J.
TITLE(s): Mesoscopic Superconducting Analogs of the Aharonov-Bohm-Casher Effect.
- In: *Physical review letters*. MAY 29 1995 v 74 n 22 Page 4519
- AUTHOR(s): Cook, Richard J. Fearn, Heidi Milonni, Peter W.
TITLE(s): Fizeau's experiment and the Aharonov-Bohm effect.
In: *American journal of physics*. AUG 01 1995 v 63 n 8 Page 705
- AUTHOR(s): Yi, J. Jeon, G. S. Choi, M. Y.
TITLE(s): Dual Aharonov-Casher effect and persistent dipole current.
In: *Physical review B: Condensed matter*. SEP 15 1995 v 52 n 11 Page 7838
- AUTHOR(s): Audretsch, Jurgen Jasper, Ulf Skarzhinsky, Vladimir D.
TITLE(s): Bremsstrahlung of relativistic electrons in the Aharonov-Bohm potential.
In: *Physical review d: particles, fields, gravitat* FEB 15 1996 v 53 n 4 Page 2178
- AUTHOR(s): Skarzhinsky, Vladimir D. Audretsch, Jurgen Jasper, Ulf
TITLE(s): Electron-positron pair production in the Aharonov-Bohm potential.
In: *Physical review d: particles, fields, gravitat* FEB 15 1996 v 53 n 4 Page 2190
-
- Time out for a summary.
- . Hooper, as well as Carr, Rognerud, Jefimenko, et al, find that a electromagnetic effect which is not shieldable, and hence difficult to distinguish from gravitation, results from equal and opposite electric currents (dipole-current), and that a similar effect can also be generated by a moving magnet or a moving electric current.
- . Recent experiments in Tampere Finland, discover a gravitational shielding effect from a levitated rotating superconductor disk. This is similar in some respects to Hooper's invention, with the equal-and-opposite electric current being generated in a superconductor disk via the Meissner effect.



. Sansbury, Volkov, Brown, Teller, Blackett, Zollner, et al, provide theoretical arguments as well as some experimental indications that equal-and-

opposite electric charge (dipole-charge) is similar, or equivalent, to a static gravitational field. And that alignment of electric dipoles in matter and in vacuum polarization, can result in a force which is not shieldable, and not easily distinguishable from gravity. Conversely, it is well known that a gravitational field, an acceleration, or a mechanical force, causes a dipole moment (polarization) to occur within a dielectric material.

. Wallace, Laithwaite, Barnett, et al, discover that gravitational and electromagnetic field effects occur due to alignment of the microscopic spin of quantum particles with the angular momentum spin axis of a larger macroscopic body.

. Aharonov and Bohm discover that an effect can occur on an electrically charged particle due to the magnetic vector potential, in regions of space where the classic Maxwell fields vanish. Originally -- on the outside of infinitely long solenoid coil (with the magnetic field cancelled by equal-and-opposite currents). Others have conducted this experiment using a toroidal coil coated with superconductor material (generating an equal-and-opposite current) to cause the Maxwell magnetic field to vanish. A similar effect, Aharonov-Casher is discovered to occur due to the electric scalar potential, in regions of space where the Maxwell electric field vanishes.

. Whittaker, and Eherenberg and Siday, have written theories which are precursors to Aharonov-Bohm, suggesting that the electromagnetic potential is a far richer and more fundamental thing than the Maxwell fields. The classical Maxwell fields are regarded as artificial abstractions. We can also note that Maxwell's theory itself, was originally much richer in variables (20 equations and 20 unknowns), before it was simplified by Gibbs and Heaviside, to the

vector formulation which we know as "Maxwell's" equations.

. Vu Ho authors a recent paper suggesting experiments relating the electromagnetic potential and the Aharonov-Bohm effect to gravitation. And in a more recent paper, using the mathematics of differential geometry and general relativity, Dr Ho demonstrates that gravity can be expressed mathematically as a coupling of two equal-and-opposite electromagnetic fields.

. Jun Liu authors recent papers suggesting that the electromagnetic potential is of paramount importance. Liu's theory predicts that "local" effects can result from the potential in regions where the Maxwell fields vanish -- a violation of the theory of invariance under electric gauge transformations. Liu theory predicts that time dilation will occur in an electric potential. Saxl, Barker, and Keller have conducted earlier experiments which demonstrate time dilation in an electric potential.

Ning Li, a consulting scientist to NASA's Marshall Space Center, who we might presume to know something, authors papers about the relationship of gravito-electric and gravitomagnetic forces to the electromagnetic potential, and methods for generation of gravitational effects with superconductor material. According to Dr Li -- "a detectable gravitomagnetic field, and in the presence of a time-dependent applied magnetic vector potential field, a detectable gravitoelectric field could be produced."

How many clues do we need? Equal-and-opposite electric sources (dipole-charges and/or dipole-currents) appear to effect the electromagnetic potential in ways which are indistinguishable from gravitation. And you know what they say about things that look like a duck.

The net sum of equal and opposite electromagnetic vectors is a zero vector, but it is NOT the same situation as no vector. For skeptics and diehards who are still having a hard time accepting the idea of electro-



gravitics, here's a simple experiment. Stand on a train track between two locomotives which are pushing on you with equal force in opposite directions. You will exhibit no net motion. None the less, you may soon begin to notice that something important is happening.

- Robert Stirnimann

Jorge Pullin Wed, 1 Feb 1995 22:55:17 -0500
(EST)

Matters of Gravity, a newsletter for the gravity community Author: Jorge Pullin (PSU), editor.

Loops, knots, gauge theories and quantum gravity Rodolfo Gambini and Jorge Pullin ; foreword by Abhay Ashtekar. New York: Cambridge University Press, 1996. Cambridge monographs on mathematical physics ISBN 0-521-47332-2 (hc)

A number of reports which have been prepared for the USAF are publicly available. These reports can be obtained from the "Defense Technical Information Center" (DTIC). Cameron Station, Alexandria VA 22304, 800-225-3842

Cravens D.L., "Electric Propulsion Study", Prepared for the Astronautics Laboratory, Air Force Space Technology Center, at Edwards AFB. August 1990. AL-TR-89-040

Mead F.B. Jr, et al, Advanced Propulsion Concepts - Project Outgrowth, AFRPL-TR-72-31, (JUN 1972).

Mead F.B. Jr, "Exotic Concepts for Future Propulsion and Space Travel", In Advanced Propulsion Concepts, 1989 JPM Specialist Session, (JANNAF) Chemical Propulsion Information Agency, CPIA Publication 528, p.93-99, (May 24, 1989).

Talley R.L, "Twenty First Century Propulsion Concept", Veritay Technology Inc, East Amherst NY. Prepared for the Phillips Laboratory, Air Force Systems Command,

Propulsion Directorate, Edwards AFB. May 1991. PL-TR-91-3009


Talley R.L., 21st Century Propulsion Concept, AFAL-TR-88-031, Apr 88.

Talley R.L., Final report on NYS contract no. (88)-166 of NYS Science and Technology Foundation with Veritay Technology, Inc., P.O. Box 305, East Amherst NY 14051.

Forward R.L., 21st Century Space Propulsion Study, AL-TR-90-030, Final Report on Contract FO4611-87-C-0029, Air Force Astronautics Lab (AFSC), (Oct 1990). --AND-- Forward,R.L., 21st Century Space Propulsion Study (Addendum), PL-TR-91-3022, Final (Addendum), OLAC Phillips Lab, formally known as Air Force Astronautics Lab (AFSC), (June 1991).

Electric Propulsion Study by Dennis L. Cravens:

TABLE OF CONTENTS	
Page	
PREFACE	1
I. INTRODUCTION	2
1.1 Background and Theoretical Developments	4
1.2 Measurement	4
6.1.3 Force Fields	8
1.4 Chirality - Odd Number Space-Like Dimensions ..	11
II. THEORIES	13
2.1 Introduction	13
2.2 General Framework of Theory	14
2.2.1 Born - Infield	17
2.2.2 Lande'	19
2.2.3 Podolsky	20
2.2.4 Corben	21
2.2.5 Flint	21
2.2.6 Ingraham	21
2.2.7 Arctan Potential	23
2.2.8 Milne	24
2.2.9 Williams	25



2.3 Development of 5-D EM Equations	27	3.10 Speed of Light in a Mass Flow	103
2.3.1 Modifications to Maxwell's Equations	33	3.11 Charged Torque Pendulum	105
2.3.2 Lorentz Forces in 5-D	36	3.12 Thermoelectric/Gravitational Effects	107
2.3.3 Wave Propagation in 5-Space	38	3.13 Binary Pulsar	107
2.3.4 Limits to Conversion Rates	40	3.14 Proton Scattering	107
2.3.5 Reduction to Newton's Laws - PPN	41	3.15 Inertial Mass Variation	107
2.3.6 Thermoelectric Potentials in Gravity Field.	43	3.16 An Improper Experiment	108
2.3.7 Field Vectors and Equations in 5-D	44		
2.4 Conservation Laws	47	IV. CONCLUSIONS AND RECOMMENDATIONS	110
2.4.1 Conservation of Energy	48		
2.4.2 Conservation of Linear Momentum	50		
2.4.3 Conservation of Angular Momentum	51		
2.4.4 Conservation of Parity	53		
2.4.5 Conservation of Pseudovectors	54		
2.4.6 Conditions for Non-Conservations	58		
2.5 Vacuum Fluctuations	60		
2.6 Quantum Considerations	62		
2.7 Compatibility of 10-D String Theories	68		
2.8 Mach's Principle	69		
2.9 Rosen's Bi-Metric Theory	72		
2.10 Non-Conservation	74		
2.11 Particles in 5-D Spaces	76		
III. EXPERIMENTS			
3.1 Approach to Selection of Experiments	78	AUTHOR(s): Zhang, Xiao-He	
3.2 Radiation Pressure	80	TITLE(s): Interactions of magnetohydrodynamic waves with gravitomagnetic fields, and their possible roles in black-hole magnetospheres.	
3.3 Biefeld-Brown Effects	83	In: Physical review. D, Particles and fields.	
3.4 Conductive Submarine	88	DEC 15 1989 v 40 n 12	
3.5 Gravitational Rotor	89	Page: 3858	
3.6 Spin Aligned Nuclei -- Magnetic and Rotational Alignment	90	AUTHOR(s): Khanna, Ramon Camenzind, Max	
3.7 Non-Inductive Coils	94	TITLE(s): The Gravitomagnetic Dynamo Effect in Accretion Disks of Rotating Black Holes.	
3.8 EM Transparency of Conductive Media	100		
3.9 Magnetic Loop	101		

In: The Astrophysical journal.
NOV 10 1994 v 435 n 2 p 2

AUTHOR(s): Casotto, S. Ciufolini, I. Vespe, F.

TITLE(s): Earth satellites and gravitomagnetic field.

In: Il nuovo cimento delle societa italiana di fisic MAY 01 1990 v 105 n 5 Page 589

AUTHOR(s): Mashhoon, Bahram Paik, Jung Ho Will, Clifford M.

TITLE(s): Detection of the gravitomagnetic field using an orbiting superconducting gravity gradiometer. Theoretical principles. In: Physical review. D, Particles and fields. MAY 15 1989 v 39 n 10 Page 2825

AUTHOR(s): Nordtvedt, K.

TITLE(s): Gravitomagnetic interaction and laser ranging to Earth satellites.

In: Physical review letters.
DEC 05 1988 v 61 n 23 Page 2647

There is a reprint of an article that appeared in "Interavia, Volume XI - No. 5, 1956" a March 23, 1956 article titled "Towards Flight without Stress or Strain... or Weight" This article has a photograph of T.T.Brown holding one of his flying disks, and another photograph of the flying disk by itself. There is some info on the operation of the electrokinetic apparatus.

The 1956 paper "The Gravitics Situation" (prepared by Gravity Rand Ltd., a division of Aviation Studies Ltd. This includes six appendices with papers by various authors including the text from T. Townsend Brown's 1929 gravitor patent.

Many documents on Gravitoelectrics/Electrogravitation refer back to the 1952 Project Winterhaven. That project is said to contain information on a Mach 3


Combat Disc. Also, have any records related to other Projects with Mr. T.T. Brown been produced. I have seen his Lab notes 1 - 3 - 4. I was looking for 2 - 5 & 6. Also, the Bahnson et al Brown lab notes during his research days at Bahnson Labs in North Carolina 1957-60 period or about. I have a poor chopped up Lab Video on the subject. I'm looking for the full video the 45 minute one. Mine is a mere 23 minutes.

I have yet to track down an original document entitled: "The Flying Saucer: The Application of the Biefeld-Brown Effect to the Solution of Space Navigation" by Mason Rose. This 50's document details how a flying saucer operates. I have a copy of a re-write and it is outstanding. And I'm also looking for a document as seen on SIGHTINGS TV entitled: "PROJECT SILVER BUG" the 1955 USAF Flying Saucer Tests. Also, seeking a copy of PROJECT WINTERHAVEN by Thomas Townsend Brown on a MACH-3 Combat Disc. The British had a stake in as well as the USAF. It is from the 1950's.

-- James Hartman, CaluNET - Future Science Admin.

The Biefeld-Brown (spelling is correct) effect is described generally as the anomalous tendency of high voltage flat capacitors to display movement towards (usually) the positive pole. Effects are most often seen at potentials above 50kv. Thomas Townsend Brown held a few patents on devices using it. It's very controversial and is part of the subject of "electrogravitics", as some say that the BB effect is actually polar gravity peeking out from behind a high electrical gradient within a dielectric. Claims are that the mass of the dielectric is a factor in the magnitude of the effect as well as the capacitance and the gradient intensity. Should be fairly easy for the home-

workshop experimenter to get a look at, but the difficulty seems to be in isolating the effect from ionic wind and simple electrostatic propulsive effects. Skeptics claim that those forces are all it ever was, but a few reports indicate that they may be wrong.



-- Rick Monteverde, Honolulu HI

The experiments involved freely suspended electrically charged capacitors, which were determined to possess angular momentum yet did not rotate. Source: Albert Einstein: Philosopher- Scientist, P. Schilpp, editor, 3rd ed., 1988, pp 522-523.

Schilpp, Paul Arthur, 1897- ed.

Albert Einstein: philosopher-scientist. [3d ed.] La Salle, Ill., Open Court [1970] xviii, 781 p. illus., facsim., ports. 25 cm. LC CALL NUMBER: QC16.E5 S3 1970

>From Richard Feynman's Lectures on Physics we learn that there is intrinsic field energy and momentum density associated with a static electro-

magnetic field configuration. When there is a change in the magnetic field, this field energy and momentum can be directly converted into kinetic energy and mechanical momentum. Feynman illustrates this with an electromagnetic carousel paradox. In this paradox, a dielectric disk (which is embedded with small charged spheres along its circumference) rotates without any apparent "counter" torque in the system. Before this rotation occurs, the dielectric disk is immersed in a static magnetic field. The subsequent rotation occurs as a consequence to reducing the previously static magnetic field to zero. The angular momentum and rotational kinetic energy comes directly from the initial static magnetic field.

"The Feynman Lectures on Physics" by Richard Feynman, R.B. Leighton, and M. Sands, Volume II p 17-6


A Report on the T. Townsend Brown Conference: "Focus on Unconventional Energies: A Symposium on Electrical Propulsion & the Technology of Electro-Gravity"

April 15-16, 1994 Philadelphia Community College, Philadelphia, PA

This conference was held in tribute to Thomas Townsend Brown and I feel that it was a great success. About 15 speakers and 80 attendees provided a brief overview of Zero Point Energy theories, Free Energy devices, electrostatics theory, and antigravity experiments and documentation. Attendees came from as far away as California and Washington. The conference program advertised the following topics: "A Review of Advanced Energy Devices: Evidence, Promises, and Dangers" by Patrick Bailey (VP INE); "Thomas Townsend Brown's Electro-Gravities Research in the 1950's" by Tom Valone (Integrity Institute); "The Role of Electro-Statics" by Charles Yost (Electric Spacecraft Journal); "Thomas Townsend Brown's Research: A Challenge to Modern Science" by Elizabeth Rauscher (Tecnic Research Laboratories); "Electro-Gravitic Theory: Explaining the Operating Principle of Brown's Electric Disks" by Paul LaViolette (The Starburst Foundation); "A Panel Discussion on Biefeld-Brown and Beyond;" "Vortices in the Zero Point Energy" by Moray King; "Design of a Compact Marx Generator Triggered by a Blumlein Capacitor" by George Hathaway; "Thomas Townsend Brown's Final Gravito-Electric Research" by Josh Reynolds (New Wave Partners); "Townsend Brown Effects Reviewed" by Ron Kovac; "Pushing the Boundaries: Electro-Hydro Dynamic Potentials ..." by Henry Monteith, and "Gravity Drop Tests" by Don Kelly (SEA). - Patrick Bailey

I have the audio tapes from the T.T. Brown conference, 11 tapes in all, and I got a lot of good information from it. - Bob Reim (reim@advantor.com)

There is a connection between Townsend Brown and UFOs. Brown was the founder of NICAP (National Investigations Committee on Aerial Phenomena) Project Skylight, and Brown served as Vice Chairman pro tempore during NICAP's organizational period in 1956.



Partial biography of Thomas Townsend Brown: 1922-23, private research laboratory, Pasadena, California; 1924-25, special electronics research, Denison University, Department of Physics; 1926-30, private research laboratory (astrophysics), Zanesville, Ohio, in collaboration with Dr. Paul Biefeld, Swazey Observatory, Granville, Ohio; 1930-33, Naval Research Laboratory (radiation and spectroscopy), Washington, DC; ... 1938, Assistant Engineering Officer (Lt. jg USNR) shakedown cruise USS NASHVILLE to Europe; 1939-40, Materials and Processes Engineer (aircraft), Glenn L. Martin Company, Baltimore; 1940-41, Officer-in-charge (Lt. USNR), Magnetic and Acoustic Minesweeping, Research and Development, Bureau of Ships, Navy Department, Washington, D. C.; 1942-43, Officer-in-charge (Lt. Comdr. USNR), Atlantic Fleet Radar Materiel School and Gyro- compass School, Naval Operating Base, Norfolk, Virginia; 1944-45, Radar Consultant, Advanced Design Section, Lockheed Aircraft Corporation, Burbank, California; ...

Also, there was a T.T. Brown on the Condon committee for UFO studies. And some of Brown's above described Navy duties are coincident with some of the times and places in stories about the Montauk Project/Philadelphia Experiment.

Quotation from a letter to William Moore from T. Townsend Brown dated 12/17/76 --

"I am still working on petroelectricity and the project is housed largely at Stanford Research Institute with additional assistance being provided by the University of California - Berkeley and the Ames Research center of NASA. Unfortunately, under the circumstances, while this project is being evaluated for funding by ERDA we should not and cannot publish details..."

"Your next question concerns the airfoils. As far as I am aware, no rf is radiated. There is, of course, a static d.c. field which accompanies the airfoils in flight."

It is very interesting to note that Townsend Brown was the pioneer in this field, and was not able to obtain very much support for his work until the 1950's. During that time, there was much discussion of gravity and antigravity within the aerospace industry and in the magazine "Aviation Week." Then the Gravity Research Group (GRG) published a detailed summary report of their review of research into "Electrostatic Motion, Dynamic Counterbary, and Barycentric Control" (i.e. "Antigravity"). This report is the last public report that any researchers have been able to find for us that deals with the physical effects of electrostatics, electrodynamics, and gravity control. (It is also worth noting that this report was found in the Wright Patterson Air Force Base Library "TL 565 A9" and was not listed in the library catalog). So, after the mid-1950's to the present, no other information regarding the technology of electrodynamics and its effect on gravity has been able to be found in any of the un-classified U.S. literature.

- Patrick Bailey

I have the FIVE (5) lab books of TT Brown's R&D at the Bahnsen Co. in Salem, N. Carolina 1958-9. I also have some other letters and drawings of the lab plus the only surviving 16mm colour film of the various stages of his work at Bahnsen Labs.

I was in contact with Dr Brown in 1983 by phone and by mail. He died of lung cancer not long after in Oct of 1985. He told me that a lot of people including Bill Morre had attributed more to his work than he had really done. In particular, he was only marginally connected with the Philadelphia Experiment as such. His main theme of R&D was dielectrics and the Biefeld-

Brown effect. He was not an electromagnetics man... only electrostatics.

>From 1983 to 1991 or so I was in frequent communication with J. Frank King who was TT Brown's boss at the Bahnsen Co. J. Frank was a good man and a good friend of mine. He, too, died in Dec 1989. Before he died I was given rights to reproduce and share letters, files, drawings, patent submissions, films etc

from his personal files on TT Brown, George Adamski, Dr Ilka, T Henry Moray and others. J Frank warned me a long time ago to take what TT Brown said with a 'grain of salt' because Townsend had a habit of 'stretching the truth' a bit to get funding which he was always in need of.... So, I warn you now in good faith: If you seek lost or hidden technology in Brown's lab notes, I don't think you will find it there; however, I am prepared to make photocopies available to you.

There are about 750 pages in all. I would need to charge you AUS\$50 per notebook which would include the air mail charges as well. In US\$ that would be about US\$38 per notebook. The film is available as are the notebooks (I think) from The Electric Spacecraft Journal in the US (Charles Yost on 704-

252-8083, FAX 202-683-3511. -- Stan Deyo

As far as I know, the last thing Brown published before his death was, "On The Possibilities of Optical-Frequency Gravitational Radiation", 2/14/1976 and 8/30/1976. I don't know where it was originally published. But you can get copy from:

Rex Research, P.O. Box 19250, Jean NV 89019

It is part of NR 046-BT2/B17-BRV "T. Brown: Petro-Voltaics" (Gravito-Electric Conversion). Most people think Brown was just into flying capacitors he was into much much more... -- Bob Paddock

Here are some titles by Townsend Brown:

"The Wizard of Electro-Gravity: The Man Who Discovered how UFOs are powered." by William L. Moore. In UFO Report magazine. Unfortunately the issue date is not on this copy, and the magazine is at work. A lot of the same information can be found in the book "The Philadelphia Experiment: Project Invisibility" by William L. Moore with Charles Berlitz. Chapter 10 "The Force Fields of Townsend Brown". These two items are the

same, I just don't know which one came first. Also there is more than one book with the title "The Philadelphia Experiment". You want the one with ISBN 0-449-20526-6. "The Townsend Brown Electro-Gravity Device: A Comprehensive Evaluation by the Office of NAVAL Research" 15 September 1952. Such as "How I Control Gravity by T. Townsend Brown" from Science and Invention Magazine Aug. 1929. "Townsend Brown and his Anti-Gravity Discs" by Gaston Burridge in Fate Magazine. No issue date is visible. "Electrical Self-Potential in Rocks" by T.Townsend Brown, some time after 1/1976, but again no source is visible. "Another Step Toward Anti-Gravity" by Gaston Burridge in The American Mercury, June 1958, p77.

"Towards Flight without Stress or Strain... or Weight" by Intel, Washington, D.C. [Doesn't make since but that is what it says.] Some one just on the list here just reinvented "The Fluid Pump" by T.Townsend Brown for the Whitehall-Rand Group, Washington DC -----

Paper: gr-qc/9207002

From:

RCAPOVI%CINVESMX.BITNET@ricevm1.rice.edu Date: Tue, 21 Jul 1992 17:52 CST Title: Remarks on Pure Spin Connection Formulations of Gravity Authors: Riccardo Capovilla and Ted Jacobson Abstract: In the derivation of a pure spin connection action functional for gravity two methods have been proposed. The first starts from a first order lagrangian formulation, the second from a hamiltonian formulation. In this note we show that they lead to identical results for the specific cases of pure gravity with or without a cosmological constant.

Paper: hep-th/9210110 (Phys. Rev. D47, R5214 (1993).) From:

pullin@mail.physics.utah.edu (Jorge Pullin) Date: Tue, 20 Oct 92 11:18:14 MDT

QUANTUM EINSTEIN-MAXWELL FIELDS: A UNIFIED VIEWPOINT FROM THE LOOP REPRESENTATION, R. Gambini, J. Pullin, 13pp. no figures. We propose a naive unification of

Electromagnetism and General Relativity based on enlarging the gauge group of Ashtekar's new variables. We construct the connection and loop representations and analyze the space of states. In the loop representation, the wavefunctions depend on two loops, each of them carrying information about both gravitation and electromagnetism. We find that the Chern-Simons form and the Jones Polynomial play a role in the model.

Paper: gr-qc/9301012

From:

porrati@MAFALDA.PHYSICS.NYU.EDU
(Massimo Porrati) Date: Wed, 13 Jan 93
20:17:21 -0500

Massive Spin-5/2 Fields Coupled to Gravity: Tree-Level Unitarity vs. the Equivalence Principle, Massimo Porrati, 6 pages. I show that the gravitational scattering amplitudes of a spin-5/2 field with mass $m \parallel M_{Pl}$ violate tree-level unitarity at energies $\sqrt{s} \approx \sqrt{mM_{Pl}}$ if the coupling to gravity is minimal. Unitarity up to energies $\sqrt{s} \approx M_{Pl}$ is restored by adding a suitable non-minimal term, which gives rise to interactions violating the (strong) equivalence principle. These interactions are only relevant at distances $d \approx 1/m$.

Paper: gr-qc/9303014

From: ISTVAN@RMK520.RMKI.KFKI.HU
Date: Wed, 10 Mar 1993 16:24:01 +0100
(WET) MAXWELL FIELDS IN SPACETIMES ADMITTING NON-NULL KILLING VECTORS, by Istvan Racz, 7 pages, PACS numbers: 04.20.Cv, 04.20.Me, 04.40.+c We consider source-free electromagnetic fields in spacetimes possessing a non-null Killing vector field, ξ^a . We assume further that the electromagnetic field tensor, F_{ab} , is invariant under the action of the isometry group induced by ξ^a . It is proved that whenever the two potentials associated with the electromagnetic field are functionally independent the entire content of Maxwell's equations is equivalent to the relation $n^a T_{ab}=0$. Since this relation is implied by Einstein's equation we argue that it is enough to solve merely Einstein's equation

for these electrovac spacetimes because the relevant equations of motion will be satisfied automatically. It is also shown that for the exceptional case of functionally related potentials $n^a T_{ab}=0$ implies along with one of the relevant equations of motion that the complementary equation concerning the electromagnetic field is satisfied.

Paper: gr-qc/9310007 (Physica Scripta 48, 649 (1993)) From: harald@nordita.dk (Harald H. Soleng) Date: Mon, 4 Oct 93 13:18:04 +0100

INVERSE SQUARE LAW OF GRAVITATION IN (2+1)-DIMENSIONAL SPACE-TIME AS A CONSEQUENCE OF CASIMIR ENERGY, H. H. Soleng, 10 pages, LaTeX, Report: UPR-0540-T, To appear in Physica Scripta. The gravitational effect of vacuum polarization in space exterior to a particle in (2+1)-dimensional Einstein theory is investigated. In the weak field limit this gravitational field corresponds to an inverse square law of gravitational attraction, even though the gravitational mass of the quantum vacuum is negative. The paradox is resolved by considering a particle of finite extension and taking into account the vacuum polarization in its interior.

Paper: gr-qc/9310019

From: rri!bri@rri.ernet.in (B.R.Iyer)

Date: Tue, 12 Oct 93 12:44:52 IST

THE FRENET SERRET DESCRIPTION OF GYROSCOPIC PRECESSION B.R.Iyer and C.V.Vishveshwara , 37 pages, Paper in Latex. The phenomenon of gyroscopic precession is studied within the framework of Frenet-Serret formalism adapted to quasi-Killing trajectories. Its relation to the congruence vorticity is highlighted with particular reference to the irrotational congruence admitted by the stationary, axisymmetric spacetime. General precession formulae are obtained for circular orbits with arbitrary constant angular speeds. By successive reduction, different types of precessions are derived for the Kerr - Schwarzschild - Minkowski spacetime family. The phenomenon is studied in the case of other interesting spacetimes, such as the De Sitter

and Godel universes as well as the general stationary, cylindrical, vacuum spacetimes.

Paper: gr-qc/9310030

From: khatsymovsky Date: Thu, 21 Oct 93 16:39:25 +0100

Can wormholes exist? V.Khatsymovsky, 10 pages, Plain LaTe_X, preprint UUITP-20/1993

Renormalized vacuum expectation values of electromagnetic stress-energy tensor are calculated in the background spherically-symmetrical metric of the wormhole's topology. Covariant geodesic point separation method of regularization is used. Violation of the weak energy condition at the throat of wormhole takes place for geometry sufficiently close to that of infinitely long wormhole of constant radius irrespectively of the detailed form of metric. This is an argument in favour of possibility of existence of selfconsistent wormhole in empty space maintained by vacuum field fluctuations in the wormhole's background.

Paper: hep-th/9402046

From: LANDI@SUHEP.PHY.SYR.EDU

Date: Tue, 08 Feb 1994 15:09:39 -0500 (EST)

GRAVITY AND ELECTROMAGNETISM IN NONCOMMUTATIVE GEOMETRY, Giovanni Landi, Nguyen Ai Viet, Kameshwar C.Wali, 1 + 11 pages, Report # SU-4240-566, We present a unified description of gravity and electromagnetism in the framework of a Z2 noncommutative differential calculus. It can be considered as a "discrete version" of Kaluza-Klein theory, where the fifth continuous dimension is replaced by two discrete points. We derive an action which coincides with the dimensionally reduced one of the ordinary Kaluza-Klein theory.

Paper: gr-qc/9404016

From: David Garfinkle Date: Sun, 10 Apr 1994 17:44:50 -0400

Generating new magnetic universe solutions from old. By David Garfinkle and M.A. Melvin. 17 pages

In this paper we apply the techniques which have been developed over the last few decades for generating nontrivially new solutions of the

Einstein-Maxwell equations from seed solutions for simple spacetimes. The simple seed spacetime which we choose is the "magnetic universe" to which we apply the Ehlers transformation. Three interesting non-singular metrics are generated. Two of these may be described as "rotating magnetic universes" and the third as an "evolving magnetic universe." Each is causally complete - in that all timelike and lightlike geodesics do not end in a finite time or affine parameter. We also give the electromagnetic field in each case. For the two rotating stationary cases we give the projection with respect to a stationary observer of the electromagnetic field into electric and magnetic components.

Paper: gr-qc/9404065 (Phys. Rev. D50 (1994) 6190) From: carroll@marie.mit.edu (Sean Carroll) Date: Sun, 1 May 1994 16:35:00 -0400

Energy-Momentum Restrictions on the Creation of Gott Time Machines, by Sean M. Carroll, Edward Farhi, Alan H. Guth, and Ken D. Olum. Plain Te_X, 41 pages incl. 9 figures. MIT-CTP #2252.

The discovery by Gott of a remarkably simple spacetime with closed timelike curves (CTC's) provides a tool for investigating how the creation of time machines is prevented in classical general relativity. The Gott spacetime contains two infinitely long, parallel cosmic strings, which can equivalently be viewed as point masses in (2+1)-dimensional gravity. We examine the possibility of building such a time machine in an open universe. Specifically, we consider initial data specified on an edgeless, noncompact, spacelike hypersurface, for which the total momentum is timelike (i.e., not the momentum of a Gott spacetime). In contrast to the case of a closed universe (in which Gott pairs, although not CTC's, can be produced from the decay of stationary particles), we find that there is never enough energy for a Gott-like time machine to evolve from the specified data; it is impossible to accelerate two particles to sufficiently high velocity. Thus, the no-CTC theorems of Tipler and Hawking are enforced in an open (2+1)-dimensional universe by a mechanism different from that which operates in a closed universe. In proving our result, we

develop a simple method to understand the inequalities that restrict the result of combining momenta in (2+1)-dimensional gravity.

Paper: gr-qc/9405050

From: MATSAS@IFT.UESP.ANSP.BR

Date: Mon, 23 May 1994 15:01 BSC (-0300 C)
 DO INERTIAL ELECTRIC CHARGES
 RADIATE WITH RESPECT TO
 UNIFORMLY ACCELERATED
 OBSERVERS?, George E.A. Matsas, 6 pages
 (REVTEX 3.0), IFT-P017/94. We revisit the long standing problem of analyzing an inertial electric charge from the point of view of uniformly accelerated observers in the context of semi-classical gravity. We choose a suitable set of accelerated observers with respect to which there is no photon emission coming from the inertial charge. We discuss this result against previous claims [F. Rohrlich, Ann. Phys. (N.Y.) vol: 22, 169 (1963)]. (This Essay was awarded a Honorable Mention for 1994 by the Gravity Research Foundation.)

Paper: gr-qc/9406032

From: wam@tdo-serv.lanl.gov (Warner A. Miller) Date: Mon, 20 Jun 94 14:44:42 MDT
 Spin Dynamics of the LAGEOS Satellite in Support of a Measurement of the Earth's Gravitomagnetism, Salman Habib, Daniel E. Holz, Arkady Kheyfets, Richard A. Matzner, Warner A. Miller and Brian W. Tolman, 16 pages, RevTeX, LA-UR-94-1289. (Part I of II, postscript figures in Part II). LAGEOS is an accurately-tracked, dense spherical satellite covered with 426 retroreflectors. The tracking accuracy is such as to yield a medium term (years to decades) inertial reference frame determined via relatively inexpensive observations. This frame is used as an adjunct to the more difficult and data intensive VLBI absolute frame measurements. There is a substantial secular precession of the satellite's line of nodes consistent with the classical, Newtonian precession due to the non-sphericity of the earth. Ciufolini has suggested the launch of an identical satellite (LAGEOS-3) into an orbit supplementary to that of LAGEOS-1: LAGEOS-3 would then experience an equal and opposite classical precession to that of LAGEOS-1. Besides

providing a more accurate real-time measurement of the earth's length of day and polar wobble, this paired-satellite experiment would provide the first direct measurement of the general relativistic frame-dragging effect. Of the five dominant error sources in this experiment, the largest one involves surface forces on the satellite, and their consequent impact on the orbital nodal precession. The surface forces are a function of the spin dynamics of the satellite. Consequently, we undertake here a theoretical effort to model the spin dynamics of LAGEOS. In this paper we present our preliminary results.

Paper: gr-qc/9407003

From: William Bruckman Date: Tue, 5 Jul 94 09:06:49 EDT

Generation of Electro and Magneto Static Solutions of the Scalar-Tensor Theories of Gravity, William Bruckman, 28 pages, LaTeX. The field equations of the scalar-tensor theories of gravitation are presented in different representations, related to each other by conformal transformations of the metric. One of the representations resembles the Jordan-Brans-Dicke theory, and is the starting point for the generation of exact electrostatic and magnetostatic exterior solutions. The corresponding solutions for each specific theory can be obtained by transforming back to the original canonical representation, and the conversions are given for the theories of Jordan-Brans-Dicke, Barker, Schwinger, and conformally invariant coupling. The electrostatic solutions represent the exterior metrics and fields of configurations where the gravitational and electric equipotential surfaces have the same symmetry. A particular family of electrostatic solutions is developed, which includes as special case the spherically symmetric solutions of the scalar-tensor theories. As expected, they reduce to the well-known Reissner-Nordstrom metric when the scalar field is set equal to a constant. The analysis of the Jordan-Brans-Dicke metric yields an upper bound for the mass-radius ratio of static stars, for a class of interior structures.

Paper: gr-qc/9407030

From: Marco SISSA +39(40)3787522 Date:
 Thu, 21 Jul 1994 15:10:04 +0200
QUANTUM ELECTROMAGNETIC WORMHOLES AND GEOMETRICAL DESCRIPTION OF THE ELECTRIC CHARGE by Marco Cavaglia 13 pages, PLAIN TEX, Report No: SISSA 92/94/A (to appear in Phys. Rev. D15).

I present and discuss a class of solutions of the Wheeler-de Witt equation describing wormholes generated by coupling of gravity to the electromagnetic field for Kantowski-Sachs and Bianchi I spacetimes. Since the electric charge can be viewed as electric lines of force trapped in a finite region of spacetime, these solutions can be interpreted as the quantum corresponding of the Ein-stein--Rosen--Misner--Wheeler electromagnetic geon.

Paper: gr-qc/9409060 (Annals of Physics vol. 240 432-458 (1995)) From: soleng@surya11.cern.ch (Harald SOLENG)
 Date: Thu, 29 Sep 94 14:01:03 +0100

Modification of the Coulomb potential from a Kaluza-Klein model with a Gauss-Bonnet term in the action, by H. H. Soleng and O. Gron, 27 pages, compressed and uuencoded postscript file with unpacking instructions; major revision to section IV.D.2 on pages 15-16 ("Corrections to the Coulomb potential at short distances") and to the figure on page 27, to be published in The Annals of Physics (NY), NORDITA 94/50 In four dimensions a Gauss-Bonnet term in the action corresponds to a total derivative, and it does not contribute to the classical equations of motion. For higher-dimensional geometries this term has the interesting property (shared with other dimensionally continued Euler densities) that when the action is varied with respect to the metric, it gives rise to a symmetric, covariantly conserved tensor of rank two which is a function of the metric and its first and second order derivatives. Here we review the unification of General Relativity and electromagnetism in the classical five-dimensional, restricted (with $g_{55} = 1$) Kaluza-Klein model. Then we discuss the modifications of the Einstein-Maxwell theory that results from adding the Gauss-Bonnet term in the action. The resulting four-

dimensional theory describes a non-linear U(1) gauge theory non-minimally coupled to gravity. For a point charge at rest, we find a perturbative solution for large distances which gives a mass-dependent correction to the Coulomb potential. Near the source we find a power-law solution which seems to cure the short-distance divergency of the Coulomb potential. Possible ways to obtain an experimental upper limit to the coupling of the hypothetical Gauss-Bonnet term are also considered.

Paper: hep-th/9410046

From: M.J. Duff Date: Fri, 7 Oct 94 13:04:15 BST

KALUZA-KLEIN THEORY IN PERSPECTIVE, M. J. Duff, 38 pages latex, NI-94-015 The Kaluza-Klein idea of extra spacetime dimensions continues to pervade current attempts to unify the fundamental forces, but in ways somewhat different from that originally envisaged. We present a modern perspective on the role of internal dimensions in physics, focussing in particular on superstring theory. A novel result is the interpretation of Kaluza-Klein string states as extreme black holes.(Talk delivered at the Oskar Klein Centenary Nobel Symposium, Stockholm, September 19-21, 1994.)

Paper: gr-qc/9509018

From: nunez@venus.fisica.unlp.edu.ar (NUNEZ Carlos) Date: Fri, 8 Sep 95 15:05:13 EST

Title: On Pseudospherically Symmetric Repulsive Gravitational Field Authors: Luis A. Anchordoqui, Graciela S. Birman, Jose D. Edelstein and Carlos Nunez Report-no: La Plata-Th 95/23 The solution of Einstein vacuum equation, for a static pseudospherically symmetric system, is presented. It describes a singular solution that produces a repulsive gravitational field with an event horizon. We analyse particle motion in such a gravitational field and comment on some interesting features of the solution. -----

"The Origin of the Electromagnetic Interaction in Einstein's Unified Field Theory", Antoci,S. General Relativity and Gravitation, Vol 23 No 1, 1991

Recently it has been shown that, if sources are appended in a certain way to the field equations of Einstein's unified theory, the contracted Bianchi identities and the field operations appear endowed with definite physical meaning. The theory looks like a gravoelectrodynamics in a polarizable Riemannian continuum. The wealth of the implied possibilities is far richer than in the so-called Einstein-Maxwell theory.

Paper: HEP-TH/9411092

From: hssong@phyy.snu.ac.kr

Date: Mon, 14 Nov 94 15:19:29 KST

Title: Factorization and polarization in linearized gravity Authors: S.Y. Choi, J.S. Shim, H.S. Song Comments: 45 pages, figures are included (uses pictex), RevTex Report-no: KEK-TH-415, HYUPT-94/10, SNUTP 94-03, We investigate all the four-body graviton interaction processes: $gX \rightarrow \gamma X$, $gX \rightarrow gX$, and $gg \rightarrow gg$, with X as an elementary particle of spin less than two in the context of linearized gravity except the spin-3/2 case. We show explicitly that gravitational gauge invariance and Lorentz invariance cause every four-body graviton scattering amplitude to be factorized. We explore the implications of this factorization property by investigating polarization effects through the covariant density matrix formalism in each four-body graviton scattering process.

Causality, electromagnetic induction, and gravitation : a different approach to the theory of electromagnetic and gravitational fields/ Oleg D. Jefimenko. Star City [West Virginia] : Electret Scientific Co., c1992. xii, 180 p. : LC CALL NUMBER: QC665.E4 J44 1992

SUBJECTS: Electromagnetic fields. Gravitational fields. Causality. Maxwell Equations.

COUNTER-GRAVITATION: The sustaining of an object in space by means of a counter-gravitational effect produced through the action of an electric field upon the object. Associated with the effects of levitation in this manner, is a simultaneous appearance of a strange luminous halo that appears at about 500,000 volts.

Sources: American Philosophical Society, Proceedings. Philadelphia, PA, years 1914-1929. Articles on Charles F. Brush's experiments.

Electrical Experimenter. "Can Electricity Destroy Gravitation?", New York, March 1918.

Electrical Experimenter. "Piggott's Electrogravitation Experiment", Vol. 8, 1920.

Hooper, William J., New Horizons in Electric, Magnetic, and Gravitational Field Theory, Principia College, Elsah, IL, 1974.

The Scientific Papers of James Clerk Maxwell. Vol. II, W.D. Niven (ed.), Constable & Co., London, 1965. "Le Sage Theory of Gravitation".

Transactions of the Academy of Science. "Nipher's Gravitation Experiments", Vol. 23, pp. 163-192+, St. Louis, 1916.

US patent No. 1,006,786, Piggott.
3,518,462, Brown.
3,610,971, Hooper.

"Journal of Propulsion and Power" of the AIAA, R.H. Woodward Waesche, Science Applications International Corporation, Editor in Chief. This Journal is devoted to the advancement of the science and technology of aerospace propulsion and power through the dissemination of original archival papers contributing to advancements in airbreathing, electric, and advanced propulsion; solid and liquid rockets; fuels and propellants; power generation and conversion for aerospace vehicles; and the application of aerospace

science and technology to terrestrial energy devices and systems. It is intended to provide readers of the Journal, with primary interests in propulsion and power, access to papers spanning the range from research through development to applications. Papers in these disciplines and the sciences of combustion, fluid mechanics, and solid mechanics as directly related to propulsion and power are solicited.

Published Bimonthly

AIAA Members \$42.00 per year (\$72.00 outside North America) Institutions \$300.00 per year (\$360.00 outside North America) -----

I have recently come into possession of a paper on magneto-gravitics and field resonance systems, presented by A.C. Holt from NASA Johnson Space Center to the American Institute of Aeronautics and Astronautics' 16th Joint Propulsion Conference, June 30-July 2, 1980. Holt presents a project using an already existing system known as the Coherent Field and Energy Resonance System (CoFERS) [probably located at Los Alamos Labs' High Magnetic Field Research Laboratory]. CoFERS utilizes a toroidal-shaped energy guide with megagauss magnetic field sources located along radius vectors equally spaced around the toroid. CoFERS is shaped like a thick flying disc. Holt goes on to say: "By converting an object's normal space-time energy pattern to an energy pattern which differs substantially from the normal pattern, the gravitational forces acting on the object are changed. The object's new pattern interacts with the surrounding space-time and virtual energy patterns, such that the interactive forces are substantially altered. The alteration of the characteristics of the continuous field of force results in the apparent motion of the object *through space-time*." [...] "Since the gravitational forces acting on the propulsion system can be quickly altered to achieve the desired motion, the *spacecraft* can make right-angle turns at very-high velocities without adversely affecting the crew or system elements. The effective gravitational field the

spacecraft/ aircraft experiences can be nearly simultaneously reoriented at a 90-degree angle, resulting in a smooth continuous motion as far as the occupants are concerned." [...] "The gravimagnetic system is perhaps best suited for use in and around ... a large mass such as the Earth." "While the gravimagnetic system is likely to be the first field-dependent propulsion system developed, the field resonance system will **bring stellar and galactic travel out of the realm of science fiction**. The field resonance system artificially generates an energy pattern which precisely matches or resonates with a virtual pattern associated with a distant space-time point. According to the model, if a fundamental or precise resonance is established, (using hydromagnetic wave fine-tuning techniques), the spacecraft will be very strongly and equally repelled by surrounding virtual patterns. At the same time, through the virtual many-dimensional structure of space-time, a very strong attraction with the virtual pattern of a distant space-time point will exist. ...this combination of very strong forces will result in the translocation of the spacecraft from its initial position through the many-dimensional virtual structure to the distant space-time point. [...] "A space-time 'jump' already appears to be supported by astrophysical research."

Should you wish the entirety of this report, "Prospects for a Breakthrough in Field Dependent Propulsion" by A.C. Holt, you can order it from AeroPlus Dispatch, 1722 Gilbreth Road, Burlingame, CA 94010; phone: (800)-662-AERO. The paper/conference number is AIAA-80-1233 (American Institute of Aeronautics and Astronautics, June 30-July 2, 1980 - 16th Annual Conference. -- Rich Boylan

There is also a great article discussing the work of A.C. Holt in the Electric Spacecraft Journal: Issue No. 5, June 30, 1992. -----

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-

<p>QC/9601024 MONTANARI@axpfe1.fe.infn.it Date: Wed, 17 Jan 1996 13:01:16 +0100 (CET) Coherent Interaction of a Monochromatic Gravitational Wave with both Matter and Electromagnetic Circuits Author(s): Enrico Montanari (1) , Pierluigi Fortini (1) ((1) University of Ferrara,INFN sezione di Ferrara, Italy) The interaction of a gravitational wave with a system made of an RLC circuit forming one end of a mechanical harmonic oscillator is investigated. We show that, in some configurations, the coherent interaction of the wave with both the mechanical oscillator and the RLC circuit gives rise to a mechanical quality factor increase of the electromagnetic signal. When this system is used as an amplifier of gravitational periodic signals a sensitivity of \$10^{-30}\$ on the amplitude of the metric could be achieved.</p> <p>GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9602004</p> <p>From: wells@cfaitamp2.harvard.edu (Jack Wells) Date: Thu, 1 Feb 1996 16:50:06 -0500 Gravitational Interaction of Spinning Bodies, Center-of-Mass Coordinate and Radiation of Compact Binary Systems Author(s): I.B. Khriplovich, A.A. Pomeransky Spin-orbit and spin-spin effects in the gravitational interaction are treated in a close analogy with the fine and hyperfine interactions in atoms. The proper definition of the center-of-mass coordinate is discussed. The technique developed is applied then to the gravitational radiation of compact binary stars. Our result for the spin-orbit correction differs from that obtained by other authors. New effects possible for the motion of a spinning particle in a gravitational field are pointed out. The corresponding corrections, nonlinear in spin, are in principle of the same order of magnitude as the ordinary spin-spin interaction.</p> <p>HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9601119</p> <p>From: gonzalez@fyma.ucl.ac.be</p>	<p>From:</p> <p>Date: Tue, 23 Jan 1996 10:03:41 +0100 (MET) Spinning Relativistic Particle in an External Electromagnetic Field Author(s): M. Chaichian , R. Gonzalez Felipe , D. Louis Martinez The Hamiltonian formulation of the motion of a spinning relativistic particle in an external electromagnetic field is considered. The approach is based on the introduction of new coordinates and their conjugated momenta to describe the spin degrees of freedom together with an appropriate set of constraints in the Dirac formulation. For particles with gyromagnetic ratio $g=2$, the equations of motion do not predict any deviation from the standard Lorentz force, while for $g \neq 2$ an additional force, which corresponds to the magnetic dipole force, is obtained.</p> <p>HIGH ENERGY PHYSICS - PHENOMENOLOGY, ABSTRACT HEP-PH/9601280</p> <p>From: MAREK@taunivm.tau.ac.il Date: 16 Jan 96 19:19 IST The Strange Spin of the Nucleon Authors: John Ellis (CERN) , Marek Karliner (Tel-Aviv Univ.) Comments: Invited Lectures at the International School of Nucleon Spin Structure, Erice, August 1995. The recent series of experiments on polarized lepton-nucleon scattering have provided a strange new twist in the story of the nucleon, some of whose aspects are reviewed in these lectures. In the first lecture, we review some issues arising in the analysis of the data on polarized structure functions, focusing in particular on the importance and treatment of high-order QCD perturbation theory. In the second lecture some possible interpretations of the "EMC spin effect" are reviewed, principally in the chiral soliton (Skyrmion) approach, but also interpretations related to the axial $U(1)$ anomaly. This lecture also discusses other indications from recent LEAR data for an $\bar{s}s$ component in the nucleon wave function, and discusses test of a model for this component. Finally, the third lecture reviews the implications of polarized structure functions measurements for experiments to search for cold dark matter particles, such as the lightest supersymmetric particle and the axion, after reviewing briefly</p>
--	--



the astrophysical and cosmological evidence for cold dark matter. -----

----- "Mechanical Propulsion From Unsymmetrical Magnetic Induction Fields" by: R.L. Schlicher A.W. Biggs W.J. Tedeschi 31st AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, July 10-12 1995

A method is presented for generating mechanical spacecraft propulsion from unsymmetrical magnetic induction fields. It is based on an unsymmetrical three-dimensional loop antenna structure driven by a repetitively-pulsed high-

current power supply. Antenna geometry is optimized for generating propulsive thrust rather than radiating electromagnetic energy. A magnetic field density gradient imbalances the magneto-mechanical forces that result from the interactions of the internal magnetic induction field with the current in the conductors of the antenna structure.

From Richard Feynman's Lectures on Physics we learn that there is intrinsic field energy and momentum density associated with a static electro-magnetic field configuration. When there is a change in the magnetic field, this field energy and momentum can be directly converted into kinetic energy and mechanical momentum. Feynman illustrates this with an electromagnetic carousel paradox. In this paradox, a dielectric disk (which is embedded with small charged spheres along its circumference) rotates without any apparent "counter" torque in the system. Before this rotation occurs, the dielectric disk is immersed in a static magnetic field. The subsequent rotation occurs as a consequence to reducing the previously static magnetic field to zero. The angular momentum and rotational kinetic energy comes directly from the initial static magnetic field.

"The Feynman Lectures on Physics" by Richard Feynman, R.B. Leighton, and M. Sands, Volume II p 17-6

"Nonlinear Electromagnetic Propulsion System and Method", R.L. Schlicher

Nineteenth Power Modulation Symposium of the IEEE, 1990 Page 139

"Classical Electrodynamics" by C.D. Jackson, 2nd Edition, John Wiley and Sons, New York, 1975

"The Feynman Lectures on Physics" Richard Feynman, R.B. Leighton, and M. Sands, Volume II p 27-9

US Patent #5142861, "Nonlinear Electromagnetic Propulsion System", R.L. Schlicher et al. 1992

Dr Peter Graneau has conducted experiments which he claims provide a demonstration of departure from classical electrodynamics at high currents levels. A force is found to exist in a direction longitudinal to current flow. Graneau ran a variety of types of experiments with a metal rod conductor immersed in a conductive fluid (mercury, or saline solution). With high amperage passing through the solution the metal rod is found to move in a longitudinal direction. There is no known explanation in conventional EM theory. This force may be similar to the force ($v \times B$) that William Hooper finds in a noninductive coil. Or Graneau's longitudinal force may be a coupling between the electromagnetic and inertial/gravitational fields, which is predicted by some 5-D unified EM/gravitational theories -- predicted to result from a divergence of the electric current vector field. Graneau's experiments should be relatively easy to duplicate. I can find no record that anyone has ever done so. Graneau has also discovered apparently anomalous forces and effects in high energy electromechanical devices such as rail guns and induction motors.

AUTHOR: Graneau, Peter.


TITLE: Ampere-Neumann electrodynamics of metals/Peter Graneau.

PUBL.: Nonantuma, MA. : Hadronic Press,

FORMAT: ix, 311 p. : ill. ; 23 cm.

DATE: 1985

SUBJECT Metals--Electric properties--History.



Free electron theory of metals--History.
Electrodynamics--History.
Electric conductors--History.
ISBN: 0911767371

AUTHOR: Graneau, Peter
TITLE: Electromagnetic Jet Propulsion in the Direction of Current Flow
In: Nature
June 18, 1982 No 295 Page 311

AUTHOR(s): Graneau, P.
TITLE(s): Ampere force calculation for filament fusion experiments.
In: Physics letters. a
MAR 22 1993 v 174 n 5/6 Page 421

AUTHOR(s): Graneau, P.
TITLE(s): Comment on "The motionally induced back-EMF in railguns".
In: Physics letters: [part A]
DEC 02 1991 v 160 n 5 Page 490

AUTHOR(s): Graneau, Peter
TITLE(s): The Difference between Newtonian and Relativistic Forces.
In: Foundations of physics letters.
OCT 01 1993 v 6 n 5 Page 491

AUTHOR(s): Graneau, P.
TITLE(s): Electrodynamic momentum measurements.
In: Journal of physics d: applied physics. DEC 01 1988 v 21 n 12 Page 1826

AUTHOR(s): Graneau, P.
TITLE(s): Far-action versus contact action.
In: Speculations in science and technology. 1990 v 13 n 3 Page 191

AUTHOR(s): Graneau, Peter
TITLE(s): Inertia's Riddle.
Summary: Inertia has been misunderstood ever since the time of Galileo says Dr. Graneau.
In: Electronics world + wireless world.
JAN 01 1990 v 96 n 1647 Page 60

AUTHOR(s): Graneau, P.

TITLE(s): Longitudinal forces in Ampere's wire-arc experiment.
In: Physics letters: [part A]
MAY 08 1989 v 137 n 3 Page 87

AUTHOR(s): Graneau, P. Thompson, D.S. Morrill, S.L.
TITLE(s): The motionally induced back-emf in railguns.
In: Physics letters: [part A]
APR 30 1990 v 145 n 8/9 Page 396

AUTHOR(s): Graneau, Peter
TITLE(s): Nonlocal Action in the Induction Motor.
In: Foundations of physics letters.
OCT 01 1991 v 4 n 5 Page 499

AUTHOR(s): Graneau, P. Graneau, N.
TITLE(s): The role of Ampere forces in nuclear fusion.
In: Physics letters: [part A]
MAY 04 1992 v 165 n 1 Page 1

AUTHOR: Graneau, Peter.
TITLE: Underground power transmission : the science, technology, and economics of high voltage cables / Peter Graneau. PUBL.: New York : Wiley,
FORMAT: x, 515 p. : ill. ; 24 cm.
DATE: 1979
SUBJECT: Electric cables
Electric power transmission
Electric lines--Underground
ISBN: 0471057576

I see that Graneau has devoted himself to the electric railgun. I looked a bit into this phenomena in 1994. I will now be able look a bit further. It seems that Graneau believes in free energy in vacuum. The railgun and the theories about it are very controversial. There is a conflict between PhD Witalis, who works for the Swedish Defense, and the established plasmaphysicists here in Uppsala. Witalis has condemned controlled hot fusion.

-- David Jonsson

MAGNETORESISTANCE IN METALS, by Pippard, A.B. Pippard provides the first systematic account of magnetoresistance in metals, the study of which has provided solid-state physicists with valuable information about electron motion in metals. The electrical resistance of a metal is usually changed when a magnetic field is applied to it, and at low temperatures the change may be very large indeed. Every metal behaves differently, and the effect has been widely used to elucidate details of electron motion in individual metals. Because there has been no systematic account of the phenomena, apart from review articles addressing special points, this book fills an obvious gap. Making no great demand on mathematical ability, it should be a valuable reference work for readers with a basic knowledge of undergraduate solid-state physics. The text is copiously illustrated with real experimental results. Cambridge Studies in Low Temperature Physics 2 1989 6 x 9 272 pp. 3 halftones 113 line diagrams Hardback 0-521-32660-5 \$84.95 (J50.00)

Albert C. Crehore published "New Electrodynamics" in 1950. In this book he described how the motion of protons in the nucleus would produce gravitational field effects. Gravitational field effects such as counter-bary are used in the mainstream effort to develop non-areodynamic non-rocket flight systems that usually referred to as "anti-gravitational". By making use of the Crehore Paradigm it is possible to derive a method of producing counter-bary. It's most likely Crehore had no knowledge of Brown's 1928 British patent for a gravitator device that would have been a macroscopic analog of a Crehore atom.

Crehore, Albert C. (Albert Cushing), b. 1868. The mystery of matter and energy; recent progress as to the structure of matter, by Albert C. Crehore ... New York, D. van Nostrand company, 1917 LC CALL NUMBER: QC173 .C8
SUBJECTS: Matter--Constitution.

Crehore, Albert C. (Albert Cushing), b. 1868. The atom, by Albert C. Crehore ... New York, D. Van Nostrand company, 1920. xvi, 161 p. diagrs. 19 cm.

LC CALL NUMBER: QC173 .C75

A man named Gerry Vassilator is an electrogravity experimenter. Last I know (in 1991) he runs an information service called MUUDO Experimental Videos

Delmar Ave
Staten Island, NY 10312.
(718)-356-9373.

There are many books available about anti-gravity and other weird science subjects from:
International Tesla Society
330-A West Uintah Street - Suite 215
Colorado Springs CO 80905-1095

The Anti-Gravity Handbook (revised ed.)
Compiled by D. Hatcher Childress
Published by - Adventures Unlimited Press
303 Main St., Kempton, Illinois 60949 USA
ISBN: 0-932813-20-8
Pub date: 1993 (First edition was in 1985) ----

Anti-gravity and the world grid / edited by David Hatcher Childress. 1st ed. Stelle, IL : Adventures Unlimited Press, c1987. 267 p. : ill. ; 26 cm. LC CALL NUMBER: BF1999 .A6386 1987
SUBJECTS: Antigravity. Grids (Cartography). Occultism. Childress, David Hatcher, 1957- ISBN: 0932813038 (pbk.) : \$12.95

The Anti-gravity handbook / compiled by D. Hatcher Childress. 1st ed. Stelle, Ill. : Adventures Unlimited Press, c1985 (1986 printing) 195 p. LC CALL NUMBER: QC178 .A58 1985
SUBJECTS: Antigravity.

Childress, David Hatcher, 1957-

ISBN: 0932813011 (pbk.) : \$12.95

Pages, Marcel J. J.

Le defi de l'antigravitation: techniques antiponderales, utilisation de l'energie de l'espace [par] M. J. J. Pages. Paris, Chiron [1974] 306 p. LC CALL NUMBER: QC178 .P23

SUBJECTS: Antigravity. Force and energy.

ISBN: 2702703097

Nipher, Francis Eugene, 1847-

Electricity and magnetism. A mathematical treatise for advanced undergraduate students. By Francis E. Nipher ... 2d ed., rev., with additions.

St. Louis, Mo., J. L. Boland book and stationery co., 1895 i.e. 1898 xi, 430 p. diagrs. 20 cm.

"My library research shows that as early as 1917, a Professor Nipher had found that the weight of substances could be reduced (become negative) by the application of electrostatic charges. (Science, Sept. 21, 1917, page 173).

Dr. Charles Brush, in a series of reports in the PROCEEDINGS OF THE AMERICAN PHILOSOPHICAL SOCIETY around 1922 found, in some well-thought-out-experiments, that weight was not only proportional to mass, but was affected by the atomic structure of the substances. For example, he found that for a given unit of mass and shape, BISMUTH falls faster than zinc or aluminum, in complete contradiction to Newton's Law of Gravity which they are still teaching in colleges today!

So far, the literature hasn't given me an answer. Incidentally, Otis Carr's work involved counter-rotating charged discs that supposedly produced thrust when they reached a certain speed in relation to the earth's rotational speed and became activated by free energy from space. Maybe he did have something." -- James E. Cox

When individual molecules are not permanently magnetized, it is possible in some cases to have a relative magnetic permeability μ which is less than one. Such a material, like hydrogen or BISMUTH, is called diamagnetic. It tends to expel magnetic field, and is repelled from regions of stronger magnetic field. The names paramagnetic and diamagnetic are sometimes confused: paramagnetic is analogous to a dielectric in an electric field, while diamagnetic is quite the opposite.


It is not possible to give a simple argument of why diamagnetism can occur. It is strictly speaking a quantum effect. However, one can see that there might be diamagnetic tendencies if electric currents can flow within molecules. An increasing magnetic field always tends to induce currents to flow in such a way as to tend to prevent the increase in the field. This is (at least temporarily) a diamagnetic kind of effect.

Thus the case where the relative magnetic permeability $\mu < 1$, is connected with the flow of electric charges in a magnetic field. There is no analogous case with electric fields since isolated magnetic poles do not, so far as is known exist. - "The New Physics" edited by Paul Davies

GE engineer Henry Wallace found unusual gravitational effects in spinning odd atomic nucleide metals. Odd atomic nucleide metals are those in which the sum of the protons are not equal to the number of neutrons, i.e. more neutrons. See US patents 3626605 and 3626606. -Ron Kita

"Also indicated in the embodiment is the orientation of the flux within the mass circuit, the latter being constructed preferably of BISMUTH."

- Henry Wallace, US patent # 3626605, Method and Apparatus for Generating a Secondary Gravitational Force Field



AUTHOR(s): Uyeda, C. Yamanaka, T. Miyako, Y.

TITLE(s): Magnetic rotation of diamagnetic oxide crystals and the origin of diamagnetic anisotropies.

In: *Physica B. Condensed matter.*
MAY 01 1995 v 211 n 1/4 Page: 342

A while back I had the need to take a peek at a copy of the periodic table of elements. So I grabbed my old, dusty college chemistry book that I could never quite bring myself to toss. It's called "Chemical Principles", published WAY back in 1970. While looking through the book, I was stunned when I came across a discussion of the possibilities of new elements. "What lies ahead for the synthesis of transuranium elements? Will there be more radioactive and extremely short-lived species such as 97 through 104? It now appears as if there is a chance of reaching a new zone of stability that might even include some non-radioactive elements. Calculations with nuclear shell models have led to the expectation that element 114, with 114 protons and 184 neutrons (both magic numbers in the new shell theory) would be an island of stability in a sea of instability."

I noted that some information was taken from an article in the April 1969 (pages 57-67) issue of "Scientific American" by Dr. Glenn Seaborg. In this article, there are excellent graphics showing the expected half-lives of all the heavyweights. They predict a fission half-life for the most stable isotope of 114 of 10 to the 16th years, and a alpha-decay half-life of 1,000 years. They didn't go into the same level of detail for 115, but it looks like the stuff would clock out considerably sooner by way of beta decay. BTW, according to the article, the proper terminology to denote an undiscovered element in a periodic column is the prefix "eka". Therefore element 115 should be eka-BISMUTH. Lose this Un-un-pentium crap! - Tom Mahood

Lazar is not the only one to theorize that the 114/115 area is stable. Check out the August

31, 1991 issue of *New Scientist* -- that respected peer-reviewed periodical. Find Glenn Seaborg's article called "The search for the missing elements." Seaborg is a renowned scientist who won the Nobel Prize for Chemistry in 1951. He and his research group at the Lawrence Berkeley Laboratory have discovered 10 of the transuranium elements. His article is very technical and interesting, and in it he has two separate graphs that show islands of stability at the 114/115 area. There is a sea of instability around these "islands." -----

AUTHOR(s): Bhattacharyya, S. Ghoshal, A. Ghatak, K.P.

TITLE(s): On the field emission from bismuth in the presence of a quantizing magnetic field.

In: *Fizika; a journal of experimental and theoretic* APR 01 1991 v 23 n 2 Page 159

AUTHOR(s): Byrne, A. P. Birkental, U. Hubel, H.

TITLE: High-Spin States in 205Bi.

In: *Zeitschrift fur Physik. A, Atomic nuclei* 1989 v 334 n 3 Page: 247

AUTHOR(s): Vezzoli, G.C. Chen, M.F. Craver, F.

TITLE(s): Magnetically-related properties of bismuth containing high Tc superconductors.

In: *Journal of magnetism and magnetic materials.* AUG 01 1990 v 88 n 3 Page 351

AUTHOR(s): Bannerjee, D. Bhattacharya, R.

TITLE(s): Magnetic Properties of Single Crystals of Bismuth Doped with Lead and Tin.

In: *Physica status solidi. b: basic research.* JAN 01 1990 v 157 n 1 Page 443

AUTHOR(s): Zhilyaev, I. N.

TITLE(s): Observation of kinetic paramagnetic effect in bismuth in a transverse magnetic field.

In: *Soviet journal of low temperature physics.* SEP 01 1988 v 14 n 9 Page 502



AUTHOR(s): Mondal, M. Banik, S.N. Ghatak, K.P.

TITLE(s): Effect of a quantizing magnetic field on the Einstein relation in bismuth.

In: Canadian journal of physics.
JAN 01 1989 v 67 n 1 Page 72

AUTHOR(s): Zheng, Q. Zeng, Z. Lai, W.

TITLE(s): The influence of Al on the electronic structure and magnetic properties of doped MnBi with huge enhancement of Kerr rotation.

In: Journal of magnetism and magnetic materials. FEB 01 1992 v 104/107 p 2 Page 1019

this novel mechanism of charge transport and comment on the effects of disorder.

Huston, David L.

The nature and possible significance of the Batamote copper-bismuth-silver anomaly, Pima County, Arizona / by David L. Huston and Paul K. Theobald. Washington : U.S. G.P.O. ; Denver, CO : For sale by the Books and Open-File Reports Section, U.S. Geological Survey, 1990. v, 19 p. : ill., maps LC CALL NUMBER: QE75 .B9 no. 1907 (ALTERNATE CLASS QE390.2.C6) SUBJECTS: Copper ores--Arizona--Batamote Mountains Region. U.S. Geological Survey bulletin ; 1907

Beck, Sherwin M.

Measured electron conversion ratios for the 1064-keV gamma ray of bismuth-207, by Sherwin M. Beck. Washington, National Aeronautics and Space Administration; [for sale by the Clearinghouse for Federal Scientific and Technical Information, Springfield, Va.] 1970. 39 p. illus. 27 cm. LC CALL NUMBER: TL521 A3525 no. 6057 SUBJECTS: Bismuth--Isotopes. Nuclear counters. NASA technical note, NASA TN D-6057

The following is an excerpt from a telephone interview between Stanton Friedman (F) and Dr. Robert Sarabacher (S). Sarabacher was a prominent, US government scientist who had a secret briefing with Canadian scientist Wilbert Smith in 1950 and told Smith that facts in a recent popular book about a UFO crash at Aztec, New Mexico were "essentially true" and that UFO's were classified by the US government 2 points higher than the H bomb. Sarabacher died in July 1986. Before Sarabacher died, Stanton Friedman did a phone interview with him. In between Friedmann's attempts to dig more UFO info out of Sarabacher, there was a lot of small talk, and since Sarabacher was fairly old, he tended to ramble a bit. However, a most interesting statement was made by Sarabacher:

F: Were you guys talking about nuclear powered flight at that time?

S: Oh, we were possibly, yes, but I held, had certain ideas see, one of the problems today, we really don't know what gravity is. We don't know and I had an idea, I'm willing to work on it in one of my theses but then my professor didn't believe me, but I had determined that BISMUTH did not obey the laws of gravity. So I thought that, "Gee, there's a leak". I might be able to get nature to tell me something.

So where exactly is Bismuth on the Periodic Table of Elements? Why directly above where 115 would fall if it exists. And the way the table works, (generally speaking) elements in the same column have similar properties. So, just what the hell was Sarabacher referring to? I don't know, but it's sure intriguing! It appears it was back when he was a grad student, in maybe the 30s or 40s. Whatever it was, it was at the very edge of the ability of equipment at the time. Does Bismuth possess any very subtle anomalous physical properties? -- Tom Mahood

In the Wallace patent, #3626606, Figs. 7A and 7B are side views of a gravity-NEUTRALIZING FLYING SAUCER, or, if anchored to the ground, a ZERO-GRAVITY CHAMBER. Each oval diagram shows a motor spinning a central disc at a very high speed, about 28,000 RPM, and also rotating two other discs sandwiched around the first disc, via gears, at a much slower speed, perhaps 2,800 RPM, in the opposite direction. The two outer discs have extensions [counter-balanced via off-center axis] that, as they rotate, alternately make contact with two wide extensions from opposite walls of the spacecraft. The central disc should have shallow spiral-shaped grooves on both sides for air-bearings, to allow the needed very close contact with the two outer discs. Each of the two outer discs has ONLY ONE [counter-balanced] extension, each one pointed opposite (180 degrees) the extension of the other disc. The most important factor making it work is

that the discs, extensions, and outer walls of the spacecraft MUST be made of any material(s) in which a very large majority of the atoms are of isotopes having "HALF-INTEGRAL ATOMIC SPINS", such as copper (3/2).

- Robert E. McElwaine

AUTHOR(s): Sun, W. Stephen, J.T. Wu, Y.
TITLE(s): Rotation-Induced Resonance and Second-Order Quadrupolar Effects on Spin Locking of Half-Integer Quadrupolar Nuclei.

In: Journal of magnetic resonance. series a. OCT 01 1995 v 116 n 2 Page: 181

AUTHOR(s): Seliger, J. Blinc, R.
TITLE(s): Orientation dependences of quadrupolar spin-lattice relaxation rates of spin-3/2 nuclei subject to a random two-site exchange in a high magnetic field: a theoretical study.

In: Journal of physics. Condensed matter : an Inst DEC 13 1993 v 5 n 50 Page: 9401

Thanks for the patent info about Wallace, I got them a few days ago and found it quite enjoyable. Actually, many people have had this notion at one time or another, I think Oleg Jefimenko wrote a book relating to this subject.

-- Keith Nagel

AUTHOR: Jefimenko, Oleg D.
TITLE: Electricity and magnetism : an introduction to the theory of electric and magnetic fields / Oleg D. Jefimenko. EDITION: 2nd ed.

PUBL.: Star City, W. Va. : Electret Scientific Co.,

DATE: 1989

SUBJECT: Electromagnetism

ISBN: 0917406081

AUTHOR: Jefimenko, Oleg D.
TITLE: Electrostatic motors; their history, types, and principles of operation (by) Oleg D. Jefimenko. With many illus., of which 57 are by David K. Walker.



PUBL.: Star City (W. Va.) Electret Scientific Co.

DATE: 1973

SUBJECT: Electrostatic apparatus and appliances

AUTHOR: Jefimenko, Oleg D.

TITLE: Causality, electromagnetic induction, and gravitation: a

different approach to the theory of electromagnetic and gravitational fields" b

PUBL: Star City [West Virginia] : Electret Scientific Co.,

DATE: 1992.

SUBJECTS: Electromagnetic fields. Gravitational fields. Causality. Maxwell Equations.

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Direct calculation of electric and magnetic forces from

In: American journal of physics.

JUL 01 1990 v 58 n 7 Page 625

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Direct calculation of the electric magnetic fields of an electric point charge moving with constant velocity. In: American journal of physics.

JAN 01 1994 v 62 n 1 Page 79

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Force exerted on a stationary charge by a moving electric current or by a moving magnet.

In: American journal of physics.

MAR 01 1993 v 61 n 3 Page 218

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Retardation and relativity: The case of a moving line charge.

In: American journal of physics.

MAY 01 1995 v 63 n 5 Page 454

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Retardation and relativity; Derivation of Lorentz-Einstein transformation from retarded integrals for electric and magnetic fields.

In: American journal of physics.

MAR 01 1995 v 63 n 3 Page 267

AUTHOR(s): Jefimenko, Oleg D.

TITLE(s): Solutions of Maxwell's equations for electric and magnetic fields in arbitrary media.

In: American journal of physics.

OCT 01 1992 v 60 n 10 Page 899

"Anti-Gravity Electronics", H. Aspden, Electroncis & Wireless World, Jan 1 1989, Vol 95 No 1635

Reinterpretation of Netwon's third law of motion suggests that it depends upon an electronic action. Electronic interaction therefore explains the paradoxical anti-gravity properties of the force precessed gyroscope.

"The Anti-Gravity Puzzle", Mark Ander, Professional Pilot, Aug 1 1989 Exploring the possibility of exceptions to Newton's inverse-square law of gravity, scientists pursue evidence in strange locations.


"The Latest Antigravity Gossip", Rock & Ice, Nov 1 1994 No 64

"Propulsion by Gyro", Eric Laithwaite, Space, Sep 1989 Vol 5 No 5 In an attempt to reveal the strange, hidden properties of gyroscopes, Professor Eric Laithwaite explains the physics behind the idea that a propulsion system could be built using gyros.

"Negative Mass in General Relativity", H. Bondi, Reviews of Modern Physics, Vol 29, July 1957, pp 423-428

"Looking for New Gravitational Forces with Antiprotons", M.M. Nieto and B.E. Bonner, Proceedings RAND Workshop on Anti Proton Science and Technology, World Scientific, Singapore, 1988 pp 328-341

"Negative and Imaginary Proper Masses", Y.P Terletskii, Paradoxes in the Theory of Relativity, Plenum, New York 1968, Chapter VI pp 83-115



"Gravitational Coupling of Negative Matter", A. Inomata and D. Peak, Nuovo Cimento, Vol B63 Sep 1969 pp 132-142

"Negative-Mass Lagging Cores of the Big Bang", B.D. Miller, Astrophysical Journal, Vol 208, Sep 1976 pp 275-285

"The Cosmological Term, the Shielding of Gravitation and the Negative Mass Hypothesis", A.A. Baranov, Izvestiya VUZ Fizika, Vol 14 Nov 1971 pp 118-120

"Negative Masses and the Energy-Sources of the Universe", Y.P. Terletskii, Experimentelle Technik der Physik, Vol 29 April 1981 pp 331-332

ELECTROMAGNETIC-GRAVITATIONAL CONVERSION CROSS SECTIONS IN EXTERNAL ELECTROMAGNETIC FIELDS

International Centre for Theoretical Physics, Trieste (Italy). LONG, HOANG NGOC SOA, DANG VAN TRAN, TUAN A. SEP. 1994 11 PAGES DE95-613589 IC-94/285 Avail: CASI HC A03/MF A01 (US Sales Only) The classical processes: the conversion of photons into gravitons in the static electromagnetic fields are considered by using Feynman perturbation techniques. The differential cross sections are presented for the conversion in the electric field of the flat condenser and the magnetic field of the solenoid. A numerical evaluation shows that the cross sections may have the observable value in the present technical scenario. CASI Accession Number: N95-30637

I have an excerpt from a paper presented under the auspices of Northrop Corp. in 1968 that gives an idea of why you would want to bother with high voltage fields. To achieve the effects described involved relatively small high voltage. The true electrogravitational effects are significant at higher E field strength. In any case, this was only the state of the art in 1968.

Their involvement in the B-2 began much later, after considerably more research.

Electroaerodynamics In Supersonic Flow
by M. S. Cahn and G. M. Andrew, Northrop Corporation, Hawthorne, California Presented at AIAA 6th Aerospace Sciences Meeting, January 22-24, 1968 - Tom Capizzi (tcpapizzi@world.std.com)

ELECTRO-AERODYNAMICS: Electric charges are applied to high-speed vehicles for the purpose of reducing air drag or eliminating sonic booms. High-speed ions are projected forward from the leading edges of the craft, the corona glow propagates forward and repels air molecules away from the oncoming surfaces, thus a shock wave cannot be mechanically produced.

Sources: Dudley, Horace C., Analog Science Fact & Fiction. "The Electric Field Rocket", November 1960.

Product Engineering. "Sonic Boom Experiments", Vol. 39, New York, pp. 35-6, March 11, 1968.

US Patent No. 3,095,167, Dudley.

The paper entitled the "U.S. Antigravity Squadron" paper appears with others in the book "ELECTROGRAVITICS SYSTEMS: Reports on a New Propulsion Methodology" edited by Thomas Valone (Washington, D.C.: Integrity Research Institute, 1994); ISBN 0-9641070-0-7.

In addition to this paper, this book also includes the following:

- 1) The 1956 paper "Electrogravitics Systems" (prepared by the Special Weapons Study Unit of Aviation Studies Ltd., a UK-based aviation industry intelligence firm). It was declassified from a confidential status some time prior to 1985 and entered the public domain as a result of a request I placed through the Wright-Patterson Air Force Base Technical Library.



2) The 1956 paper "The Gravities Situation" (prepared by Gravity Rand Ltd., a division of Aviation Studies Ltd. This includes six appendices with papers by various authors including the text from T. Townsend Brown's 1929 gravitor patent.

3) A paper by Banesh Hoffman entitled "Negative Mass as a Gravitational Source of Energy in the Quasistellar Radio Sources.

4) A collection of diagrams copied from various patents by T. Townsend Brown.

You may order a copy from:
Starburst Publications, 1176 Hedgewood Lane,
Schenectady NY 12309, USA

Also available from Starburst Publications is the book "Subquantum Kinetics: The Alchemy of Creation" (ISBN 0-9642025-0-6). Subquantum kinetics is a new approach to microphysical theory that utilizes concepts from the fields of nonlinear chemical kinetics, irreversible thermodynamics, and general system theory, replacing the current mechanistic foundation of physics with a reaction-kinetic model. This new approach resolves a number of problems that plague classical and modern physics also may provide some insights into the electrogravitic connection that Brown was researching. In particular, chapter 9 gives some background information on Townsend Brown's electrogravitics.

Scott, W.B. "Black World engineers, scientists encourage using highly classified technology for civil applications." Aviation Week & Space Technology, March 9, 1992, pp. 66,67.

Brown, T.T. "How I Control Gravity." Science and Invention Magazine, August 1929. Reprinted in Psychic Observer 37(1) pp.14 - 18.

Burridge, G. "Another Step Towards Antigravity." The American Mercury 86(6) (1958):77 - 82.

Sigma, Rho, "Ether Technology: A Rational Approach to Gravity Control." Lakemont, GA: CSA Printing & Bindery, 1977, p. 44-49, quoting a letter from T. Townsend Brown dated February 14, 1973.

Intel. "Towards Flight Without Stress or Strain...Or Weight." Intervia Magazine 11(5) (1956):373-374

Rose, M. "The Flying Saucer: The Application of the Biefeld-Brown Effect to the Solution of the Problems of Space Navigation." University for Social reesearch, April 8, 1952.

LaViolette, P.A. "An Introduction to Subquantum Kinetics: Part Journal of General Systems, Special Issue on Systems Thinking in Physics" 11(1985):295-328.

LaViolette, P.A. "Subquantum Kinetics: The Alchemy of Creation." Schenectady, NY, 1994.

LaViolette, P.A. "Beyond the Big Bang: Ancient Myth and the Science of Continuous Creation." Rochester, VT:Inner Traditions Intl., 1994.

LaViolette, P.A. "A Theory of Electrogravitics." Electric Spacecraft Journal, Issue 8, 1993, pp. 33 - 36.

LaViolette, P.A. "A Tesla Wave Physics for a Free Energy Universe." Proceedings of the 1990 International Tesla Symposium, Colorado Springs, CO: International Tesla Society, 1991, pp. 5.1 - 5.19.

Aviation Studies (International) Ltd. "Electrogravitic Systems: An Examination of Electrostatic Motion, Dynaimc Counterbary and Barycentric Control." Report GRG 013/56 by Aviation Studies, Special Weapons Study Unit, London, February 1956. (Library of



Congress No. 3,1401,00034,5879; Call No. TL565.A9).

LaViolette, P. "Electrogravitics: Back to the Future." Electric Spacecraft Journal, Issue 4, 1992, pp. 23 - 28.

LaViolette, P. "Electrogravitics: An Energy-Efficient Means of Spacecraft Propulsion." Explore 3 (1991): 76 - 79; idea No. 100159 submitted to NASA's 1990 Space Exploration Outreach Program.

Aviation Studies (International) Ltd. "The Gravitics Situation". prepared by Gravity Rand Ltd. - a divison of Aviation Studies, London, December 1956.

Northrup Studying Sonic Boom Remedy." Aviation Week & Space Technology, Jan. 22, 1968, p.21.

Rhodes, L. "Ex-NASA Expert Says Stealth Uses Parts from UFO." Arkansas Democrat, Little Rock, AR., April 9, 1990.

Scott, W.B. "Inside the Stealth Bomber" Tab/Aero Books: New York, 1991.

One of the most famous researchers in this area is John Searl, who noticed that spinning metal would accumulate electrons on the rim, possibly through some kind of centrifugal thrust. The initial test was a metal disk attached to a breakaway coupling driven by a gasoline engine. It was carried out in the country and as the disk reached higher and higher speeds, tremendous electrostatic forces were generated which were estimated at 10 to the 6th volts! That's when the disk began to glow blue, broke the coupling, rose to about 30 feet, continued to accelerate, turned pink and shot off into space. Searl claims this happened with many of his early tests and as a result he lost the device each time. Later he learned how to control the device. The neat thing about it, the Searl disk is self-propelling using a magnetic drive.

-- Jerry Decker

THE SEARL EFFECT

(The Introduction)

[To contact WCVE write to:
23 Sesame Street
Richmond VA, 23235
or phone: 804-320-1301
or fax: 804-320-8729]

ANTIGRAVITY: The Dream made Reality

[The Story of John R. R. Searl]

by John A. Thomas Jr.

Published by Direct International Science Consortium 13 Blackburn, Low Strand, Grahame Park Estate, London NW95NG England

Available in this country through John A. Thomas, Jr. 373 Rock Beach Rd.
Rochester, NY 14617-1316
Phone: (716) 467-2694

Thomas, John A., Jr.

Antigravity : the dream made reality : the story of John R.R. Searl / by John A. Thomas Jr. London : Direct International Science Consortium, c1993. 1 v. (various pagings) : ill. (some col.) ; 28 cm. LC CALL NUMBER: QC178 .T46 1993

SUBJECTS: Searl, John R. R. (John Roy Robert), 1932-. Antigravity. ISBN: 1898827990 (spiral)

NEXUS Magazine

Volume 2, Number 17

P.O. Box 177

Kempton, IL 60946

Phone: (815) 252-6464


Fax: (815) 253-6300

Extraordinary Science

Volume VI. Issue 2

ISSN 1043-3716

Aspden, Harold A. (1989). "The Theory of the Gravitation Constant," Physical Essays, Vol. 2, No. 2, pages 173-179.



Aspden, Harold A. (1991). "The Theory of Antigravity," Physical Essays, Vol. 4, no. 1, pages 13-19.

Electrogravitic Systems: Reports on a New Propulsion Methodology by Thomas Valone, M.A., P.E.

The Anti-Gravity Handbook
by D. Hatcher Childress

Ether-Technology: A Rational Approach to Gravity-Control by Rho Sigma

Sigma, Rho.

Forschung in Fesseln : das Ratsel d. Elektro-Gravitation / von Rho Sigma. Wiesbaden-Schierstein : Ventla-Verl., 1972. 272 p. : ill. ; 21 cm. LC CALL NUMBER: TL789 .S524 1972

SUBJECTS: Unidentified flying objects. Gravitation. ISBN: DM24.00

There is also a good book written by an Aerospace Engineer who worked for General Electric, named John Ackerman. The book is called "To Catch a Flying Star". It is available from Univelt, Inc., P.O. Box 28130, San Diego, CA. 92128 ISBN 0-912183-03-9.

AUTHOR: Ackerman, John.
TITLE: To catch a flying star : a scientific theory of UFOs / by John Ackerman ; with a forward by Walter H. Andrus, Jr. PUBL.: San Diego, Calif. : Univelt,
DATE: 1989
SUBJECT: UNIDENTIFIED FLYING OBJECTS, SPACECRAFT, UFOs, INTERSTELLAR SPACECRAFT, INTERSTELLAR TRAVEL LONG DURATION SPACE FLIGHT, ASTRONAUTICS FLIGHT CONTROL
ISBN: 0912183039 (pbk.)

ELECTRIC SPACECRAFT JOURNAL \$24

yr. US P.O. BOX 18387 \$29 yr.

Canada/Mexico

Asheville, North Carolina 28814 \$39

yr. other countries

I highly recommend this semi-pro publication. Buy all the back issues too! It's not just 'counterbary', but has articles on energy anomalies, Tesla, unconventional hobby projects, unconventional physics, etc.

SPACE ENERGY NEWS The Space Energy Association is dedicated

PO Box 11422 to the pioneering work of several sci-

Clearwater FL 34616 entists and inventors, including Nikola

Suscription info:\$35, 4 issu/yr Telsa, Viktor Schauberger, T. Henry Moray, Editors: Donald A. Kelly T.T. Brown, Alfred Hubbard, T.J.J. See,

Michael MarinoErwin Saxl, Hans Coler and others.


"Antigravity" by R.M. Santilli

The Institute for Basic Research,

PO Box 1577

Palm Harbor FL 34682

Recently I had read a book from Hans A. Nieper with the title "Konversion von Schwerkraft-Feld-Energie" (transformation of gravity field energy). This book tells from transformers of the types Fluxtransformer (electrodynamic system) and Capacitor Discharge (solid state method). My problem is that I find nowhere informations about these machines. So I hope that YOU can give me hints whre I can find informations (books, files,articles ...) about these machines and theories. I am also grateful for your opinions about these theory and these machines. -- Michael Bell Berlin, Germany



Revolution in Technik, Medizin, Gesellschaft. English. Revolution in technology, medicine and society : conversion of gravity field energy / Hans A. Nieper. Extended ed. in English, 1. ed. Oldenburg : MIT Verlag, 1985. 384 p. : ill. ; 22 cm.

LC CALL NUMBER: TJ163.7 .R4813 1985
 SUBJECTS: Power (Mechanics)--Congresses. Translation of: Revolution in Technik, Medizin, Gesellschaft. Cover title: Dr. Nieper's Revolution in technology, medicine, and society. Includes proceedings of the Symposium on Energy Technology, Hannover, Nov. 27-28, 1980, and the First International Symposium on Non-Conventional Energy Technology, Oct. 23-24, Toronto, Oct. 23-24, 1981. ISBN: 392518807X

Nieper, Hans A. Zur Theorie der Schwerkraftwirkungen. In "Revolution in Technik Medizin Gesellschaft",

Bearden, T. E. Maxwell's lost Unified Field Theory of Electromagnetics and Gravitation. In "New Energy Technology", pg. 25. Published by The Planetary Association for Clean Energy, nc. Ottawa/Hull, Canada.

Bearden, Thomas E. (1988). "Maxwell's Original Quaternion Theory Was a Unified Field Theory of Electromagnetics and Gravitation," Proceedings of the International Tesla Society, 1988, ITS Books.

Moretti, Angelo. Possibility of Non-Zero Mass in Synchrotron Radiation. In "What Physics for the next century?" pg. 397 - Inediti No. 59, Societ Editrice Andromeda, Bologna

Gunnufson, Craig. Neuere Neutrinomessungen aus der Sonne unterstuetzen eine neue Theorie. Lecture held at a congress on Gravity Field Energy in Toronto, Oct. 1981. In Nieper, "Revolution in Technik Medizin Gesellschaft", Illmer Verlag, Hannover

Seike, Shinichi. Lecture held at Energy Symposium in Hannover, November 1980. In Nieper, Revolution in Technik Medizin Gesellschaft. -----

Anyone ever see Stan Deyo's book Cosmic Conspiracy? He talked about an ElectroGravitics society I think it was.

These things must have some sort of high voltage resonating circuitry, in a round shape obviously. They can recharge over high power lines. They can apparently become invisible. They also must be able to change their mass so they can accelerate at very high rates.

For anyone interested, he wrote a second book called "The Vindicator Scrolls" which contains more information.

"Space Warps: A Review of One Form of Propulsionless Transport," _Journal of the British Interplanetary Society_ 42 (Nov. 1989): 533-542.

"Negative Matter Propulsion," _Journal of Propulsion and Power_ 1 (Jan.-Feb. 1990): 28-37.

Vonsovskii, S, Ferromagnetic Resonance, 1966. Feynman, R, Feynman Lectures on Physics, v2, 1964 Chikazumi, S, Physics of Magnetism, 1964 Soohoo, R,


Microwave Magnetics, 1988

Herlach, F, Strong and Ultrastrong Magnetic Fields, 1985 -----

[1] The Feynman Lectures on Physics, 1963, v1 chp13 p8.

[2] Bottcher, C., Theory of Electric Polarization, 1973, v1 pp71,289.

[3] Albert Einstein: Philosopher-Scientist, 1949, pp522-523.



Feynman's Lectures Vol II Chapter 10, Page 10-8, describes an electrostatic effect that may be related to the Biefeld-Brown effect. Feynman shows that a force results on a dielectric due to the gradient of the square of the electrical field.

There have been quite a few people who have also looked at 'gravitationless' universes, such as:

Ralph Juergens, 'Reconciling Celestial Mechanics', Pensee Fall 1992. C E R Bruce, A New Approach to Astrophysics and Cosmogony, London 1944 Problems of Atmospheric and Space Electricity, Elsevier, 1965 Eric Crew, Electricity in Astronomy, SIS Review, Vol 1 No 1-4. Earl Milton, Electric Stars in a Gravity-Less Electrified Cosmos SIS Review, Vol V, No 1.

Ian Tresman
London, UK

AUTHOR: Driscoll, R.B.
TITLE: Comments on the paper "Gravitational lift via the Coriolis force" by Leon R. Dragone.
In: Hadronic journal.
JUL 01 1988 v 11 n 4 Page: 177

Paper: gr-qc/9503060
From: linet@ccr.jussieu.fr (Bernard LINET)
Date: Thu, 30 Mar 1995 14:55:07 +0200
Title: Vacuum polarization induced by a uniformly accelerated charge Author: B. Linet
Report-no: GCR-941003
We consider a point charge fixed in the Rindler coordinates which describe a uniformly accelerated frame. We determine an integral expression of the induced charge density due to the vacuum polarization at the first order in the fine structure constant. In the case where the acceleration is weak, we give explicitly the induced electrostatic potential.


Paper: gr-qc/9504023
From: Mathias PILLIN Date: Mon, 17 Apr 1995 10:43:50 +0900
Title: Pure spin-connection formulation of gravity and classification of energy-momentum tensors

Author: Mathias PILLIN
Report-no: YITP/U-95-12
It is shown how the different irreducibility classes of the energy-momentum tensor allow for a pure spin-connection formulation. Ambiguities in this formulation especially concerning the need for constraints are clarified.

Paper: gr-qc/9504041
From: SHORE@crnvma.cern.ch
Date: Tue, 25 Apr 95 17:22:56 SET
Title: "Faster than Light" Photons in Gravitational Fields -- Causality, Anomalies and Horizons

Authors: G.M. Shore
Report-no: SWAT-95/70
A number of general issues relating to superluminal photon propagation in gravitational fields are explored. The possibility of superluminal, yet causal, photon propagation arises because of Equivalence Principle violating interactions induced by vacuum polarisation in QED in curved spacetime. Two general theorems are presented: first, a polarisation sum rule which relates the polarisation averaged velocity shift to the matter energy-momentum tensor and second, a 'horizon theorem' which ensures that the geometric event horizon for black hole spacetimes remains a true horizon for real photon propagation in QED. A comparison is made with the equivalent results for electromagnetic birefringence and possible connections between superluminal photon propagation, causality and the conformal anomaly are exposed.

Paper: hep-th/9506035
From: Gary Gibbons Date (revised): Sun, 27 Aug 95 11:31:39 BST Title: Electric-Magnetic Duality Rotations in Non-Linear Electrodynamics



Authors: G W Gibbons, D A Rasheed -- To appear in Nucl Phys B Report-no: DAMTP preprint # R95/46.

We show that there is a function of one variable's worth of Lagrangians for a single Maxwell field coupled to gravity whose equations of motion admit electric-magnetic duality.

Paper: gr-qc/9506053

From: ESPOSITO@napoli.infn.it

Date: Mon, 26 Jun 1995 10:24:36 +0200 (CET-DST) Title: Euclidean Maxwell Theory in the Presence of Boundaries Author: Giampiero Esposito

Comments: 18 pages, plain-tex, to appear in: Heat-Kernel Techniques and Quantum Gravity, Discourses in Mathematics and Its Applications, No. 4, edited by S.A. Fulling (Texas A&M University, College Station, Texas, 1995)

Report-no: DSF preprint 95/31

This paper describes recent progress in the analysis of relativistic gauge conditions for Euclidean Maxwell theory in the presence of boundaries. The corresponding quantum amplitudes are studied by using Faddeev-Popov formalism and zeta-function regularization, after expanding the electromagnetic potential in harmonics on the boundary 3-geometry. This leads to a semiclassical analysis of quantum amplitudes, involving transverse modes, ghost modes, coupled normal and longitudinal modes, and the decoupled normal mode of Maxwell theory.

Paper: gr-qc/9507050

From: Luis Octavio Pimentel Date: Tue, 25 Jul 1995 11:19:44 -0500 (CDT) Title: Electromagnetic Field in Some Anisotropic Stiff Fluid Universes Authors: Pimentel L O Report-no: UAMI-AG-95-29

The electromagnetic field is studied in a family of exact solutions of the Einstein equations whose material content is a perfect fluid with stiff equation of state ($p = \epsilon$). The field equations are solved exactly for several members of the family. -----


El Escorial Summer School on Gravitation and General Relativity 1992: Rotating objects and relativistic physics: Proceedings of the El Escorial Summer School on Gravitation and General Relativity (1992) Held at El Escorial, Spain, 24-28 August 1992 / F.J. Chinea, L.M. Gonzalez-Romero, eds. Berlin ; New York : Springer-Verlag, c1993. 302p. LC CALL NUMBER: QC178 .E36 1992

SUBJECTS: Gravitational fields. General relativity. Astrophysics Chinea, F. J. (Francisco Javier), 1949-

Gonzalez-Romero, L. M. (Luis Manuel), 1962- ISBN: 354057364X (Berlin : acid-free paper) : DM90.00 038757364X (New York : acid-free paper) : \$62.00 -----

"Propulsion Techniques: Action and Reaction", Peter J. Turchi, editor, Ohio State University This is the first of three volumes devoted to space propulsion part of a new series of titles with articles taken from the pages of Aerospace America. The three volume collection of over 150 articles rescues the insights, concerns and dreams of dozens of space propulsion experts for the next generation of aerospace scientists and engineers. Written by well-known figures in space propulsion, including Werner von Braun, Martin Summerfield, Ernst Stuhlinger and Jerry Grey, these books provide readily accessible source material for design courses in astronautical engineering. This first volume surveys the technologies of rocketry in the traditional categories of liquid, solid, hybrid, nuclear and electric propulsion. Historical trends and cycles are displayed in each category as articles describe concepts and progress from the early visions of Goddard, Oberth and Tsiolkovsky to proposed (and re-proposed) ideas for advanced space thrusters. In addition to descriptions of rocket engines of various types, including photon and laser propulsion, associated technologies for propellants and space-electrical power systems are discussed.

Spring 1995, 350 pp, illus, Paperback



ISBN 1-56347-115-9
(Available from the AIAA)

I have a book from Russia that may interest you. It's in Russian and it's called "Experimental Gravity", and is jointly authored by a father and son, S.M. Poliakov and O.M. Poliakov. It describes "gyro-gravity" and "ferromagnetic-gravity" and also how to produce gravity. It's 130 pages and contains a lot of pictures and diagrams and equations (that's about all I understand from it). To judge from the pictures the Russians must have conducted a lot of research in this topic (Many different devices are on the pictures). The question is now: How do I get a translator? -- David Jonsson

INTRODUCTION TO EXPERIMENTAL GRAVITONICS Abstract of book by S.M.Poliakov and O.S.Poliakov

The experiments part covers the following subjects: 1. Light-beam curvature and optical-radiation frequency shift is created and investigated in an artificial nonhomogeneous gravitational field.

A new gravitational effect, named "quadrature" frequency shift in the curved light beam is predicted and calculated. 2. Magnetostriiction is at last explained as a secondary gravitational effect. An equation derived for magnetostriiction permits to calculate the magnetostriiction curve.

3. The propagation velocity of gravitational radiation (generated by a laboratory source) was measured for "quadrupole" - 9×10^{20} cm/s or squared light velocity.

4. It was demonstrated that gravitation is only one of NONLINEAR-MECHANICS EFFECT, that can be created in mechanical system or in ferromagnetic.

The book was published at the author's expense in 1991. Most powerful experimental result described in this book is more than 1200 grammes of pulsed G-force. Several mechanical systems and systems using ferrites are detailed here.

Second edition in English is ready for copy process (disket's text). Editors and investors are interested in joint project for publication can get direct contact with Dr. Poliakov by address: Moscow area, 141120, FRIAZINO, 60-let str., 1-167. Phone 7-095-4658822.

Alexander V. Frolov
P.O.Box 37, St.-Petersburg, 193024, Russia.
E-mail: alex@frolov.spb.ru

--
Super-weapon designer Edward Teller has written an article entitled: "Electromagnetism and Gravitation", Proc. Nat. Acad. Sci. USA Vol 74, No 4. P. 2664-2666. Teller's article is referenced in the book by the Poliakov brothers about Russian experiments in gravitation control.

I wrote already about Poliakov's book "Experimental Gravitonics". He wrote it in 1991 and published in Russian. Now he have English version as MSDOS text on diskett. If you wish help for Dr.Poliakov, write for him and buy book in Russian or copy of text in English. Here is more information.

-- Alex Frolov

"Experimental Gravitonics" Spartak M. Poliakov, Oleg S. Poliakov Russia 141120 Moscow area, Friazino, 60-let SSSR str., 1 - 167.

TABLE OF CONTENTS

Chapter 1

New notions of things forgotten long ago @1
Is the "light barrier" penetrable? p.7
@2 Energy relations and the mechanism of "C-barrier" penetration. p.10


Chapter 2

Microstructural models of the photon and electron @1 What do we know about the photon, an electromagnetic-radiation quantum? p.21

@2 Uniquanta parameters p.23

1. Uniquanta spin (postulated) p.23

a. Linear polarization Fig.8a p.25




b. Circularly polarized nonrotating photon Fig.8b. p.26 c. Circularly polarized rotating photon. P.26 2. Magnetic moment of a uniquantum (postulated) p.27 3. Uniquantum gravitational mass (postulated) p.27 4. Equivalent charge and radius of a uniquantum. p.27 5. Tangential velocity of uniquantum rotation p.29 6. Uniquantum magnetic field p.30
7. Gravitational constant of the uniquantum p.30 8. Gravitational field of the uniquantum p.31 @3 Photon model p.32
@4 Phenomenological microstructural model of the electron p.35 @5 Derivation of approximate gravitational equations of practical interest p.45

part 2

Experimental verification of new gravitational equations p.55 Chapter 1
Experimental verification of mechanical gravitational equations @1 Problems pertaining to the velocity of gravitational-radiation propagation p.56
@2 Principles of determining the propagation velocity of the unknown radiation from the measured momentum of recoil. p.59 @3 Description of the experimental set-up p.64
@4 Gyroscopic multipole "Buket" (Bouget) p.68 @5 Investigation of the effects of dummy shock rotation p.77 @6 The effect of shock braking of the rotating gyroscope p.79 @7 Quadrupole generator of directional gravitational radiation "Yoilka" (Fir) p.80
1. Swing of dummies p.82
2. Swing of gyroscopes p.83
3. Precession of dummies p.84
4. Precession of gyroscopes p.85
@8 Experimental results for the quadrupole generator p.86 @9 Mathematical model of the quadripole generator p.89 @10 Questions of practical application of the results p.107 Chapter 2
Experimental verification of the natural relation between magnetism and gravitation, corollaries from the microstructural model of electron p.111 @1 Gravitational interpretation of magnetostriction p.114 Experimental results p.117
@2 Magnetostrictive curvature of optical beam p.118 Choice of the material for

magnetogravioptical investigations p.121
Magnetic-gap parameters p.123
Experimental methods p.123
@3 Gravioptical effects in GRT p.126
@4 Gravitational frequency shift of optical relation in a nonhomogeneously -magnetized ferromagnetic material p.128
@5 Quadratic gravioptical effect p.130
@6 Some fantastic possibilities opening for modern fundamental science p.131 @7 Generator of short gravitational pulses (by ferromagnetics) p.134 @8 Problem of gravitational receiver p.137 Conclusion p.139

1. H.E.Puthoff, The energetic vacuum: implications for energy research, Speculations in science and technology, vol.13, No.3, p.247.
2. Thomas Valone, Inertial propulsion... Newsletter of Planetary Association for Clean Energy, vol.7 No.1, p.6-12. Published by PACE, Inc. 100 Bronson Av., Suite 1001, Ottawa, Ontario K1R 6G8, Canada.
3. Re NASA electrostatic levitation experiments and Thomas Townsend Brown's research look the Newsletter of Planetary Association for Clean Energy, vol.7 No.4.p.7. July, 1994. "Electrogravitics developments" reprinted from NEXUS Special.
4. "The Swiss Methernitha-Linden Converter", p.3-6. Space Energy Newsletter, June 1993, vol.4 No.2. Published by Space Energy Association, P.O.Box 11422, Clearwater, FL 34616, USA. My understanding of this electrostatic machine allows development of a simple scheme: self-rotating thanks to electrostatic forces disk and ordinary electrical generator connected with axis of disk.
5. Conception of Edmund Whittaker (papers of 1903-1904) is developed by T.E.Barden in his book "Gravitobiology", published by Tesla Book Co., P.O.Box 121873, Chula Vista, CA 91912, USA.
6. P.D. Ouspensky, A New Model of the Universe, New York, 1971.p.433 in Russian edition of 1993.



7. Example of joint demonstration of "gravity/chronal/over-unit power" effect is invention of Ivan Stepanovitch Filimonenko of 1960. His version of cold fusion system produced: 1. Heat power 2. Motive force without fling back of mass 3. Influence on time-period of half-decay. Article of N.E.Zaev published in "Izobretatel i Razionalizator", Russia, No.1 1995, p.8-9.

8. Alexander V. Frolov, The Application of Potential Energy for Creation of Power, New Energy News, vol.2, No.1, May 1994. Published by Institute for New Energy, P.O.Box 58639, Salt Like City, UT 84158-8639, USA.

9. V.V.Lensky, General for Many-Polarity, Irkutsk, Russia, 1986. Published in Russian by Irkutsk University.

10. N.A.Kozyrev, Selected works, 1991, published by University of St.-Petersburg, in Russian.

= BEARDEN, THOMAS E., AND WALTER ROSENTHAL (1991). "ON A TESTABLE UNIFICATION OF ELECTROMAGNETICS, GENERAL RELATIVITY, AND QUANTUM MECHANICS," 26TH IECEC, VOL. 4, PAGES 487-492, AND ASSOCIATION OF DISTINGUISHED AMERICAN SCIENTISTS.

BEARDEN, THOMAS (1992). "A REDEFINITION OF THE ENERGY ANSATZ, LEADING TO A FUNDAMENTALLY NEW CLASS OF NUCLEAR INTERACTIONS," 27TH IECEC, PAGES 4.303-4.310, AND ASSOCIATION OF DISTINGUISHED AMERICAN SCIENTISTS.

GROTZ, TOBY (1992). "THE USE OF MIRROR IMAGE SYMMETRY IN COIL WINDING, APPLICATIONS AND ADVANTAGES IN MAGNETIC FIELD GENERATION," 27TH IECEC, PAGES 4.311-4.313.

HATHAWAY, GEORGE D. (1991). "FROM ANTI-GRAVITY TO ZERO-POINT ENERGY: A TECHNICAL REVIEW OF ADVANCED PROPULSION CONCEPTS," AIDAA/AIAA/DGLR/JSASS 22ND INTERNATIONAL ELECTRIC PROPULSION CONFERENCE, VIAREGGIO, ITALY.

LAVIOLETTE, PAUL A. (1991). "SUBQUANTUM KINETICS: EXPLORING THE CRACK IN THE FIRST LAW," 26TH IECEC, VOL. 4, PAGES 352-357.

SURGALLA, LYNN A. (1991). "NONLINEAR DYNAMICS: MATHEMATICAL PHYSICS FOR 21ST CENTURY TECHNOLOGY (A TUTORIAL FOR ENGINEERS)," 26TH IECEC, VOL. 4, PAGES 394-399.

VALONE, THOMAS (1991). "NON-CONVENTIONAL ENERGY AND PROPULSION METHODS," 26TH IECEC, VOL. 4, PAGES 439-444.

DEPALMA, BRUCE (1991). "MAGNETISM AS A DISTORTION OF A PRE-EXISTENT PRIMORDIAL ENERGY FIELD AND THE POSSIBILITY OF EXTRACTION OF ELECTRICAL ENERGY DIRECTLY FROM SPACE," 26TH IECEC, VOL. 4, PAGES 429-432.

VALONE, THOMAS (1991). "THE ONE-PIECE FARADAY GENERATOR: RESEARCH RESULTS," 26TH IECEC, VOL. 4, PAGES 473-478.

AUTHOR :Znidarsic, Frank
 TITLE :Elementary antigravity / Frank Znidarsic. LANGUAGE :ENGLISH
 PUBLISHED :New York : Vantage Press ; 1989 PHYSICAL DESC :53 p. ; 21 cm.
 SUBJECT :Gravitation, Antigravity

Title: THE SOURCE OF INERTIAL AND GRAVITATIONAL MASS Author: Frank

Znidarsic P.E. FZNIDARS@GPU.COM, ELECTRICAL ENGINEER WITH THE PENN ELECTRIC CO. JOHNSTOWN, Pa. COMMENTS: 8 PAGES ASCII TXT FILE. NEW ENERGY PROJECT, Submitted to the Canadian Journal of Physics, July 94 resubmitted Jan 95 Texted in the DOS editor. (Available at Elektromagnum web site)

Title: GENESIS OF AND ZERO POINT ENERGY

Author: FZNIDARSIC@GPU.COM, F. Znidarsic Electrical Engineer with the Pennsylvania Electric Co. Johnstown Pa. Comments: ASCII *.TXT, 5 PAGES, Texted on DOS editor submitted to the Canadian Journal of Physics July 94, Resubmitted Jan 95

Report_no: Special Energy Prog. (Available at Elektromagnum web site)

The Jackson text is entirely wrong if it does discuss magnetic fields without a potential - the electromagnetic 4-potential A always applies to the conservation of 4-momentum (energy and momentum) by electromagnetic interactions.

See GRAVITATION by Misner, Thorne, and Wheeler.

The causative agent of gravitational gradients (Potential) was first enumerated G. L. Le Sage in 1784. H. A. Lorentz and G. H. Darwin evaluated Le Sage's postulate mathematically and rejected it when, as a result of their evaluation, it was found that although the postulated process could describe all observed gravitational phenomena (and inherently results in GR), the mechanism REQUIRED a continuous overall LOSS of energy (apparently disappearing into matter, in a clear violation of the conservation of energy). I could go on to suggest that mining this concept and process could lead one to significant "pay dirt" in terms of understanding how GR relates to QM.

The kinetic model of Le Sage does lead to a valid model for physical processes. The key is Super-Fluid theory. Both GR and QM are fully

described as well as why each dominates at the scales they do. Helmholtz proved that a superfluid vortex ring is infinitely stable and without a boundary discontinuity can not be created or destroyed. Kelvin mathematically proved that the equations that describe small linear disturbances in a superfluid vortex sponge are IDENTICAL to the equations that describe the propagation of light through space. A EXCELLENT book on this topic is "A History of the Theories of Aether and Electricity" by Sir Edmund Whittaker, Dover Reprint 1989 and I can not give any reference a higher recommendation. Why is this information not known, that sir is a very long and interesting story. - Paul Stowe

For a much more current model, than LeSage, of a sink-source interpretation of gravity, see O.C. Hilgenberg's "Gravitation, Tromben, und Wellen in bewegten Medien" (1931), GieSmann & Bartsch. It's in German. It's in the National Union Catalogue. If you don't read German, then Carl Frederick Krafft's "Ether and Matter" (1945), Dietz Printing Co., contains some portions translated from Hilgenberg's work. Hilgenberg developed a quantum numbering system of the atoms based upon Krafft's ether-vortex atom model, entitled "Quantenzahlen, Wirbelring-Atommodelle und Heliumsechserring- Aufbauprinzip des Periodensystems der chemischen Elemente" Which means roughly: A Quantum Number, Vortex Atom model and Hexagonal-ring construction principle of the periodic system of the chemical elements. Krafft saw the quantization of energy as a logical consequence of a system of particles consisting of, basically, pumps, which could take in and give out energy and ether/space at limited rates, based on rotations of the various rings.

In his model, as opposed to LeSage's, the flux terminates in what he called equipotential zones, much as one would see if two jets of water were to collide coaxially. So, the intake of two bodies "squirting" out ether, will be on the back sides of both bodies, and hence, they are pushed together. LeSage saw each body



acting as shields to the ultramundane particles racing in all directions in space (cosmic rays foreseen in the 18th century!) Thus the two bodies move into each others 'shadow'.

Louis Kevran's work on low energy transmutations of the elements was predicted by Krafft's model. And there was a man named Nemos who claimed to have developed a television type microscope not limited by the optical paths of standard microscopes, and I have a photo he took which shows, it is claimed, the nucleus of an iron atom, as a collection of vortices arranged peripherally (non-coaxially). It's like looking down on pearl-like smoke rings connected by vibrating jets. It's an amazing photo(?) Krafft's prediction was that the proton and electron would be double vortex structures, and the neutron a triple vortex structure. The picture seems to confirm that. So much for my two bits. The work of Krafft is not heavily laden with math. The beauty of it to me is it can be visualized. (Note: Nemo's "microscope" may be based on the technology very recently developed which is known as magnetic microscopy.)

I have a neat little book written by Carl Krafft, back in the 30s. He was an avid developer of an ether-vortex theory of atomic and gravitational forces. His theory was underpinned by the writings of O.C. Hilgenberg and Hermann Fricke of Germany, pre-Nazi. Hilgenberg's views included a vertical ether sink as the cause of gravity, and the consequent development of mass in the exterior of the earth. Hence the earth expanded over time, with periodic explosive expansions and contractions. Meanwhile, Krafft developed the idea of combining vortices in face to face rolling contact, which provides four basic forms: single vortex=neutrino; double vortex, with rolling contact drawing ether into the periphery-proton; with rolling contact into the poles-electron; then the neutron which is three vortices combining forming a neutral, polarizable particle drawing ether into one end, out the sides, in the sides out the other end. Krafft's books were all self published.

-- Roger Cathey

Is there anyone out there who is familiar with modern Kaluza-Klein theory? Or even with the kind Kaluza got Einstein to buy into in 1921 or whenever it was...? This is supposed to unify gravity and electromagnetics in a five-dimensional model. Would it not predict the kind of effects Brown was experimenting with, if true?

-John Sangster

Kaluza and Klein had the idea of extending GR to 5 dimensions. When they did Maxwell's eqns. just sort of pop out. Unfortunately the weak and strong nuclear forces don't. So people try expanding GR into 10, 11 or even 26. It's interesting to do the reverse. Expand Maxwell's eqns. into 5 dimensions. I did it and soon I'll (hopefully) give some details. But essentially you get two out of three of Newton's gravitational equations. You get something similar to the third eqn. but I haven't convinced myself that mine is a generalization of Newton's.

-Ray Cote, KSC

Kaluza-Klein theories are an attempt to give a general relativistic explanation for all the forces of nature not just gravity. The original Kaluza-Klein idea was hatched in 1920 by Kaluza and in 1926 by Klein. The basic idea is that there are 4 space dimensions and 1 time dimension (a 5 dimensional manifold) rather than the 3 space dimensions and 1 time dimensions that there appear to be. Then generalizing Einsteins field equations to this 5 dimensional space (and making the assumption that one of the dimensions gets "curled" up or "compactified") you find that you get Einsteins 4D field equations plus Maxwell's equations. The way that this happens is one of the most beautiful aspects of this theory, and it really makes you (or at least it makes me) think that there should be at least some aspect of this theory that has some correspondence in reality.

Kaluza-Klein theories were revived in the mid seventies by A. Chodos (I think) who showed how you can incorporate all the interactions that we know about now (the strong and the weak

forces which weren't unified with gravity in Kaluza's original theory, which only unified EM and gravity) by taking spacetime to be 11 dimensional (10 space dimensions and 1 time).

I can't really think of a good laymans intro to Kaluza-Klein, but there is a Physics Report in 1985 and 1986 that gives a review of Kaluza-Klein theories by Bailin and Love (?). And there is a Frontiers of Physics book edited by T. Appelquist which reprints most of the important articles on the subject. - Doug Singleton

"The Possibly Unifying Effect of the Dynamic Theory", May 1983, by P.E. Williams

This is part of a series of works by Williams. The novel aspect of the work it that Williams starts from thermodynamics instead of the usual general relativistic and Newtonian approach. Williams develops the 5-D field equations and the neo-coulombic potential. The equations allow for inductive coupling between the electric and gravitational fields.

-- Dennis Cravens

Kaku, Michio.

Hyperspace : a scientific odyssey through parallel universes, time warps, and the 10th dimension / Michio Kaku. New York : Oxford University Press, LC CALL NUMBER: QC793.3.F5 K35 1994 *CIP SUBJECTS: Kaluza-Klein theories. Superstring theories. Hyperspace. ISBN: 0195085140 (alk. paper)

THE INTERACTION OF MAGNETIZATIONS WITH AN EXTERNAL ELECTROMAGNETIC FIELD AND A TIME-DEPENDENT MAGNETIC AHARONOV-BOHM EFFECT Joint Inst. for Nuclear Research, Dubna (USSR). AFANASEV, G.N. NELHIEBEL, M. STEPANOVSJKIJ, YU. P. AB(Technische Univ., Vienna, Austria.) AC(Academy of Sciences of the Ukraine, Kharkov, Ukraine.) 1994 20 PAGES DE95-613463 JINR-E-2-94-

297 Avail: CASI HC A03/MF A01 (US Sales Only) We investigate how the choice of the magnetization distribution inside the sample affects its interaction with the external electromagnetic field. The strong selectivity to the time dependence of the external electromagnetic field arises for the particular magnetizations. This can be used for the storage and ciphering of information. We propose a time-


dependent Aharonov-Bohm-like experiment in which the phase of the wave function is changed by the time-dependent vector magnetic potential. The arising time-dependent interference picture may be viewed as a new channel for the information transfer.

CASI Accession Number: N95-30368

AUTHOR :Doughty, Noel A. (Noel Arthur)
TITLE :Lagrangian interaction : an introduction to relativistic symmetry in electrodynamics and gravitation / Noel A. Doughty. PUBLISHED :Sydney ; Readwood City, Calif. : Addison-Wesley, c1990. DESC :xix, 569 p. : ill. ; 23 cm.
SUBJECT :Electrodynamics, Gravitation, Relativity, Symmetry

GENERALIZED HALL ACCELERATION FOR SPACE PROPULSION SASOH, AKIHIRO AATohoku University, Sendai, Japan In: International Symposium on Space Technology and Science, 18th, Kagoshima, Japan, May 17-22, 1992. Vols. 1 & 2 . A95-82299 Tokyo, Japan ISTS Editorial Board 1992 6 PAGES 1992 p. 403-408

The operation characteristics of electric propulsion devices which utilize Hall effect have been generalized. The electrostatic acceleration is enhanced by thermoelectric effect; an ion kinetic energy can be higher than that associated with the electrostatic potential. Depending on the extent of this effect, there exist two acceleration modes, an electrostatic and an electrostatic/electrothermal hybrid one, the latter characterized by low voltage.



ABSTRACTS OF THE AMERICAN MATHEMATICAL SOCIETY:

vol. 12 (1991)p.572 Abstract *91T-81-200 by Alexander Abian: "The inertia of Time and the energy spent on moving Time forward"

vol. 13 (1992)p.344 Abstract *92T-81-79 by Alexander Abian: "The universal Time"

vol. 15 (1994)p.437 Abstract *94T-81-92 by Alexander Abian "Time has inertia. Equivalence of Time and mass"

vol. 15 (1994)p.585 Abstract *94T-81-164 by Alexander Abian "Time has inertia. Equivalence of Time and mass. How to measure the mass of Time"

Look up a paper published by Miguel Alcubierre in Classical and Quantum Gravity 11 (1994) pp. L73-L77. It's titled "The Warp-Drive: Hyper-Fast Travel within General Relativity". If you can follow a lot of math (or at least the gist of it) it is fascinating. -Ian McBride

"The Warp drive: hyper-fast travel within general relativity" by: Miguel Alcubierre
Department of Physics and Astronomy,
University of Wales, College of Cardiff, PO Box 913, Cardiff CF1 3YB, UK

Article taken from the May 1994 issue of "Classical and Quantum Gravity", a scientific magazine which you are not likely to find at your local newsagent.

ABSTRACT. It is shown how, within the framework of general relativity and without the introduction of wormholes, it is possible to modify a spacetime in a way that allows a spaceship to travel with an arbitrarily large speed. By a purely local expansion of spacetime behind the spaceship and an opposite contraction in front of it, motion faster than the speed of light as seen by observers outside the disturbed region is possible. The resulting distortion is reminiscent of the 'warp drive' of science fiction. However,

just as happens with wormholes, exotic matter will be needed in order to generate a distortion of spacetime like the one discussed here. -----

Anomalous Info Nexus

S P A C E D R I V E S PO Box 228
Kingston Springs, TN

Introductory Reading List U S A
37082-0228

Anomalous Info Nexus, 615.952.5638,
3/12/24/96/14.4 Kbps v.32bis, for Space Drive
Info, Files, and graphics.

Correy, Lee, STAR DRIVER, Del Ray Books,
New York, 1976, # 28994 (Fiction)

Clarke, Arthur C., PROFILES OF THE FUTURE, Bantam Books, New York 1964, # H2734, pp. 46-60, 235

Clarke, Arthur C., RENDEVOUS WITH RAMA, Ballantine Books New York 1974, # 24175, pp. 113-4, 207-8, 265-6 (Fiction)

Davis, William O., Jr., "The Fourth Law Of Motion," ANALOG, May 1962, pp. 83-104

Dean, Norman L., "System For Converting Rotary Motion Into Unidirectional Motion," (Dean Drive) U.S. Patent # 2,886,976

Electric Spacecraft Journal, 73 Sunlight Dr, Leicester, NC 28748, 704.683.0313 Voice / 704.683.3511 FAX / 615.952.5638 BBS Published since 1991, Quarterly, Subscription Rate: \$24/Yr (Only U.S. publication dedicated to Space Drives R&D)

Forward, Robert L., "Spin Drive To The Stars," ANALOG, Apr 1981, pp. 64-70

Harrison, Harry, THE DALETH EFFECT, Berkley SF Books, New York, 1977, # S188O, (Fiction)

Kidd, Alexander D. (Aka Sandy), "Gyroscopic Apparatus", U.S. Patent # 5,024,112

Pournelle, Jerry, A STEP FARTHER OUT, Ace Books, New York, 1983, #78586, pp. 170-187, 229-238

Sellings, Arthur, THE QUY EFFECT, Berkley SF Books, New York, 1967, # X1350, (Fiction)

Stine, G. Harry, "Detesters, Phasers, and Dean Drives," ANALOG, Jun 1976, pp. 68-80

Thornson, Brandson R., "Apparatus For Developing A Propulsive Force," U.S. Patent # 4,631,971

David Jonsson Voice&Fax +46-18-24 51 52
P.O Box 353 Postal giro 499 40 54-7
S-751 06 UPPSALA Internet E-mail
t89djo@tdb.uu.se SWEDEN ++++++Cold
EMISSION before the end of the
century+++++

AUTHOR: Terletskii, IAkov Petrovich,
1912-

Paradoksy teorii otnositelnosti.
Russian/English TITLE: Paradoxes in the
theory of relativity, by Yakov P.


Terletskii. With a foreword by Banesh
Hoffmann. PUBL.: New York, Plenum
Press, 1968

NOTES: Translation of Paradoksy teorii
otnositelnosti.

QUANTUM FLUCTUATIONS AND
SEMICLASSICAL GRAVITY THEORY
PH.D. THESIS Tufts Univ., Medford, MA.
KUO, CHUNG-I. 1994, 119 PAGES Avail:
Univ. Microfilms Order No. DA9419336
Semiclassical gravity theory should serve as a
working model before the final theory of
quantized gravity is known, or as an
approximation for manageable calculations
even when the final theory is known. We deal
with the important issue of the applicability of
the semiclassical theory of gravity, specifically
considering the effects of quantum fluctuations

of the matter fields and the induced metric
perturbations. The quantum fields with
negative energy densities are proposed to be
the cases where the semiclassical theory is no
longer valid. We start with a discussion of the
basic notions and developments of
semiclassical gravity theory, and continue with
a discussion of the establishment of a
meaningful measurement of the deviation from
semiclassical theory. The measure is a
normalized dispersion of the energy density.
The non-positive-definiteness of the energy
density of quantum fields is derived and
discussed. Important cases, like squeezed
states and the Casimir effect, which exhibit
negative energy densities, are discussed and
the deviations from semiclassical theory are
checked. A test particle method using a
generalized Langevin equation is formulated
for the physical description of systems for
which the semiclassical theory can not be used.
Quantum fields around straight infinite cosmic
string is another example of where the negative
energy densities may arise. We examine the
validity of semiclassical theory for this case.
The Casimir force due to the zero-point
fluctuations of the electromagnetic fields in the
presence of a conducting plate is another case
where the quantum fluctuations are large and
naive classical consideration should be
modified. Using the test particle method, we
are able to show that it is a relaxation
phenomenon and that a notion of effective
temperature can be associated with it. CASI
Accession Number: N95-29527

ELECTROMAGNETIC-GRAVITATIONAL
CONVERSION CROSS SECTIONS IN
EXTERNAL ELECTROMAGNETIC FIELDS
International Centre for Theoretical Physics,
Trieste (Italy). LONG, HOANG NGOC SOA,
DANG VAN TRAN, TUAN A. SEP. 1994 11
PAGES DE95-613589 IC-94/285 Avail: CASI
HC A03/MF A01 (US Sales Only) The
classical processes: the conversion of photons
into gravitons in the static electromagnetic
fields are considered by using Feynman
perturbation techniques. The differential cross
sections are presented for the conversion in the
electric field of the flat condenser and the



magnetic field of the solenoid. A numerical evaluation shows that the cross sections may have the observable value in the present technical scenario. CASI Accession Number: N95-30637

THE PHYSICS OF TACHYONS. 3:
TACHYON ELECTROMAGNETISM
DAWE, ROSS L. HINES, KENNETH C.
University of Melbourne, Parkville, Australia
1994 34 PAGES Australian Journal of Physics
(ISSN 0004-9506) vol 47, no 4 1994 p 431-
464 Research supported by the ARC and the
University of Melbourne HTN-95-01061
A new formulation of the theory of tachyons
using the same two postulates as in special
relativity is applied to electro-magnetism.
Tachyonic transformations of the
electromagnetic fields E and B are rigorously
derived from Maxwell's equations and are
shown to be the same as for bradyonic
transformations. Tachyonic tranformations of
current density, charge density, scalar and
vector potentials are also derived and
discussed. Tachyonic optics and the four-
potential of a moving tachyonic charge are also
discussed, along with generalized four-vector
transformations and electromagnetic four-
tensors in extended relativity. Use is made of a
switching principle to show how tachyons
automatically obey the law of conservation of
electric charge in any inertial reference frame,
even though the observed tachyon electric
charge is not an invariant between observers.
The electromagnetic field produced by a
charged tachyon takes the form of a Mach
cone, inside which the electromagnetic field is
real and detectable, while outside the cone the
field generated by the tachyon is imaginary
and undetectable.

CASI Accession Number: A95-90247

Franklin, Allan, 1938-
The rise and fall of the "Fifth Force" :
discovery, pursuit, and justification in modern
physics / Allan Franklin. New York :
American Institute of Physics, c1993. 141 p.
ill. ; 25 cm. LC CALL NUMBER: QC6 F673
1993

SUBJECTS: Physics--Methodology.
Gravitation.

Michlo, George, 1942-
The push of gravity / George Michlo ;
illustrated by Warwick Humphries. 1st ed.
New York : Vantage Press, c1993. xv, 101 p. :
ill. ; 24 cm. LC CALL NUMBER: QC178
.M49 1993
SUBJECTS: Gravitation.
ISBN: 0533091330


Doughty, Noel A. (Noel Arthur)
Lagrangian interaction : an introduction to
relativistic symmetry in electrodynamics and
gravitation / Noel A. Doughty. Sydney ;
Readwood City, Calif. : Addison-Wesley,
c1990. xix, 569 p. : ill. ; 23 cm. LC CALL
NUMBER: QC631 .D68 1990
SUBJECTS: Electrodynamics. Gravitation.
Relativity. Symmetry. ISBN: 0201416255
(U.S.) : \$33.95

Alexander, S.
Gravity and inertia : the mechanism / by S.
Alexander. Santa Barbara, Calif. : G.E.C.
Research, c1985. 64 p. : ill. ; 23 cm. LC CALL
NUMBER: QC178 .A44 1985
SUBJECTS: Gravitation. Inertia (Mechanics)
ISBN: 0939525054

Harrigan, Gregory Leo, 1919-
The great gravity myth / Gregory Leo
Harrigan. 2nd ed., rev. and enl. Minneapolis :
Shanty Press, 1991. p. cm. LC CALL
NUMBER: QC178 .H28 1991 *CIP - NOT
YET IN LC* SUBJECTS: Gravitation--
Miscellanea. Serendipity in science. ISBN:
0916403033 (lib. bdg.) : \$8.95

Soldano, B. A.
A new look at Maxwell's equations and the
permittivity of free space / by B.A. Soldano.
Greenville, S.C., U.S.A. : Grenridge Pub.,
1982. 50 p. LC CALL NUMBER: QB341 .S65
1982
SUBJECTS: Gravitation. Maxwell equations.

Gallimore, J. G.
Transverse paraphysics : the new science of
space, time, and gravity control J.G.



Gallimore. Millbrae, Calif. : Tesla Book Co., 1982. ix, 359 p. LC CALL NUMBER: QC173.59.S65 G35 1982
SUBJECTS: Space and time. Gravitation.

Mancini Ridolfini, Niccolo.
Elettricità e magnetismo; rotazione elettromagnetica gravitazionale. Bologna, L. Cappelli, 1931. vii, 506 p. illus. 25 cm. LC CALL NUMBER: QC518.M36
SUBJECTS: Electromagnetic theory. Space and time. Gravitation.

King, Moray B.
Tapping the zero-point energy / Moray B. King. Provo, UT : Paraclete Pub., c1989. iii, 169 p. : ill. ; 22 cm.
LC CALL NUMBER: QC178.K5575 1989
SUBJECTS: Antigravity. Radiation. Electromagnetics. Force and energy. ISBN: 0962335606 : \$9.95

The Large N expansion in quantum field theory and statistical physics : from spin systems to 2-dimensional gravity / editors, Edouard Brezin, Spenta R. Wadia. Singapore ; River Edge, NJ : World Scientific, c1993. xiv, 1130 p. LC CALL NUMBER: QC174.45.L37 1993

SUBJECTS: Quantum field theory. String models. Gauge fields (Physics) ISBN: 9810204558

Magnetic susceptibility of superconductors and other spin systems / edited by Robert A. Hein, Thomas L. Francavilla, and Donald H. Liebenberg. New York : Plenum Press, c1991. xx, 606 p. : ill. ; 26 cm. LC CALL NUMBER: QC611.97.M34 M34 1991

SUBJECTS: Superconductors--Magnetic properties--Magnetic susceptibility United States. Office of Naval Research. Office of Naval Research Workshop on Magnetic Susceptibility of Superconductors and Other Spin Systems (1991 : Coolfont, W. Va.) "Proceedings of the Office of Naval Research Workshop on Magnetic Susceptibility of Superconductors and Other Spin Systems, held May 20-23, 1991, in Coolfont, Berkeley Springs, West Virginia"--T.p. verso. ISBN: 0306441977

Spin waves and magnetic excitations / volume editors, A.S. Borovik-Romanov, S.K. Sinha. Amsterdam ; New York : North-Holland ; New York, N.Y. : Sole distributors for the U.S.A. and Canada, Elsevier Science Pub. Co., 1988. LC CALL NUMBER: QC762.S66 1988
SUBJECTS: Spin waves. Magnons. Dielectrics--Magnetic properties. Metals--Magnetic properties. Modern problems in condensed matter sciences ; v. 22 ISBN: 0444870687 (v. 1)

Aono, Osamu, 1937-
Rotation of a magnetic field / Osamu Aono and Ryo Sugihara. Nagoya, Japan : Institute of Plasma Physics, Nagoya University, 1986. 6 p. ; 30 cm. LC CALL NUMBER: QC717.6.N35 no. 792 (ALTERNATE CLASS QC754.2.M3)
SUBJECTS: Magnetic fields. Electrodynamics. Research report (Nagoya Daigaku. Purazumu Kenkyūjo) ; IPPJ-792.

Handbook of electron spin resonance : data sources, computer technology, relaxation, and ENDOR / edited by Charles P. Poole, Jr. and Horacio A. Farach. New York : American Institute of Physics, c1994. x, 660 p. : ill. ; 25 cm. LC CALL NUMBER: QC762.H32 1994
SUBJECTS: Electron paramagnetic resonance. Electron nuclear double resonance spectroscopy. Relaxation phenomena. ISBN: 1563960443 (acid-free)

Mims, W. B.
The linear electric field effect in paramagnetic resonance / W. B. Mims. Oxford : Clarendon Press, 1976. 339 p. : ill. ; 24 cm. LC CALL NUMBER: QC762.M55
SUBJECTS: Electron paramagnetic resonance. Electric fields. ISBN: 0198519443 : L9.75

Morrison, Clyde A. (Clyde Arthur), 1926-Angular momentum theory applied to interactions in solids / C.A. Morrison. Berlin ; New York : Springer-Verlag, c1988. 159 p. ; 25 cm. LC CALL NUMBER: QD475.M68 1988

SUBJECTS: Crystal field theory. Angular momentum. ISBN: 0387189904 (U.S. : pbk.)

----- Mirman, R.

Massless representations of the Poincare Group : electromagnetism, gravitation, quantum mechanics, geometry / R. Mirman. Commack, N.Y. : Nova Science Publishers, 1995. p. cm.

LC CALL NUMBER: QC20.7.G76 M57 1995
 CIP - NOT YET IN LC SUBJECTS: Representations of groups. Poincare series. Electromagnetism. Gravitation. Quantum theory. Geometry. Mathematical physics. ISBN: 1560722592

Antunez de Mayolo, Santiago, 1887-1967. The neutral element base of matter and probable cause of gravitation / Santiago Antunez de Mayolo. Lima, Peru : Universidad Nacional Mayor de San Marcos, 1948. 36 p. : ill. ; 22 cm.

LC CALL NUMBER: MLCS 94/12050 (Q)
 SUBJECTS: Nuclear physics. Matter-Constitution. Translation of "Work presented in Spanish to the IV. South American Chemistry Congress, Santiago, Chile, March 1948."

Ciufolini, Ignazio, 1951-
 Gravitation and inertia / Ignazio Ciufolini and John Archibald Wheeler. Princeton, N.J. : Princeton University Press, c1995. xi, 498 p.
 LC CALL NUMBER: QC173.59.G44 C58 1995

SUBJECTS: Geometrodynamics. General relativity. Gravitation. Inertia. ISBN: 0691033234 (acid-free paper)

TITLE: Excalibur Briefing
 AUTHOR: Thomas E. Bearden
 COPYRIGHT DATE: 1980, 1988
 PUBLISHER: Strawberry Hill Press/A Walnut Hill Book ISBN# 0-89407-060-6
 PURCHASED FROM: Tesla Book Co. or Fry's INC. INQ. COMMENTS

According to the front and rear covers this book explains paranormal phenomena and the interaction of mind and matter. There are 4 chapters plus a glossary and bibliography. 332 pages, 42 photographs, and 40 illustrations. Chapter one is called... A Sampling of Specific Paranormal Phenomena. Some of the subjects in this chapter are...Remote Viewing The

Moray Radiant Energy Device, Thought Photography, Pavlita's Psychotronic Generators, UFO's, Kirlian Photography, Psychic Surgery Chapter Two is called...A Theoretical Background for Understanding PT, UFO's and PSI Phenomena...Some of the subjects are.... Unexplained Mysteries of Physics, Two Slit Experiment, Radionics, Biofields and Maverick Worlds Chapter Three is called...New Military Applications of PSI Research Some of the subjects covered are...Background to Psychotronic Research in the U.S and the U.S.S.R., Radiation of the U.S. Embassy, Hyperspace Howitzer operation, Virtual States and Hyperspaces, Feynman diagrams The Neurophone, Soviet Woodpecker signals The last chapter covers Soviet Phase Conjugate Directed Energy Weapons (Weapons that use time reversed Electromagnetic Waves) The Glossary is about 30 pages long and is very useful.

Bearden, T. E. (Thomas E.), 1930-
 Excalibur briefing / Thomas E. Bearden ; foreword by John White ; special drawings by Hal Crawford. San Francisco : Strawberry Hill Press, c1980. LC CALL NUMBER: BF1999 .B387 1980
 SUBJECTS: Occultism. Unidentified flying objects. ISBN: 0894070150 (pbk.) : \$8.95

TITLE: Magnetism:An Introductory Survey
 AUTHOR: E.W. Lee
 COPYRIGHT DATE: 1963,1970
 PUBLISHER: Dover Publications Inc. New York ISBN# 0-486-24689-2
 PURCHASED FROM: Lindsay Publications
 COMMENTS
 Paperback, 280 pages, Some photographs and Illustrations Some of subjects covered are....Atomic Theory of Matter, Earth's Magnetism History of Magnetism, Magnetism in Scientific Research, Paramagnetism and Diamagnetism

TITLE: Tesla: The Lost Inventions
 AUTHOR: George Trinkaus

COPYRIGHT DATE: 1988
 PUBLISHER: High Voltage Press
 ISBN# N/A
 PURCHASED FROM: Lindsay Publications
 COMMENTS
 Paper, 33 Pages, 42 Illustrations. Describes Tesla's lost inventions in plain, easy to understand English. According to the author, patents are hard to understand. In the illustrations he shows the patent number. Some of the inventions include... Disk Turbine
 Rotary Engine
 Magnifying Transmitter
 Transport
 Free Energy Receiver

----- TITLE: Tesla: Man out of Time
 AUTHOR: Margaret Cheney
 COPYRIGHT DATE: 1981
 PUBLISHER: Laurel Book by Dell Publishing Co. ISBN# 0-440-39077-X
 PURCHASED FROM: Tesla Book Co. or Lindsay Publications COMMENTS
 Paperback 320 Pages, 8 Pages of Rare Photographs Good biography of Tesla. 30 Chapters plus Reference Notes. Chapter 29 deals with Tesla's papers and what may have happened to them after he died.

----- TITLE: The Cosmic Conspiracy
 AUTHOR: Stan Deyo
 COPYRIGHT DATE: 1978
 PUBLISHER: West Australian Texas Trading ISBN# 0-908477-00-7
 PURCHASED FROM: Tesla Book Co.
 COMMENTS
 Paperback 200 Pages. The book is divided into 3 sections with each section containing at least 6 chapters. Plus there are 7 Appendices. At the end of each section there is a Suggested reading list for that section. Section One deals with research into Electro- Gravitic Propulsion Tesla, Weather Warfare, Conspiracies. Section Two deals with Mysticism and Numerology, mystery schools, Illuminati Section Three deals with Religious Ideas In the appendices there are articles on Townsend Brown, Einstein's Relativity error, Michelson Morley experiment, Electro-Dynamic Propulsion, Practical Ion Craft. Very fascinating book.


----- TITLE: The Philadelphia Experiment: Project Invisibility AUTHOR: William L. Moore, Charles Berlitz COPYRIGHT DATE: 1979
 PUBLISHER: Fawcett Crest New York
 ISBN# 0-449-24280-3
 PURCHASED FROM: Waldenbooks
 COMMENTS
 Paperback 288 pages.
 Book on the alleged Navy experiment to make a ship invisible; to radar or optically or both. Supposedly the ship not only became invisible but dematerialized and rematerialized at a distant location, then re-materialized at the original location (Philadelphia Navy Yard). Plus there were severe side effects to the crew members. Some of the crew were said to have disappeared into another dimension. Some never to return. Chapter 9 (The Unexpected Key) is very interesting because it describes an interview with a scientist who was involved with the Philadelphia experiment when it was being planned.

The best introduction to dyads and dyadic analysis in electromagnetism in my opinion is the following book:

Hollis C Chen, Theory of Electromagnetic Waves: A Coordinate-Free Approach (1983, McGraw-Hill; 1992, TechBooks).

It's already a proven fact that angular momentum will generate an opposing force to gravity in the way you describe. This is a purely General Relativistic effect. There's an article in the 1988 Foundations of Physics "An Exact Solution to Einstein's Field Equations: Gravitational Force Can Also Be Repulsive!"

It requires an immensely huge angular momentum to get any decent repulsion, such that you're not going to get it by any mechanical means



For those of you who are not familiar with the obscure aspects of General Relativity, hopefully this will steer you in the right direction for further research and knowledge. Non-Newtonian gravitational fields, which may be either attractive or repulsive, can be generated from three effects. These are that of rotating masses, moving masses, or fluctuating masses relative to a stationary, non-rotating body. These effects are similar to centrifugal, Coriolis, and other inertial forces and were first described by W. de Sitter in 1916 and Hans Thirring in 1918. Dr. Robert L. Forward published his Guidelines to Antigravity in March 1963 in the American Journal of Physics. Dr. Forward is an expert in General Relativity and Gravity Research and studied under Weber at the University of Maryland. In his guidelines article, he discusses the dipole effect of gravity as predicted by General Relativity. Unfortunately, the forces generated are extremely weak without very dense mass or extremely high angular velocities. I suggest that everyone with an interest in such aspects obtain a copy of this article and read it through before passing any judgements as to these forces existing or being generated!

-- Phillip Carpenter

Might a mass (gravitational charge) in motion also produce another type of field much like a magnetic one?

Something like this is "gravitomagnetic effect" is theoretically predicted. If you were in such a field, it would simply give the impression that you were in a locally rotating frame of reference, so moving objects would experience coriolis forces, even when you were not rotating relative to distant reference points. As the effect is of the order of $v_1 v_2/c^2$ where v_1 is the speed of the gravitational source and v_2 is the speed of the test object, it is extremely small and has not yet been measured.

Note also that a rotating massive object is expected to give rise to a similar field in the same way as a current loop gives rise to a magnetic field. This is known as the Lense-Thirring effect. A first-order Special Relativity approximation (which only applies for a locally inertial frame of reference where space


isn't significantly curved) is simply that the rotation field is $(v_1 \times g)/c^2$ where g is the Newtonian acceleration vector v_1 is the velocity of the source object. The acceleration that field generates for a body moving with velocity v_2 is $v_2 \times (v_1 \times g)/c^2$. Note for comparison that the magnetic field is $B = (v_1 \times E)/c^2$ so the magnetic force is $q v_2 \times (v_1 \times E)/c^2$. The gravitational rotation field calculated in this way is equal to $2w$ where w is the apparent angular velocity of rotation. It is hoped that "conscience-guided" satellite experiments may confirm this effect within a few years, but at present there are too many other disturbances which make it too difficult to measure such a small effect. The rotation field, whether caused by a linearly moving mass or a rotating object, only affects moving masses. However, there is of course a much stronger associated acceleration field which affects all masses. From the subjective point of view, the acceleration field may appear to be partly linear acceleration and partly "centrifugal" force associated with rotary motion, but this is a higher-order effect.

-- Jonathan Scott

Some scientists in Boulder, CO (USA) have succeeded in cooling down matter into the elusive Bose-Einstein condensate. The kinetic energy of the atoms in this state have been removed. If you could maintain this state in stable form and spin it, the angular momentum would repel the earth and lift many times its own mass. Outside of the atmosphere, this could produce the desired gravitational dipole effect.

-

Bonaldi, M., et al., "Inertial and Gravitational Experiments With Superfluids: A Progress Report," Proceedings of the Fourth Marcel Grossmann Meeting on General Relativity, Elsevier Science Publishers B.V., 1985, pp. 1309-1317.



Title: ANGULAR MOMENTUM PARADOXES WITH SOLENOIDS AND MONOPOLES In: Phys.Lett.118B:385,1982
 Date/Source: August 1982
 Fermilab Library: FERMILAB-PUB-82/53-THY -- Preprint -- Available

Title: Long range effects in asymptotic fields and angular momentum of classical field electrodynamics Date/Source: February 1995
 Fermilab Library: CALL NUMBER DESY-95-035 -- Preprint -- Available

Title: Angular Momentum
 Authors: Brink, D. M. (David Maurice), and G.R. Satchler Date/Source: Oxford : Clarendon Press ; New York : Oxford University Press, 1993.
 Fermilab Library: CALL NUMBER QC793.3.A5 B75 1993 -- Book -- Available

AUTHOR(s): Hayasaka, Hideo Takeuchi, Sakae
 TITLE: Gravitation and Astrophysics.
 Summary: Anomalous weight reduction on a gyroscope's right rotations around the vertical axis on the Earth.
 In: Physical review letters.
 DEC 18 1989 v 63 n 25 Page 2701

AUTHOR(s): Starzhinskii, V.M.
 TITLE: An exceptional case of motion of the Kovalevskaia gyroscope.
 In: PMM, Journal of applied mathematics and mechanic 1983 v 47 n 1 Page 134

From: sphinx@world.std.com (John Sangster, SPHINX Technologies) Subject: Weight Reduction in Spinning Masses Date: Fri, 3 Nov 1995 06:04:35 GMT

Recently Hideo Hayasaka and Sakae Takeuchi of the Engineering Faculty at Tohoku University in Japan have published an experimental result of this sort. They found

that gyroscopes spinning clockwise as seen from above, at their location, exhibited a decrease in relative mass of 5.07×10^{-5} and 4.22×10^{-5} respectively for the two gyroscope configurations studied. (Weight was multiplied by $1-e$ where e is the relative factors given above, if I haven't botched up in my arithmetic.) The effect as plotted in the paper I saw appears to be perfectly linear to within reasonable experimental error, thus giving a rotational velocity at which the weight would go to zero which I made out to be 3.27 MHz (million rotations per second) in the first case and 3.95 MHz in the second.

That was with CLOCKWISE rotation as seen from above. With COUNTERclockwise rotation, the same experimental setup showed ZERO EFFECT. Zip. Nada. Nichts. Nyechevo. You get the idea. For one thing, this result makes it almost certain that they are NOT dealing with bad lab technique. Not to mention the fact that they spent nearly a year and a half going over and over their setup and trying to answer all objections by the reviewers of their Physical Review Letters paper (it eventually appeared in PRL (63 2701)). As far as I know, nobody has published a theoretical model that accounts for these observations. The idea of a physical phenomenon that appears only in one direction of rotation is rather unprecedented. I know of only one other mathematical/physical phenomenon that does this, and I'm trying to understand how the two might be related, but without success as yet.

-- John Sangster

Physicist Alex Harvey wrote an article about the Hayakawa-Taguechi experiment. The article was published in:

Nature, Aug 23 1990, Vol 346 Page 705

You'll also find other references there. Harvey shows mathematically that an angular momentum vector aligned antiparallel to the local gravitational field violates the equivalence principle. He also shows that the path of a spinning body under gravity need not be geodesic. Here are two "holes" in GR that seem to account for the behavior of H & T's



gyros. New experiments should be designed to force the asymmetry to appear, as predicted by theory, rather than passively leave the results to chance.

There is a dimensional error of Hayasaka and Takeuchi which CAN be corrected by supplying a quantity that restores proper dimensionality. In simplest terms, H and T's result looks like: { deltaN = - (proportionality constant) m w r } where deltaN is the weight change in Newtons, m is the mass of the rotor in kg, w is the rotation frequency in angular units and r is the radius of the rotor in meters. The units of the missing quantity are radians per second. The rotation, w, has already been counted. The missing quantity is the precession, Wp. With clockwise rotation, the vector J points down the spin axis, while the precession vector, Wp, points up the spin axis. Physicist Alex Harvey, writing about H and T's results, confirmed that there is no (symmetrical) weight gain, no effect at all, with counter-clockwise rotation, J (up). In this case, says Harvey, "[J] is parallel to the gravitational field."

-- laradex3@sj.znet.com

AUTHOR(s): Harvey, Alex

TITLE(s): Complex Transformation of the Kasner Metric.

In: General relativity and gravitation.

OCT 01 1989 v 21 n 10 Page 1021

AUTHOR(s): Harvey, Alex

TITLE(s): Cosmological models.

In: American journal of physics.

OCT 01 1993 v 61 n 10 Page 901

AUTHOR(s): Harvey, Alex

TITLE(s): Identities of the scalars of the four-dimensional

Riemannian manifold.

In: Journal of mathematical physics.

JAN 01 1995 v 36 n 1 Page 356

AUTHOR(s): Harvey, Alex

TITLE(s): Will the Real Kasner Metric Please Stand Up.

In: General relativity and gravitation.

DEC 01 1990 v 22 n 12 Page 1433

>Maybe I've missed it, but I've looked seriously, and there seems to be no information in undergraduate or graduate level physics reference books which mentions the relationship between macroscopic and microscopic angular momentum -- much less provides any analysis or explanation linking quantum angular momentum to macroscopic angular momentum.

You're catching on. The subject of compound angular momentum, or internal and external angular momentum, or intrinsic and extrinsic angular momentum has been a repressed subject for about 2 and half decades. Add to that list, spherical pendulums, Coriolis effect, except as applied to ballistics and meteorology as used by the US military, and Shafer's pendulum, that neat little device used as the artificial horizon of aircraft.

>How does quantum angular momentum become organized from a microscopic to a macroscopic level? Has anyone ever published any work about this? I can't find any.

There isn't any that I know of, though back in the late fifties, there was a fellow named Edward Condon at the University of Colorado who was fairly proficient on the subject. So much so that he wrote the rotational dynamics section, called noninertial dynamics at the time, of the reference "The Handbook of Physics" which he also co-edited (Chapter 5). I don't recall offhand who the publisher was (Harcourt/Brace?), though it was endorsed by the American Institute of Physics. Later, when Mr Condon was the head of the USAF project 'Blue Book', he labored to suppress his own work when the directive was handed down from the Navy's Turtle Island project.

-- James Youlton

In the Barnett effect a long iron cylinder, when rotated at high speed about its longitudinal axis, is found to develop a measurable component of magnetization, the value of which is proportional to the angular speed. The effect is attributed to the influence of the

impressed rotation upon the revolving electronics systems due to the mass property of the unpaired electrons within the atoms. -- Henry Wallace

Barnett, S.J., "Magnetization By Rotation," The American Physical Society, Second Series, vol. VI, No. 2, Jun., 1915, pp. 171-172.

Barnett, S.J., "Magnetization By Rotation," The Physical Review, Second Series, vol. VI, No. 4, Oct., 1915, pp. 239-270.

The Barnett Effect is known to me as the effect of a change in volume of a magnetic material in response to a change in its magnetization strength. If a ferrite material is exposed to a higher magnetization field (more current through the coil) the ferrite will change in volume. I was not aware that this has anything to do with alignment to a spinning axis. For further information about this aspect of the Barnett effect, see: Ref. Handbook of Magnetic Phenomena, by Harry S Burk, Van Nostrand Reinhold 1986 Page 262. -- William Clymer

Magnetic systems with competing interactions : frustrated spin systems / edited by H.T. Diep. Singapore ; River Edge, N.J. : World Scientific, c1994. xiv, 335 p. : ill. ; 24 cm.

LC CALL NUMBER: QC754.2.S75 M34 1994

SUBJECTS: Magnetization. Rotational motion. Spin waves. Ferromagnetism.

CONTENTS:

Nonlinear phenomena and chaos in magnetic materials / P.E. Wigen -- Some nonlinear effects in magnetically ordered materials / H. Suhl -- Spin-wave instability processes in ferrites / M. Chen & C.E. Patton -- Spin-wave dynamics in a ferrimagnetic sphere: experiments and models / P.H. Bryant, D.C. Jeffries, & K. Nakamura -- Spin-wave auto-oscillations in YIG spheres driven by parallel pumping and subsidiary resonance / S.M.

Rezende & A. Azevedo -- Strong chaos in magnetic resonance / M. Warden -- Magnetostatic modes in thin films / R.D. McMichael & P.E. Wigen -- Fractal properties in magnetic crystal / H. Yamazaki -- Spin-wave envelope solitons in magnetic films / A.N. Slavin, B.A. Kalinikos, & N.G. Korshikov. ISBN: 9810210051

Hence the Wilson-Blackett proportionality between the angular momentum of planets, stars etc and their magnetic moment. For more information see Science News Aug 6 '94 p82. -

AUTHOR(s): Bloxham, Jeremy Gubbins, David

TITLE(s): The Evolution of the Earth's Magnetic Field.

Summary: The origin of the field has fascinated more than a dozen generations of physicists. Molten iron in the outer core, driven by convection and influenced by the earth's rotation, acts as a dynamo that generates the field. Now historical records of magnetic-field changes yield new insights into the process and into how the field may behave in the future.

In: Scientific American. DEC 01 1989 v 261 n 6 Page 68

AUTHOR(s): Malov, I.F.

TITLE(s): Angle between the magnetic field and the rotation axis in pulsars.

In: Soviet astronomy.

MAR 01 1990 v 34 n 2 Page 189

AUTHOR(s): Marsheva, N. M.

TITLE(s): Permanent rotation of a heavy rigid body in a magnetic field.

In: Moscow university mechanics bulletin. 1989 v 44 n 1

AUTHOR(s): Vitale, S. Bonaldi, M. Falferi, P.

TITLE: Magnetization by rotation and gyromagnetic gyroscopes.



Summary: We discuss how the general phenomenon of magnetization by rotation may be used probe the angular velocity of the laboratory with respect to a local frame of inertia. We show that gyroscope with no moving parts based on this pheno-
In: Physical review B: Condensed matter. JUN 01 1989 v 39 n 16 p B Page 11993

CONDENSED MATTER THEORY,
ABSTRACT COND-MAT/9509141 From:
Erwin Frey Date: Fri, 22 Sep 1995 09:43:52
+0200

Critical Dynamics of Magnets

Authors: Erwin Frey , Franz Schwabl (TU Muenchen) Comments: Review article (154 pages, figures included) We review our current understanding of the critical dynamics of magnets above and below the transition temperature with focus on the effects due to the dipole--dipole interaction present in all real magnets. Significant progress in our understanding of real ferromagnets in the vicinity of the critical point has been made in the last decade through improved experimental techniques and theoretical advances in taking into account realistic spin-spin interactions. We start our review with a discussion of the theoretical results for the critical dynamics based on recent renormalization group, mode coupling and spin wave theories. A detailed comparison is made of the theory with experimental results obtained by different measuring techniques, such as neutron scattering, hyperfine interaction, muon--spin-resonance, electron--spin--resonance, and magnetic relaxation, in various materials. Furthermore we discuss the effects of dipolar interaction on the critical dynamics of three--dimensional isotropic antiferromagnets and uniaxial ferromagnets. Special attention is also paid to a discussion of the consequences of dipolar anisotropies on the existence of magnetic order and the spin--wave spectrum in two--dimensional ferromagnets and antiferromagnets. We close our review with a formulation of critical dynamics in terms of nonlinear Langevin equations.


Paper: cond-mat/9501029
From: Kazuhiro Kuboki Date: Mon, 09 Jan 1995 10:40:11 EST

Title: Proximity-induced time-reversal symmetry breaking at Josephson junctions between unconventional superconductors
Author: Kazuhiro Kuboki and Manfred Sigrist
We argue that a locally time-reversal symmetry breaking state can occur at Josephson junctions between unconventional superconductors. Order parameters induced by the proximity effect can combine with the bulk order parameter to form such a state. This property is specifically due to the intrinsic phase structure of the pairing wave function in unconventional superconductors. Experimental consequences of this effect in high-temperature superconductors are examined.

Paper: cond-mat/9501088
From: David Benedict Bailey Date: Thu, 19 Jan 1995 11:34:10 -0800 (PST) Title: Gapless Time-Reversal Symmetry Breaking Superconductivity Authors: A. M. Tikofsky and D. B. Bailey We consider a layered superconductor with a complex order parameter whose phase switches sign from one layer to the next. This system is shown to exhibit gapless superconductivity for sufficiently large interlayer pairing or interlayer hopping. In addition, this description is consistent with experiments finding signals of time-reversal symmetry breaking in high-temperature superconductors only at the surface and not in the sample bulk.

Paper: cond-mat/9501133
From: ioffe@physics.rutgers.edu (Lev Ioffe)
Date: Mon, 30 Jan 95 08:59:22 EST
Title: On the spin density wave transition in a two dimensional spin liquid.

Authors: B. L. Altshuler, L. B. Ioffe, A. I. Larkin, A. J. Millis. Strongly correlated two dimensional electrons are believed to form a spin liquid in some regimes of density and temperature. As the density is varied, one expects a transition from this spin liquid state to a spin density wave antiferromagnetic state. In this paper we show that it is self-consistent to assume that this transition is second order



and, on this assumption, determine the critical behavior of the $2p_F$ susceptibility, the NMR rates T_1 and T_2 and the uniform susceptibility. We compare our results to data on high T_c materials.

Paper: gr-qc/9502041

From: Barry Haddow Date: Fri, 24 Feb 1995 18:59:15 (GMT)

Title: Purely Magnetic Spacetimes

Author: Barry Haddow (Trinity College, Dublin, Ireland) Purely magnetic spacetimes, in which the Riemann tensor satisfies $R_{abcd}u^bu^d=0$ for some unit timelike vector u^a , are studied. The algebraic consequences for the Weyl and Ricci tensors are examined in detail and consideration given to the uniqueness of u^a . Some remarks concerning the nature of the congruence associated with u^a are made.

Paper: cond-mat/9502103

From: deb@rri.ernet.in (Debnarayan Jana) Date: Fri, 24 Feb 95 11:23:21+050

Title: Universal Diamagnetism of Charged Scalar Fields Authors: Debnarayan Jana

We show that charged scalar fields are always diamagnetic, even in the presence of interactions and at finite temperatures. This generalises earlier work on the diamagnetism of charged spinless bosons to the case of infinite degrees of freedom.

"CP Violation and Antigravity Revisited", G. Chardin, Nuclear Physics, Jun 7 1993, Vol 558

"Equivalence Principle Violation, Antigravity and Anyons Induced by Gravitational Chern-Simons Couplings", S. Deser, Classical and Quantum Gravity, 1992, Vol 9 Supp

"The Arguments Against Antigravity and the Gravitational Acceleration of Anti-Matter", Michael Martin, Physics Reports, Jul 1 1991, Vol 205

"Empirical Limits to Antigravity", Ericson & Richter, Europhysics Letters, Feb 15 1990, Vol 11 no 4

"Chern-Simons Quantizations of (2+1) Anti-de Sitter Gravity on a Torus", K. Ezawa, Classical and Quantum Gravity, Feb 1 1995 Vol 12 No 2

"Green's Function for Anti-de Sitter Space Gravity", Gary Kleppe, Physical Review d: Particles, Fields, Gravity; Dec 15 1994 Vol 50 No 12

"Lowest Eigenvalues of the Energy Operator for Totally Anti Symmetric Massless Fields of the N-Dimensional Anti-de Sitter Group", R.R. Metsaev, Classical and Quantum Gravity, Nov 1 1994, Vol 11 No 11

"The Positivity of Energy for Asymptotically Anti-de Sitter Spacetimes", E. Woolgar, Classical and Quantum Gravity, Jul 1 1994, Vol 11 No 7

"Vacuum Polarization Near Asymptotically Anti-de Sitter Black Holes in Odd Dimensions", Shiraishi & Maki, Classical and Quantum Gravity, Jul 1 1994, Vol 11 No 7

"Strong Anti Gravity: Life in the Shock Wave", Fabbrichesi & Roland, Nuclear Physics B, Dec 21 1992, Vol 388 No 2

"Global Solutions of Yang-Mills Equations on Anti-de Sitter Spacetime", Choquet-Bruhat, Classical and Quantum Gravity, Dec 1 1989, Vol 6 No 12

"The Scalar Wave Equation on Static de Sitter and Anti-de Sitter Spacetimes", D. Polarski, Classical and Quantum Gravity, Jun 1 1989

"Lehman Representation of the Spinor Two-Point Function in Anti-de Sitter Space", E. Gath, Classical and Quantum Gravity, May 1 1989, Vol 6 no 5

Dr. Bernhard Haisch has modeled inertial mass as deriving from an accelerated body's interaction with the zero point field (ZPF), consonant with a large body of refereed physics literature.

Haisch in Feb 1994 Phys. Rev. A
 Science vol 263 p 612
 Scientific American vol 270, p 30
 New Scientist 25 Feb 1995 p 30

"Gravity as a Zero-Point-Fluctuation Force,"
 H.E. Puthoff, Physical Review A: General
 Physics. Mar 1 1989, Vol39 No 5

The 4 February 1994 issue of Science magazine has an article about a new theory of inertia. A recent paper by Bernhard Haisch, Alfonso Rueda and Hal Puthoff in the 1 Feb 1994 issue of Physical Review A, based on earlier work by Andrei Sakharov, derives inertia from quantum electromagnetic vacuum fluctuations. The idea is that if inertia is due to some strange quantum EM effects, it might be understood and controlled, and even neutralized. Haisch is at the Lockheed Palo Alto laboratories, Rueda, at Cal. State. Long Beach, and Puthoff at the Institute for Advanced Studies in Austin Texas. Needless to say, this new theory is serious, but very controversial physics. A test is planned later this year at the SLAC linear accelerator by exposing a high energy electron beam to terawatt laser. Keep tuned!

-- John H. Chalmers Jr

A recent controversial theory of Austin Institute for Advanced Study physicist Hal Puthoff and his collaborators Haisch and Rueda appears to explain gravity as not an intrinsic property of matter but as an indirect consequence of Maxwellian electromagnetic radiation, namely that (as earlier suggested by the late Russian dissenter Sakharov) gravity is a "shadow effect" similar to the Casimir Effect of quantum electrodynamics. Bass points out that if the Haisch-Puthoff-Rueda theory is correct then Hodowanec's idea of tapping the earth's gravity field in some electromagnetic way not hitherto suggested is conceivable. - Joel McClain

Puthoff and his collaborators have gone so far as to use SED (Stochastic Electro-Dynamics) to _explain_ both gravitational & inertial mass and to show their equivalence, and to derive Newton's $F = Ma$, and to derive Mach's principle (without which Einstein admitted that no theory of gravity could claim to be complete), and to derive Dirac's "cosmological numerical coincidences" as inevitabilities, and to derive Newtonian gravity, and to derive the Newton-Cavendish parameter G!!! -- Robert Bass

It is an amazing coincidence that the total Newtonian gravitational potential energy of any object due to all masses in the universe is equal in magnitude to its total energy, at least to within a small factor, considering that this involves an expression involving multiple factors of the order of 10 to the 40th power. This was pointed out by Dirac in his Large Numbers Hypothesis, and used as part of a beautiful illustrative theory by Dennis Sciama [1], in which he constructs a theory of gravity closely analogous to the classical theory of electromagnetism, and shows that inertia can be directly attributed to the gravitational effect of accelerating relative to the gravitational potential sources of the whole universe (or indeed of accelerating the whole universe relative to the object, because in Sciama's theory, the two points of view are equivalent). This theory is obviously consistent with Mach's Principle (which is effectively that inertial motion is in some sense relative to the rest of universe). Sciama's theory is only a simplified approximation, but it is so neat that it seems likely that some similar principle must apply also within General Relativity. However, one of its most basic implications is that the gravitational "constant" G would depend on the distribution of matter in the universe, which seems to be in direct conflict with GR. I personally think GR is probably not quite right. -- Jonathan Scott

[1] D.W.Sciama, "On the Origin of Inertia", M.N.R.A.S. Vol. 113, p34, 1953.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9412012 From: "Haret Rosu" Date: 3 Dec 94 19:36:00 CST

Classical and quantum inertia: a heuristic introduction, Author(s): Haret C. Rosu

Report: IFUG-27/94,

Comments: 20 pages, LaTex 11pt, no figures. A non-technical discussion of the problem of inertia is provided both in classical physics and in the quantum world. After briefly reviewing the classical formulations (WEP, EEP, and SEP), I pass to a presentation of the equivalence statements for quantum vacuum states. One can also find a number of related comments and suggestions.

Krech, Michael.

The Casimir effect in critical systems / Michael Krech. Singapore ; River Edge, NJ : World Scientific, c1994. x, 253 p. : ill. ; 23 cm.

LC CALL NUMBER: QC173.4.C74 K74 1994

SUBJECTS: Critical phenomena. Casimir effect. ISBN: 9810218451

Cavity quantum electrodynamics/edited by Paul R. Berman. Boston : Academic Press, c1994. xvi, 464 p. : ill. ; 24 cm. LC CALL NUMBER: QC446.2 .C38 1994

SUBJECTS: Quantum optics. Quantum electrodynamics. Casimir effect. ISBN: 0120922452 (alk. paper)

Long-range Casimir forces : theory and recent experiments on atomic systems Edited by Frank S. Levin and David A. Micha. New York : Plenum Press, c1993. LC CALL NUMBER: QC680 .L63 1993

SUBJECTS: Casimir effect.

ISBN: 0306443856

Physics in the making : essays on developments in 20th century physics: in honour of H.B.G. Casimir on the occasion of his 80th birthday/ edited by A. Sarlemijn and M.J. Sparnaay. Amsterdam : North-Holland; New York, N.Y., U.S.A. : Sole distributors for the U.S.A. and Canada, Elsevier Science Pub. Co., 1989. xiv, 361 p. : ill. ; 23 cm. LC CALL NUMBER: QC7 .P48 1989

SUBJECTS: Casimir, H. B. G. (Hendrik Brugt Gerhard), 1909- Casimir, H. B. G. (Hendrik Brugt Gerhard), 1909- Sarlemijn, Andries, 1936-

Sparnaay, M. J. (Marcus Johannes)

ISBN: 0444881212

Edwards-Casimir Quantum Vacuum Drive -- A hypothetical drive exploiting the peculiarities of quantum mechanics by restricting allowed wavelengths of virtual photons on one side of the drive (the bow of the ship); the pressure generated from the unrestricted virtual photons toward the aft generates a net force and propels the drive. ----

CONDENSED MATTER, ABSTRACT COND-MAT/9505108 From:

moraes@guinness.ias.edu (Fernando Moraes) Date: Tue, 23 May 95 17:12:35 EDT

Enhancement of the magnetic moment of the electron due to a topological defect

Author(s): Fernando Moraes (Institute for Advanced Study, Princeton) In the framework of the theory of defects/three-dimensional gravitation, it is obtained a positive correction to the magnetic moment of the electron bound to a disclination in a dielectric solid. With the disclination modelled as a parallel plate casimir effect.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9212077 From:

milton@phyast.nhn.uoknor.edu (Kim Milton) Date: Fri, 11 Dec 92 16:13:13 CST

MAXWELL-CHERN-SIMONS CASIMIR EFFECT, KIMBALL A. MILTON, DEPARTMENT OF PHYSICS AND ASTRONOMY, UNIVERSITY OF OKLAHOMA In odd-dimensional spaces, gauge invariance permits a Chern-Simons mass term for the gauge fields in addition to the usual Maxwell-Yang-Mills kinetic energy term. We study the Casimir effect in such a (2+1)-dimensional Abelian theory. For the case of parallel conducting lines the result is the same as for a scalar field. For the case of

circular boundary conditions the results are completely different, with even the sign of the effect being opposite for Maxwell-Chern-Simons fields and scalar fields. We further examine the effect of finite temperature. The Casimir stress is found to be attractive at both low and high temperature. Possibilities of observing this effect in the laboratory are discussed.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9303038 PHYS. REV. D 48, 776 (1993)
FROM: LFORD@PEARL.TUFTS.EDU

Date: Wed, 31 Mar 1993 17:47 EDT
MOTION OF INERTIAL OBSERVERS THROUGH NEGATIVE ENERGY, BY L.H. FORD AND THOMAS A. ROMAN,

Recent research has indicated that negative energy fluxes due to quantum coherence effects obey uncertainty principle-type inequalities of the form $\|\Delta E\|, \{\Delta \tau\} \approx 1$. Here $\|\Delta E\|$ is the magnitude of the negative energy which is transmitted on a timescale $\Delta \tau$. Our main focus in this paper is on negative energy fluxes which are produced by the motion of observers through static negative energy regions. We find that although a quantum inequality appears to be satisfied for radially moving geodesic observers in two and four-dimensional black hole spacetimes, an observer orbiting close to a black hole will see a constant negative energy flux. In addition, we show that inertial observers moving slowly through the Casimir vacuum can achieve arbitrarily large violations of the inequality. It seems likely that, in general, these types of negative energy fluxes are not constrained by inequalities on the magnitude and duration of the flux. We construct a model of a non-gravitational stress-energy detector, which is rapidly switched on and off, and discuss the strengths and weaknesses of such a detector.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9304008 PHYS. REV. D 47, 4510 (1993).
FROM: LFORD@PEARL.TUFTS.EDU
Date: Tue, 6 Apr 1993 12:56 EDT

SEMICLASSICAL GRAVITY THEORY AND QUANTUM FLUCTUATIONS, BY CHUNG-I KUO AND L. H. FORD.

We discuss the limits of validity of the semiclassical theory of gravity in which a classical metric is coupled to the expectation value of the stress tensor. It is argued that this theory is a good approximation only when the fluctuations in the stress tensor are small. We calculate a dimensionless measure of these fluctuations for a scalar field on a flat background in particular cases, including squeezed states and the Casimir vacuum state. It is found that the fluctuations are small for states which are close to a coherent state, which describes classical behavior, but tend to be large otherwise. We find in all cases studied that the energy density fluctuations are large whenever the local energy density is negative. This is taken to mean that the gravitational field of a system with negative energy density, such as the Casimir vacuum, is not described by a fixed classical metric but is undergoing large metric fluctuations. We propose an operational scheme by which one can describe a fluctuating gravitational field in terms of the statistical behavior of test particles. For this purpose we obtain an equation of the form of the Langevin equation used to describe Brownian motion.

HIGH ENERGY PHYSICS - PHENOMENOLOGY, ABSTRACT HEP-PH/9307258
From: langfeld@ptsun1.tphys.physik.uni-tuebingen.de (Kurt Langfeld)
Date: Tue, 13 Jul 93 08:04:30 +0200

CASIMIR EFFECT OF STRONGLY INTERACTING SCALAR FIELDS, BY K. LANGFELD, F. SCHMUSER, AND H. REINHARDT

Non-trivial ϕ^4 -theory is studied in a renormalisation group invariant approach inside a box consisting of rectangular plates and where the scalar modes satisfy periodic boundary conditions at the plates. It is found that the Casimir energy exponentially approaches the infinite volume limit, the decay rate given by the scalar condensate. It therefore essentially differs from the power law of a free theory. This might provide experimental

access to properties of the non-trivial vacuum. At small interplate distances the system can no longer tolerate a scalar condensate, and a first order phase transition to the perturbative phase occurs. The dependence of the vacuum energy density and the scalar condensate on the box dimensions are presented.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC 9310007 PHYSICA SCRIPTA 48, 649 (1993) FROM: harald@nordita.dk (Harald H. Soleng)

Date: Mon, 4 Oct 93

INVERSE SQUARE LAW OF GRAVITATION IN (2+1)-DIMENSIONAL SPACE-TIME AS A CONSEQUENCE OF CASIMIR ENERGY, H. H. SOLENG, The gravitational effect of vacuum polarization in space exterior to a particle in (2+1)-dimensional Einstein theory is investigated. In the weak field limit this gravitational field corresponds to an inverse square law of gravitational attraction, even though the gravitational mass of the quantum vacuum is negative. The paradox is resolved by considering a particle of finite extension and taking into account the vacuum polarization in its interior.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9312069 From: segui@cc.unizar.es

Date: Thu, 9 DEC 93 13:50 GMT

A MODIFIED SCHWINGER'S FORMULA FOR THE CASIMIR EFFECT, M.V. COUGO-PINTO, C. FARINA AND ANTONIO J. SEGUI-SANTONJA

After briefly reviewing how the (proper-time) Schwinger's formula works for computing the Casimir energy in the case of "scalar electrodynamics" where the boundary conditions are dictated by two perfectly conducting parallel plates with separation "a" in the Z-axis, we propose a slightly modification in the previous approach based on an analytical continuation method. As we will see, for the case at hand our formula does not need the use of Poisson summation to get a (renormalized) finite result.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9401123 From: segui@cc.unizar.es

Date: Tue, 25 JAN 94 21:47 GMT

SCHWINGER'S METHOD FOR THE MASSIVE CASIMIR EFFECT, BY M.V. COUGO-PINTO, C. FARINA AND A.J. SEGUI-SANTONJA

We apply to the massive scalar field a method recently proposed by Schwinger to calculate the Casimir effect. The method is applied with two different regularization schemes: the Schwinger original one by means of Poisson formula and another one by means of analytical continuation.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9405060 From: Shtykov Nikolay

Date: Tue, 10 May 94 17:40:50 JST

THE FINITE VACUUM ENERGY FOR SPINOR, SCALAR AND VECTOR FIELDS, N.SHTYKOV We compute the one-loop potential (the Casimir energy) for scalar, spinor and vectors fields on the spaces \mathbb{R}^{m+1} , \mathbb{S}^N , \mathbb{CP}^2 . As a physical model we consider spinor electrodynamics on four-dimensional product manifolds. We examine the cancelation of a divergent part of the Casimir energy on even-dimensional spaces by means of including the parameter M in original action. For some models we compare our results with those found in the literature.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9408172 From: LFORD@PEARL.TUFTS.EDU

DECOHERENCE AND VACUUM FLUCTUATIONS, L.H. FORD, TUFTS UNIVERSITY The interference pattern of coherent electrons is effected by coupling to the quantized electromagnetic field. The amplitudes of the interference maxima are changed by a factor which depends upon a double line integral of the photon two-point function around the closed path of the electrons. The interference pattern is sensitive to shifts in the vacuum fluctuations in regions from which the electrons are excluded. Thus

this effect combines aspects of both the Casimir and the Aharonov-Bohm effects. The coupling to the quantized electromagnetic field tends to decrease the amplitude of the interference oscillations, and hence is a form of decoherence. The contributions due to photon emission and to vacuum fluctuations may be separately identified. It is to be expected that photon emission leads to decoherence, as it can reveal which path an electron takes. It is less obvious that vacuum fluctuations also can cause decoherence. What is directly observable is a shift in the fluctuations due, for example, to the presence of a conducting plate. In the case of electrons moving parallel to conducting boundaries, the dominant decohering influence is that of the vacuum fluctuations. The shift in the interference amplitudes can be of the order of a few percent, so experimental verification of this effect may be possible. The possibility of using this effect to probe the interior of matter, e.g., to determine the electrical conductivity of a rod by means of electrons encircling it is discussed. (Presented at the Conference on Fundamental Problems in Quantum Theory, University of Maryland, Baltimore County, June 18-22, 1994.)

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9410043 PHYS. REV. D 51, 4277(1995). FROM: FORD@TUHEP.PHY.TUFTS.EDU Date: Fri, 28 Oct 1994 20:33 EST

AVERAGED ENERGY CONDITIONS AND QUANTUM INEQUALITIES, L.H. FORD AND THOMAS A. ROMAN

Connections are uncovered between the averaged weak (AWEC) and averaged null (ANEC) energy conditions, and quantum inequality restrictions on negative energy for free massless scalar fields. In a two-dimensional compactified Minkowski universe, we derive a covariant quantum inequality-type bound on the difference of the expectation values of the energy density in an arbitrary quantum state and in the Casimir vacuum state. From this bound, it is shown that the difference of expectation values also obeys AWEC and ANEC-type integral conditions. In contrast, it is well-known that the stress tensor

in the Casimir vacuum state alone satisfies neither quantum inequalities nor averaged energy conditions. Such difference inequalities represent limits on the degree of energy condition violation that is allowed over and above any violation due to negative energy densities in a background vacuum state. In our simple two-dimensional model, they provide physically interesting examples of new constraints on negative energy which hold even when the usual AWEC, ANEC, and quantum inequality restrictions fail. In the limit when the size of the space is allowed to go to infinity, we derive quantum inequalities for timelike and null geodesics which, in appropriate limits, reduce to AWEC and ANEC in ordinary two-dimensional Minkowski spacetime. We also derive a quantum inequality bound on the energy density seen by an inertial observer in four-dimensional Minkowski spacetime. The bound implies that any inertial observer in flat spacetime cannot see an arbitrarily large negative energy density which lasts for an arbitrarily long period of time.


GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9411053 From: "Haret Rosu" Date: 20 Nov 94 21:15:00 CST

On the assignment of frequency spectra to quantum vacuum effects, Author: Haret C. Rosu, Report: IFUG-25/94, I discuss in an introductory manner, i.e., in the form of comments on available references, the problem of assigning frequency spectra to such fundamental effects like Casimir, Hawking, Unruh, and squeezing effects. This may help to clarify their differences as well as their similarities.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9411056 From: ulvi@tapir.Caltech.EDU (Ulvi Yurtsever) Date: Mon, 21 Nov 94 15:56:11 -0800

The averaged null energy condition and difference inequalities in quantum field theory, by: Ulvi Yurtsever

Recently, Larry Ford and Tom Roman have discovered that in a flat cylindrical space,



although the stress-energy tensor itself fails to satisfy the averaged null energy condition (ANEC) along the (non-achronal) null geodesics, when the "Casimir-vacuum" contribution is subtracted from the stress-energy the resulting tensor does satisfy the ANEC inequality. Ford and Roman name this class of constraints on the quantum stress-energy tensor "difference inequalities." Here I give a proof of the difference inequality for a minimally coupled massless scalar field in an arbitrary two-dimensional spacetime, using the same techniques as those we relied on to prove ANEC in an earlier paper with Robert Wald. I begin with an overview of averaged energy conditions in quantum field theory.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9502024 From:

MANKO@napoli.infn.it

Date: Mon, 27 Feb 1995 16:32:21 +0200
(CET) Deformation of Partical Distribution Functions due to Q-nonlinearity and Nonstationary Casimir Effect, Author: V. I. Man'ko The geometrical phase is shown to be integral of motion. Deformation of particle distribution function corresponding to nonstationary Casimir effect is expressed in terms of multivariable Hermite polynomials. Correction to Planck distribution due to q-nonlinearity is discussed.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9503001 From:

onofrio%38619.hepnet@Csa4.LBL.Gov

Date: Wed, 1 Mar 95 08:23:43 PST
Detecting Casimir Forces through a Tunneling Electromechanical Transducer Authors: Roberto Onofrio , Giovanni Carugno We propose the use of a tunneling electromechanical transducer to dinamically detect Casimir forces between two conducting surfaces. The maximum distance for which Casimir forces should be detectable with our method is around \$1 \mu m\$, while the lower limit is given by the ability to approach the surfaces. This technique should permit to study gravitational forces on the same range of distances, as well as the vacuum friction provided that very low dissipation mechanical resonators are used.

CONDENSED MATTER THEORY,
ABSTRACT COND-MAT/9505023 From:
moraes@guinness.ias.edu (Fernando Moraes)
Date: Fri, 5 May 95 09:35:57 EDT

Casimir effect around disclinations

Author: Fernando Moraes (Institute for Advanced Study, Princeton) This communication concerns the structure of the electromagnetic quantum vacuum in a disclinated insulator. It is shown that a nonzero vacuum energy density appears when the rotational symmetry of a continuous insulating elastic medium is broken by a disclination. An explicit expression is given for this Casimir energy density in terms of the parameter describing the disclination.

CONDENSED MATTER THEORY,
ABSTRACT COND-MAT/9505108 From:
moraes@guinness.ias.edu (Fernando Moraes)
Date: Tue, 23 May 95 17:12:35 EDT

Enhancement of the magnetic moment of the electron due to a topological defect Author: Fernando Moraes (Institute for Advanced Study, Princeton) In the framework of the theory of defects/three-dimensional gravitation, it is obtained a positive correction to the magnetic moment of the electron bound to a disclination in a dielectric solid.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9506005 From: JAEKEL Marc
Date: Wed, 7 Jun 1995 16:30:40 +0200

Mechanical Effects of Radiation Pressure Quantum Fluctuations Authors: Marc-Thierry Jaekel (Laboratoire de Physique Théorique de l'Ecole Normale Supérieure), Serge Reynaud (Laboratoire Kastler-Brossel) As revealed by space-time probing, mechanics and field theory come out as complementary descriptions for motions in space-time. In particular, quantum fields exert a radiation pressure on scatterers which results in mechanical effects that persist in vacuum. They include mean forces due to quantum field fluctuations, like Casimir forces, but also fluctuations of these forces and additional forces linked to motion. As in classical electron theory, a moving scatterer is submitted to a radiation reaction force which

modifies its motional response to an applied force. We briefly survey the mechanical effects of quantum field fluctuations and discuss the consequences for stability of motion in vacuum and for position fluctuations.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9506006 From: JAEKEL Marc
 Date: Wed, 7 Jun 1995 16:58:17 +0200

Quantum Fluctuations and Inertia

Authors: Marc-Thierry Jaekel (Laboratoire de Physique Théorique de l'Ecole Normale Supérieure), Serge Reynaud (Laboratoire Kastler-Brossel) Vacuum field fluctuations exert a radiation pressure which induces mechanical effects on scatterers. The question naturally arises whether the energy of vacuum fluctuations gives rise to inertia and gravitation in agreement with the general principles of mechanics. As a new approach to this question, we discuss the mechanical effects of quantum field fluctuations on two mirrors building a Fabry-Perot cavity. We first put into evidence that the energy related to Casimir forces is an energy stored on field fluctuations as a result of scattering time delays. We then discuss the forces felt by the mirrors when they move within vacuum field fluctuations, and show that energy stored on vacuum fluctuations contributes to inertia in conformity with the law of inertia of energy. As a further consequence, inertial masses exhibit quantum fluctuations with characteristic spectra in vacuum.

QUANTUM PHYSICS, ABSTRACT
QUANT-PH/9506023 From:

claudia@cromwell.physics.uiuc.edu (Claudia C Eberlein) Date: Thu, 15 Jun 95 11:13:57 - 0500

Sonoluminescence as quantum vacuum radiation Author: Claudia Eberlein (Dept of Physics, UIUC, Urbana, IL) Sonoluminescence is explained in terms of quantum radiation by moving interfaces between media of different polarizability. It can be considered as a dynamic Casimir effect, in the sense that it is a consequence of the imbalance of the zero-point fluctuations of the electromagnetic field during the non-inertial motion of a boundary. The transition amplitude from the vacuum into a


two-photon state is calculated in a Hamiltonian formalism and turns out to be governed by the transition matrix-element of the radiation pressure. Expressions for the spectral density and the total radiated energy are given.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT
HEP-TH/9508086 From: eli@ecm.ub.es (Emili Elizalde)

Date: Fri, 18 Aug 1995 10:14:50 +0200

A precise definition of the Casimir energy, Authors: K. Kirsten , E. Elizalde The somehow arbitrary definition of the Casimir energy corresponding to a quantum system in a d -dimensional ultrastatic spacetime ---profusely used in the last years--- which has been criticized sometimes for adopting without a sound argument the minimal subtraction scheme, is shown to be completely equivalent to the definition stemming naturally from the concept of functional determinant through the zeta-function prescription. This is done by considering the theory at finite temperature and by defining then the Casimir energy as its energy in the limit $T \rightarrow 0$. The ambiguity in the coefficient $C_{d/2}$ is understood to be a result of the necessary renormalization of the free energy of the system. As an example, the Casimir energy corresponding to a general $(1+2)$ -dimensional toroidal spacetime with flat spatial geometry, parametrized by the corresponding Teichmüller parameters, and its precise dependence on these parameters is obtained under the form of an analytic function.

Ernest G. Cullwick. In his book "Electromagnetism and Relativity", published in 1957, was one of the first to provide an analysis of the probable coupling between EM and inertial fields. Cullwick realized that Maxwell's equations and most existing theories of electrodynamics assume that the mass of an electron is zero. At Maxwell's time this was a reasonable assumption. But it is well known today that electrons have mass, and therefore an inertial momentum is always associated with an electric current. Cullwick suggested in his analysis that coupling terms between EM and inertia may be very small, but would likely



appear sometime in the future as we go to higher current densities. And he was one of the first scientists to predict some of the odd effects which can now seen with superconductors. Cullwick was also one of the first to identify and attempt an analysis of the relativistic paradoxes and unusual effects which occur in a rotating EM field. His work still stands today as one of the only existing efforts to consider the problem of a rotating EM field.

AUTHOR: Cullwick, E. G. (Ernest Geoffrey), 1903-

TITLE: Electromagnetism and relativity : with particular reference to moving media and electromagnetic induction / by E. G. Cullwick.

EDITION 2d ed.

PUBL.: New York : J. Wiley,

DATE: 1959 (2nd Edition)

SUBJECT: Electromagnetic theory, Relativity (Physics)

AUTHOR: Cullwick, E. G. (Ernest Geoffrey), 1903-

TITLE: The fundamentals of electromagnetism by E.G. Cullwick.

EDITION 3rd ed.

PUBL.: London, Cambridge U.P.,

DATE: 1966 (3rd Edition)

SUBJECT: Electromagnetism

AUTHOR: Cullwick, E. G. (Ernest Geoffrey), 1903-

TITLE: The fundamentals of electromagnetism; a restatement for engineering students and others of physical and theoretical principles in accordance with modern scientific thought, by E. Geoffrey Cullwick ... With an appendix and numerous examples on the recently adopted M.K.S. system of practical units ...

PUBL.: New York, The Macmillan company; Cambridge, Eng., The University press,

DATE: 1939


SUBJECT: Electromagnetism

If you work out the metric for EM waves circulating in a cavity you get some strange results. There is a preliminary discussion of this effect in the article by Houshang Ardavan, 'Gravitational Waves from Electromagnetic Waves' in the book "Classical General Relativity," edited by W.B. Bonner, I.N. Islam and M.A.H. MacCollum (Cambridge Univ. Press, 1984).

It is something I have seen done. At the point in an annular cavity where the phase velocity goes from less than c to greater than c, a term shows up in the derived metric of the system that looks like a source term. On the other hand you have assumed that the metric is source free in the EM region of the cavity. So you get a solution which contradicts the hypothesis that went into building the solution. You get something which is possibly unphysical. Now Einstein's equation and the associated geometry is pretty tricky and it is easy to get unphysical solutions. The final arbitors of whether a solution is satisfactory or not is physical reasonability and self consistancy (these are almost the same thing). The cavity problem seems very physically reasonable initially, but ends with a self-consistancy problem which appears to be unphysical. Also, Cauchy's theorem does not apply to this case since it becomes a mixed type problem (elliptic and hyperbolic PDEs), so the Hawking singularity theorems don't a priori apply. It is something very interesting, but to publish it with out being scoffed at would take a lot of work and possibly inventing some new math. -- Jim McClune, University of Missouri

ROTATING FIELDS IN GENERAL RELATIVITY, by Islam, J.N. Begins with a short introduction to the relevant aspects of general relativity. This is followed by a detailed derivation of the Wehl-Lewis-Papapetrou form of the stationary axially symmetric metric. The Kerr and Tomimatsu-Sato forms of the rotating interior and exterior solutions of the Einstein equations are then considered. Subject: physics
1985 6 X 9 122 pp. 4 diagrams

Hardback 0-521-26082-5 \$47.95 (J7.99)



>If an EM field is somehow rotated extremely fast, shouldn't all matter be repelled from its center? -kgo.

How fast do you want it rotated? It's fairly simple to construct a system to produce rotating EM waves at whatever rotational velocity you wish by feeding a pair of broadside dipole arrays with quadrature phased waves. It is quite simple to construct a system that would have a rotational velocity of C within the uniform field area. It might also be fairly easy to do this with a Hemholtz coil arrangement as well, but the broadside array will be much easier to do at easily engineerable frequencies. Some really interesting paradoxes come about when the rotational frequency is high enough so that the rotational velocity exceeds C within the uniform field area of the arrays or within the hemholtz coils. -- Robert Shannon

Ehrenfest Paradox (Ehrenfest, 1909) --

The special relativistic "paradox" involving a rapidly rotating disc. Since any radial segment of the disc is perpendicular to the direction of motion, there should be no length contraction of the radius; however, since the circumference of the disc is parallel to the direction of motion, it should contract.

Question -- by Kung Lo (October 1995):

Take a rigid disk of radius R and spin it up to angular velocity . As seen by an observer S that is at rest in the center of the disk, the radius is still R, but the circumference is contracted by the Lorentz effect. How is this possible?

More physically, if a fixed ring is just outside the spinning disk and placed with equally spaced markers on the rim of the disk and on the fixed ring, I know by symmetry that, when one marker on the disk is aligned with a marker on the ring, all pairs of markers must be aligned. This contradicts the fact that, for observer S, the distance between successive markers on the disk is reduced by the Lorentz factor.

Answer -- provided by David Djajaputra (November 1995): It seems that the rotating disk paradox (it turned out to be Ehrenfest's paradox) has been extensively analyzed by many people (including Einstein himself, who developed general relativity to answer this problem, as one author speculates...). This I found from a nice paper :

O. Gron, "Relativistic description of a rotating disk" Am. J. Phys. V43, 869 (1975), and all the references therein.

The key sentence in Gron's paper is at the end of Section IV: "By definition a Born rigid motion of a body leaves lengths unchanged, when measured in the body's proper frame . (...) A Born rigid motion is not a material property of the body, but the result of a specific program of forces designed to set the body in motion without introducing stresses. (...) A transition of the disk from rest to rotational motion, while it satisfies Born's definition of rigidity, is a kinematic impossibility"

With this kinematics the radius is R and the circumference is as measured by observer S (lab frame), but an observer riding on the disk will measure a distance R to the center and a distance around the circumference (he can do this measurement by slowly walking around the spinning disk with a meter tape). This is consistent with the usual Lorentz contraction . The point is that this is NOT a Born rigid motion. There is much more in Gron's paper. -- Vittorio Celli

Several key phrases keep popping up regarding rotating fields, powerful magnetic pulsed fields, and 90 degree cross-field phase shifts. For example, Preston Nicholes describes a device known as a Delta T antenna in the Montauk series of books. The Delta T antenna is described as a pyramidal structure, but lets just take two square loops, placed at 90 degrees to each other, and feed these two loops with an RF signal, also with a 90 degree phase shift, we will produce a rotating magnetic field



within the loops (these loops share a common center point, and each loop is in a plane 90 degrees from the other) The speed of rotation of this magnetic field is a direct function of the frequency of the applied RF signal. At the center of the antenna, the rotational velocity is zero, but as you move out from the center, and rotational velocity increases. At some distance from center would reach the speed of light, dependant of the frequency used. One could imagine that the rotational velocity of this rotating magnetic field could reach the speed of light within the antenna structure itself if a way could be found to make the antenna much larger than a normally resonant antenna would be for that same frequency. At several hundred megahertz, a two meter per side square loop would have a rotational velocity well in excess of the speed of light within the antenna structure itself.

What effect would there be at the boundary where the rotational velocity reached, and then exceeded the speed of light. How could the magnetic field even propagate to the center of the antenna structure if it would have to move faster than light to reach that space? If hemholtz coils were used instead of loops, the magnetic field strength would be uniform inside the structure, how could the field strength be uniform if there is not sufficient time for the field to propagate through the space inside the structure itself?

Could such an effect actually generate a wormhole like phenomena, at energy levels far below that of neutron stars and such? As the causal mechanism, the magnetic field, is in rotation, would this describe a traversable worm hole as has been postulated in relationship to rotating black holes? -- Robert Shannon

Aono, Osamu, 1937-

Rotation of a magnetic field / Osamu Aono and Ryo Sugihara. Nagoya, Japan : Institute of Plasma Physics, Nagoya University, 1986. 6 p. ; 30 cm. LC CALL NUMBER: QC717.6 .N35 no. 792 (ALTERNATE CLASS QC754.2.M3)

SUBJECTS: Magnetic fields.


Electrodynamics. Research report (Nagoya Daigaku. Purazumu Kenkyujo) ; IPPJ-792. ----

Let me clear this up a bit, the two coils are acting as antennae already, producing the rotating field by vector summation of the radiated quadrature phased EM waves. The loops would be operating as the driven elements of a cubical antenna, not as coils as such. If you prefer, substitute the two loop antenna with a pair of crossed dipoles at 90 degrees, this will also produce the rotating field, but the center will be occupied by the dipoles rather than be open as with loop antenna or by using sets of broadside arrays. Note that this is not the same as the rotational speed reaching c inside the "uniform field" area, as described earlier. It's simple a tool to understand the generation of the rotating field and the relationship between applied frequency and the resultant rotational speed. Rather than loop elements, in practice you might use a phased array of dipole elements that produces a constant phase plane wave, not unlike a pair of hemholtz coils produced a uniform field within the coil sets. Four of these "broadside arrays" would form the four sides of a cube, inside of which you could induce the fast rotating fields from the radiated EM waves. In all cases, the driven elements are launching EM waves at c. Only the vector sum of the two (of four) quadrature fields is in rotation, which leads us back to the question of what happens as the rotational velocity of the sum of these EM fields reaches c within the field generator, and there is not sufficient time for the fields to propagate across the $V_r = c$ boundary?

This is the point where two different physicists have tried to lead me down the garden path of "red shifted magnetic fields". I'm not sure I'm ready to buy that concept just yet.

-- Robert Shannon

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9601034 From: Tevian Dray Date: Mon, 22 Jan 1996 10:57:03 PST
The Rotating Quantum Vacuum



Author(s): Paul C. W. Davies , Tevian Dray , Corinne A. Manogue Report-no: ADP 95-43/M36 (University of Adelaide) We derive conditions for rotating particle detectors to respond in a variety of bounded spacetimes and compare the results with the folklore that particle detectors do not respond in the vacuum state appropriate to their motion. Applications involving possible violations of the second law of thermodynamics are briefly addressed.

I'm also saying that a pair of crossed coils will start behaving differently when the driving frequency is so high that the field lines near them try to exceed the speed of light. At low frequencies the coils create a rotating magnetic field. At high frequencies they send out radio waves having a rotating field vector (circularly polarized waves, in other words.) WITHIN the volume of the coils the fields still rotate, at least until the frequency is raised so high that the coils are many wavelengths across. At these frequencies the fields in the center of the crossed coils would be of complex shape, maybe some kind of contracting spiral. (Which is interesting, because at very high frequencies there would be a "hot spot" at the exact center of the crossed coils.) -- Robert Shannon

On similar topic: anyone ever heard of the "CFA antenna" flap in the UK? CFA is for "crossed-field antenna." There were a bunch of articles and letters to the editor in EWW, "Electronics and Wireless World," the British engineering mag. The CFA-believers thought they had discovered a way to make 1-foot antennas which were efficient at 100-meter wavelengths. The key to the CFA was to create the e- and b-fields separately: feed both a coil-loop and a pair of capacitor-spheres with separate high-current and high-voltage signals respectively, orient them 90deg to produce a broadside wave, shift the phases with L/C networks to form the proper EM wave (90? zero?), and then obtain a powerful EM emission from a tiny antenna. There was a great quantity of argument and name-calling over this, all done in slow-motion over many months of letters in the letters-to-the-editor

column. Then it just died away. Either the pro-CFA side couldn't prove that it worked, or nobody believed the proof they did find.


-- William Beaty

And some comments about rotating EM fields by Dr Dennis Cravens in a report titled "Electric Propulsion Study", done for the Astronautics Laboratory at Edwards AFB. August 1990. Dennis Cravens was formerly with SAIC Corp, and is now working with CETI in development of cold fusion. Anyway, here's some things he says in the electric propulsion report about the "peculiarities" of a rotating magnet:

ROTATION OF MAGNETS - There is a continuing debate in physics as to the reality of the magnetic field. The prime question is whether the axial magnetic field of a bar magnet rotates with the magnet or is stationary. The Faraday homopolar generator dates back to the 1830s. DePalma, Tewari, and others have attempted to utilize the Faraday generator to produce more power than needed to run it. Most objective reviews of the work have, however, failed to see such effects.

It is doubtful that these claims will be independently validated and even more doubtful that they will lead directly to a propulsive system. However, the work on homopolar generators as high current devices is reasonable and may be useful for ground uses. The angular momentum complications seem to rule the system out for any practical space applications.

SEARL EFFECT - The Searl Effect is a separate issue from homopolar generator above. Searl has claimed to produce disk levitation by rapidly rotating magnets. There have been claims of anti-gravity, high electric fields, perpetual motion, inertial loss, and gas ionization. All these claims come from Searl or those supportive of his work and no outside witnesses are available. Searl has not supplied any technical data or specifics of the operation in any easily referenced source. It is not recommended that his work be experimentally



followed by the USAF. It is worth noting however, that a rotating magnet does have some definite theoretical peculiarities.

Through the years there have been many interesting developments concerning the Faraday Homopolar generator. DePalma has claimed to get more energy out than is supplied to the the generator. None of the claims seem to withstand careful examination and no machine has ever been made self driving. The underlying reason that such claims continue to surface is that rotating magnetic fields are extremely difficult to handle within existing theories. This is because for a rotating frame there is a distance (removed from the axis) which is travelling at velocities greater than c. Although the distance is not withing any real physical object, it's existence within the mathematical development greatly complicates any calculations.

DePalma B.E., "Electro-Mechanical Device for the Amplification of Electrical Power", The New Age Science Magazine, No 7, 1980

Tewari P., "Generation of Electrical Power from Absolute Vacuum by High Speed Rotation of Conducting Magnetic Cylinder", Tech. Rep. Dept. of Atomic Energy, Bombay India, 1985

Searl, J.R.R., British provisional patent specification #57578, 1970 -----

These articles are indicative of studies of EM waves and rotating bodies. It appears that when EM waves pass through rotating dielectrics some unusual effects are predicted. This may lead to some interesting future technology.

-- Dr Dennis Cravens

"Some Remarks on Scattering by a Rotating Dielectric Cylinder", D. Schreiber, Journal of EM Waves and Applications, Vol 2 No2 1988

"Rotating Bodies and Electrodynamics in a Rotating Reference Frame", I.B. Zeldovich and L.V. Rozhavskii, Radiofizka Vol 29 No 9, 1986 -----

-- Here's an interesting news brief from Infinite Energy magazine, July/Aug 1995, Dr Eugene Mallove - editor. (603)-228-4516

A bombshell paper has just been published in the American Journal of Physics, Vol 63 No 8, August 1995, pages 694-705, "Maxwell's Equations in a Rotating Medium: Is There a Problem?" by Gerald N. Pellegrini and Arthur R. Swift (the latter of the Dept of Physics and Astronomy, University of Massachusetts, Amherst MA)"

The paper is a direct challenge to Special Relativity. It proves one of two things about a classic 1913 experiment of Wilson and Wilson that was used to verify the prediction of relativity that "a moving magnetic dipole develops an electric dipole moment.' The conclusion of the paper is that Special Relativity does NOT agree with this experiment -- and no one has ever challenged the quality of the experiment.

Peregrinni told Infinite Energy that he thinks that all of relativity as well as Maxwell's equations as descriptors of EM radiation are now called into question. -----

-

The origin of the Montauk Project dates back to 1943 when radar invisibility was being researched aboard the USS Eldridge. As the Eldridge was stationed at the Philadelphia Navy Yard, the events concerning the ship have commonly been referred to as the "Philadelphia Experiment." The objective of this experiment was to make the ship undetectable to radar and while that was achieved, there was a totally unexpected and drastic side effect. The ship became invisible to the naked eye and was removed from time and space as we know it. It went into 10-dimensional hyper-space. For further info into this, read the book called "Hyperspace" by Dr.

Michio Kaku. A movie called "The Philadelphia Experiment" was made, but delayed for two years as the Pentagon was able to halt its release. After the war, research continued under the tutelage of Dr. John Von Neumann who had directed the technical aspects of the Philly Experiment. A massive human factor study was also begun at Brookhaven National Labs on Long Island, New York -- known as the Phoenix Project. The Montauk Project culminated on August 12, 1983. A full blown time portal was fully functioning, but things were out of control and the project was crashed. An unauthorized video has been widely distributed regarding this story and several lectures have been given on the Montauk Project. One science reporter for the New York times started the project but tacked off when he discovered to his own surprise that the Montauk Project was indeed real.

Three books have been released by Preston Nichols, who was involved in the Project, and Peter Moon. They are

1) The Montauk Project: Experiments in Time - 1992
 2) Montauk Revisited: Adventures in Synchronicity - 1994
 3) Pyramids of Montauk Explorations in Consciousness-1995
 This coming year, 1996, the next book will be out and the title will be "Montauk Reconciled"

-- Richard Frager

HIGH ENERGY PHYSICS -
 PHENOMENOLOGY, ABSTRACT HEP-
 PH/9412234 From: Michael Martin Nieto 505-
 667-6127 Date: Mon, 5 Dec 94 09:52:27 -
 0700

THEORETICAL MOTIVATION FOR
 GRAVITATION EXPERIMENTS ON
 ULTRA-LOW ENERGY ANTIPROTONS
 AND ANTIHYDROGEN

Authors: Michael Martin Nieto , T. Goldman ,
 John D. Anderson , Eunice L. Lau, J. Perez-
 Mercader

Comments: Written version of invited contribution to LEAP'94: Third Biennial Conference on Low-Energy Antiproton Physics. We know that the generally accepted theories of gravity and quantum mechanics are fundamentally incompatible. Thus, when we

try to combine these theories, we must beware of physical pitfalls. Modern theories of quantum gravity are trying to overcome these problems. Any ideas must confront the present agreement with general relativity, but yet be free to wonder about not understood phenomena, such as the dark matter problem and the anomalous spacecraft data which we announce here. This all has led some "intrepid" theorists to consider a new gravitational regime, that of antimatter. Even more "daring" experimentalists are attempting, or considering attempting, the measurement of the gravitational force on antimatter, including low-energy antiprotons and, perhaps most enticing, antihydrogen.

HIGH ENERGY PHYSICS - EXPERIMENT,
 ABSTRACT HEP-EX/9412018 From:
 PHILLIPS@hep.phy.duke.edu

Date: Fri, 30 Dec 1994 16:03:31 -0500 (EST)
 A Technique for Directly Measuring the Gravitational Acceleration of Antihydrogen, By: Thomas J. Phillips, Duke University Durham Comments: Written version of invited contribution to LEAP'94: Third Biennial Conference on Low-Energy Antiproton Physics. The gravitational force on antimatter has never been directly measured. A method is suggested for measuring the acceleration of antimatter \overline{g} by measuring the deflection of a beam of neutral antihydrogen atoms in the Earth's gravitational field. While a simple position measurement of the beam could be used, a more efficient measurement can be made using a transmission interferometer. A 1% measurement of \overline{g} should be possible from a beam of about 100,000 atoms, with the ultimate accuracy being determined largely by the number of antihydrogen atoms that can be produced. A method is suggested for producing an antihydrogen beam appropriate for this experiment.

HIGH ENERGY PHYSICS -
 PHENOMENOLOGY, ABSTRACT HEP-
 PH/9509336 From: Michael Martin Nieto 505-
 667-6127 Date: Tue, 19 Sep 95 14:08:11 -
 0600



Antimatter Gravity and Antihydrogen Production Authors: Michael H. Holzscheiter , T. Goldman , Michael Martin Nieto Certain modern theories of gravity predict that antimatter will fall differently than matter in the Earth's gravitational field. However, no experimental tests of gravity on antimatter exist and all conclusions drawn from experiments on matter depend, at some level, on a specific model. We have proposed a direct measurement that would compare the gravitational acceleration of antiprotons to that of negatively charged hydrogen ions. Substantial progress towards the development of this experiment has been achieved. Based on our work a number of alternative proposals for measuring "g" on both charged and neutral antimatter have been made. We summarize the present status of our experiment and also discuss the steps that would be necessary to produce antihydrogen in an environment suitable for gravity measurements.

Hi Robert, I have one reference for you. The book is called "Suppressed Inventions and other Discoveries". It's an anthology edited by Jonathon Eisen. Authors include: Dr. Brian O'Leary, Christopher Bird, Jeanne Manning, Barry Lynes, and others. As well as Townsend Brown, the inventors/doctors (as well as inventions the book also covers various cancer treatments which have had research suppressed) who are discussed include Naessons, RifeHoxsey, Gerson, Tesla, Brown, Reich and others.

The book covers free energy, various "unfree" though different motive technologies, cancer cures which have worked but not seen approval by the AMA, Roswell, the Mars face, and as a delight to conspiracy buffs, there are also chapters on how various Government bodies have suppressed these technologies, as well as how the AMA came to be all powerful in the field of suppressing alternate treatments.

The book is published by:
Auckland Institute of Technology Press
Private Bag 92006

Auckland, New Zealand

ISBN No. 0-9583334-7-5

For further research, consult the following sources:

Fer-de-Lance by T.E. Beardon
Tesla Book Company
P.O. Box 121873
Chula Vista, CA 91912 USA

Leading Edge Research Group
(Leading Edge Journal #77 12/94)
P.O. Box 7530 Ste 58
Yelm, Washington 98597 USA

Nexus Magazine
P.O. Box 66
8400 AB Gorredijk
The Netherlands
Tel/Fax: 31-(0)5133-5567

The information on the electrogravitics reference list which is of particular interest to me are the Laithwaite and Wallace references. I think my work (Electrical-Dipole Theory of Gravitation) explains what they were observing and why. Here are some additional references. -- Ralph Sansbury


Fischbach, Sudarsky, Szafer, Talmadge, and Aronson in "Reanalysis of the Eotvos Experiment" (Phys Rev Let vol 56 p3 6/1/86)

J.H. Pratt and G.B. Airy 1855 Phil Trans v145

Fredrich Zollner, Explanation of Universal Gravitation through the Static Action of Electricity and the General Importance of Weber's Laws, 1882

Immanuel Velikovsky, Cosmos without Gravitation, 1964

V. A. Bailey In the May 14 , 1960 issue of Nature



- P.M.S. Blackett In the May 17, 1947 issue of Nature
- T. Gold in a later issue (April 2, 1949) of Nature
- Henry Wallace US patent number 3 626 605
- P.S. Wesson Phys Rev D v23 p1730 (1981)
- Sansbury R.N. Electrical Engineering Times (12/28/87)
- Sansbury R.N. US patent number 4,355,195
- Sansbury R.N. Rev. Sci. Instr. (3/85)
- Bartlett D.F. Rev.Sci. Instr. (10/90)
- Peter Graneau, Nature v295 1982 p311
- Weiskopf M.C., Carrico, Gould, Lipworth and Stein, Physical Review Letters 1968, vol21 p1645
- Coles and Good, Physical Review 1946 p979
- Kaufmann W. p502 in World of the Atom by H. Bourse and L. Motz
- W.J. Duffin, Electricity and Magnetism Wiley 1973
- R.A. Tricker, Early Electrodynamics Pergamon Oxford 1965 -----
-

Paper: gr-qc/9410019
 From: Peter Marzlin Date: Mon, 17 Oct 94
 12:50:28 +0100
 THE DIPOLE COUPLING OF ATOMS AND LIGHT IN GRAVITATIONAL FIELDS, Karl-Peter Marzlin, 10 pages, LaTeX
 The dipole coupling term between a system of N particles with total charge zero and the electromagnetic field is derived in the presence of a weak gravitational field. It is shown that the form of the coupling remains the same as in flat space-time if it is written with respect to the proper time of the observer and to the measurable field components. Some remarks

concerning the connection between the minimal and the dipole coupling are given.

The level of difficulty in the above paper is well beyond my grasp. But what is clear is that it presents an analysis which strongly suggests that the textbook wavefunctions for electrons within atomic matter can be best described by the dipole coupling rather than the coulomb gauge. The paper also relates the dipole coupling to a weak gravitational field. The last paragraph of the paper provides substance to the idea that gravity is at least in part, an electric dipole phenomena. Here is the last paragraph:

"It is interesting to make a comparison of the present results with the well known formal equivalence between the Maxwell field in curved space and in a dielectric medium (23). In this approach one defines a formal dielectric displacement vector to describe the influence of gravity on the Maxwell field. In the absence of particles, i.e. for vanishing polarisation P, the formal electric displacement agrees with the vector delta defined above. Also the coupling of the Poynting vector to the rotation occurs in the energy of the formal Maxwell field."

The paper referenced (23) is:

A.M. Volkov, A.A. Izmest'ev, and G.V. Skrotskii, Soviet Physics JETP 32, page 686, (1971)

Note: There are a variety of other theories and experiments which attempt to show that a static gravitational field is identical to that which results from electric dipole moments -- a polarisation of the vacuum. And conversely, it is well known that if you accelerate a dielectric material, or in "equivalence" subject a dielectric material to a gravitational field or other mechanical force -- an electric field due to dipole moment (polarisation P) will be generated within the material. This effect is especially prevalent in structured crystal dielectrics (piezoelectric materials), which are used as transducers in accelerometer sensors. You can also find piezoelectric material, and

conversion of mechanical force to a high voltage electric field, in push-button spark igniters used on gas grills and cigarette lighters.

Here's a thought. To enlighten those folks who continue to stubbornly try to debunk the evident relationship between gravitation and electromagnetics -- insert one of these spark igniters in a neuro-sensitive body cavity, and click it as many times as necessary.

One issue with the electrostatic dipole hypothesis is that once the magnetic effects of spin etc have been considered there is no evidence of such dipoles inside atomic nuclei and electrons. However if magnetic properties of nuclei and electrons can be represented in terms of electrostatic dipoles as recent experiments and theoretical discussion seem to indicate then this objection is avoided. The dipole can be produced by a negatively oriented particle orbiting a positive central particle so that the combination has a net positive charge (see Rev Sci Instr Mar 1985 and Geomagnetism: Gravity Measured by Magnetic Materials, ICP Press, Box 492 NY NY 10185 \$25US 1994 by R Sansbury) An added benefit: the observed quadrupole in nuclei and electrons makes more sense in a physically real Taylor expansion by the inclusion of an observed dipole term as well; that is the dipole term is not observed because its effects are wrongly attributed to another cause, magnetism; thus magnetism is properly regarded as a derived apparently separate force like the Coriolis sideways force on bodies moved radially on a rotating platform.

-- Ralph Sansbury

About electric dipole precession. The article "Electricity" in Britannica includes a resonance equation for dipole precession in dielectrics. It was identical in form to the one used in magnetic resonance, except for the obvious differences in units. Dielectric precession (resonance) frequencies were in the optical range.


Brown didn't use resonance; but he did use a steady frequency. His frequency, too, would damp out if it were discontinued. Greater results than Brown's could probably be achieved with lasers. But I doubt you'll find a better description of dielectric dipole resonance. The Britannica article gives the mathematics.

van der Waals force (J.D. van der Waals) -- Forces responsible for the non-ideal behavior of gases, and for the lattice energy of molecular crystals. There are three causes: dipole-dipole interaction; dipole-induced dipole moments; and dispersion forces arising because of small instantaneous dipoles in atoms.

"The Electric Dipole Moment of the Electron", Bernreuther & Suzuki, Reviews of Modern Physics, April 1991 vol 63 no 2 -- An electron or any other elementary particle can possess an electric moment (EDM) only by virtue of an interaction that violates parity and time-reversal invariance. The question of whether an electron EDM exists is thus related directly to the unsolved problem of CP violation. According to the standard model, in which CP violation is accounted for in terms of the Kobayashi-Maskawa matrix, the electron EDM is predicted to be far too small to be observed experimentally. However, a number of alternative theoretical models of CP violation predict larger values of the electron EDM. These models are of special interest now, when experimental limits on the electron EDM are improving substantially.

"The Electron Electric Dipole Moment for a CP-violating Neutral Higgs Sector", J.F. Gunion, Physics Letters: Part 8, Nov 8 1990

"New Experimental Limit on the Electron Electric Dipole Moment", Abdullah & Commins, Physical Review Letters, Nov 5 1990



"The Standard Model Prediction for the Electric Dipole Moment of the Electron", F. Hoogeveen, Nuclear Physics B, Sep 10 1990

"Electric Dipole Moment of the Electron and the Neutron", S.M Barr, Physical Review Letters, July 2 1990, Vol 65 No 1

"Effective Hamiltonian for the Electric Dipole Moment of the Neutron", Boyd, Gupta & Trivedi, Physics Letters: Part 8, May 24 1990

"A search for the Electric Dipole Moment of the Neutron", K.F. Smith, Physics Letters: Part 8, Jan 4 1990, Vol 234 No 1/2

"Interpretation of the Neutron Electric Dipole Moment: Possible Relationship to Epsilon", Booth, Briere & Sachs, Physical Review D Jan 1 1990, Vol 41 No 1

"Inclusion of the Toroidal-Moment Contribution in the Probability of the Electric Dipole Transition", R.G. Nazmitdinov, Soviet Journal of Nuclear Physics, Sep 1 1990, Vol 53 No 2 -----

But what is the thing in atomic nuclei that collectively produces the gravitational field of the Earth etc. and which causes individual nuclei to react in the prescribed manner? The hypothesis proposed is that atomic nuclei contain small electrostatic dipoles ($10^{-37} C.m$) with radial and longitudinal components transverse to the west to east spinning direction of the Earth etc. Such dipoles explain the nuclear magnetic moment and electrostatic quadrupole moment inferred from the hyperfine spectra emitted by some excited atoms and the deflection of molecules such as orthohydrogen in a magnetic field (but not parahydrogen because the magnetic moments are anti parallel in pairs and cancel)

The Cavendish measurement of the horizontal gravitational force between two lead spheres instead of being attributed to the small masses of each can be attributed to the small horizontal component of the radial force, directed to the center of the Earth, due to the


mass of the Earth on each of the small masses. That is gravity is not a property of mass per se but only of spinning mass.

The atomic nuclei of all elements, except iron, cobalt, and nickel primarily, tend to line up in the direction of the surrounding atomic nuclei when the bulk object of which they are a part is moved but in the case of the magnetic elements the bulk material must also move to complete the required alignment, hence the north south and downward movement of a magnetized steel compass needle. Hence the Wilson-Blackett proportionality between the angular momentum of planets, stars etc and their magnetic moment where the constant of proportionality is the square root of the gravitational constant divided by the speed of light. For more information see Science News Aug 6 '94 p82. - Ralph Sansbury

Edward Teller, "Electromagnetism and Gravitation", Proceedings of the National Academy of Science, Vol 74 No 4, Pages 2664-2666.

In this paper Dr Teller suggests some clues about the coupling between electromagnetism and gravitation. In the first part of his paper Teller describes how an electric field due to polarization can be induced in a dielectric material which is subject to angular or linear acceleration, or if subject to a gravitational field. In the second part of the paper Teller describes, using purely dimensional analysis, how a magnetic field might be produced by a spinning mass. He also comments that the magnitude of this magnetic field might be exceedingly small, and notes that a "numerical" factor could exist which might act to increase the magnitude of the field.

(Note: It is speculated by others that alignment of microscopic particles with the macroscopic spin axis of the earth, could result in a large "numerical" factor. Fact is, the earth does have a fairly large measurable magnetic field, about which there are a variety of theories as to the origin.)



Paper: hep-th/9506049

From: HORIE@dipmza.physik.Uni-Mainz.DE

Date: Thu, 08 Jun 1995 11:23:23 +0100

Title: New Insight into the Relation between Torsion and Electromagnetism Author: Kenichi Horie (Mainz Univ.)

Report-no: MZ/TH 95-16

In several unified field theories the torsion trace is set equal to the electromagnetic potential. Using fibre bundle techniques we show that this is no leading principle but a formal consequence of another geometric relation between space-time and electromagnetism.

HIGH ENERGY PHYSICS - THEORY, ABSTRACT HEP-TH/9409018 From:

HORIE@VIPMZW.physik.Uni-Mainz.DE

Date: Sat, 03 Sep 1994 10:27:48 +0100

GEOMETRIC INTERPRETATION OF ELECTROMAGNETISM IN A GRAVITATIONAL THEORY WITH SPACE-TIME TORSION BY KENICHI HORIE, INSTITUT FUR PHYSIK, JOHANNES GUTENBERG-

-UNIVERSITAT MAINZ, D--55099 MAINZ, GERMANY, A complete geometric unification of gravity and electromagnetism is proposed by considering two aspects of torsion: its relation to spin established in Einstein-Cartan theory and the possible interpretation of the torsion trace as the electromagnetic potential. Starting with a Lagrangian built of Dirac spinors, orthonormal tetrads, and a complex rather than a real linear connection we define an extended spinor derivative by which we obtain not only a very natural unification, but can also fully clarify the nontrivial underlying fibre bundle structure. Thereby a new type of contact interaction between spinors emerges, which differs from the usual one in Einstein-Cartan theory. The splitting of the linear connection into a metric and an electromagnetic part together with a characteristic length scale in the theory strongly suggest that gravity and electromagnetism have the same geometrical origin.

"Gauge Invariant Electromagnetic Coupling with Torsion Potential", Richard T. Hammond, General Relativity and Gravitation, Vol 23 No 11 1991 Electromagnetism is coupled to torsion in a gauge invariant manner by relaxing minimal coupling and introducing into the Lagrangian a term bilinear the electromagnetic field tensor and its torsion potential. The resulting coupling between electromagnetism and torsion is examined and a solution corresponding to traveling coupled waves is given. Since torsion is usually regarded as resulting from the spin of a body, this might establish a classical relationship between charge and spin. The results suggest that the effect should be looked for in high intensity electric fields of low frequency.

"Detecting Torsion from Massive Electrodynamics", L.C. Garcia de Andrade, and M. Lopes, General Relativity and Gravitation, Vol 25 No 11 1993 A new method of detecting torsion in the case of massive electrodynamics is proposed. Several authors have proposed methods for the detection of torsion in theories of the Einstein-Cartan type, and also in theories where the torsion field propagates. These theories are based on the studies of Dirac test particles, which have spin like the electron, and the gyroscope-like precession of these atomic particles. The interaction energy between the torsion vector Q, and an electric dipole p, is given by (p dot Q).

AUTHOR(s): de Andrade, L.C. Garcia

TITLE(s): Electron gyroscopes to test torsion gravity?

In: Il nuovo cimento delle societa italiana di fisica OCT 01 1994 v 109 n 10 Page: 1123

AUTHOR: De Sabbata, Venzo.

TITLE: Spin and Torsion in Gravitation by Venzo de Sabbata, and C. Sivaram.

PUBL.: Singapore ; River Edge, NJ : World Scientific,

FORMAT: xii, 313 p. : ill. ; 23 cm.

DATE: 1994

SUBJECTS: Torsion, Gravitation

AUTHOR: De Sabbata, Venzo.

TITLE: Introduction to Gravitation
 by Venzo de Sabbata and Maurizio Gasperini.
 PUBL.: Singapore ; Philadelphia : World Scientific,
 FORMAT: ix, 346 p. : ill. ; 23 cm.
 DATE: 1985
 SUBJECTS: General relativity, Torsion, Gravitation

AUTHOR: NATO Advanced Study Institute on Cosmology and Gravitation (1979: Bologna, Italy)
 TITLE: Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity
 Edited by Peter G. Bergmann and Venzo De Sabbata. PUBL.: New York : Plenum Press : NATO Scientific Affairs Division,
 FORMAT: ix, 510 p. : ill. ; 26 cm.
 DATE: 1980
 SERIES: NATO Advanced Study Institutes Series v 58 Series B Physics

CONFERENCE :International Conference on Magnetic and Electric Resonance and Relaxation (1962: Eindhoven)

TITLE :Magnetic and electric resonance and relaxation; proceedings of the XIth Colloque Ampere, Eindhoven, July 2-7, 1962. PUBLISHED :Amsterdam, New York, North-Holland Pub. Co.; Interscience Publishers, 1963.

DESC :xi,789p. illus.,diags.,tables. 24cm.

The Lorentz-Dirac equation is a purely classical expression for the electromagnetic force on a point charge, including the self-force from the particle's own radiation. It's a strange equation, with solutions that are manifestly unphysical under certain circumstances. If you want to know more about it, you might want to look at:

S. Parrott, Relativistic Electrodynamics and Differential Geometry, Springer-Verlag, 1987.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9403058 PHYS. REV. D50 (1994 3867) carroll@marie.mit.edu (Sean Carroll) Tue, 29 Mar 1994 19:57:32 -0500

CONSEQUENCES OF PROPAGATING TORSION IN CONNECTION-DYNAMIC THEORIES OF GRAVITY, BY SEAN M. CARROLL AND GEORGE B. FIELD, 16 PAGES PLUS ONE FIGURE (PLAIN TEX), MIT-CTP #2291.

We discuss the possibility of constraining theories of gravity in which the connection is a fundamental variable by searching for observational consequences of the torsion degrees of freedom. In a wide class of models, the only modes of the torsion tensor which interact with matter are either a massive scalar or a massive spin-1 boson. Focusing on the scalar version, we study constraints on the two-dimensional parameter space characterizing the theory. For reasonable choices of these parameters the torsion decays quickly into matter fields, and no long-range fields are generated which could be discovered by ground-based or astrophysical experiments.

GENERAL RELATIVITY & QUANTUM COSMOLOGY, ABSTRACT GR-QC/9304047 From: KUBYSHIN%EBUBEBCM1.BITNET@FRMO P11.CNUSC.FR Date: Sun, 02 May 93 12:55:30 BCN

INVARIANT CONNECTIONS WITH TORSION ON GROUP MANIFOLDS AND THEIR APPLICATION IN KALUZA-KLEIN THEORIES, KUBYSHIN YU.A., MALYSHENKO V.O. AND MARIN RICOY D. Invariant connections with torsion on simple group manifolds S are studied and an explicit formula describing them is presented. This result is used for the dimensional reduction in a theory of multidimensional gravity with curvature squared terms on M^4 times S. We calculate the potential of scalar fields, emerging from extra components of the metric and torsion, and analyze the role of the torsion for the stability of spontaneous compactification.



Subject: Antigravity in Jane's
From: "Terry Colvin"

"All those interested in advanced propulsion concepts should check out Jane's Defence Weekly, 10 June 1995. An article discusses anti-gravity schemes and shows drawings of sauceroid vehicles from British Aerospace among others. Area 51 is mentioned, as well as an unclassified paper done for the USAF by Science Applications International Corp. in 1990. The subject was [Electric Propulsion], a[n] euphemism for anti-gravity according to Jane's. Michael Flora"

Anti-Gravity for Real -- Discussed in Jane's Defence Weekly

Jane's Defence Weekly is a most respected journal in the defense industry. Jane's has often been the first to break the news about secret development of radically new technologies and equipment.

Jane's Defence Weekly of 10 June 1995, has an article about advanced aerospace technologies, written by Nick Cook. The idea of anti-gravity is taken seriously and is auspiciously present throughout the article -- including three artist renditions of future anti-gravity based craft.

The Jane's article commences with a mention of anti-gravity technology, and also ends with a few paragraphs discussing anti-gravity. In between is the bulk of the article, which consists of discussion of "conventional" subjects, including: Hypersonics, Gas Turbine Inrements, The Super Cockpit, and Stealth.

At the start of the Jane's article there is some information from the Gravity Rand Report on Electrogravitics which was done for the USAF in 1956, and was recently declassified. Here's an excerpt from the beginning of the Jane's article.

Take this example from a specialist US aviation magazine in 1956. "We're already working with equipment to cancel out gravity," Lawrence D Bell, founder of the company that bears his name was quoted as saying. Bell, apparently, was not the only one working in this field. Others said to be seeking to master this arcane 'science' included the Glenn L Martin Company, Convair, Lear, and Sperry Gyroscope. Within a few years we were assured, aircraft, cars, submarines and power stations would all be driven by this radical new propulsion technology. Sadly it was not to be.

Here's the ending section of the Jane's article.

BEYOND 2001

Groom Lake Nevada is the epicentre of classified USAF research into Stealth and other exotic aerospace technologies. Several years after the collapse of the Soviet threat, activity and investment at this remote, highly secret air base (so secret its presence is, as yet, unacknowledged by the US government) is still on the increase. While research into less sensitive technologies such two-dimensional thrust-vectoring and advanced short take-off and vertical landing (ASTOVL) are pursued in the open at nearby Edwards AFB in California, Groom Lake is set to hang onto its secrets. The USAF's recent confiscation of 1600 acres of public land bordering the facility is consistent with the Pentagon's desire to maintain its lead in quantum leap technologies -- some of which, according to well qualified observers in and around the Nevada area, defy current thinking into the predicted direction of aerospace engineering.

That aerospace companies continue to look at highly radical alternative air vehicle concepts is evidence of the ongoing quest for breakthrough designs. Glimpses into this world are rare, but provide some insight into likely 21st century research activity. The 1990 unclassified 'Electric Propulsion Study' (a quest for antigravity propulsion system by another name) conducted by the USA's Science Application International Corp (SAIC) on behalf of USAF's then Astronautics Laboratory at Edwards AFB shows that USAF



visionaries are still being given free reign. Until recently BAe (British Aerospace) also provided internal resources for its own anti-gravity studies and even went so far as to outline this thinking with artists' concepts -- a case of Lawrence Bell's vision perhaps being not so wide of the mark after all.

Before he died, Ben Rich, who headed Lockheed's Skunk Works from 1975-1991, was quoted as saying: "We have some new things. We are not stagnating. What we are doing is updating ourselves, without advertising. There are some new programmes, and there are certain things -- some of them 20 to 30 years old -- that are still breakthroughs and appropriate to keep quiet about. Other people don't have them yet."

Thirty years from now, we may still not know the half of what is currently being tested in and around Groom Lake.

Copyright 1995, Jane's Defence Weekly, All rights reserved.

/* The above information is transmitted under the "Fair Use" rulings of the 1976 Copyright Act for NON-profit academic and general information purposes. */

 AUTHOR(s):
 McIntosh, C.B.G. Arianrhod, R. Wade, S.
 TITLE(s): Electric and magnetic Weyl tensors: classification and analysis.
 In: Classical and quantum gravity.
 JUN 01 1994 v 11 n 6 Page 1555

AUTHOR(s): Arianrhod, R. Lun, A.W.-C. McIntosh, C.B.G.
 TITLE(s): Magnetic curvatures.
 In: Classical and quantum gravity.
 SEP 01 1994 v 11 n 9 Page 2331

AUTHOR(s): Arianrhod, R. McInthosh, C.B.G.
 TITLE(s): Principle null directions of Petrov type I Weyl spinors: geometry and symmetry.
 In: Classical and quantum gravity.

AUG 01 1992 v 9 n 8 Page 1969

AUTHOR(s): Hoenselaers, C. Perjes, Z.
 TITLE(s): Multipole moments of axisymmetric electrovacuum spacetimes.
 In: Classical and quantum gravity.
 OCT 01 1990 v 7 n 10 Page 1819

AUTHOR(s): de Felice, Fernando Yu, Yunqiang Fang, Jing
 TITLE(s): Relativistic charged spheres.
 In: Monthly notices of the royal astronomical society NOV 01 1995 v 277 n 1 Page: L17

AUTHOR(s): de Felice, Fernando
 TITLE(s): Dynamics on a rotating disk.
 In: Physical review. A, Atomic, molecular, and optical physics NOV 01 1995 v 52 n 5 Page 3452

AUTHOR(s): de Felice, Fernando Yu, Yunqiang Coriasco, Sandro
 TITLE(s): The Lynden-Bell and Katz Definition of Gravitational Energy: Applications to Singular Solutions. In: General relativity and gravitation.
 AUG 01 1994 v 26 n 8 Page 813

AUTHOR(s): Cavaglia, Marco de Alfaro, Vittorio de Felice, Fernando
 TITLE(s): Anisotropic wormhole: Tunneling in time and space.
 In: Physical review d: particles, fields, gravitation JUN 15 1994 v 49 n 12 Page 6493

AUTHOR(s): de Felice, Fernando
 TITLE(s): Rotating frames and measurements of forces in general relativity.
 In: Monthly notices of the royal astronomical society SEP 15 1991 v 252 n 2 Page 197

AUTHOR(s): Hammond, Richard
 TITLE(s): Tetrad Formulation of Gravity with a Torsion Potential.
 In: General relativity and gravitation.
 NOV 01 1994 v 26 n 11 Page 1107

AUTHOR(s): Hammond, Richard
 TITLE(s): Spin, Torsion, Forces.
 In: General relativity and gravitation.
 MAR 01 1994 v 26 n 3 Page 247



AUTHOR(s): Hammond, Richard T.
 TITLE(s): Gauge Invariant Electromagnetic Coupling with Torsion Potential.

In: General relativity and gravitation.
 NOV 01 1991 v 23 n 11 Page 1195

AUTHOR(s): Hammond, Richard T.
 TITLE(s): Magnetic Charge Type Equations from Torsion.
 In: General relativity and gravitation.
 SEP 01 1991 v 23 n 9 Page 973

AUTHOR(s): Hammond, Richard T.
 TITLE(s): Dynamic Torsion from a Linear Langrangian.
 In: General relativity and gravitation.
 APR 01 1990 v 22 n 4 Page 451

AUTHOR(s): Ringermacher, H.I.
 TITLE(s): An electrodynamic connection.
 In: Classical and quantum gravity.
 SEP 01 1994 v 11 n 9 Page 2383

AUTHOR(s): Anandan, J. Hagen, C.R.
 TITLE(s): Neutron acceleration in uniform electromagnetic fields.
 In: Physical review. A, Atomic, molecular, and opt OCT 01 1994 v 50 n 4 Page 2860

AUTHOR(s): Anandan, J.
 TITLE(s): Relativistic gravitation and superconductors.
 In: Classical and quantum gravity.
 JUN 01 1994 v 11 n 6A Page 23

AUTHOR(s): Georgiou, A.
 TITLE(s): Rotating Einstein-Mazwell fields: smoothly matched exterior and interior spacetimes with charged dust and surface layer.
 In: Classical and quantum gravity.
 JAN 01 1994 v 11 n 1 Page 167

AUTHOR(s): Unnikrishnan, C.S.
 TITLE(s): Experimental gravitation in India: progress and challenges.
 In: Classical and quantum gravity.
 JUN 01 1994 v 11 n 6A Page 195

AUTHOR(s): Cowsik, R. Tandon, S.N. Unnikrishnan, C.S.

TITLE(s): Limit on the strength of intermediate-range forces coupling to isospin.
 In: Physical review letters.
 NOV 07 1988 v 61 n 19 Page 2179

AUTHOR(s): Banerjee, A. Panigrahi, D. Chatterjee, S.
 TITLE(s): Evolution of Kaluza-Klein inhomogeneous model with a cosmological constant.
 In: Journal of mathematical physics.
 JUL 01 1995 v 36 n 7 Page 3619

AUTHOR(s): Chatterjee, S. Panigrahi, D. Banerjee, A.
 TITLE(s): Inhomogeneous Kaluza-Klein cosmology.
 In: Classical and quantum gravity.
 FEB 01 1994 v 11 n 2 Page 371

It might interest antigravity researchers to know (for those not already aware) that Professor ER Laithwaite, a respected British electrical engineer, has been doing work on this very subject for decades, but when he tried to demonstrate the viability of his theories to his peers their closed minds closed ranks and ridiculed his efforts as fantasy. Laithwaite lost cred with the scientific community and had to rely just on one or two close associates in semi-secrecy.

A recent (a year or two ago) series of TV programmes in Britain (on controversial scientific discoveries that have yet to accepted by the scientific establishment as worthy of further research and funding) ran an episode on Laithwaite. He claimed that gyroscopes could transfer mass.

I know of one book he wrote: Transport Without Wheels published by Paul Elek in 1977 ISBN 0236400665 (info from an old note I made) though this is NOT specifically about his antigravity theories (I remember that it concentrated on propulsion via electrical rails)



I would be most interested in learning about anything he (or anyone else) might have written specifically on his antigravity work.

- George Szaszvari

"Propulsion by Gyro", Eric Laithwaite, Space, Sep 1989 Vol 5 No 5 In an attempt to reveal the strange, hidden properties of gyroscopes, Professor Eric Laithwaite explains the physics behind the idea that a propulsion system could be built using gyros.

-

RS Electrogravitic References: Part 16 of 19.

AUTHOR(s): Ljubicic, A. Zovko, N.
TITLE(s): Lorentzian component of the fifth force.
In: Fizika B.

JAN 01 1992 v 1 n 1 Page: 1

AUTHOR(s): Bertotti, B. Sivaram, C.
TITLE(s): Radiation of the \leftrightarrow field.
In: Il Nuovo cimento della Societa italiana di fisic NOV 01 1991 v 106 n 11 Page: 1299

AUTHOR(s): Fujii, Y.
TITLE(s): The Theoretical Background of the Fifth Force.
In: International journal of modern physics. a, pa AUG 20 1991 v 6 n 20 Page: 3505

AUTHOR(s): Mannheim, Philip D.
TITLE(s): General Relativity and Fifth Force Experiments.
In: Astrophysics and space science.
JUL 01 1991 v 181 n 1 Page: 55

AUTHOR(s): Cho, Y.M. Park, D.H.
TITLE(s): Fifth Force from Kaluza-Klein Unification.
In: General relativity and gravitation.
JUL 01 1991 v 23 n 7 Page: 741

AUTHOR(s): Fujii, Y.
TITLE(s): Locally varying particle masses due to a scalar fifth-force field.
In: Physics letters: [Part B]
FEB 14 1991 v 255 n 3 Page: 439

AUTHOR(s): Hagiwara, Yukio
TITLE(s): No gravimetric evidence for the fifth force?

Summary: TEXT IN JAPANESE
In: Chigaku zasshi =
1991 v 100 n 3 Page: 429

AUTHOR(s): Cho, Y.M. Park, D.H.
TITLE(s): Higher-dimensional unification and fifth force.
In: Il nuovo cimento delle societa italiana di fisic AUG 01 1990 v 105 n 8/9 Page: 817

AUTHOR(s): Sardanashvily, G.
TITLE(s): The Gauge Model of the Fifth Force (E,SUM).
In: Acta physica Polonica, B.
AUG 01 1990 v 21 n 8 Page: 583

AUTHOR(s): Schimdt, H.-J.
TITLE(s): Fifth force, dark matter, and fourth-order gravity.
In: Europhysics letters.
AUG 01 1990 v 12 n 7 Page: 667

AUTHOR(s): de Sabbata, Venzo Sivaram, C.
TITLE(s): Fifth Force as a Manifestation of Torsion.
In: International journal of theoretical physics.
JAN 01 1990 v 29 n 1 Page: 1

AUTHOR(s): Timoshenko, E.G.
Sardanashvily, G.A.
TITLE(s): Gauge model for the fifth force.
In: Moscow university physics bulletin.
1990 v 45 n 4 Page: 73

AUTHOR(s): Hagiwara, Yukio
TITLE(s): The fifth force-doubt about newton's gravitational law
Summary: TEXT IN JAPANESE
In: Chigaku zasshi =
1990 v 99 n 3 n 904 Page: 263

AUTHOR(s): Gasperini, M.
TITLE(s): Phenomenological consequences of a direct fifth force coupling to photons.



In: Physical review. D, Particles and fields. NOV 15 1989 v 40 n 10 Page: 3525

AUTHOR(s): Gasperini, M.

TITLE(s): Fifth force and the gravimagnetic hypothesis..

In: Physics letters: [part A]

OCT 02 1989 v 140 n 6 Page: 271

AUTHOR(s): Fayet, P.

TITLE(s): The fifth force charge as a linear combination of baryonic, leptonic (or B-L) and electric charges.

In: Physics letters: [Part B]

AUG 17 1989 v 227 n 1 Page: 127

AUTHOR(s): Mufti, A. Kwong, N.H. Schaudt, K.J.

TITLE(s): Search for the fifth force using Gauss's law.

In: Physics letters: [part A]

JUL 31 1989 v 139 n 3 / 4 Page: 115

AUTHOR(s): Bizzeti, P.G.

TITLE(s): Search for a Composition-Dependent Fifth Force.

Summary: A differential accelerometer consisting of a solid sphere floating freely inside a stratified saline solution has been used to search for a composition-dependent force, originated by a mountain relief. No evidence of such a force has been obtained.

In: Physical review letters.

JUN 19 1989 v 62 n 25 Page: 2901

AUTHOR(s): T.M. Aliev, Dobroliubov, M.I. Ignatiev, A. Yu.

TITLE(s): Do Kaon decays constrain the fifth force?

In: Physics letters: [Part B]

APR 20 1989 v 221 n 1 Page: 77

AUTHOR(s): Riveros, C. Loguidice, E. A. Vucetich, H.

TITLE(s): On differential fifth force measurements.

In: Physics letters: [part A]

APR 17 1989 v 136 n 7/8 Page: 343

AUTHOR(s): Kuroda, K. Mio, N.

TITLE(s): A Free Fall Interferometer to Search for a Possible Fifth Force.

In: IEEE transactions on instrumentation and measure APR 01 1989 v 38 n 2 Page: 189

AUTHOR(s): Faller, J. E. Fischbach, E. Fujii, Y.

TITLE(s): Precision Experiments to Search for the Fifth Force.

In: IEEE transactions on instrumentation and measure APR 01 1989 v 38 n 2 Page: 180

AUTHOR(s): Stubbs, C. W. Adelberger, E. G. Heckel, B. R.

TITLE(s): Gravitation and Astrophysics: Limits on composition-dependent interactions using a laboratory source: Is there a "fifth force" coupled to isospin? In: Physical review letters. FEB 06 1989 v 62 n 6 Page: 609

TITLE(s): Alternate source of fifth force challenged.

In: Science news.

OCT 01 1988 v 134 n 14 Page: 214

TITLE(s): The stimulation of the fifth force.

In: Nature.

SEP 29 1988 v 335 n 6189 Page: 393

One of the first "scientific" DOGMAS fed to new physics students is the doctrine about "projectile motion". Students are given several formulas or equations from which they can precisely calculate how high and far a projectile will travel given its initial speed and angle from the ground. But the results are NOT so absolute as students are led to believe, even if they take into account air resistance and Coriolis effects.

Recent experiments have shown that if the projectile is SPINNING at HIGH SPEED, (at least 27,000 RPM), [axis of spin coinciding with line of projection], the projectile will travel HIGHER and FARTHER than predicted by Newtonian mechanics! Similarly, experiments with falling gyroscopes have



shown that a gyroscope whose enclosed rotor is spinning at high speed (about 27,000 RPM) falls AT A DIFFERENT RATE than when the same gyroscope falls with rotor NOT spinning. The AMOUNT of DEVIATION might depend on the MATERIAL COMPOSITION of the projectile or rotor, as suggested in the text of U.S. Patent #3,626,605, "Method and Apparatus for Generating a Secondary Gravitational Force Field", by Henry W. Wallace, Dec. 14, 1971.

These DEVIATIONS are EASILY REPRODUCABLE, and effectively DISPROVE the OVER-HYPED "General Theory of Relativity" which states that gravity results from a "warping or distortion of space" caused by the MERE PRESENCE of mass. - Robert McElwaine

The late Henry W Wallace died April 1994. Fellow researchers at GE were not "happy" with his research regarding gravitational fields. An interesting article was written in The New Scientist circa 1980 about Wallace's work.

-- Ron Kita

Henry Wallace was an engineer at General Electric about 25 years ago, and developed some incredible inventions relating to the underlying physics of the gravitational field. Few people have heard of him or his work.

US Patent #3626605 -- "Method and Apparatus for Generating a Secondary Gravitational Force Field"
Awarded to Henry Wm Wallace of Ardmore PA Dec 14, 1971

US Patent #3626606 -- "Method and Apparatus for Generating a Dynamic Force Field"
Awarded to Henry Wm Wallace of Ardmore PA Dec 14, 1971

US Patent #3823570 -- "Heat Pump" (based on technology similar to the above two inventions)

Awarded to Henry Wm Wallace of Freeport NY July 16, 1973

Wallace discovered that a force field, similar or related to the gravitational field, results from the interaction of relatively moving masses. He built machines which demonstrated that this field could be generated by spinning masses of elemental material having an odd number of nucleotides -- i.e. a nucleus having a multiple half-integral value of h-bar, the quantum of angular momentum. Wallace used bismuth or copper material for his rotating bodies and "kinnemassic" field concentrators. Aside from the immense benefits to humanity which could result from a better understanding of the physical nature of gravity, and other fundamental forces, Wallace's inventions could have enormous practical value in countering gravity or converting gravitational force fields into energy for doing useful work. So, why has no one heard of him? One might think that the discoverer of important knowledge such as this would be heralded as a great scientist and nominated for dynamite prizes. Could it be that his invention does not work? Anyone can get the patents. Study them -- Wallace -- General Electric -- detailed descriptions of operations -- measurements of effects -- drawings and models -- it is authentic. If you're handy you can even build it yourself. It does work.

So what is going on?

One explanation I've heard is that Wallace ran up against the politics of science, as dictated in the late 1960's by the power-block at Princeton, who were primarily interested in promoting the ideas of their main man, Einstein, and the gravitation-is-geometry paradigm. Maybe there is some truth to this story. Nowadays, there seems to be a piss-pot full of theoretical physicists working on abstract geometrical theories and other absurdly difficult mental masturbations, while no one seems to have made any effort to provide a theoretical explanation of the physics of a nuts-and-bolts invention which could have enormous practical value. Maybe we can blame it on the Princeton folks, but I'm more inclined to believe that our defense industry black project community has confiscated and

suppressed knowledge of Wallace's discoveries. All done of course under the most honorable and sacred banner of national security. Well, it's been 25 years. We ought to be real secure by now. Isn't it way past time for some trickle down benefits to real people? ---

There are two paragraphs about the Wallace inventions in the Electric Propulsion Study by Dr Dennis Cravens, prepared in 1991. Cravens had this to say about Wallace's work:

ROTATIONAL ALIGNMENT - Nuclei can also be aligned by rotation. Henry Wallace claimed some unusual effects assigned to electromagnetic and gravitational couplings. This was in three US patents (3823570, 3626605, and 362606). The assertion was that the application of a rotational force on a material of half-integral spin would result in a reorientation of the nuclear structure and could be utilized for "altering its gravitational attraction toward other bodies, separation of isotopes by distinguishing between nuclei according to their nucleon content..." The patents are written in a very believable style which includes part numbers, sources for some components, and diagrams of data.

Attempts were made to contact Wallace using patent addresses and other sources but he was not located nor is there a trace of what became of his work. However, should the work be real it may furnish a novel experimental approach to experimental design. The concept can be somewhat justified on general relativistic grounds since rotating frames of time varying fields are expected to emit gravitational radiation. Even if the work does not give a direct gravitational coupling it may furnish a new method for nuclear spin alignment.

An article about the Wallace patents appeared in the British magazine "New Scientist" in February 1980. This was written nearly ten years after Wallace was awarded his patents. Here's a paragraph from the article.

"Although the Wallace patents were initially ignored as cranky, observers believe that his invention is now under serious but secret investigation by the military authorities in the US. The military may now regret that the patents have already been granted and so are available for anyone to read."


I know -- it's a tease. And the rest of the article is the same way. It provides barely enough information to jab your psyche a little, and not nearly enough to get you off your comfortable ass. And who knows who the anonymous party of "observers" are, who believe that a secret investigation is underway by the military -- or whether these observers even exist at all. None the less, the New Scientist has a fairly well established track record for accurate identification of new science trends and issues. And, while the editors of this British journal may be prone to enjoyment of gossip and innuendo, it generally turns out to be grounded in truth.

NUCLEAR SPIN SELECTIVITY OF CHEMICAL REACTIONS A.L.Buchachenko, N.N.Semenov Institute of Chemical Physics, Russian Academy of Sciences

4 Ul. Kosygina, 117334 Moscow, Russian Federation, Fax: +7(095)938-2156 A property of spin selective reactions to sort the nuclei according to their spin and orientation, is discussed. The separation of spin (magnetic) and spinless (nonmagnetic) nuclei forms the basis for the magnetic isotope effect, the separation of nuclei according to their orientation and creation of nuclear alignment in reaction products is a basis for the chemically induced nuclear polarisation phenomenon. Bibliography - 50 references.

Received 14 February 1995

NUCLEAR THEORY, ABSTRACT NUCL-TH/9601046 From:
spevak@TAUPHY.TAU.AC.IL
Date: Tue, 30 Jan 1996 17:18:34 +0200



Collective T- and P- Odd Electromagnetic Moments in Nuclei with Octupole Deformations

Author(s): N. Auerbach , V.V. Flambaum , V. Spevak Parity and time invariance violating forces produce collective P- and T- odd moments in nuclei with static octupole deformation. Collective Schiff moment, electric octupole and dipole and also magnetic quadrupole appear due to the mixing of rotational levels of opposite parity and can exceed single-particle moments by more than a factor of 100. This enhancement is due to two factors, the collective nature of the intrinsic moments and the small energy separation between members of parity doublets. The above moments induce T- and P- odd effects in atoms and molecules. Experiments with such systems may improve substantially the limits on time reversal violation.

The Hughes-Drever experiment was conducted in 1959-1960 independently by Vernon Hughes and collaborators at Yale University, and by Ron Drever at Glasgow University. In the Glasgow version, the experiment examined the ground state of the lithium-7 nucleus in an external magnetic field. The state has total angular momentum quantum number $3/2$, and thus is split into four equally spaced levels by the magnetic field. When the nucleus undergoes a transition between a pair of adjacent levels, the photon emitted has the same energy or frequency, no matter which pair of levels was involved. The result is a single narrow spectral line. Any external perturbation of the nucleus that is associated with a preferred direction in space, such as the motion of the Earth relative to the mean rest frame of the universe, will destroy the equality of the energy spacing between the four levels, since the nuclear wave functions of the four levels have different spacial dependencies relative to the magnetic field. Using nuclear magnetic resonance techniques, the experiments set a limit on the separation or spread in frequency of line that corresponded to a limit on anisotropy or bidirectional dependence in the energy of the nucleus at the level of one part in 10^{23} . -- Clifford Will,

Chapt 2 of The New Physics, edited by Paul Davies -----

Magnetic resonance in its various forms, NMR, EPR, and EFR, are all applied to relatively small specimens and, with the exception of EFR, are rarely applied to magnetic materials. EFR means Electron Ferromagnetic Resonance, and the best intro to this subject is by Vonsovskii. Curiously, there is no published data on EFR for large ferromagnetic specimens. A literature search at a campus of the University of California revealed nothing. F. Herlach has said that there is an 'open' literature and a 'closed' literature concerning magnetic research.

-- Larry Adams

A body which is spinning within a larger macroscopic body which is also spinning will tend to align the axis of its angular momentum with the angular momentum of the larger body.

For example, a gyroscope located on the earth, unless it is in a frictionless gimbal, with always try to precess due to the rotation of the earth into alignment with the earth's polar axis, at which point it will no longer precess due to earth rotation.

Another example, a cylinder of magnetic material spinning around its longitudinal axis will develop a magnetic field proportional to its angular velocity (Barnett Effect), because the angular momentum of the electrons in the material will attempt to precess and come into alignment with the macroscopic axis of the spinning cylinder, which also brings into alignment the magnetic moment of the electrons, some of which have unpaired spins (ferromagnetic), resulting in generation of a macroscopic magnetic field. Similarly, it is known that a static magnetic field itself contains angular momentum -- and spinning the source of the static field, whether a magnet or DC current loop, will result in a corresponding increase or decrease in the field strength.

Another example is the inventions of Henry Wallace. Wallace found that if you spin a

material which has an odd number of nucleotides, i.e. having an "unpaired" value of angular momentum, resulting in a nucleus with a multiple integer of a one-half value of quantum momentum. The spin in the nucleus will begin to line up with the macroscopic spin axis, and will create an unusual force field related to gravity -- which he call a "kinemassic" field.

Maybe I've missed it, but I've looked seriously, and there seems to be no information in undergraduate or graduate level physics reference books which mentions the relationship between macroscopic and microscopic angular momentum -- much less provides any analysis or explanation linking quantum angular momentum to macroscopic angular momentum. Why not? How does quantum angular momentum become organized from a microscopic to a macroscopic level? Has anyone ever published any work about this? I can't find any.

Date: Sun, 5 Nov 1995

From: James Youlton To: Robert Stirniman
Re: Angular Momentum and the Barnett Effect

On Wed, 1 Nov 1995, Robert Stirniman wrote:
>Maybe I've missed it, but I've looked seriously, and there seems to be no information in undergraduate or graduate level physics reference books which mentions the relationship between macroscopic and microscopic angular momentum -- much less provides any analysis or explanation linking quantum angular momentum to macroscopic angular momentum.

You're catching on. The subject of compound angular momentum, or internal and external angular momentum, or intrinsic and extrinsic angular momentum has been a repressed subject for about 2 and half decades. Add to that list, spherical pendulums, Coriolis effect, except as applied to ballistics and meteorology as used by the US military, and Shafer's pendulum, that neat little device used as the artificial horizon of aircraft.

>How does quantum angular momentum become organized from a microscopic to a macroscopic level? Has anyone ever published any work about this? I can't find any.

There isn't any that I know of, though back in the late fifties, there was a fellow named Edward Condon at the University of Colorado who was fairly proficient on the subject. So much so that he wrote the rotational dynamics section, called noninertial dynamics at the time, of the reference "The Handbook of Physics" which he also co-edited (Chapter 5). I don't recall offhand who the publisher was (Harcourt/Brace?), though it was endorsed by the American Institute of Physics. Later, when Mr Condon was the head of the USAF project 'Blue Book', he labored to suppress his own work when the directive was handed down from the Navy's Turtle Island project.


-- James Youlton

Condon directed a government UFO project, but was never the head of Blue Book. That position was held, for most or perhaps all of Blue Book's life, by an Air Force Officer Named Edward Ruppelt. Blue Book was shut down in 1969, shortly after the report of the project Condon directed, "Scientific Study of Unidentified Flying Objects". -- Jim Giglio

AUTHOR: Cousins, Frank W.
TITLE: The anatomy of the gyroscope : a report in 3 parts comprising a literature and patent survey directed to the gyroscope and its applications / by Frank W. Cousins ; edited by John L. Hollington.
PUBL.: Neuilly-sur-Seine, France : North Atlantic Treaty Organization, Advisory Group for Aerospace Research and Development,
FORMAT: 296 p. (in various pagings) ; 30 cm.
DATE: 1988
SERIES: AGARDograph no. 313

AUTHOR: Leimanis, E. (Eugene)
TITLE: The general problem of the motion of coupled rigid bodies

- about a fixed point.
 PUBL.: Berlin, New York, Springer-Verlag,
 FORMAT: xvi, 337 p. illus. 24 cm.
 DATE: 1965
 SERIES: Springer tracts in natural philosophy. v. 7
 SUBJECT Dynamics, Rigid Gyroscopes, Two-body problem, Astrodynamics
- AUTHOR(s): de Andrade, L.C. Garcia
 TITLE: Electron gyroscopes to test torsion gravity?
 In: Il nuovo cimento delle societa italiana di fisic OCT 01 1994 v 109 n 10 Page 1123
- AUTHOR(s): Abe, Hiroshi Yoshida, Tetsuo Turuga, Kikuo
 TITLE: Piezoelectric-ceramic cylinder vibratory gyroscope.
 In: Japanese journal of applied physics. part 1, r SEP 01 1992 v 31 n 9B Page 3061
- AUTHOR(s): Case, William B. Shay, Michael A.
 TITLE: On the interesting behavior of a gimbal-mounted gyroscope.
 In: American journal of physics. JUN 01 1992 v 60 n 6 Page 503
- AUTHOR(s): Zhuravlev, V.F.
 TITLE: Nutational self-oscillation of a free gyroscope.
 In: Mechanics of solids. 1992 v 27 n 6 Page 11
- AUTHOR(s): Chang, C.O. Chou, C.S.
 TITLE: Partially Filled Nutation Damper for a Freely Processing Gyroscope.
 In: Journal of guidance, control, and dynamics. SEP 01 1991 v 14 n 5 Page 1046
- AUTHOR(s): Chang, C.O. Chou, C.S. Liu, L.Z.
 TITLE: Stability analysis of a freely precessing gyroscope carrying a mercury ring damper.
 In: Journal of sound and vibration. MAY 08 1991 v 146 n 3 Page 491
- AUTHOR(s): Imanishi, Akira Maruyama, Koichi Midorikawa, Shoichi
 TITLE: Observation against the Weight Reduction of Spinning Gyroscopes.
 In: Journal of the physical society of japan. APR 01 1991 v 60 n 4 Page 1150
- AUTHOR(s): Petry, Walter
 TITLE: Angular Momentum and Gyroscope in Flat Space-Time Theory of Gravitation.
 In: Astrophysics and space science. JAN 01 1991 v 175 n 1 Page 1
- AUTHOR(s): Zhivkov, A.I.
 TITLE: Geometry of invariant manifolds of a gyroscope in the field of a quadratic potential.
 In: Mathematics of the USSR: Izvestija. 1991 v 37 n 1 Page 227
- AUTHOR(s): Hayashi, Kenji Shirafuji, Takeshi
 TITLE: Frame-Dragging Precession of Orbiting Gyroscopes in New General Relativity and Possible Violation of Equivalence Principle.
 In: Progress of theoretical physics. DEC 01 1990 v 84 n 6 Page 1074
- AUTHOR(s): El-Sabaa, F.M.
 TITLE: On the Periodic Motion of a Gyroscope Supported by Cardan Gimbal.
 In: al-Majallah al-Arabiyah lil-ulum wa-al-handasa JUL 01 1990 v 15 n 3 Page 495
- AUTHOR(s): Moffat, J.W. Brownstein, J.R.
 TITLE: Spinning test particles and the motion of a gyroscope according to the nonsymmetric gravitation theory. In: Physical review. D, Particles and fields. MAY 15 1990 v 41 n 10 Page 3111
- AUTHOR(s): Nitschke, J.M. Wilmarth, P.A.
 TITLE: Null result for the weight change of a spinning gyroscope.
 In: Physical review letters. APR 30 1990 v 64 n 18 Page 2115



AUTHOR(s): Faller, J. E. Hollander, W. J. Nelson, P. G.
 TITLE(s): Gyroscope-weighing experiment with a null result.
 In: Physical review letters.
 FEB 19 1990 v 64 n 8 Page 825

AUTHOR(s): Rumyantsev, V.V.
 TITLE: Stability of permanent rotations of a nonsymmetric liquid-filled gyroscope.
 In: Mechanics of solids.
 1990 v 25 n 6 Page 1

AUTHOR(s): Panayotounakos, D.E. Theocaris, P.S.
 TITLE: On the Decoupling and the Solutions of the Euler Dynamic Equations Governing the Motion of a Gyroscope. In: Zeitschrift fur angewandte Mathematik und Mechanik 1990 v 70 n 11 Page 489

AUTHOR(s): Hayasaka, Hideo Takeuchi, Sakae
 TITLE: Gravitation and Astrophysics.
 Summary: Anomalous weight reduction on a gyroscope's right rotations around the vertical axis on the Earth.
 In: Physical review letters.
 DEC 18 1989 v 63 n 25 Page 2701

AUTHOR(s): Laithwaite, Eric
 TITLE: Propulsion by Gyro.
 Summary: In an attempt to reveal the strange, hidden properties of gyroscopes, Professor Eric Laithwaite explains the physics behind the idea that a propulsion system could be built using gyros.
 In: Space. SEP 01 1989 v 5 n 5 Page 36

AUTHOR(s): Vitale, S. Bonaldi, M. Falferi, P.
 TITLE: Magnetization by rotation and gyromagnetic gyroscopes.
 Summary: We discuss how the general phenomenon of magnetization by rotation may be used probe the angular velocity of the laboratory with respect to a local frame of inertia. We show that gyroscope with no moving parts based on this pheno-

In: Physical review B: Condensed matter. JUN 01 1989 v 39 n 16 p B Page 11993

AUTHOR(s): Aspden, H.
 TITLE: Anti Gravity Electronics.
 Summary: Reinterpretation of Newton's Third Law of Motion suggests that it depends upon and electronic action. Electronic interaction therefore explains the paradoxical anti-gravity properties of the force processed gyroscope. In: Electronics & Wireless World.
 JAN 01, 1989 v 95 n 1635 Page 29

AUTHOR(s): Sachs, Mendel
 TITLE: The Precessional Frequency of a Gyroscope in the Quaternionic Formulation of General Relativity. In: Foundations of physics.
 JAN 01 1989 v 19 n 1 Page 105

AUTHOR(s): Medvedev, A.V.
 TITLE: Motion of a rapidly run-up gyroscope acted upon by a constant moment in a resistive medium.
 In: Mechanics of solids.
 1989 v 24 n 2 Page 21

AUTHOR(s): Starzhinskii, V.M.
 TITLE: An exceptional case of motion of the Kovalevskaia gyroscope.
 In: PMM, Journal of applied mathematics and mechanic 1983 v 47 n 1 Page 134

AUTHOR: Gray, Andrew, 1847-1925.
 TITLE: A treatise on gyrostatics and rotational motion; theory and applications.
 PUBL.: New York, Dover Publications
 FORMAT: 530 p. illus. 22 cm.
 DATE: 1959

QA861:P4
 Perry, John
 SPINNING TOPS AND GYROSCOPIC MOTION. By John Perry. Dover, 1957. 102 pages

Articles and Books by Kip Thorne:



AUTHOR: Misner, Charles W.
 TITLE: Gravitation (by) Charles W. Misner,
 Kip S. Thorne (and) John
 Archibald Wheeler.
 PUBL.: San Francisco, W. H. Freeman
 FORMAT: xxvi, 1279 p. illus. 26 cm.
 1973
 SUBJECT: Astrophysics, General relativity,
 Gravitation

GENERAL RELATIVITY & QUANTUM
 COSMOLOGY, ABSTRACT GR-
 QC/9308009 THE QUANTUM
 PROPAGATOR FOR A
 NONRELATIVISTIC PARTICLE IN THE
 VICINITY OF A TIME MACHINE DALIA S.
 GOLDWIRTH, MALCOLM J. PERRY, TSVI
 PIRAN AND KIP S. THORNE.

We study the propagator of a non-relativistic, non-interacting particle in any non-relativistic "time-machine" spacetime of the type shown in Fig. 1: an external, flat spacetime in which two spatial regions, V_- at time t_- and V_+ at time t_+ , are connected by two temporal wormholes, one leading from the past side of V_- to the future side of V_+ and the other from the past side of V_+ to the future side of V_- . We express the propagator explicitly in terms of those for ordinary, flat spacetime and for the two wormholes; and from that expression we show that the propagator satisfies completeness and unitarity in the initial and final "chronal regions" (regions without closed timelike curves) and its propagation from the initial region to the final region is unitary. However, within the time machine it satisfies neither completeness nor unitarity. We also give an alternative proof of initial-region-to-final-region unitarity based on a conserved current and Gauss's theorem. This proof can be carried over without change to most any non-relativistic time-machine spacetime; it is the non-relativistic version of a theorem by Friedman, Papastamatiou and Simon, which says that for a free scalar field, quantum mechanical unitarity follows from the fact that the classical evolution preserves the Klein-Gordon inner product.

AUTHOR(s): Thorne, Kip S.
 TITLE(s): Gravitational-wave bursts with
 memory: The Christodoulou
 effect.

In: Physical review. D, Particles and fields.
 JAN 15 1992 v 45 n 2 Page 520

AUTHOR(s): Apostolatos, Theocharis A.
 Thorne, Kip S.
 TITLE(s): Rotation halts cylindrical,
 relativistic gravitational
 collapse.

In: Physical review. D, Particles and fields.
 SEP 15 1992 v 46 n 6 Page 2435

AUTHOR(s): Echeverria, Fernando
 Klinkhammer, Gunnar Thorne, Kip S.

TITLE(s): Billiard balls in wormhole
 spacetimes with closed timelike
 curves: Classical theory.

In: Physical review. D, Particles and fields.
 AUG 15 1991 v 44 n 4 Page 1077

AUTHOR(s): Eich, Chris Zimmermann,
 Mark E. Thorne, Kip S.

TITLE(s): Giant and supergiant stars
 with degenerate neutron cores.

In: The astrophysical journal.
 NOV 01 1989 v 346 n 1 p 1 Page 277

AUTHOR(s): Frolov, Valery P. Thorne, Kip S.

TITLE(s): Renormalized stress-energy
 tensor near the horizon of a
 slowly evolving, rotating black hole

Summary: The renormalized expectation
 value of the stress-energy
 tensor $\tau_{\mu\nu}$ of a quantum field in an arbitrary
 quantum state near the future horizon of a
 rotating (Kerr) black hole is derived in two
 very different ways: One derivation (restricted
 for simplicity to a massless scalar field) makes
 use of traditional techniques of quantum field
 theory in curved spacetime, augmented by a
 variant of the "ETA formalism" for handling
 superradiant modes. In: Physical review. D,
 Particles and fields. APR 15 1989 v 39 n 8
 Page 2125

AUTHOR(s): Kim, Sung-Won Thorne, Kip S.

TITLE(s): Do vacuum fluctuations prevent the creation of closed timelike curves?
 In: Physical review. D, Particles and fields. JUN 15 1991 v 43 n 12 Page 3929

AUTHOR: Thorne, Kip S.
 TITLE: Black holes and time warps : Einstein's outrageous legacy / Kip S. Thorne.
 PUBL.: New York : W.W. Norton,
 FORMAT: 619 p. : ill. ; 24 cm. 1994
 SERIES: Commonwealth Fund Book Program (Series)
 SUBJECT: Relativity, Astrophysics, Physics--Philosophy, Black holes

AUTHOR: Harrison, B. Kent.
 TITLE: Gravitation theory and gravitational collapse (by) B. Kent
 Harrison, Kip S. Thorne, Masami Wakano (and) John Archibald Wheeler.
 PUBL.: Chicago, University of Chicago Press
 FORMAT: xvii, 177 p. illus. 25 cm. 1965
 SUBJECT: Astrophysics, Gravitation

AUTHOR: Thorne, Kip S.
 TITLE: Gravitational radiation : a new window onto the universe / Kip S. Thorne, William R. Kenan, Jr.
 PUBL.: Cambridge, (Cambridgeshire) ; New York : Cambridge University Press,
 DATE: 1988
 SUBJECT: Gravitational radiation

AUTHOR: Braginskii, V. B. (Vladimir Borisovich
 TITLE: Quantum measurement / Vladimir B. Braginsky and Farid Ya. Khalili ; edited by Kip S. Thorne.
 PUBL.: Cambridge (England) ; New York, NY, USA : Cambridge University Press,
 SUBJECT: Quantum theory, Physical measurements

AUTHOR: Braginskii, V. B. (Vladimir Borisovich)
 Sistemy s maloi dissipatsiei. English

TITLE: Systems with small dissipation / V.B. Braginsky, V.P. Mitrofanov, V.I. Panov ; edited by Kip S. Thorne and Cynthia Eller ; translated by Erast Gliner. PUBL.: Chicago : University of Chicago Press,
 FORMAT: xii, 145 p. : ill. ; 24 cm.
 DATE: 1985
 SUBJECT: Harmonic oscillators--Design and construction. Physical measurements
 NOTES: Translation of: Sistemy s maloi dissipatsiei.
 Includes index.

Title: Black holes : the membrane paradigm / edited by Kip S. Thorne, Richard H. Price, Douglas A. Macdonald. Date/Source: New Haven : Yale University Press, 1986. -----

SELECTED "MEMORABLE" PAPERS FROM THE AMERICAN JOURNAL OF PHYSICS:

E. U. Condon, "Where Do We Live? Reflections on Physical Units and the Universal Constants," 2 (2), 63-69 (1934).


V. F. Weisskopf, "On the Theory of the Electric Resistance of Metals," 11 (1), 1-12 (1943).

E. U. Condon and P. E. Condon, "Effect of Oscillations of the Case on the Rate of a Watch," 16 (1), 14-16 (1948).

G. F. Pake, "Fundamentals of Nuclear Magnetic Resonance Absorption I," 18 (7), 438-452 (1950); "Fundamentals of Nuclear Magnetic Resonance Absorption II," 18 (8), 473-486 (1950).


J. H. Van Vleck, "Landmarks in the Theory of Magnetism," 18 (8), 495-509 (1950).

Herbert Goldstein, "The Classical Motion of a Rigid Charged Body in a Magnetic Field," 19 (2), 100-109 (1951).



- J. C. Slater, "The Electron Theory of Solids," 19 (6), 368-374 (1951).
- L. I. Schiff, "Quantum Effects in the Radiation from Accelerated Relativistic Electrons," 20 (8), 474-478 (1952).
- Kerson Huang, "On the Zitterbewegung of the Dirac Electron," 20 (8), 479-484 (1952).
- F. Keffer, H. Kaplan, and Y. Yafet, "Spin Waves in Ferromagnetic and Antiferromagnetic Materials," 21 (4), 250-257 (1953).
- F.M. Purcell, "Nuclear Magnetism," 22 (1), 1-8 (1954).
- H. G. Dehmelt, "Nuclear Quadrupole Resonance," 22(3), 110-120 (1954).
- C. Kittel, "The Effective Mass of Electrons in Crystals," 22 (5), 250-252 (1954).
- E. J. Zimmerman, "Numerical Coincidences in Microphysics and Cosmology," 23(31), 136-141 (1955).
- W. H. Furry, "Lorentz Transformation and the Thomas Precession," 21 (8), 517-525(1955).
- C. E. Chase, "Ultrasonic Propagation in Liquid Helium," 24 (3), 136-155(1956).
- E. C. Watson, "On the Relations Between Light and Electricity" (a translation of Reinrich Hertz's Heidelberg lecture of 1889), 25 (6), 335-343 (1957).
- E. M. Purcell, "Gravitation Torsion Balance," 25 (6), 393-394 (1957).
- Leonard Eisenbud, "On the Classical Laws of Motion," 26 (3), 144- 159 (1958).
- P. Morrison, "Approximate Nature of Physical Symmetries," 26 (6), 358-368 (1958).
- Bela G. Kolossvary, "Eotvos Balance," 27 (5), 336-343 (1959).
- Leon N. Cooper, "Theory of Superconductivity," 28 (2), 91-101 (1960).
- R. H. Dicke, "Eotvos Experiment and the Gravitational Red Shift," 28 (4), 344-347 (1960).
- Gerald Holton, "On the Origins of the Special Theory of Relativity," 28 (7), 627-636 (1960).
- F. Rohrlich, "Self-Energy and the Stability of the Classical Electron," 28 (7), 639-643 (1960).
- P. W. Bridgman, "Significance of the Mach Principle," 29 (1), 32-36 (1961).
- Robert Weinstock, "Laws of Classical Motion. What's F? What's m? What's a?" 29 (10), 698-702 (1961).
- T. Gold, "The Arrow of Time," 30 (6), 403-410 (1962).
- Robert W. Brehme, "A Geometric Representation of Galilean and Lorentz Transformations," 30 (7), 489-496 (1962).
- David H. Frisch and James H. Smith, "Measurement of Relativistic Time Dilation Using u-Mesons," 31 (5), 342-355 (1963).
- R. H. Dicke, "Cosmology, Mach's Principle and Relativity," 31 (7), 500- 509 (1963).
- R. S. Shankland, "Michelson-Morley Experiment," 32 (1), 16-35 (1964).
- Philip Morrison, "Less May Be More," 32 (6), 441-457 (1964).
- A. L. Schawlow, "Measuring the Wavelength of Light with a Ruler," 33 (11), 922-923 (1965).
- Arthur Komar, "Foundations of Special Relativity and the Shape of the Big Dipper," 33 (12), 1024-1027 (1965).

- R. H. Romer, "Angular Momentum of Static Electromagnetic Fields," 34 (9), 772-778 (1966); "Electromagnetic Angular Momentum," 35 (5), 445-446(1967).
- A. Gamba, "Physical Quantities in Different Reference Systems According to Relativity," 35 (2), 83-89 (1967).
- Emerson M. Pugh and George E. Pugh, "Physical Significance of the Poynting Vector in Static Fields," 35 (2), 153-156 (1967).
- R. H. Dicke, "Gravitation and Cosmic Physics," 35 (7), 559-566 (1967).
- O. L. Brill and B. Goodman, "Causality in the Coulomb Gauge," 35 (9), 832-837 (1967).
- Martin S. Tiersten, "Force, Momentum Change, and Motion," 37 (1), 82-87 (1969).
- W. H. Furry, "Examples of Momentum Distributions in the Electromagnetic Field and in Matter," 37 (6), 621-636 (1969).
- Gerald Holton, "Einstein and the 'Crucial' Experiment," 37 (10), 968-982 (1969)
- Herman Erlichson, "Aharonov-Bohm Effect and Quantum Effects on Charged Particles in Field-Free Regions," 38 (2), 162-173 (1970).
- F. O. Schulz-DuBois, "Foucault Pendulum Experiment by Kamerlingh Onnes and Degenerate Perturbation Theory," 38 (2), 173-188 (1970).
- John Clarke, "The Josephson Effect and e/h ," 38 (9), 1071-1095 (1970).
- Timothy H. Boyer, "Energy and Momentum in Electromagnetic Field for Charged Particles Moving with Constant Velocities," 39 (3), 257-270 (1971).
- Roger Y. Tsien, "Pictures of Dynamic Electric Fields," 40 (1), 46-56 (1972).
- S. Chandrasekhar, "On the 'Derivation' of Einstein's Field Equations," 40 (2), 224-234 (1972).
- Barry R. Holstein and Arthur R. Swift, "The Relativity Twins in Free Fall," 40 (5), 746-750 (1972).
- Henry Pierce Stapp, "The Copenhagen Interpretation," 40 (8), 1098-1116 (1972).
- N. Bloembergen, "The Concept of Temperature in Magnetism," 41 (3), 325-331 (1973).
- Julian Schwinger, "Precession Tests of General Relativity -- Source Theory Derivations," 42 (6), 307-510 (1974).
- Julian Schwinger, "Spin-Precession -- A Dynamical Discussion," 42 (6), 510-513 (1974).
- Allan Franklin, "Principle of inertia in the Middle Ages," 44 (6), 529-545 (1976).
- Hans C. Ohanian, "What is the principle of equivalence?" 45 (10), 903-909 (1977).
- Frederik J. Belinfante, "Can individual elementary particles have individual properties?" 46 (4), 329-336 (1978).
- Timothy H. Boyar, "Electrostatic potential energy leading to an inertial mass change for a system of two point charges," 46 (4), 383-385 (1978).
- E. J. Konopioski, "What the electromagnetic vector potential describes," 46 (5), 499-502 (1978).
- Sidney D. Drell, "When is a particle?" 46 (6), 597-606 (1978).
- Timothy H. Boyer, "Electrostatic potential energy leading to a gravitational mass change for a system of two point charges," 47 (2), 129-131(1979).



Barry R. Holstein and Arthur R. Swift, "Elementary derivation of the radiation field from an accelerated charge," 49 (4), 346-347 (1981).

P. C. Peters, "Where is the energy stored in a gravitational field?" 49(6), 564-569 (1981).

Robert H. Romer, "Motion of a sphere on a tilted turntable," 49 (10), 985-986 (1981).

H. Richard Crane, "Short Foucault pendulum. A way to eliminate the precession due to ellipticity," 49 (11), 1004-1006 (1981).

M. Danos, "Bohm-Aharonov effect. The quantum mechanics of the electrical transformer," 50 (1), 64-66 (1982).

Frank S. Crawford, "Elementary derivation of the magnetic flux quantum," 50 (6), 514-516 (1982).

Robert Weinstock, "Dismantling a centuries-old myth: Newton's Principia and inverse-square orbits," 50 (7), 610-617 (1982).

Frank S. Crawford, "Superball and time-reversal invariance," 50 (9), 856 (1982).

Robert C. Hilborn, "Einstein coefficients, cross sections, f values, dipole moments, and all that," 50 (11), 982-986 (1982).

Allen I. Janis, "Simultaneity and special relativistic kinematics," 51 (3), 209-213 (1983).

Hans C. Ohanian, "On the approach to electro- and magnetostatic equilibrium," 51 (11), 1020-1022 (1983).

N. David Mermin, "Relativity without light," 52 (2), 119-124 (1984).

Timothy H. Boyer, "Lorentz-transformation properties for energy and momentum in electromagnetic systems," 53 (2), 167-171 (1985).

Tyler A. Abbott and David J. Griffiths, "Acceleration without radiation," 53 (12), 1203-1211 (1985).

Victor F. Weisskopf, "Search for Simplicity: Maxwell, Rayleigh, and Mt. Everest," 54 (1), 13-14 (1986).

M. W. P. Strandberg, "Special relativity completed: The source of some 2s in the magnitude of physical phenomena," 54 (4), 321-331 (1986).

Hans C. Ohanian, "What is spin?" 54 (6), 500-505 (1986).

L. Lederman, "Unification, grand unification, and the unity of physics," 54 (7), 594-600 (1986).

E. T. Osypowski and M. G. Olason, "Isynchronous motion in classical mechanics," 55 (8), 720-725 (1987).

W. M. Saslow, "Electromechanical implications of Faraday's law: A problem collection," 55 (11), 986-993 (1987).

Michael S. Morris and Kip S. Thorne, "Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity," 56 (5), 395-412 (1988).


Mark A. Heald, "Energy flow in circuits with Faraday emf," 56 (6), 540-547 (1988).

N. David Mermin, "The amazing many colored relativity engine," 56 (7), 600-611 (1988).

Timothy H. Bayer, "The force on a magnetic dipole," 56 (8), 688-692 (1988).

A. R. Janah, T. Padmanabhan, and T. P. Singh, "On Feynman's formula for the electromagnetic field of an arbitrarily moving charge," 56 (11), 1036-1038 (1988).

Harold S. Zapsolsky, "On electric fields produced by steady currents," 56 (12), 1137-1141 (1988).



M. Kugler, "Motion in noninertial systems: theory and demonstrations," 57 (3), 247-251 (1989).

W. Zimmermann, Jr., "A wave-packet description of the motion of a charged particle in a uniform magnetic field," 57 (7), 593-598 (1989).

Ray Skinner and John A. Weil, "An introduction to generalized functions and their application to static electromagnetic point dipoles, including hyperfine interactions," 57 (9), 777-791 (1989).

S. Washburn, "Conductance fluctuations in loops of gold," 57 (12) 1069-1078 (1989).

Hans Dehmelt, "Less is more. Experiments with an individual atomic particle at rest in free space," 58 (1), 17-27 (1990).

Robert J. Birgenau, "Novel magnetic phenomena and high-temperature superconductivity in lamellar copper oxides," 58 (1), 28-40 (1990).

Freeman J. Dyson, "Feynman's proof of the Maxwell equations," 58 (3), 209-211(1990). See also comments by Norman Dombey, Robert W. Brehme, James L. Anderson, and I. E. Farquhar, 59(l), 85-87 (1991).

Cyrus S. MacLatchy and Hugh A. Chipman, "A dynamic method of measuring the charge induced on a conductor," 58 (9), 811-816 (1990).

G. Matteucci, "Electron wavelike behavior: A historical and experimental introduction," 58(12), 1143-1147 (1990). -----

SELECTED ARTICLES FROM APEIRON MAGAZINE.

Apeiron Magazine (apeiron@aei.ca)
4405 St-Dominique
Montreal, Quebec H2W 2B2 Canada.

Apeiron Number 1 (September 1987)
* Henrik Broberg (Stockholm) – Particle Mass in a Cosmological Perspective

* Toivo Jaakkola (University of Helsinki Observatory) – Mach's Principle and Properties of Local Structure

Apeiron Number 2 (February 1988)
* Jean-Claude Pecker/Jean-Pierre Vigier (University of Paris) – A Possible Tired-Light Mechanism

Apeiron Number 3 (August 1988)
* D.F. Roscoe (University of Sheffield) – Gravitation as an Inertial Process
* Amitabha Ghosh (Indian Institute of Technology, Kanpur) – Velocity-Dependent Inertial Induction: A Case for Experimental Observation

Apeiron Number 5 (Fall 1989)
* S.V.M. Clube (University of Oxford) – Lorentzian Gravity and Cosmology

Apeiron Number 8 (Autumn 1990)
* Jacques Trempe (Montreal, Quebec) – Laws of Light Propagation in Galilean Space-Time
* Thomas E. Phipps Jr. (Urbana, Illinois) – Weber-type Laws of Action-at-a-Distance in Modern Physics

Apeiron Numbers 9-10 (Winter-Spring 1991) * S.V.M. Clube (Astrophysics Department, Oxford University) – Mass Inflation as a Recurring Property of Matter in Astrophysical Situations
* Amitabha Ghosh (Indian Institute of Technology, Kanpur) – Velocity Dependent Inertial Induction: A Possible Tired-Light Mechanism * David Roscoe (Department of Applied Mathematics, Sheffield University) – Gravity out of Inertia
* Henrik Broberg (Djursholm, Sweden) – Mass, Energy, Space * Toivo Jaakkola (University Observatory, Helsinki) – Electrogravitational Coupling: Empirical and Theoretical Arguments

Apeiron Number 12 (February 1992)



* Zaman Akil (Kuwait City) – On the Constant of Gravitation * Andr   K.T. Assis (State University of Campinas, Brazil) – On Hubble's Law of Redshift, Olbers' Paradox and the Cosmic Background Radiation

* S.C. Tiwari (Banaras Hindu University, Varanasi, India) – The Nature of Time

Apeiron Number 13 (June 1992)

* Andr   K.T. Assis (State University of Campinas, Brazil) – On the Absorption of Gravity

* D.F. Roscoe (University of Sheffield) – The Equivalence Principle as a Consequence of the Third Law

* Joop F. Nieland (Arles sur Tech, France) – Vacuum Refraction Theory of Gravitation

* H.E. Wilhelm (University of Utah) – Explanation of Anomalous Unipolar Induction in Corotating Conductor-Magnet Arrangements by Galilean Electrodynamics

Apeiron Number 14 (September 1992)

* Thomas E. Phipps Jr. (Urbana, Illinois) – Lorentz Contraction of the Coulomb Field: An Experimental Proposal * Peter Huber (Germanistisches Seminar, Heidelberg University) – Does the Velocity of Light Decrease?

Apeiron Number 15 (February 1993)

* H.E. Wilhelm (University of Utah) – Galilei Covariant Electrodynamics of Moving Media with Applications to the Experiments of Fizeau and Hoek

* S.X.K. Howusu (University of Jos, Nigeria) – The Confrontation between Relativity and the Principle of Reciprocal Action * Henrik Broberg (Norwegian Telecom, Oslo) – On the Kinetic Origin of Mass

Aperion Number 16 (June 1993)

* C.I. Mocanu (Polytechnical Institute of Bucharest, Romania) – Is Thomas Rotation a Paradox?

* Zu Shaozhi and Xu Xiangqun (Beijing Control Device Research Institute) – On the Relativity of Simultaneity * Martin Kokus (Hopewell, PA) and A.O. Barut, University of Colorado – Suggestion for Unifying Two

Types of Quantized Redshift of Astronomical Bodies

Apeiron Number 17 (October 1993)

* Thomas E. Phipps Jr. (Urbana, Illinois) – Ampere Tension and Newton's Laws

* S.X.K. Howusu (University of Jos, Nigeria) – General Mechanics of a Photon in the Gravitational Field of a Stationary Homogeneous Spherical Body

Apeiron Number 18 (February 1994)

* Toivo Jaakkola (Tuorla Observatory, Turku, Finland) – Radiative and Dynamical Implications of Electrogravity * Peter Huber (Heidelberg University) – The Cosmological Redshift as a Virtual Effect of Gravitation

* H.E. Wilhelm (University of Utah) – Fitzgerald Contraction, Larmor Dilatation, Lorentz Force, Particle Mass and Energy as Invariants of Galilean Electrodynamics

* Adolphe Martin (Longueuil, Quebec) – Einstein to Galilean Relativity

Apeiron Number 19 (June 1994)

* S.H. Kim (University of Texas at Arlington) – A Non-Lorentzian Force Stronger than the Lorentz Force

* Constantin Antonopoulos (National Technological University of Athens) – The Semantics of Absolute Space * P. Graneau (Northeastern University, Arlington) and A.K.T. Assis (University of Campinas, Brazil) – Kirchhoff on the Motion of Electricity in Conductors

* Peter F. Browne (University of Manchester) – Newtonian Cosmology with Renormalized Zero-Point Radiation


Apeiron Number 20 (October 1994)

* V.A. Kuligin, G.A. Kulagina and M.V. Korneva (University of Voronezh, Russia) – Epistemology and Special Relativity * G. Galeczki (University of Kieln) – Physical Laws and the Theory of Special Relativity

* T. Chang (University of Alabama, Huntsville) – A Formulation of the Gravitational Equation of Motion

Aperion Volume 2, Number 1 (January 1995)

* Paul Marmet (University of Ottawa) – Origin



of the 3 K Radiation * Wen-Xiu Li (University of Science and Technology of China) – On the Relativity of Lengths and Times

Apeiron Volume 2, Number 2 (April 1995)

* Amitabha Ghosh (Indian Institute of Technology, Kanpur) – Dynamical Inertial Induction and the Potential Energy Problem * Halton C. Arp (Max-Planck-Institut für Astrophysik) – New Light on Redshift Periodicities

Apeiron Volume 2, Number 3 (July 1995)

* P.F. Browne (University of Manchester) – de Sitter Cosmology Reinterpreted

----- Articles and books by Robert L. Forward:

AUTHOR(s): Cramer, John G. Forward, Robert L. Landis, Geoffrey A. Visser, M. Benford, G. TITLE: Natural wormholes as gravitational lenses.

In: Physical Review-D: Particles, Fields, Gravitation Traversable wormholes, energy condition, time machines MAR 15 1995 v 51 n 6 Page 3117

AUTHOR(s): Forward, R.L. TITLE: Statite: A Spacecraft That Does Not Orbit. In: Journal of spacecraft and rockets. SEP 01 1991 v 28 n 5 Page 606

AUTHOR(s): Forward, Robert TITLE: Advanced Space Propulsion. In: The Journal of social, political, and economic s Wint 1990 v 15 n 4 Page 387

AUTHOR(s): Forward, Robert L. TITLE: Light-Levitated Geostationary Cylindrical Orbits: Correction and Expansion. In: The Journal of the astronautical sciences. JUL 01 1990 v 38 n 3 Page 335

AUTHOR(s): Forward, Robert L. TITLE: Advanced space propulsion. In: Aerospace america. JUL 01 1990 v 28 n 7 Page 60

AUTHOR(s): Forward, Robert L.

TITLE: Grey Solar Sails.
In: The Journal of the astronautical sciences. APR 01 1990 v 38 n 2 Page 161

AUTHOR(s): Forward, Robert

TITLE: The power of negative matter: Does matter with a negative mass exist somewhere in the cosmos? If it does, it would make the perfect space-drive. In: New scientist.

MAR 17 1990 v 125 n 1708 Page 54

AUTHOR(s): Forward, R.L.

TITLE: Solar Photon Thruster.
In: Journal of spacecraft and rockets. JUL 01 1990 v 27 n 4 Page 411

AUTHOR(s): Forward, R.L.

TITLE: Negative matter Propulsion.
In: Journal of propulsion and power. JAN 01 1990 v 6 n 1 Page 28

AUTHOR(s): Forward, R.L.

TITLE: Space Warps: A Review of One Form of Propulsionless Transport.
In: JBIS; Journal of the British Interplanetary Society NOV 01 1989 v 42 n 11 Page 533

AUTHOR: Forward, Robert L.

TITLE: Negative Matter Propulsion
In: July 1988 AIAA Joint Propulsion Conference, Journal of Propulsion and Power Vol 6 no 1 pp 28-37

AUTHOR(S): Robert L. Forward and Joel Davis.

TITLE: Mirror Matter: Pioneering Antimatter Physics
Source: New York : Wiley, 1988.
Series: Wiley science editions

AUTHOR: Forward, Robert L.

TITLE: Spin Drive To The Stars
In: ANALOG, Apr 1981, pp. 64-70

AUTHOR: Forward, R.L.

TITLE: Far Out Physics
In: Analog Science Fiction/Science Fact Vol 95, August 1975 pages 147-166

TITLE: Bibliography of interstellar travel and communication: April 1977
 AUTHOR: Mallove, Eugene F. and Forward, Robert L.

TITLE: Camelot 30K / 1993
 AUTHOR: Forward, Robert L.

TITLE: Dragon's egg ; Starquake / 1994
 AUTHOR: Forward, Robert L.

TITLE: The flight of the dragonfly / 1985
 AUTHOR: Forward, Robert L.

TITLE: Indistinguishable from magic : speculations and visions of the future / 1995
 AUTHOR: Forward, Robert L.

TITLE: Marooned on Eden / 1993
 AUTHOR: Forward, Robert L.

TITLE: Martian rainbow / 1991
 AUTHOR: Forward, Robert L.

TITLE: Mirror matter : pioneering antimatter physics / 1988
 AUTHOR: Forward, Robert L.

TITLE: A national space program for interstellar exploration. 1975
 AUTHOR: Forward, Robert L.

TITLE: Ocean under the ice / 1994
 AUTHOR: Forward, Robert L.

TITLE: Rescued from paradise / 1995
 AUTHOR: Forward, Robert L.

TITLE: Return to Rocheworld / 1993
 AUTHOR: Forward, Robert L.

TITLE: Rocheworld / 1990
 AUTHOR: Forward, Robert L.

TITLE: Starquake / 1986
 AUTHOR: Forward, Robert L.

TITLE: Timemaster / 1992
 AUTHOR: Forward, Robert L.

Dr. Harold Aspden recently retired after serving many years as IBM's patent agent in Europe. He is the discoverer of the "Aspden Effect" or rotational inertia in spinning magnets (NEN, Jan. & Feb. 1995). His Ph.D. thesis involved demonstrations of anomalies in magnetic fields

Books and articles by Harold Aspden:


AUTHOR: Aspden, Harold.
 TITLE: Gravitation / by Harold Aspden.
 PUBL.: Southampton, Eng. : Sabberton Publications,
 FORMAT: 78 p. ; 22 cm.
 DATE: 1975
 SUBJECT Gravitation
 ISBN: 0850560055. 0850560063

AUTHOR: Aspden, Harold.
 TITLE: Modern aether science.
 PUBL.: Southampton, Eng., Sabberton Publications
 FORMAT: 165 p. illus. 22 cm.
 DATE: 1972
 SUBJECT Ether (Space)
 ISBN: 0850560039 0850560047 (pbk)

AUTHOR: Aspden, Harold.
 TITLE: The need for a new theory of gravitation, by H. Aspden.
 PUBL.: Southampton (Hants.), Sabberton Publications,
 FORMAT: (1), 4 p. 22 cm.
 DATE: 1966
 SERIES: His Aether science paper no. 1
 SUBJECT Gravitation
 Relativity (Physics)

AUTHOR: Aspden, Harold.
 TITLE: Physics unified / by Harold Aspden.
 PUBL.: Southampton : Sabberton Publications,
 FORMAT: xi, 206 p. : ill. ; 22 cm.
 DATE: 1980
 SUBJECT Unified field theories
 ISBN: 0850560101

AUTHOR: Aspden, Harold.
 TITLE: Physics without Einstein.
 PUBL.: Southampton, Sabberton Publications,



FORMAT: xiii, 224 p. illus. 23 cm.

DATE: 1969

SUBJECT Physics

ISBN: 850560012

AUTHOR(s): Aspden, H.

TITLE(s): Anti Gravity Electronics.

Summary: Reinterpretation of Newton's Third Law of Motion suggests that it depends upon an electronic action. Electronic interaction therefore explains the paradoxical anti-gravity properties of the force processed gyroscope. In: Electronics & wireless world.

JAN 01 1989 v 95 n 1635 Page: 29

AUTHOR(s): Aspden, H.

TITLE(s): Conservative hadron interactions exemplified by the creation of the kaon.

In: Hadronic journal.

MAY 01 1989 v 12 n 3 Page: 101

AUTHOR(s): Aspden, H.

TITLE(s): The Harwen energy radiation generation.

In: Speculations in science and technology. 1990 v 13 n 4 Page: 295

AUTHOR(s): Aspden, H.

TITLE(s): Instantaneous electrodynamic potential with retarded energy transfer.

In: Hadronic journal.

NOV 01 1988 v 11 n 6 Page: 307

AUTHOR(s): Aspden, H.

TITLE(s): Speculations in energy: editorial introduction.

In: Speculations in science and technology. 1990 v 13 n 4 Page: 243

AUTHOR(s): Aspden, H.

TITLE(s): The theory of the proton constants.

In: Hadronic journal.

JUL 01 1988 v 11 n 4 Page: 169

Selected Publications by LEONARD PARKER

* On the Magnetic Moment of a Charged Particle in a Changing Magnetic Field, Nuovo Cimento 40B 99 (1965). * Equivalence Principle and Motion of a Gyroscope, Physical Review 175, 1658 (1968).

* Motion in a Schwarzschild Field I. Precession of a Moving Gyroscope, American Journal of Physics 37, 309 (1969). * Faster-Than-Light Inertial Review 188, 2287 (1969).

* Special Relativity and Diagonal Transformations (with G. Schmieg), American Journal of Physics 38, 218 (1970). * Quantized Matter Fields and the Avoidance of Singularities in General Relativity (with S. A. Fulling), Physical Review D7, 2357 (1973).

* Metric of Two Spinning Charged Sources in Equilibrium (with R. Ruffini and D. Wilkins), Physical Review D7, 2874 (1973). * Quantized Scalar Fields in a Closed Anisotropic Universe (with B. L. Hu and S. A. Fulling), Physical Review D8, 2377 (1973). * Adiabatic Regularization of the Energy-Momentum Tensor of a Quantized Field in Homogeneous Spaces (with S. A. Fulling), Physical Review D9, 341 (1974).

* Renormalization in the Theory of a Quantized Scalar Field Interacting With a Robertson-Walker Spacetime (with S. A. Fulling), Annals of Physics (N.Y.) 87, 176-203 (1974). * Adiabatic Analysis and Renormalization in Semiclassical Gravitation Theory (with S. A. Fulling), Bulletin American Physical Society 19 108 (1974).

* Conformal Energy-Momentum Tensor in Curved Space-Time: Adiabatic Regularization and Renormalization (with S. A. Fulling and B. L. Hu), Physical Review D10, 3905 (1974).

* Solution of the Einstein-Maxwell Equations for Two Unequal Spinning Sources in Equilibrium (with R. A. Kobiske), Physical Review D10, 2321 (1974).

* The Interaction of Gravity with Quantized Fields (with S. A. Fulling), award winning essay, Gravity Research Foundation Awards, 1974.

* New Developments in the Theory of Gravity Interacting With a Quantized Field, Journal of General Relativity and Gravitation 6, 21 (1975).



- * Nonlinear Gravitational Effects and Magnetic Monopoles, Physical Review Letters 34, 412 (1975).
- * Quantized Fields and Particle Creation in Curved Space-Time (66 pages), in Proceedings of the Second Latin American Symposium on Relativity and Gravitation (Universidad Simon Bolivar, Caracas, 1976).
- * The Production of Elementary Particles by Strong Gravitational Fields (120 pages), in Proceedings of the Symposium on Asymptotic Properties of Space-Time (Plenum Publishing Corp., New York, 1977), editors, P. Esposito and L. Witten.
- * Angular Momentum and Dirac Charge Quantization in Curved Spacetime, (with J. Friedman and S. Mayer), Physical Review D17, 1957 (1978).
- * Applied Quantum Gravity: Applications of the Semiclassical Theory, in On the Path of Albert Einstein, edited by B. Kursunoglu, A. Perlmutter, and L. F. Scott (Plenum Press, New York, 1979), pp. 145-166.
- * Quantized Matter Fields ... (with S. A. Fulling), reprinted in Cosmology, Selected Reprints, edited by L. C. Shepley and A. A. Strassenburg (American Association of Physics Teachers, Stony Brook, N.Y., 1979).
- * On Renormalization of ϕ^4 Field Theory in Curved Spacetime I, (with T. S. Bunch and P. Panangaden), Journal of Physics A13, 901 (1980)
- * One-Electron Atom in Curved Spacetime, Physical Review Letters 44, 1559 (1980).
- * The One-Electron Atom as a Probe of Spacetime Curvature, Physical Review D22, 1922 (1980).
- * Self-Forces and Atoms in Gravitational Fields, Physical Review D24, 535 (1981).
- * The Atom as a Probe of Curved Spacetime, Journal of General Relativity and Gravitation 13, 307 (1981).
- * Gravitational Perturbation of the Hydrogen Spectrum (with L. O. Pimentel), Physical Review D25, 3180 (1982).
- * Quantum Gravity 2 (book review), Science 217, 346 (1982).
- * Remote Quantum Mechanical Detection of Gravitational Radiation, (with T. K. Leen and L. O. Pimentel), General Relativity and Gravitation 15, 761-776 (1983).
- * Renormalization and Scaling of Non-Abelian Gauge Fields in Curved Space-Time, in Gauge Theory and Gravitation, edited by K. Kikkawa, N. Nakanishi, and H. Nariai (Springer-Verlag, Berlin, 1983), pp. 96-100.
- * Gravitational Perturbations of the Hydrogen Atom, in Proceedings of the Third Marcel Grossmann Meeting on General Relativity, edited by Hu Ning (Science Press and North-Holland Publishing Co., New York, 1983).
- * Effective Couplings at High Curvature, in Proceedings of the Workshop on Induced Gravitation, Erice, Italy (Sept. 1983).
- * Renormalization Group Analysis of Grand Unified Theories in Curved Spacetime (with D. J. Toms), Physical Review D29, 1584-1608 (1984).
- * Effective Couplings of Grand Unified Theories in Curved Spacetime, (with D. J. Toms), Physical Review Letters 52, 1269 (1984).
- * Some Cosmological Aspects of Quantum Gravity, in Quantum Theory of Gravity, edited by S. M. Christensen (Adam Hilger Ltd., Bristol, 1984), pp. 89-102 (invited paper for volume in honor of 60th birthday of B. S. DeWitt).
- * Curvature Dependence of Renormalized Coupling Constants, Foundations of Physics 14, 1121 (1984) (invited paper for issue in honor of 75th birthday of Nathan Rosen).
- * Effective Coupling Constants and GUT's in the Early Universe, in Inner Space/Outer Space, The Interface Between Cosmology and Particle Physics, edited by E. W. Kolb et al. (University of Chicago Press, Chicago, 1985).
- * Models of Rapidly Rotating Neutron Stars (with J. L. Friedman and R. Ipser), Nature 312, 255 (1984).
- * Gravity and Grand Unified Theories (with D. J. Toms), General Relativity and Gravitation 17, 167 (1985).
- * New Form for the Coincidence Limit of Feynman Propagator, or Heat Kernel, in Curved Spacetime (with D. J. Toms), Physical Review D31, 953 (Rapid Communications) (1985).
- * Explicit Curvature Dependence of Coupling Constants (with D. J. Toms), Physical Review D31, 2424 (1985).
- * Proof of Summed Form of Proper-Time Expansion for Propagator in Curved Space-Time (with I. Jack), Physical Review D31, 2439 (1985).
- * Curvature-Induced Asymptotic Freedom (with E. Calzetta and I. Jack), Physical Review



Letters 55, 1241 (1985). * Renormalization Group and Nonlocal Terms in the Curved-Spacetime Effective Action (with D. J. Toms), Physical Review D32, 1409 (1985).

* Rapidly Rotating Neutron Star Models (with J. L. Friedman and J. R. Ipser), Astrophysical Journal 304, 115-139 (1986). * Quantum Gauge Fields at High Curvature (with E. Calzetta and I. Jack), Physical Review D33, 953-977 (1986). * Renormalization Group Methods in Curved Spacetime (with D. J. Toms) in Quantum Field Theory and Quantum Statistics, edited by I. A. Batalin, C. J. Isham, and G. A. Vilkowisky (invited paper for volume in honor of 60th birthday of Soviet Physicist E. J. Fradkin) (Adam Hilger Ltd., Bristol, 1988). * Gravitational Particle Production in the Formation of Cosmic Strings, Phys. Rev. Letters 59, 1369 (1987). * Gravitational Singularities and Two-Body Interactions, in Nonlinear Phenomena in Relativity and Cosmology, editors, J. R. Buchler et al. (New York Academy of Sciences, Vol. 631, New York, 1991), 31-39.

* Ultrarelativistic Bose-Einstein Condensation in the Einstein Universe and Energy Conditions (with Yang Zhang), Phys. Rev. D44, 2421-2431 (1991).

* Relativistic Condensate as a Source for Inflation (with Yang Zhang), Phys. Rev. D47, 416 (1993).

* Einstein Equations with Quantum Corrections Reduced to Second Order, Phys. Rev. D47, 1339 (1993) (with J. Simon). * Physical Solutions of Semiclassical General Relativistic Models with Quantum Corrections, in Topics on Quantum Gravity and Beyond, Essays in Honor of Louis Witten on His Retirement, editors, F. Mansouri and J. J. Scanio, (World Scientific, Singapore, 1994).

* Atomic Spectra in the Gravitational Field of a Collapsing Prolate Spheroid (with D. Vollick and I. Redmount), Phys. Rev. D (submitted 1994).

* Physical Distinction Among Alternative Vacuum States in Flat Spacetime (with S. Winters-Hilt and I. Redmount), Phys. Rev. D (submitted 1994).

Publications by Matt Visser
(visser@kiwi.wustl.edu)
<http://www.physics.wustl.edu/~visser/homepage.html>

Charge nonconserving decays in ordinary matter. Physical Review D24 (1981) 2542--2544.

Concerning the mass of the photon.
Physics Letters B109 (1982) 373--374.

Aspects of supersymmetry breaking.
Lawrence Berkeley Laboratory Report, LBL-18189, September 1984. [Ph.D. Thesis]

The topological degree for supersymmetric chiral models. Physical Review D32 (1985) 510--512.

Some generalizations of the O'Raifeartaigh model. Journal of Physics A18 (1985) L979--L982.

An exotic class of Kaluza-Klein models.
Physics Letters B158 (1985) 22--25.

A supergravity model without elementary gauge singlets. Physics Letters B160 (1985) 77--80.

Number of massless fermion families in superstring theory. With Itzhak Bars Physics Letters B163 (1985) 118--122.

Tuning the cosmological constant in N=1 supergravity. Physics Letters B165 (1985) 289--291.

Tree level mass spectra in the observable sector. Nuclear Physics B271 (1986) 53--60.

Fermion families in superstring theory. With Itzhak Bars Proceedings of the Oregon meeting, 1985 Annual meeting of the Division of Particles and Fields of the American Physical Society, edited by: R. Hwa, (World Scientific, Singapore, 1986), pp. 829--834.



Feeble intermediate range forces from higher dimensions. With Itzhak Bars Physical Review Letters 57 (1986) 25--28.

Feeble forces and gravity. With Itzhak Bars General Relativity and Gravitation 19 (1987) 219--223. [This essay was awarded Second Prize in the 1986 Essay Competition sponsored by the Gravity Research Foundation.]

Feeble forces. With Itzhak Bars in: Proceedings of the 23rd International Conference in High Energy Physics, Berkeley, July 1986, edited by: S. C. Loken, (World Scientific, Singapore, 1987), volume 2, pp. 1032--1037.

A guide to data in elementary particle physics. (as part of the Berkeley Particle Data Group collaboration) Lawrence Berkeley Laboratory Report, LBL--90 (Revised), UC--34D, September 1986.

Is the 'missing mass' really missing? General Relativity and Gravitation 20 (1988) 77--81. [This essay was awarded an honorable mention in the 1987 Essay Competition sponsored by the Gravity Research Foundation.]

Determinants of conformal wave operators in four dimensions. With Steve Blau and Andreas Wipf Physics Letters B209 (1988) 209--213.

Zeta functions and the Casimir energy. With Steve Blau and Andreas Wipf) Nuclear Physics B310 (1988) 163--180.

Determinants, Dirac operators, and one-loop physics. With Steve Blau and Andreas Wipf International Journal of Modern Physics A4 (1989) 1467--1484.

A classical model for the electron. Physics Letters A139 (1989) 99--102.

Traversable wormholes: Some simple examples. Physical Review D39 (1989) 3182--3184.

Traversable wormholes from surgically modified Schwarzschild spacetimes. Nuclear Physics B328 (1989) 203--212.

Wormholes, baby universes, and causality. Physical Review D41 (1990) 1116--1124.

Quantum mechanical stabilization of Minkowski signature wormholes. Physics Letters B242 (1990) 24--28. [Based on a talk given at the 'Wormshop', Fermilab, May 1989.]

Quantum wormholes in Lorentzian signature. in: Proceedings of the Rice meeting, 1990 meeting of the Division of Particles and Fields of the American Physical Society, edited by: B. Bonner and H. Miettinen, (World Scientific, Singapore, 1990), volume 2, pp. 858--860.

Canonically quantized gravity: Disentangling the supermomentum and superhamiltonian constraints. Physical Review D42 (1990) 1964--1972.

Quantum wormholes. Physical Review D43 (1991) 402--409.

Wheeler-DeWitt quantum gravity in (2+1) dimensions. in: Beyond the Standard Model II: Proceedings of the International Conference in High Energy Physics, Norman, Oklahoma, 1--3 November 1990, edited by: K. A. Milton, R. Kantowski, and M. A. Samuel, (World Scientific, Singapore, 1991), pp. 354--359.

Wheeler wormholes and topology change: A minisuperspace analysis. Modern Physics Letters A6 (1991) 2663--2667. [Based on an essay that was awarded an honorable mention in the 1990 Essay Competition sponsored by the Gravity Research Foundation.]

Analytic results for the effective action. With Steve Blau and Andreas Wipf International Journal of Modern Physics A6 (1991) 5409--5433.



Dirty black holes: Thermodynamics and horizon structure. *Physical Review D* 46 (1992) 2445--2451.

>From wormhole to time machine:
Remarks on Hawking's chronology protection conjecture. *Physical Review D* 47 (1993) 554--565.

van Vleck determinants:
Geodesic focussing and defocussing in Lorentzian spacetimes. *Physical Review D* 47 (1993) 2395--2402.

Hawking radiation: A particle physics perspective. *Modern Physics Letters A* 8 (1993) 1661--1670.

Dirty black holes: Entropy versus area.
Physical Review D 48 (1993) 583--591.

Dirty black holes: Entropy as a surface term.
Physical Review D 48 (1993) 5697--5705.

Lateral wave contributions to the low--altitude radar propagation factor. With Gerald Gilbert and Eric Raiten
Radio Science 29 (1994) 483--494.

Hawking's chronology protection conjecture:
Singularity structure of the quantum stress-energy tensor. *Nuclear Physics B* 416 (1994) 895--906.

van Vleck determinants: Traversable wormhole spacetimes. *Physical Review D* 49 (1994) 3963--3980.

Natural wormholes as gravitational lenses.
With John Cramer, Robert Forward, Michael Morris, Gregory Benford, and Geoffrey Landis
Physical Review D 51 (1995) 3117--3120.

Scale anomalies imply violation of the averaged null energy condition. *Physics Letters B* 349 (1995) 443--447.


Lorentzian Wormholes --- from Einstein to Hawking. Full length technical monograph.
American Institute of Physics Press, July 1995.

Acoustic propagation in fluids:
An unexpected example of Lorentzian geometry. gr-qc/9311028; gr-qc@xxx.lanl.gov.
Submitted for publication.

Thin-shell wormholes: Linearization stability
With Eric Poisson)
gr-qc/9506083; gr-qc@xxx.lanl.gov
Submitted for publication.

New book by Matt Visser. Published 1995
"Lorentzian Wormholes---from Einstein to Hawking", by Matt Visser (Washington University in St. Louis). American Institute of Physics Press (Woodbury, New York). ISBN 1-56396-394-9
412 pages (including index and 38 figures); hardback; US\$59.00 (US\$47.20 for APS/AIP members). To order---Voice: 1-800-809-2247; FAX: 1-802-864-7626. Table of contents:

Chapter 0 Preface	vii
Part i --- Background	1
Chapter 1 Introduction	3
Chapter 2 General relativity	9
Chapter 3 Quantum field theory	31
Chapter 4 Units and natural scales	39
Part ii -- History	43
Chapter 5 The Einstein--Rosen bridge	45
Chapter 6 Spacetime foam	53
Chapter 7 The Kerr wormhole	75
Chapter 8 The cosmological constant	81
Chapter 9 Wormhole taxonomy	89
Chapter 10 Interregnum	95
Part iii - Renaissance	97
Chapter 11 Traversable wormholes	99
Chapter 12 Energy conditions	115
Chapter 13 Engineering considerations	137
Chapter 14 Thin shells: Formalism	153
Chapter 15 Thin shells: Wormholes	165
Chapter 16 Topological censorship	195
Part iv -- Time Travel	201
Chapter 17 Chronology: Basic notions	203
Chapter 18 From wormhole to time machine	227
Chapter 19 Response to the paradoxes	249
Part v --- Quantum Effects	277
Chapter 20 Semiclassical quantum gravity	279



Chapter 21 van Vleck determinants: Formalism 295 Chapter 23 Singularity structure 333 Chapter 24 Minisuperspace wormholes 347 Part vi -- Reprise 367 Chapter 25 Where we stand 369 ----- Other articles about wormholes:	TITLE(s): Quantum electromagnetic wormholes and geometrical description of the electric charge. In: Physical review d: particles, fields, gravitat OCT 15 1994 v 50 n 8 Page 5087
AUTHOR(s): Kar, Sayan Sahdev, Deshdeep TITLE(s): Restricted class of traversable wormholes with traceless matter. In: Physical review d: particles, fields, gravitat AUG 15 1995 v 52 n 8 Page 2030	AUTHOR(s): Singh, Harvendra TITLE(s): Magnetic wormholes and vertex operators. In: Physical review d: particles, fields, gravitat OCT 15 1994 v 50 n 8 Page 5033
AUTHOR(s): Wang, Anzhong Letelier, Patricio S. TITLE(s): Dynamical Wormholes and energy Conditions. In: Progress of theoretical physics. JUL 01 1995 v 94 n 1 Page 137	AUTHOR(s): Li, Xin-zhou TITLE(s): Dimensionally continued wormhole solutions. In: Physical review d: particles, fields, gravitat SEP 15 1994 v 50 n 6 Page 3787
AUTHOR(s): Clement, Gerard TITLE(s): Wormhole cosmic strings. In: Physical review d: particles, fields, gravitat JUN 15 1995 v 51 n 12 Page 6803	AUTHOR(s): Marugan, G.A. Mena TITLE(s): Bases of wormholes in quantum cosmology. In: Classical and quantum gravity. SEP 01 1994 v 11 n 9 Page: 2205
AUTHOR(s): Cramer, John G. Forward, Robert L. Landis, Geoffrey A. TITLE(s): Natural wormholes as gravitational lenses. In: Physical review d: particles, fields, gravitat MAR 15 1995 v 51 n 6 Page 3117	AUTHOR(s): Ridgway, S. Alexander TITLE(s): S1xS2 wormholes and topological charge. In: Physical review d: particles, fields, gravitat JUL 15 1994 v 50 n 2 Page 892
AUTHOR(s): Agnese, A.G. La Camera, M. TITLE(s): Wormholes in the Brans-Dicke theory of gravitation. In: Physical review d: particles, fields, gravitat FEB 15 1995 v 51 n 4 Page 2011	AUTHOR(s): Cavaglia, Marco de Alfaro, Vittorio de Felice, Fernando TITLE(s): Anisotropic wormhole: Tunneling in time and space. In: Physical review d: particles, fields, gravitat JUN 15 1994 v 49 n 12 Page 6493
AUTHOR(s): Cadoni, Mariano Cavaglia, Marco TITLE(s): Cosmological and wormhole solutions in low-energy effective string theory. In: Physical review d: particles, fields, gravitat NOV 15 1994 v 50 n 10 Page 6435	AUTHOR(s): Cotsakis, Spiros Leach, Peter Flessas, George TITLE(s): Minimally coupled scalar field wormholes. In: Physical review d: particles, fields, gravitat JUN 15 1994 v 49 n 12 Page 6489
AUTHOR(s): Cavaglia, Marco	AUTHOR(s): Visser, Matt TITLE(s): van Vleck determinants: Traversable wormhole spacetimes. In: Physical review d: particles, fields, gravitat APR 15 1994 v 49 n 8 Page 3963



AUTHOR(s): Hochberg, David Kephart, Thomas W.

TITLE(s): Can Semi-Classical Wormholes Solve the Cosmological Horizon Problem?

In: General relativity and gravitation.

FEB 01 1994 v 26 n 2 Page 219

AUTHOR(s): Kar, Sayan Sahdev, Deshdeep Bhawal, Biplab

TITLE(s): Scalar waves in a wormhole geometry.

In: Physical review d: particles and fields. JAN 15 1994 v 49 n 2 Page 853

AUTHOR(s): Khatsymovsky, V.

TITLE(s): Can wormholes exist?

In: Physics letters. [part b].

JAN 13 1994 v 320 n 3 / 4 Page 234

AUTHOR(s): Kim, S.W. Lee, H. Kim, S.K.

TITLE(s): (2+1)-dimensional Schwarzschild-de Sitter wormhole.

In: Physics letters: [part A]

DEC 20 1993 v 183 n 5 / 6 Page 359

AUTHOR(s): Liu, Liao

TITLE(s): Wormhole created from vacuum fluctuation.

In: Physical review d: particles and fields.

DEC 15 1993 v 48 n 12 Page R5463

AUTHOR(s): Gonzalez-Diaz, P.F.

TITLE(s): The conjugate gauge wormhole vertex.

Electrogravitics Systems: An examination of electrostatic motion, dynamic counterbary and barycentric control.¹

Prepared by: Gravity Research Group Aviation Studies (International) Limited
Special Weapons Study Unit
29-31 Cheval Place, Knightsbridge
London, S.W.7. England
Report GRG- 013/56 February 1956.
AF Wright Aeronautical Laboratories Wright-Patterson Air Force Base Technical Library
Dayton, Ohio 45433 TL 565 A9
Bar Code: 3 1401 00034 5879

Contents

Introductory Notes.....	1
Discussion.....	3
Conclusions.....	18
Appendix I - Aviation Report extracts.....	21
Appendix II - Electrostatic Patents.....	33

Electrogravitics Systems

An examination of electrostatic motion, dynamic counterbary and barycentric control. It has been accepted as axiomatic that the way to offset the effects of gravity is to use a lifting surface and considerable molecular energy to produce a continuously applied force that, for a limited period of time, can remain greater than the effects of gravitational attraction. The original invention of the glider and evolution of the briefly self-sustaining glider, at the turn of the century led to progressive advances in power and knowledge. This has been directed to

¹ The current article was not prepared by the Air Force but by an organization called Gravity Research Group, Special Weapons Study Unit, a subdivision of Aviation Studies (International) Ltd. from the UK Aviation Studies was a research group ran by two young aerospace analysts Richard Worcester and John Longhurst, who in mid-1950 also published an aerospace and defense industry newsletter. [Venik]

refining the classic Wright Brothers' approach. Aircraft design is still fundamentally as the Wrights adumbrated it, with wings, body, tails, moving or flapping controls, landing gear and so forth. The Wright biplane was a powered glider, and all subsequent aircraft, including the supersonic jets of the nineteen-fifties are also powered gliders. Only one fundamentally different flying principle has so far been adopted with varying degrees of success. It is the rotating wing aircraft that has led to the jet lifters and vertical pushers, copters, ducted fans and lift induction turbine propulsion systems. But during these decades there was always the possibility of making efforts to discover the nature of gravity from cosmic or quantum theory, investigation and observation, with a view to discerning the physical properties of aviation's enemy. It has seemed to Aviation Studies that for some time insufficient attention has been directed to this kind of research. If it were successful such developments would change the concept of sustentation, and confer upon a vehicle qualities that would now be regarded as the ultimate in aviation. This report summarizes in simple form the work that has been done and is being done in the new field of electrogravitics. It also outlines the various possible lines of research into the nature and constituent matter of gravity, and how it has changed from Newton to Einstein to the modern Hlavaty concept of gravity as an electromagnetic force that may be controlled like a light wave. The report also contains an outline of opinions on the feasibility of different electrogravitics systems and there is reference to some of the barycentric control and electrostatic rigs in operation. Also included is a list of references to electrogravitics in successive Aviation Reports since a drive was started

by Aviation Studies (International) Limited to suggest to aviation business eighteen months ago that the rewards of success are too far-reaching to be overlooked, especially in view of the hopeful judgement of the most authoritative voices in microphysics. Also listed are some relevant patents on electrostatics and electrostatic generators in the United States, United Kingdom and France. [Signed] Gravity Research Group 25 February 1956

Discussion

Electrogravitics might be described as a synthesis of electrostatic energy used for propulsion - either vertical propulsion or horizontal or both - and gravitics, or dynamic counterbary, in which energy is also used to set up a local gravitational force independent of the earth's. Electrostatic energy for propulsion has been predicted as a possible means of propulsion in space when the thrust from a neutron motor or ion motor would be sufficient in a dragless environment to produce astronomical velocities. But the ion motor is not strictly a part of the science of electrogravitics, since barycentric control in an electrogravitics system is envisaged for a vehicle operating within the earth's environment and it is not seen initially for space application. Probably large scale space operations would have to await the full development of electrogravitics to enable large pieces of equipment to be moved out of the region of the earth's strongest gravity effects. So, though electrostatic motors were thought of in 1925, electrogravitics had its birth after the War, when Townsend Brown sought to improve on the various proposals that then existed for electrostatic motors sufficiently to produce some visible manifestation of sustained motion.

Whereas earlier electrostatic tests were essentially pure research, Brown's rigs were aimed from the outset at producing a flying article. As a private venture he produced evidence of motion using condensers in a couple of saucers suspended by arms rotating round a central tower with input running down the arms. The massive-k situation was summarized subsequently in a report, Project Winterhaven, in 1952. Using the data some conclusions were arrived at that might be expected from ten or more years of intensive development - similar to that, for instance, applied to the turbine engine. Using a number of assumptions as to the nature of gravity, the report postulated a saucer as the basis of a possible interceptor with Mach 3 capability. Creation of a local gravitational system would confer upon the fighter the sharp-edged changes of direction typical of motion in space. The essence of electrogravitics thrust is the use of a very strong positive charge on one side of the vehicle and a negative on the other. The core of the motor is a condenser and the ability of the condenser to hold its charge (the k-number) is the yardstick of performance. With air as 1, current dielectrical materials can yield 6 and use of barium aluminate can raise this considerably, barium titanium oxide (a baked ceramic) can offer 6,000 and there is promise of 30,000, which would be sufficient for supersonic speed. The original Brown rig produced 30 fps on a voltage of around 50,000 and a small amount of current in the milliamp range. There was no detailed explanation of gravity in Project Winterhaven, but it was assumed that particle dualism in the subatomic structure of gravity would coincide in its effect with the issuing stream of electrons from the electrostatic energy source to produce counterbary. The Brown work probably remains a realistic

approach to the practical realization of electrostatic propulsion and sustentation. Whatever may be discovered by the Gravity Research Foundation of New Boston a complete understanding and synthetic reproduction of gravity is not essential for limited success. The electrogravitics saucer can perform the function of a classic lifting surface - it produces a pushing effect on the under surface and a suction effect on the upper, but, unlike the airfoil, it does not require a flow of air to produce the effect. First attempts at electrogravitics are unlikely to produce counterbary, but may lead to development of an electrostatic VTOL vehicle. Even in its developed form this might be an advance on the molecular heat engine in its capabilities. But hopes in the new science depend on an understanding of the close identity of electrostatic motivating forces with the source and matter of gravity. It is fortuitous that lift can be produced in the traditional fashion and if an understanding of gravity remains beyond full practical control, electrostatic lift might be an adjunct of some significance to modern thrust producers. Research into electrostatics could prove beneficial to turbine development, and heat engines in general, in view of the usable electron potential round the periphery of any flame. Materials for electrogravitics and especially the development of commercial quantities of high-k material is another dividend to be obtained from electrostatic research even if it produces no counterbary. This is a line of development that Aviation Studies, Gravity Research Group is following. One of the interesting aspects of electrogravitics is that a breakthrough in almost any part of the broad front of general research on the intranuclear processes may be translated into a meaningful advance towards the feasibility of electrogravitics systems. This

demands constant monitoring in the most likely areas of the physics of high energy sub-nuclear particles. It is difficult to be overoptimistic about the prospects of gaining so complete a grasp of gravity while the world's physicists are still engaged in a study of fundamental particles - that is to say those that cannot be broken down any more. Fundamental particles are still being discovered - the most recent was the Segre-Chamberlain-Wiegand attachment to the bevatron, which was used to isolate the missing anti-proton, which must - or should be presumed to - exist according to Dirac's theory of the electron. Much of the accepted mathematics of particles would be wrong if the anti-proton was proved to be non-existent. Earlier Eddington has listed the fundamental particles as:- e. The charge of an electron m. The mass of an electron. M. The mass of a proton. h. Planck's constant c. The velocity of light. G. The constant of gravitation, and [Greek letter, small lambda]. The cosmical constant. It is generally held that no one of these can be inferred from the others. But electrons may well disappear from among the fundamental particles, though, as Russell says, it is likely that e and m will survive. The constants are much more established than the interpretation of them and are among the most solid of achievements in modern physics.

Gravity may be defined as a small scale departure from Euclidean space in the general theory of relativity. The gravitational constant is one of four dimensionless constants: first, the mass relation of the nucleon and electron. Second is $(e^*e)/(h^*c)$ [equation form], third, the Compton wavelength of the proton, and fourth is the gravitational constant, which is the ratio of the electrostatic to the gravitational attraction

between the electron and the proton. One of the stumbling blocks in electrogravitics is the absence of any satisfactory theory linking these four dimensionless quantities. Of the four, moreover, gravity is decidedly the most complex, since any explanation would have to satisfy both cosmic and quantum relations more acceptably and intelligibly even than in the unified field theory. A gravitational constant of around 10.E-39 [equation form] has emerged from quantum research and this has been used as a tool for finding theories that could link the two relations. This work is now in full progress, and developments have to be watched for the aviation angle. Hitherto Dirac, Eddington, Jordan and others have produced differences in theory that are too wide to be accepted as consistent. It means therefore that (i) without a cosmical basis, and (ii) with an imprecise quantum basis and (iii) a vague hypothesis on the interaction, much remains still to be discovered. Indeed some say that a single interacting theory to link up the dimensionless constants is one of three major unresolved basic problems of physics. The other two main problems are the extension of quantum theory and a more detailed knowledge of the fundamental particles. All this is some distance from Newton, who saw gravity as a force acting on a body from a distance, leading to the tendency of bodies to accelerate towards each other. He allied this assumption with Euclidean geometry, and time was assumed as uniform and acted independently of space. Bodies and particles in space normally moved uniformly in straight lines according to Newton, and to account for the way they sometimes do not do so, he used the idea of a force of gravity acting at a distance, in which particles of matter cause in others an acceleration proportional to their mass, and inversely proportional to the square of the

distance between them. But Einstein showed how the principle of least action, or the so-called cosmic laziness means that particles, on the contrary, follow the easiest path along geodesic lines and as a result they get readily absorbed into space-time. So was born non-linear physics. The classic example of non- linear physics is the experiment in bombarding a screen with two slits. When both slits are open particles going through are not the sum of the two individually but follows a non-linear equation. This leads on to wave-particle dualism and that in turn to the Heisenberg uncertainty principle in which an increase in accuracy in measurement of one physical quantity means decreasing accuracy in measuring the other. If time is measured accurately energy calculations will be in error; the more accurate the position of a particle is established the less certain the velocity will be; and so on. This basic principle of the acausality of microphysics affects the study of gravity in the special and general theories of relativity. Lack of pictorial image in the quantum physics of this interrelationship is a difficulty at the outset for those whose minds remain obstinately Euclidean. In the special theory of relativity, space-time is seen only as an undefined interval which can be defined in any way that is convenient and the Newtonian idea of persistent particles in motion to explain gravity cannot be accepted. It must be seen rather as a synthesis of forces in a four dimensional continuum, three to establish the position and one the time. The general theory of relativity that followed a decade later was a geometrical explanation of gravitation in which bodies take the geodesic path through space-time. In turn this means that instead of the idea of force acting at a distance it is assumed that space, time, radiation and particles are linked and variations in them from gravity



are due rather to the nature of space. Thus gravity of a body such as the earth instead of pulling objects towards it as Newton postulated, is adjusting the characteristics of space and, it may be inferred, the quantum mechanics of space in the vicinity of the gravitational force. Electrogravitics aims at correcting this adjustment to put matter, so to speak, 'at rest'.

One of the difficulties in 1954 and 1955 was to get aviation to take electrogravitics seriously. The name alone was enough to put people off. However, in the trade much progress has been made and now most major companies in the United States are interested in counterbary. Groups are being organised to study electrostatic and electromagnetic phenomena. Most of industry's leaders have made some reference to it. Douglas has now stated that it has counterbary on its work agenda but does not expect results yet awhile. Hiller has referred to new forms of flying platform, Glenn Martin say gravity control could be achieved in six years, but they add that it would entail a Manhattan District type of effort to bring it about. Sikorsky, one of the pioneers, more or less agrees with the Douglas verdict and says that gravity is tangible and formidable, but there must be a physical carrier for this immense trans-spatial force. This implies that where a physical manifestation exists, a physical device can be developed for creating a similar force moving in the opposite direction to cancel it. Clarke Electronics state they have a rig, and add that in their view the source of gravity's force will be understood sooner than some people think. General Electric is working on the use of electronic rigs designed to make adjustments to gravity - this line of attack has the advantage of using rigs already in existence for other defence work. Bell also has an experimental rig

intended, as the company puts it, to cancel out gravity, and Lawrence Bell has said he is convinced that practical hardware will emerge from current programs. Grover Leoning is certain that what he referred to as an electro-magnetic contra-gravity mechanism will be developed for practical use. Convair is extensively committed to the work with several rigs. Lear Inc., autopilot and electronic engineers have a division of the company working on gravity research and so also has the Sperry division of Sperry-Rand. This list embraces most of the U.S. aircraft industry. The remainder, Curtiss-Wright, Lockheed, Boeing and North American have not yet declared themselves, but all these four are known to be in various stages of study with and without rigs. In addition, the Massachusetts Institute of Technology is working on gravity, the Gravity Research Foundation of New Boston, the Institute for Advanced Study at Princeton, the CalTech Radiation Laboratory, Princeton University and the University of North Carolina are all active in gravity. Glenn L. Martin is setting up a Research Institute for Advanced Study which has a small staff working on gravity research with the unified field theory and this group is committed to extensive programs of applied research. Many others are also known to be studying gravity, some are known also to be planning a general expansion in this field, such as in the proposed Institute for Pure Physics at the University of North Carolina. A certain amount of work is also going on in Europe. One of the French nationalized constructors and one company outside the nationalized elements have been making preliminary studies, and a little company money has in one case actually been committed. Some work is also going on in Britain where rigs are now in existence. Most of it is private venture work, such as

that being done by Ed Hull a colleague of Townsend Brown who, as much as anybody, introduced Europe to electrogravitics. Aviation Studies' Gravity Research Group is doing some work, mainly on k studies, and is sponsoring dielectric investigations. One Swedish company and two Canadian companies have been making studies, and quite recently the Germans have woken up to the possibilities. Several of the companies have started digging out some of the early German papers on wave physics. They are almost certain to plan a gravitics program. Curiously enough the Germans during the war paid no attention to electrogravitics. This is one line of advance that they did not pioneer in any way and it was basically a U.S. creation. Townsend Brown in electrogravitics is the equivalent of Frank Whittle in gas turbines. This German overlooking of electrostatics is even more surprising when it is remembered how astonishingly advanced and prescient the Germans were in nuclear research. (The modern theory of making thermonuclear weapons without plutonium fission initiators returns to the original German idea that was dismissed, even ridiculed. The Germans never went very far with fission, indeed they doubted that this chain would ever be made to work. The German air industry, still in the embryo stage, has included electrogravitics among the subjects it intends to examine when establishing the policy that the individual companies will adopt after the present early stage of foreign licence has enabled industry to get abreast of the other countries in aircraft development.

It is impossible to read through this summary of the widening efforts being made to understand the nature of matter of gravity without sharing the hope that many groups now have, of major theoretical

breakthroughs occurring before very long. Experience in nucleonics has shown that when attempts to win knowledge on this scale are made, advances are soon seen. There are a number of elements in industry, and some managements, who see gravity as a problem for later generations. Many see nothing in it all and they may be right. But as said earlier, if Dr. Vaclav Hlavaty thinks gravity is potentially controllable that surely should be justification enough, and indeed inspiration, for physicists to apply their minds and for management to take a risk. Hlavaty is the only man who thinks he can see a way of doing the mathematics to demonstrate Einstein's unified field theory - something that Einstein himself said was beyond him. Relativity and the unified field theory go to the root of electrogravitics and the shifts in thinking, the hopes and fears, and a measure of progress is to be obtained only in the last resort from men of this stature. Major theoretical breakthroughs to discover the sources of gravity will be made by the most advanced intellects using the most advanced research tools. Aviation's role is therefore to impress upon physicists of this calibre with the urgency of the matter and to aid them with statistical and peripheral investigations that will help to clarify the background to the central mathematical and physical puzzles. Aviation could also assist by recruiting some of these men as advisers. Convair has taken the initiative with its recently established panel of advisers on nuclear projects, which include Dr. Edward Teller of the University of California. At the same time much can be done in development of laboratory rigs, condenser research and dielectric development, which do not require anything like the same cerebral capacity to get results and make a practical contribution As gravity is likely to be




linked with the new particles, only the highest powered particle accelerators are likely to be of use in further fundamental knowledge. The country with the biggest tools of this kind is in the best position to examine the characteristics of the particles and from those countries the greatest advances seem most likely. Though the United States has the biggest of the bevatron - the Berkeley bevatron is 6.2 bev - the Russians have a 10 bev accelerator in construction which, when it is completed, will be the world's largest. At Brookhaven a 25 bev instrument is in development which, in turn, will be the biggest. Other countries without comparable facilities are of course at a great disadvantage from the outset in the contest to discover the explanation of gravity. Electrogravitics, moreover, unfortunately competes with nuclear studies for its facilities. The clearest thinking brains are bound to be attracted to localities where the most extensive laboratory equipment exists. So, one way and another, results are most likely to come from the major countries with the biggest undertakings. Thus the nuclear facilities have a direct bearing on the scope for electrogravitics work. The OEEC report in January made the following points:- The U.S. has six to eight entirely different types of reactor in operation and many more under construction. Europe has now two different types in service. The U.S. has about 30 research reactors plus four in Britain,, two in France. The U.S. has two nuclear-powered marine engines. Europe has none, but the U.K. is building one. Isotope separation plants for the enrichment of uranium in the U.S. are roughly 11 times larger than the European plant in Britain. Europe's only heavy water plant (in Norway) produces somewhat less than one-twentieth of American output. In 1955 the number of technicians employed in

nuclear energy work in the U.S. was about 15,000; there are about 5,000 in Britain, 1,800 in France, and about 19,000 in the rest of Europe. But the working party says that pessimistic conclusions should not be drawn from these comparisons. European nuclear energy effort is unevenly divided at the moment, but some countries have notable achievements to their credit and important developments in prospect. The main reason for optimism is that, taken as a whole, "Europe's present nuclear effort falls very far short of its industrial potential". Though gravity research, such as there has been of it, has been unclassified, new principles and information gained from the nuclear research facilities that have a vehicle application is expected to be withheld. The heart of the problem to understanding gravity is likely to prove to be the way in which the very high energy sub-nuclear particles convert something, whatever it is, continuously and automatically into the tremendous nuclear and electromagnetic forces. Once this key is understood, attention can later be directed to finding laboratory means of duplicating the process and reversing its force lines in some local environment and returning the energy to itself to produce counterbary. Looking beyond it seems possible that gravitation will be shown to be a part of the universal electro-magnetic processes and controlled in the same way as a light wave or radio wave. This is a synthesis of the Einstein and Hlavaty concepts. Hence it follows that though in its initial form the mechanical processes for countering gravity may initially be massive to deal with the massive forces involved, eventually this could be expected to form some central power generation unit. Barycentric control in some required quantity could be passed over a distance by a form of radio wave. The prime energy

source to energise the waves would of course be nuclear in its origins. It is difficult to say which lines of detailed development being processed in the immediate future is more likely to yield significant results. Perhaps the three most promising are: first, the new attempt by the team of men led by Chamberlain working with the Berkeley bevatron to find the anti-neutron, and to identify more of the characteristics of the anti-proton² and each of the string of high energy particles that have been discovered during recent operation at 6.2 bev. A second line of approach is the United States National Bureau of Standards program to pin down with greater accuracy the acceleration values of gravity. The presently accepted figure of 32.174 feet per second per second is known to be not comprehensive, though it has been sufficiently accurate for the limited needs of industry hitherto. The NBS program aims at re-determining the strength of gravity to within one part of a million. The present method has been to hold a ball 16 feet up and chart the elapsed time of descent with electronic measuring equipment. The new program is based on the old, but with this exceptional degree of accuracy it is naturally immensely more difficult and is expected to take 3 years. A third promising line is the new technique of measuring high energy particles in motion that was started by the University of California last year. This involves passing cosmic rays through a chamber containing a mixture of gas, alcohol and water vapour. This creates charged atoms,


or positive ions, by knocking electrons off the gas molecules. A sudden expansion of the chamber results in a condensation of water droplets along the track which can be plotted on a photographic plate. This method makes it easier to assess the energy of particles and to distinguish one from the other. It also helps to establish the characteristics of the different types of particle. The relationship between these high energy particles and their origin, and characteristics have a bearing on electrogravitics in general. So much of what has to be discovered as a necessary preliminary to gravity is of no practical use by itself. There is no conceivable use, for instance, for the anti-proton, yet its discovery even at a cost of \$9-million is essential to check the mathematics of the fundamental components of matter. Similarly it is necessary to check that all the nuclear ghosts that have been postulated theoretically do in fact exist. It is not, moreover, sufficient, as in the past, only to observe the particles by radiation counters. In each instance a mechanical maze has to be devised and attached to a particle accelerator to trap only the particle concerned. Each discovery becomes a wedge for a deeper probe of the nucleus. Many of the particles of very high energy have only a fleeting existence and collisions that give rise to them from bevatron bombardment is a necessary prerequisite to an understanding of gravity. There are no shortcuts to this process. Most of the major programs for extending human knowledge on gravity are being conducted with instruments already in use for nuclear research and to this extent the cost of work exclusively on gravitational examinations is still not of major proportions. This has made it difficult for aviation to gauge the extent of the work in progress on gravity research.

² The reaction is as follows: protons are accelerated to 6.2 bev, and directed at a target of copper. When the proton projectile hits a neutron in one of the copper atoms the following emerge: the two original particles (the projectile and the struck neutron) and a new pair of particles, a proton and anti-proton. The anti-proton continues briefly until it hits another proton, then both disappear and decay into mesons



Conclusions

1. No attempts to control the magnitude or direction of the earth's gravitational force have yet been successful. But if the explanation of gravity is to be found in the as yet undetermined characteristics of the very high energy particles it is becoming increasingly possible with the bevatron to work with the constituent matter of gravity. It is therefore reasonable to expect that the new bevatron may, before long, be used to demonstrate limited gravitational control.
2. An understanding and identification of these particles is on the frontiers of human knowledge, and a full assessment of them is one of the major unresolved puzzles of the nucleus. An associated problem is to discover a theory to account for the cosmic and quantum relations of gravity, and a theory to link the gravitational constant with the other three dimensionless constants.
3. Though the obstacles to an adequate grasp of microphysics still seem formidable, the transportation rewards that could follow from electrogravitics are as high as can be envisaged. In a weightless environment, movement with sharp-edged changes of direction could offer unique manoeuvrability.
4. Determination of the environment of the anti-proton, discovery of the anti-neutron and closer examination of the other high energy particles are preliminaries to the hypothesis that gravity is one aspect of electromagnetism that may eventually be controlled like a wave. When the structure of the nucleus becomes clearer, the influence of the gravitational force upon the nucleus and the nature of its behaviour in space will be more readily understood. This is a great advance on the Newtonian concept of gravity acting at a distance.
5. Aviation's role appears to be to establish facilities to handle many of the peripheral and statistical investigations to help fill in the background on electrostatics.
6. A distinction has to be made between electrostatic energy for propulsion and counterbary. Counterbary is the manipulation of gravitational force lines; barycentric control is the adjustment to such manipulative capability to produce a stable type of motion suitable for transportation.
7. Electrostatic energy sufficient to produce low speeds (a few thousand dynes, has already been demonstrated. Generation of a region of positive electrostatic energy on one side of a plate and negative on the other sets up the same lift or propulsion effect as the pressure and suction below and above a wing, except that in the case of electrostatic application no airflow is necessary.
8. Electrostatic energy sufficient to produce a Mach 3 fighter is possible with megavolt energies and a k of over 10,000.
9. k figures of 6,000 have been obtained from some ceramic materials and there are prospects of 30,000.



10. Apart from electrogravitics there are other rewards from investment in electrostatic equipment. Automation, autonetics and even turbine development use similar laboratory facilities.
11. Progress in electrogravitics probably awaits a new genius in physics who can find a single equation to tie up all the conflicting observations and theory on the structure and arrangement of forces and the part the high energy particles play in the nucleus. This can occur any time, and the chances are improved now that bev. energies are being obtained in controlled laboratory conditions.

Appendix I

Extracts From Aviation Report Anti-Gravitation Research

The basic research and technology behind electro-anti-gravitation is so much in its infancy that this is perhaps one field of development where not only the methods but the ideas are secret. Nothing therefore can be discussed freely at the moment. Very few papers on the subject have been prepared so far, and the only schemes that have seen the light of day are for pure research into rigs designed to make objects float around freely in a box. There are various radio applications, and aviation medicine departments have been looking for something that will enable them to study the physiological effects on the digestion and organs of an environment without gravity. There are however long term aims of a more revolutionary nature that envisage equipment that can defeat gravity. Aviation Report 20 August 1954

MANAGERIAL POLICY FOR ANTI-GRAVITICS

The prospect of engineers


devising gravity-defeating equipment - or perhaps it should be described as the creation of pockets of weightless environments - does suggest that as a long term policy aircraft constructors will be required to place even more emphasis on electro-mechanical industrial plant, than is now required for the transition from manned to unmanned weapons. Anti-gravitics work is therefore likely to go to companies with the biggest electrical laboratories and facilities. It is also apparent that anti-gravitics, like other advanced sciences, will be initially sponsored for its weapon capabilities. There are perhaps two broad ways of using the science - one is to postulate the design of advanced type projectiles on their best inherent capabilities, and the more critical parameters (that now constitutes the design limitation) can be eliminated by anti-gravitics. The other, which is a longer term plan, is to create an entirely new environment with devices operating entirely under an anti-gravity envelope. Aviation Report 24 August 1954 THE GREATER THE EASIER Propulsion and atomic energy Trends are similar in one respect: the more incredible the long term capabilities are, the easier it is to attain them. It is strange that the greatest of nature's secrets can be harnessed with decreasing industrial effort, but greatly increasing mental effort. The Americans went through the industrial torture to produce tritium for the first thermonuclear experiment, but later both they and the Russians were able to achieve much greater results with the help of lithium 6 hydride. The same thing is happening in aviation propulsion: the nuclear fuels are promising to be tremendously powerful in their effect, but excessively complicated in their application, unless there can be some means of direct conversion as in the strontium 90 cell. But lying behind and

beyond the nuclear fuels is the linking of electricity to gravity, which is an incomparably more powerful way of harnessing energy than the only method known to human intellect at present - electricity and magnetism. Perhaps the magic of barium aluminium oxide will perform the miracle in propulsion that lithium 6 hydride has done in the fusion weapon. Certainly it is a well-known material in dielectrics, but when one talks of massive-k, one means of course five figures. At this early stage it is difficult to relate k to Mach numbers with any certainty, but realizable k can, with some kinds of arithmetic, produce astounding velocities. They are achievable, moreover, with decreasing complexity, indeed the ultimate becomes the easiest in terms of engineering, but the most hideous in terms of theory. Einstein's general theory of relativity is, naturally, and important factor, but some of the postulates appear to depend on the unified field theory, which cannot yet be physically checked because nobody knows how to do it. Einstein hopes to find a way of doing so before he dies.

Aviation Report 31 August 1954

GRAVITICS FORMULATIONS All indications are that there has still been little cognizance of the potentialities of electrostatic propulsion and it will be a major undertaking to re-arrange aircraft plants to conduct large-scale research and development into novel forms of dielectric and to improve condenser efficiencies and to develop the novel type of materials used for fabrication of the primary structure. Some extremely ambitious theoretical programs have been submitted and work towards realization of a manned vehicle has begun. On the evidence, there are far more definite indications that the incredible claims are realizable than there was, for instance, in supposing that uranium fission would result in a bomb. At

least it is known, proof positive, that motion, using surprisingly low k, is possible. The fantastic control that again is feasible, has not yet been demonstrated, but there is no reason to suppose the arithmetic is faulty, especially as it has already led to a quite brisk example of actual propulsion. That first movement was indeed an historic occasion, reminiscent of the momentous day at Chicago when the first pile went critical, and the phenomenon was scarcely less weird. It is difficult to imagine just where a well-organised examination into long term gravitics prospects would end. Though a circular platform is electrostatically convenient, it does not necessarily follow that the requirements of control by differential changes would be the same. Perhaps the strangest part of this whole chapter is how the public managed to foresee the concept though not of course the theoretical principles that gave rise to it, before physical tests confirmed that the mathematics was right. It is interesting also that there is no point of contact between the conventional science of aviation and the New: it is a radical offshoot with no common principles. Aerodynamics, structures, heat engines, flapping controls, and all the rest of aviation is part of what might be called the Wright Brothers era - even the Mach 2.5 thermal barrier piercers are still Wright Brothers concepts, in the sense that they fly and they stall, and they run out of fuel after a short while, and they defy the earth's pull for a short while. Thus this century will be divided into two parts - almost to the day. The first half belonged to the Wright Brothers who foresaw nearly all the basic issues in which gravity was the bitter foe. In part of the second half gravity will be the great provider. Electrical energy, rather irrelevant for propulsion in the first half becomes a kind of catalyst to motion, in the second half of



the century. Aviation Report 7 September 1954

ELECTRO-GRAVITICS PARADOX

Realization of electro-static propulsion seems to depend on two theoretical twists and two practical ones. The two theoretical puzzles are: first, how to make a condenser the centre of a propulsion system, and second is how to link the condenser system with the gravitational field. There is a third problem, but it is some way off yet, which is how to manipulate kva for control in all three axes as well as for propulsion and lift. The two practical tricks are first how, with say a Mach 3 weapon in mind, to handle 50,000 kva within the envelope of a thin pancake of 35 feet in diameter and second how to generate such power from within so small a space. The electrical power in a small aircraft is more than is a fair sized community the analogy being that a single rocketjet can provide as much power as can be obtained from the Hoover Dam. It will naturally take as long to develop electro-static propulsion as it has taken to coax the enormous power outputs from heat engines. True there might be a flame in the electro-gravitic propulsion system, but it would not be a heat engine - the temperature of the flame would be incidental to the function of the chemical burning process. The curious thing is that though electro-static propulsion is the antithesis of magnetism,³ Einstein's unified field theory is an attempt to link gravitation with electro-magnetism. This all-embracing theory goes on logically from the general theory of relativity, that gives an ingenious geometrical interpretation of the concept of force which is mathematically consistent with gravitation but fails in the case of electro-magnetism, while the special theory of

relativity is concerned with the relationship between mass and energy. The general theory of relativity fails to account for electro-magnetism because the forces are proportional to the charge and not to the mass. The unified field theory is one of a number of attempts that have been made to bridge this gap, but it is baffling to imagine how it could ever be observed. Einstein himself thinks it is virtually impossible. However Hlavaty claims now to have solved the equations by assuming that gravitation is a manifestation of electro-magnetism. This being so it is all the more incredible that electro-static propulsion (with kva for convenience fed into the system and not self-generated) has actually been demonstrated. It may be that to apply all this very abstruse physics to aviation it will be necessary to accept that the theory is more important than this or that interpretation of it. This is how the physical constants, which are now regarded as among the most solid of achievements in modern physics, have become workable, and accepted. Certainly all normal instincts would support the Einstein series of postulations, and if this is so it is a matter of conjecture where it will lead in the long term future of the electro-gravitic science.

Aviation Report, 10 September 1954

Electro-Gravitic Propulsion Situation

Under the terms of Project Winterhaven the proposals to develop electro-gravitics to the point of realizing a Mach 3 combat type disc were not far short of the extensive effort that was planned for the Manhattan District.⁴ Indeed the drive to develop the new prime mover is in some respects rather similar to the experiments

³ Though in a sense this is true, it is better expressed in the body of this report than it was here in 1954.

⁴ The proposals, it should be added, were not accepted.

that led to the release of nuclear energy in the sense that both involve fantastic mathematical capacity and both are sciences so new that other allied sciences cannot be of very much guide. In the past two years since the principle of motion by means of massive-k was first demonstrated on a test rig, progress has been slow. But the indications are now that the Pentagon is ready to sponsor a range of devices help further knowledge. In effect the new family of TVs would be on the same tremendous scope that was envisaged by the X-1, 2, 3, 4 and 5 and D.558s that were all created for the purpose of destroying the sound barrier - which they effectively did, but it is a process that is taking ten solid years of hard work to complete. (Now after 7 years the X-2 has yet to start its tests and the X-3 is still in performance testing stage). Tentative targets now being set anticipate that the first disc should be complete before 1960 and it would take the whole of the 'sixties to develop it properly, even though some combat things might be available ten years from now. One thing seems certain at this stage, that the companies likely to dominate the science will be those with the biggest computers to work out the ramifications of the basic theory. Douglas is easily the world's leader in computor capacity, followed by Lockheed and Convair. The frame incidentally is indivisible from the engine". If there is to be any division of responsibility it would be that the engine industry might become responsible for providing the electrostatic energy (by, it is thought, a kind of flame) and the frame maker for the condenser assembly which is the core of the main structure.

Aviation Report 12 October 1954

Gravitics Study Widening

The French are now understood to be pondering the most effective way of entering the field of electro-gravitic propulsion systems. But not least of the difficulties is to know just where to begin. There are practically no patents so far that throw very much light on the mathematics of the relation between electricity and gravity. There is, of course, a large number of patents on the general subject of motion and force, and some of these may prove to have some application. There is, however, a series of working postulations embodied in the original Project Winterhaven, but no real attempt has been made in the working papers to go into the detailed engineering. All that had actually been achieved up to just under a year ago was a series of fairly accurate extrapolations from the sketchy data that has so far been actually observed. The extrapolation of 50 mph to 1,800 mph, however, (which is what the present hopes and aspirations amount to) is bound to be a rather vague exercise. This explains American private views that nothing can be reasonably expected from the science yet awhile. Meanwhile, the NACA is active, and nearly all the Universities are doing work that borders close to what is involved here, and something fruitful is likely to turn up before very long. Aviation Report 19 October 1954 GRAVITICS STEPS Specification writers seem to be still rather stumped to know what to ask for in the very hazy science of electro-gravitic propelled vehicles. They are at present faced with having to plan the first family of things - first of these is the most realistic type of operational test rig, and second the first type of test vehicle. In turn this would lead to sponsoring of a combat disc. The preliminary test rigs which gave only feeble propulsion have been somewhat improved, but of course the speeds reached so far are only those more associated with what is attained on the

roads rather than in the air. But propulsion is now known to be possible, so it is a matter of feeding enough KVA into condensers with better k figures. 50,000 is a magic figure for the combat saucer - it is this amount of KVA and this amount of k that can be translated into Mach 3 speeds. Meanwhile Glenn Martin now feels ready to say in public that they are examining the unified field theory to see what can be done. It would probably be truer to say that Martin and other companies are now looking for men who can make some kind of sense out of Einstein's equations. There's nobody in the air industry at present with the faintest idea of what it is all about. Also, just as necessary, companies have somehow to find administrators who know enough of the mathematics to be able to guess what kind of industrial investment is likely to be necessary for the company to secure the most rewarding prime contracts in the new science. This again is not so easy since much of the mathematics just cannot be translated into words. You either understand the figures, or you cannot ever have it explained to you. This is rather new because even things like indeterminacy in quantum mechanics can be more or less put into words. Perhaps the main thing for management to bear in mind in recruiting men is that essentially electro-gravitics is a branch of wave technology and much of it starts with Planck's dimensions of action, energy and time, and some of this is among the most firm and least controversial sections of modern atomic physics. Aviation Report 19 November 1954. ELECTROGRAVITICS PUZZLE Back in 1948 and 49, the public in the U.S. had a surprisingly clear idea of what a flying saucer should, or could, do. There has never at any time been any realistic explanation of what propulsion agency could make it do those things, but its

ability to move within its own gravitation field was presupposed from its manoeuvrability. Yet all this was at least two years before electro-static energy was shown to produce propulsion. It is curious that the public were so ahead of the empiricists on this occasion, and there are two possible explanations. One is that optical illusions or atmospheric phenomena offered a preconceived idea of how the ultimate aviation device ought to work. The other explanation might be that this was a recrudescence of Jung's theory of the Universal Mind which moves up and down in relation to the capabilities of the highest intellects and this may be a case of it reaching a very high peak of perception. But for the air industries to realize an electro-gravitic aircraft means a return to basic principles in nuclear physics, and a re-examination of much in wave technology that has hitherto been taken for granted. Anything that goes any way towards proving the unified field theory will have as great a bearing on electro-gravitics efforts as on the furtherance of nuclear power generally. But the aircraft industry might as well face up to the fact that priorities will in the end be competing with the existing nuclear science commitments. The fact that electro-gravitics has important applications other than for a weapon will however strengthen the case for governments to get in on the work going on. Aviation Report 28 January 1955 MANAGEMENT NOTE FOR ELECTRO-GRAVITICS The gas turbine engine produced two new companies in the U.S. engine field and they have, between them, at various times offered the traditional primes rather formidable competition. Indeed GE at this moment has, in the view of some, taken the Number Two position. In Britain no new firms managed to get a footing but one, Metro-Vick, might have done if it had put

its whole energies into the business. It is on the whole unfortunate for Britain that no bright newcomer has been able to screw up competition in the engine field as English Electric have done in the airframe business. Unlike the turbine engine, electro-gravitics is not just a new propulsion system, it is a new mode of thought in aviation and communications, and it is something that may become all-embracing. Theoretical studies of the science unfortunately have to extend right down to the mathematics of the meson and there is no escape from that. But the relevant facts wrung from the nature of the nuclear structure will have their impact on the propulsion system, the airframe and also its guidance. The airframe, as such, would not exist, and what is now a complicated stressed structure becomes some convenient form of hard envelope. New companies therefore who would like to see themselves as major defence prime contractors in ten or fifteen years time are the ones most likely to stimulate development. Several typical companies in Britain and the U.S. come to mind - outfits like AiResearch, Raytheon, Plessey in England, Rotax and others. But the companies have to face a decade of costly research into theoretical physics and it means a great deal of trust. Companies are mostly overloaded already and they cannot afford it, but when they sit down and think about the matter they can scarcely avoid the conclusion that they cannot afford not to be in at the beginning. Aviation Report 8 February 1955 ELECTRO-GRAVITICS BREAKTHROUGHS Lawrence Bell said last week he thought that the tempo of development leading to the use of nuclear fuels and anti-gravitational vehicles (he meant presumably ones that create their own gravitational field independently of the earth's) would accelerate. He added that the breakthroughs now feasible will

advance their introduction ahead of the time it has taken to develop the turbojet to its present pitch. Beyond the thermal barrier was a radiation barrier, and he might have added ozone poisoning and meteorite hazards, and beyond that again a time barrier. Time however is not a single calculable entity and Einstein has taught that an absolute barrier to aviation is the environmental barrier in which there are physical limits to any kind of movement from one point in space-time continuum to another. Bell (the company not the man) have a reputation as experimentalists and are not so earthy as some of the other U.S. companies; so while this first judgement on progress with electrogravitics is interesting, further word is awaited from the other major elements of the air business. Most of the companies are now studying several forms of propulsion without heat engines though it is early days yet to determine which method will see the light of day first. Procurement will open out because the capabilities of such aircraft are immeasurably greater than those envisaged with any known form of engine. Aviation Report 15 July 1955

THERMONUCLEAR-

ELECTROGRAVITICS INTERACTION The point has been made that the most likely way of achieving the comparatively low fusion heat needed - 1,000,000 degrees provided it can be sustained (which it cannot be in fission for more than a microsecond or two of time) - is by use of a linear accelerator. The concentration of energy that may be obtained when accelerators are rigged in certain ways make the production of very high temperatures feasible but whether they could be concentrated enough to avoid a thermal heat problem remains to be seen. It has also been suggested that linear accelerators would be the way to develop the high electrical energies needed for

creation of local gravitational systems. It is possible therefore to imagine that the central core of a future air vehicle might be a linear accelerator which would create a local weightless state by use of electrostatic energy and turn heat into energy without chemical processes for propulsion. Eventually - towards the end of this century - the linear accelerator itself would not be required and a ground generating plant would transmit the necessary energy for both purposes by wave propagation. Aviation Report 30 August 1955 POINT ABOUT THERMONUCLEAR REACTION REACTORS The 20 year estimate by the AEC last week that lies between present research frontiers and the fusion reactor probably refers to the time it will take to tap fusion heat. But it may be thought that rather than use the molecular and chemical processes of twisting heat into thrust it would be more appropriate to use the now heat source in conjunction with some form of nuclear thrust producer which would be in the form of electrostatic energy. The first two Boeing nuclearjet prototypes now under way are being designed to take either molecular jets or nuclear jets in case the latter are held up for one reason or another. But the change from molecular to direct nuclear thrust production in conjunction with the thermonuclear reactor is likely to make the aircraft designed around the latter a totally different breed of cat. It is also expected to take longer than two decades, though younger executives in trade might expect to live to see a prototype. Aviation Report 14 October 1955 ELECTROGRAVITICS FEASIBILITY Opinion on the prospects of using electrostatic energy for propulsion, and eventually for creation of a local gravitational field isolated from the earth's has naturally polarized into the two opposite extremes. There are those who

say it is nonsense from start to finish, and those who are satisfied from performance already physically manifest that it is possible and will produce air vehicles with absolute capabilities and no moving parts. The feasibility of a Mach 3 fighter (the present aim in studies) is dependent on a rather large k extrapolation, considering the pair of saucers that have physically demonstrated the principle only achieved a speed of some 30 fps. But, and this is important, they have attained a working velocity using very inefficient (even by today's knowledge) form of condenser complex. These humble beginnings are surely as hopeful as Whittle's early postulations. It was, by the way, largely due to the early references in Aviation Report that work is gathering momentum in the U.S. Similar studies are beginning in France, and in England some men are on the job full time. Aviation Report 15 November 1955 ELECTRO-GRAVITICS EFFORT WIDENING Companies studying the implications of gravitics are said in a new statement, to include Glenn Martin, Convair, Sperry-Rand, Sikorsky, Bell, Lear Inc. and Clark Electronics. Other companies who have previously evinced interest include Lockheed Douglas and Hiller. The remainder are not disinterested, but have not given public support to the new science - which is widening all the time. The approach in the U.S. is in a sense more ambitious than might have been expected. The logical approach, which has been suggested by Aviation Studies, is to concentrate on improving the output of electrostatic rigs in existence that are known to be able to provide thrust. The aim would be to concentrate on electrostatics for propulsion first and widen the practical engineering to include establishment of local gravity forcefields, independent of those of the earth's, to provide unfettered vertical

movement as and when the mathematics develops. However, the U.S. approach is rather to put money into fundamental theoretical physics of gravitation in an effort first to create the local gravitation field. Working rigs would follow in the wake of the basic discoveries. Probably the correct course would be to sponsor both approaches, and it is now time that the military stepped in with big funds. The trouble about the idealistic approach to gravity is that the aircraft companies do not have the men to conduct such work. There is every expectation in any case that the companies likely to find the answers lie outside the aviation field. These would emerge as the masters of aviation in its broadest sense. The feeling is therefore that a company like A. T. & T. is most likely to be first in this field. This giant company (unknown in the air and weapons field) has already revolutionized modern warfare with the development of the junction transistor and is expected to find the final answers to absolute vehicle levitation. This therefore is where the bulk of the sponsoring money should go. Aviation Report 9 December 1955

APPENDIX II ELECTROSTATIC PATENTS [This following table has been retyped to fit this 60 column file format] [The report has the dates before the titles on the same line.] [All other formatting and spelling is exact.] ELECTROSTATIC MOTORS (a) American Patents still in force. 2,413,391 Radio Corp. America Power Supply System 20-6-42/31-12-46 2,417,452 Ratheon Mfg. Co. Electrical System 17-1-44/18- 3-47 2,506,472 W.B. Smits Electrical Ignition Apparatus 3-7-46Holl/ 2- 5-50 2,545,354 G.E.C. Generator 16-3-50/13- 3-51 (- Engl.P.676,953) 2,567,373 Radio Corp. America El'static Generator 10-6-49/11- 9-51 2,577,446 Chatham Electronics El'static

Voltage Generator 5-8- 50/ 4-12-51
2,578,908 US-Atomic Energy C. El'static
Voltage Generator 26-5-47/18-12-51
2,588,513 Radio Corp. America El'static
High-Voltage Generator 10-6-49/11- 3-52
2,610,994 Chatham Electronics El'static
Voltage Generator 1-9-50/16- 9-52
2,662,191 P. Okey El'static Machine 31-7-52/8 -12-53 2,667,615 R.G. Brown
El'static Generator 30-1-52/26- 1-54
2,671,177 Consolidated Engg. Corp
El'static Charging App's 4-9-51/2 - 3-54
2,701,844 H.R. Wasson El'static Generator
of Electricity 8-1250/ 8- 2-55 2,702,353
US-Navy Miniature Printed Circuit
Electrostatic Generator 17-7-52/15- 2-55
(b) British patents still in force. 651,153
Metr.-Vickers Electr.Co. Voltage
Transformation of electrical energy 20-5-48/14- 3-51 651,295 Ch.F.Warthen sr.
(U.S.A.) Electrostatic A.C. Generator 6- 8-48/14- 3-51 731,774 "Licentia" El'static
High-Voltage Generator 19-9-52 & 21-11-52Gy/ 15- 6-55 (c) French patents still in force. 753,363 H. Chaumat Moteur electrostatique utilisant l'energie cinetique d'ions gazeux 19-7-32/13-10-33 . 749,832
H. Chaumat Machine electrostatique a excitation independante 24-1-32/29- 7-33
[This table has been retyped to fit this 60 column file format] [The report has the titles at the end of each line, w/ o ()] [All other formatting and spelling is exact.] The following patents derive from P. Jolivet (Algiers), marked "A" and from N.J. Felici, E. Gartner (Centre National des Recherches Scientifique - CRNS -) later also by R. Morel, M. Point etc. (S.A. des Machines Electrostatiques -SAMES-) and of Societe' d'Appareils de Controle et d'Equipment des Moteurs -SACEM-), marked "G " (because the development was centred at the University Grenoble. Mark of Application England America France Germany (Title) Applicant Date ---

----- G 8-11-44) 993,017 14- 8-45) 637,434
2,486,140 56,027 860,649 (Electrostatic Influence Machine G 17-11-44 639,653
2,523,688 993,052 815,667 (Electrostatic Influence Machine A 28- 2-45 912,444
(Inducteurs de Machines el'static) G 3- 3-
45 643,660 2,519,554 995,442 882,586
(El'static Machines A 8- 6-45 915,929
(Machines electrocstatiques a flasques) A
16- 8-45 918,547 (Generatrice el'statique)
G 20- 9-45) 998,397 21- 9-45) 643,664
2,523,689 837,267 (Electrostatic Machines) A
4- 2-46 923,593 (Generatrice el'statique) G
17- 7-46 643,579 2,530,193
1002,031 811,595 (Generating Machines)
G 20- 2-47 671,033 2,590,168 (Ignition device)
G 21-3 -47 2,542,494 655,474 Re-
23,560 944,574 860,650 (El'static Machines) G
6- 6-47 645,916 2,522,106
948,409 810,042 (El'static Machines) A
16- 6-47 947,921 (Generatrice el'statique)
G 16- 1-48 669,645 2,540,327 961,210
810,043 (El'static Machines) G 21- 1-49
669,454 2,617,976 997,991 815,666
(El'static Machines) G 7- 2-49 675,649
2,649,566 1010,924 870,575 (El'static Machines) G
15- 4-49 693,914 2,604,502
1011,902 832,634 (Commutators for electrical machine) G
9-11-49 680,178
2,656,502 1004,950 850,485 (El'static Generate) G
9-10-50) 702,494 2,675,516
1030,623 20- 2-51) (El'static Generate) G
29-11-50) 702,421 1028,596 20- 2-51)
(El'static Generate) G 21-11-51 719,687
1051,430 F10421 (El'static Machines) G
20- 8-52 731,773 2,702,869 938,198
(El'static Machines) G 6-11-52 745,489
(El'static Generator) G 12- 2-53 745,783
(Rotating El'static Machines) G 8- 1-52
715,010 2,685,654 1047,591 (Rotating El'static Machines producing a periodical discharge.) Appl'n.No. ----- G 27- 2-54
5726/55 (El'static Machines) G 8- 3-54
6790/ 55 (El'static Machines) G 28- 1-55
2748/56 (El'static Machines) NOTE:- ALL
THE LISTED PATENTS ARE STILL IN


FORCE "The Flying Saucer" The Application of the Biefeld-Brown Effect to the Solution of the Problems of Space Navigation by Mason Rose, Ph.D., President University for Social Research (1952) Published in Science and Invention, August 1959, and Psychic Observer, Vol. XXXVII, No. 1 Documents Menu The scientist and layman alike encounter a primary difficulty in understanding the Biefeld-Brown effect and it's relation to the solution of the flying saucer mystery. A proper interpretation of this theory is prevented because both scientist and layman are conditioned to think in electromagnetic concepts, whereas the Biefeld-Brown effect relates to electrogravitation. Their lack of awareness is justifiable, however, because the data on electrogravitation, inasmuch as it is a comparatively recent and unpublished development, has limited availability and circulation. Townsend Brown, the discoverer of electrogravitational coupling, is the only known experimental scientist in this new area of scientific development as of this writing. Thus, anyone wanting to understand electrogravitation and its applications to astronautics must dismiss the principles of electromagnetics in order to grasp the essentially different principles of electrogravitation. Electrogravitational effects do not obey the known principles of electromagnetism. Electrogravitation must be understood as an entirely new field of scientific investigation and technical development. The most efficient method of effecting an understanding of electrogravitation is to review the evolutionary development of electromagnetism. From the smallest atom to the largest galaxy, the universe operates on three basic forces, namely Electricity, Magnetism and Gravitation. These forces can be represented as follows: Taken separately, these forces are of no real

practical use. Electricity by itself is static electricity and therefore functionless. It will make your hair stand on end, but that is about all. Magnetism by itself has very few practical applications aside from the magnetic compass, and gravity simply keeps objects and people pinned to the earth. However, when they are used to work in combination with each other, almost endless technical applications come into being. Currently, our total electrical development is based on the coupling of electricity with magnetism, which provides the basis for the countless uses we make of electricity in modern societies. Faraday conducted the first productive empirical experiment with electromagnetism around 1830, and Maxwell did the basic theoretical work in 1865. The application of electromagnetism to microscopic and sub-microscopic particles was accomplished by Max Planck's work in quantum physics about 1890; and then in 1905 Einstein came forward with relativity, which dealt with gravitation as applied to celestial bodies and universal mechanics. It is principally out of the work of these four great scientists that our electrical developments, ranging from the simple lightbulb to the complexities of nuclear physics, have emerged. In 1923, Dr. Biefeld, Professor of Physics and Astronomy at Dennison University and a former classmate of Einstein in Switzerland, suggested to his protoge, Townsend Brown, certain experiments which led to the discovery of the Biefeld-Brown effect, and ultimately to the electrogravitational energy spectrum (in actuality, it was Brown who first observed the effect and brought it to the attention of Dr. Biefeld, who suggest further experiments to determine the origin of and enhance the effect - Juniper). Biefeld wondered if an electrical condenser, hung by a thread, would have a tendency to

move when it was given a heavy electrical charge. Townsend Brown provided the answer. There is such a tendency. After 28 years of investigation by Brown into the coupling effect between electricity and gravitation, it was found that for each electromagnetic phenomenon there exists an electrogravitational analogue. This means, from the technical and commercial viewpoint, potentialities for future development and exploitation are as great or greater than the present electrical industry. When one considers that electromagnetism is basic to the telephone, telegraph, radio, television, radar, electric generators and motors, power production and distribution, and is an indispensable adjunct to transportation of all kinds, one can see that the possibility of a parallel, but different development in electrogravitation has almost unlimited prospects. The initial experiments conducted by Townsend Brown, concerning the behavior of a condenser when charged with electricity, had the characteristic of simplicity which has marked most other great scientific advancements. The first startling revelation was that if placed in free suspension with the poles horizontal, the condenser, when charged, exhibited a forward thrust toward the positive poles. A reversal of polarity caused a reversal of the direction of thrust. The experiment was set up as follows: The antigravity effect of vertical thrust is demonstrated by balancing a condenser on a beam balance and then charging it. After charging, if the positive pole is pointed upward, the condenser moves up. If the charge is reversed and the positive pole pointed downward, the condenser thrusts down. The experiment is conducted as follows: These two simple experiments demonstrate what is now known as the Biefeld-Brown effect. It is the first and, to the best of our knowledge, the only method of affecting a gravitational field by


electrical means. It contains the seeds of control of gravity by man. The intensity of the effects is determined by five factors, which are: 1. The separation of the plates of the condenser, the closer the plates, the greater the effect. 2. The ability of the material between the plates to store the electrical energy in the form of elastic stress. A measure of this ability is called the "K" of the material. The higher the "K," the greater the Biefeld-Brown effect. 3. The area of the plates, the greater the area giving the greater effect. 4. The voltage difference between the plates; more voltage, more effect. 5. The mass of the material between the plates; the greater the mass, the greater the effect. It is this fifth point which is inexplicable from the electromagnetic viewpoint and which provides the connection with gravitation. On the basis of further experimental work from 1923 to 1926; Townsend Brown in 1926, described what he called a "space car." This was a revolutionary method of terrestrial and extra-terrestrial flight, presented for experiment while motor propelled planes were yet in a primitive stage. This engineering feat by Townsend Brown was all the more remarkable when we consider such a machine produces thrust with no moving parts, does not use any aerodynamic principles of flight, and has neither control surfaces, or a propeller. Townsend Brown had discovered the secret of how the flying saucers fly years before and such objects were reported. Now the basic differences between electromagnetism and electrogravity have been described and the basic principles of the Biefeld-Brown effect have been outlined, we are finally ready to understand the principles of astronautics or the conquest of space. The earth creates and is surrounded with a gravitational field which approaches zero as we go far into space. This field presses objects and

people to the earth's surface; hence it presses a saucer object to the earth. However, through the utilization of the Biefeld-Brown effect, the flying saucer can generate an electrogravitational field of its own which modifies the earth's field. This field acts like a wave, with the negative pole at the top of the wave and the positive pole at the bottom, the saucer travels like a surfboard on the incline of a wave that is kept continuously moving by the saucer's electrogravitational generator. Since the orientation of the field can be controlled, the saucer can thus travel on its own continuously generated wave in any desired angle or direction of flight. Since the saucer always moves towards its positive pole, the control of the saucer is accomplished by varying the orientation of the positive charge. Control, therefore, is gained by switching charges rather than by the control surfaces. Since the saucer is traveling on the incline of a continually moving wave which it generates to modify the earth's gravitational field, no mechanical propulsion is necessary. Once we understand that the horizontal and vertical controls are obtained by shifting the positive pole which turns the field, then we are in a position to extrapolate a finished saucer design. The method of controlling the flight of the saucer is illustrated by the following simple diagrams showing the charge variations necessary to accomplish all directions of flight. The saucer's edge would contain a number of conductor segments, and the saucer would turn in any direction simply by shifting the positive and negative charges to appropriate positions along its edge. The vertical thrust would be regulated by varying the charge on top of the saucer, the amount of thrust being regulated by the amount of charge generated. In all probability, flying saucers do not utilize external controls for



direction, nor do they have any visible means of propulsion. Flying saucers travel using the Biefeld-Brown electrogravitational effect, and hence do not utilize any of the standard aerodynamic principles of an airfoil. Flying saucers cannot be understood from the traditional principles of aeronautical engineering; however, the older points of view are useful for critical theoretical analysis and empirical testing. Before UFO's were ever seen and validly reported, Townsend Brown developed a captive flying saucer - a scale model saucer with a free bearing going around a stationary pole. Brown did not start with round objects, in fact, the first object that he flew was a triangle, the next a square, then a square with the edges cut off, and finally a round shaped saucer. Eventually, experiments proved the saucer shape most effective. Changes were made for empirical reasons. Having solved the problem of horizontal thrust, Townsend Brown developed a profile shape which would be most efficient to navigate the electrogravitational field for maximum vertical thrust. The final profile that developed was the shape illustrated here: The first report of a disc-shaped object in the sky dates back to the sixteenth century. At long intervals during the centuries since then have come other reports. Most of them are undoubtedly unreliable as observations, distorted by telling and retelling. But in these older reports, as well as in the very numerous series which has accumulated since 1947, there is a teasing common thread concerning appearance and behavior which makes any certainties about the unreality of flying saucers very insecure. One of the great difficulties in substantiation of these reports is that, in both appearances and behavior, these objects seem to be simple scientific impossibilities. Here are some of the reasons advanced by technical men to

prove the impossibility of devices such as the reports describe: 1. The reports reveal, in most cases, no method of propulsion which can be understood. There are no propellers in any of the reports. Some of the reports describe a long flame jet trailing behind a cigar shaped object. But this flame is orange-red in color, indicating an inefficient combustion which would make it ineffective as a reaction jet such as propels rockets and jet planes. No other known physical laws seemed capable of explaining the observed motion of the objects. 2. The reports describe a range of speed and acceleration from stationary hovering to speeds greater than present day rockets can deliver, and the changes of rate of motion, the accelerations, are far beyond the capacities of any known man-made vehicles. Flight experts point out that such accelerations would impose impossible stresses on any human or human like occupants. therefore, they say, the reports must be false or erroneous. many of the reports concern night sightings and describe a glow, usually of blue or violet color, around a periphery of the objects. Physicists have noted that such a glow is characteristic of a very high voltage electrical discharge, but add that this suggests no means of explaining the appearance or behavior of the objects described in the reports. 4. The description of shapes and performance seems to indicate a complete or almost complete disregard of aerodynamic principles. The objects seem not to need the support of air as a plane does, nor to depend on the lift provided by properly designed surfaces moved rapidly through an air medium. These are weighty arguments PROVIDED THE ASSUMPTIONS BEHIND THEM ARE CORRECT. As I have previously indicated, the observed motion of condenser has been labeled the Biefeld-Brown effect. Studying this effect, Brown




pointed out in 1923 that this tendency of a charged condenser to move might easily grow into a new and basically different method of propulsion. By 1926 he had described a "space car" utilizing this new principle. By 1928 he had built working models of a boat propelled in this manner. By 1938 he had shown that his specially designed condensers not only moved, but had certain interesting effects on plants and animals. All of this, while very exciting, is for most of us just a repetition and reinforcement of the rapid scientific development so characteristic of our age. But then came the unexpected Townsend Brown, working in his laboratory, building models and trying endless variations in size, shape and design of his charged condensers, made a flying saucer which flew around a maypole, before flying saucers became a newspaper topic. And the reasons listed above, which led the specialists to reject the reports of observed saucers, proved to be both explicable and necessary to their operation under the electrogravitational principle. Let us look at the four main objectives in a new light:

1. No understood method of propulsion. The saucer made by Brown have no propellers, no jets, no moving parts at all. They create a modification of the gravitational field around themselves, which is analogous to putting them on the incline of a hill. They like a surfboard on a wave. The surfboard moves without propellers or jets to, but it is confined to the direction and speed of the water wave. The electrogravitational saucer creates its own "hill," which is a local distortion of the gravitational field, then it takes this "hill" with it in any chosen direction at any rate.
2. The second objection concerned the tremendous accelerations which on the basis of previous technology, would subject any animal occupants to unbearable stresses. But, says Brown, the

occupants of one of his saucers would feel no stress at all, no matter how sharp the turn or how great the acceleration. This is because the ship and all the occupants and the load are all responding equally to the wavelike distortion of the local gravitational field. In an airplane the propeller pumps air backward and, by reaction, the plane moves forward. The reaction thrust on the propeller is transferred to the frame of the aircraft. This frame then shoves the load and occupants forward *contrary to their natural tendency to move at a constant rate in a constant direction*. But in the saucer no such transfers of thrust from one member to another occurs. The entire assembly moves in unison in response to the locally modified gravitational field. The nearest analogy in our experience is going down in an elevator. When the elevator starts down, it is not necessary for the elevator to shove on our bodies, both elevator and passengers share a gravitational tendency to move down. They do so without and shoving or any stresses between elevator and passengers.

3. Townsend Brown's saucers require a highly charged leading edge - the positive pole. But such a charged pole produces an electrical corona. In the largest models made, this develops a decided bluish-violet glow easily in the darkness or a dim light. A full scale ship operating on this principle would be expected to produce a spectacular corona effect visible for many miles.
4. The outlines and shape of Brown's saucers were the result of electrogravitational considerations, not the result of wind tunnel tests of aerodynamic designs. For they move, not on the lift of air, but on the lift of a modified gravitational field. In operating flying saucers such aerodynamic considerations would have to be taken into account to reduce drag and friction, but not to produce lift and thrust.
5. And, finally,




when Brown turned his attention to improved ways of generating high voltages, the most promising new method involved the use of a flame jet to convey negative charges astern. This flame was relatively inefficient as a generator if it was adjusted for the best combustion of the fuel. But if it was adjusted to an orange-red color, indicating incomplete combustion of fuel, it conveyed the charges very effectively and set up the required negative space charge behind the ship. The reasons advanced by the experts to "explain away" the saucer reports, when seen from a new and different viewpoint appear to be the specific reasons why they can operate, on electrogravitational rather than electromagnetic principles. The next opinion which must be corrected is the idea of overly intensified supersonic vibration. The Townsend Brown experiments indicate that the positive field which is traveling in front of the saucer acts as a buffer wing which starts moving the air out of the way. This immaterial electrogravitational field acts as an entering wedge which softens the supersonic barrier, thus allowing the material leading edge to enter into a softened pressure area. Diagramed, this would be illustrated as follows: It should be noted that in a jet plane or guided missile the extra weight added to create the Biefeld-Brown electrogravitational effect would be compensated for by the added thrust created by the movement of the plane toward the positive field created in front of the leading edge. As we have previously stated, for every known electromagnetic effect there is an analogous electrogravitational effect but electrogravitational applications and results differ from those of electromagnetic. This presupposes that an entire new electrogravitational industry comparable to the present electromagnetic

industry will emerge from the theoretical formulations and empirical experiments of Townsend Brown. INDEXED: BROWN United States Patent Office 3,187,206 Patented June 1, 1965

3,187,206 ELECTROKINETIC APPARATUS Thomas Townsend Brown, Walkertown, N.C., assigner, by mesne assignments, to Electrokinetics, Inc., a corporation of Pennsylvania Filed May 9, 1958, Ser. No. 734,342 23 Claims (Cl. 310-5) This invention relates to an electrical device for producing a thrust by the direct operation of electrical fields. I have discovered that a shaped electrical field may be employed to propel a device relative to its surroundings in a manner which is both novel and useful. Mechanical forces are created which move the device continuously in one direction while the masses making up the environment move in the opposite direction. When the device is operated in a dielectric fluid medium, such as air, the forces of reaction appear to be present in that medium as well as on all solid material bodies making up the physical environment. In a vacuum, the reaction forces appear on the solid environmental bodies, such as the walls of the vacuum chamber. The propelling force however is not reduced to zero when all environmental bodies are removed beyond the apparent effective range of the electrical field. By attaching a pair of electrodes to opposite end of a dielectric member and connecting a source of high electrostatic potential to these electrodes, a force is produced in the direction of one electrode provided that electrode is of such configuration to cause the lines-of-force to converge steeply upon the other electrode. The force, therefore, is in a direction from the region of high flux density toward the region of low flux density, generally in the di-

rection through the axis of the electrodes. The thrust produced by such a device is present if the electrostatic field gradient between the two electrodes is non-linear. This non-linearity of gradient may result from a difference in the configuration of the electrodes, from the electrical potential and/or polarity of adjacent bodies, from the shape of the dielectric member, from a gradient in the density, electric conductivity, electric permittivity and magnetic permeability of the dielectric member or a combination of these factors. A basic device for producing force by means of electrodes attached to a dielectric member is disclosed in my Patent 1,974,483. In one embodiment disclosed in my patent, an electrostatic motor comprises devices having a number of radially directed fins extended from one end of the dielectric body and a point electrode on the opposite end of the dielectric body. When this device is supported in a fluid medium, such as air, and a high electrostatic potential is applied between the two electrodes, a thrust is produced in the direction of the end to which the fins are attached. Other electrostatic devices for producing thrust are disclosed and described in detail in my British Patent 300,- 311, issued August 15, 1927. Recent investigations in electrostatic propulsion have led to the discovery of improved devices for producing thrust by the use of electrical vectorial forces. Accordingly, it is the primary object of this invention to provide an improved electrical device for producing thrust. It is another object of this invention to provide a device for producing modulated thrust in response to varying electrical signals, which device produces a greater effect than the prior type devices mentioned above. It is another object of this invention to provide a device which shapes or concentrates electrostatic flux to

produce an improved thrust. Broadly, the invention relates to shaping an electrical field to produce a force upon the device that shapes the field. The electrical field is shaped by the use of an electrode of special configuration whereby the electric lines-of-force are made to converge at a distance from the electrode. One illustrative embodiment of this invention which satisfies the above requirement is an arcuate surface, or alternatively, a system of wires, tubes or plates embedded in a dielectric surface and forming a directive array. One such highly-charged electrode electrode acting within and upon an ambient of differential electrical potential will move in response to the forces created by shaping of the electrostatic field. If a smaller electrode is added at or near the focus of the field-shaping electrode, and mechanically attached to that electrode, both electrodes as a system will move in a direction of the larger or field-shaping electrode. As is mentioned above, the field-shaping electrode alone, when charged with respect to its electric ambient, will move or possess a force in the direction of its apex. If another electrode carrying a different charge is added at or near the focal point of the field-shaping electrode, then the field becomes more concentrated, i.e. shaped to a greater degree and the resulting thrust is greater than that which exists when the field-shaping electrode alone is employed. Briefly in accordance with aspects of this invention, an electrode is connected on each end of a dielectric member and one of the electrodes defines a large area flat or preferably arcuate surface which is curved in such a direction to produce, usually in co-operation with the other electrode, a shaped electrostatic field. Advantageously, if the arcuate electrode is in the form of a parabola or hyperbola, the length of the



dielectric member may be such that the other electrode is located in the region of the focus of the parabola or hyperbola, as the case may be. If the arcuate electrode is hemispherical, the other electrode is located near the center of the hemi-sphere. In accordance with other aspects of this invention the dielectric member supporting the two electrodes may have electrical conductivity and/or dielectric constant which varies progressively between its ends so that the dielectric member contributes to the non linearity of the field gradient and causes a greater thrust to be developed. In accordance with still other aspects of this invention, an annular electrode member is secured to an electrode mounted in the region of the axis of the annular electrode. If the second electrode is located at the center of the annular electrode and the two electrodes are energized, such force is not detected. However, if the second or innermost electrode is displaced from the center of the annular electrode in the region of the axis of the annular electrode and the electrodes are energized, then thrust will be produced by the two electrodes. The annular electrode may either be a flat ring, a toroid, or a section of a cylinder. In accordance with still other aspects of this invention, tapered dielectric members having electrodes secured to opposite edges thereof may be employed to produce a thrust in response to the application of potentials to these electrodes. The thrust produced by these tapered dielectric members maybe further augmented by embedding massive particles, such as lead oxide, in the wedges, which particles are usually more concentrated near the points of the wedges. Accordingly, it is a feature of this invention to provide an electrical device for producing thrust which includes a dielectric member and electrodes supported at each end of the dielectric

member, one of which electrodes is located in the region of the focal point of the arc of the arcuate surface electrode. It is another feature of this invention to provide a device for producing thrust having a dielectric member and a pair of electrodes secured to opposite ends of the dielectric rod or member, one of which electrodes defines a parabolic or hyperbolic surface, the other electrode being located in the region of the focus of said surface. It is another feature of this invention to employ an insulating rod or member between the two electrodes, which rod or member has a varying dielectric constant, said dielectric constant progressively increasing or decreasing along the length of the dielectric member. It is still another feature of this invention to employ a rod or member connected between the two electrodes across which an electrostatic potential is applied, which rod or member has a varying electrical conductivity, said conductivity progressively increasing or decreasing along the length of the dielectric member. It is another feature of this invention to employ a single electrode having an arcuate surface and to connect a source of potential to the arcuate surface which is opposite in polarity to the potential of the masses comprising the environment of the arcuate surface. It is still a further feature of this invention to employ an arcuate electrode as a device for producing thrust and to apply a varying electrical signal to the arcuate electrode. It is still another feature of this invention to employ a wedge of dielectric material having electrodes on opposite ends thereof to produce a thrust in response to the application of electrical potentials. It is still a further feature of this invention to employ a tapered dielectric material having massive particles embedded therein to produce a thrust in response to the application of potentials to the electrodes

secured to the dielectric member. It is still a further feature of this invention to employ an annular ring electrode and a second electrode secured to the annular electrode in the region of the axis of the annular electrode in the region of the axis of the annular electrode to produce a thrust in response to the application of electrical potentials thereto. These and other various objects and features of this invention will be apparent from consideration of the following description when read in connection with the accompanying drawing wherein: FIGURE 1 is a view in elevation of one illustrative embodiment of this invention; FIGURE 2 is a view in elevation, partly in section, of another illustrative embodiment of this invention; FIGURE 3 is a graphical representation of the field gradient between the electrodes of one illustrative example of this invention in which distance from one electrode is plotted as the abscissa whereas flux density is plotted as the ordinate; FIGURE 4 is a perspective view of another illustrative embodiment of this invention; FIGURE 5 and 6 are perspective views of still another illustrative embodiment of this invention; FIGURE 7 is an end view of another illustrative embodiment of this invention employing a pair of devices of the type disclosed in FIGURE 6, which devices are mounted and serially connected in a single array; FIGURES 8A, 8B and 8C are views in elevation, partly in section, of still other illustrative embodiments of this invention. Referring to FIGURE 1, there is depicted an insulating member 10 having an arcuate electrode 12 mounted on one end thereof and a second electrode 14 mounted on the opposite end thereof. A source of direct current voltage 13 is connected to electrodes 12 and 14 through conductors 15 and 17, respectively. I have discovered that if two electrodes are mounted on opposite end of a dielectric member,


and a field emanates from these electrodes which produces a linear gradient through the dielectric member as shown by the dotted line 30 of FIGURE 3, then no thrust is produced by the dielectric member. However, if the field is distorted to produce a non-linear gradient such as graphically represented by line 32 in FIGURE 3, then a thrust will be produced, which thrust will be related to the degree of non-linearity of the field gradient. One way to produce a gradient which varies non linearly is to shape one of the electrodes in a form of an arcuate surface such as 12. However, numerous other ways to influence the field gradient will be disclosed below. Electrode 14 represents a substantial mass and it has been found that best results are obtained if the surface area of electrode 14 is greater than the surface area of the end of rod 10. In one particular example, a spherical electrode having a diameter greater than the diameter of rod 10, produced very satisfactory results. Advantageously, the dielectric member 10, may be employed to increase the non-linearity of the field gradient. For example, the dielectric member may be of material having a uniform relative dielectric constant and be tapered in the direction of electrode 14 such that the member 10 in the region of electrode 12 has a much greater cross-sectional area than the end of member 10 which is connected to electrode 14. An equivalent result may be obtained if the member 10 is of uniform diameter but has a dielectric of graduated density or which comprises a material having a progressively different electrical conductivity or dielectric constant, or alternatively the electrical conductivity, varies from a low value in the region of electrode 14 to a high value in the region of electrode 12. The arcuate electrode 12 may be either a stitched wire surface or a solid conducting surface. In the case of

stitched wire surface, the wires are very close together so that when an electrical potential is applied to these wires, they act substantially in the same manner as a conductive surface. Arcuate electrode 12 will produce a thrust when a potential is applied to the electrode 12 which is opposite in polarity to the potential of the bodies in the region of electrode 12. Such a thrust will be produced even though the dielectric member 10 and the electrode 14 are eliminated from the structure. However, the thrust produced by the charged arcuate electrode 12 when actuating alone is less than the thrust produced by the combined device, that is, employing the dielectric member 10 and the oppositely charged electrode 14. Referring now to FIGURE 2, there is depicted another illustrative embodiment of this invention in which field-shaping is accomplished. In the embodiment of FIGURE 2, the planar electrode 18 is connected to a hemispherical electrode 14 by means of a dielectric rod 10. When a source of electrical potential (not shown) is connected through wires 15 and 17 to electrodes 18 and 14, respectively, a field gradient will be produced between electrodes 18 and 14, which field gradient varies in accordance with the graph represented by the solid line 32 of FIGURE 3. In this particular embodiment, as well as in the embodiment of FIGURE 1, the non-linearity of the field gradient is further augmented by the use of a connecting rod 10 which is a dielectric with progressively different dielectric constant between electrodes 18 and 14. A similar result may be produced by the use of a rod 10 having electrical conductivity which varies progressively between electrodes 18 and 14. Referring now to FIGURE 4 there is depicted still another illustrative embodiment of this invention in which a thrust is produced in response to the

application of electrical potentials. A frusto-conical surface 25 comprising a metal or having a metal surface to be used on an electrode is connected to a tapered member 27. The tapered member 27 is frusto-conical and is primarily of non-conductive material but contains granules of semi-conducting material, which granules are concentrated near the tip 28. Mounted on tip 28 is a half-wave radiator 29 which may be in the form of a disk. It is noted that the axis of member 27 coincides with the axis of member 25. When a source of potential is connected to electrodes 25 and 29, a thrust is produced in the direction of the arrow 31 regardless of the polarity of the applied voltage. However, a greater thrust is produced if the electrode 25 is positive with respect to electrode 29. Alternating current voltages may also be applied to electrodes 25 and 29 and the potential may be either superimposed upon or substituted for the direct current voltages. Preferably, the frequency of the applied A.C. voltage is such that the diameter of the disk 29 constitutes a half-wavelength of the applied voltage. Referring now to FIGURE 5 there is disclosed a tapered member 32 which is non-conductive material and may contain particles of semi-conducting material in a manner similar to member 27. The semi-conducting material contained in member 32 and in member 27 may be any convenient form of massive particles such as lead oxide. Along one surface of member 32 is an electrode 34 while along the opposite surface is another electrode 36. When a potential is applied to these electrodes, preferably of a polarity such that electrode 36 is positive with respect to electrode 34, a thrust is produced in the direction of the arrow 37. In the devices disclosed in both of FIGURES 4 and 5, the thrust produced by the electrodes is augmented by the varying cross-sectional area

of the non-conductive member connecting the electrodes and is further augmented by the voltage gradient produced by the embedded particles, which voltage gradient is greater than that which would be introduced by a tapered non-conductive member without embedded particles. Referring now to FIGURE 6 there is depicted a bank of members 32 such as disclosed in FIGURE 5 in which like electrodes 36A through 36E are secured together by a connector in any convenient form, such as plate 38. Each of these members 32A through 32E produces a thrust in the direction of the arrow 37A and the resultant force is equal to the sum of the thrust produced by the individual members 32 in response to the application of potentials to the electrodes 34A-34E and 36. In FIGURE 7 there is depicted a pair of banks of members, such as depicted in FIGURE 6, in which the electrodes are serially connected. In this particular instance, a plate or other member 40 comprises an electrode on which are mounted an array of members 32F through 32J. A second electrode 42 is secured between electrodes 32F through 32J and electrodes 32K through 32P. A third electrode 44 is connected to the electrode 34 on each of members 32K through 32P. It is to be noted that electrodes 40 and 44 are connected to a source of one potential while electrode 42 is connected to a source of the opposite potential. The thrust produced by this array is in the direction of arrow 37B and the manner in which this thrust is produced is similar to that explained in connection with FIGURES 5 and 6, although it would appear that electrode 42 will experience a natural attraction for electrodes 40 and 44. A non-linear field gradient is produced between these electrodes by the varying cross-sectional area of members 32 and by the presence of semi-conducting particles in

members 32. This non-linear field gradient gives rise to the thrust, as mentioned above. Referring now to FIGURES 8A, 8B and 8C there is depicted other illustrative embodiments of this invention. In FIGURE 8A a toroid member 43 has an electrode 50 supported at its center by means of insulating rod 52. If the electrode 50 and the toroid member 43 are both conducting surfaces defining electrodes and these electrodes are connected to sources of opposite potential, no thrust will be developed by the device. If, however, as depicted in FIGURE 8B electrode 50 is translated along the axis of generation of toroid or annular member 48 and again supported by non-conductive members 52, this device will experience a downward thrust, as indicated by arrow 53, in response to the application of potentials of either polarity. It is believed that this force is produced by the annular configuration of electrode 48 and the off central location of electrode 50. In the instance of FIGURE 8C, electrode 50 is positioned beneath the center of electrode 48 and positioned on the axis of generation of electrode 48. When potentials are applied to electrodes 48 and 50 in FIGURE 8C, a thrust is produced in an upward direction, as indicated by arrow 54. Here again the field gradient is produced by the configuration of electrode 48 and the location of electrode 50 with respect to electrode 48. From the foregoing discussion, it is also apparent that a combination of a curved electrode, a supporting member of varying cross-sectional area, and a second electrode supported by the connecting member will produce a thrust along the axis of the curved electrode when potentials are applied to the electrodes. Similarly, a thrust may be developed between plane electrodes of unequal areas which are connected by a member of varying cross-



sectional area. The thrust developed by this last mentioned device is further increased by the introduction of semi-conductive particles in the non-conducting member, which particles are more concentrated in the region of the smaller electrode than in the region of the larger electrode. Further, these tapered members having planar electrodes connected to opposite surfaces may be stacked in vertical arrays and connected in parallel, or they may be stacked in vertical arrays connected in series with similar vertical arrays. In applying potentials to these various embodiments, it has been found that the rate at which the potential is applied often influences the thrust. This is especially true where the dielectric members of high dielectric constant are used and the charging time is a factor. In such cases, the field gradient changes as the charge is built up. In such cases where initial charging currents are high, dielectric materials of high magnetic permeability likewise exhibit thrust with time. One advantageous manner of applying potential is that of employing potentials which vary cyclically. It is thus apparent that one embodiment of this invention embodies a pair of electrodes mounted on an insulating member, one of which electrodes defines an arcuate surface to produce an improved thrust in response to the application of direct current potentials. It is also apparent that this thrust is augmented by increasing the non-linearity of the field gradient by a progressively-changing characteristic of the dielectric member connecting these electrodes. This non-linearity of field may be produced by a gradient in electric conductivity, electric permittivity and/or magnetic permeability along the length of the member, or it may result from a change in the cross-sectional area of the rod which rod has otherwise uniform characteristics. While I have

shown and described various embodiments of my invention, it is understood that the principles thereof may be extended to many and varied types of machines and apparatus. The invention therefore is not to be limited to the details illustrated and described herein. I claim: 1. A device for producing thrust comprising a field shaping surface formed of stitched, closely spaced conductors and having a dielectric material therebetween to define a smooth surface, a dielectric member connected to said field shaping surface and an electrode on the end of said dielectric member remote from said field shaping surface and means for applying electrical potential between said electrode and said closely spaced conductors. 2. A device for producing thrust in accordance with claim 1 wherein said dielectric member has a dielectric constant which varies progressively between said electrode and said surface means. 3. A device for producing thrust comprising an electrode having a relatively large surface area, an electrode 7 positioned in the region of the axis of generation of said surface and having a relatively small surface area, dielectric means connecting said electrodes and means for applying a varying electrical potential to said electrodes. 4. A device in accordance with claim 3 wherein said dielectric means exhibits a dielectric constant which varies progressively from a relatively high value in the region of the large electrode to a relatively low value in the region of said small electrode. 5. A device in accordance with claim 3 wherein said dielectric means has an electrical conductivity which varies progressively between said electrodes. 6. A device for producing thrust comprising a planar electrode, a second electrode positioned in the region of the axis of generation of said planar electrode and having a surface area smaller than the surface area of said planar

electrode, a dielectric member connecting said electrodes and means for applying a high electrostatic potential to said electrodes. 7. A device in accordance with claim 6 wherein said dielectric member is tapered from the planar electrode towards the smaller electrode. 8. A device in accordance with claim 6 wherein said dielectric member has a conductivity which varies progressively from a relatively high value near the planar electrode to a relatively low value near the smaller electrode. 9. A device for producing thrust in response to the application of electrical potentials to the electrodes thereof comprising a first electrode, a second electrode having a relatively large planar surface area with respect to said first electrode and means including a connecting member supporting said electrodes in spaced relationship for producing a varying field gradient between said electrodes. 10. A device in accordance with claim 9 wherein said connecting member has a varying cross-section. 11. A device in accordance with claim 9 wherein said connecting member tapers between said electrodes. 12. A device in accordance with claim 9 wherein said first and second electrodes are flat electrodes of unequal area. 13. A device according to claim 9 including means for applying a varying electrical potential to said electrodes. 14. A device in accordance with claim 9 wherein said connecting member has a dielectric constant which varies between electrodes. 15. A device in accordance with claim 14 wherein said first electrode is a frusto-conical surface and wherein said connecting member extends along the axis of generation of said first electrode. 16. A device in accordance with claim 14 wherein said first electrode defines a frusto-conical surface. 17. A device in accordance with claim 9 wherein said connecting member comprises semi-

conducting particles whereby said connecting member is given a conductivity gradient. 18. A device in accordance with claim 15 wherein said second electrode is a disk-shaped radiator and wherein the potentials applied to said electrodes are alternating current potentials, the diameter of said disk-shaped electrode being equal to a half-wave length of the alternating current potential. 19. A device in accordance with claim 15 wherein said connecting member contains semiconducting particles which are more concentrated in the region of the disk radiator than in the region adjacent said first electrode. 20. A device for producing thrust in response to the application of electrical potentials to the electrodes thereof comprising an annular electrode, a second electrode, and insulating means connecting said electrodes whereby thrust is produced along the axis of generation of said annular electrode in response to the application of electrical potentials thereto. 21. A device in accordance with claim 20 wherein said annular electrode comprises a toroidal surface. 22. A device in accordance with claim 20 wherein said second electrode is mounted on the axis of generation of said annular electrode. 23. A device in accordance with claim 22 wherein said second electrode is displaced from the center of said annular electrode whereby a thrust is developed along said axis in a direction from said second electrode towards that annular electrode in response to the application of electrical potentials thereto. References Cited by the Examiner

United States Patents 1,974,483 9/34
Brown _____ 310-5

Foreign Patents 1,003,484 11/51 France
Milton O. Hirshfield, Primary Examiner.
Oris L. Rader, David X. Sliney,

Examiners. Research and Preliminary Engineering Space Vehicle Program
Subject: Research and Preliminary Engineering for Space Vehicle Program
Part I - Propulsion Part II - Navigational and Flight Control Instruments Part III - Communication and Remote Control Systems Part IV - Materials of Construction

Synopsis

This proposal recommends a long-range coordinated program for the research and preliminary engineering leading to the construction of prototype space vehicles. The proposal stresses the importance of research on gravitation and on the relationships between electrodynamics and gravitation. It calls attention to certain possibilities inherent in such a program of development. The program would be divided and conducted in steps, as follows: (a) Preliminary investigations of physical methods, (b) Engineering development, (c) Advance design, (d) Construction of operating prototypes.

Scope Of Interest

(a) Propulsion The program anticipates the use of electric methods of propulsion and control, and specifically excludes the consideration of jet or rocket propulsion devices, per se, except as a means for generating the required high electrical potential. Emphasis is placed on studies of the physical relationships between electrodynamics and gravitation leading to the development of methods of electrogravitic propulsion and steering control. (b) Navigational and Flight Control Instruments Problems of space navigation are to be examined, particularly with the view of applying electrogravitic principles in the design of flight

instruments to indicate: 1. Gravitational Vector (insensitive to acceleration), 2. Acceleration (insensitive to gravity), 3. Gravitational Gradient (insensitive to acceleration), 4. Gravitational Potential of Space, 5. Electrical Potential of Space, 6. Space Speed (absolute ether drift). (c) Communication and Remote Control Systems Applications of electrogravitic induction to communications and remote control are to be developed. Use of gravitational radiation is the objective. Basic tests with electrically-shielded capacitors and massive high-K dielectrics are proposed. Methods are to be extended into full-scale communication systems. Such systems, while similar to electromagnetic (radio) systems may be found to offer many advantages - such as higher penetrability, elimination, elimination of "shaded" areas, higher velocity of wave propagation and a wholly new spectrum of channels. (d) Materials of Construction Continuation of the research of the late Charles Francis Brush on the "non-equivalence of mass and weight" is recommended (see appendix). Further confirmation of the Brush findings may be provided by the existence of gravitational isotopes, as distinguished from mass isotopes. Procedures for isolating gravitational isotopes in common aircraft metals, with the object of creating super-light alloys. Methods of beneficiation are suggested for enriching the content of lighter gravitational isotopes in common aircraft metals, with the object of creating super-light alloys. The spontaneous evolution of heat, observed by Brush and Harrington, appears to be one of the characteristics of lighter gravitational isotopes, and may serve as a tracer in the steps of beneficiation. Studies are proposed to determine the source of the energy and to investigate possible uses of said heat. This program is to include a study of the



rare earth metals and their alloys, and also the metal tantalum, regarded as potential aircraft materials. In nature, most of the rare earth metals (and tantalum) indicate mean values of specific gravity having large negative anomalies, and this property makes them interesting as probable rich sources of gravitational isotopes required in the manufacture of super-light materials of construction.

Program Of Research And Development

Group A - Electrodynamic- Gravitational Field Relationships

Purpose: Generation of quasi-gravitation by electrical means, quantitative measurements and derivation of equations.
Abstract: One of the basic relationships between the electrodynamic field and the gravitational field appears to be revealed "during the process of charging or discharging electric capacitors".
Proposal: Confirming experiments are proposed in which two or more large high-voltage capacitors are associated spatially with a short-period geophysical gravimeter. Careful observations are to be made of the momentary gravitational anomalies induced in the region which accompany the change in electrical state. Studies are proposed of the effects of varying total capacitance, rate-of-change of electric charge, mass of dielectric materials, specific inductive capacity of said materials and whether the effects are vector or scalar. These investigations shall be directed toward the derivation of a satisfactory mathematical expression including all of the above factors.

Group B - Propagation Of Gravitational Waves

Purpose: Transmission and reception of electro-gravitational waves for purposes of communication and remote control.

Abstracts: Preliminary experiments have indicated the existence of an inductive inter-action between two independent shielded capacitors. In these experiments, a discharging capacitor induces a voltage in an adjacent capacitor, and the effect appears to penetrate electromagnetic shielding. Theoretically, this effect of one capacitor upon another appears to be electrogravitic in nature and constitutes evidence of a new type of wave propagation. It is believed that this form of inductive transmission may eventually be utilized in a completely new method of wireless communication. Proposal: It is proposed that progressively larger-scale and longer-range transmissions be conducted. Beginning with untuned systems, laboratory tests are proposed to explore the basic electrogravitic relationships between simple systems of capacitors. Then, progressing to tuned systems, and pulsed (radar) applications, large- scale out-of-door demonstrations are suggested. Such demonstrations shall be conducted between suitably protected transmitting and receiving vaults (preferably underground) which are thoroughly shielded against electromagnetic radiation. Appropriate studies of wave attenuation due to transmission through large masses of earth may then be undertaken. Similar studies of wave attenuation in sea water are also prosed. These studies are to be supported by fundamental research on the nature of electrogravitic induction (See appendix for outline).

Group C - Ponderomotive Forces In Solid Dielectrics

Purpose: Isolation and measurement of electrogravitic forces in solid dielectrics.
Abstract: Investigations conducted by Biefeld and Brown point to the existence of a hitherto unrecognized ponderomotive

force in all ferroelectrics under changing electric stress. This force appears to be a function of the specific inductive capacitance and the mass of the dielectric material, as well as high voltage and current factors. Recent availability of the massive barium titanate (high-K) dielectrics and other dielectrics of this class give promise of developing these forces to the point where they may become of practical importance in specific propulsion applications. Proposal: A survey of dielectric materials revealing this effect is proposed. Beginning with a critical analysis, using the Townsend Brown Differential Electrometer (an instrument developed at the Naval Research Laboratory and at the University of Pennsylvania), studies are proposed of the forces developed in mica, glass, marble, phenolics and dielectrics in general and then, in particular, the newer ceramic dielectrics. This work is to be augmented by basic determinations of the Biefeld-Brown effect in vacuum. (See appendix). It is proposed that, after suitably active materials are selected, scale models of other rotary and linear "motors" be constructed and tested. With the necessary engineering data then at hand, a motor to weigh approximately 500 lbs. may be constructed to propel a model ship. This is proposed as a practical demonstration of one of the forms of electrogravitic drive. Ether drift and space-couple observations, including specifically a repetition of the classic Trouton-Noble experiment (but using massive dielectrics) are suggested as being of interest not only for their contribution to basic knowledge of the nature of space but as bearing upon the principle of operation of space speed indicators (See appendix). Low temperature experiments in physics of the solid state (using the liquid-helium cryostat) are highly recommended but are

expensive. These experiments, however, may be so designed as to provide answers to many questions relative to the fundamental nature of gravitation. They are to embrace such subjects as the "Anomalous Mass of the Electron in Metals" and the "Behavior of Super-Cooled Dielectrics". The availability of the liquid-helium cryostat would enable the project to engage continuously in low-temperature work which could contribute enormously to our knowledge of solid state physics.

Group D - Reactive Forces In Fluid Dielectrics

Purpose: Development of high speed electrokinetic propulsive systems for spacecraft. Abstract: Studies of boundary forces (where electrodes are in contact with fluid dielectrics) reveal the existence of a "complex" of inter-acting forces, some of which are purely electrostatic, some electromagnetic and some which could be electrogravitic. The tentative theory requires these electrogravitic forces to be present whenever a mass of dielectric material is charged and moving, and to increase in proportion to the volume of the fluid which is charged and moved. Hence, it is, in a sense, the juxtaposition of the elements of the static form of the capacitor described in Group C experiments, and provides what may be described as an electrokinetic propulsive system, with direct applications to high speed aircraft and spacecraft. Proposal: It is proposed that electrically-charged circular airfoils be mathematically analyzed and improved. Starting with 2 ft. discs at 50 KV, the steps of the development should include 4 ft. discs at 150 KV and a 10 ft. disc at 500 KV. Careful measurements are to be made of both static and dynamic thrust. Studies are also proposed wherein the discs are adapted for vertical lift (levitation) as well

as for horizontal thrust and this feature may be incorporated in the design of the 10 ft. experimental model. It is proposed that studies likewise be made of various methods for obtaining the required high voltages, and these studies should include the development and evaluation of the capacitor voltage multiplier and the "flame-jet" electrostatic generator to provide up to 15 million volts (See appendix). This work is to be augmented by the engineering studies on the relative efficiency of propulsion of electrified discs in air at reduced pressure or in vacuum and at various voltages.

Group E - The Spontaneous Generation Of Heat In Certain Complex Silicates, Lavas And Clays.

Purpose: To establish the existence of the positive effects observed by Brush and Harrington, determine the origin of the energy represented and extend the observations into the rare earth /and other/ elements. Abstract: The discovery by Charles Francis Brush of an unexplained heating effect in certain materials is strikingly reminiscent of the discovery of radioactivity by Henri Becquerel which led to the isolation of radium by the Curies. Recent studies have indicated a certain parallelism between the radioactive elements and the rare earth elements, revealing the possibility of a spontaneous release of energy (in this instance in the form of heat alone) by the rare earth elements. Thus, the newly discovered "thermoactivity" may bear the same relationship with the rare earth elements as "radioactivity" bears with the radioactive elements. In all probability the source of the energy represented in thermoactivity will be found in the complex unstable electronic shells of the rare earth atoms, and not in the nuclei as in radioactivity.

The anomalous gravitational properties of the rare earth elements and their widespread but tenuous occurrence in nature point to other parallels with the radioactive elements such as decay of activity and critical limits of mass. Proposal: 1. Conduct an organized examination of materials (complex silicates, lavas, and clays) known to exhibit spontaneous heating. 2. Conduct a field search for additional materials. 3. Attempt to isolate and purify materials showing spontaneous heating. 4. Extend the examinations into the rare earth metals and tantalum. 5. Conduct mathematical and theoretical studies of gravitational isotopes as distinguished from mass isotopes. 6. Study the mechanism of spontaneous heat generation and the decay of the effect. 7. Study the effects of ionization, electric and magnetic fields upon the effect. 8. Study methods of beneficiation of materials for intensification of the heating effect.

Group F - Navigational And Flight Control Instruments

1. Gravitational Vector (stable vertical) 2. Accelerometer (inertial gradient) 3. Gravitational Gradient (gravity) Purpose: Engineering development and design, adaptation for operation in conjunction with servo mechanisms for actuating flight control devices. Abstract: The physical principles underlying the operation of these instruments have evolved from the findings of C. F. Brush (see appendix - references). Quite recently, careful studies of the records of the Brush experiments have been made. The conclusions, while still highly controversial, leave no doubt that the Brush concept, if supported, can provide answers to many difficult and, as yet unsolved, problems in navigational and flight control instrumentation. The Brush findings may be expressed simply: "the ratio of mass to weight is not the same for

all kinds of matter, as has been supposed, and the mass-weight ratio is not constant even in the same kind of matter". Based on the hypothesis of the non-equivalence of mass and weight, the principles of three instruments have evolved as follows: 1. Gravitational vector (stable vertical) Two equal (inertial) masses of unequal weight are utilized in a balanced pivoted device. Such a device may be said to be inertially symmetrical and gravitationally asymmetrical. The pivoted system will orient itself to the vector of gravity and yet remain insensitive to the inertial effects of acceleration and centrifugal force. 2. Accelerometer (inertial gradient) Two unequal (inertial) masses of equal weight disposed in a pivoted device as above, but with spiral spring (or the like) tending to restore the movable system to zero position. The pivoted element responds quantitatively to inertial effects of acceleration and centrifugal force but is insensitive to gravity. 3. Gravitational Gradient (gravity) Such an instrument is the reverse of the accelerometer above and structurally resembles the gravitational vector indicator except that it includes a spring which is adjusted to restore the indication to zero in the absence of a gravitational field. The movable element responds quantitatively to gravity (g) only and is completely insensitive to all inertial effects such as acceleration or centrifugal force. Proposal: It is proposed that operating models of the three types of instruments be constructed and tested on a centrifugal carriage. Materials for the dipoles of the pivoted systems are to be selected from beneficiated gravitational isotopes (both light and heavy) developed under Group E. These prototype instruments are to be adaptable for operation in conjunction with telemetric circuits or servo mechanisms for actuating flight control devices.

IMPLEMENTATION (a) Laboratory facilities To be supplied directly by the parent company or its subsidiaries. (b) Personnel Members of the regular research and engineering staff of the parent company and its subsidiaries, with services of such technical or operational consultants as may be necessary. (c) Sub-contracts To be awarded only when it is impractical or uneconomical to perform the work with company facilities. CONDENSED SCHEDULE AND ESTIMATED COST 1955 - October through December (3 months only) Organizational work and planning (\$ 3,000.00) Establishment of technical reference library on gravitation and allied subjects (\$ 5,000.00) Preliminary beneficiation of gravitational isotopes (\$ 12,500.00) Engineering design of flight control instruments (\$ 12,500.00) Quantitative tests of Biefeld_Brown effects in solid dielectrics (\$ 9,500.00) Basic tests on propagation of gravitational waves for communication purposes (\$15,000.00) Total (\$ 57,500.00) 1956 - 1958 (3 years) Detailed schedule and expenditures to be determined during the preliminary organizational and planning stage. Respectfully submitted. T. Townsend Brown Consultant Washington D. C. December 1, 1955 Condensed Outlines of Proposed Fundamental Research SECTION A - ANOMALOUS BEHAVIOR OF MASSIVE HIGH-K DIELECTRICS: I - General Description. II - Investigation of the Biefeld-Brown Effect (a) Basic pendulum experiment 1. Effects of mass 2. Effects of K in fluid media 3. High vacuum tests 4. Mathematical relationships involved, derivation of equation (b) Series-capacitor experiments: 1. Effects of mass 2. Effects of K in internal dielectric 3. Effects of K in external fluid dielectric 4. External electrostatic forces 5. Effects of shielding (c) Variations of ponderomotive forces

with time: 1. Studies of possible causes 2. Design of continuous recording instrument III - Variations of K with electrostatic potential and/or gravitational potential (a) Studies of hypothesis; mathematical relationships involved. (b) Implications of effect of electrostatic potential and gravitational potential upon: 1. speed of light 2. rates of nuclear fission (c) Measurement of change in potential difference in capacitors with change in electrostatic potential and/or gravitational potential IV - Studies of massive high-K dielectrics (a) Theory of dielectrics, sources of polarization (b) Detailed examination of titanium oxide, barium titanate, lithium thallium tartrate - looking toward increasing values of K (c) Measurement of ponderomotive forces developed by series capacitors containing high-K dielectrics: 1. With applied potential in steady state 2. With changing potential 3. Effects of varying rate-of-change (d) Potential differences developed in polarized materials with change in overall electrostatic potential 1. Effects of mass 2. Effects of K (e) Stability of electrets: 1. Anomalous rates of voltage decay 2. Diurnal variations V - Analysis of Electrified Disc-airfoils (a) Theoretical considerations (b) Thrust measurements: 1. In air at reduced pressure 2. In hard vacuum 3. In fluid dielectrics of various K 4. Effects of viscosity of fluid dielectrics SECTION B - ELECTROGRAVITIC INDUCTIVE EFFECTS: I - General Description II - Investigation of the Fernando Sanford Effect (a) Repeat experiments (b) Series-capacitor experiments (c) Theoretical considerations III - Studies of Potential Variations (a) In large insulated masses (b) Effects of mass (c) In capacitors (d) Electrical potential vs.

gravitational potential (e) The mountain effect (f) Centrifugal potential effects IV - Studies of Voltage Gradients in Dielectric Materials (a) Long-series capacitors (b) Effects of mass (c) Directional gravitational effects V - Polar Capacitors (a) Shift of potential of the mid-point (b) Directional effects (c) Effects of elevation from earth (d) Tests below earth surface VI - Short-period Gravity Meter (a) Design and Construction (b) Detection of gravitational waves REFERENCE: Terrestrial Electricity, Fernando Sanford, Professor Emeritus of Physics - Stanford University. Stanford University Press. SECTION C - DETERMINATIONS OF TIDAL EFFECTS ON BROWN DIFFERENTIAL ELECTROMETER I - General Description II - Analysis of Zanesville and Philadelphia observations (a) Solar component (b) Sidereal component (c) Lunar components, correlations with: 1. Lunar hour angle 2. Angular distance, phase 3. Distance from earth 4. Altitude of moon 5. Right ascension of moon (d) Studies of combined effects (e) Detailed comparison of Zanesville and Philadelphia observations (f) Comparison with simple tidal curves: 1. Ocean tides 2. Atmospheric tides, barometric (Maris effect) (g) Secular changes (h) Correlation with other natural variables III - Analysis of California observations (a) Secular changes (b) Solar, lunar and Sidereal components (c) 75th meridian and 120th meridian observations (d) Regional vs world-wide variations (e) Local variations (f) Correlation with other factors IV - Analysis of Fernando Sanford Records (a) Solar, lunar and sidereal components (b) Comparison with Zanesville and Philadelphia records (c) Comparison with atmospheric electric gradient and earth current records V - Analysis of Section A electrometer observations (Current Program) (a) Studies

related to gravitational and electrical variables (b) Secular changes (c) Comparison with former records VI - Analysis of Section D thermoactivity observations (Current Program) (a) Diurnal variations (b) Secular changes (c) Comparison with Section V records SECTION D - GRAVITATIONAL ISOTOPES I - Investigation of the Charles Francis Brush Effects (a) Impairment of gravitational acceleration: 1. In complex silicates, lavas and clays 2. In barium aluminate, barium titanate and other high-K materials (b) Persistent generation of heat: 1. Calorimetric analysis 2. Mass effect, particle size 3. Diurnal variations (c) Correlations between (a) and (b) II - Studies of Gravitational Isotopes as Distinguished from Mass Isotopes (a) Definitions (b) Theoretical considerations (c) Gravitational periodic table of the elements: 1. Specific gravities with positive and negative anomalies 2. The Lanthanide contraction 3. Parallels between the lanthanide and actinide series of elements (d) Spontaneous evolution of heat: 1. Parallels between thermoactivity and radioactivity 2. Decay of thermoactivity, increase of gravitational mass, computations of half-life 3. Exponential increase in thermoactivity with total (localized) mass 4. Possibilities of "critical mass" effects (e) Determination of origin of energy: 1. Unstable electron shells 2. Dirac "holes" 3. Possibilities of negative mass. Lofting properties 4. Effects of electric and magnetic fields 5. Effects of changes in electric and/or gravitational potential 6. Diurnal and secular variations in thermoactivity and/or weight III - Beneficiation of Gravitational Isotopes (a) Occurrence in nature: 1. In all elements 2. In rare-earth elements 3. In special cases (b) Nascent gravitational isotopes: 1. Enrichment following chemical or nuclear reactions 2. Presence

in reaction products of nuclear reactors 3. Breeder technique (c) Methods of beneficiation: 1. Settling and centrifuging 2. Settling and thermal diffusion IV - Possible Uses of Gravitational Isotopes (a) Super-light (and super-heavy) fractions for: 1. Materials of construction (alloys) 2. Sensitive elements of navigational instruments (b) Contra-terrene (negative gravitational mass) possibilities as (fixed lift) lofting agents: 1. Materials of construction for aircraft and spacecraft 2. Lofting "capsules" (c) As a source of heat 1. Building materials (and the like) where slight warming effect is desired 2. Steam generation (similar to but less energetic than nuclear fuels) 3. Explosives REFERENCES: Brush, C.F., Physical Review, 31, p 1113(A), 32, p 633 abstract; Proc. Amer. Philosophical Soc. Vol.IX No. 2, 1921; Vol. LXVII No. 2, 1928; Vol LXVIII No. 1, 1929; Journal of Franklin Inst., Vol. 206, No. 1, 1928. Harrington, E.A., Nat'l Bu. of Standards, Proc. Amer. Philosophical Soc., Vol. LXXII, No. 5, 1933.


Gravitational Periodic Table Of The Elements indicating parallel relationships 1 Hydrogen GROUP I GROUP II 2 Helium 10 Neon 3 Lithium
 11 Sodium 4 Beryllium
 12 Magnesium 5 Boron 13
 Aluminum 6 Carbon 14
 Silicon 7 Nitrogen 15
 Phosphorus 8 Oxygen 16
 Sulfur 9 Flourine 17
 Chlorine GROUP III GROUP IV 18 Argon 36 Krypton 19 Potassium.
 37 Rubidium 20 Calcium
 38 Strontium 21 Scandium
 39 Yttrium 22 Titanium
 40 Zirconium 23 Vanadium
 41 Niobium 24 Chromium
 42 Molybdenum 25 Manganese.
 43 Technicium 26 Iron

.. 44 Ruthenium	27 Cobalt
45 Rodium	28 Nickel 46
Palladium	29 Copper 47
Silver	30 Zinc 48 Cadmium
31 Gallium	49 Indium	32
Germanium. 50 Tin	33
Arsenic	51 Antimony	34
Selenium	52 Tellurium	35
Bromine	53 Iodine GROUP	
V GROUP VI	54 Zenon	86
Radon	55 Caesium	87
Francium	56 Barium	88
Radium	57 Lanthanum	89
Actinium	58 Cerium	90
Thorium	59 Praseodymium	
91 Proactinium	60 Neodymium	
.... 92 Uranium	61 Promethium	
.... 93 Neptunium	62 Samarium	
.... 94 Plutonium	63 Europium	
.... 95 Americium	64 Gadolinium	
.... 96 Curium	65 Terbium	
.... 97 Berkelium	66 Dysprosium	
.... 98 Californium	67 Holmium	
.... 99 Einsteinium	68 Erbium	
.... 100 Fermium	69 Thulium	
101 70 Ytterbium	102 71	
Lutecium	103 72 Halfnium	
..... 104 73 Tantalum		
.... 105 74 Tungsten	106 75	
Rhenium	107 76 Osmium ..	
..... 108 Newtonium	77 Iridium ..	
..... 109 78 Platinum		
.. 110 79 Gold	111 80	
Mercury	112 81 Thallium ..	
..... 113 82 Lead		
114 83 Bismuth	115 84	
Polonium	116 85 Astatine ..	
..... 117 Research and Prelim.		
Engineering Space Vehicle Program		
General Objectives:		
In the following outlines of fundamental research, specific details have been referred to for the purpose of imparting a clear understanding of the nature of the proposed investigations. These preliminary outlines should not be construed as limiting the		

scope or delineating special interest. The goal of the project is a carefully integrated and adequately financed study of gravitation, embracing every relationship between gravitation and electrodynamics. It is to be remarked that the problem of relating gravitation to electrodynamics and the quantum theory is one which has taxed the ingenuity of some of the best mathematical brains for the last 30 years. So far, no very complete or satisfactory resolution of the matter has been found. Yet we are not completely in the dark with regard to it and the situation today is far from discouraging. The so-called "red-shift" produced by gravitation, and even the deviation of light by stars, are phenomena which are concerned with a relationship between gravitation and electrodynamics. Even though they are cosmological in extent, the magnitudes of these phenomena are small. Effects recently discovered in massive dielectrics point to the existence of hitherto unsuspected gravitational relationships and appear to have brought the matter for the first time into the realm of terrestrial experimentation. No one can deny the possibility that, as a result of this and other discoveries, a concept may result which so revolutionizes all our previous thoughts on gravity, electrodynamics and quantum theory as to render the story of the inter-relationships of these fields one of consistency and satisfaction. Such a program as herein outlined is necessarily of long-range. Unquestionably, there will be found many productive avenues of exploration which cannot be described in detail or even foreseen at the present time. Policy: The project must adopt a policy of inviting assistance from able physicists interested in the special problems involved. It must not fail to take into account and investigate


any phenomena which bears even remotely upon the subject. For example, in the study of physical properties of dielectrics, low-temperature research may be highly fruitful. Electrodynamic phenomena occur at low- temperatures which are completely unknown at room temperature. The possibilities of discovering wholly unsuspected gravitational effects below the superconductivity threshold, at temperatures approaching absolute zero, appear to be worth the costs involved. Library: The establishment of an adequate reference library on gravitation and related subjects, for the accumulation of technical information and to serve as liaison with academic institutions throughout the world, is a requirement of utmost importance particularly at the beginning of the program. Highest Priority: No one can guarantee results in research. No one can predict the direction the research will take. It is the express purpose of this project to obtain the technical answers as rapidly as possible by forming a coordinated program in which the best minds and all necessary laboratory facilities are brought together. It is the sincere hope that, in this way, a century of normal evolution in science, especially toward a better understanding of the nature of gravitation, may be compressed into from 5 to 10 years. Such a program is expensive but, as it was with the atomic bomb project in America, money was traded to gain a far more valuable commodity - time. So it may also be with man's ultimate conquest of space. As a necessary and inevitable prerequisite, a concerted study of gravitation is clearly indicated. We are forced to the conclusion that a research program organized specifically for this purpose can no longer be neglected. The Trouton-Noble Experiment (with massive dielectrics): The experiment concerns itself with an electromagnetic torque operating on a

charged condenser which moves with uniform velocity in a direction inclined to the normal of its surface. According to the theory of relativity, compensating effects, in this case having to do with the effect of the dielectric materials in the condenser on the torque aforesaid. In the days before universal acceptance of the theory of relativity, there was a reason to believe that measurements of the rotation of such a condenser as the above, when supported by some suspension, would serve to determine the velocity of the earth's motion through space. If, for a moment, we put ourselves in the mind of one who does not accept the theory of relativity in its entirety or wishes to test its validity further, the torque described above and possible rotation resulting from it becomes matters of experimental interest. A situation of great interest centers around the effect of the dielectric materials in the condenser on the torque. Now it appears that the original calculation of the torque is completely erroneous; and it appears that if the torque had been calculated correctly, invoking the same fundamental principles as were invoked in the earlier calculations, it would have been found to depend only upon the potential difference between the plates of the condenser and to be independent of the dielectric constant. However, a more refined analysis of the situation, which does not simply average the properties of the polarized molecules into a representation in terms of a dielectric constant, reveals that there may be a contribution to the torque which depends on the nature of the molecular dipoles, and in a manner which is not expressible in terms of the dielectric constant. The above conclusions were reached by Kennard and Swann independently by different processes of mathematical analysis. They have rendered the Trouton-Noble experiment one of considerable interest to



a person who had any doubts about the theory of relativity, and the interest would be enhanced by the bearing of the nature of the dielectric material upon the outcome of the experiment. Mass of the Electrons in Metals: This experiment has to do with the observations of momentum in a ring of conducting material carrying a current at the instant when the material is carried from the super-conducting to the non-superconducting state. Briefly, the above experiment envisages a metal ring in which a current of electricity has been produced by the creation of a magnetic field passing through the ring when all is at a temperature such that the super-conducting state prevails. Under such conditions, the current will continue practically indefinitely. If we now raise the temperature, the super-conductivity will disappear at a certain critical temperature, and the annular momentum of the electric current will be shared with the ordinary material of the ring in such a way as to give an angular rotation to the latter. The ring is, of course, to be envisaged as supported by a suspension and the angular rotation observed will depend upon the stiffness of this suspension. An interesting feature of the experiment lies in the fact that the sensitivity is greatest when the cross-section of the wire of the ring is smallest. The limiting conditions which determine the ultimate sensitivity are based upon the requirement that when the energy of the current is dissipated and the metal of the ring passes through the super-conducting state, the heat evolved shall not be sufficient to burn up the apparatus. The fundamental theoretical interest of the experiment lies in the fact that the angular rotation obtained depends upon the electronic mass, and theoretical considerations have been presented to support the belief that this electronic mass may be different for the electrons in a

metal than for the electrons in a free state. Most authorities on quantum theory are of the opinion that the effective mass of the electron in a metal is the same as that for an electron in a free state. However, even those who support this view are in favor of performing the experiment because of the complexity of the theoretical considerations involved. The most fundamental requirement is, of course, a means of producing liquid helium, and this implies a cryostat. If a cryostat were obtained, the potentialities of an enormous amount of other work in solid state physics would be provided for. Investigation of High-K Dielectrics at Low Temperatures: Research in solid state phenomena with special relation to dielectrics of high-K is of great current interest. Investigation of the properties of substances of high-K should be made in the realms of breakdown resistance, ferro-electrets, hysteresis, and allied phenomena. Special interest attaches also to the characteristics of electrets as such and to the conditions necessary to secure high activity of such electrets over long periods of time. In all the foregoing work, low temperature researches involving the cryostat would be of fundamental importance; for although the dielectrics are not usually used at low temperatures, many of the characteristics which determine their behavior at ordinary temperatures can be examined more readily by experiments performed at low temperatures. A survey of the literature on low temperature phenomena shows a large amount of work which has been carried out on the properties of paramagnetic salts, whereas the properties of dielectric materials have hardly been investigated at all. The reasons for this difference in emphasis are essentially understood. At liquid helium temperatures, the system of magnetic moments in most of the common paramagnetic salts is still in a thermally



disordered state so that its magnetic properties are still varying with temperature in an interesting fashion. In addition, since the technique of adiabatic demagnetization of a paramagnetic salt is the sole means, at the present time, of producing temperatures well below 1oK, it is only natural that a great amount of effort has been spent in the elucidation of the properties of these materials. Most normal dielectric materials show a negligible variation of their dielectric properties with temperature, especially in the liquid helium region. This may be seen by looking at the main sources of polarization in a dielectric, namely: 1. The electric polarizability, which arises from the fact that the outer electrons of an atom can be displaced with respect to the nucleus by an external electric field thereby creating a dipole moment. This is a property of the particular atom under consideration and is independent of temperature. 2. The ionic polarizability, arising from the displacement of positive ions with respect to negative ions in an ionic crystal. In most materials, this type of polarizability varies only very slowly with temperature, leading to a slight variation of dielectric constant with temperature. That this is not always the case is the reason for the present proposal. 3. Polarization due to the alignment of molecules with permanent dipole moments. In solids where the molecule is not free to rotate, this effect is absent. In recent years, a number of ferroelectric compounds have been discovered which are practically completely analogous in their dielectric behavior to ferromagnetic materials. Thus they show a Curie temperature, above which the dielectric constant follows a Curie-Weiss law and below which they exhibit spontaneous electrical polarization and hysteresis properties. Barium titanate (BaTiO_3) is the best known of these

compounds. Most of these compounds have Curies temperatures which are fairly high. Two compounds are known which have very low Curie temperatures. These are Potassium Tartrate (KTiO_3) and Lithium Thallium Tartrate ($\text{LiTiC}_4\text{H}_4\text{O}_6\text{H}_2\text{O}$), with Curie temperatures at 13.2oK and 10oK respectively. The existence of these very low Curie temperatures has created an additional interest in the study of dielectrics at the low temperatures obtainable with a Collins Helium cryostat. In addition to the intrinsic value of a program on the properties of dielectrics at low temperatures, it is conceivable that it might be possible to provide another means of producing temperatures lower than 1oK other than adiabatic demagnetization. If one had a ferroelectric material with a Curie temperature well below 1oK, then by the adiabatic, reversible depolarization of the material, it should be possible to produce a cooling effect (electro-caloric effect). Since the equipment involved in this process is somewhat simpler than in the corresponding magnetic case, it would be of considerable interest to investigate its feasibility. This method is not applicable below the Curie temperature since the presence of hysteresis and spontaneous polarization introduces irreversible heating effects upon applying or removing an external electric field. The Curie temperature of BaTiO_3 can be decreased by reducing the lattice parameter either by the addition of strontium or by application of external pressure. Presumably, this technique can be used to decrease the Curie temperature of KTiO_3 or $\text{LiTiC}_4\text{H}_4\text{O}_6\text{H}_2\text{O}$. An understanding of the factors which influence the Curie temperature and of the range of Curie temperatures in different crystals is important for the development of a basic

theory of ferroelectricity. In summary, it appears that a program on the properties of dielectrics at low temperatures can contribute substantially to an understanding of solids. The starting point for this program should logically be an investigation of KTiO_3 and $\text{LiTiC}_4\text{H}_4\text{O}_6\cdot\text{H}_2\text{O}$ as well as structurally similar crystals and their solid solutions with each other. Electromagnetic Equations for the Super-Conductive State: Among the many interesting phenomena which occur at low temperatures, superconductivity has long held the attention of experimentalist and theoretician alike since its discovery by H. Kammerlingh Onnes in 1911. With the discovery of the Meissner effect in 1933, the basic experimental behavior necessary for the development of an electrodynamic theory of super-conductivity has been established. F. and H. London, in 1935, developed a set of equations which describe the macroscopic electrodynamic behavior of super-conductors in a quantitative manner to the present time. One experiment which would shed considerable light on the correctness of these equations has been suggested by F. London. This experiment involves a study of the magnetic properties of a rotating sphere. The theory of this experiment is worked out in complete detail by F. London. The following is a physical description of the nature of this experiment. Consider a sphere of radius R . If we start with the sphere at rest below its super-conducting transition temperature and bring it into motion with uniform angular velocity (w), then by considering the super-conducting electrons are perfectly free it can be deduced that the sphere should become magnetized upon rotation. The reason for this is as follows: When the sphere is initially set into motion, the electrons - being perfectly free

from interaction with the crystal lattice - will not move with the sphere, and a current is set up due to their relative motion. This changing current in turn induces an electric field within the sphere which acts on the electrons in such a manner as to accelerate them in the direction of rotation of the sphere. The final result is that when the sphere has reached a constant angular velocity the super-conducting electrons everywhere move with the sphere except for a narrow layer at the surface, where they lag behind slightly to produce a small current. This result was predicted on the basis of a free electron theory before the development of the theory of F. and H. London. The London theory predicts the same result for the rotating sphere, except that it makes an additional prediction. F. London states that the rotating sphere will have a magnetic moment independent of the prehistory of the sphere. In particular, if a rotating sphere is cooled below its transition temperature while rotating, the sphere will acquire the same magnetic moment as it would upon starting from rest below its transition temperature and being brought to the same angular velocity. On the basis of a free electron theory of super-conductivity, it is difficult to understand how a sphere which is already rotating will suddenly acquire a magnetic moment upon being cooled below its transition temperature. In this case, the electrons move with the sphere above the transition temperature due to their finite interaction with the lattice (finite resistance). That they should suddenly lag behind to produce a magnetic moment on cooling below the transition temperature seems surprising. The magnetic moment predicted for the rotating sphere is small, but should be measurable with sufficiently careful experimental technique. This experiment would constitute a fundamental



method of testing the basic assumptions of
the London theory.

ELECTROGRAVITICS SYSTEMS

**REPORTS ON A NEW PROPULSION
METHODOLOGY**

Edited by

Thomas Valone, M.A., P.E.

Foreword by *Elizabeth Rauscher, Ph.D.*

Integrity Research Institute
Washington, DC 20005

for
T. TownsendBrown

**Electrogravitics Systems: Reports on a New
Propulsion Methodology**

Edited by Thomas Valone

First Edition, April, 1994

Second Edition, November, 1995

Third Edition, April 1999

Fourth Edition, January, 2001

"The U.S. Antigravity Squadron"
Copyright © 1993 by Paul A. LaViolette

Cover illustration Copyright © 1993
by Paul A. LaViolette

ISBN 0-9641070-0-7

published by
INTEGRITY RESEARCH INSTITUTE
1220 L Street NW, Suite 100-232
Washington, DC 20005
202-452-7674
800-295-7674
www.integrity-research.org

sendforfree catalog

CONTENTS

Foreword.....	4
<i>Elizabeth Rauscher, Ph.D.</i>	
Editor's Introduction.....	6
<i>Thomas Valone, U.A., P.E.</i>	
Electrogravitics Systems.....	11
<i>Aviation Studies, Ltd.</i>	
The Gravities Situation.....	42
<i>Gravity Rand Ltd., Div. of Aviation Studies, Ltd.</i>	
Negative Mass As a Gravitational Source of Energy in the Quasi-Stellar Radio Sources.....	73
<i>Banesh Hoffman</i>	
The U.S. Antigravity Squadron.....	78
<i>Paul LaViolette</i>	
Appendix.....	97
<i>Collection of T. T. Brown's Patents</i>	

FOREWORD

by Elizabeth Rauscher, Ph.D., *Professor of Nuclear and Astrophysics, University of Nevada*

Electromagnetism is bipolar, i.e. it attracts and repels. We can shield X-rays, gamma rays, radio waves, etc., but what about gravity? Gravity appears to have only one polarity — attraction! We have balloons, planes, and rockets that overcome gravity but can we build a shield against gravity? Roger Babson, a good friend of Thomas Edison, established the Gravity Research Foundation in 1948 at Edison's suggestion. So what would anti-gravity "look like"? Let us explore these issues:

Standard physical models include *four* fundamental forces in Nature. They are the nuclear force, the electromagnetic force, the weak, nuclear decay force, and gravitational force. The nuclear force and the gravitational force have the similar property of being attractive only. What of anti-matter ~ does it rise in a gravitational field? Such an experiment was attempted at the Stanford Linear

Accelerator Center, Stanford, CA without confirmed results.

In 1971, I published a book and several papers on a ten dimensional geometric model of quantum gravity in which I treated the four major force fields on an "equal footing" in such a manner as to consider them as *bi* or dual polar, having both attraction and repulsion.

T. Townsend Brown, who I met in 1981, led me to replicate his research on some properties of electrostatics, capacitance and anomalous current flows on unique materials. Unlike the current view, electrostatic phenomena are very complex. How does this work relate to the ideas of UFO propulsion (an early interest of T.T. Brown)? Certainly he has presented the scientific community with many questions we need to investigate.

I have also theoretically examined a five and eight dimensional geometry which includes the Kaluza Klein geometry which is an abstract formalism relating electromagnetism to the gravitational field. This model interested Albert Einstein in the 1930's. There is a long path between theoretical concepts, romantic wishes and preliminary experiments to detailed experimental verification and actual designed technology.

Let us re-examine Brown's works and rethink some of the issues which he has suggested to us. Science is an ongoing process, not a fixed set of facts, ever changing and developing.

Prof. Elizabeth A. Rauscher, Ph.D.

Editor's INTRODUCTION

"Newton attempted to explain the force of gravity on two hypotheses: the existence of a medium, or ether, and action at a distance. The first hypothesis he rejected as being physically absurd, the second as contrary to reason. Newton had, therefore, no theory of gravity. However, his long and sustained effort to understand gravity was not without at least one serious consequence. For involvement in the ether theory obscured from Newton the universal character of the inverse-square relation, and delayed for twenty years his final formulation of the law of gravity."*

Many people have had the opportunity to meet and learn from T. Townsend Brown before he passed away about a decade ago. I was fortunate to correspond with him after I returned from the Gravity Field Energy Conference in Germany, 1980, where his name was mentioned a lot. He was a friendly but quiet person who learned some of his magic from Dr. Paul Alfred Biefeld, a physicist at the California Institute for Advanced Studies. "In 1923, Biefeld discovered that a heavily charged electrical condenser moved toward its positive pole when suspended in a gravitational field. He assigned Brown to study the effect as a research project." (*How to Build a Flying Saucer*, Pawlicki, Prentice-Hall, 1981)

The *Electric Spacecraft Journal* also notes that Brown's effect may have an earlier precedent as well, citing an article by Nipher which appeared in the March, 1918 issue of the *Electrical Experimenter* (p.743). Whoever deserves credit, the question today is the credibility of T. T. Brown and whether his effect is significant for propulsion.

It can be argued that if the two main reports of this anthology are historically valid, Brown may have been debriefed after his release from military consultation service. The interesting fact about the first report is that under the black streak, prominently seen across the original title page, the word "CONFIDENTIAL" actually was written. Dr. Paul LaViolette notes that the *Electrogravitics Systems* report may be obtained from the Wright-Patterson AFB technical library, even though it is not actually listed on their library computer. He found that it is not available from any other library in the U.S. Paul deserves our gratitude for retrieving the first report.

To authenticate the discovery of these two reports, I have included a copy of the 1956 *Gravitics Situation Order Form* at the end of this

* Evans, *American Journal of Physics*, 26, 619, 1958

Introduction, as well as copies of the original title pages for *Electrogravitics Systems*. Aviation Studies is also still in business today.

I feel that it is important to mention that even though Talley has, in 1991, attempted a replication of Brown's work, up to 19 kV (and 33 kV with anomalous breakdown effects) for Edwards AFB, we see in these uncovered reports, mention of the "magic number" of 50,000, with a target wattage of 50,000 kW and a dielectric K value of 50,000. Brown is also quoted as recommending heating of the cathode.

Many people have tried low voltage replication (e.g. Dr. Harold Puthoff...!? kV) but these reports indicate that higher voltage is better, up to a maximum effect.

Cravens, also a recent military consultant, gives Brown a high rating for practicality and propulsion (p. 79 of his report).* He also notes that older high voltage supplies most likely had time-varying voltage regulation signals superimposed upon the DC electrostatic field. This fact, along with similar Hutchison and Searl effect signals, may point the way toward an overlooked detail in the race to replicate T.T. Brown.

Also, *Science* magazine notes that Puthoff, Haisch and Rueda present a new theory of inertia which they say raises a "provocative notion that inertia, once understood, might be controlled" (Vol. 263, Feb., 1994). This underscores our return to T.T. Brown and his research because of its simplicity.

I have included an extra article by Dr. Banesh Hoffman, the biographer of Einstein, because he also theorized about negative mass, the concept which Gravity Rand Ltd. believes holds a key to understanding the Biefeld-Brown effect.

What can be said about Dr. LaViolette's well-researched sleuth about the B-2 aircraft? It is a provocative thesis that presents a plausible modern application of Brown's research, explaining many of the B-2 characteristics. As more facts are uncovered, we will consider future additions to the Electrogravitics Series.

Thomas Valone, M.A., P.E.
Physicist and Prof. Engineer

* D.L. Cravens, "Electric Propulsion Study," prepared for the: Astronautics Lab (AFSC), Space Systems Division, Edwards AFB, CA 93523, Aug. 1990, AL-TR-89-040, #ADA 227121

AVIATION STUDIES (INTERNATIONAL) LIMITED
Aviation House, 66 Sloane St., London S.W.1. Tel. SLO 0577/8 Cables AVIALEG LONDON

THE GRAVITICS SITUATION

A new report of interest both to management and to engineers will be available in Mid-December. The section for engineers is in three parts. First is an assault on the quantum mechanical approach to the existence of negative mass; this is supported by Mozer's paper who in turn looks to Schrodinger's time independent equation with the center of mass motion removed. A second approach is electrogavitics, which is Townsend Brown's work, and the report contains a complete specification of Brown's gravitator device and a survey of this branch of activity. Third is an outline of possible ways of finding the link between gravitational fields and nuclear energy which draws on Beaumont's complex spin, Deser, Arnowitt and Bondi. This section is supported by Deser and Arnowitt's proposition, as they expressed it. This part is perhaps the most important of all.

For management there is a section dealing with the impact of various approaches on company policy; this includes a rough estimation of the value of current work. This section also includes a glossary of terms in an attempt to prevent the dictionary running away with the science; together with Gravity Rand observations. (Fee \$30: £10)

To: Gravity Rand Ltd.
Aviation House, 66 Sloane Street,
London, S.W.1.
England.

Please send us your report, "THE GRAVITICS SITUATION".

- Payment should be made out to AVIATION STUDIES (INTERNATIONAL) LIMITED.
- We enclose \$ (£).....
- Invoice us later

Address: (to be used for mailing).....

3 1401 00034 5879

Technicar
Daytona Beach

ELECTROGRAVITICS SYSTEMS

An examination of electrostatic motion,
dynamic counterforce and barycentric control.

AVIATION STUDIES (INTERNATIONAL) LTD

TL
565
A9



3 1401 00034 5879

DATE DUE

ALL BOOKS SUBJECT TO RECALL AFTER 4 WEEKS

11 JUL 1984

26 AUG 84

ALA FIRM



TECHNICAL LIBRARY

Wright - Patterson
Air Force Base, Ohio 45433

ELECTROGRAVITICS SYSTEMS

An examination of electrostatic motion, dynamic counterbary and barycentric control.

**AVIATION STUDIES (INTERNATIONAL)
LTD.**

Prepared by:
Gravity Research Group
Special Weapons Study Unit
Aviation Studies (International) Limited, London.
29-31 Cheval Place, Knightsbridge
London, S,W,7 England

Report GRG-013/56 February 1956

CONTENTS

Introductory Notes

Discussion

Conclusions

APPENDIX

I - Aviation Report Extracts

II - Electrostatic Patents

ELECTROGRAVITICS SYSTEMS

An examination of electrostatic motion, dynamic counterbary and barycentric control

It has been accepted as axiomatic that the way to offset the effects of gravity is to use a lifting surface and considerable molecular energy to produce a continuously applied force that, for a limited period of time, can remain greater than the effects of gravitational attraction. The original invention of the glider, and evolution of the briefly self-sustaining glider, at the turn of the century led to progressive advances in power and knowledge. This has been directed to refining the classic Wright Brothers' approach. Aircraft design is still fundamentally as the Wrights adumbrated it, with wings, body, tails, moving or flapping controls, landing gear and so forth. The Wright biplane was a powered glider, and all subsequent aircraft, including the supersonic jets of the nineteen-fifties are also powered gliders. Only one fundamentally different flying principle has so far been adopted with varying degrees of success. It is the rotating wing aircraft that has led to the jet lifters and vertical pushers, coleopters, ducted fans and lift induction turbine propulsion systems.

But during these decades there was always the possibility of making efforts to discover the nature of gravity from cosmic or quantum theory, investigation and observation, with a view to discerning the physical properties of aviations' enemy.

It has seemed to Aviation Studies that for some time insufficient attention has been directed to this kind of research. If it were successful such developments would change the concept of sustentation, and confer upon a vehicle qualities that would now be regarded as the ultimate in aviation.

This report summarizes in simple form the work that has been done and is being done in the new field of electrogravitics. It also outlines the various possible lines of research into the nature and constituent matter of gravity, and how it has changed from Newton to Einstein to the modern Hlavaty concept of gravity as an electromagnetic force that may be controlled like a light wave.

The report also contains an outline of opinions on the feasibility of different electrogravitics systems and there is reference to some of the barycentric control and electrostatic rigs in operation.

Also included is a list of references to electrogravitics in successive Aviation Reports since a drive was started by Aviation Studies (International) Limited to suggest to aviation business eighteen months ago that the rewards of success are too far-reaching to be overlooked, especially in view of the hopeful judgement of the most authoritative voices in microphysics. Also listed are some relevant patents on electrostatics and electrostatic generators in the United States, United Kingdom and France.

Gravity Research Group

DISCUSSION

Electrogravitics might be described as a synthesis of electrostatic energy use for propulsion - either vertical propulsion or horizontal or both - and gravitics, or dynamic counterbary, in which energy is also used to set up a local gravitational force independent of the earth's.

Electrostatic energy for propulsion has been predicted as a possible means of propulsion in space when the thrust from a neutron motor or ion motor would be sufficient in a dragless environment to produce astronomical velocities. But the ion motor is not strictly a part of the science of electrogravitics, since barycentric control in an electrogravitics system is envisaged for a vehicle operating within the earth's environment and it is not seen initially for space application. Probably large scales space operations would have to await the full development of electrogravitics to enable large pieces of equipment to be moved out of the region of the earth's strongest gravity effects. So, though electrostatic motors were thought of in 1925, electrogravitics had its birth after the War, when Townsend Brown sought to improve on the various proposals that then existed for electrostatic motors sufficiently to produce some visible manifestation of sustained motion. Whereas earlier electrostatic test were essentially pure research. Brown's rigs were aimed from the outset at producing a flying article. As a private venture he produced evidence of motion using condensers in a couple of saucers suspended by arms rotating round a central tower with input running down the arms. The massive-k situation was summarized subsequently in a report, Project Winterhaven, in 1952. Using the data some conclusions were arrived at that might be expected from ten or more years of intensive development - similar to that, for instance, applied to the turbine engine. Using a number of assumptions as to the nature of gravity, the report postulated a saucer as the basis of a possible interceptor with Mach 3 capability. Creation of a local gravitational system would confer upon the fighter the sharp-edged changes of direction typical of motion in space.

The essence of electrogravitics thrust is the use of a very strong positive charge on one side of the vehicle and a negative on the other. The core of the motor is a condenser and the ability of the condenser to hold its charge (the K-number) is the yardstick of performance. With air as 1, current dielectrical materials can yield 6 and use of barium aluminate can raise this considerable,

barium titanium oxide (a baked ceramic) can offer 6,000 and there is promise of 30,000, which would be sufficient for supersonic speed.

The original Brown rig produced 30 fps on a voltage of around 50,000 and a small amount of current in the milliamp range. There was no detailed explanation of gravity in Project Winterhaven, but it was assumed that particle dualism in the subatomic structure of gravity would coincide in its effect with the issuing stream of electrons from the electrostatic energy source to produce counterbary. The Brown work probably remains a realistic approach to the practical realization of electrostatic propulsion and sustentation. Whatever may be discovered by the Gravity Research Foundation of New Boston a complete understanding and synthetic reproduction of gravity is not essential for limited success. The electrogravitics saucer can perform the function of a classic lifting surface - it produces a pushing effect on the under surface and a suction effect on the upper, but, unlike the airfoil, it does not require a flow of air to produce the effect.

First attempts at electrogravitics are unlikely to produce counterbary, but may lead to development of an electrostatic VTOL vehicle. Even in its developed form this might be an advance on the molecular heat engine in its capabilities. But hopes in the new science depend on an understanding of the close identity of electrostatic motivating forces with the source and matter of gravity. It is fortuitous that lift can be produced in the traditional fashion and if an understanding of gravity remains beyond full practical control, electrostatic lift might be an adjunct of some significance to modern thrust producers. Research into electrostatics could prove bénéficiai to turbine development, and heat engines in general, in view of the usable electron potential round the periphery of any flame. Material for electrogravitics and especially the development of commercial quantities of high-k material is another dividend to be obtained from electrostatic research even if it produces no counterbary. This is a line of development that Aviation Studies' Gravity Research Group is following.

One of the interesting aspects of electrogravitics is that a breakthrough in almost any part of the broad front of general research on the intranuclear processes may be translated into a meaningful advance toward the feasibility of electrogravitics systems. This demands constant monitoring in the most likely areas of the physics of high energy sub-nuclear particles. It is difficult to be overoptimistic about the prospects of gaining so complete a grasp of gravity while the world's physicists are still engaged in a study of fundamental

particles - that is to say those that cannot be broken down any more. Fundamental particles are still being discovered - the most recent was the Segre-Chamberlain-Wiegand attachment to the bevatron, which was used to isolate the missing anti-proton, which must - or should be presumed to - exist according to Dirac's theory of the electron. Much of the accepted mathematics of particles would be wrong if the anti-proton was proved to be non-existent. Earlier Eddington has listed the fundamental particles as:

e. The charge of an electron.

m. The mass of an electron.

M. The mass of proton.

h. Planck's constant

c. The velocity of light.

G. The constant of gravitation, and

lambda. The cosmical constant.

It is generally held that no one of these can be inferred from the others. But electrons may will disappear from among the fundamental particles, though, as Russell says, it is likely that *e* and *m* will survive. The constants are much more established than the interpretation of them and are among the most solid of achievements in modern physics.

* * * *

Gravity may be defined as a small-scale departure from Euclidean space in the general theory of relativity. The gravitational constant is one of four dimensionless constants: first, the mass relation of the nucléon and electron, second is e^2/hc , third, the Compton wavelength of the proton, and fourth is the gravitational constant, which is the ratio of the electrostatic to the gravitational attraction between the electron and the proton.

One of the stumbling blocks in electrogravitics is the absence of any

satisfactory theory linking these four dimensionless quantities. Of the four, moreover, gravity is decidedly the most complex, since any explanation would have to satisfy both cosmic and quantum relations more acceptably and intelligibly even than the unified field theory. A gravitational constant of around 10^{-39} has emerged from quantum research and this has been used as a tool for finding theories that could link the two relations. This work is now in full progress, and developments have to be watched for the aviation angle. Hitherto Dirac, Eddington, Jordan and others have produced differences in theory that are too wide to be accepted as consistent. It means therefore that (i) without a cosmical basis, and (ii) with an imprecise quantum basis and (iii) a vague hypothesis on the interaction, much remains still to be discovered. Indeed some say that a single interacting theory to link up the dimensionless constants is one of three major unresolved basic problems of physics. The other two main problems are the extension of quantum theory and a more detailed knowledge of the fundamental particles.

All this is some distance from Newton, who saw gravity as a force acting on a body from a distance, leading to the tendency of bodies to accelerate towards each other. He allied this assumption with Euclidean geometry, and time was assumed as uniform and acted independently of space. Bodies and particles in space normally moved uniformly in straight lines according to Newton, and to account for the way they sometimes do not do so, he used the idea of a force of gravity acting at a distance, in which particles of matter cause in others an acceleration proportional to their mass, and inversely proportional to the square of the distance between them.

But Einstein showed how the principle of least action, or the so-called cosmic laziness means that particles, on the contrary, follow the easiest path along geodesic lines and as a result they get readily absorbed into space-time. So was born non-linear physics. The classic example of non-linear physics is the experiment in bombarding a screen with two slits. When both slits are open particles going through are not the sum of the two individually but follows a non-linear equation. This leads on to wave-particle dualism and that in turn to the Heisenberg uncertainty principle in which an increase in accuracy in measurement of one physical quantity means decreasing accuracy in measuring the other. If time is measured accurately energy calculations will be in error; the more accurate the position of a particle is established the less certain the velocity will be; and so on. This basic principle of causality of microphysics affects the study of gravity in the special and general theories of relativity. Lack of pictorial image in the quantum physics of this

interrelationship is a difficulty at the outset for those whose minds remain obstinately Euclidean.

In the special theory of relativity, space-time is seen only as an undefined interval which can be defined in any way that is convenient and the Newtonian idea of persistent particles in motion to explain gravity cannot be accepted. It must be seen rather as a synthesis of forces in a four dimensional continuum, three to establish the position and one the time. The general theory of relativity that followed a decade later was a geometrical explanation of gravitation in which bodies take the geodesic path through space-time. In turn this means that instead of the idea of force acting at a distance it is assumed that space, time, radiation and particles are linked and variations in them from gravity are due rather to the nature of space.

Thus gravity of a body such as the earth, instead of pulling objects towards it as Newton postulated, is adjusting the characteristics of space and, it may be inferred, the quantum mechanics of space in the vicinity of the gravitational force. Electrogravitics aims at correcting this adjustment to put matter, so to speak, 'at rest'.

* * * *

One of the difficulties in 1954 and 1955 was to get aviation to take electrogravitics seriously. The name alone was enough to put people off. However, in the trade much progress has been made and now most major companies in the United States are interested in counterbary. Groups are being organized to study electrostatic and electromagnetic phenomena. Most of industry's leaders have made some reference to it. Douglas has now stated that it has counterbary on its work agenda by does not expect results yet awhile. Hiller has referred to new forms of flying platform, Glenn Martin say gravity control could be achieved in six years, but they add that it would entail a Manhattan District type of effort to bring it about. Sikorsky, one of the pioneers, more or less agrees with the Douglas verdict and says that gravity is tangible and formidable, but there must be a physical carrier for this immense trans-spatial force. This implies that where a physical manifestation exists, a physical device can be developed for creating a similar force moving in the opposite direction to cancel it. Clarke Electronics state they have a rig, and add that in their view the source of gravity's force will be understood sooner than some people think. General Electric is working on the use of electronic

rigs designed to make adjustments to gravity - this line of attack has the advantage of using rigs already in existence for other defence work. Bell also has an experimental rig intended, as the company puts it, to cancel out gravity, and Lawrence Bell has said he is convinced that practical hardware will emerge from current programs. Grover Leoning is certain that what he referred to as an electro-magnetic contra-gravity mechanism will be developed for practical use. Convair is extensively committed to the work with several rigs. Lear Inc., autopilot and electronic engineers have a division of the company working on gravity research and so also has the Sperry division of Sperry-Rand. This list embraces most of the U.S. aircraft industry. The remainder, Curtiss-Wright, Lockheed, Boeing and North American have not yet declared themselves, but all these four are known to be in various stages of study with and without rigs.

In addition, the Massachusetts Institute of Technology is working on gravity, the Gravity Research Foundation of New Boston, the Institute for Advanced Study at Princeton, the CalTech Radiation Laboratory, Princeton University and the University of North Carolina are all active in gravity. Glenn L. Martin is setting up a Research Institute of Advanced Study which has a small staff working on gravity research with the unified field theory and this group is committed to extensive programs of applied research. Many others are also known to be studying gravity, some are known also to be planning a general expansion in this field, such as in the proposed Institute for Pure Physics at the University of North Carolina.

A certain amount of work is also going on in Europe. One of the French nationalized constructors and one company outside the nationalized elements have been making preliminary studies, and a little company money has in one case actually been committed. Some work is also going on in Britain where rigs are now in existence. Most of it is private venture work, such as that being done by Ed Hull a colleague of Townsend Brown who, as much as anybody, introduced Europe to electrogravitics. Aviation Studies' Gravity Research Group is doing some work, mainly on k studies, and is sponsoring dielectric investigations.

One Swedish company and two Canadian companies have been making studies, and quite recently the Germans have woken up to the possibilities. Several of the companies have started digging out some of the early German papers on wave physics. They are almost certain to plan a gravities program. Curiously enough the Germans during the war paid no attention to

electrogravitics. This is one line of advance that they did not pioneer in any way and it was basically a U.S. creation. Townsend Brown in electrogravitics is the equivalent of Frank Whittle in gas turbines. This German overlooking of electrostatics is even more surprising when it is remembered how astonishingly advanced and prescient the Germans were in nuclear research. (The modern theory of making thermonuclear weapons without plutonium fission initiators returns to the original German idea that was dismissed, even ridiculed. The Germans never went very far with fission, indeed they doubted that this chain would ever be made to work.) The German air industry, still in the embryo stage, has included electrogravitics among the subjects it intends to examine when establishing the policy that the individual companies will adopt after the present early stage of foreign license has enabled industry to get abreast of the other countries in aircraft development.

* * * *

It is impossible to read through this summary of the widening efforts being made to understand the nature of matter of gravity without sharing the hope that many groups now have, of major theoretical breakthroughs occurring before very long. Experience in nucleonics has shown that when attempts to win knowledge on this scale are made, advances are soon seen. There are a number of elements in industry, and some managements, who see gravity as a problem for later generations. Many see nothing in it all and they may be right. But as said earlier, if Dr. Vaclav Hlavaty thinks gravity is potentially controllable that surely should be justification enough, and indeed inspiration, for physicists to apply their minds and for management to take a risk. Hlavaty is the only man who thinks he can see a way of doing the mathematics to demonstrate Einstein's unified field theory - something that Einstein himself said was beyond him. Relativity and the unified field theory go to the root of electrogravitics and the shifts in thinking, the hopes and fears, and a measure of progress is to be obtained only in the last resort from men of this stature.

Major theoretical breakthroughs to discover the sources of gravity will be made by the most advanced intellects using the most advanced research tools. Aviations role is therefore to impress upon physicists of this calibre with the urgency of the matter and to aid them with statistical and peripheral investigations that will help to clarify the background to the central mathematical and physical puzzles. Aviation could also assist by recruiting some of these men as advisers. Convair has taken the initiative with its

recently established panel of advisers on nuclear projects, which include Dr. Edward Teller of the University of California. At the same time much can be done in development of laboratory rigs, condenser research and dielectric development, which do not require anything like the same cerebral capacity to get results and make a practical contribution.

As gravity is likely to be linked with the new particles, only the highest powered particle accelerators are likely to be of use in further fundamental knowledge. The country with the biggest tools of this kind is in the best position to examine the characteristics of the particles and from those countries the greatest advances seem most likely.

Though the United States has the biggest of the bevatrons - the Berkeley bevatron is 6.2 bev - the Russians have a 10 bev accelerator in construction which, when it is completed, will be the world's largest. At Brookhaven a 25 bev instrument is in development which, in turn, will be the biggest. Other countries without comparable facilities are of course at a great disadvantage from the outset in the contest to discover the explanation of gravity. Electrogravitics, moreover, unfortunately competes with nuclear studies for its facilities. The clearest thinking brains are bound to be attracted to localities where the most extensive laboratory equipment exists. So, one way and another, results are most likely to come from the major countries with the biggest undertakings. Thus the nuclear facilities have a direct bearing on the scope for electrogravitics work.

The OEEC report in January made the following points:

The U.S. has six to eight entirely different types of reactors in operation and many more under construction. Europe has now two different types in service.

The U.S. has about 30 research reactors plus four in Britain, two in France.

The U.S. has two nuclear-powered marine engines. Europe has none, but the U.K. is building one.

Isotope separation plants for the enrichment of uranium in the U.S. are roughly 11 times larger than the European plant in Britain.

Europe's only heavy water plant (in Norway) produces somewhat less than one-twentieth of American output.

In 1955 the number of technicians employed in nuclear energy work in the U.S. was about 15,000; there are about 5,000 in Britain, 1,800 in France, and about 1,000 in the rest of Europe. But the working party says that pessimistic conclusions should not be drawn from these comparisons. European nuclear energy effort is unevenly divided at the moment, but some countries have notable achievements to their credit and important developments in prospect. The main reason for optimism is that, taken as a whole, iEurope's present nuclear effort falls very short of its industrial potential.[†]

Though gravity research, such as there has been of it, has been unclassified, new principles and information gained from the nuclear research facilities that have a vehicle application is expected to be withheld.

The heart of the problem to understanding gravity is likely to prove to be the way in which the very high energy sub-nuclear particles convert something, whatever it is, continuously and automatically into the tremendous nuclear and electromagnetic forces. Once this key is understood, attention can later be directed to finding laboratory means of duplicating the process and reversing its force lines in some local environment and returning the energy to itself to produce counterbary. Looking beyond it seems possible that gravitation will be shown to be a part of the universal electro-magnetic processes and controlled in the same way as a light wave or radio wave. This is a synthesis of the Einstein and Hlavaty concepts. Hence it follows that though in its initial form the mechanical processes for countering gravity may initially be massive to deal with the massive forces involved, eventually this could be expected to form some central power generation unit. Barycentric control in some required quantity could be passed over a distance by a form of radio wave. The prime energy source to energize the waves would of course be nuclear in its origins.

It is difficult to say which lines of detailed development being processed in the immediate future is more likely to yield significant results. Perhaps the three most promising are: first, the new attempt by the team of men led by Chamberlain working with the Berkeley bevatron to find the anti-neutron, and to identify more of the characteristics of the anti-proton and each of the string of high energy particles that have been discovered during recent operation at

6.2 bev.*

A second line of approach is the United States National Bureau of Standards program to pin down with greater accuracy the acceleration value of gravity. The presently accepted figure of 32.174 feet per second per second is known to be not comprehensive, though it has been sufficiently accurate for the limited needs of industry hitherto. The NBS program aims at re-determining the strength of gravity to within one part of a million. The present method has been to hold a ball 16 feet up and chart the elapsed time of descent with electronic measuring equipment. The new program is based on the old, but with this exceptional degree of accuracy it is naturally immensely more difficult and is expected to take 3 years.

A third promising line is the new technique of measuring high energy particles in motion that was started by the University of California last year. This involves passing cosmic rays through a chamber containing a mixture of gas, alcohol and water vapor. This creates charged atoms, or positive ions, by knocking electrons off the gas molecules. A sudden expansion of the chamber results in a condensation of water droplets along the track which can be plotted on a photographic plate. This method makes it easier to assess the energy of particles and to distinguish one from the other. It also helps to establish the characteristics of the different types of particle. The relationship between these high energy particles; and their origin and characteristics, have a bearing on electrogravitics in general.

So much of what has to be discovered as a necessary preliminary to gravity is of no practical use by itself. There is no conceivable use, for instance, for the anti-proton, yet its discovery even at a cost of \$9-million is essential to check the mathematics of the fundamental components of matter. Similarly it is necessary to check that all the nuclear ghosts that have been postulated theoretically do in fact exist. It is not, moreover, sufficient, as in the past, only to observe the particles by radiation counters. In each instance a mechanical maze has to be devised and attached to a particle accelerator to trap only the particle concerned. Each discovery becomes a wedge for a deeper probe of the nucleus. Many of the particles of very high energy have only a fleeting

*The reaction is as follows: protons are accelerated to 6.2 bev, and directed at a target of copper. When the proton projectile hits a neutron in one of the copper atoms the following emerge: the two original particles (the projectile and the struck neutron) and a new pair of particles, a proton and anti-proton. The anti-proton continues briefly until it hits another proton, then both disappear and decay into mesons.

existence and collisions that give rise to them from bevatron bombardment is a necessary prerequisite to an understanding of gravity. There are no shortcuts to this process.

Most of the major programs for extending human knowledge on gravity are being conducted with instruments already in use for nuclear research and to this extent the cost of work exclusively on gravitational examinations is still not of major proportions. This has made it difficult for aviation to gauge the extent of the work in progress on gravity research.

CONCLUSIONS

1. No attempts to control the magnitude or direction of the earth's gravitational force have yet been successful. But if the explanation of gravity is to be found in the as yet undetermined characteristics of the very high energy particles it is becoming increasingly possible with the bevatron to work with the constituent matter of gravity. It is therefore reasonable to expect that the new bevatron may, before long, be used to demonstrate limited gravitational control.
2. An understanding and identification of these particles is on the frontiers of human knowledge, and a full assessment of them is one of the major unresolved puzzles of the nucleus. An associated problem is to discover a theory to account for the cosmic and quantum relations of gravity, and a theory to link the gravitational constant with the other three dimensionless constants.
3. Though the obstacles to an adequate grasp of microphysics still seem formidable, the transportation rewards that could follow from electrogravitics are as high as can be envisaged. In a weightless environment, movement with sharp-edged changes of direction could offer unique manoeuvrability.
4. Determination of the environment of the anti-proton, discovery of the anti-neutron and closer examination of the other high energy particles are preliminaries to the hypothesis that gravity is one aspect of electromagnetism that may eventually be controlled like a wave. When the structure of the nucleus becomes clearer, the influence of the gravitational force upon the nucleus and the nature of its behavior in space will be more readily understood. This is a great advance on the Newtonian concept of gravity acting at a distance.
5. Aviation's role appears to be to establish facilities to handle many of the peripheral and statistical investigations to help fill in the background on electrostatics.
6. A distinction has to be made between electrostatic energy for propulsion

and counterbary. Counterbary is the manipulation of gravitational force lines; barycentric control is the adjustment to such manipulative capability to produce a stable type of motion suitable for transportation.

7. Electrostatic energy sufficient to produce low speeds (a few thousand dynes) has already been demonstrated. Generation of a region of positive electrostatic energy on one side of a plate and negative on the other sets up the same lift or propulsion effect as the pressure and suction below and above a wing, excepts that in the case of electrostatic application no airflow is necessary.
8. Electrostatic energy sufficient to produce a Mach 3 fighter is possible with megavolt energies and a k of over 10,000.
9. k figures of 6,000 have been obtained from some ceramic materials and there are prospects of 30,000.
10. Apart from electrogravitics there are other rewards from investment in electrostatic equipment. Automation, autonetics and even turbine development use similar laboratory facilities.
11. Progress in electrogravitics probably awaits a new genius in physics who can find a single equation to tie up all the conflicting observations and theory on the structure and arrangement of forces and the part the high energy particles play in the nucleus. This can occur any time, and the chances are improved now that bev. energies are being obtained in controlled laboratory conditions.

APPENDIX I

EXTRACTS FROM AVIATION REPORT

ANTI-GRAVITATION RESEARCH

The basic research and technology behind electro-anti-gravitation is os much in its infancy that this is perhaps one field of development where not only the methods but the ideas are secret. Nothing therefore can be discussed freely at the moment. Very few papers on the subject have been prepared so far, and the only schemes that have seen the light of day are for pure research into rigs designed to make objects float around freely in a box. There are various radio applications, and aviation medicine departments have been looking for something that will enable them to study the physiological effects on the digestion and organs of an environment without gravity. There are however long term aims of a more revolutionary nature that envisage equipment that can defeat gravity.

Aviation Report 20 August 1954

MANAGERIAL POLICY FOR ANTI-GRAVITICS

The prospect of engineers devising gravity-defeating equipment - or perhaps it should be described as the creation of pockets of weightless environments - does suggest that as a long term policy aircraft constructors will be required to place even more emphasis on electro-mechanical industrial plant, than is now required for the transition from manned to unmanned weapons. Anti-gravitics work is therefore likely to go to companies with the biggest electrical laboratories and facilities. It is also apparent that anti-gravitics, like other advanced sciences, will be initially sponsored for its weapon capabilities. There are perhaps two broad ways of using the science - one is to postulate the design of advanced type projectiles on their best inherent capabilities, and the more critical parameters (that now constitutes the design limitation) can be eliminated by anti-gravitics. The other, which is a longer term plan, is to create an entirely new environment with longer term plan, is to create an entirely new environment with devices operating entirely under an anti-gravitic envelope.

Aviation Report 24 August 1954

THE GREATER THE EASIER

Propulsion and atomic energy trends are similar in one respect: the more incredible the long term capabilities are, the easier it is to attain them. It is strange that the greatest of nature's secrets can be harnessed with decreasing industrial effort, but greatly increasing mental effort. The Americans went through the industrial torture to produce tritium for the first thermonuclear experiment, but later both they and the Russians were able to achieve much greater results with the help of lithium 6 hydride.

The same thing is happening in aviation propulsion: the nuclear fuels are promising to be tremendously powerful in their effect, but excessively complicated in their application, unless there can be some means of direct conversion as in the strontium 90 cell. But lying behind and beyond the nuclear fuels is the linking of electricity to gravity, which is an incomparably more powerful way of harnessing energy than the only method known to human intellect at present - electricity and magnetism. Perhaps the magic of barium aluminum oxide will perform the miracle in propulsion that lithium 6 hydride has done in the fusion weapon.

Certainly it is a well-known material in dielectrics, but when one talks of massive-k, one means of course five figures. At this early stage it is difficult to relate k to Mach numbers with any certainty, but realizable k can, with some kinds of arithmetic, produce astounding velocities. They are achievable, moreover, with in terms of engineering, but the most hideous in terms of theory. Einstein's general theory of relativity is, naturally, an important factor, but some of the postulates appear to depend on the unified field theory, which cannot yet be physically checked because nobody knows how to do it. Einstein hopes to find a way of doing so before he dies.

Aviation Report 31 August 1954

GRAVITICS FORMULATIONS

All indications are that there has still been little cognizance of the potentialities of electrostatic propulsion and it will be a major undertaking to re-arrange aircraft plants to conduct large-scale research and development into novel forms of dielectric and to improve condenser efficiencies and to develop the novel type of materials used for fabrication of the primary structure. Some extremely ambitious theoretical programs have been submitted and work

towards realization of a manned vehicle has begun. On the evidence, there are far more definite indications that the incredible claims are realizable than there was, for instance, in supposing that uranium fission would result in a bomb. At least it is known, proof positive, that motion, using surprisingly low k , is possible. The fantastic control that again is feasible, has not yet been demonstrated, but there is no reason to suppose the arithmetic is faulty, especially as it has already led to quite brisk example of actual propulsion. That first movement was indeed an historic occasion, reminiscent of the momentous day at Chicago when the first pile went critical, and the phenomenon was scarcely less weird. It is difficult to imagine just where a well-organized examination into long term gravitics prospects would end. Though a circular planform is electrostatically convenient, it does not necessarily follow that the requirements of control by differential changes would be the same. Perhaps the strangest part of this whole chapter is how the public managed to foresee the concept, though not of course the theoretical principles that gave rise to it, before physical tests confirmed that the mathematics was right. It is interesting also that there is no point of contact between the conventional science of aviation and the New: it is a radical offshoot with no common principles. Aerodynamics, structures, heat engines, flapping controls, and all the rest of aviation is part of what might be called the Wright Brothers era - even the Mach 2.5 thermal barrier piercers are still Wright Brothers concepts, in the sense that they fly and they stall, and they run out of fuel after a short while, and they defy the earth's pull for a short while. Thus this century will be divided into two parts - almost to the day. The first half belonged to the Wright Brothers who foresaw nearly all the basic issues in which gravity was the bitter foe. In part of the second half, gravity will be the great provider. Electrical energy, rather irrelevant for propulsion in the first half becomes a kind of catalyst to motion in the second half of the century.

Aviation Report 7 September 1954

ELECTRO-GRAVITICS PARADOX

Realization of electro-static propulsion seems to depend on two theoretical twists and two practical ones. The two theoretical puzzles are: first, how to make a condenser the center of a propulsion system, and second is how to link the condenser system with the gravitational field. There is a third problem, but it is some way off yet, which is how to manipulate kva for control in all three axes as well as for propulsion and lift. The two practical tricks are first how, with say a Mach 3 weapon in mind, to handle 50,000 kva within the envelope

of a thin pancake of 35 feet in diameter and second how to generate such power from within so small a space. The electrical power in a small aircraft is more than in a fair sized community the analogy being that a single rocketjet can provide as much power as can be obtained from the Hoover Dam. It will naturally take as long to develop electro-static propulsion as it has taken to coax the enormous power outputs from heat engines. True there might be a flame in the electro-gravitic propulsion system, but it would not be a heat engine - the temperature of the flame would be incidental to the function of the chemical burning process.

The curious thing is that though electro-static propulsion is the antithesis of magnetism,* Einstein's unified field theory is an attempt to link gravitation with electro-magnetism. This all-embracing theory goes on logically from the general theory of relativity, that gives an ingenious geometrical interpretation of the concept of force which is mathematically consistent with gravitation but fails in the case of electro-magnetism, while the special theory of relativity is concerned with the relationship between mass and energy. The general theory of relativity fails to account for electro-magnetism because the forces are proportional to the charge and not to the mass. The unified field theory is one of a number of attempts that have been made to bridge this gap, but it is baffling to imagine how it could ever be observed. Einstein himself thinks it is virtually impossible. However Hlavaty claims now to have solved the equations by assuming that gravitation is a manifestation of electro-magnetism.

This being so it is all the more incredible that electro-static propulsion (with kva for convenience fed into the system and not self-generated) has actually been demonstrated. It may be that to apply all this very abstruse physics to aviation it will be necessary to accept that the theory is more important than this or that interpretation of it. This is how the physical constants, which are now regarded as among the most solid of achievements in modern physics, have become workable, and accepted. Certainly all normal instincts would support the Einstein series of postulations, and if this is so it is a matter of conjecture where it will lead in the long term future of the electro-gravitic science.

Aviation Report 10 September 1954

*Though in a sense this is true, it is better expressed in the body of this report than it was here in 1954.

ELECTRO-GRAVITIC PROPULSION SITUATION

Under the terms of Project Winterhaven the proposals to develop electro-gravitics to the point of realizing a Mach 3 combat type disc were not far short of the extensive effort that was planned for the Manhattan District*. Indeed the drive to develop the new prime mover is in some respects rather similar to the experiments that led to the release of nuclear energy in the sense that both involve fantastic mathematical capacity and both are sciences so new that other allied sciences cannot be of very much guide. In the past two years since the principle of motion by means of massive-k was first demonstrated on a test rig, progress has been slow. But the indications are now that the Pentagon is ready to sponsor a range of devices to help further knowledge. In effect the new family of TVs would be on the same tremendous scope that was envisaged by the X-1, 2, 3, 4 and 5 and D.558s that were all created for the purpose of destroying the sound barrier - which they effectively did, but it is a process that is taking ten solid years of hard work to complete. (Now after 7 years the X-2 has yet to start its tests and the X-3 is still in performance testing stage). Tentative targets now being set anticipate that the first disc should be complete before 1960 and it would take the whole of the 'sixties to develop it properly, even though some combat things might be available ten years from now.

One thing seems certain at this stage, that the companies likely to dominate the science will be those with the biggest computers to work out the ramifications of the basic theory. Douglas is easily the world's leader in computer capacity, followed by Lockheed and Convair. The frame incidentally is indivisible from the engine. If there is to be any division of responsibility it would be that the engine industry might become responsible for providing the electrostatic energy (by, it is thought, a kind of flame) and the frame maker for the condenser assembly which is the core of the main structure.

Aviation Report 12 October 1954

GRAVITICS STUDY WIDENING

The French are now understood to be pondering the most effective way of entering the field of electro-gravitic propulsion systems. But now least of the

"The proposals, it should be added, were not accepted.

difficulties is to know just where to begin. There are practically no patents so far that throw very much light on the mathematics of the relation between electricity and gravity. There is, of course, a large number of patents on the general subject of motion and force, and some of these may prove to have some application. There is, however, a series of working postulations embodied in the original Project Winterhaven, but no real attempt has been made in the working papers to go into the detailed engineering. All that had actually been achieved up to just under a year ago was a series of fairly accurate extrapolations from the sketchy data that has so far been actually observed. The extrapolation of 50 mph to 1,800 mph, however, (which is what the present hopes and aspirations amount to) is bound to be a rather vague exercise. This explains American private views that nothing can be reasonable expected from the science yet awhile. Meanwhile, the NACA is active, and nearly all the Universities are doing work that borders close to what is involved here, and something fruitful is likely to turn up before very long.

Aviation Report 19 October 1954

GRAVITICS STEPS

Specification writers seem to be still rather stumped to know what to ask for in the very hazy science of electro-gravitic propelled vehicles. They are at present faced with having to plan the first family of things - first of these is the most realistic type of operational test rig, and second the first type of test vehicle. In turn this would lead to sponsoring of a combat disc. The preliminary test rigs which gave only feeble propulsion have been somewhat improved, but of course the speeds reached so far are only those more associated with what is attained on the roads rather than in the air. But propulsion is now known to be possible, so it is a matter of feeding enough KVA into condensers with better k figures. 50,000 is a magic figure for the combat saucer - it is this amount of KVA and this amount of k that can be translated into Mach 3 speeds.

Meanwhile Glenn Martin now feels ready to say in public that they are examining the unified field theory to see what can be done. It would probably be truer to say that Martin and other companies are now looking for men who can make some kind of sense out of Einstein's equations. There's nobody in the air industry at present with the faintest idea of what it is all about. Also, just as necessary, companies have somehow to find administrators who know enough of the mathematics to be able to guess what kind of industrial

investment is likely to be necessary for the company to secure the most rewarding prime contracts in the new science. This again is not so easy since much of the mathematics just cannot be translated into words. You either understand the figures, or you cannot ever have it explained to you. This is rather new because even things like indeterminacy in quantum mechanics can be more or less put into words.

Perhaps the main thing for management to bear in mind in recruiting men is that essentially electro-gravitics is a branch of wave technology and much of it starts with Planck's dimensions of action, energy and time, and some of this is among the most firm and least controversial sections of modern atomic physics.

Aviation Report 19 November 1954

ELECTRO-GRAVITICS PUZZLE

Back in 1948 and 49, the public in the U.S. had a surprisingly clear idea of what a flying saucer should, or could, do. There has never at any time been any realistic explanation of what propulsion agency could make it do those things, but its ability to move within its own gravitation field was presupposed from its manoeuvrability. Yet all this was at least two years before electrostatic energy was shown to produce propulsion. It is curious that the public were so ahead of the empiricists on this occasion, and there are two possible explanations. One is that optical illusions or atmospheric phenomena offered a preconceived idea of how the ultimate aviation device ought to work. The other explanation might be that this was a recrudescence of Jung's theory of the Universal Mind which moves up and down in relation to the capabilities of the highest intellects and this may be a case of it reaching a very high peak of perception.

But for the air industries to realize and electro-gravitic aircraft means a return to basic principles in nuclear physics, and a re-examination of much in a wave technology that has hitherto been taken for granted. Anything that goes any way towards proving the unified field theory will have as great a bearing on electro-gravitics efforts as on the furtherance of nuclear power generally. But the aircraft industry might as well face up to the fact that priorities will in the end be competing with the existing nuclear science commitments. The fact that electro-gravitics has important applications other than for a weapon will however strengthen the case for governments to get in on the work going on.

Aviation Report 28 January 1955

MANAGEMENT NOTE FOR ELECTRO-GRAVITICS

The gas turbine engine produced two new companies in the U.S. engine field and they have, between them, at various times offered the traditional primes rather formidable competition. Indeed GE at this moment has, in the view of some, taken the Number Two position. In Britain no new firms managed to get a footing, but one, Metro-Vick, might have done if it had put its whole energies into the business. It is on the whole unfortunate for Britain that no bright newcomer has been able to screw up competition in the engine field as English Electric have done in the airframe business.

Unlike the turbine engine, electro-gravitics is not just a new propulsion system, it is a new mode of thought in aviation and communications, and it is something that may become all-embracing. Theoretical studies of the science unfortunately have to extend right down to the mathematics of the meson and there is no escape from that. But the relevant facts wrung from the nature of the nuclear structure will have their impact on the propulsion system, the airframe and also its guidance. The airframe, as such, would not exist, and what is now a complicated stressed structure becomes some convenient form of hard envelope.

New companies therefore who would like to see themselves as major defence prime contractors in ten or fifteen years time are the ones most likely to stimulate development. Several typical companies in Britain and the U.S. come to mind - outfits like AiResearch, Raytheon, Plessey in England, Rotax and others. But the companies have to face a decade of costly research into theoretical physics and it means a great deal of trust. Companies are mostly overloaded already and they cannot afford it, but when they sit down and think about the matter they can scarcely avoid the conclusion that they cannot afford not to be in at the beginning.

Aviation Report 8 February 1955

ELECTRO-GRAVITICS BREAKTHROUGHS

Lawrence Bell said last week he thought that the tempo of development leading to the use of nuclear fuels and anti-gravitational vehicles (he meant presumably ones that create their own gravitational field independently of the earth's) would accelerate. He added that the breakthroughs now feasible will advance their introduction ahead of the time it has taken to develop the turbojet

to its present pitch. Beyond the thermal barrier was a radiation barrier, and he might have added ozone poisoning and meteorite hazards, and beyond that again a time barrier. Time however is not a single calculable entity and Einstein has taught that an absolute barrier to aviation is the environmental barrier in which there are physical limits to any kind of movement from one point in space-time continuum to another. Bell (the company not the man) have a reputation as experimentalists and are not so earthy as some of the other U.S. companies; so while this first judgement on progress with electrogravitics is interesting, further word is awaited from the other major elements of the air business. Most of the companies are now studying several forms of propulsion without heat engines though it is early days yet to determine which method will see the light of day first. Procurement will open out because the capabilities of such aircraft are immeasurably greater than those envisaged with any known form of engine.

Aviation Report 15 July 1955

THERMONUCLEAR-ELECTROGRAVITICS INTERACTION

The point has been made that the most likely way of achieving the comparatively low fusion heat needed - 1,000,000 degrees provided it can be sustained (which it cannot be in fission for more than a microsecond or two of time) - is by use of a linear accelerator. The concentration of energy that may be obtained when accelerators are rigged in certain ways make the production of very high temperatures feasible but whether they could be concentrated enough to avoid a thermal heat problem remains to be seen. It has also been suggested that linear accelerators would be the way to develop the high electrical energies needed for creation of local gravitational systems. It is possible therefore to imagine that the central core of a future air vehicle might be a linear accelerator which would create a local weightless state by use of electrostatic energy and turn heat into energy without chemical processes for propulsion. Eventually - towards the end of this century - the linear accelerator itself would not be required and a ground generating plant would transmit the necessary energy for both purposes by wave propagation.

Aviation Report 30 August 1955

POINT ABOUT THERMONUCLEAR REACTION REACTORS

The 20 year estimate by the AEC last week that lies between present research frontiers and the fusion reactor probably refers to the time it will take to tap fusion heat. But it may be thought that rather than use the molecular and chemical processes of twisting heat into thrust it would be more appropriate to use the new heat source in conjunction with some form of nuclear thrust producer which would be in the form of electrostatic energy. The first two Boeing nuclearjet prototypes now under way are being designed to take either molecular jets or nuclear jets in case the latter are held up for one reason or another. But the change from molecular to direct nuclear thrust production in conjunction with the thermonuclear reactor is likely to make the aircraft designed around the latter a totally different breed of cat. It is also expected to take longer than two decades, though younger executives in trade might expect to live to see a prototype.

Aviation Report 14 October 1955

ELECTROGRAVITICS FEASIBILITY

Opinion on the prospects of using electrostatic energy for propulsion, and eventually for creation of a local gravitational field isolated from the earth's has naturally polarized into the two opposite extremes. There are those who say it is nonsense from start to finish, and those who are satisfied from performance already physically manifest that it is possible and will produce air vehicles with absolute capabilities and no moving parts. The feasibility of a Mach 3 fighter (the present aim in studies) is dependent on a rather large k extrapolation, considering the pair of saucers that have physically demonstrated the principle only an achieved speed of some 30 fps. But, and this is important, they have attained a working velocity using a very inefficient (even by today's knowledge) form of condenser complex. These humble beginnings are surely as hopeful as Whittle's early postulations.

It was, by the way, largely due to the early references in Aviation Report that work is gathering momentum in the U.S. Similar studies are beginning in France, and in England some men are on the job full time.

Aviation Report 15 November 1955

ELECTRO-GRAVITICS EFFORT WIDENING

Companies studying the implications of gravitics are said, in a new statement, to include Glenn Martin, Convair, Sperry-Rand, Sikorsky, Bell, Lear Inc. and Clark Electronics. Other companies who have previously evinced interest include Lockheed, Douglas and Killer. The remainder are not disinterested, but have not given public support to the new science - which is widening all the time. The approach in the U.S. is in a sense more ambitious than might have been expected. The logical approach, which has been suggested by Aviation Studies, is to concentrate on improving the output of electrostatic rigs in existence that are known to be able to provide thrust. The aim would be to concentrate on electrostatics for propulsion first and widen the practical engineering to include establishment of local gravity forcefields, independent of those of the earth's, to provide unfettered vertical movement as and when the mathematics develops.

However, the U.S. approach is rather to put money into fundamental theoretical physics of gravitation in an effort first to create the local gravitation field. Working rigs would follow in the wake of the basic discoveries. Probably the correct course would be to sponsor both approaches, and it is now time that the military stepped in with big funds. The trouble about the idealistic approach to gravity is that the aircraft companies do not have the men to conduct such work. There is every expectation in any case that the companies likely to find the answers lie outside the aviation field. These would emerge as the masters of aviation in its broadest sense.

The feeling is therefore that a company like A.T.& T. is most likely to be first in this field. This giant company (unknown in the air and weapons field) has already revolutionized modern warfare with the development of the junction transistor and is expected to find the final answers to absolute vehicle levitation. This therefore is where the bulk of the sponsoring money should go.

Aviation Report 9 December 1955

APPENDIX II

ELECTROSTATIC PATENTS

ELECTROSTATIC MOTORS

(a) American patents still in force.

2,413,391	Radio Corp.America	20-6-42/31-12-46	Power Supply System
2,417,452	Ratheon Mfg.Co.	17-1-44/18- 3-47	Electrical System
2,506,472	W.B. Smits	3-7-46Holl/ 2- 5-50	Electrical Ignition Apparatus
2,545,354	G.E.C.	16-3-50/13- 3-51	Generator
(-Engl.P. 676,953)			
2,567,373	Radio Corp.America	10-6-49/11- 9-51	El' static Generator
2,577,446	Chatham Electronics	5-8-50/ 4-12-51	El' static Voltage Generator
2,578,908	US-Atomic Energy C.	28-5-47/18-12-51	El' static Voltage Generator
2,588,513	Radio Corp.America	10-6-49/11- 3-52	El' static High-Voltage Generator
2,610,994	Chatham Electronics	1-9-50/16- 9-52	El' static Voltage Generator
2,662,181	P.Okey	31-7-52/8 -12-53	El' static Machine
2,667,615	R.G. Brown	30-1-52/26- 1-54	El' static Generator
2,671,177	Consolidated Engg.Corp	4-9-51/2 - 3-54	El' static Charging App's.
2,701,844	H.R. Wasson	8-1280/ 8- 2-55	El' static Generator of Electricity
2,702,353	US-Navy	17-7-52/15- 2-55	Miniature Printed Circuit Electrostatic Generator

(b) British patents still in force.

651,153	Metr.-Vickers Electr.Co.	26-5-48/14- .3-51	Voltage Transformation of electrical energy.
651,295	Ch.P.Warthen sr.(U.S.A.)	6-8-48/14- 3-51	Electrostatic A.C. Generator
731,774	"Licentia"	19-9-52 & 21-11-52Gy/15- 6-55	El'-static High-Voltage Generator

(c) French patents still in force

753,363	H.Chaumat	19-7-32/13-10-33	Moteur électrostatique utilisant l'énergie cinétique d'ions gazeux
749,832	H.Chaumat	24-1-33/29- 7-33	Machine électrostatique à excitation indépendante

The following patents derive from P.Jolivet (Algiers), marked "A" and from N.J.Felici, E.Gartner (Centre National des Recherches Scientifique - CRNS -) later also by R.Morel, M.Point etc.- (3. A, des Machines Electrostatiques -SAMES-) and d' Société d'Appareils de Contrôle et d'Equipment des Moteurs -SACEM-), marked "G" (because the development was centred at the University Grenoble.

Mark of Applicant	Application Date	England	America	France	Germany	Title
G	8-11-44) 14- 8-45)	637,434	2,486,140	993,017 56,027	860,649	Electrostatic Influence Machine
G	17-11-44	639,653	2,523,688	903,052	815,667	Electrostatic Influence Machine
A	28- 2-45			912,444		Inducteurs de Machines el'static
G	3- 3-45	643,660	2,519,554	905,442	882,586	El'static Machines
A	8- 6-45			915,029		Machines electrostatiques a flasques
A	16- 8-45			918,547		Generatrice el'statische.
G	20- 9-45) 21- 9-45)	643,664	2,523,689	998,397 56,356	837,267	Electrostatic Machines
A	4- 2-46			923,593		Generatrice el'statische
G	17- 7-46	643,579	2,530,193	1002,031	811,595	Generating Machines
G	20- 2-47	671,033	2,590,168			Ignition device
G	21-3 -47	655,474	2,542,494 Re-23,560	944,574	860,650	El'static Machines
G	6- 6-47	645,916	2,522,106	948,409	810,042	El'static Machines
A	16- 6-47			947,921		Génératrice él'statische.
G	16- 1-48	669,645	2,540,327	961,210	810,043	El'static Machines
G	21- 1-49	669,454	2,617,976	997,991	815,666	El'static Machines
G	7- 2-49	675,649	2,649,566	1010,924	870,575	El'static Machines
G	15- 4-49	693,914	2,604,502	1011,902	832,634	Commutators for electrical machine
G	0-11-49	680,178	2,656,502	1004,950	850,485	El'static Generate
G	9-10-50) 20- 2-51)	702,494	2,675,516	1030,623		El'static Generate
G	29-11-50) 20- 2-51)	702,421		1028,596		El'static Generate
G	21-11-51	710,687		1051,430	F10421	El'static Machines
G	20- 8-52	731,773	2,702,869		938,198	El'static Machines
G	6-11-52	745,489				El'static Generator
G	12- 2-53	745,783				Rotating El'static Machines
G	8- 1-52	715,010	2,685,654	1047,501		Rotating El'static Machines producing a periodical discharge
G	27- 2-54	Appl' n. No. 5726/55				El'static Machines
G	8- 3-54	6790/55				El'static Machines
G	28- 1-55	2748/56				El'static Machines

NOTE:- ALL THE LISTED PATENTS ARE STILL IN FORCE

THE GRAVITICS SITUATION

December 1956

Gravity Rand Ltd.
66 Sloane Street
London S W 1

Theme of the science for 1956-1970:
SERENDIPITY

Einstein's view: "It may not be an unattainable hope that some day a clearer knowledge of the processes of gravitation may be reached; and the extreme generality and detachment of the relativity theory may be illuminated by the particular study of a precise mechanism."

CONTENTS

I - Engineering note on present frontiers of knowledge

II - Management note on the gravitics situation

III - Glossary

IV - References

V - Appendix

Appendix I: Summary of Townsend Brown's original specification or an apparatus for producing force or motion

Appendix II: Mozer's quantum mechanical approach to the existence of negative mass and its utilization in the construction of gravitationally neutralized bodies

Appendix III: Gravity effects (Beams)

Appendix IV: A link between Gravitation and nuclear energy (Deser and Arnowitt)

Appendix V: Gravity/Heat interaction (Wickenden)

Appendix VI: Weight-mass anomaly (Perl)

Thanks to the Gravity Research Foundation for Appendix II to VI

I. Engineering note on present frontiers of knowledge

Gravitics is likely to follow a number of separate lines of development: the best known short term proposition is Townsend Brown's electrostatic propulsion by gravitators (details of which are to be found in the Appendix I). An extreme extrapolation of Brown's later rigs appears to suggest a Mach 3 interceptor type aircraft. Brown called this basically force and motion, but it does not appear to be the road to a gravitational shield or reflector. His is the brute force approach of concentrating high electrostatic charges along the leading edge of the periphery of a disk which yields propulsive effect. Brown originally maintained that his gravitators operate independently of all frames of references and it is motion in the absolute sense relative to the universe as a whole. There is however no evidence to support this. In the absence of any such evidence it is perhaps more convenient to think of Brown's disks as electrostatic propulsion which has its own niche in aviation. Electrostatic disks can provide lift without speed over a flat surface. This could be an important advance over all forms of airfoil which require induced flow; and lift without air flow is a development that deserves to be followed up in its own right, and one that for military purposes is already envisaged by the users as applicable to all three services. This point has been appreciated in the United States and a program in hand may now ensure that development of large sized disks will be continued. This is backed up by the U.S. Government, but it is something that will be pursued on a small scale. This acceptance follows Brown's original suggestion embodied in Project Winterhaven. Winterhaven recommended that a major effort be concentrated on electrogravitics based on the principle of his disks. The U.S. Government evaluated the disks wrongly, and misinterpreted the nature of the energy. This incorrect report was filed in an official assessment, and it took some three years to correct the earlier misconception. That brings development up to the fairly recent past, and by that time it was realized that no effort on the lines of Winterhaven was practical, and that more modest aims should be substituted. These were re-written around a new report which is apparently based on newer thoughts and with some later patents not yet published which form the basis of current U.S. policy. It is a matter of some controversy whether this research could be accelerated by more money but the impression in Gravity Rand is that the base of industry is perhaps more than adequately wide. Already companies are specializing in evolution of particular components of an electrogravitics disk. This implies that the science is in the same state as the ICBM namely

that no new breakthroughs are needed, only intensive development engineering. This may be an optimistic reading of the situation, it is true that materials are now available for the condensers giving higher k figures than were postulated in Winterhaven as necessary, and all the ingredients necessary for the disks appear to be available. But industry is still some way from having an adequate power source, and possessing any practical experience of running such equipment.

The long term development of gravity shields, absorbers, and magic metals' appears at the moment however to be a basically different problem, and work in this is not being sponsored (officially, that is), so far as is known. The absorber or shield could be intrinsically a weapon of a great power, the limits of which are difficult to foresee. The power of device to undermine the electrostatic forces holding the atom together is a destructive by-product of military significance, hi unpublished work Gravity Rand has indicated the possible effect of such a device for demolition. The likelihood of such work being sponsored in small countries outside the U.S. is slight, since there is general lack of money and resources and in all such countries quick returns are essential.

Many people hold that little or no progress can be made until the link in the Einstein unified field theory has been found. This is surely a somewhat defeatist view, because although no all-embracing explanation of the relationship between the extraordinary variety of high energy particles continually being uncovered is yet available much can be done to pin down the general nature of anti-gravity devices.

There are several promising approaches one of them is the search for negative mass, a second is to find a relationship between gravity and heat, and a third is to find the link between gravitation and the coupled particles. Taking the first of these: negative mass, the initial task is to prove the existence of negative mass, and Appendix II outlines how it might be done, this is Mozer's approach which is based on the Schroedinger time independent equation with the center of mass motion removed. As the paper shows this requires some 100 bev. which is beyond the power of existing particle accelerators: however the present Russian and American nuclear program envisage 50 bev. bevatrons within a few years and at the present rate of progress in the nuclear sciences it seems possible that the existence of negamass will be proved by this method of a Bragg analysis of the crystal structure or disproved. If negamass is established the precise part played by

the subnuclear particles could be quickly determined. Working theories have been built up to explain how negative masses would be repelled by positive masses and pairs would accelerate gaining kinetic energy until they reach the speed of light and then assume the role of the high energy particles. It has been suggested by Ferrell that this might explain the role of neutrino, but this seems unlikely without some absence of rest mass or charge of the neutrino makes it especially intriguing. Certainly further study of the neutrino would be relevant to gravitational problems. If therefore the aircraft industry regards antigravity as part of its responsibilities it cannot escape the necessity of monitoring high-energy physics of the neutrino. There are two companies definitely doing this but little or no evidence that most of the others know even what a neutrino is.

The relationship between electrical charges and gravitational forces however will depend on the right deductions being drawn from excessively small abnormalities (see Appendix VI). First clues to such small and hitherto unnoticed effects will come by study of the unified field theory: such effects may be observed in work on the gravithermals, and interacting effects of heat and gravity. This as Beams says (see Appendix III) is due to results from the alignment of the atoms. Gravity tensions applied across the ends of a tube filled with electrolyte can produce heat or be used to furnish power. The logical extension of this is an absorber of gravity in the form of a flat plate and the gravitational flux acting on it (its atomic and molecular structure, its weight density and form are not at this stage clear) would lead to an increase in heat of the mass of its surface and subsurface particles.

The third approach is to aim at discovering a connection between nuclear particles and the gravitational field. This also returns to the need for interpreting microscopic relativistic phenomena at one extreme in terms of microscopic quantum mechanical phenomena at the other. Beaumont in suggesting a solution recalls how early theory established, rough and ready assumptions of the characteristics of electron spin before the whole science of the atomic orbital was worked out. These were based on observation, and they were used with some effect at a time when data was needed. Similar assumptions of complex spin might be used to link the microscopic to the macroscopic. At any rate there are some loose ends and complex spin to be tied up, and these could logically be sponsored with some expectation of results by companies wondering how to make a contribution.

If a real spin or rotation is applied to a planar geoid the gravitational

equipotentials can be made less convex, plane or concave. These have the effect of adjusting the intensity of the gravitational field at will which is a requirement for the gravity absorber. Beaumont seemed doubtful whether external power would have to be applied to achieve this but it seems reasonable to suppose that power could be fed into the system to achieve a bénéficiai adjustment to the gravitational field, and conventional engineering methods could ensure that the weightlessness from the spin inducer. The engineering details of this naturally still in the realms of conjecture but at least it is something that could be worked out with laboratory rigs, and again the starting point is to make more accurate observations of small effects. The technique would be to accept any anomalies in nature, and from them to establish what would be needed to induce a spin artificially.

It has been argued that the scientific community faces a seemingly impossible task in attempting to alter gravity when the force is set up by a body as large as this planet and that to change it might demand a comparable force of similar planetary dimensions. It was scarcely surprising therefore that experience had shown that while it has been possible to observe the effects of gravity it resisted any form of control or manipulation. But the time is fast approaching when for the first time it will be within the capability of engineers with bevatrons to work directly wit particles that, it is increasingly accepted contribute to the source of gravitation, and whilst that in itself may not lead to an absorber of gravity, it will at least throw more light on the sources of the power.

Another task is solution (see Appendix IV) of outstanding equations to convert gravitational phenomena to nuclear energy. The problem, still not yet solved, may support the Bondi-Hoyle theory that expansion of the universe represents energy continually annihilated instead of being carried to the boundaries of the universe. This energy loss manifests itself in the behavior of the heparin and K-particles which would, or might, form the link between the microcosm and macrocosm. Indeed Deser and Arnowitt propose that the new particles are a direct link between gravitationally produced energy and nuclear energy. If this were so it would be the place to begin in the search for practical methods of gravity-manipulation. It would be realistic to assume that the K-particles are such a link. Then a possible approach might be to disregard objections which cannot be explained at this juncture until further unified field links are established. As in the case of the spin and orbital theories, which were naive in the beginning, the technique might have to accept the apparent forces and make theory fit observation

until more is known.

Some people feel that the chances of finding such a unified field theory to link gravity and electrodynamics are high yet think that the finding of a gravity shield is slight because of the size of the energy source, and because the chances of seeing unnoticed effect seem slender. Others feel the opposite and believe that a link between the Einstein general relativistic and Quantum Theory disciplines. Some hope that both discoveries may come together while a few believe that a partial explanation of both may come about the same time, which will afford sufficient knowledge of gravitational fields to perfect an interim type of absorber using field links that are available. This latter seems the more likely since it is already beginning to happen. There is not likely to be any sudden full explanation of the microcosm and macrocosm, but one strand after another joining them will be fashioned as progress is made towards quantizing the Einstein theory.

II. Management Note on the Gravitics Situation

The present anti-gravity situation as one of watching and waiting by the large aircraft prime contractors for lofting inventions or technological breakthroughs. Clarence Birdseye in one of his last utterances thought that an insulator might be discovered by accident by someone working on a quite different problem and in 500 years gravity insulators would be commonplace. One might go further than Birdseye and say that principles of the insulator would, by then, be fundamental to human affairs, it would be as basic to the society as the difference today between the weight of one metal and another. But at same time it would be wrong to infer from Birdseye a remark that a sudden isolated discovery will be the key to the science. The hardware will come at a time when the industry is ready and waiting for it. It will arrive after a long period of getting accustomed to thinking in terms of weightlessness and naturally it will appear after the feasibility of achieving it in one form or another has been established in theory. (But this does not mean that harnessed forces will be necessarily fully understood at the outset.)

The aim of companies at this stage must therefore surely be to monitor the areas of progress in the world of high energy physics which seem likely to lead to establishment of the foundations of anti-gravity. This means keeping a watchful eye on electrogravitics, magnetogravitics, gravitics-isotopes and electrostatics in various forms for propulsion or lévitation. This is not at the present stage a very expensive business, and investment in

laboratory man-hours is necessary only when a certain line of reasoning which may look promising comes to a dead-end for lack of experimental data or only when it might be worth running some laboratory tests to bridge a chasm between one part of a theory and another, or in connecting two or more theories together. If this is right, anti-gravity is in a state similar to nuclear propulsion after the NEPA findings, yet before the ANP project got under way. It will be remembered that was the period when the Atomic Energy Commission sponsored odd things here and there that needed doing. But it would be misleading to imply that hardware progress on electrostatic disks is presently so far as nuclear propulsion was in that state represented by ANP. True the NEPA-men came to the conclusion that a nuclear-propelled aircraft of a kind could be built, but it would be only a curiosity. Even at the time of the Lexington and Whitman reports it was still some way from fruition, the aircraft would have been more than a curiosity but not competitive enough to be seriously considered.

It is not in doubt that work on anti-gravity is in the realm of the longer term future. One of the tests of virility of an industry is the extent to which it is so self confident of its position that it can afford to sponsor R&D which cannot promise a quick return. A closing of minds to anything except line of development that will provide a quick return to a sign of either a straitlaced economy or of a pure lack of prescience, (or both).

Another consideration that will play its part in managerial decision is that major turning points in anti-gravity work are likely to prove far removed from the tools of the aircraft engineer. A key instrument for example that may determine the existence of negamass and establish posimass negamass interaction is the super bevatron. It needs some 100 bev gammas on hydrogen to perform a Bragg analysis of the elementary particle structure by selective reflection to prove the existence of negamass. This value is double as much the new Russian bevatron under construction and it is 15 times as powerful as the highest particle accelerations in the Berkeley bevatron so far attained. Many people think that nothing much can be done until negamass has been observed. If industry were to adopt this approach it would have a long wait and a quick answer at the end. But the negamass-posimass theory can be further developed, and, in anticipation of its existence, means of using it in a gravitationally neutralized body could be worked out. This moreover is certainly not the only possible approach a breakthrough may well come in the interaction between gravitational action and heat. Theory at the moment suggests that if gravity could produce heat the effect is limited at the moment

to a narrow range (see Appendix V). But the significant thing would be establishment of a principle.

History may repeat itself thirty years ago, and even as recently as the German attempts to produce nuclear energy in the war, nobody would have guessed that power would be unlocked by an accident at the high end of the atomic table. All prophecies of atomic energy were concerned with how quickly means of fusion could be applied at the low end. In anti-gravity work, and this goes back to Birdseye, it may be an unrelated accident that will be the means of getting into the gravitational age. It is a prime responsibility of management to be aware of possible ways of using theory to accelerate such a process. In other words, serendipity.

It is a common thought in industry to look upon the nuclear experience as a precedent for gravity, and to argue that gravitics will similarly depend on the use of giant tools, beyond the capabilities of the air industry; and that companies will edge into the gravitational age on the coat tails of the Government, as industry has done, or is doing, in nuclear physics. But this over looks the point that the two sciences are likely to be different in their investment. It will not need a place like Hanford or Savannah River to produce a gravity shield or insulator once the know how has been established. As a piece of conceptual engineering the project is probably likely to be much more like a repetition of the turbine engine. It will be simple in its essence, but the detailed componentry will become progressively more complex to interpret in the form of a stable flying platform and even more intricate when it comes to applying the underlying principles to a flexibility of operating altitude ranging from low present flight speeds at one extreme to flight in a vacuum at the other, this latter will be the extreme test of its powers. Again the principle itself will function equally in a vacuum. Townsend Brown's saucers could move in a vacuum readily enough, but the supporting parts must also work in a vacuum. In practice they tend to give trouble, just as gas turbines bits and pieces start giving trouble in proportion to the altitude gained in flight.

But one has to see this rise in complexity with performance and with altitude attainment in perspective; eventually the most advanced capability may be attained with the most extremely simple configurations. As is usual however in physics developments the shortest line of progress is a geodesic, which may in turn lead the propulsion trade into many roundabout paths as being the shortest distance between aims and achievement.

But aviation business is understandably interested in knowing precisely how to recognize early discoveries of significance, and the Gravity Rand report is intended to array and outline some of the more promising lines. One suggestion frequently made is that propulsion and levitation may be only the last though most important of a series of others, some of which will have varying degrees of gravitic element in their constitution. It may be that the first practical application will be in the greater freedom of communications offered by the change, in wave technique that it implies. A second application is to use the wave technique for anti-submarine detection, either airborne or seaborne. This would combine the width of horizon in search radar with the underwater precision of Magnetic Airborne Detection, and indeed it may have the range of scatter transmissions. Chance discoveries in the development of this equipment may lead to the formulation of new laws which would define the relationship of gravity in terms of usable propulsion symbols. Exactly how this would happen nobody yet knows; and what industry and government can do at this stage is to explore all the possible applications simultaneously, putting pressure where results seem to warrant it.

In a paper of this kind it is not easy to discuss the details of the wave technique in communications, and the following are some of the theories, briefly stated which require no mathematical training to understand, which it would be worth management keeping an eye on. In particular watch should be made of quantitative tests on lofting, and beneficiation of material. Even quite small beneficiation ratios are likely to be significant. There are some lofting claims being made of 20% and more, and the validity of these will have to be weighed carefully. Needless to say much higher ratios than this will have to be attained. New high k techniques and extreme k materials are significant. High speeds in electrostatic propulsion of small discs will be worth keeping track of (by high speed one means hundreds of rpm.h.) and some of these results are beginning to filter through for general evaluation. Weight mass anomalies, new oil-cooled cables, interesting negavolt gimmicks, novel forms of electrostatic augmentation with hydrocarbon and on hydrocarbon fuels are indicative, new patents under the broadest headings of force and motion may have value, new electrostatic generator inventions could tip the scales and unusual ways of turning condensers inside-out, new angular propulsion ideas for barycentric control; and generally certain types of saucer configuration are valuable pointers to ways minds are working.

Then there is the personnel reaction to such developments. Managements are in the hands of their technical men, and they should beware of technical teams who are dogmatic at this stage. To assert electro-gravitics is nonsense is as unreal as to say it is practically extant. Management should be careful of men in their employ with a closed mind or even partially closed mind on the subject.

This is a dangerous age when not only is anything possible, but it is possible quickly a wise Frenchman once said, "You have only to live long enough to see everything and the reverse of everything" and that is true in dealing with very advanced high energy physics of this kind. Scientists are not politicians they can reverse themselves once with acclaim, twice even with impunity. They may have to do so in the long road to attainment of this virtually perfect air vehicle. It is so easy to get bogged down with problems of the present; and whilst policy has to be made essentially with the present in mind, and in aviation a conservative policy always pays, it is management's task and duty to itself to look as far ahead as the best of its technicians in assessing the posture of the industry.

HI**GLOSSARY**

Gravithermals:	alloys which may be heated or cooled by gravity waves. (Lover's definition).
Thermistors: Electrade:	materials capable of being influenced by gravity.
Gravitator:	a plurality of cell units connected in series: negative and positive electrodes with an interposed insulating member (Townsend Brown's definition).
Lofting:	the action of lévitation where gravity's force is more than overcome by electrostatic or other propulsion.
Beneficiation:	the treatment of an alloy or substance to leave it with an improved mass-weight ratio.
Counterbary:	this, apparently, is another name for lofting.
Barycentric control:	the environment for regulation of lofting processes in a vehicle.
Modulation:	the contribution to lofting conferred on a vehicle by treatment of the substance of its construction as distinct from that added to it by outside forces. Lofting is a synthesis of intrinsic and extrinsic agencies.
Absorber; insulator:	these terms, there is no formal distinction between them as yet, are based on an analogy with electromagnetism. This is a questionable assumption since the similarity between electromagnetic and gravitational fields is valid only in some respects such as both having electric and magnetic elements. But the difference in coupling strengths, noted by many experimenters, is fundamental to the science. Gravity moreover may turn out to be the only nonquantized field in nature, which would make it basically unique.

	The borrowing of terms from the field of electromagnetism is therefore only a temporary convenience. Lack of Cartesian representation makes this a baffling science for many people.
Negamass:	proposed mass that inherently has a negative charge.
Posimass:	mass the observed quantity positively charged.
Shield:	a device which not only opposes gravity (such as an absorber) but also furnishes an essential path along which or through which, gravity can act. Thus, whereas absorbers, reflectors and insulators can provide a gravitationally neutralized body, a shield would enable a vehicle or sphere to fall away, in proportion to the quantity of shielding material.
Screening:	gravity screening was implied by Lanczos. It is the result of any combination of electric or magnetic fields in which one or both elements are not subject to varying permeability in matter.
Reflector:	a device consisting of material capable of generating buoyant forces which balance the force of attraction. The denser the material the greater the buoyancy force. When the density of the medium the result will be gravitationally neutralized. A greater density of material assumes a lofting role.
Electrogravitics:	The application of modulating influences in an electrostatic propulsion system.

Magnetogravitics:	the influence of electromagnetic and meson fields in a reflector.
Boson Fields:	these are defined as gravitational electromagnetic and meson fields (Metric tensor).
Fermion Fields:	these are electrons, neutrinos, muons, nucléons and V particles (Spinors).
Gravitator cellular body:	two or more gravitator cells connected in series within a body (Townsend Brown's definition).

Mackenzie **Physical Review**, 2 pp. 321-43.

Eotvos, Pekar and Fekete, **Annalen der Physik** 68. (1922) pp.11-16.

Heyl, Paul R. **Scientific Monthly**. 47. (1938) p. 115.

Austin, Thwing, **Physical Review** 5, (1897), pp. 494-500.

Shaw, **Nature** (April 8, 1922) p. 462. Proc. Roy. Soc. 102, (Oct. 6, 1922), p. 46.

Brush, **Physical Review**, 31, p. 1113 (A).

Wold, **Physical Review**, 35, p. 296 (Abstract).

Majorana, **Attidella Reale, Accademie die Lincei**, 28, (1919), pp. 160,221, 313,416,480,29 (1920) pp. 23, 90, 163, 235, Phil Mag. 39. (1920) p. 288.

Schneiderov, **Science**, (May 7, 1943), 97, sup. p. 10.

Brush, **Physical Review** 32, P. 633, (abstract).

Lanczos, **Science**, 74, (Dec. 4, 1931) sup. p. 10.

Eddington, **Report on the Relativity Theory of Gravitation** (1920) Fleetway Press, London.

W.D. Fowler et al, **Phys. Rev.** 93, 861, 1954.

R.L. Arnowitt and S. Deser, **Phys. Rev** 92 1061, 1953.

R.L. Amowitt Bull, **APS** 94 798. 1954 S. Deser, **Phys. Rev.** 93, 612, 1954.

N. Schein, D.M. Haskin and R.G. Classer.
Phys. Rev. 95, 855, 1954.

R.L. Arnowitt and S. Peser, unpublished Univ. of California
Radiation Laboratory Report, 1954.

H. Bondi and T. Gold, **Mon. Not. R. Astr. Soc.** 108, 252, 1948
F. Hoyle, **Mon. Not. R. Astr. Soc.** 108, 372, 1948.

B. S. De Wirt, New Directions for Research in the Theory of
Gravitation, **Essay on Gravity**, 1953.

C.H. Bondi. **Cosmology**, Cambridge University Press, 1952.

F.A.B. Pirani and A. Schild, **Physical Review** 79, 986 (1950)

Bergman, Penfield, Penfield Schiller and Zatzkis, **Phys. Rev.**
80, 81 (1950).

B.S. DeWitt. **Physical Review** 85, 653 (1952).

See, for example D. Boh, **Quantum Theory**,
New York, Prentice-Hall, Inc., (1951) Chapter 22.

A. Pais, **Proceedings of the Lorentz Kamerlingh
Onnes Conference**, Leyden, June 1953.

For the treatment of spinors in a unified field theory see W.
Pauli. **Annalen der Physik**, 18, 337 (1933) See also B.S.
DeWitt and C.M. DeWitt, **Physical Review**, 87, 116 (1952).

Carter, F.L., The Quantum Mechanical Electromagnetic Approach to
Gravity, **Essay on Gravity** 1953.

On Negative mass in the Theory of Gravitation Prof. J. M.
Luttinger **Essay on Gravity** 1951.

**SUMMARY OF TOWNSEND BROWN'S ORIGINAL
PATENT SPECIFICATION**

This invention relates to a method of controlling gravitation and for deriving power therefrom, and to a method of producing linear force or motion. The method is fundamentally electrical.

The invention also relates to machines or apparatus requiring electrical energy that control or influence the gravitational field or the energy of gravitation; also to machines or apparatus requiring electrical energy that exhibit a linear force or motion which is believed to be independent of all frames of reference save that which is at rest relative to the universe taken as a whole, and said linear force or motion is furthermore believed to have no equal and opposite reaction that can be observed by any method commonly known and accepted by the physical science to date.

Such a machine has two major parts A and B. These parts may be composed of any material capable of being charged electrically. Mass A and mass B may be termed electrodes A and B respectively. Electrode A is charged negatively with respect to electrode B, or what is substantially the same, electrode B is charged positively with respect to electrode A, or what is usually the case, electrode A has an excess of electrons while electrode B has an excess of protons.

While charged in this manner the total force of A toward B is the sum of force g (due to the gravitational field), and force e (due to the imposed electrical field) and force x (due to the resultant of the unbalanced gravitational forces caused by the electro-negative charge or by the presence of an excess of electrons of electrode A and by the electro-positive charge or by the presence of an excess of protons on electrode B).

By the cancellation of similar and opposing forces and by the addition of

similar and allied forces the two electrodes taken collectively possess a force $2x$ in the direction of B. This force $2x$, shared by both electrodes, exists as a tendency of these electrodes to move or accelerate in the direction of the force, that is, A toward B and B away from A. Moreover any machine or apparatus possessing electrodes A and B will exhibit such a lateral acceleration or motion if free to move.

In this specification, I have used terms as "gravitator cells" and gravitator cellular body" which are words of my own coining in making reference to the particular type of cell I employ in the present invention. Wherever the construction involves the use of a pair of electrodes, separated by an insulating plate or member, such construction complies with the term gravitator cells, and when two or more gravitator cells are connected in series within a body, such will fall within the meaning of gravitator cellular body.

The electrodes A and B are shown as having placed between them an insulating plate or member C of suitable material, such that the minimum number of electrons or ions may successfully penetrate it. This constitutes a cellular gravitator consisting of one gravitator cell.

It will be understood that, the cells being spaced substantial distance apart, the separation of adjacent positive and negative elements of separate cells is greater than the separation of the positive and negative elements of any cell and the materials of which the cells are formed being the more readily affected by the phenomena underlying my invention than the mere space between adjacent cells, any forces existing between positive and negative elements of adjacent cells can never become of sufficient magnitude to neutralize or balance the force created by the respective cells adjoining said spaces. The uses to which such a motor, wheel, or rotor may be put are practically limitless as can be readily understood without further description. The structure may suitably be called a gravitator motor of cellular type.

In keeping with the purpose of my invention an apparatus may employ the electrodes A and B within a vacuum tube. Electrons, ions, or thermions can migrate readily from A to B. The construction may be appropriately termed an electronic, ionic, or thermionic gravitator as the case may be.

In certain of the last named types of gravitator units, it is desirable or necessary to heat to incandescence the whole or a part of electrode A to obtain better emission of negative thermions or electrons or at least to be able to

control that emission by variation in the temperature of said electrode A. Since such variations also influence the magnitude of the longitudinal force or acceleration exhibited by the tube, it proves to be a very convenient method of varying this effect and of electrically controlling the motion of the tube. The electrode A may be heated to incandescence in any convenient way as by the ordinary methods utilizing electrical resistance or electrical induction.

Moreover, in certain types of the gravitator units, now being considered, it is advantageous or necessary also to conduct away from the anode or positive electrode B excessive heat that may be generated during the operation. Such cooling is effected externally by means of air or water cooled flanges that are in thermo connection with the anode, or it is effected internally by passing a stream of water, air, or other fluid through a hollow anode made specially for that purpose.

The gravitator motors may be supplied with the necessary electrical energy for the operation and resultant motion thereof from sources outside and independent of the motor itself. In such instances they constitute external or independently excited motors. On the other hand the motors when capable of creating sufficient power to generate by any method whatsoever all the electrical energy required therein for the operation of said motors are distinguished by being internal or self-excited. Here, it will be understood that the energy created by the operation of the motor may at times be vastly in excess of the energy required to operate the motor. In some instances the ratio may be even as high as a million to one. Inasmuch as any suitable means for supplying the necessary electrical energy, and suitable conducting means for permitting the energy generated by the motor to exert the expected influences on the same may be readily supplied, it is now deemed necessary to illustrate details herein. In said self-excited motors the energy necessary to overcome the friction or other resistance in the physical structure of the apparatus, and even to accelerate the motors against such resistance, is believed to be derived solely from the gravitational field or the energy of gravitation. Furthermore, said acceleration in the self-excited gravitator motor can be harnessed mechanically so as to produce usable energy or power, said usable energy or power, as aforesaid, being derived from or transferred by the apparatus solely from the energy of gravitation.

The gravitator motors function as a result of the mutual and unidirectional forces exerted by their charged electrodes. The direction of these forces and the resultant motion thereby produced are usually toward to positive electrode.

This movement is practically linear. It is this primary action with which I deal.

As has already been pointed out herein, there are two ways in which this primary action can accomplish mechanical work. First, by operating in a linear path as it does naturally, or second, by operating in a circular path. Since the circle is the most easily applied of all the geometric figures, it follows that the rotary form is the most important.

There are three general rules to follow in the construction of such motors. First, the insulating sheets should be as thin as possible and yet have a relatively high puncture voltage. It is advisable also to use paraffin saturated insulators on account of their high specific resistance. Second, the potential difference between any two metallic plates should be as high as possible and yet be safely under the minimum puncture voltage of the insulator. Third, there should in most cases be as many plates as possible in order that the saturation voltage of the system might be raised well above the highest voltage limit upon which the motor is operated. Reference has previously been made to the fact that in the preferred embodiment of the invention herein disclosed the movement is towards the positive electrode. However, it will be clear that motion may be had in a reverse direction determined by what I have just termed saturation voltage by which is meant the efficiency peak or maximum of action for that particular type of motor: the theory, as I may describe it, being that as the voltage is increased, the force or action increases to a maximum which represent the greatest action in a negative to positive direction. If the voltage were increased beyond that maximum the action would decrease to zero and thence to the positive to negative direction.

The rotary motor comprises broadly speaking, an assembly of a plurality of linear motors fastened to or bent around the circumference of a wheel. In that case the wheel limits the action of the linear motors to a circle, and the wheel rotates in the manner of a fireworks pin wheel.

I declare that what I claim is:

1. A method of producing force or motion which comprises the step of aggregating the predominating gravitational lateral or linear forces of positive and negative charges which are so cooperatively related as to eliminate or practically eliminate the effect of the similar and opposing forces which said charges exert.

2. A method of producing force or motion, in which a mechanical or structural part is associated with at least two electrodes or the like, of which the adjacent electrodes or the like have charges of differing characteristics, the resultant, predominating, unidirectional gravitational force of said electrodes or the like being utilized to produce linear force or motion of said part.
3. A method according to Claim 1 or 2, in which the predominating force of the charges or electrodes is due to the normal gravitational field and the imposed electrical field.
4. A method according to Claim 1, 2 or 3, in which the electrodes or other elements bearing the charges are mounted, preferably rigidly, on a body or support adapted to move or exert force in the general direction of alignment of the electrodes or other charge bearing elements.
5. A machine or apparatus for producing force or motion, which includes at least two electrodes or like elements adapted to be differently charged, so relatively arranged that they produce a combined linear force or motion in the general direction of their alignment.
6. A machine according to Claim 5 in which the electrodes or like elements are mounted, preferably rigidly on a mechanical or structural part, whereby the predominating uni-directional force obtained from the electrodes or the like is adapted to move said part or to oppose forces tending to move it counter to the direction in which it would be moved by the action of the electrodes or the like.
7. A machine according to Claim 5 or 6, in which the energy necessary for charging the electrodes or the like is obtained either from the electrodes themselves or from an independent source.
8. A machine according to Claim 5, 6 or 7, whose force action or motive power depends in part on the gravitational field or energy of gravitation which is controlled or influenced by the motion of the electrodes or the like.
9. A machine according to any of Claim 5 to 8, in the form of a motor including a gravitator cell or a gravitator cellular body, substantially as described.
10. A machine according to Claim 9, in which the gravitator cellular body or

an assembly of the gravitator cells is mounted on a wheel-like support, whereby rotation of the latter may be effected, said cells being of electronic, ionic or thermionic type.

11. A method of controlling or influencing the gravitational field or the energy of gravitation and for deriving energy or power therefrom comprising the use of at least two masses differently electrically charged, whereby the surrounding gravitational field is affected or distorted by the imposed electrical field surrounding said charged masses, resulting in a uni-directional force being exerted on the system of charged masses in the general direction of the alignment of the masses, which system when permitted to move in response to said force in the above mentioned direction derives and accumulates as the result of said movement usable energy or power from the energy of gravitation or the gravitational field which is so controlled, influenced, or distorted.
12. The method of and the machine or apparatus for producing force or motion, by electrically controlling or influencing the gravitational field or energy of gravitation.

APPENDIX II

A Quantum Approach To The Existence of Negative Mass and Its Utilization in the Construction of Gravitationally Neutralized Bodies.

Since the overwhelming majority of electrostatic quantum mechanical effects rely for their existence on an interplay of attractive and repulsive forces arising from two types of charge, few if any fruitful results could come from a quantum mechanical investigation of gravity unless there should be two types of mass. The first type, positive mass (hereafter denoted as posimass) retains all the properties attributed to ordinary mass, while the second type, negative mass (hereafter noted as negamass) differs only in that its mass is an inherently negative quantity.

By considering the quantum mechanical effects of the existence of these two types of mass, a fruitful theory of gravity will be developed. Theory will explain why negamass has never been observed, and will offer a theoretical foundation to experimental methods of detecting the existence of negamass and utilizing it in the production of gravitationally neutralized bodies.

To achieve these results recourse will be made to Schroedinger's time-independent equation with the center of mass motion removed. This equation is:

$$-\frac{h^2}{8\pi G} \nabla^2 \psi + V(\vec{r}) = E \psi$$

where all the symbols represent the conventional quantum mechanical quantities. Particular attention will be paid to the reduced mass:

$$\frac{1}{M} = \frac{1}{m_1} + \frac{1}{m_2}$$

where m_1 and m_2 are the masses of the two interacting bodies.

One can approach the first obstacle that any theory of negamass faces, namely the explanation of why negamass has never been observed, by a consideration of how material bodies would be formed if a region of empty

space were suddenly filled with many posimass and negamass quanta. To proceed along these lines, one must first understand the nature of the various possible quantum mechanical interactions of posimass and negamass.

Inserting the conventional gravitational interaction potential into Schroedinger's equation and solving for the wave function ψ , yields the result that the probability of two posimass quanta being close together is greater than the possibility of their being separated. Hence, there is said to be an attraction between pairs of posimass quanta. By a similar calculation it can be shown that while the potential form is the same, two negamass quanta repel each other. This arises from the fact that the reduced mass term in Schroedinger's equation is negative in this latter case. The type of posimass-negamass interaction is found to depend on the relative sizes of the masses of the interacting posimass and negamass quanta, being repulsive if the mass of the negamass quantum is greater in absolute value than the mass of the posimass quantum, and attractive in the opposite case. If the two masses are equal in absolute value, the reduced mass is infinite and Schroedinger's equation reduces to:

$$(Y - E)\psi = 0$$

Since the solution $\psi = 0$ is uninteresting physically, it must be concluded that $V = E$, and hence there is no kinetic energy of relative motion. Thus, highest energy state. Thus, independent of, and in addition to the attractive posimass-posimass gravitational interaction there is a repulsive quantum mechanical exchange interaction between pairs of posimass quanta, when the system is in state E_{odd} . The result of these two oppositely directed interactions is the two posimass quanta are in stable equilibrium at some separation distance. Since this equilibrium occurs between all posimass pairs in an elementary particle, a necessary consequence of the existence of negamass is that when in the first excited state elementary particles have a partial crystalline structure.

This theoretical conclusion is capable of experimental verification by performing a Bragg analysis of the elementary particle structure through shining high energy gamma rays on hydrogen. Part of the gamma ray energy will be utilized in lowering the system from energy E_{even} to E_{odd} , and if selective reflection is observed, it will constitute a strong verification of the existence of negamass. An order of magnitude calculation shows that, if the equilibrium distance between pairs of posimass quanta is one one

millionth the radius of an electron, 100 bev gamma rays will be required to perform this experiment.

Having discussed why negamass has never been observed, and having derived an experimental test of its existence, it is next desirable to develop an experimental method utilizing negamass in the production of gravitationally neutralized bodies by further consideration of some ideas previously advanced. It has been pointed out that if a source of negamass is present, a posimass sphere continues to absorb negamass quanta until equilibrium is reached as a result of the reduced mass becoming infinité. Because the sphere thus produced is practically massless, and because the gravitational interaction between two bodies is proportional to the product of their respective masses, it follows that the sphere is practically unaffected by the presence of other bodies. And thus the problem of making gravitationally neutralized bodies is reduced to the problem of procuring a source of negamass quanta. This will be the next problem discussed.

The binding energy of a negamass quantum in a posimass sphere may be obtained as one of the eigenvalue solutions to Schroedinger's Equation. If the negamass quanta in a body are excited to energies in excess of this binding energy by shining sufficient energetic gamma rays on the body, these negamass quanta will be emitted and a negamass source will thus be obtained.

To estimate the gamma ray energy required to free a negamass quantum from a posimass body, certain assumptions must be made concerning the size and mass of a posimass and negamass quanta. Since these quantities are extremely indefinite, and since the whole theory is at best qualitative, attempting to estimate the energy would be a senseless procedure. Suffice it to that because of the intimate sub-elementary nature of the posimass-negamass interaction, it seems reasonable to assume that quite energetic gamma rays will be required to break this strong bond.

To briefly review what has been shown, a quantum mechanical theory of negamass has been developed, based on the assumptions that gravitational interactions obey the laws of quantum mechanics and that all possible interactions of negamass and posimass with themselves and each other follow the well known inverse square law. This theory explains the experimental fact that negamass has never been observed, and outlines plausible experimental methods of determining the existence of negamass

and utilizing it in the construction of gravitationally neutralized bodies. While these experimental methods may perhaps be put on the realm of practicality at the present, there is every reason to hope that they will be performable in the future. At the time, the while there is an interaction potential between the equal mass posimass and negamass quanta, it results in no relative acceleration and thus, no mutual attraction or repulsion. While much could be said about the philosophical implications of the contradiction between this result and Newton's 2nd Law, such discussion is out of the scope of the present paper, and the author shall, instead, return with the above series of derivations to a consideration on the construction of material bodies in a region suddenly filled with many posimass and negamass quanta.

Because of the nature of posimass-posimass and negamass-negamass interactions, the individual posimass quanta soon combine into small posimass spheres, while nothing has, as yet, united any negamass quanta. Since it is reasonable to assume that a posimass sphere weighs more than a negamass quantum in absolute value, it will attract negamass quanta and begin to absorb them. This absorption continues until the attraction between a sphere and the free negamass quanta becomes zero due to the reduced mass becoming infinité. The reduced mass becomes infinité when the sphere absorbs enough posimass and negamass quanta equal to the negative of the mass of the next incoming negamass quantum. Thus, the theory predicts that all material bodies after absorbing as many negamass quanta as they can hold, weigh the same very small amount regardless of size.

Since this prediction is in violent disagreement with experimental fact, one must conclude that the equilibrium arising as a result of the reduced mass becoming infinite has not yet been reached. That is, assuming that negamass exists at all, there are not enough negamass quanta present in the universe to allow posimass spheres to absorb all the negamass they can hold. One is thus able to explain the experimental fact that negamass has never been observed by deriving the above mechanism in which the smaller amounts of negamass that may be present in the universe are strongly absorbed by the greater amounts of posimass, producing bodies composed of both posimass and negamass, but which have a net positive, variable, total mass.

Having thus explained why negamass has never been observed in the

pure state, it is next desirable to derive an experimental test if the existence of negamass through considering the internal quantum mechanical problem of small amounts of negamass in larger posimass spheres. One is able to gain much physical insight into this problem by simplifying it to the qualitatively similar problem of one negamass quantum in the field of two posimass quanta that are fixed distances apart. Further simplification from three dimensions to one dimension and replacement of the posimass quanta potentials by square barriers, yields a solution in which the ground state energy E_0 of the negamass quantum in the field of one posimass quantum, is split into two energy levels in the field of the two posimass quanta. These two levels correspond to even and odd parity solutions of the wave equation where E lies higher and E_{odd} lower than E_0 . The magnitudes of the differences $E_{\text{even}} - E_0$ and $E_0 - E_{\text{odd}}$ depend on the separation distance between the two posimass quanta, being zero for infinite separation and increasing as this separation distance is decreased.

Since the energy of a system involving negamass tends to a maximum in the most stable quantum mechanical configuration, the negamass quantum will normally be in state E_{even} . When the system is excited into state E_{odd} the negamass quantum will favor the situation in which the two posimass quanta are as far apart as possible, since E_{odd} increases with increasing separation distance between the two posimass quanta, and the system tends towards the plausibility of the existence of negamass and the theory behind the construction of gravitationally neutralized bodies from it, will meet their final tests.

SUMMARY PARAGRAPH

A quantum mechanical theory of negative mass is developed based on the assumption that gravitational interactions obey the laws of quantum mechanics, and that all possible interactions of negative and positive mass with themselves and each other follow the well known inverse square law. This theory explains the experimental fact that negative mass has never been observed, and outlines plausible experimental methods of determining the existence of negamass and utilizing it in the construction of gravitationally neutralized bodies.

Prof. F. Mozer

GRAVITY EFFECTS

The order of magnitude of the heat given off by an alloy as a result of the separation by gravity tension can be reliably estimated. Suppose we assume that an alloy of half tin and half lead completely fills a tube 5 meters long and 100 cm cross section which is maintained accurately at a temperature 277°C. At this temperature the alloy is liquid. Suppose next that the tube is raised from a horizontal plane into a vertical position i.e. to a position where its length is parallel to the direction of gravity. If then the alloy is free from convection as it would be if it is maintained at uniform temperature and if it is held in this position for several months, the percentage of tin at the bottom of the tube will decrease while the relative amount at the top will increase. A simple calculation shows that the concentration of tin at the top is about one tenth of one percent greater than at the bottom and that approximately one calorie of heat is given off in the separation progress. If after several months the tube is again placed so that its length is in a horizontal plane, the tin and lead will remix due to the thermal agitation of the atoms and heat is absorbed by the alloy.

Another interesting effect occurs when an electrolyte is subjected to gravity tension. Suppose a five meter glass tube is filled with a water solution of say barium chloride and the electrical potential between its ends is measured first when the length of the tube is parallel to the horizontal and second, when its length is practically zero when the tube is horizontal and approximately eighty five microvolts when it is vertical. This effect was discovered by Des Coudres in 1892. If a resistor is attached across the ends when the tube is vertical, heat of course is produced. If the tube is maintained at constant temperature the voltage decreases with time and eventually vanishes. The effect is believed to result from the fact that the positively charged barium ions settle faster than the lighter negatively charged chlorine ions as a result of gravity tension.

In conclusion we have seen that gravity tension effects an alloy in such a way that it gives off heat. This phenomenon results from the alignment of the atoms and from their separation by the gravitational field, the contribution of the latter being larger than that of the former. Also the gravity tension sets up a potential across the ends of a tube filled with an

electrolyte and this potential when applied across an external circuit may produce heat or drive an electric motor to furnish power. Several other small thermal effects possibly may arise from gravity tension in addition to those discussed above but space is not available to consider them in this essay. Also studies of the effect of gravitational fields and their equivalent centrifugal fields upon matter will no doubt be of great value in the future.

J.W. Beams

APPENDIX IV

A LINK BETWEEN GRAVITATION AND NUCLEAR ENERGY

More quantitatively we propose the following field equations describing the above phenomena:

$$-kT_{\mu\nu} = R_{\mu\nu} + \frac{1}{2}Rg_{\mu\nu} + C_{\mu\nu}[\Phi, \Psi]$$

$$\left[\frac{1}{i}\gamma^\mu \partial_\mu + m + \lambda \sigma^{\mu\nu} K_{\mu\nu}(x) \right] \Psi = 0$$

with a similar equation for Φ . In the above, Ψ represents the hyperon wave functions, and Φ the K-particle quantized field operators. The first three terms in the first equation are the usual structures in the Einstein General Relativity. The last term, $C_{\mu\nu}$, is the "creation" tensor⁸ which is to give us our conversion from gravitational to nuclear energy. It is like $T_{\mu\nu}$ in being an energy-momentum term. In the second equation ∂_μ represents the covariant derivative while γ^μ is a generalized Dirac matrix arranged so that the second equation is indeed covariant under the general group of coordinate transformations. The $\sigma^{\mu\nu}K_{\mu\nu}$ term will automatically include the higher hyperon levels. $C_{\mu\nu}$ is a functional of the hyperon and K-field variables Ψ and Φ . As can be seen these equations are coupled in two ways: first the creation term $C_{\mu\nu}$ depends upon the field variables Ψ and Φ while the gravitational metric tensor $g_{\mu\nu}$ enters through the covariant, derivative, etc. λ is a new universal constant giving the scale of the level spacings of the hyperons. Rigorously speaking the field equations should be, of course, second quantized. For purposes of obtaining a workable first approximation it is probably adequate to take expectation values and solve the semi-classical equations. The creation tensor $C_{\mu\nu}$, must be a bilinear integral of the Φ and Ψ fields and may have cross terms as well of the form $\int \Phi \bar{\Psi} \Psi dx$. These equations will indeed be difficult to solve, but upon solution will give the distribution of created energy and hence lead eventually to the more practical issues desired.

APPENDIX V

GRAVITY/HEAT INTERACTION

Let us suppose that we have to investigate the question whether gravitational action alone upon some given substance or alloy can produce heat. We do not specify its texture, density nor atomic structure. We assume simply the flux of gravitational action followed by an increase of heat in the alloy.

If we assume a small circular surface on the alloy then the gravitational flux on it may be expressed by Gauss' Theorem and it is $4\pi n M$, where M represents the mass of all sub-surface particles. The question is, can this expression be transformed into heat. We will assume it can be. Now recalling the relativity law connecting mass and energy:

$$M = m_0 + \frac{T/c^2}{c} \quad (\text{by Einstein})$$

where: T = kinetic energy

m_0 = initial mass

c = velocity of light

we set $4\pi M = m_0 + \frac{m_0 v^2}{2c^2}$.

But v^2/c^2 is a proper fraction; hence $M = m_0 + m_0/2k$

In the boundary case $v = c$, then $M = m_0 (1 + 1/k)$.

For all other cases, $4\pi M = m_0 \left(\frac{k+1}{k} \right)$ where $k = 0$

Strictly, M should be preceded by a conversion factor $1/k$ but if inserted, it does not alter the results. Thus if gravity could produce heat, the effect is limited to a narrow range, as this result shows.

It merits stress that in a gravitational field the flow lines of descent are Geodesies.

J. Wickenden

APPENDIX VI

WEIGHT-MASS ANOMALY

There is a great need for a precise experimental determination of the weight to mass ratio of protons or electrons. Since the ratio for a proton plus an electron is known already, the determination of the ratio for either particle is sufficient. The difficulty of a direct determination of the gravitational deflection of a charged particle in an experiment similar to the neutron or neutral atom experiment is due to electrical forces being much greater than gravitational field forces. Thus stray electrons or ions which are always present on the walls of an apparatus can exert sufficient force to completely mask the gravitational force. Even if the surface charges are neglected, image charges of the electron beam itself and self-repulsion in the beam may obscure at the gravitational deflection. An additional problem is the earth's magnetic field. Electrons of even a few volts energy will feel a force due to the earth's field a thousand billion times larger than the gravitational deflection. This last problem is avoided in a static measurement of the ratio such as a weighing of ionized matter. However, this last method has the additional difficulty of requiring a high proportion of ionized to unionized matter in the sample being weighed. Of course all these problems can be resolved to some extent but it is questionable if an experiment of either of the above types can be designed in which all the adverse effects can simultaneously be sufficiently minimized. Probably a completely new type of experiment will have to be devised to measure the weight to mass ratio of the proton or electron. Such a measurement may detect a deviation from the law of constant weight to mass ratio. If such an anomaly can be shown to exist, there is the possibility of finding a material which would be acted upon in an unusual manner in a gravitational field.

Martin L. Perl*

*Editor's Note: Martin L. Perl is the recipient of the 1995 Nobel Prize in physics.

**NEGATIVE MASS AS A GRAVITATIONAL
SOURCE OF ENERGY IN THE QUASI-STELLAR
RADIO SOURCES**

Banesh Hoffman

Queens College of the City University of
New York

Flushing, N.Y. 11367

Essay Awarded the First Prize
by
Gravity Research Foundation
1964

ABSTRACT

Based on the asymmetry of gravitational radiation with respect to positive and negative mass, a physical process is proposed whereby negative mass could be generated inside certain celestial objects. This process could then account for the prodigious amounts of energy- radiated by the recently discovered quasi-stellar radio sources,* it could provide an alternative to the synchrotron theory of the continuous spectrum of their emitted radiation. It could also account for the existence of cosmic ray showers of extremely high energy.

Negative mass may or may not exist. If it does, the according to both Newtonian mechanics and Einstein's general theory of relativity, it behaves in a most astonishing manner. For example, by the principle of equivalence the ratio of gravitational to inertial mass must be positive for all mass. Therefore, as is well known, positive mass attracts negative as well as positive mass, while negative mass repels both types of mass. Consequently, if a mass is placed near a mass $-m$, the two move in the same direction with ever-increasing speed, the negative mass chasing the positive. At first this seems to contradict the law of conservation of energy. But the particle of negative mass acquires negative energy as its speed increases, and the total energy- of the two particles remains constant.

So disconcerting is the behavior of negative mass that when Dirac discovered the negative-mass solutions of his relativistic equations of the electron he was dismayed. But by imagining almost all of the negative energy states filled, he was able to treat the occasional vacancies as particles of positive mass and positive charge, thus arriving at the concept of anti-matter. Anti-matter is now treated as having positive mass in its own right rather than as being an absence of negative mass.

We propose here to take the idea of negative mass seriously, a major reason for doing so being the desperate theoretical situation into which physics has been thrust by the anomalous behavior of the recently-discovered

* Now called "quasars" — Editor

quasi-stellar radio sources.¹ The idea of negative mass is extremely natural in the general theory of relativity. Indeed, one can exclude negative mass from Einstein's theory only by an ad hoc assumption extraneous to the theory.

According to Einstein's theory, gravitational waves are theoretically possible. It may well be that they are not generated by bodies in free fall; but if they exist they should be generated when matter is strongly influenced by non-gravitational forces, in the special cases that have been studied², these waves carry away energy and cause a corresponding diminution in the mass of the radiating body. But there is a peculiar asymmetry about the energy transported by the gravitational waves: the energy is positive whether the mass of the source is positive or negative. Since this apparently arises from the Minkowskian signature of space-time it would seem to be of a fundamental nature. It has significant consequences. For unlike electromagnetic waves, which do not transport charge though their source is charge, gravitational waves are produced by mass and transport mass (in the form of energy). So if a body of mass m gives off gravitational waves of energy $c^2 Am$, its mass is reduced to $m - Am$, but if the mass of the emitter is $-m$ it becomes $-(m + Am)$, which is a greater amount of negative mass than before.

Let us assume that there is a quantum law, analogous to the law of conservation of baryon number, that prevents particles of positive rest mass from decaying into particles of negative rest mass. And let us assume that under extreme and unusual conditions such as might occur inside the quasi-stellar radio sources, and only under such conditions, the law is violated, so that "forbidden" transitions can occur, these transitions being linked to the weakest known interaction, gravitation, and requiring the emission of gravitational quanta of positive energy.

Consider what might happen near the center of an extremely hot and massive celestial object. The high pressure and temperature will subject particles there to intense nuclear forces which, being nongravitational, could cause strong emission of gravitational radiation. Particles of positive rest mass emitting gravitational quanta of sufficient size will turn into particles of negative rest mass; and further emission by these particles will only increase the amount of their negative rest mass. Some of the gravitational radiation may be reabsorbed by matter of negative rest mass, but much of it will escape to the less central regions of the celestial object where conditions

do not favor such transitions, and there it will undergo "Compton" collisions, giving up energy that will ultimately be converted to electromagnetic and other radiation.

Since negative masses react perversely to all except gravitational forces, they will be less buoyant than positive masses. (I am indebted to Dr. Ivor Robinson for this remark.) But negative mass is self-repelling. So, as it accumulates centrally, it will tend to cause instability, and part of it may ultimately escape from the quasi-stellar objects in eruptions that could be the cause of the characteristic jets associated with some of these objects.

It is precisely because the quasi-stellar object emits amounts of energy so prodigious as to defy conventional explanation that the concept of negative mass deserves to be taken seriously. For when negative mass is reduced within such objects positive energy is made available, and the greater the amount of negative mass created, the greater amount of positive energy released. Nor need positive mass always be used up when such energy is released; for if a particle of negative mass is sufficiently accelerated it can radiate further gravitational quanta, thus sending out more positive energy while falling to a yet lower negative energy state. (Since the inertia increases, the process need not diverge.) The violet shift caused by the negative mass will be negligible compared with the Hubble shift. Moreover, should any negative mass escape, whether in a jet or less spectacularly, the quasi-stellar object will merely be left with greater positive energy than before. Thus the mechanism outlined here could well account for the extraordinary flux of energy from these objects.

The electromagnetic radiation ultimately generated in the charged outer layers by the gravitational radiation would have a wide range of frequencies, and this could account for the remarkable intensity observed in the radio region. (The observed polarization is small, and it neither rules out the proposed mechanism nor confirms that synchrotron radiation is a major component.)

What happens to particles of negative mass that escape? They are unlikely to be present in interstellar space insufficient amounts to affect significantly our estimates of the average density of the universe. Since they repel all matter, they cannot form negative-mass stars. If the universe is such that negative-mass particles can, on balance, "escape to infinity" there will be an effect of continual creation of positive energy in the

observed region.

Since negative mass is not antimatter, a charged particle of rest mass $-m$ meeting a particle of rest mass $+m$ and the same (not opposite) charge could form with it a stable, loosely bound system whose mass would be merely that of the electromagnetic binding energy. Such an entity would acquire enormous speeds in extended intergalactic fields too weak to dissociate it. If it dissociated due to collision near the earth the $+m$ component could cause a cosmic ray shower of enormous energy. The energy of some observed showers is puzzlingly high.

If a charged particle of negative mass were detected in, say, a cloud chamber and mistakenly interpreted as a particle of opposite charge and positive mass, there would be seeming discrepancies in such things as the energy and charge balances of reactions that might occur. It would be interesting to see if corroborative evidence for the existence of particles of negative mass could be found by the cosmic ray experimenters, for such a discovery would have momentous importance.

REFERENCES

1. Proceedings of the International Symposium on Gravitational Collapse, Dallas (to appear).
2. For an account of the work of Bondi, Brill, Weber, and Wheeler, on this point see D.R. Brill, Annals of Physics 7 (1959), p. 466. See also H. Bondi et al, Proc. Roy. Soc. 269A (1962) p. 21.

THE U.S. ANTIGRAVITY SQUADRON

by Paul A. LaViolette, Ph.D.

Abstract

Electrogravitic (antigravity) technology, under development in U.S. Air Force black R&D programs since late 1954, may now have been put to practical use in the B-2 Advanced Technology Bomber to provide an exotic auxiliary mode of propulsion. This inference is based on the recent disclosure that the B-2 charges both its wing leading edge and jet exhaust stream to a high voltage. Positive ions emitted from its wing leading edge would produce a positively charged parabolic ion sheath ahead of the craft, while negative ions injected into its exhaust stream would set up a trailing negative space charge with a potential difference in excess of 15 million volts. According to electrogravitic research carried out by physicist T. Townsend Brown, such a differential space charge would set up an artificial gravity field that would induce a reactionless force on the aircraft in the direction of the positive pole. An electrogravitic drive of this sort could allow the B-2 to function with over-unity propulsion efficiency when cruising at supersonic velocities.

For many years rumors circulated that the U. S. was secretly developing a highly advanced radar-evading aircraft. Rumor turned to reality in November of 1988, when the Air Force unveiled the B-2 Advanced Technology Bomber. Although military spokesmen provided the news media with some information about the craft's outward design and low radar and infrared profile, there was much they were silent about. However, several years later, some key secrets about the B-2 were leaked to the press. On March 9, 1992, *Aviation Week and Space Technology* magazine made the surprising disclosure that the B-2

electrostatically charges its exhaust stream and the leading edges of its wing-like body.(1) Those familiar with the electrogravitics research of American physicist T. Townsend Brown will quickly realize that this is tantamount to stating that the B-2 is able to function as an antigravity aircraft.

Aviation Week obtained their information about the B-2 from a small group of renegade West Coast scientists and engineers who were formerly associated with black research projects. In making these disclosures, these scientists broke a code of silence that rivals the Maria's. They took the risk because they felt that it was important for economic reasons that efforts be made to declassify certain black technologies for commercial use. Two of these individuals said that their civil rights had been blatantly abused (in the name of security) either to keep them quiet or to prevent them from leaving the tightly controlled black R&D community.

Several months after *Aviation Week* published the article, black world security personnel went into high gear. That sector of the black R & D community received very strong warnings and as a result, the group of scientists subsequently broke off contact with the magazine.



Figure 1. The B-2 Stealth Bomber in flight.
(U. S. Air Force photo)

Clearly, the overseers of black R & D programs were substantially concerned about the information leaks that had come out in that article.

To completely understand the significance of what was said about the B-2, one must first become familiar with Brown's work. Beginning in the mid 1920's, Townsend Brown discovered that it is possible to create an artificial gravity field by charging an electrical capacitor to a high-voltage.(2) He specially built a capacitor which utilized a heavy, high charge-accumulating (high k-factor) dielectric material between its plates and found that when charged with between 75,000 to 300,000 volts,




Figure 2. Gravitational force developed by Brown's charged high-voltage capacitor.

it would move in the direction of its positive pole (Figure 2). When oriented with its positive side up, it would proceed to lose about 1 percent of its weight.(3, 4) He attributed this motion to an electrostatically-induced gravity field acting between the capacitor's oppositely charged plates. By 1958, he had succeeded in developing a 15 inch diameter model saucer that could lift over 110 percent of its weight!(5) Brown's experiments had launched a new field of investigation which came to be known as electrogravitics, the technology of controlling gravity through the use of high-voltage electric charge.

As early as 1952, an Air Force major general witnessed a demonstration in which Brown flew a pair of 18-inch disc airfoils suspended from opposite ends of a rotatable arm. When electrified with 50,000 volts, they circuited at a

speed of 12 miles per hour.(6) About a year later, he flew a set of 3 foot diameter saucers for some Air Force officials and representatives from a number of major aircraft companies. When energized with 150,000 volts, the discs sped around the 50-foot diameter course so fast that the subject was immediately classified. *Interavia* magazine later reported that the discs would attain speeds of several hundred miles per hour when charged with several hundred thousand volts.(7)

Brown's discs were charged with a high positive voltage on a wire running along their leading edge and a high negative voltage on a wire running along their trailing edge. As the wires ionized the air around them, a dense cloud of positive ions would form ahead of the craft and corresponding cloud of negative ions would form behind the craft (Figure 3). Brown's research indicated that, like the charged plates of his capacitors, these ion clouds induced a gravitational force directed in the minus to plus direction. As the disc moved forward in response to its self-generated gravity field, it would carry with it its positive and negative ion clouds and their associated electrogravity gradient. Consequently, the discs would ride their advancing gravity wave much like surfers ride an ocean wave.




Figure 3. A side view of one of Brown's circular flying discs showing the location of its ion charges and induced gravity field.

Dr. Mason Rose, one of Townsend's colleagues, described the discs' principle of operation as follows:(8)

The saucers made by Brown have no propellers, no jets, no moving parts at all. They create a modification of the gravitational field around themselves, which is analogous to putting them on the incline of a hill. They act like a surfboard on a wave...The electro-gravitational saucer creates its own "hill," which is a local distortion of the gravitational field, then it takes this "hill" with it in any chosen direction and at any rate.

The occupants of one of [Brown's] saucers would feel no stress at all, no matter how sharp the turn or how great the acceleration. This is because the ship, the occupants and the load are all responding equally to the wavelike distortion of the local gravitational field.

Although skeptics at first thought that the discs were propelled by more mundane effects such as the pressure of negative ions striking the positive electrode, Brown later carried out vacuum chamber tests which proved that a force was present even in the absence of such ion thrust. He did not offer a theory to explain this nonconventional electrogravitic phenomenon; except to say that it was predicted neither by general relativity nor by modern theories of electromagnetism. However, recent advances in theoretical physics provide a rather straightforward explanation of the principle. According to the novel physics of subquantum kinetics, gravity potential can adopt two polarities, instead of one.(9 - 13) Not only can a gravity field exist in the form of a matter-attracting gravity potential well, as standard physics teaches, but it can also exist in the form of a matter-repelling gravity potential hill. Moreover, it predicts that these gravity polarities should be directly matched with electrical polarity: positively charged particles such as protons generating gravity wells and negatively charged particles such as electrons generating gravity hills.*

Consequently, subquantum kinetics predicts that the negative ion cloud behind Brown's disc should form a matter-repelling gravity hill while the positive ion cloud ahead of the disc should form a matter-attracting gravity well. As increasing voltage is applied to the disc, the gravity potential hill and well become increasingly prominent and the gravity potential gradient between them increasingly steep, hi Rose's terminology, the craft would find itself on the incline of a gravitational "hill." Since gravity force is known to increase in accordance with the steepness of such a gravity potential slope, increased voltage would induce an increasingly strong gravity force on the disc and

* Thus contrary to conventional theory, the electron produces a matter-repelling gravity field. Electrically neutral matter remains gravitationally attractive because the proton's G-well marginally dominates the electron's G-hill.

would act in the direction of the positive ion cloud. The disc would behave as if it were being tugged by a very strong gravitational field emanating from an invisible planet-sized mass positioned beyond its positive pole.

Early in 1952, Brown had put together a proposal, code named "Project Winterhaven," which suggested that the military develop an antigravity combat saucer with Mach 3 capability. The 1956 intelligence study entitled *Electrogravitics Systems: An Examination of Electrostatic Motion, Dynamic Counterbary and Barycentric Control*, prepared by the private aviation intelligence firm Aviation Studies (International) Ltd., indicates that as early as November 1954 the Air Force had begun plans to fund research that would accomplish Project Winterhaven's objectives.(14 - 16) The study, originally classified "confidential," mentions the names of more than ten major aircraft companies which were actively involved in electrogravitics research in an attempt to duplicate or extend Brown's seminal work. Additional information is to be found in another aviation intelligence report entitled *The Gravitics Situationen*) Since that time much of the work in electro-antigravity has proceeded in Air Force black projects on a relatively large scale.

One indication that Brown's electrogravitics ideas were being researched by the aerospace industry surfaced in January 1968. At an aerospace sciences meeting held in New York, Northrop officials reported that they were beginning wind tunnel studies to research the aerodynamic effects of applying high-voltage charges to the leading edges of aircraft bodies.(18) They said that they expected that the resulting electrical potential would ionize air molecules upwind of the aircraft and that the resulting repulsive electrical forces would condition the air stream so as to lower drag, reduce heating and soften the supersonic boom."

Although this sonic cushion effect is purely electrostatic, Northrop apparently got the idea for investigating this effect directly from Brown, for his electrokinetic flying disc patent explains that the positively charged, leading-edge electrode would produce just this effect. Brown states:(6)

By using such a nose form, which at present appears to be the best suited for flying speeds approaching or exceeding the speed of sound, I am able to produce an ionization of the atmosphere in the immediate region of this foremost portion of the mobile vehicle. I believe that this ionization facilitates piercing the sonic barrier and minimizes the abruptness with which the transition takes place in passing from subsonic velocities to

* Although the author of that article speculated that Northrop might be negatively charging the aircraft's leading edge, the sonic barrier effects can also be accomplished with a positive charge, as Brown originally suggested.

supersonic velocities.

Also, in his 1952 paper on Brown's saucers, Dr. Rose stated: (8)

The Townsend Brown experiments indicate that the positive field which is traveling in front of the saucer acts as a buffer wing which starts moving the air out of the way. This...field acts as an entering wedge which softens the supersonic barrier.

Interestingly, in 1981 the Pentagon contracted the Northrop Corporation to work on the highly classified B-2 Advanced Technology Bomber. Northrop's past experience in airframe electrostatics must have been a key factor contributing to its winning of this contract, *for Aviation Week* reported that the B-2 uses "electrostatic field-generating techniques" in its wing leading edges to help it minimize aerodynamic turbulence and thereby reduce its radar cross section.(l) The same article mentions that the B-2 also charges its jet engine exhaust stream, which has the effect of rapidly cooling its exhaust and thereby remarkably reducing its thermal signature.

Although these disclosures were framed in the context of enhancing the B-2's radar invisibility, in fact, they are part of the B-2's antigravitic drive capability. With a positively charged, wing leading edge and a negatively charged, exhaust stream (Figure 4), the B-2 would function essentially as an electrogravitic aircraft. Just as in Townsend Brown's flying discs, the positive and negative ion clouds would produce a locally altered gravity field that would cause the B-2 to feel a forward-directed gravitic force.

The design is also very similar to the saucer craft that Brown described in his electrokinetic generator patent (Figure 5).(19) The craft Brown proposed was to be powered by a flame-jet generator, a high-voltage power supply that had the advantage of being both efficient and relatively lightweight (Figure 6). His generator design utilizes a jet engine with an electrified needle mounted in the exhaust nozzle to produce negative ions in the jet's exhaust stream. The negatively ionized exhaust is then discharged through a number of nozzles at the rear of the craft. As the minus charges leave the craft in this manner, an increasingly greater potential difference develops between the jet engine body and the negatively charged exhaust cloud behind the craft. By electrically insulating the engines and *conveying* their positive charges forward to a wire running along the vehicle's leading edge, the required positively charged ion cloud is built up at the front of the vehicle. A metallic surface or wire grid positioned near the exhaust stream exit collects some of the high-voltage




Figure 4. The profile of the B-2 as seen from above. The plane measures 69 feet from front to back and 172 feet from wing tip to wing tip. Cowlings on either side of the cockpit feed large amounts of intake air to the flame-jet high-voltage generators enclosed within its body."




Figure 5. A version of the flying disc design which Brown proposed for development under Project Winterhaven.")




Figure 6. A high-voltage flame-jet generator.

electrons and this recycled power is used to run the exhaust ionizers. Brown estimated that such a generator could produce potentials as high as 15 million volts across his craft.

Rumors circulating among people close to the project allege that the B-2 does utilize antigravity technology. A similar claim was publicly made in 1990 by Bob Oechsler, an ex-NASA mission specialist.(20) So, our conjecture that the B-2 incorporates an electrogravitic drive, appears to be substantially correct. If this is true, the B-2 is the first mass-produced antigravity vehicle to be displayed to the public! It is the final realization of the kind of craft that Brown had proposed in Project Winterhaven, and that the 1956 Aviation Studies report had disclosed was beginning to be developed by the military in late 1954. The secrecy that has so tightly surrounded the B-2 most likely does not as much concern its radar-evading technology, as it does its antigravity propulsion technology, although both are probably closely intertwined. The use of such nonconventional propulsion technology would explain the B-2's high price tag, which has averaged well over a billion dollars per plane.

Although the black world scientists mentioned nothing about electrogravitics in their *Aviation Week* disclosure about the B-2, they did admit to the existence of very "dramatic, classified technologies," applicable to "aircraft control and propulsion." They were especially hesitant to discuss these projects, noting that they are "very black." One of them commented, "Besides, it would take about 20 hours to explain the principles, and very few people would understand them anyway." Apparently, what he meant is that this aircraft control and propulsion technology is based on physics principles that go beyond what is currently known and understood by most academic physicists. Indeed, by all normal standards, electrogravitics is a very exotic propulsion science.

The B-2's body design raises suspicions that the aircraft is indeed an electrogravitic vehicle. A primary design criterion for an electrogravitic craft is that it have a large horizontally disposed surface area so as to permit the development of a sufficiently strong antigravity lift force. As Townsend Brown's experiments demonstrated, such an aircraft need not necessarily be disc-shaped n triangular and square-shaped forms also exhibit antigravity lift when electrified n although disc shapes give the best performance. The triangular platforms used in the B-2 and other advanced stealth aircraft may have been deemed better for reasons of their much lower radar cross section.

Interestingly, one of the central features of the B-2's classified technology is the makeup of its hull's outer surface. Authorities tell us that the hull is composed of a highly classified, radar-absorbing material (RAM). Ceramic

dielectrics are a likely choice for the B-2 since, unlike many "lossy" dielectrics which function as radar wave absorbers, they are lossless, and hence, transparent to radar waves. More importantly, ceramic dielectrics also have the ability to store large amounts of high-voltage charge. Could the B-2's outer RAM layer be fabricated from an advanced, high-k, high-density dielectric ceramic, a material capable of exerting an enormous electrogravitic lift force when charged? Knowing that high-voltage charge is applied to the B-2's surface, the notion that its body surface is designed to function as a giant, high-voltage capacitor does not seem so far fetched. Given that the outer RAM skin is a key component of the B-2's highly secret, electrogravitic propulsion system, it is not surprising that special care would be taken to keep its composition secret.

Evidence that the B-2 might indeed use a high-density ceramic RAM comes from information leaked by the above-mentioned black world scientists, who disclosed information about the development of low-radar-observability, dielectric ceramics made from powdered depleted uranium.(1) The material is said to have approximately 92% of the bulk density of uranium, which would give it a specific gravity of about 17.5, as opposed to about 6 for barium titanate dielectrics. Thus, this new material has about three times the density of the high-k ceramics that were being tested in the 1950's, and hence would develop at least three times the electrogravitic pull.

The B-2's positively charged leading edge, another key component of its propulsion technology, was also a matter of special concern to Northrop designers. According to *Aviation Week*, the bomber's leading edges posed a particularly challenging production problem on the first aircraft. The leading-edge ionizer is most probably a conductive strip that runs along the B-2's sharp prow and is electrically charged to upwards of many millions of volts. As the B-2 moves forward, its electrified leading edge deflects the approaching air stream to either side, so that a large fraction of the generated positive ions are carried away from its body surface and are prevented from immediately contacting and neutralizing the negative ions in the B-2's exhaust stream. As a result, the B-2 is able to build up very large space charges ahead of and behind itself, which subject it to a large gravitational potential gradient. This artificially produced gravitational gradient would steepen as the B-2 attains higher speeds and deflects its positive ions outward with increasing force. Hence the B-2's electrogravitic drive would operate more efficiently when the craft was moving at higher speeds.

Best results would be obtained when the B-2 was traveling at supersonic speeds. Positive ions from its leading edge would become entrained in the upwind, sonic shock front and flow away from the craft through that sonic

boundary layer, to later converge on the negatively charged exhaust stream. Although military sources claim that the B-2 is a subsonic vehicle, in all likelihood, it is capable of supersonic flight. Probably, this capability was not disclosed in order to avoid raising curiosity about how the craft generates the required thrust.

Both in subsonic and supersonic flight, the deflected positive ions would form an ellipsoidal sheath as they circuited around the B-2 (Figure 7). The B-2's forward, positive ion sheath would act very much like an extended positively-charged electrode surface. Thus, the electrogravitic force propelling the B-2 would arise not just from the leading-edge electrode, but also from the entire positively-charged forward ion sheath. The positive and negative ion space charge distributions would very much resemble the charge configuration that Brown employed in some of his later electrogravitic experiments. Compare Figure 7, with the arcuate electrogravitic device shown in Figure 8, which is copied from one of Brown's patents.(21) Brown noted that he obtained a greater electrogravitic thrust when the positive electrode was curved and made much larger than his negative electrode. (He also stated that the apparatus produced greater thrust if the dielectric constant of its intervening dielectric rod increased along its length so as to produce a nonuniform electric field relative to the positively charged canopy. This describes precisely the characteristic of the B-2's trailing exhaust stream. At increasing distances from the craft, the negative ions progressively slow down and become more concentrated, thereby creating a nonlinear electrogravitic field along the length of the exhaust stream.) Thus, the B-2's ion sheath is optimally configured as an electrogravitic drive.

As seen in Figure 9, each of the B-2's leading edges is segmented into eight sections separated from one another by ten-centimeter wide struts. Quite possibly, the struts electrically isolate the sections from one another so that they may be individually electrified. In this way, through proper control of the applied voltage, it would be possible to gravitically steer the craft. Brown had suggested a similar idea as a way of steering his saucer craft.

The leading-edge sections positioned in front of the air scoops most likely are sparingly electrified so as to prevent positive ions from entering the engine ducts and neutralizing the negative ions being produced there.




Figure 7. A side view of the B-2 showing the shape of its electrically charged Mach 2 supersonic shock and trailing exhaust stream. Solid arrows show the direction of ion flow; dashed arrows show the direction of the gravity gradient induced around the craft.

These two nonelectrified leading-edge sections would be ideal places to mount forward-looking radar antennae since the ion plasma sheath produced by the other leading-edge sections would form a barrier that would interfere with radar signal transmission. In fact, the B-2's two Hughes Aircraft radar units are mounted precisely in these leading-edge locations: right in front of the air intakes.

The 1956 *Electrogravitics Systems* report suggested that there be a division of responsibility in the program to develop a Mach-3 electrogravitic aircraft, that the "condenser assembly which is the core of the main structure" be developed by an airframe manufacturer and that the flame-jet generator




Figure 8. An electrogravitic thrust-producing device described in one of Brown's patents.




Figure 9. A view of the B-2 with a cutaway showing the arrangement of its flame-jet generators.

which provides the electrostatic energy for the craft be developed by companies specializing in jet engine technology.(22) Consistent with that suggestion, we find that Northrop, a company experienced in aircraft electrostatics, was contracted to develop the B-2's airframe and that General Electric, a company experienced in the development of jet engines and superconducting electrical generators was contracted to develop the B-2's engines. In fact, the report mentions GE as one of the companies involved in early electrogravitics work. Also, it is known that in 1959 Townsend Brown conducted electrogravitics vacuum chamber experiments at the GE Aerospace facilities.

Authorities claim that the Stealth is powered by four General Electric F-1 18-GE-100 jet engines similar to those used in the F-16 fighter. However, the absence of a forceful exhaust thrust from the B-2, especially at take off, leads us to suspect that the B-2 does not operate on a conventional jet-thrust principle. Rather, its engines more likely are designed to function as flame-jet high-voltage generators. The propulsive force lofting the craft then, would come not so much from the jet exhaust thrust, as it would from the electrogravitic force field electrically powered by the jet's generators. Such flame-jet generators also would account for the presence of ions that *Aviation Week* says are present in the B-2's exhaust stream. As in Brown's saucer, the engine body would acquire a high positive charge as it exhausted negative ions. Presumably, the engine is electrically insulated from the aircraft hull and its positive charges are conducted forward to power the leading-edge ionizers.

The B-2's GE engines are reported to each be capable of putting out 19,000 pounds of thrust. Consequently, all four engines together should provide the B-2 with a total output of about 140,000 horsepower which translates into an electric power output of about 25 megawatts, assuming a 30 percent conversion efficiency.* By comparison, the November 1954 *Aviation Report* concluded that a 35-foot diameter electrogravitic combat disc would need to have access to about 50 megawatts of power in order to attain Mach 3 flight speeds.(22) So, it appears that the magnitude of the B-2's power output is in the right ballpark.

When the B-2 was unveiled in 1988, one Air Force official commented that the B-2 uses a system of baffles to mix cool intake air with its hot exhaust gases so as to cool the gases and thereby make them less visible to

* This horsepower estimate is based upon the assumption that the jets would be able to propel the craft to a velocity of about 600 miles per hour (Mach 0.8).

infrared-guided missiles. Although IR invisibility might be one side-benefit, most likely the real purpose for diluting the exhaust is to greatly increase the flow volume and hence the ability of the exhaust stream to eject negative charges from the craft. Much of the air entering the B-2's intake scoops probably bypasses the flame-jet intake and is mixed with the jet's hot ionized exhaust (Figure 9). Additional ions would be injected into the combined airstream by means of additional downstream electrodes. This augmented volume of ionized gases would then discharge through the two rectangular exhaust nozzles positioned near the rear of the B-2's wing. As the exhaust streams leave the nozzles they contact the titanium coated overwing exhaust ducts, portrayed in Figure 9. These open duct sections may function as rear electrical grids that collect million volt electrons from the exhaust streams and recycle them to power the exhaust and wing air ionizers. This would be done in the same fashion as Brown had suggested in his patent (Figure 6).

As the exhaust leaves the craft, it passes over trailing-edge exhaust deflectors, flaps which can be swiveled so as to direct the exhaust stream either up or down for flight control. This accomplishes more than just vectoring of the exhaust thrust; it also changes the direction of the electrogravitic force vector. When the exhaust is deflected downward, negative charges are directed below the craft. As a result, the electrogravitic force on the craft becomes vectored upward as well as forward. When the exhaust stream is deflected upward, its negative ions are directed above the craft, resulting in an electrogravitic force that is directed downward as well as forward. Thus, by controlling these flaps, the B-2 is able to control its field so as to induce either a gain or loss of altitude.

Once the B-2 was going fast enough, it would receive sufficient air flow through its scoops that it could maintain a relatively high flow rate of ionized exhaust even with its engine combustion thrust substantially reduced. Since hot exhaust is not essential to its operation, the high-voltage generator could just as well run on cool intake air with the flame jets entirely shut off. As Brown points out in his electrokinetic generator patent, "It is to be understood that any other fluid stream source might be substituted for the combustion chamber and fuel supply."(19)

In such a "coasting mode," where jet combustion is entirely shut off, the B-2 would be able to fly for an indefinitely long period of time with essentially zero fuel consumption, powering itself primarily with energy tapped from its self-generated gravitational gradient. For example, during coasting, the kinetic energy of the scooped air stream would arise entirely from the craft's own forward motion, this motion, in turn, being due to the pull of the electrogravitic propulsion field. The kinetic energy of this ionized air stream is responsible

for linearly accelerating negative ions down the B-2's exhaust ducts and hence for creating the multi-megavolt potential difference relative to the positively charged engine body. The craft's high-voltage electron collector grids (the overwing exhaust ducts) recover a portion of this power to run the craft's ionizers. Provided that this power drain is not excessive and that the plane's propulsive gravity field can be adequately maintained, the craft would be able to achieve a state of perpetual propulsion. Such perpetual motion behavior is possible in devices having the capability to manipulate their own gravity field.

It should be added here that when the B-2 flies at a sufficiently high velocity, such that the flow rate of its scooped air exceeds many times the exhaust flow rate from its jet turbines, the electrical power output of its mixed exhaust will be comparably larger, perhaps exceeding 100 megawatts.

When the B-2 was first unveiled, critics had suggested that it could not risk flying at high altitudes because it might create vapor trails that would be visible to an enemy. Edward Aldridge Jr., the secretary of the Air Force, was asked whether that problem had been solved. He replied "Yes, but we're not going to disclose how." Clearly, to explain how the B-2 could travel at high altitude with its engines essentially shut off and producing no vapor trail, he would have to disclose the vehicle's nonconventional mode of propulsion. Incidentally, in such a coasting mode, the B-2's waste heat output also would be greatly reduced, hence lessening its chance of being detected with infrared sensors.

The B-2's Emergency Power Units (EPUs) probably play a key role in assisting such high-altitude flight. According to Bill Scott,(23) each EPU consists of a small self-contained gas turbine powered by hydrazine, a liquid that rapidly decomposes into gases when activated by a catalyst. The expanding gases are made to drive a turbine which in turn drives an electrical generator. Public disclosures state that the purpose of the EPU is to supply electric power to the craft should the B-2's four jet engines happen to flame out, or its four electrical generators happen to simultaneously fail. More likely, they were designed to function as auxiliary generators capable of operating at high-altitudes (or even in space) where the air would be too thin to sustain normal jet combustion. At high-altitude the decomposed hydrazine gases would take the place of scooped air as the medium for transporting ions from the craft. That is, after passing through the EPU, these gases would be electrified and expelled from the craft in the same fashion as would the jet exhaust. Townsend Brown noted that his electrogravitic propulsion system could run just as well using a compressed gas source such as carbon dioxide as the ion carrying medium as it could using the exhaust from a jet engine.

Is all this just idle speculation? Or, could the B-2 really be the realization

of one of mankind's greatest dreams ~ an aircraft that has mastered the ability to control gravity.

NOTES

1. Scott, W. B. "Black World engineers, scientists encourage using highly classified technology for civil applications." *Aviation Week & Space Technology*, March 9, 1992, pp. 66-67.
2. Brown, T. T. "How I control gravity." *Science and Invention Magazine*, August 1929. Reprinted in *Psychic Observer* 37(1) pp. 14-18.
3. Burridge, G. "Another step toward anti-gravity." *The American Mercury* 86(6)(1958):77-82.
4. Moore, W. L., and Berlitz, C. *The Philadelphia Experiment: Project Invisibility*. New York: Fawcett Crest, 1979, Ch. 10.
5. Rho Sigma, *Ether Technology: A Rational Approach to Gravity Control* Lakemont, GA: CSA Printing & Bindery, 1977, p. 44-49, quoting a letter from T. Brown dated February 14, 1973.
6. Brown, T. T. "Electrokinetic apparatus." U.S. patent #2,949,550 (filed July 1957, issued August 1960).
7. Intel. "Towards flight without stress or strain...or weight." *Interavia Magazine* 11(5) (1956):373-374.
8. Rose, M. "The flying saucer: The application of the Biefeld-Brown effect to the solution of the problems of space navigation." *University for Social Research*, April 8, 1952.
9. LaViolette, P. A. "An introduction to subquantum kinetics: Part n. An open systems description of particles and fields." In *International Journal of General Systems, Special Issue on Systems Thinking in Physics* 11 (1985):295-328.
10. LaViolette, P. A. *Subquantum Kinetics: The Alchemy of Creation*. Schenectady, NY, 1994.
11. LaViolette, P. A. *Beyond the Big Bang: Ancient Myth and the Science of Continuous Creation*. Rochester, VT: Inner Traditions Intl., 1994.
12. LaViolette, P. A. "A theory of electrogravitics." *Electric Spacecraft Journal*, Issue 8, 1993, pp. 33 - 36.
13. LaViolette, P. A. "A Tesla wave physics for a free energy universe." *Proceedings of the 1990 International Tesla Symposium*, Colorado Springs, CO: International Tesla Society, 1991, pp. 5.1 - 5.19.
14. Aviation Studies (International) Ltd. *Electrogravitics Systems: An examination of electrostatic motion, dynamic counterbary and barycentric control* Report GRG 013/56 by Aviation Studies, Special Weapons Study

- Unit, London, February 1956. (Library of Congress No. 3,1401,00034,5879; CallNo.TL565.A9).
15. LaViolette, P. "Electrogravitics: Back to the future." *Electric Spacecraft Journal*, Issue 4, 1992, pp. 23-28.
 16. LaViolette, P. "Electrogravitics: An energy-efficient means of spacecraft propulsion." *Explore* 3 (1991): 76-79; idea No. 100159 submitted to NASA's 1990 Space Exploration Outreach Program.
 17. Aviation Studies (International) Ltd. *The Gravitics Situation*, prepared by Gravity Rand Ltd. _ a division of Aviation Studies, London, December 1956.
 18. "Northrop studying sonic boom remedy." *Aviation Week & Space Technology*, Jan. 22, 1968, p 21.
 19. Brown, T. T. "Electrokinetic generator." U.S. patent #3,022,430 (filed July 1957, issued February 1962).
 20. Rhodes, L. "Ex-NASA expert says Stealth uses parts from UFO." *Arkansas Democrat*, Little Rock, AR, April 9, 1990.
 21. Brown, T. T. "Electrokinetic apparatus." U.S. patent #3,187,206 (filed May 1958, issued June 1965).
 22. Aviation Studies, *Electrogravitics Systems*, op. cit., pp. 21-27.
 23. Scott, W.B. *Inside the Stealth Bomber*. Tab/Aero Books: New York, 1991.

Editor's Note: For those wishing to do more research in electrogravitics, the following are recommended:

The Townsend Brown Electro-Gravity Device. A comprehensive evaluation by the Office of Naval Research, with accompanying documents, issued Sept. 15, 1952. Every page was marked "confidential" until the classification of the report was cancelled by ONR. Detailed report with a surprise at the end. #612 22 pg. \$6

T. T. Brown's Electrogravitics Research, by Thomas Valone, P.E. A collection of papers including a one on electrogravitics with diagrams and references, another one with details of the Townsend Brown Notebooks, repeated Brown experiment, and one that summarizes the Bahnsen Lab film/video with still photos. #603 28 pg. \$5

SubQuantum Kinetics: The Alchemy of Creation, by Dr Paul LaViolette. A masterpiece of theoretical work predicting electrogravitic forces and more with a whole chapter devoted to electrogravitics theory. #605 208 pg. book \$15

>>>TURN TO LAST PAGE OF BOOK FOR ORDERING INFORMATION<<<

APPENDIX

A collection of electrogravitics patents' cover sheets
and relevant articles


Note: complete patents may be ordered for \$3 each from:

U. S. Patent Office
Box 9
Washington, DC 20231

Complete patents may also be viewed at www.uspto.gov



Thomas Townsend Brown



This Drawing is a Reproduction of the Original on a reduced scale]

Dated this 15th day of August, 1927.

For the Applicant,
HERBERT HADDAN & Co.,
Chartered Patent Agents,
31 and 32, Bedford Street, Strand,
London, W.C. 2.

Redhill: Printed for His Majesty's Stationery Office, by Lowe & Malcolmson, Ltd.—1928.




Fig. 8.




Fig. 10.








Fig. 10.



[This Drawing is a reproduction of the Original on a reduced scale]



PATENT SPECIFICATION



Application Date: Aug. 15, 1927. No. 27,452 / 27.

300,311

Complete Accepted: Nov. 15, 1928.

COMPLETE SPECIFICATION.

A Method of and an Apparatus or Machine for Producing Force or Motion.

I, THOMAS TOWNSEND BROWN, of 15, Eighth Street, in the City of Zanesville, State of Ohio, United States of America, a citizen of the United States of America, do hereby declare the nature of this invention and in what manner the same is to be performed, to be particularly described and ascertained in and by the following statement:

This invention relates to a method of controlling gravitation and for deriving power therefrom, and to a method of producing linear force or motion. The method is fundamentally electrical.

The invention also relates to machines or apparatus requiring electrical energy that control or influence the gravitational field or the energy of gravitation; also to machines or apparatus requiring electrical energy that exhibit a linear force or motion which is believed to be independent of all frames of reference save that which is at rest relative to the universe taken as a whole, and said linear force or motion is furthermore believed to have no equal and opposite reaction that can be observed by any method commonly known and accepted by the physical science to date.

The invention further relates to machines or apparatus that depend for their force action or motive power on the gravitational field or energy of gravitation that is being controlled or influenced as above stated; also, to machines or apparatus that depend for their force action or motive power on the linear force or motion exhibited by such machines or apparatus previously mentioned.

The invention further relates to machines and apparatus that derive usable energy or power from the gravitational field or from the energy of gravitation by suitable arrangement, using such machines and apparatus as first above stated as principal agents.

To show the universal adaptability of my novel method, said method is capable of practical performance and use in connection with motors for automobiles, space cars, ships, railway locomotion, prime movers for power installations, aeronautics. Still another field is the use

of the method and means enabling the same to function as a gravitator weight changer. Specific embodiments of the invention will be duly disclosed through the medium of the present Specification.

Referring to the accompanying drawings, forming part of this Specification:

Figure 1 is an elevation, with accompanying descriptive data, broadly illustrating the characteristic or essential elements associated with any machine or apparatus in the use of which the gravitational field or the energy of gravitation is utilized and controlled, or in the use of which linear force or motion may be produced.

Figure 2 is a similar view of negative and positive electrodes with an interposed insulating member, constituting an embodiment of the invention.

Figure 3 is a similar view of a cellular gravitator composed of a plurality of cell units connected in series, capable of use in carrying the invention into practice.

Figure 4 is an elevation of positive and negative electrodes diagrammatically depicted to indicate their relation and use when conveniently placed and disposed within a vacuum tube.

Figures 5 and 5' are longitudinal sectional views showing my gravitator units embodied in vacuum tube form wherein heating to incandescence is permitted as by electrical resistance or induction at the negative electrode; and also permitting, where desired, the conducting of excessive heat away from the anode or positive electrode by means of air or water cooling devices.

Figure 6 is an elevation of an embodiment of my invention in a rotary or wheel type of motor utilizing the cellular gravitators illustrated in Figure 3.

Figure 7 is a view similar to Figure 6 of another wheel form or rotary type of motor involving the use of the gravitator units illustrated in Figure 5, or Figure 5'.

Figure 8 is a perspective view partly in section of the cellular gravitator of Figure 3 illustrating the details thereof.

Figures 9, 10 and 10' are detail views of the cellular gravitator.

Figure 11 is a view similar to Figure 3

[Price 1/-]

with the same idea incorporated in a rotary motor.

Figures 12 and 13 are detail views thereof.

5 The general showing in Figure 1 will make clear how my method for controlling or influencing the gravitational field or the energy of gravitation, or for producing linear force or motion, is utilized by 10 any machine or apparatus having the characteristics now to be pointed out.

Such a machine has two major parts A and B. These parts may be composed of any material capable of being charged 15 electrically. Mass A and mass B may be termed electrodes A and B respectively. Electrode A is charged negatively with respect to electrode B, or what is substantially the same, electrode B is charged 20 positively with respect to electrode A, or what is usually the case, electrode A has an excess of electrons while electrode B has an excess of protons.

While charged in this manner the total 25 force of A toward B is the sum of force g (due to the normal gravitational field), and force e (due to the imposed electrical field) and force x (due to the resultant of the unbalanced gravitational forces 30 caused by the electro-negative charge or by the presence of an excess of electrons on electrode A and by the electro-positive charge or by the presence of an excess of protons on electrode B).

35 Likewise the total force of B toward A is the sum of force g (due to the normal gravitational field), and force e (due to the imposed electrical field), minus force x (due to the resultant of unbalanced 40 gravitational forces caused by the electro-negative charge or by the presence of an excess of electrons on electrode A and by the electro-positive charge or by the presence of an excess of protons on electrode 45 B).

By the cancellation of similar and opposing forces and by the addition of similar and allied forces the two electrodes taken collectively possess a force $2x$ in the direction of B. This force $2x$ shared by both electrodes exists as a tendency of these electrodes to move or accelerate in the direction of the force, that is, A toward B and B away from A. Moreover any machine or apparatus possessing electrodes A and B will exhibit such a lateral acceleration or motion if free to move. Such a motion is believed to be due to the direct control and influence 50 of the energy of gravitation by the electrical energy which exists in the unlike electrical charges present on the affected electrodes. This motion seems to possess no equal or opposite motion that is detectable by the present day mechanics.

It is to be understood that in explaining the theory underlying my invention I am imparting by best understanding of that theory, derived from practical demonstration by the use of appropriate apparatus made in keeping with the teachings of the present Specification. The practice of the method, and apparatus aiding in the performance of the method, have been successful as herein disclosed, 70 and the breadth of my invention and discovery is such as to embrace any corrected or more refined theory that may be found to underlie the phenomena which I believe myself to be the first to discover 75 and put to practical service.

In this Specification I have used terms as "gravitator cells" and "gravitator cellular body" which are words of my own coining in making reference to the particular type of cell I employ in the present invention. Wherever the construction involves the use of a pair of electrodes, separated by an insulating plate or member, such construction complies with the term gravitator cells; and when two or more gravitator cells are connected in series within a body, such will fall within the meaning of gravitator cellular body.

90 In Figure 2 the electrodes A and B are shown as having placed between them an insulating plate or member C of suitable material, such that the minimum number of electrons or ions may successfully penetrate it. This constitutes a cellular gravitator consisting of one gravitator cell.

A cellular gravitator, consisting of more than one cell, will have the cell units connected in series. This type is illustrated 105 in Figure 3, D being insulating members and E suitable conducting plates. It will be readily appreciated that many different arrangements for cell units, each possessing distinct advantages, may be resorted to.

One arrangement, such as just referred to, is illustrated in Figure 6 of the drawings. Here the cells designated F are grouped in spaced relation and placed evenly around the circumference of a wheel G. Each group of cells F possesses a linear acceleration and the wheel rotates as a result of the combined 120 forces. It will be understood that, the cells being spaced substantial distances apart, the separation of adjacent positive and negative elements of separate cells is greater than the separation of the positive 125 and negative elements of any cell, and the materials of which the cells are formed being the more readily affected by the phenomena underlying my invention than the mere space between adjacent cells, any 130

forces existing between positive and negative elements of adjacent cells can never become of sufficient magnitude to neutralize or balance the force created by the respective cells adjoining said spaces. The uses to which such a motor, wheel, or rotor may be put are practically limitless, as can be readily understood, without further description. The structure 10 may suitable be called a gravitator motor of cellular type.

In keeping with the purpose of my invention an apparatus may employ the electrodes A and B within a vacuum tube. This aspect of the invention is shown in Figures 4 and 5. In Figure 4 the electrodes A and B are such as are adapted to be placed within a vacuum tube II (Fig. 5), the frame and mounting being well within the province of the skilled artisan. Electrons, ions, or thermions 20 can migrate readily from A to B. The construction may be appropriately termed an electronic, ionic, or thermionic gravitator as the case may be.

In certain of the last named types of gravitator units, it is desirable or necessary to heat to incandescence the whole or a part of electrode A to obtain better 30 emission of negative thermions or electrons or at least to be able to control that emission by variation in the temperature of said electrode A. Since such variations also influence the magnitude of the 35 longitudinal force or acceleration exhibited by the tube, it proves to be a very convenient method of varying this effect and of electrically controlling the motion of the tube. The electrode A may 40 be heated to incandescence in any convenient way as by the ordinary methods utilizing electrical resistance or electrical induction, an instance of the former being shown at J (Fig. 5) and an instance of 45 the latter at J' (Fig. 5), the vacuum tube in Fig. 5 being designated H¹.

Moreover in certain types of the gravitator units, now being considered, it is advantageous or necessary also to conduct 50 away from the anode or positive electrode B excessive heat that may be generated during the operation of tube II or H¹. Such cooling is effected externally by means of air or water cooled flanges that 55 are in thermo connection with the anode, or it is effected internally by passing a stream of water, air, or other fluid through a hollow anode made especially for that purpose. Air cooled flanges are 60 illustrated at K (Fig. 5) and a hollow anode for the reception of a cooling liquid or fluid (as air or water) is shown at K' (Fig. 5). These electronic, ionic, or thermionic gravitator units may be 65 grouped in any form productive of a de-

sired force action or motion. One such form is the arrangement illustrated in Figures 7 where the particular gravitator units in question are indicated at L, disposed around a wheel or rotary motor 70 similarly to the arrangement of the gravitator motor of cellular type shown in Figure 6, the difference being that in Figure 7, the electronic, ionic, or thermionic gravitator units are utilized. This motor may appropriately be designated as a gravitator motor of the electronic, ionic, or thermionic type, respectively. 75

The gravitator motors of Figures 6 and 7 may be supplied with the necessary electrical energy for the operation and resultant motion thereof from sources outside and independent of the motor itself. In such instances they constitute external or independently excited motors. On the other hand the motors when capable of creating sufficient power to generate by any method whatsoever all the electrical energy required therein for the operation 80 of said motors are distinguished by being internal or self-excited. Here, it will be understood that the energy created by the operation of the motor may at times be vastly in excess of the energy required 85 to operate the motor. In some instances the ratio may be even as high as a million to one. Inasmuch as any suitable means for supplying the necessary electrical energy, and suitable conducting means 90 for permitting the energy generated by the motor to exert the expected influence on the same may be readily supplied, it is now deemed necessary to illustrate details herein. In said self-excited motors the 95 energy necessary to overcome the friction or other resistance in the physical structure of the apparatus, and even to accelerate the motor against such resistance, is believed to be derived solely from the 100 gravitational field or the energy of gravitation. Furthermore, said acceleration in the self-excited gravitator motor can be harnessed mechanically so as to produce 105 usable energy or power, said usable energy or power, as aforesaid, being derived from or transferred by the apparatus 110 solely from the energy of gravitation. 115

The gravitator motors function as a result of the mutual and unidirectional 120 forces exerted by their charged electrodes. The direction of these forces and the resultant motion thereby produced are usually toward the positive electrode. This movement is practically linear. It 125 is this primary action with which I deal.

As has already been pointed out herein, there are two ways in which this primary action can accomplish mechanical work. First, by operating in a linear path as it 130

does naturally, or second, by operating in a curved path. Since the circle is the most easily applied of all the geometric figures, it follows that the rotary form is the important. While other forms may be built it has been considered necessary to explain and illustrate only the linear and rotary forms.

The linear form of cellular gravitator is illustrated in detail in Figures 8, 9 and 10. It is built up of a number of metallic plates alternated or staggered with sheets of insulating material (Fig. 3). Each pair of plates so separated by insulation act as one gravitator cell, and each plate exhibits the desired force laterally. The potential is applied on the end plates and the potential difference is divided equally among the cells. Each metallic plate in the system possesses a force usually toward the positively charged terminus, and the system as a whole moves or tends to move in that direction. It is a linear motor, and the line of its action is parallel to the line of its electrodes.

There are three general rules to follow in the construction of such motors. First, the insulating sheets should be as thin as possible and yet have a relatively high puncture voltage. It is advisable also to use paraffin-saturated insulators on account of their high specific resistance. Second, the potential difference between any two metallic plates should be as high as possible and yet be safely under the minimum puncture voltage of the insulator. Third, there should, in most cases, be as many plates as possible in order that the saturation voltage of the system might be raised well above the highest voltage limit upon which the motor is operated. Reference has previously been made to the fact that in the preferred embodiment of the invention herein disclosed the movement is towards the positive electrode. However, it will be clear that motion may be had in a reverse direction determined by what I have just termed "saturation voltage", by which is meant the efficiency peak or maximum of action for that particular type of motor; the theory, as I may describe it, being that as the voltage is increased the force or action increases to a maximum which represents the greatest action in a negative-to-positive direction. If the voltage were increased beyond that maximum the action would decrease to zero and thence to the positive-to-negative direction.

Referring more specifically to Figs. 8, 9, and 10, red fiber end plates 1 act as supports and end insulators, and the first metallic plate 2 (for example aluminum) is connected electrically, through the fiber

end plate, with the terminal 5. The second insulating sheet 3 is composed, for example, of varnished cambric sometimes known as "empire cloth". The relative size and arrangement of the metallic plate and insulating sheets are best seen in Figures 9 and 10. A paraffin filler 6 is placed between adjacent insulating sheets and around the edges of the metallic plates (Fig. 10a) and 6 represents a thin paraffin coating over the whole motor proper. 7 and 8 indicate successive layers of "empire cloth" or similar material, and 9 is a binding tape thereon. A thin film of a substance such as black spirit varnish 10 protects and insulates the entire outer surface. A phosphor bronze safety gap element 11 is connected electrically with the terminal (not shown) opposite to the terminal 5. A safety gap element corresponding with the element 11 is electrically connected with the terminal 5, but it has not been shown, in order better to illustrate interior parts. The purpose of the safety gaps is to limit the voltage imposed on the motor to the predetermined maximum and to prevent puncture.

The rotary motor (Figs. 11, 12 and 13), comprises broadly speaking, an assembly of a plurality of linear motors, fastened to or bent around the circumference of a wheel. In that case the wheel limits the action of the linear motors to a circle, and the wheel rotates in the manner of a fireworks pin wheel. The illustrations I have given are typical. The forms of Figures 6 and 7 have been defined. In Figure 11, the insulating end disk 1a has an opening 2a therethrough for an extension of the shaft 12. The disk 1a is secured to a suitable insulating motor shell, by fiber bolts or screws in any convenient manner, there being another of these disks at the opposite end of the shell, in the same manner as the opposite end plates 1 in Figure 8. The cells are built upon an insulating tube 11a disposed about the shaft-space 3a. Thick insulating wedges 4a separate the four linear motors illustrated. These thick insulating wedges, so-called, are substantially greater in body than the aggregate insulating sheets of the units. In some instances, however, dependent upon materials employed for the charged elements and the insulating members, this need not necessarily be the case. In each motor of this circular series of motors, there are the alternate sheets of insulation 5a associated with the alternate metallic plates 6a; paraffin fillers 7a along the edges of the plates 6a and between the insulating sheets 5a being employed similarly to the use of paraffin in Figure 8. 130

The rotary motor is encircled by metallic (preferably copper) collector rings 10a, which are connected with the end metallic plates of the separate linear motors at 9a and 13 (Fig. 12), one of these connections 9 being shown in detail where the insulating tube is cut away at 8 (Fig. 11).

It is unnecessary herein to illustrate a housing or bearings because any insulated housing and good ball bearings, conveniently supplied, will complete the motor. The potential is applied to the safety gap mounted on the housing and thence is conducted to the collector rings 10 of the motor by means of sliding brushes.

While I have in the foregoing Specification outlined, in connection with the broader aspects of my invention, certain forms and details, I desire it understood that specific details have been referred to for the purpose of imparting a full and clear understanding of the invention, and not for purposes of limitation, because it should be apparent that many changes in construction and arrangement, and many embodiments of the invention, other than those illustrated, are possible without departing from the spirit of the invention or the scope of the appended claims.

Having now particularly described and ascertained the nature of my said invention and in what manner the same is to be performed, I declare that what I claim is:—

1. A method of producing force or motion, which comprises the step of aggregating the predominating gravitational lateral or linear forces of positive and negative charges which are so co-operatively related as to eliminate or practically eliminate the effect of the similar and opposing forces which said charges exert.

2. A method of producing force or motion, in which a mechanical or structural part is associated with at least two electrodes or the like, of which the adjacent electrodes or the like have charges of differing characteristics, the resultant, predominating, uni-directional gravitational force of said electrodes or the like being utilized to produce linear force or motion of said part.

3. A method according to Claim 1 or 2, in which the predominating force of the charges or electrodes is due to the normal gravitational field and the imposed electrical field.

4. A method according to Claim 1, 2 or 3, in which the electrodes or other elements bearing the charges are mounted, preferably rigidly, on a body or support adapted to move or exert force in the general direction of alignment of the electrodes or other charge-bearing elements.

5. A machine or apparatus for producing force or motion, which includes at least two electrodes or like elements adapted to be differently charged, so relatively arranged that they produce a combined linear force or motion in the general direction of their alignment.

6. A machine according to Claim 5, in which the electrodes or like elements are mounted, preferably rigidly, on a mechanical or structural part, whereby the predominating uni-directional force obtained from the electrodes or the like is adapted to move said part or to oppose forces tending to move it counter to the direction in which it would be moved by the action of the electrodes or the like.

7. A machine according to Claim 5 or 6, in which the energy necessary for charging the electrodes or the like is obtained either from the electrodes themselves or from an independent source.

8. A machine according to Claim 5, 6 or 7, whose force action or motive power depends in part on the gravitational field or energy of gravitation, which is controlled or influenced by the action of the electrodes or the like.

9. A machine according to any of Claims 5 to 8, in the form of a motor including a gravitator cell or a gravitator cellular body, substantially as described.

10. A machine according to Claim 9, in which the gravitator cellular body or an assembly of the gravitator cells is mounted on a wheel-like support, whereby rotation of the latter may be effected, said cells being of electronic, ionic or thermionic-type.

11. A method of controlling or influencing the gravitational field or the energy of gravitation and for deriving energy or power therefrom comprising the use of at least two masses differently electrically charged, whereby the surrounding gravitational field is affected or distorted by the imposed electrical field surrounding said charged masses, resulting in a uni-directional force being exerted on the system of charged masses in the general direction of the alignment of the masses, which system when permitted to move in response to said force in the above mentioned direction derives and accumulates as the result of said movement usable energy or power from the energy of gravitation or the gravitational field which is so controlled, influenced, or distorted.

12. The method of and the machine or apparatus for producing force or motion, by electrically controlling or influencing the gravitational field or energy of gravitation, substantially as hereinbefore described with reference to the accompanying drawings.

Sept. 25, 1934.






T. T. BROWN

1,974,483

ELECTROSTATIC MOTOR

Filed Feb. 7, 1930

2 Sheets-Sheet 1



Inventor
Thomas T. Brown

284 *Milam & Milam*
Attorneys

Aug. 16, 1960

T. T. BROWN

2,949,550

ELECTROKINETIC APPARATUS

Filed July 3, 1957

2 Sheets-Sheet 1

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5



INVENTOR
THOMAS TOWNSEND BROWN

BY

Watson, Cole, Kindle & Watson

ATTORNEYS

Jan. 23, 1962

T. T. BROWN

3,018,394

ELECTROKINETIC TRANSDUCER

Filed July 3, 1957

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5



INVENTOR
THOMAS TOWNSEND BROWN
BY
Watson, Cole, Grindle & Watson
ATTORNEYS

Feb. 20, 1962

T. T. BROWN

3,022,430

ELECTROKINETIC GENERATOR

Filed July 3, 1957

2 Sheets-Sheet 1

FIG. 1




FIG. 2





FIG. 3



INVENTOR
THOMAS TOWNSEND BROWN
BY

Staten, Lea, Hendry & Staten
ATTORNEYS

June 1, 1965




T. T. BROWN

3,187,206

ELECTROKINETIC APPARATUS

Filed May 9, 1958

2 Sheets-Sheet 1



INVENTOR
Thomas T. Brown
BY

Hatton, Cole, Henkle & Hatton
Attorneys

Jan. 3, 1967

T. T. BROWN

3,296,491

METHOD AND APPARATUS FOR PRODUCING IONS AND
ELECTRICALLY-CHARGED AEROSOLS

Filed Sept. 19, 1961

3 Sheets-Sheet 1




FIG. 1




FIG. 2




FIG. 3b




FIG. 3a




FIG. 5




FIG. 4




FIG. 6

INVENTOR

THOMAS TOWNSEND BROWN

BY Watson, Cole, Grindle & Watson
ATTORNEYS

June 30, 1970

T. T. BROWN

3,518,462

FLUID FLOW CONTROL SYSTEM

Filed Aug. 21, 1967

2 Sheets-Sheet 1




Fig. 1






Fig. 2

A.C.
SIGNAL
INPUT

INVENTOR.

THOMAS TOWNSEND BROWN

BY Fulwider, Patton, Bailey,
Lee, and Utche

ATTORNEY

DATE-EVENT CHART FOR T.T. BROWN

- 1905 March 18, IT Brown born in Zanesville, Ohio.
- 1921 TTB discovers motion of the Coolidge X-ray tube caused by high voltage.
- 1922 Attended Cal-Tech; left school and joined the Navy.
- 1923 Attended Kenyon College, Ohio.
- 1924 Attended Denison University, Gambier, Ohio where he met Prof. Biefeld (Ph.D., Zurich, Switzerland, 1900).
- 1926 Joined Swazey Observatory in Ohio, directed by Prof. Paul Alfred Biefeld.
- 1928 Maximum thrust measured on "Gravitor," 1% of total weight.
- 1929 Wrote article for *Science and Inventions*, "How I Control Gravitation."
- 1930 Left Swazey Observatory, signed on to Naval Research Lab, Washington, DC.
- 1932 U.S. Navy Department, staff physicist, International Gravity Expedition to West Indies.
- 1933 Staff physicist, Johnson-Smithsonian Deep Sea Expedition; joined Naval Reserve; soil engineer for Federal Emergency Relief Administration; joined FCC as an administrator.
- 1939 Lieutenant in Navy Reserve, moved to Maryland as a materials engineer for Glen L. Martin (Aircraft) Co., called into Navy Bureau of Ships as an officer in magnetic and acoustic mine research and development
- 1940 May have joined the "Philadelphia Experiment."
- 1942 Lt. Commander, Commanding Officer of U.S. Navy Radar School at Norfolk, VA.
- 1943 Collapsed from exhaustion, retired from Navy, 6 months recuperation. Gravitor not recognized.
- 1944 Position with Lockheed-Vega, radar consultant
- 1945 Went to Hawaii, continued research: (Presented ideas to Admiral Arthur W. Radford, Commander in Chief of U.S. Pacific Fleet, who later became Joint Chief of Staff/Eisenhower, 1953-57. No apparent interest was shown).
- 1952 Started Winterhaven Project in Cleveland, Ohio.
Set up Townsend Brown Foundation office in Menlo Park, CA and demonstrated "flying saucer"-shaped disks electrically propelled around a "maypole."
- 1955-56 Presentations in England of his ideas.
- 1955-57 Leesburg, VA from Oct. 1955 to Feb. 1957; Umatillo, FL, Nov. 1957.
- 1956 Project work in France with his electrogravitation ideas for SNCASCO. Brown formed UFO study group, NICAP, in Washington, DC.
- 1957-60 Hired as consultant for Whitehall-Rand Project under auspices of Bahnsen Labs (Agnew Bahnsen) to do antigravity research and development
- 1958 Brown formed Rand International, Ltd.
- 1958-67 Brown indicated he made no notes in his notebooks.
- 1959 Consultant with GJE. Space Center at King of Prussia, PA.
- 1960s Physicist for Electro Kinetics at Bala Cynwyd, PA. Also semi-retired.
- 1967 T.T. Brown Notebook: October 23, 1967 in Santa Monica, CA.
- 1970 T.T. Brown Notebook: May 31, 1970 at Stanford University Hospital
- 1973 T.T. Brown Notebook: indicates presence on Catalina Island, March 1973-Sep. 1973.
- 1975 Honolulu, Hawaii working on "rock electricity."
- 1976-77 Sunnyvale, CA and University of Calif/Berkeley: rock electricity work.
- 1979(7) Consultant at Stanford Research Institute.
- 1980 University of North Carolina at Chapel Hill, Project Coordinator for XERXES Project
- 1983 Project XERXES continued at California State University (Los Angeles); Brown residing at Avalon, California, as his health was more troublesome. ; •'» >
- 1985 Brown died at Avalon.

T.T. Brown Experiment Replicated

Larry Deavenport

Larry Deavenport is a hobbyist in electronics who attended the Texas State Tech Institute. He spent 20 years in maintenance at a copper refining plant

The first time I recall hearing T. Townsend Brown's name was in a grade school science class during the early 1960s. At that time, NASA was in the process of developing ion and plasma rockets.

Like Nikola Tesla before him, Brown was a pioneer. At an early age, he was to make a remarkable discovery that would lead to the establishment of a field of research called electrogravitics.

While still a student at Denison University in Granville, Ohio, Brown noticed movement from wires of opposite polarity which would come in close proximity to one another. These wires had a tendency to attract while in the presence of very high voltage. Brown reported his observation to his physics professor, Paul Biefeld, who encouraged him to do further experiments. The research that followed led to the Biefeld-Brown effect and six U.S. patents for Brown.

In one of his experiments, Brown mounted condenser plates on a rotor. When voltage was applied, the rotor turned in a set direction. If the polarity of the voltage was reversed, the rotor would spin in the opposite direction.

The experiment which resulted in U.S. Patent #2,949,550, is perhaps Brown's most well-known. A large disk-shaped negative plate follows a positively charged wire (electrode) separated by a dielectric insulator. When two of these disks are suspended from a rotor mechanism and high voltage is applied, movement follows which is in the direction of the positive electrode, as illustrated in Figure 1.




Fig. 1 T.T. Brown's "electrokinetic apparatus." Figure 6 from U.S. Patent #2,949,550, granted Aug. 16, 1960.

To order any of the items mentioned in the article, contact Bob Iannini at Information Unlimited, P.O. Box 716, Amherst, NH 03031-0716. Tel 603-673-4730.

The IU catalog is \$1. The plan is part GRA1, and sells for \$15. The kit is part GRA1K, and sells for \$99.50. The 12V, 3amp power supply is part GRA10, and is made and priced to order.

The replication

During the summer of 1994, I decided to replicate Brown's rotor experiment. I machined two $2\frac{3}{4}$ -inch by $\frac{1}{8}$ -inch thick aluminum disks, with a tapered edge.

Using a 16-inch by 2-inch wide piece of balsa wood, I built a crude, but workable, rotor to suspend the disks. Small pieces of #36 steel wire, separated by insulators, were attached to the rotor as condensers. A strip of $\frac{3}{8}$ -inch copper was formed into a ring and glued to a $1\frac{1}{2}$ -inch by 18-inch tall PVC support for brush contact. The rotor was then suspended on top of the support with a glass door roller used for the bearing. The bearing was later replaced with a needle spindle because the original caused drag problems by creating too much surface area, which prevented the rotor from turning.

The power supply was a 0-15 volt input oscillator with a

25kV, 800 microamp output at 25kHz pulsed DC. The circuit diagram is illustrated in Figure 2.

When I applied the voltage, the disks behaved as if they wanted to turn, but they did not. I made several adjustments between the positive and negative electrodes of the disk, but still there was no rotation.

At the same time, I was experimenting with other power supplies. In one of these other trials, a 7000VAC, 5 milliamp transformer with six 10,000V capacitors and six 20,000V, 2 watt diodes were used. The six capacitors and six diodes were soldered together to form a Cockcroft-Walton generator. This was set with spark gaps between them, and run in parallel with the original power supply. When the power supplies were turned on, the higher current transformer with the Cockcroft-Walton DC ladder began to amplify the action of

the lesser powered 25kV, 25Hz pulsed DC supply. The aluminum disks began to resonate with every aluminum box in the room. The disks behaved as if they wanted to move up and down, but the rotor did not turn.

It was several months before I tried any more experiments. Early this year, I was discussing the Brown experiment and the problems I was having with a friend. I showed him the rotor and mentioned the problem of having too much bearing surface drag from too large a bearing. We agreed that the bearing should be smaller, and decided to try a needle or small spindle. The changes were made, and on the first weekend in February, I started experimenting again. The new setup is shown in figures 3 and 4.

This time, the small rotor turned more freely; however, it needed to be balanced. Bearing surface drag and rotor balance will greatly affect the outcome of the experiment. Also very important is the size of the rotor bearing and how well the unit is balanced in conjunction with the size of the powersupply.

The rotor began to turn slowly at first, at about 15 rpm. Several days later, I repeated some experiments with the rotor, and found that a ground connection was loose. When this was corrected, the rotor speed increased to about 35 rpm.

During this time, I contacted Dr. Paul LaViolette who suggested making the center of the disks more blunt instead of tapered. On my new 4-inch disks, shown in Figure 5, I took this into consideration.

One of the later experiments with the disks used a much larger power supply-0-50 volts DC, 0-15 amp input. The voltage was increased to





Fig. 3. The author's "magnetopole" rotor device.




Fig. 4. Rotor illustration and detail.




Fig. 4
Enlarged view of the disk electrode.

various levels and special notice was taken as to what happened on the high voltage end. The original power supply or oscillator was rated for 6-15 volts with a 25 kHz DC pulsed output. It was originally purchased about 10 years ago for the Hagen device (U.S. Patent #3,120,363). In this experiment, a platform about 12 inches in diameter is negatively charged, while a smaller circular platform containing very small wires is positively charged. The slow moving mass of air and charged ions are supposed to generate lift. Regardless, this power supply worked well on Brown's experiment.

I also contacted Bob Laninini, the owner of Information Unlimited and the manufacturer of this device. He first thought that due to the age of the power supply, a pulsed rippling effect might be generated from the capacitor. Since I do not own an oscilloscope, and have only a linear frequency counter built into my meter, there is no proof that this may have happened. However, after conducting tests using other capacitors (a 63 volt, 100,000 μ F and a 63 volt, 15,000 μ F) in place of the original 20 volt, 10,000 μ F capacitor, I ruled out the rippling effect. If there had been a pulsed ripple produced from the capacitor, it would not have been present in a larger capacitor.

In the higher voltage experiments, the speed of the rotor increased significantly, from 35 rpm to about 60 rpm. This is after the power supply had reached 24 volts, according to the VOM reading. I checked with an analog meter for verification of input. This would have doubled the output of the high voltage to 50,000 volts and the current to about 1.5 millamps. The rotor turned at a maximum speed of 60 rpm.

Several factors were responsible for the first breakdown between electrodes on the disks. High voltage power transistors had reached their maximum wattage, causing overheating, and the needle bearing on the rotor prevented the rotor from reaching high speeds.

Another experiment involved adding the Crockett-Walton voltage ladder to the high voltage terminal of the power supply. This


stepped up the voltage, but dropped the microamp rating down. The rotor turned at about half the speed it normally turned, at 25kV. This indicates that voltage and current rise must remain proportional to maintain the effect.

An analysis of Brown's experiments

- 1 Voltage and amperage must be raised proportionally to maintain the electrogravitic effect.
- 2 As the voltage and current are raised, dielectric strength (K) or spacing between positive and negative electrodes must be increased.
- 3 Power or frequency of charging time must be increased proportionally as the size of the area is increased.
- 4 Changing the size of either electrode will affect the efficiency of ionic flow, if the dielectric and power source remain constant. The ratio in the size of the positive to the negative plate is directly proportional to the amount of voltage and current applied through a dielectric medium. With a smaller, high voltage low microamp power supply, the electrode on the positive side may have to be larger. On a higher voltage milliamps system, the positive electrode may need to be smaller.
- 5 Geometric configurations of shape may enhance or diminish the efficiency of the electromagnetic effect.
- 6 The electromagnetic effect increases in proportion to voltage, amperage, dielectric strength (the K factor) and area, when all of these increase together. It is lost when an arcover or breakdown occurs.
- 7 Some writers have, in the past, insinuated that electronic levitation is a product of very high frequencies and high voltages. Although a significant power increase may be present in this state, my experiments with the Biefeld-Brown effect show that it is determined by the conditions thus described. Certainly, frequency, time lag and phase angle, along with all the formulas dealing with resonance would affect the efficiency of an AC system if it were used, but frequency and high voltage by themselves do not produce this effect.

In conclusion, I believe that the electrogravitic force is ever present, be it at one volt, one amp, one ohm, or 50,000 volts and 1 milliamp with a high dielectric strength. It is simply more noticeable at the high voltage electrostatic state, which is the opposite of the electromagnetic state.

Contact the author, Larry Deavenport, at 1823 Lawson Lane, Amarillo, TX 79106.



Disc speed v ([]) and propulsion efficiency s (o) as a function of average DC voltage V plotted logarithmically from ONR test data. Filled points are data from a separate test carried out by T. T. Brown. (*Subquantum Kinetics*, LaViolette, p. 173)

The Antigravity Research of T. Townsend Brown

Many readers may recall the name of I. Townsend Brown (mentioned briefly in the previous article) as it was historically associated with UFOs. Brown was the first director of the now-defunct—but once very active—National Investigations Committee on Aerial Phenomena (NICAP), a Washington, D.C.-based lobby group that was very instrumental in the 1960s for getting congressional leaders interested in the UFO mystery. But what the majority may not realize was that Brown was also a very highly acclaimed physicist who held many patents and worked on a variety of highly classified projects.

It may well have been that Brown served as a very important "link" between the UFOlogical community and work being done in the 40s and 50s on what in scientific circles is best known as "Electrogravitics


Systems." There are even some who suggest that Brown's work in this field may have been assisted by his knowledge of the crash landing of a spaceship near Roswell, New Mexico. Because of his close dealings with the government and many private aerospace firms (both in the United States and abroad) he would have easily had access to knowledge considered to be on the "cutting edge."

According to Tomas Valpro, a licensed professional engineer and head of the Integrity Research Institute (1377 K Street, N.W., Washington, D.C. 20005), some of Brown's early patterns showed designs that look very much like the familiar "flying saucer" seen throughout North America during this time period, in fact, in the group's recently published report titled *Electrogravitics Systems—Reports on a New*


Propulsion Methodology, Valone sees a very strong tie between Brown's discoveries and the development of the B-2 Stealth Bomber.

«There is," he says, "substantial evidence that the electrogravitics research of the 1950s actually resulted in the B-2 Stealth Bomber auxiliary propulsion system." In the report, an article by Dr. Paul LaViolette is summarized further reach these conclusions:

1. The B-2 charges the leading edges of its wing-like body, with high voltage;
2. The B-2 is shaped just like IT. Brown suggested an electro-gravitic craft should look, for maximum charge separation;
3. Northrup tested leading-edge charging in 1968;
4. T.T. Brown suggested that the craft should be powered by a flame-jet generator like the



A version of the flying disc design that Brown proposed for development under Project Winterhaven.



© 1993 Paul LaViolette

A high-voltage flame-jet generator.

- B-2 engine.
- 5. Aviation Week admits the existence of "dramatic, classified technologies" applicable to "aircraft control and propulsion" on the B-2;
- 6. Aviation Week also disclosed that the ceramic RAM on the B-2 outer skin is powdered depleted uranium, which just happens to have a dielectric constant of three times that of the high-K dielectrics tested in the 1950s (barium titanate oxide);
- 7. The B-2's Emergency Power Units (EPU) can work at high altitudes or even in space, driving an electrical generator;
- 8. Edward Aldridge, the Secretary of the Air Force, admits that the B-2 creates no vapor trail at high altitudes.
- 9. The decomposed gases from

the EPU's function as the ion-carrying medium, according to T.T. Brown.


From this point it is not hard to reach out even further and safely conclude that a great deal of what is going on inside Area 51 (Nevada's top secret military base) is based upon the work of Brown and others who long ago saw another way to reach the stars besides utilizing rockets that need very heavy fuel payloads.

After reading the available literature, even seasoned scientists have had to agree that Brown's propulsion methods may be our best bet in reaching out to the planets in our solar system that we might possibly be able to visit even now if these discoveries were not being kept under lock and key by those few who would deem it necessary to utilize this valuable knowledge for their

own private gain.

John Searl's Work

British engineer John Searl claims to have built over 50 versions of his own "flying saucer," and has been receiving media attention—no matter how limited—for a number of years with his claims of having developed a noiseless, propulsion-free, levity disc that requires no fuel and flies vertically when taking off and landing. Searl even makes the bold claim that one of his craft actually flew around the world several times in the early 1970s...apparently without detection. It's claimed that "ultra-high voltage electro-static force fields are developed by segmented rings rotating in the disc's periphery. The craft's direction is controlled by varying voltage around the edge, thus producing unbelievable speed and agility."



The Searl Levity Disk...a man-made Flying Saucer goes aloft.

Et si le B-2 n'était pas seulement un avion ?

Selon des révélations dans les milieux de la recherche et de la construction aéronautique aux Etats-Unis, le bombardier « furtif » de Northrop expérimenterait des techniques d'« électrogravité ».

Si un avion est, par définition, un appareil de navigation aérienne plus lourd que l'air se déplaçant dans l'atmosphère en obéissant aux lois de l'aérodynamique, alors le bombardier stratégique américain B-2A *Spirit* construit par Northrop, vedette furtive du dernier Salon du Bourget, est bien plus qu'un avion. Et il apparaîtrait, dans cette logique, à aucune famille officiellement connue de moyens de transport aérien.

Pour mieux comprendre la nature des révélations qui filtrent actuellement sur les étranges particularités de cette machine volante, un détour par les coulisses de l'histoire s'impose.

Tandis que le B-2 était officiellement présenté au public, en novembre 1988, l'US Air Force déclarait que les technologies développées pour le mettre au point « allaient révolutionner l'industrie aérospatiale américaine ». Cette phrase prophétique passa inaperçue.

BLACK PROGRAMME

Trois ans plus tard, William B. Scott, rédacteur en chef de l'hebdomadaire *Aviation Week & Space Technology* était approché par un groupe d'ingénieurs travaillant de longue date sur des « programmes noirs » du Pentagone. Décidés à accélérer le processus de déclassification des technologies dont les civils méritaient, à



B-2 présente des particularités qui lui valent le surnom de « savonnette volante ».

(Photo AFP)

leurs yeux, de bénéficier, ces « renégats » lui dévoilèrent informellement quelques-uns

Thèse hardie

Alexander Szames est un jeune journaliste scientifique feu de technologies « exotiques », c'est-à-dire de celles qui deviennent, parfois, la normalité de demain après avoir paru « impossibles » au moment où elles ont commencé à faire parler d'elles. Sa thèse sur le bombardier B-2, fondée sur des études et des révélations anglo-saxonnes, est pour le moins hardie. Une hypothèse prémature ? Pas étrange, cependant, à des recherches qui sont effectivement menées sur de nouveaux modes de propulsion. La réponse appartient à l'avenir.

*L'Alsace, 10 Mars 1996, p 6
Mulhouse, France*

des plus noirs secrets de la chauve-souris géante. Dont Scott se fit l'écho : le B-2 charge d'électricité positive ses bords d'attaque, un dispositif spécial lui permet d'électriser négativement les gaz brûlés en sortie de tuyère et sa cellule est tapissée d'un matériau diélectrique (non conducteur d'électricité) à base de poudre d'uranium apparu capable d'absorber les ondes radio-

Personne ne fut choqué autre mesure par ces révélations. Et pourtant, ce qui semble être l'un des plus grands secrets technologiques de la guerre froide venait de tomber... dans le domaine public.

Le B-2 est bien plus furtif qu'il n'y paraît. En électrisant ses bords d'attaque, il laisse l'air silencieux et crée un « parapluie » de plasma modelé sur sa propre onde de choc. Ce phénomène se matérialise sous forme d'un halo lumineux

de couleur rose-orange. Ses bénéfices sont immédiats : le plasma agit comme un bouclier anti-radar (il emprisonne une bonne partie des ondes électromagnétiques) et rend plus fluide l'atmosphère alentour. Le B-2 devient une véritable savonnette volante.

CANONS À IONS

Les moteurs du bombardier sont tout aussi exotiques. Conçus par General Electric (GE), ils servent effectivement à « pousser » l'appareil. Mais au décollage seulement. En régime de croisière, ils changent de fonction et font plutôt office de... canons à ions négatifs. Pourquoi ? Pour mieux diluer les gaz chauds dans l'atmosphère, c'est-à-dire réduire la signature infra-rouge ? Cette explication semble logique, mais n'est pas forcément la seule valable. « Bien des aspects du processus de furtivité (du B-2) restent classifiés », indique l'US Air Force.

apparent d'antigravité), étudiée avec rage depuis le milieu des années 50 dans la quasi-totalité des établissements de recherche et développement occidentaux. Ce dont témoigne un rapport sur les « Systèmes électrogravifiques », récemment déterré par le Dr. LaViolette et établi en 1955 à l'initiative d'une société de consultation britannique, Aviation Studies.

ARCHIVES SECRÈTES

Si le B-2 ne constitue qu'un malgré aspect des percées technologiques accomplies pendant la guerre froide, un pan entier de l'histoire du XX^e siècle reste à découvrir. Peut-être en saurons-nous plus en octobre prochain, lorsque le Pentagone rendra publiques, comme prévu par le décret d'*Aviation Week*, le B-2 est par nature un condensateur : ses bords d'attaque lui servent d'anode (électrode positive), ses flux de combustion de cathode aérodynamique (électrode négative) et sa cellule de diélectrique.

Quant au B-2, ce fascinant appareil appartient déjà à la préhistoire. En dévoilant, dans *New World Vista*, sa dernière étude scientifique, l'US Air Force prévoyait l'arrivée rapide d'avions-caméléons. Électriques, certes... mais totalement invisibles. Y compris à l'œil nu !

Alexandre SZAMES

« Sous-litre et intérêts sont de la rédaction.

Book Review

Electrogravitics Systems: Reports on a New Propulsion Methodology

THOMAS VALONE, ED.

REVIEWED BY LESLEE KULBA

The authors who contributed to *Electrogravitics Systems: Reports on a New Propulsion Methodology* assert that electrogravitics is not a dead-end field of research.

The 111-page book presents information indicating that anti-gravity has been and is being seriously investigated by leading aircraft industries as well as governments. An underlying theme is that T.T. Brown propulsion, once developed, will usher in an age of flight so revolutionary it will make all previous aviation, from the Wright brothers to space shuttles, constitute the Stone Age of flight.

This book can be appreciated by anyone who is interested in electrogravitics. It contains basic information for the neophyte (such as glossaries, patent lists and basics on T.T. Brown research), as well as clippings and information which make a case for the reality of electrogravitics technology.

Structurally, the book consists of four papers: "Electrogravitics Systems," from Aviation Studies, Ltd.; "The Gravitics Situation," from Gravity Rand Ltd., Div. of Aviation Studies, Ltd.; "Negative Mass As a Gravitational Source of Energy in the Quasi-Stellar Radio Sources," by Banesh Hoffman; and "The U.S. Antigravity Squadron," by


Paul LaViolette. All papers encourage serious research.

The book is thought-provoking. "What is gravity?" it asks; a question Newton and Einstein did not satisfactorily explain. Experts have told us: "It just is."

Electrogravitics Systems looks beyond, and asks how fast is gravity if not an instantaneous action-at-a-distance. Does it travel in waves? Does gravitational force, like other forces, generate heat? If so, how much?

The book makes some interesting points. For example, gravity may be the only force that is not quantized. It also states that all fundamental forces (e.g., electricity and magnetism) appear to exist in the form of opposites (poles). So, why shouldn't gravity have its opposite—negative gravity? Since gravitational force is dependent only on two or more masses and the absolute value of the distances separating them, the quest for antigravity then takes on the form of a quest for antimatter, not an uncommon task for modern physicists.

Questions are raised throughout the book, and solutions remain distant, though it is pointed out that the most satisfactory theo-



ries usually are simplistic in explanation and application.

Having made a theoretical case for electrogravitics, the book also makes a historical one. Hints of electrogravitics in the history of aviation, revealed through developments and statements made by major aircraft industries in articles from *Aviation Report* in the mid-1950s, are reprinted. T.T. Brown's work is described in detail.

With optimism, the book repeatedly states that the breakthroughs in antigravity devices are almost guaranteed to come from big industry which can pour grand sums of money into extensive research and development facilities.

The paper by Paul LaViolette was an intriguing speculation that the B-2 stealth bomber operated on T.T. Brown's principle of propulsion. Statements from government and ex-government workers and officials were shown to fit in nicely with this possibility. LaViolette argues that several disclosed as well as probable technological details of this classified design are consistent with design specifications for a would-be T.T. Brown aircraft.

Electrogravitics Systems was published by the Integrity Research Institute, 1377 K Street NW, Suite 204, Washington, D.C. 20005. It retails for \$15.

PUBLICATIONS available from Integrity Research Institute

T. T. Brown's Electrogravitics Research, by Thomas Valone, P.E. A collection of papers including a one on electrogravitics with diagrams and references, another one with details of the Townsend Brown Notebooks, repeated Brown experiment, and one that summarizes the Bahnsen Lab film/video with still photos. #603 28 pg. \$5

The Townsend Brown Electro-Gravity Device. A comprehensive evaluation by the Office of Naval Research, with accompanying documents, issued Sept. 15, 1952. Every page was marked "confidential" until the classification of the report was cancelled by ONR. Detailed report with a surprise at the end. #612 22 pg. \$6

Electrogravitics Radio Interview. Just before the 1994 T. T. Brown Symposium in Philadelphia, 21st Century's Radio Hieronymus & Co. conducted a radio interview with Valone about the speakers. Topics covered are energy, propulsion, electrogravitics and current research in the field. #604 1 hour audiotape \$5

Thomas Townsend Brown: Bahnsen Lab 1958-1960. This rare find, supplied by Sev Bonnie, a friend of the Bahnsen family, is a documentary recorded originally in 8 mm film. Though silent, it is in color and reveals short segments of all of the Brown-Bahnsen laboratory experiments. Specially designed craft levitate with electrogravitic propulsion. #606 approx. 1 hour NTSC VMS videotape \$20

SubQuantum Kinetics: The Alchemy of Creation, by Dr. Paul LaViolette. A masterpiece of theoretical work predicting electrogravitic forces and more. An entire chapter is devoted to electrogravitics. Topics include the creation of matter, fields, and forces based on systems science. #605 208 pg. book \$15

Free Energy and Antigravity Propulsion. Hosted by Tom Valone. Over the past few decades, there has been a considerable production of new and exiting energy and propulsion inventions which are emerging. A multimedia presentation to a conference audience in Mt. Shasta, CA. #607 1/2 hour NTSC VMS video \$25

Future Energy: Proceedings of the First International Conference on Future Energy. Papers from the 1999 three-day COFE symposium featuring more than 14 scientists, most with visual presentations of their breakthroughs in energy. Also includes invited papers. FREE upon request with #802 purchase: Talking CD-ROM with 20 hours of digital audio lectures (usually \$49). #802 250 pg. book \$40

Call, write, or email for a complete newsletter/catalog on Future Energy!

ORDERING INFORMATION:

Shipping cost is \$2 for U.S. residents. Add \$3 for Canada. Add \$5 for postage overseas. Send check, money order or charge card number to Integrity Research Institute, 1220 L St. NW Suite 100-232, Washington, DC 20005. Call 800-295-7674 or 202-452-7674 for catalog or to place an order. Email: iri@erols.com Visit: www.integrity-research.org

The Huntfor Zero Point

The Hunt for Zero Point

*Inside the Classified World of
Antigravity Technology*



Nick Cook

BROADWAY BOOKS

New York



THE HUNT FOR ZERO POINT. Copyright © 2001 by High Frontier Productions, Ltd. All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval system, without written permission from the publisher.

For information, address Broadway Books, a division of Random House, Inc.,
1540 Broadway, New York, NY 10036.

A British edition of this book was published in 2001 by Century, The Random House Group Limited, 20 Vauxhall Bridge Road, London, SW1V 2SA, United Kingdom.

Broadway Books titles may be purchased for business or promotional use or for special sales. For information, please write to: Special Markets Department, Random House, Inc.,
1540 Broadway, New York, NY 10036.

PRINTED IN THE UNITED STATES OF AMERICA.

BROADWAY BOOKS and its logo, a letter B bisected on the diagonal, are trademarks of Broadway Books, a division of Random House, Inc.

Visit our website at www.broadwaybooks.com

First Broadway Books edition published 2002

Library of Congress Cataloging-in-Publication Data
Cook, Nick.

The hunt for zero point: inside the classified world of antigravity technology /

Nick Cook.— 1st ed.

p. cm.

Includes bibliographical references and index.

1. Research aircraft—History. 2. Antigravity—Research—History. 3. Aeronautics, Military—Research—History. 4. Military weapons—Research—History. 5. Defense information, Classified—History. I. Title.

TL567.R47 .C 2002

629.13'07'2073—dc21

2001052668

FIRST EDITION

ISBN 0-7679-0627-6

1 3 5 7 9 1 0 8 6 4 2

In memory of Julian Cook, inventor, and Harry Hawker, fighter pilot

Per ardua ad astra

And for my children, Lucy and William, that one day they or their
children may see the stars more closely

Author's Note and Acknowledgments

Not everyone who has assisted me in *The Hunt For Zero Point* has wished to be identified.

There are four people—"Amelia Lopez," "Lawrence Cross," "Daniella Abelman" and "Dr. Dan Marckus"—whose identities I have deliberately blurred. They, of course, know who they are and I gratefully acknowledge their help.

It goes without saying that I could not have interpreted the mass of raw science data in this book without the exceptional skills of Dan Marckus. His knowledge, wisdom, humor and friendship have been an inspiration to me.

Everyone else is exactly as identified in the text.

The people and institutions I can thank freely are listed below:

My colleagues at *Jane's Defence Weekly* and *Interavia*, Tom Valone of the Integrity Research Institute, Paul LaViolette Ph.D., Dr. Kathleen Bunten, Isabel Best, Joel Carpenter, the Imperial War Museum, Philip Henshall, Graham Ennis, Rowland White, Guy Norris, Jeremy Bartlett, Michael Collins, Guy Rigby, Kevin Bexley, Stephen Parker, Bill Zuk, Yvonne Kinkaid, the Office of (U.S.) Air Force History, David Windle, Chris Gibson, Tony Frost, the late Prof. Brian Young, Dr. Ron Evans, BAE Systems, Garry Lyles, George Schmidt, Marc Millis, the U.S. National Aeronautics and Space Administration, Lockheed Martin and the Lockheed Martin Skunk Works, Dr. Evgeny Podkletnov, Dr. Hal Puthoff, the late Ben Rich, Bill Sweetman, Glenn Campbell, Mark Farmer and the Groom Lake Interceptors, Mike Grigsby, James Cook, David Monaghan, George Muellner, Marian Swedija, Irving Baum, Ken Jones, Charles T. Schaeffer, the 16th Armored Division veterans' association, Dr. Amy Schmidt, the U.S. National Archives Records Administration, Ned Bedessem, U.S. Army Center of Military History, the Bundesarchiv at Freibourg, the U.K. Public Records Office, Maurizio Verga, James Holland, Callum Coats, Don Kelly, Bob Widmer, Boyd Bushman, George Hathaway, John Hutchison and John B. Alexander.

My particular thanks go to:

Mark Booth, Hannah Black, Anna Cherrett and the whole team at Century, Charlie Conrad, Claire Johnson, Chava Boylan, Becky Cole, and the whole Broadway Books team in New York, Bill Rose for his outstanding picture research, Igor Witkowski for taking me to the Wenceslas Mine and for sharing his tremendous insight into German secret weapons, Tom Agoston for his groundbreaking book *Blunder!*, the Schauberger family—Frau Ingeborg, Joerg and Ingrid—for their kindness and hospitality at Bad Ischl, Brue Richardson for her constancy, serenity and guidance, Barry and Fiona Shaw for putting up with me (and the rest) during a myriad of writing weekends, Dr. Patrick G. Bailey and Charlee Trantino for their sterling work on the index, and Mark Lucas for being a damn fine friend and agent.

Last, but not least, I would like to thank Ali, my family and hers. This book would have been quite impossible without them.

Prologue

The dust devils swirled around my Chevrolet Blazer, catching the early evening light. I watched Sheriff's Deputy Amelia Lopez clamber out of her Chrysler Le Baron and stare for a moment in the direction the plane had gone down ten years ago.

I grabbed my rucksack. By the time I looked up again she was already striding toward the peak.

Over a rusted barbed-wire fence and we were into the scrub—with new traces of green at its tips from the spring rains. Beyond lay the edge of the Sequoia National Forest, a huge expanse of protected park and woodland.

We left the broken fence posts behind and our cars were lost against the sunset. I looked for other traces of a human presence, but found none, even though we were only twelve miles from Bakersfield, California, a city of four hundred thousand people on the edge of the Sierra Nevadas.

Amelia Lopez' peaked cap and the firearm on her gunbelt were clearly silhouetted as she moved along the jagged edge of the ridgeline.

As I sucked down the warm thin air and wiped the sweat out of my eyes, I tried to picture her as she must have been on that sweltering July night ten years earlier; the night she'd been out partying with her college friends at a campsite near the Kern River.

It was in the breaking hours of July 11, 1986. Just as she was settling into her sleeping bag the jet went supersonic somewhere in the black sky overhead.

The pressure wave of the sonic boom hit the campsite like a clap of thunder, sending a shower of embers from the campfire into the night sky.

Amelia was too startled to say a thing; then, the entire horizon was flash-lit by an enormous explosion, the flames shooting skyward as the plane plowed into Saturday Peak ten miles away.

She told me it had sparked a dozen brushfires on the edge of the forest; that it took more than a hundred Forest Service and local firefighters to

put it out. Her only thought was that this wasn't an aircraft at all, but a hydrogen bomb.

Within hours, every newspaper in the state, and a whole lot more besides, had a reporter heading for these foothills with a brief to find out what had hit the ground. Amelia Lopez, a law student at the state university in Sacramento, had been one of several witnesses quoted in the papers that had covered the story, which was how I'd traced her.

She and her friends hadn't gone more than five miles toward the impact point when one of them noticed a figure on the trail up ahead. The scrub either side of her erupted with movement and the next thing she knew her face was in the dirt and she had a boot in her back and a gun at her head.

Out of the corner of her eye she saw that they were soldiers—not California National Guard, as might have been expected in an environmental emergency, but SWAT-types brandishing assault rifles, night-vision systems and a shit-load of threats about government property and national security.

Two of her friends started on about their rights under the Constitution, that this was public land and there wasn't a person on earth who could tell them to get off it. But to Lopez their protests registered as white noise on the edge of a persistent and piercing alarm. These soldiers were unlike any she'd ever seen.

She screamed for her friends to shut up, but once the screaming began, she couldn't stop it. She screamed and she yelled and she flailed against the pressure in her back, until the next thing she felt was the slap from her roommate that brought her around. When she finally understood what she was being told, it was that the soldiers were gone.

None of them said a word as they doubled back to the campsite. When they reached it, still numb with shock, they found a bunch of reporters on-site getting statements from other witnesses. Somebody shoved a tape recorder in her face and started asking questions and before she knew it she'd given her name and stammered something about an atomic blast.

As for the rest, she and her friends said nothing.

Amelia Lopez kept a lid on her feelings for the next two and half years, until November 1988, in fact, when the outgoing Reagan administration revealed the existence of the F-117A Stealth Fighter, an aircraft that had been flying in secret squadron service out of a classified air base in Nevada for over five years. In that time, she learned, it had crashed twice: and on one of those occasions—on the night of July 10 -11, 1986, to be precise—she'd had the grave misfortune to be there.

The troops had been part of a Pentagon "red team" flown in by helicopter to secure the crash site at all costs.

When she got my message, she'd been reluctant to meet up at first, but when I finally persuaded her to talk, she found it difficult to stop.

We reached the crash site soon after the sun dipped below the edge of the mountain. The summit was only 2,000 feet above us, but here the ground was even and covered in a crusty layer of dirt. The plants and trees were younger than the vegetation we'd passed on the way up. But that was the only real clue something had happened here.

Amelia Lopez sat on a rock and slowly removed her mirror shades before pouring bottled springwater over her face. I felt her eyes follow me as I moved between clumps of vegetation, kicking over rocks and sifting the sand, even though there was nothing to see.

Lopez bent down and ran her fingers through the soil. "I read they sieved the dirt for a thousand yards out from the impact point," she said. "Those guys were damned thorough. A few weeks after they left it was like *nothing* ever happened here." She paused a moment before adding: "You being an expert, I imagine you knew that."

It was framed as a question and I grappled for something to say, conscious that she'd brought me here for any insight I could provide into the events of that night.

I said nothing, so she turned to me and said: "Are you gonna tell me what is really going on here?"

Overhead, an eagle cried. As I watched it wheel on the updrafts I hoped that she wouldn't press me for an answer, because I didn't know what to tell her.

Standing here in this place, I was filled with the old feeling. It was almost impossible to articulate, but it left you with a taste in the mouth, some innate sense, that however far you dug, however many people you interviewed or questioned, you were simply scratching the surface of the sprawling U.S. defense-industrial base. What had happened here, the events that had imprinted themselves onto the landscape in a moment or two of madness a decade or so earlier, were almost tangible, even though there was no physical evidence—no fragments amidst the thin soil and the rocks—to suggest anything out of the ordinary had occurred.

These people were thorough; Lopez herself had said it. But they left something behind, something you couldn't see or touch—and it was that trace, that echo of past deeds, that had brought me here.

The Stealth Fighter was real enough. As a reporter, I'd covered it from the inside out. Yet as a piece of technology it was more than two decades

old, almost every detail of it in the open now. But strip away the facts and the feeling persisted.

I got it when I went to U.S. government defense laboratories and on empty windblown hangar floors in parched, little-known corners of the country. I got it at press conferences in power-soaked corridors of the Pentagon. But most of all I got it when I stared into the eyes of the people who worked on those programs.

What I got back was a look. Individually, it said nothing, but collectively it told me there was a secret out there and that it was so big no one person held all the pieces. I knew, too, that whatever it was, the secret had a dark heart, because I could sense the fear that held it in place.

It was impossible to tell Lopez any of this, of course, because it was simply a feeling. But as I headed back for the car, I knew the trip hadn't been a waste.

At long last, the secret had an outline.

Through half-closed eyes, I could almost reach out and touch it.

Chapter 1

From the heavy-handed style of the prose and the faint handwritten "1956" scrawled in pencil along the top of the first page, the photocopied pages had obviously come from some long-forgotten schlock popular science journal.

I had stepped away from my desk only for a few moments and somehow in the interim the article had appeared. The headline ran: "The G-Engines Are Coming!"

I glanced around the office, wondering who had put it there and if this was someone's idea of a joke. The copier had cut off the top of the first page and the title of the publication with it, but it was the drawing above the headline that was the giveaway. It depicted an aircraft, if you could call it that, hovering a few feet above a dry lakebed, a ladder extending from the fuselage and a crewmember making his way down the steps dressed in a U.S.-style flight suit and flying helmet—standard garb for that era. The aircraft had no wings and no visible means of propulsion.

I gave the office another quick scan. The magazine's operations were set on the first floor. The whole building was open-plan. To my left, the business editor was head-down over a proof-page checking copy. To her right was the naval editor, a guy who was good for a windup, but who was currently deep into a phone conversation and looked like he had been for hours.

I was reminded of a technology piece I'd penned a couple of years earlier about the search for scientific breakthroughs in U.S. aerospace and defense research. In a journal not noted for its exploration of the fringes of paranormality, nor for its humor, I'd inserted a tongue-in-cheek reference to gravity—or rather to antigravity, a subject beloved of science-fiction writers.

"For some U.S. aerospace engineers," I'd said, "an antigravity propulsion system remains the ultimate quantum leap in aircraft design." The implication was that antigravity was the aerospace equivalent of the holy grail: something longed for, dreamed about, but beyond reach—and likely always to remain so.

Somewhat the reference had escaped the sub-editors and, as a result, amongst my peers, other aerospace and defense writers on the circuit, I'd taken some flak for it. For Jane's, the publishing empire founded on one man's obsession with the detailed specifications of ships and aircraft almost a century earlier, technology wasn't something you joked about.

The magazine I wrote—and still write—for, *Jane's Defence Weekly*, documented the day-to-day dealings of the multibillion-dollar defense business. *JDW*, as we called it, is but one of a portfolio of products detailing the ins and outs of the global aerospace and defense industry. If you want to know about the thrust-to-weight ratio of a Chinese combat aircraft engine or the pulse repetition frequency of a particular radar system, somewhere in the Jane's portfolio of products there is a publication that has the answers. In short, Jane's was, and always has been, about facts. Its motto is: Authoritative, Accurate, Impartial.

It was a huge commercial intelligence-gathering operation; and provided they had the money, anyone could buy into its vast knowledge base.

I cast a glance at the bank of sub-editors' work-stations over in the far corner of the office, but nobody appeared remotely interested in what was happening at my desk. If the subs had nothing to do with it, and usually they were the first to know about a piece of piss-taking that was going down in the office, I figured whoever had put it there was from one of the dozens of other departments in the building and on a different floor. Perhaps my anonymous benefactor had felt embarrassed about passing it on to me?

I studied the piece again.

The strapline below the headline proclaimed: "By far the most potent source of energy is gravity. Using it as power, future aircraft will attain the speed of light." It was written by one Michael Gladych and began: "Nuclear-powered aircraft are yet to be built, but there are research projects already under way that will make the super-planes obsolete before they are test-flown. For in the United States and Canada, research centers, scientists, designers and engineers are perfecting a way to control gravity—a force infinitely more powerful than the mighty atom. The result of their labors will be antigravity engines working without fuel—weightless airliners and space ships able to travel at 170,000 miles per second."

On any other day, that would have been the moment I'd have consigned it for recycling. But something in the following paragraph caught my eye.

The gravity research, it said, had been supported by the Glenn L.



The G-Engines Are Coming!

By far the most potent source of energy is gravity. Using it as power future aircraft will attain the speed of light.

By MICHAEL GLADYCH

■ Nuclear-powered aircraft are yet to be built, but there are research projects already under way that will make the super-planes obsolete before they are test-flown. For in the United States and Canada research centers, scientists, designers and engineers are perfecting a

which there has been no escape. "What goes up must come down," they said. The bigger the body the stronger the gravity-attraction it has for other objects . . . the larger the distance between the objects, the lesser the gravity pull. Defining those rigid rules was as

way to earth.

This discovery gave modern scientists a new hope. We already knew how to make magnets by coiling a wire around an iron core. Electric current running through the coiled wire created a magnetic field and it could be switched on

Martin Aircraft Company, Bell Aircraft, Lear "and several other American aircraft manufacturers who would not spend millions of dollars on science fiction." It quoted Lawrence D. Bell, the founder of the plane-maker that was first to beat the sound barrier. "We're already working on nuclear fuels and equipment to cancel out gravity." George S. Trimble, head of Advanced Programs and "Vice President in charge of the G-Project at Martin Aircraft," added that the conquest of gravity "could be done in about the time it took to build the first atom bomb."

A little further on, it quoted "William P. Lear, the chairman of Lear Inc., makers of autopilots and other electronic controls." It would be another decade before Bill Lear went on to design and build the first of the sleek business jets that still carry his name. But in 1956, according to Gladych, Lear had his mind on other things.

"All matter within the ship would be influenced by the ship's

gravitation only," Lear apparently said of the wondrous G-craft. "This way, no matter how fast you accelerated or changed course, your body would not feel it any more than it now feels the tremendous speed and acceleration of the earth." The G-ship, Gladych explained, could take off like a cannon shell, come to a stop with equal abruptness and the passengers wouldn't even need seat belts. This ability to accelerate rapidly, the author continued, would make it ideal as a space vehicle capable of acceleration to a speed approaching that of light.

There were some oblique references to Einstein, some highly dubious "facts" about the nature of subatomic physics and some speculation about how various kinds of "antigravity engines" might work.

But the one thing I kept returning to were those quotes. Had Gladych made them up or had Lawrence Bell, George S. Trimble and William "Bill" Lear really said what he had quoted them as saying?

Outside, the rain beat against the double-glazed windows, drowning the sound of the traffic that crawled along the London to Brighton road and the unrelenting hum of the air conditioning that regulated the temperature inside.

The office was located in the last suburb of the Greater London metropolis; next stop the congested joys of the M25 ring road and the M23 to Gatwick Airport. The building was a vast redbrick two-story bunker amid between-the-wars gray brickwork and pebbledash. The rain acted like a muslin filter, washing out what little ambient color Coulsdon possessed. In the rain, it was easy to imagine that nothing much had changed here for decades.

As aviation editor *of JDW*, my beat was global and it was pretty much unstructured. If I needed to cover the latest air-to-surface weapons developments in the U.S.A., I could do it, with relatively few questions asked. My editor, an old pro, with a history as long as your arm in publishing, gave each of us, the so-called "specialists" (the aviation, naval and land systems editors), plenty of rope. His only proviso was that we filed our expenses within two of weeks of travel and that we gave him good, exclusive stories. If I wanted to cover an aerospace and defense exhibition in Moscow, Singapore or Dubai, the funds to do so were almost always there.

As for the job itself, it was a mixture of hard-edged reporting and basic provision of information. We reported on the defense industry, but we were part of it, too—the vast majority of the company's revenue coming from the same people we wrote about. Kowtowing was a no-no, but so was kicking down doors. If you knew the rules and played by them you could access almost any part of the global defense-industrial complex.

In the course of a decade, I'd visited secret Russian defense facilities and ultrasensitive U.S. government labs. If you liked technology, a bit of skulduggery and people, it was a career made in heaven. At least 60 percent of the time I was on the road. The bit I liked least was office downtime.

Again, I looked around for signs that I was being set up. Then, satisfied that I wasn't, but feeling self-conscious nonetheless, I tucked the Gladych article into a drawer and got on with the business of the day. Another aerospace and defense company had fallen prey to post—Cold War economics. It was 24 hours before the paper closed for press and the news editor was yelling for copy.

Two days later, in a much quieter moment, I visited the Jane's library. It was empty but for the librarian, a nice man way past retirement age who used to listen to the BBC's radio lunchtime news while gazing out over the building's bleak rear lot.

In the days before the Internet revolution, the library was an invaluable resource. Fred T. Jane published his first yearbook, *Jane's Fighting Ships*, in 1898; and in 1909 the second, *Jane's All The World's Aircraft*, quickly built on the reputation of the former as a reference work *par excellence* for any and all information on aeronautical developments. Nigh on a century later, the library held just about every book and magazine ever put out by the company and a pile of other reference works besides.

I scanned the shelves till I found what I was looking for.

The *Jane's All The World's Aircraft* yearbook for 1956 carried no mention of antigravity experiments, nor did successive volumes, but that came as no great surprise. The yearbooks are the aerospace equivalent of *Burke's Peerage* or the *Guinness Book of Records*: every word pored over, analyzed and double-checked for accuracy. They'd have given anti-gravity a very wide berth.

For a story like this, what I was looking for was a news publication.

I looked along the shelves again. Jane's had gotten into the magazine publishing business relatively recently and the company's copies of *Flight International* and *Aviation Week* ran back only a few years. But it did have bound volumes of *Interavia Aerospace Review* from before the Second World War. And it was on page 373 in the May 1956 edition of this well-respected publication, in amongst advertisements for Constellation airliners, chunky-looking bits of radar equipment and (curiously for an aviation journal) huge "portable" Olivetti typewriters, that I found a feature bylined "Intel, Washington, D.C." with the headline:

"Without Stress or Strain . . . or Weight." Beneath it ran the strapline: "The following article is by an American journalist who has long taken a keen interest in questions of theoretical physics and has been recommended to the Editors as having close connections with scientific circles in the United States. The subject is one of immediate interest, and *Interavia* would welcome further comment from knowledgeable sources."

The article referred to something called "electro-gravitics" research, whose aim was to "seek the source of gravity and its control." This research, "Intel" stated, had "reached a stage where profound implications for the entire human race are beginning to emerge."

I read on, amused by the tone and wondering how on earth the article had come to be accepted in a mainstream aerospace journal.

"In the still short life of the turbojet airplane [by then, 1956, little more than a decade], man has had to increase power in the form of brute thrust some twenty times in order to achieve just twice the speed. The cost in money in reaching this point has been prodigious. The cost in highly specialized man-hours is even greater. By his present methods man actually fights in direct combat the forces that resist his efforts. In conquering gravity he would be putting one of his most competent adversaries to work for him. Antigravities is the method of the picklock rather than the sledgehammer."

Not only that, the article stated, but antigravity could be put to work in other fields beyond aerospace. "In road cars, trains and boats the headaches of transmission of power from the engine to wheels or propellers would simply cease to exist. Construction of bridges and big buildings would be greatly simplified by temporary induced weightlessness etc. Other facets of work now under way indicate the possibility of close controls over the growth of plant life; new therapeutic techniques, permanent fuelless heating units for homes and industrial establishments; new sources of industrial power; new manufacturing techniques; a whole field of new chemistry. The list is endless . . . and growing."

It was also sheer fantasy.

Yet, for the second time in a week I had found an article—this time certainly in a publication with a solid reputation—that stated that U.S. aerospace companies were engaged in the study of this "science." It cited the same firms mentioned by Gladych and some new ones as well: Sperry-Rand and General Electric among them. Within these institutions, we were supposed to believe, people were working on theories that could not only make materials weightless, but could actually give them

"negative weight"—a repulsive force that would allow them to loft away "contra-gravitationally." The article went further. It claimed that in experimentation conducted by a certain "Townsend T. Brown" weights of some materials had already been cut by as much as 30 percent by "energizing" them and that model "disc airfoils" utilizing this technology had been run in a wind tunnel under a charge of a hundred and fifty kilovolts "with results so impressive as to be highly classified."

I gazed out over the slate rooftops. For *Interavia* to have written about antigravity, there had to have been something in it. The trouble was, it was history. My bread-and-butter beat was the aerospace industry of the 1990s, not this distant cozy world of the fifties with its heady whiff of jet-engine spirit and the developing Cold War.

I replaced the volume and returned to my desk. It should have been easy to let go, but it wasn't. If people of the caliber quoted by Gladych and *Interavia* had started talking about antigravity anytime in the past ten years I would have reported it—however skeptical I might be on a personal level. Why had these people said the things they had with such conviction? One of them, George S. Trimble, had gone so far as to predict that a breakthrough would occur in around the same time it took to develop the atomic bomb—roughly five years. Yet, it had never happened. And even if the results of "Townsend T. Brown's" experiments had been "so impressive as to be highly classified," they had clearly come to naught; otherwise, by the '60s or '70s the industry would have been overtaken by fuelless propulsion technology.

I rang a public relations contact at Lockheed Martin, the U.S. aerospace and defense giant, to see if I could get anything on the individuals Gladych had quoted. I knew that Lawrence Bell and Bill Lear were both dead. But what about George S. Trimble? If Trimble was alive—and it was a long shot, since he would have to be in his 80s—he would undoubtedly confirm what I felt I knew to be true; that he had been heavily misquoted or that antigravity had been the industry's silly-season story of 1956.

A simple phone call would do the trick.

Daniella "Dani" Abelman was an old media contact within Lockheed Martin's public affairs organization. Solid, reliable and likable, she'd grown up in the industry alongside me, only on the other side of the divide. Our relationship with the information managers of the aerospace and defense world was as double-edged as the PR/reporter interface in any other industry. Our job was to get the lowdown on the inside track and, more often than not, it was bad news that sold. But unlike our

The Hunt for Zero Point

national newspaper counterparts, trade press hacks have to work within the industry, not outside it. This always added an extra twist to our quest for information. The industry comprised hundreds of thousands of people, but despite its size, it was surprisingly intimate and incestuous enough for everyone to know everyone else. If you pissed off a PR manager in one company, even if it was on the other side of the globe, you wouldn't last long, because word would quickly get around and the flow of information would dry up.

But with Abelman, it was easy. I liked her. We got on. I told her I needed some background on an individual in one of Lockheed Martin's "heritage" companies, a euphemism for a firm it had long since swallowed whole.

The Glenn L. Martin Company became the Martin Company in 1957. In 1961, it merged with the American-Marietta Company, becoming Martin-Marietta, a huge force in the Cold War U.S. defense electronics industry. In 1994, Martin-Marietta merged again, this time with Lockheed to form Lockheed Martin. The first of the global mega-merged defense behemoths, it built everything from stealth fighters and their guided weapons to space launchers and satellites.

Abelman was naturally suspicious when I told her I needed to trace an ex-company employee, but relaxed when I said that the person I was interested in had been doing his thing more than 40 years ago and was quite likely dead by now.

I was circumspect about the reasons for the approach, knowing full well if I told her the real story, she'd think I'd taken leave of my senses.

But I had a bona fide reason for calling her—and one that legitimately, if at a stretch, involved Trimble: I was preparing a piece on the emergence of the U.S. aerospace industry's "special projects" facilities in the aftermath of the Cold War.

Most large aerospace and defense companies had a special projects unit; a clandestine adjunct to their main business lines where classified activities could take place. The shining example was the Lockheed Martin "Skunk Works," a near-legendary aircraft-manufacturing facility on the edge of the California high desert.

For 50 years, the Skunk Works had sifted Lockheed for its most highly skilled engineers, putting them to work on top secret aircraft projects.

Using this approach it had delivered some of the biggest military breakthroughs of the 20th century, among them the world's first Mach 3 spyplane and stealth, the art of making aircraft "invisible" to radar and other enemy sensor systems.

But now the Skunk Works was coming out of the shadows and, in the

process, giving something back to its parent organization. Special projects units were renowned for bringing in complex, high-risk defense programs on time and to cost, a skill that had become highly sought after by the main body of the company in the austere budget environment of the 1990s.

Trimble, I suggested, might be able to provide me with historical context and "color" in an otherwise dry business story. "Advanced Programs," the outfit he was supposed to have worked for, sounded a lot like Martin's version of the Skunk Works.

Abelman said she'd see what she could do, but I wasn't to expect any short-order miracles. She wasn't the company historian, she said dryly, but she'd make a few inquiries and get back to me.

I was surprised when she phoned me a few hours later. Company records, to her surprise—and mine—said that Trimble was alive and enjoying retirement in Arizona. "Sounds hard as nails, but an amazing guy by all accounts," she breezed. "He's kinda mystified why you want to talk to him after all this time, but seems okay with it. Like you said, it's historical, right?"

"Right," I said.

I asked Abelman, while she was at it, for all the background she had on the man. History or not, I said, trying to keep it light, I liked to be thorough. She was professional enough to sound less than convinced by my newfound interest in the past, but promised she'd do her best. I thanked her, then hung up, feeling happy that I'd done something about it. A few weeks, a month at the outside, the mystery would be resolved and I could go back to my regular beat, case closed.

Outside, another bank of gray storm clouds was rolling in above rooftops that were still slick from the last passing shower.

I picked up my coat and headed for the train station, knowing that somewhere between the office and my flat in central London I was going to get soaked right through.

The initial information came a week later from a search through some old files that I'd buried in a collection of boxes in my basement: a company history of Martin Marietta I'd barely remembered I'd acquired. It told me that in 1955 Trimble had become involved in something called the Research Institute for Advanced Studies, RIAS, a Martin spin-off organization whose brief was to "observe the phenomena of nature . . . to discover fundamental laws . . . and to evolve new technical concepts for the improvement and welfare of mankind."

Aside from the philanthropic tone, a couple of things struck me as

fishy about the RIAS. First off, its name was as bland as the carefully chosen "Advanced Development Projects"—the official title of the Skunk Works. Second, was the nature and caliber of its recruits. These, according to the company history, were "world-class contributors in mathematics, physics, biology and materials science."

Soon afterward, I received a package of requested information from Lockheed Martin in the mail. RIAS no longer existed, having been subsumed by other parts of the Lockheed Martin empire. But through an old RIAS history, a brochure published in 1980 to celebrate the organization's "first 25 years," I was able to glean a little more about Trimble and the outfit he'd inspired. It described him as "one of the most creative and imaginative people that ever worked for the Company."

I read on.

From a nucleus of people that in 1955 met in a conference room at the Martin Company's Middle River plant in Maryland, RIAS soon developed a need for its own space. In 1957, with a staff of about 25 people, it moved to Baltimore City. The initial research program, the brochure said, was focused on NASA and the agency's stated goal of putting a man on the moon. But that wasn't until 1961.

One obvious question was, what had RIAS been doing in the interim? Mainly math, by the look of it. Its principal academic was described as an expert in "topology and nonlinear differential equations."

I hadn't the least idea what that meant.

In 1957, the outfit moved again, this time to a large mansion on the edge of Baltimore, a place chosen for its "campus-like" atmosphere. Offices were quickly carved from bedrooms and workshops from garages.

It reminded me of accounts I'd read of the shirtsleeves atmosphere of the early days of the Manhattan Project when Oppenheimer and his team of atom scientists had crunched through the physics of the bomb.

And that was the very same analogy Trimble had used. The conquest of gravity, he'd said, would come in the time it took to build the bomb.

I called a few contacts on the science and engineering side of Lockheed Martin, asking them, in a roundabout kind of way, whether there was, or ever had been, any part of the corporation involved in gravity or "counter-gravitational" research. After some initial questions on their part as to why I should be interested, which I just about managed to palm off, the answer that came back was a uniform "no." Well, almost. There was a guy, one contact told me, a scientist who worked in the combat aircraft division in Fort Worth who would talk eloquently about the mysteries of Nature and the universe to anyone who would listen. He'd

also levitate paper clips on his desk. Great character, but a bit of a maverick.

"Paper clips?" I'd asked. "A maverick scientist levitating paper clips on his desk? At Lockheed Martin? Come on."

My source laughed. If he hadn't known better, he'd have said I was working up a story on antigravity.

I made my excuses and signed off. It was crazy, possibly dangerous stuff, but it continued to have me intrigued.

I called an old friend who'd gained a degree in applied mathematics. Tentatively, I asked whether topology and nondifferential linear equations had any application to the study of gravity.

Of course, he replied. Topology—the study of shape in physics—and nonlinear equations were the standard methods for calculating gravitational attraction.

I sat back and pieced together what I had. It didn't amount to much, but did it amount to something?

In 1957, George S. Trimble, one of the leading aerospace engineers in the U.S. at that time, a man, it could safely be said, with a background in highly advanced concepts and classified activity, had put together what looked like a special projects team; one with a curious task.

This, just a year after he started talking about the Golden Age of Antigravity that would sweep through the industry starting in the 1960s.

So, what went wrong?

In its current literature, the stuff pumped out in press releases all the time, the U.S. Air Force constantly talked up the "vision": where it was going to be in 25 years, how it was going to wage and win future wars and how technology was key.

In 1956, it would have been as curious as I was about the notion of a fuelless propulsion source, one that could deliver phenomenal performance gains over a jet; perhaps including the ability to accelerate rapidly, to pull hairpin turns without crushing the pilot and to achieve speeds that defied the imagination. In short, it would have given them something that resembled a UFO.

I rubbed my eyes. The dim pool of light that had illuminated the Lockheed-supplied material on Trimble and RIAS had brought on a nagging pain in the back of my head. The evidence was suggesting that in the mid-'50s there had been some kind of breakthrough in the anti-gravity field and for a small window in time people had talked about it freely and openly, believing they were witnessing the dawn of a new era, one that would benefit the whole of mankind.

Then, in 1957, everyone had stopped talking about it. Had the military woken up to what was happening, bringing the clamps down?

Those in the know, outfits like Trimble's that had been at the forefront of the breakthrough, would probably have continued their research, assembling their development teams behind closed doors, ready for the day they could build real hardware.

But of course, it never happened.

It never happened because soon after Trimble, Bell and Lear made their statements, sanity prevailed. By 1960, it was like the whole episode never took place. Aerospace development continued along its structured, ordered pathway and antigravity became one of those taboo subjects that people like me never, ever talked about.

Satisfied that everything was back in its place and as it should be, I went to bed.

Somewhere in my head I was still tracking the shrill, faraway sounds of the city when the phone rang. I could tell instantly it was Abelman. Separated by an ocean and five time zones, I heard the catch in her breathing.

"It's Trimble," she said. "The guy just got off the phone to me. Remember how he was fine to do the interview? Well, something's happened. I don't know who this old man is or what he once was, but he told me in no uncertain terms to get off his case. He doesn't want to speak to me and he doesn't want to speak to you, not now, not ever. I don't mind telling you that he sounded scared and I don't like to hear old men scared. It makes me scared. I don't know what you were really working on when you came to me with this, Nick, but let me give you some advice. Stick to what you know about; stick to the damned present. It's better that way for all of us."

Chapter 2

In 1667, Newton mathematically deduced the nature of gravity, demonstrating that the same force that pulls an apple down to earth also keeps the moon in its orbit and accounts for the revolutions of the planets. But today, we are still thwarted in attempts to measure it with any great precision. In lab experiments carried out since the 1930s, G has consistently defied efforts to be measured to more than a few decimal places.

This was what the reference books told me as I plowed through a stack of standard science works in the musty, gothic surroundings of the local library.

It was intensive work. Science was something I'd come to associate with the grind of exams. It didn't feel like the beginnings of a journalistic investigation.

I continued to scratch notes. But Newton openly stated that he had no idea what gravity actually was. All he knew was that it had to be caused by something.

The idea that a body may act on another through the vacuum of space over huge distances "without the mediation of anything else . . . is to me so great an absurdity that I believe no man can ever fall into it. Gravity must be caused . . . but whether this agent be material or immaterial I have left to the consideration of my reader."

I glanced up. The librarian, who'd been waiting to catch my eye, nodded toward the clock on the wall behind her. I looked around and realized I was the only person in the room. I'd lost track; it was a Saturday and the library closed early.

Outside, the rain had given way to the starlit sky of a passing cold front. I pulled up my collar and started down the street, dodging puddles that shimmered under the streetlights. The anomaly over gravity's measurement and the uncertainties over its causes only served to tell me how incomplete my knowledge of physics was.

I reached the edge of the common. The lights of my home street were faintly visible through the trees. I thought about my late-night call from

Abelman. In the week since her approach to Trimble and his initial favorable response to the idea of an interview, it very much appeared that somebody had gotten to him. And then I thought about what I had just learned. If we had no real understanding of gravity, how could people say with such certainty that *antigravity* could not exist?

In 1990, the U.S. Air Force had been looking at developing a weapon capable of firing a "plasma bullet"—a doughnut-shaped ring of ionized gas—at 10,000 kilometers per second.

Shiva Star was capable of generating and holding up to 10 megajoules of electrical energy and a potential 10 trillion watts—three times as much as the entire U.S. electricity grid carried in a year. At the time that I visited Shiva, which was located within the USAF's directed energy research laboratory at Kirtland Air Force Base in New Mexico, program engineers had been readying to fire a plasma bullet sometime in 1995. The purpose of the bullet was to destroy incoming Russian nuclear warheads and, despite some fierce technological challenges, the program engineers were confident they could do it. But several years later, when I returned to Kirtland, it was like the plasma bullet project never existed. Engineers had difficulty even recalling it.

Officially, it had been terminated on cost grounds. But this made little sense. The program had been budgeted at \$3.6 million per year for five years. Eighteen million bucks to produce a true quantum leap weapon system. Few people I spoke to bought the official version. Somewhere along the way, they said, Shiva must have delivered. Somewhere along the line, the program had gone black.

I coupled this knowledge with what the antigravity proponents had been saying in 1956. If you could find a way of shielding objects from the effects of gravity, the military, let alone the economic ramifications would be enormous.

Aircraft propulsion seems to have progressed little in appreciable terms since the advent of the jet engine in the 1940s. Incremental improvements for decades have been of the order of a few percentage points. The fastest aircraft in the world—officially, at least—is still the Lockheed Blackbird, designed in the late 1950s, first flown in 1962 and retired in 1990.

Amongst my peers, there had been speculation since the late 1980s about the existence of a secret replacement for the Blackbird, a mythical plane called Aurora that supposedly flew twice as fast and on the edges of space.

I had no direct evidence of Aurora, but then I'd never gone looking for it either. On balance, though, I felt *something* had been developed.

In 1992, circumstantial evidence of Aurora's existence was strengthened when *Jane's Defence Weekly* carried a detailed sighting of a massive triangular-shaped aircraft spotted in formation with USAF F-111 bombers and an air-refueling tanker above an oil rig in the North Sea. The sighting was credible because it was witnessed by a highly trained aircraft recognition expert in the Royal Observer Corps who happened to be on the rig at the time.

Shiva had been my first brush with the "black" world, the Pentagon's hidden reservoir of defense programs—projects so secret that officially they did not exist. Since then, I'd felt the presence of other deep black projects, but only indirectly.

Looking for the black world was like looking for evidence of black holes. You couldn't see a black hole, no matter how powerful your telescope, because its pull sucked in everything around it, including the light of neighboring stars. But astronomers knew that black holes existed because of the intense friction they generated on their edges. It was this that gave them away.

Forty years ago, the people in charge of the Air Force's hidden budget would have been quick to see the extraordinary implication of Trimble's message; that there would be no limit—no limit at all—to the potential of an antigravity aerospace vehicle.

The black world would have thrilled to the notion of a science that did not exist.

As I crossed from the park back into the glare of the streetlights, I knew I was quite possibly staring at a secret that had been buried more than 40 years deep.

I called Lawrence Cross, an aerospace journalist from the circuit, an ex-Jane's man, now a bureau chief for a rival publication in Australia.

Cross and I had spent long hours ruminating on the existence of U.S. Air Force black programs and the kind of technologies the Air Force might be pursuing in ultrasecrecy.

I liked Cross, because he had his feet squarely on the ground, was a hell of a good reporter, but wasn't your average dyed-in-the-wool-type hack. It had been a while since we had spoken.

The phone rang for ages. It was ten at night my time, eight in the morning his; and it was the weekend. I could hear the sleep in his voice when he finally picked up the handset. In the background, a baby was

crying. Cross had three kids under six years old. Ninety percent of the time he looked completely exhausted.

He remained quiet as I sketched out the events of the past weeks. I told him about the article, Trimble's initial willingness to be interviewed, then the phone call from Abelman and her insistence, on Trimble's say-so, that I drop the whole business.

"This is interesting," he said, stifling a yawn, "but why the long-distance call?"

"I wanted to tap you on one of your case studies."

"Uh-huh." He sounded wary, more alert suddenly. "Which one?"

"Belgium," I said. "Wasn't there some kind of flap there a year or two back?"

"You could say that. Hundreds of people reported seeing triangular-shaped craft all over the country in two 24-hour waves—one in 1989, the other in 1990. The Belgian Air Force even scrambled F-16s to intercept them. Why the sudden interest?"

"You once told me that those craft might have been the result of some kind of secret U.S. development effort."

Cross laughed. "Maybe I did. But I've had time to study the official reports since—the ones put out by the Belgian government. Those craft were totally silent. They hovered, often very close to witnesses, and they never made a sound. You may find my take on this hard to swallow, but there is no technology—no technology on earth—that could produce that kind of performance."

"Didn't the Belgian press try to tag the sightings to Aurora?"

"Yes, but you and I know that that's crazy. Even if Aurora is real, don't tell me it can remain stationary one moment and fly Mach 7 the next. And without making a sound. Belgian radar tracked these things. The tapes show that they pulled turns of around 20 to 40 g—enough to kill a human pilot." He paused. "You're not seriously suggesting what I think you're suggesting, are you?"

"A 40-year U.S. development effort, in the black, to make antigravity technology a reality? Why not? They were talking about it openly in 1956, Lawrence, then it dropped off the scope. Completely and utterly, like somebody orchestrated the disappearance. It makes me want to consider the possibility, at least, that someone achieved a breakthrough and the whole thing went super-classified."

"And now you've got the bit between your teeth?"

"Something like that, yes."

For a long moment, Cross fell silent. Then I heard him light a cigarette. In the background, I heard his wife calling him. Then he

cupped the receiver, because I caught his voice, muffled, telling her he'd be there in a minute. The baby was still bawling its lungs out.

"If you break cover on this," he said, "you'll blow everything. For yourself, I mean."

"Come on," I said, "it's a story. It may be an old story, but I'll apply the rules that I would on any other. If there's any truth in it, the answers will pop out. They usually do."

"That's so bloody naive. If there is any truth in it, which I doubt, they'll already know you're interested and that's not going to help you one little bit. They'll stand in your way, like they may have done already with this old man . . . Trimble? If there isn't any truth ink, then you're just going to look like a fucking idiot."

"They, Lawrence? Who's they?"

"The security people. The keepers of the secrets. The men-in-black. You know who I mean."

I didn't. To my ears, it sounded more than a little insane.

"I'm going to bide my time," I said, returning to the reason for the call, "do this at my own pace. For the moment, there's no need for me to break cover. Right now, all I have to do is conduct some low-level research and keep my eyes and ears open when I'm out there in the field. No one has to know about any of this, Lawrence. All I'm asking for in the meantime is a little help. Some of your knowledge. A few facts."

"Listen," he said, "there are no facts in this field; the whole business, if you want to know, is riven with disinformation, much of it, in my opinion, deliberately orchestrated. Sooner or later, you're going to have to surface and when you do, some of that crazy UFO spin is going to rub off on you. That happens and you'll never eat lunch in this great industry of ours again. Do you understand what I'm saying?"

And then his tone softened. "I've got to go now, but if you really are hell-bent on taking this forward, you might want to try an outfit in Washington, D.C., called the Integrity Research Institute. They have a handle on some of this material. Just promise me you'll keep my name out of it, okay?" And with that, he hung up.

During a trawl of the Library of Congress in the summer of 1985, Dr. Paul LaViolette, a researcher badly bitten by the gravity bug, tripped over a reference in the card indexing system to a report called *Electrogravitics Systems—An Examination of Electrostatic Motion, Dynamic Counterbary and Barycentric Control*. The words immediately alerted LaViolette, because he knew they were synonyms for something that science said was impossible: antigravity.

LaViolette had become interested in antigravity during his study for a Ph.D. in systems science from Portland State University. His curiosity was aroused further when he realized that the electrogravitics report was missing from its rightful place in the library's stacks—"like it had been lifted," he told me later.

When LaViolette asked the librarian if she could try to locate a copy elsewhere, what began as a simple checking procedure soon developed into a full-blown search of the interlibrary loan network, all to no avail.

Convinced he would never see a copy, LaViolette gave up, but weeks later the librarian called to tell him that she had managed to track one down. It had been buried in the technical library of Wright-Patterson Air Force Base in Dayton, Ohio.

There were no other copies known to exist across the whole of America.

"Whatever's in it," she told LaViolette dryly, "it must be pretty exotic."

When LaViolette finally got the Air Force to release the report, exotic seemed about right. *Electrogravitics Systems*, drafted in February 1956, the same year Trimble and his colleagues began pronouncing publicly on antigravity, contained details of what appeared to be a Mach 3 antigravity aerospace vehicle designed as a fighter-interceptor for the U.S. Air Force.

The fact that the report had been located at the Air Force's premier research and development facility instantly made *Electrogravitics Systems* an intriguing piece of work. That the USAF also appeared to have sat on it for almost 40 years only served to underscore its importance, LaViolette and his associates felt.

I had managed to track LaViolette through the Integrity Research Institute. The institute sat on the edges of a sub-culture of researchers who relentlessly picked over the antigravity issue. Type "antigravity" into a Net search engine and an outpouring of conspiracy-based nonsense on the government suppression of antigravity technology usually popped up onscreen. In the sober world of defense journalism, I'd never remotely envisaged a day when I'd have to involve myself in such matters.

Now, recalling Cross' warnings about professional suicide, I was reluctant to take the plunge, but realized that I had no choice.

I spoke to a colleague of LaViolette's called Tom Valone who mercifully never pressed me on the reasons why I was so interested in the antigravity business. A few days later, a copy of *Electrogravitics Systems* duly arrived through the mail.

Electrogravitics Systems had not been prepared by the Air Force, it turned out, but by an outfit called the Gravity Research Group, Special Weapons Study Unit, a subdivision of Aviation Studies (International) Limited, a British organization.

I did some checking. Aviation Studies had been a think tank run by Richard "Dicky" Worcester and John Longhurst, two talented young aerospace analysts who in the mid-1950s produced a cyclo-styled newsletter that had a reputation for close-to-the-knuckle, behind-the-scenes reporting on the aerospace and defense industry.

From the start, the *Electrogravitics Systems* report adopted a lofty tone: "Aircraft design is still fundamentally as the Wrights adumbrated it, with wings, body, tails, moving or flapping controls, landing gear and so forth. The Wright biplane was a powered glider, and all subsequent aircraft, including supersonic jets of the 1950s are also powered gliders."

The writers went on to say that insufficient attention had been paid to gravity research and that the "rewards of success are too far-reaching to be overlooked."

But while it quickly became clear that the report was not authored by the USAF, or apparently even commissioned by it, it did appear to be remarkably well informed about the progress of gravity—and antigravity—research in the United States and elsewhere; something that LaViolette and Valone felt might have accounted for its burial for so long in the Wright-Patterson technical library.

The word "antigravity" was not used in the report, presumably because this pop-scientific term was felt to be too lurid for the report's paying subscribers. To those less fussed, however, the term "electrogravitics" seemed pretty much synonymous with antigravity, even by Aviation Studies' own definition:

"Electrogravitics might be described as a synthesis of electrostatic energy used for propulsion—either vertical propulsion or horizontal or both—and gravitics, or dynamic counterbary, in which energy is also used to set up a local gravitational force independent of that of the earth."

Counterbary is subsequently defined as "the action of lévitation where gravity's force is more than overcome by electrostatic or other propulsion."

Antigravity, then.

The report echoed what had already been made clear in the Gladych and *Interavia* articles: that antigravity was of real interest to just about every U.S. aerospace company of the day.

"Douglas has now stated that it has counterbary on its work agenda, but does not expect results yet awhile. Hiller has referred to new forms

of flying platform. Glenn Martin says gravity control could be achieved in six years, but they add it would entail a Manhattan Project type of effort to bring it about. Sikorsky, one of the pioneers, more or less agrees with the Douglas verdict . . . Clarke Electronics state they have a rig, and add that in their view the source of gravity's force will be understood sooner than some people think. General Electric is working on electronic rigs designed to make adjustments to gravity—this line of attack has the advantage of using rigs already in existence for other defense work. Bell also has an experimental rig intended, as the company puts it, to cancel out gravity, and Lawrence Bell has said that he is convinced practical hardware will emerge from current programs. Grover Leoning is certain that what he referred to as an electromagnetic contra-gravity mechanism will be developed for practical use. Convair is extensively committed to the work with several rigs. Lear Inc., autopilot and electronic engineers, have a division of the company working on gravity research and so also has the Sperry division of Sperry-Rand. This list embraces most of the U.S. aircraft industry. The remainder, Curtiss-Wright, Lockheed, Boeing and North American have not yet declared themselves, but all these four are known to be in various stages of study with and without rigs."

Everybody, in other words, had had a finger in the pie.

But where, I asked, was the evidence 40 years on? It wasn't simply that the results of all this activity had failed to yield hardware; no corporate history of any these companies that I'd ever read in all my years as a journalist in the industry even hinted that gravity, antigravity, electrogravitics, counterbary—call it what you will—had ever been of interest to any of them.

And then, on page three of the *Electrogravitics Systems* report I saw the mention of Thomas Townsend Brown—the "Townsend T. Brown" of the *Interavia* piece in the Jane's library.

The report referred to Brown's "Project Winterhaven." Apparently, Brown had invented a whole new approach to the mechanics of flight.

This notion of electrogravitic lift supposedly worked on the principle that a plate-like object charged positively on one side and negatively on the other would always exhibit thrust toward the positive pole, i.e. from negative to positive. If the plate is mounted horizontally and the positive pole is uppermost, the object will in effect lose weight, because it will want to rise skyward.

I was in no rush to judge the validity of the physics. But I'd worked in the business long enough to know that the military would be quick to

dismiss anything that wasn't practical. So the key question for me was how the military had viewed Brown's work.

I called the Integrity Research Institute and asked Tom Valone to send me as much information as his organization possessed on Brown.

Then, over the next few weeks, my life assumed a pattern. In the dead hours, long after the phone stopped ringing, I'd head down into the basement, fire up the computer and hit the Net, staying on-line deep into the night.

My great regret was that I couldn't contact George S. Trimble directly. Had I done so, I knew that Abelman would have gone ballistic. She'd told me to stay away from him and she had the power to ensure that I became an outcast if I didn't. Lockheed Martin was a large company and its word would spread quickly.

In that respect, Cross was right. Pretty soon, no one would want to talk to a technology hack who was running around asking madcap questions about antigravity.

Better, then, to do what I was doing; keep a lid on it. Here, in the silence of the night, I could roam the Internet and remain anonymous. Besides, it really did seem to be a case of reviewing the evidence and following up the clues—clues that had apparently never registered with the experts who had been reporting on all this for years.

That in itself was seductive. Little by little, I could feel myself being pulled in.

Chapter 3

Thomas Townsend Brown was born into a prominent family from Zanesville, Ohio, in 1905, two years after the Wright Brothers, residents of nearby Dayton, propelled their "Flyer," the first aircraft capable of sustained powered flight, into the sky at Kitty Hawk, North Carolina. From an early age, Brown exhibited traits that would later come to mark his work as a scientist. As a child, he built a workshop in his backyard and at the age often was already receiving signals from across the Atlantic on a homemade radio-receiver. At 16, he was broadcasting from his own radio station.

For all his prowess as an inventor, Brown appears to have been a somewhat recalcitrant student. In 1922, he was enrolled at the California Institute of Technology (Cal Tech), but soon fell into a disagreement with his teachers over the time allowed by the Institution for laboratory work, something he lived for. After failing his first year exams in chemistry and physics, he persuaded his father to sponsor the construction of a laboratory of his own. With no expense spared, a lab was installed on the second floor of their new house in Pasadena.

The quid pro quo, apparently, was that "T.T." had to receive home tuition to boost his grades. Brown subsequently reported that he made considerable progress in chemistry, while still finding time to devise an X-ray spectrometer for astronomical measurements. It was at this point that he began to develop his first theories about electrogravitation. In essence, Brown believed that gravitation might be a form of radiation, much like light, with a "push" as opposed to a "pull" effect.

"Word of this got out among my classmates," he related in a document called "A Short Autobiography," which I found on the Net, "and although shunned and made fun of by the professors, I was nevertheless called to the attention of Dr. Robert Andrew Millikan, Director of Cal Tech and, incidentally, my first physics teacher, who explained to me in great detail, why such an explanation of gravitation was utterly impossible and not to be considered."

In 1923, Brown transferred to Kenyon College, Gambier, Ohio, and

the following year switched to Ohio's Denison University where he came under the tutelage of physicist and astronomer Dr. Paul Alfred Biefeld, a former classmate of Einstein's in Switzerland. In earlier experimentation, Brown had made the startling discovery that a Coolidge X-ray tube exhibited thrust when charged to high voltage. It took Brown a while to realize that the motion was not caused by the X rays themselves, but by the electricity coursing through the tube. Brown went on to develop a device he called the "Gravitor," an electrical condenser sealed in a Bakelite case, that would exhibit a one percent weight gain or a one percent weight loss when connected to a 100-kilovolt power supply.

In 1929, Brown wrote up his discoveries in a paper entitled *How I Control Gravitation* and speculated as to how his invention might one day be used:

"The Gravitor, in all reality, is a very efficient motor. Unlike other forms of motor, it does not in any way involve the principles of electromagnetism, but instead it utilizes the principles of electro-gravitation.

"A simple gravitor has no moving parts, but is apparently capable of moving itself from within itself. It is highly efficient for the reason that it uses no gears, shafts, propellers or wheels in creating its motive power. It has no internal mechanical resistance and no observable rise in temperature. Contrary to the common belief that gravitational motors must necessarily be vertical-acting, the gravitor, it is found, acts equally well in every conceivable direction."

This ability to manipulate the force in all axes opened up the Gravitor's potential to aviation. Brown's research led him to the development of a shape that was the most efficient in the production of electro-gravitational lift: that of a perfect disc or saucer.

The way he proposed to control the craft was by dividing the disc into segments, each of which could be selectively charged. By moving the charge around the rim of the saucer, it would, Brown said, be possible to make it move in any direction.

I had to remind myself that this was the 1920s.

The aviation industry was still absorbing the technical lessons of the First World War, the biplane was king and fighters, typified by the Boeing Model ISA, a leading U.S. design of the mid-1920s, were struggling to attain top speeds of 160 mph.

But although Brown's principles seemed to have been confirmed by observation—something, after all, was causing his condenser plates to move, however much the naysayers denigrated the possibility that it was due to an electrogravitic reaction—he had no way of generating and maintaining an electrical charge that would keep a small model craft in the air, let alone to maneuver it. This force was calculated to be in the

region of 50 kilovolts, 200 times more than the charge that came from my wall socket.

Yet these were early days. What was astounding was that Brown had developed a concept for an air vehicle, shaped in the form of a disc, years before anyone had coined the term "flying saucer."

This, I thought, had to be more than happenstance, especially as the military were showing signs that they considered Brown to be not just a visionary, but a practical man of invention; someone they could turn to to develop nuts-and-bolts hardware.

In September 1930, Brown joined the U.S. Navy Reserve, but instead of going to sea was given orders to report to the Naval Research Laboratory (NRL) in Washington, D.C. In due course, he transferred to the NRL's "Heat and Light Division" and here he carried on the experiments that he started in Ohio.

Experiments were conducted which seemed to prove the concept of gravitation which he had first postulated at Cal Tech in 1923.

In 1932, Brown served as staff physicist on the Navy's Gravity Expedition to the West Indies. But by 1933, the effects of the Great Depression were forcing cutbacks within the Navy and Brown, suddenly finding himself out of a job at the NRL, was driven to seek work in industry, which he interleaved with his duties as a reserve naval officer.

With war clouds gathering in Europe, Brown was drafted full-time into the Navy and assigned to acoustic and magnetic minesweeping research, a field for which he seemed to have a great natural aptitude. He developed a method for maintaining the buoyancy of minesweeping cables, designed to trigger magnetic mines with a strong electrical force field, while simultaneously shielding them from blast effects. Brown took out a patent on the idea, which was immediately classified.

By 1940, he was appointed head of the Navy Bureau of Ships' mine-sweeping research and development activity and it was during this period that he experimented with "degaussing"—a method for canceling a ship's magnetic field. This was a critical breakthrough as it was a ship's magnetic field that triggered a new breed of German mines, weapons that the Nazis had developed in quantity.

There was no question that Brown was by now well plugged into the military's research and development community, and that his opinions were given great credit.

It is ironic, then, that it is at this point that his credibility is most open to question, thanks to his supposed association with the so-called "Philadelphia Experiment."

In this, according to legend, the U.S. Navy spirited one of its warships, complete with its crew complement, into another dimension.

The Philadelphia Experiment was made famous in a book of the same name by Charles Berlitz and William Moore, writers who have forged names for themselves with tales of paranormal mysteries. Even they confessed that it was "questionable whether Brown was really ever very heavily involved in the Philadelphia Experiment project"—a comment that assumes this incident took place at all.

But the association is ingrained. Run a check with a search engine and Brown is up there, his name highlighted, nine times out of ten, right alongside it.

I tried to maneuver around this obstacle, this blip in Brown's otherwise blemishless wartime record, but whichever way I turned, it wouldn't go away.

Its effect was powerful and immediate. Up until this moment, I'd trawled cyberspace and decades-old documents in the growing conviction that this unusually gifted engineer was an important, long-overlooked link in the antigravity mystery highlighted by the Gladych and *Interavia* articles.

But the instant the Philadelphia Experiment started to be mentioned in the same breath as Brown, it made me want to drop him like a stone.

It was just as Cross had said it would be. I could feel the association, with its whiff of paranoia and conspiracy, threatening to act like a contaminant on my thinking.

Better to avoid it altogether.

Everything had been going so well. Brown was university-educated, he'd worked in the aviation industry, he'd been embraced by the military and he'd done classified work. And then, the Philadelphia Experiment had come along.

Shit.

The substance of the story is that while experimenting with techniques in 1943 to make navy ships invisible to radar through the use of intense electromagnetic fields, a destroyer, the USS *Eldridge*, disappeared from its berth in Philadelphia and reappeared moments later at a Navy yard in Norfolk, Virginia, 250 miles away. During this time-lapsed interval, the crew were supposedly transported into a "parallel dimension," an experience that drove many of them insane.

Brown's credentials at the time, coupled with his refusal in the years since altogether to dismiss the Philadelphia Experiment as hokum, have helped to stoke the myth. And so it has stuck to him like glue.

In the silence of the basement, confronted by this dead end, I began to wonder about that.

If the original authorities on the story, Berlitz and Moore, had had their own doubts about Brown's involvement in the Philadelphia Experiment, how had he come to be linked so inextricably to it?

And then I thought of something else; something Cross had said. That this whole field was riven with disinformation, some of it, in his opinion, deliberately managed. The Soviets had even coined a term for it: *disinformatsiya*.

During World War Two and the long decades of the Cold War the Russian military had used *disinformatsiya* to achieve key tactical and strategic objectives.

Far from avoiding the Philadelphia Experiment, which was what my every professional instinct yelled I should be doing, I came to the uncomfortable conclusion that I ought to do precisely the opposite.

Tentatively, I began to surf the Net for obscure websites that told the story.

There are two versions of what the Philadelphia Experiment was trying to achieve. One holds that the Navy was testing a method that would make ships invisible to radar, the other was that it was attempting to develop some kind of *optical* cloaking device as well; that by generating an intense electromagnetic field around the *Eldridge*, it would distort both light and radar waves in its vicinity, rendering the ship invisible to both sensor systems and the human eye.

To do this, the legend stated, the *Eldridge* was equipped with tons of electronic equipment. These, the story had it, included massive electrical generators, several powerful radio frequency transmitters, thousands of power amplifier tubes, cabling to distribute the energy around the ship and special circuitry to tune and modulate the fields.

During the first test, which is supposed to have taken place in July 1943, the *Eldridge* is said to have become invisible at its berth, the only sign that it was there being a trough of displaced water beneath the hull. Though classified a success, the ship remaining invisible for around 15 minutes, the crew experienced side effects, including nausea, disorientation and memory loss.

If we can for the moment assume that there's some truth in the story this would have been quite unsurprising to those in charge of the science, since the nerve impulses by which the brain operates are signaled electrically. In effect, just as earlier iterations of the equipment had demonstrated on enemy mines, the field generators would have

"degaussed" the brains of anyone within the area of influence, causing them temporary and, in some cases, permanent damage.

In such circumstances, I could conceive of reasons why the Navy might want to plant misleading stories about the *Eldridge*—the more so, since it involved stealth.

It remains highly sensitive technology, but the use of electromagnetic fields in generating radar stealth is a technique well known to military science today.

But during World War Two—if Brown had discovered a means of shielding U.S. Navy ships from enemy radar (or even for marginally reducing their radar signature)—it would have rung right off the classification scale.

The last thing the Navy would have wanted was sailors running around complaining of headaches and revealing how they had come by them.

The next experiment supposedly went even further. In October 1943, after the Navy had replaced the first crew, a second test on the *Eldridge* was carried out.

This time, soon after the generators were switched on, the ship shimmered for a few seconds, remaining visible in outline only. Then there was a blinding flash and it vanished altogether. The legend states that it was transported briefly to a berth in Norfolk, Virginia, before making its way back to the yard at Philadelphia.

When investigators went on board, they found that some of the crew had been swallowed by the ether that had momentarily consumed the *Eldridge*. These men were never seen again.

Those that did make it back were either made dangerously ill by their adventure, experiencing intense nausea and headaches for years, or were driven mad by it.

Weirdest of all, five of the crew were found fused into the metallic structure of the ship, which had materially transmuted to accommodate them.

These were the broad "facts" as presented to Berlitz and Moore, who were first alerted to the mystery by a man who claimed to have been on the *Eldridge*. They were supported by an eccentric source identified variously as Carlos Allende and Carl Allen, who claimed to have witnessed the whole thing from the deck of the SS *Andrew Furuseth*, a merchant marine ship berthed close to the *Eldridge* in the Philadelphia navy yard.

This was in the mid-1970s, more than 30 years after the purported events. When the book came to be written, Brown became an inextricable part of the myth.

%
3fc
%

The only other thing I knew about disinformation, certainly as it had been practiced by the Soviets, was that it worked best when mixed in with a little truth.

Intuitively, therefore, I felt Brown must have been somewhere in the vicinity of the "experiment" when it was supposed to have happened. Beyond that, I could draw no other conclusions, so I returned to the documented facts of his career.

In 1942, he was appointed head of the Atlantic Fleet Radar Materiel School and Gyrocompass School in Norfolk, Virginia, a position that would have made him privy to some of the most highly classified technical secrets of the day. Whatever work he was engaged on there, it appears to have taken its toll, since the following year he suffered a nervous breakdown and was discharged from the Navy.

There is something neatly synergistic in this unfortunate development, which appears to have been real enough, with the mystery surrounding the USS *Eldridge*. One suggestion is that whatever work Brown was engaged in at the time, be it in the radar or minesweeping field, it caused him temporary memory loss.

While it stretched credulity to believe in stories of optical invisibility, tele-transportation and parallel dimensions, I did find it possible to envisage a scenario in which the legend of the *Eldridge* had grown out of the secrecy of Brown's legitimate work in the radar field; in much the same way that the supposed ability of RAF night fighter pilots to see in the dark by eating carrots stemmed from a childishly simple British ruse to protect its radar secrets at around the same time.

Despite Brown's illness—and whatever really happened in the Philadelphia and Norfolk shipyards—it appears to have had no long-term detrimental effect on his standing in the eyes of the Navy. In 1944, he went back to work as a radar consultant in Burbank, California, at Lockheed's Vega Division, which was responsible for development of the Navy's PV-2 Harpoon and P-2V Neptune antisubmarine patrol bombers.

At war's end, Brown moved to Pearl Harbor, Hawaii, where he once again resumed the antigravity work that had driven his research efforts as a young man.

It seemed to me from this period—long before the story of the Philadelphia Experiment emerged to complicate the picture—as if two portraits of Brown had been painted and were now in circulation: one portraying him as a mildly eccentric inventor with some harebrained ideas about negating the forces of gravity; the other showing him to be a

man responsible for some of the most highly classified research of the war.

Seen through this ambivalent prism, the story of the Philadelphia Experiment has helped to perform a very important function.

By 1980, it had managed to tip Brown over the edge; make him a wholly discredited figure in the eyes of science.

That left me with the uncomfortable feeling that the story had been carefully stage-managed. If so, why? And why so long after the supposed events had taken place?

The next occasion Brown is reputed to have demonstrated his Gravitor and tethered flying discs was to Admiral Arthur W. Radford, Commander in chief of the U.S. Pacific Fleet, in Hawaii in 1945. Brown had by now left California and was resident in Hawaii, where he worked temporarily as a consultant at the Pearl Harbor Navy Yard.

His demonstrations supposedly failed to impress the Navy, but there is no known official record of its reaction. One account is that it "refused funding for further research because of the negative opinion of other scientists." Another relates the story of his room being broken into after the Pearl Harbor demonstration and the theft of his notebooks. According to this variant, which was related by a close friend of Brown's, Josh Reynolds, the Navy returned the books days later, adding that it remained uninterested. The reason given was that the "effect" was not due to electrogravitation, but to "ionic wind"—this, despite Brown's view that he had conclusively proven otherwise with experiments under oil back in the 1920s.

From 1945 until 1952, little is known about Brown's activities, but based on what happened next, it is clear that along the way he returned from Hawaii to Los Angeles, where he established the Townsend Brown Foundation.

In 1952, the Foundation received an unannounced visit by Major General Victor E. Bertrandias of the U.S. Air Force. I learned of this thanks to a recovered transcript of a phone conversation between Bertrandias and another USAF general called Craig, whose job description was not immediately clear.

The transcript had been forwarded to me by Tom Valone at the Integrity Research Institute. It made clear that Bertrandias was astounded by what he witnessed.

"It sounds terribly screwy, but Friday I went down with Lehr, a man named Lehr, to a place called the Townsend Brown Foundation, and believe it or not, I saw a model of a flying saucer," Bertrandias reported.

"No," Craig replied, apparently without irony.

"I thought I should report it," Bertrandias told him. "There was a lot of objection to my getting in there by the party that took me and the thing frightened me—frightened me for the fact that it is being held or conducted by a private group. I was in there from about one-thirty until about five in the afternoon and I saw these two models that fly and the thing has such a terrific impact that I thought we ought to find out something about it—who these people are and whether the thing is legitimate. If it ever gets away I say it is in the stage in which the atomic development was in the early days."

"I see," General Craig replied. The man's urbane delivery earmarked him, to me at least, as someone big in Air Force intelligence.

"I was told that I was not to say anything about it," Bertrandias burbled on, "but I'm afraid that all that I heard made me believe in it and these were not schoolboys. It was conducted in rather an elaborate office in Los Angeles. I thought I should report it to you."

"Yes, well I'll look into it and see what I can find out," Craig told him, shortly before hanging up.

It seems that what Bertrandias had stumbled upon, a little before the Townsend Brown Foundation was ready to pitch it formally to the military, was Project Winterhaven—the distillation of all Brown's ideas into a blueprint for a manned antigravity fighter, built in the shape of a disc and capable of Mach 3, twice the speed of the leading jet-powered interceptor of the day.

Years later, LaViolette had found extensive references to Winterhaven in the sole remaining copy of the *Electrogravitics Systems* report the Congressional librarian had tracked down to the technical library at Wright-Patterson Air Force Base.

I was rapidly coming full circle.

I needed to see the report on Project Winterhaven, but remembering the difficulty LaViolette had had in tracking down the one remaining copy of *Electrogravitics Systems*, I couldn't begin to think how or where I would lay my hands on one. I imagined, given the Air Force's evident excitement about Brown's work, that if there were any copies of *Winterhaven* left, they were buried somewhere deep.

I rang Valone in Washington—my third call in as many weeks—and this time got the question I'd been dreading. Was Jane's running some kind of investigation into antigravity?

I told him it wasn't; that my interest in Brown's work was born of personal curiosity, nothing more, and that I'd appreciate it if he'd keep the matter to himself.

Valone seemed quite unfazed by this request. In the murky world of antigravity research, I realized it was probably par for the course.

The conversation drifted inevitably toward Townsend Brown and *Winterhaven*. It was then that I mentioned how I'd give my eyeteeth to know what was in it.

A week later, to my absolute astonishment, a copy of *Winterhaven* (registered copy No. 36) landed on my desk. Attached to the front page was a note from Valone. Thanks to the Freedom of Information Act, there was no need for me and my teeth to part company, he said. I smiled. The man was a one-stop shop.

He assured me, too, that no one else bar him and LaViolette would ever know that I was interested in the subject matter.

Nagged by the ease with which the military had relinquished its grip on *Winterhaven*, I started reading.

In his pitch to the military, on page six, Brown had written: "The technical development of the electrogravitic reaction would usher in a new age of speed and power and of revolutionary new methods of transportation and communication.

"Theoretical considerations would predict that, because of the sustained acceleration, top limits of speed may be raised far beyond those of jet propulsion or rocket drive, with possibilities of eventually approaching the speed of light in 'free space.' The motor which may be forthcoming will be essentially soundless, vibrationless and heatless."

I made notes. This was language that was similar in tone to that expressed by George S. Trimble and his quoted contemporaries in the 1956 Gladych article. Trimble, who 40 years later had canceled my interview with him, because he appeared—to Abelman's mind, at least—to be too afraid to talk.

The attributes of the technology were also right in line with the characteristics exhibited by the flying triangles observed over Belgium.

The little model discs that had so impressed General Bertrandias "contain no moving parts and do not necessarily rotate while in flight," the *Winterhaven* report stated. "In atmospheric air they emit a bluish-red electric coronal glow and a faint hissing sound."

Project *Winterhaven*, then, offered a systematic approach for the establishment of a U.S. antigravity program—echoing the origins of the U.S. atomic bomb project a decade earlier and the path that Trimble had advocated in 1956.

In the report, Brown recommended starting modestly with 2-foot discs charged at 50 kilovolts, then proceeding to 4-foot discs at 150 kV

and finally to a 10-foot disc at 500 kV, with careful measurements being made along the way of the thrust generated.

This was a sensible, structured approach. It showed that Winterhaven was ready to transition from the drawing board to the technology demonstration phase.

Full-blown development wouldn't be far behind.

Brown had also formulated a method for generating the required high voltages for a free-flying disc—the problem that had earlier stymied its practical development into a manned aircraft—by means of a "flame-jet generator." This was a jet engine modified to act as an electrostatic generator capable of providing up to 15 million volts to the skin of the craft. The *Interavia* article, I remembered, had talked of trials involving 3-foot discs run in a 50-foot diameter air course under a charge of 150 kV "with results so impressive as to be highly classified."

Brown's flying saucer now had a power source.

In one way and another things seemed to be gaining momentum.

But it was short-lived.

Days after General Bertrandias' conversation with his spook colleague General Craig, the documents trail showed that the Air Force Office of Special Investigations, AFOSI, was well and truly onto the Brown case. AFOSI was the Air Force equivalent of the FBI—an indication that Craig had taken Bertrandias' interpretation of his visit to the Brown Foundation seriously. Once again, it was Valone who furnished me with the relevant papers.

They showed that AFOSI uncovered a copy of a report prepared by one Willoughby M. Cady of the Office of Naval Research, the ONR, in Pasadena, California, entitled: "An Investigation Relative to the Townsend Brown Foundation."

The report, which was forwarded by AFOSI to Major General Joseph F. Carroll, Deputy Inspector General of the USAF, made disappointing reading all around.

"Mr. Brown claims that a gravitational anomaly exists in the neighborhood of a charged condenser," Cady wrote in his conclusions, adding: "This effect has not been well documented by Mr. Brown."

Cady went on to say that there was no electrogravitational link, merely a disturbance of the air around the model saucers stimulated by the electrical charge and it was this that was causing them to be propelled upward and forward.

Any hope that this technology might be of use in aerospace propulsion was damned by Cady. "If the efficiency of conversion of fuel energy into electrical energy is 21 percent, the overall efficiency of propulsion of the

model flying saucers by the electric wind is 0.3 percent. This compares with about 25 percent for a propeller-driven airplane and 15 percent for a jet airplane."

If Cady's conclusions were clear to me, they would have been crystal clear to the Navy and the Air Force. Brown's flying saucer would have had trouble getting off the ground, let alone cruising at Mach 3. The whole "science" of electrogravitics, Cady was telling his superiors, was a waste of time, effort and money.

The Navy and its old foe the Air Force would have concluded that they were both far better off sticking with jets.

Even under the auspices of a deeply classified program, an arrangement by which one branch of the armed forces might be unaware of projects taking place in another, it was hard to see Brown's work surviving such an appraisal.

On September 22, 1952, six months after General Bertrandias' telephone call to General Craig, documents showed that the Air Force downgraded its interest in the work of the Townsend Brown Foundation from "confidential" to "unclassified."

This explained why the Pentagon's archives had contained a declassified copy of *Winterhaven*.

Despite the view of LaViolette, Valone and many others that this was the moment Project Winterhaven went super-classified, forming the basis of a number of "black programs" in the antigravity field from the mid-1950s onward, I found this hard to believe. Although LaViolette's theory dovetailed neatly with the ringing silence that followed the zealous rhetoric of George Trimble and his colleagues in the U.S. aerospace industry on the subject of antigravity in 1956, there was no evidence at all to support it.

Nor, crucially, did LaViolette's thesis explain Brown's behavior from this period onward. For although it was obvious that he knew how to keep a secret—his wartime work would have instilled in him the need for the rigors of secrecy—if the ONR report was merely a blind, his subsequent actions went way beyond attempts to lead investigators away from a buried program within the U.S. Navy or Air Force or both. The energy with which he now set out to prove his electrogravitational theory seemed, in fact, to stem specifically from the comprehensive rejection of his experiments by the U.S. military. For the rest of his life, Brown, who was universally described as quiet, unassuming, honest and likable, behaved like a man driven to prove his point.

In 1955, he went to work for the French aerospace company

SNCASO—Société Nationale de Constructions Aéronautiques du Sud-Ouest.

During this one-year research period, he ran his discs in a vacuum. If anything, they worked better in a vacuum, something that prompted SNCASO, which was interested in exploiting Brown's work for possible space applications, to offer him an extension of his contract. But in 1956, SNCASO merged with its counterpart Sud-Est and its new bosses saw little future in the space business. They wanted to build aeroplanes; real ones, with wings and jet engines.

His contract terminated, Brown returned to America, where he helped found the National Investigations Committee on Aerial Phenomena (NICAP), an unofficial study group set up to analyze the growing body of UFO sightings across the American continent and elsewhere. It was Brown's belief that the study of UFOs, many of which exhibited characteristics similar to his discs, could shed light on their propulsion methodology and this, in turn, could be exploited for science and space travel. The trouble was, of course, that this alienated him even more from the mainstream.

It also further alienated him from me. As I'd immersed myself in Brown's life and work, I'd wondered more and more if his "discovery" of electrogravitics had had something to do with the emergence of the whole UFO phenomenon in the '40s, if these "alien" craft had in fact been top secret aerospace vehicles propelled by a power source that science even today refused to recognize. This, however, did not gel with a man who went on to found a UFO study group.

In 1957, Brown was hired as a consultant to continue his antigravity work for the Bahnson Company of North Carolina and in 1959 he found himself consulting for the aerospace propulsion giant General Electric.

I found little corroborating evidence for Brown's activities during these years.

When he went into semiretirement in the mid-1960s, Valone and LaViolette saw this as a signal that, in effect, he had been bought off by the military, especially as he hardly touched electrogravitics again. His last great interest involved a series of ultimately successful attempts to draw stored electrical energy—albeit in minute quantities—from common or garden rocks . . .

Unquestionably a highly gifted and unusual man, Brown died in relative obscurity in 1983.

In trying to draw lessons from his story, lessons that might perhaps help explain the sudden outpouring of interest in antigravity by America's leading aerospace designers in the late 1950s, I found myself

trying to put the pieces of the puzzle together. The problem was, however I went about it, they wouldn't form into any recognizable picture.

If the Navy had developed Brown's experiments into a fully fledged antigravity program, why had its premier research arm, the ONR, gone out of its way to dismiss them? And if this was an elaborate ruse to throw others off the scent—including its old rival the U.S. Air Force—why did Brown continue in his work, eventually taking it to another country? If electrogravitics was classified, this would have handed it on a plate to another nation, which was madness.

There was, conceivably, a chance that Brown's work had been hijacked by the U.S. military without his knowledge at the end of World War Two, but this too seemed unlikely. Given Brown's wartime clearances, it would have been simpler to have sworn him to secrecy after the Hawaii experiments, had the Navy been that impressed with them.

There was another, remote possibility—that Brown's work had been rejected by the military, not because it was hokey or crazy, but because its principles were already known to them—and, perhaps, therefore already the subject of advanced development activity. Had this been the case, it might well have explained why, a few years later, Bell, Convair, Martin and so many other companies—equally ignorant of this activity—aired their views on antigravity unchecked for a number of months, as recorded in the *Interavia* article, until someone, somewhere ordered them to silence.

As I kicked this notion around, I found sympathy for it from an unlikely source.

In the appendix section of a compendium of UFO sightings, I found a secret memorandum dated September 23, 1947, from Lieutenant General Nathan Twining, head of the U.S. Army Air Forces' Air Materiel Command (AMC), to Brigadier General George Schulgen, a senior USAAF staff officer in Washington.

The memorandum had been declassified and released into the public domain only in the late 1970s. I'd found it during another late-night trawl of my burgeoning archive, a cup of strong coffee in my hand to keep me awake.

In the memo, Twining states in his "considered opinion" that the rash of UFO sightings in America during the summer of 1947—the first real wave, as it turned out—had been "something real and not visionary or fictitious."

He continued: "There are objects . . . approximating the shape of a disc, of such appreciable size as to appear to be as large as a man-made

aircraft." These discs exhibited a number of common characteristics, amongst them a metallic or light-reflecting surface, an absence of any jet trail and no propulsion sound.

In addition, they were capable of "extreme rates of climb, maneuverability (particularly in roll), and action which must be considered evasive when sighted."

For an Air Force officer—a senior one, even—to admit to the reality of UFOs was quite an admission, but it was not unique. Other then-classified memos from other USAAF officers of the day show them to have been similarly perplexed by the disc sightings phenomenon across America from mid-1947 onward.

UFO researchers had seized on these documents as evidence that the USAAF (and from 1947 its successor, the USAF) recognized the reality of UFOs while officially pooh-poohing them—a blanket denial that remains policy to this day.

My attention, however, had been drawn to something quite else.

As an aerospace analyst, and in light of the Brown research, I found the key part of Twining's memo in what followed. UFO researchers had concentrated entirely on the main thrust of the memo: Twining's acknowledgment of the reality of UFOs.

The part that interested me, the postscript, added almost as an afterthought, had been totally overlooked. It so absorbed me that I never noticed the pencil on which I set down my cup of lukewarm coffee. It tipped over and the drink spilled, spreading across the open pages like ink on a blotter. I swore, mopped it up as best I could and returned to the text. The stain, as poor luck would have it, had covered the paragraph I was interested in, making the task of rereading it a painfully slow affair. I uttered the words aloud, enunciating them slowly so I knew I had read it right.

"It is possible," Twining wrote, "within the present U.S. knowledge—provided extensive detailed development is undertaken—to construct a piloted aircraft which has the general description of the object above which would be capable of an approximate range of 7,000 miles at subsonic speeds."

As head of Air Materiel Command, the branch of the USAAF that assumed responsibility for all USAAF aircraft development, Twining—who would have written the memo under the best advice from his subordinates—should have known what he was talking about. The problem was that 50 years this side of Twining's memo, at the end of the 20th century, I still knew of no combat aircraft capable of the performance characteristics Twining had elucidated. Even the Lockheed

e. There is a possibility that some of the incidents may be caused by natural phenomena, such as meteors.

d. The reported operating characteristics such as extreme rates of climb, maneuverability (particularly in roll), and action which must be considered erratic when sighted or contacted by friendly aircraft and radar, lend belief to the possibility that some of the objects are controlled either manually, automatically or remotely.

e. The apparent common description of the objects is as follows:

(1) Metallic or light reflecting surface.

(U) 7352

SECRET:

SECRET:

Basis for CO, AFM, AF to DC, AFM, AFM Operations Concerning Flying Discs".

(2) Absence of trail, except in a few instances when the object apparently was operating under high performance conditions.

(3) Circular or elliptical in shape, flat on bottom and domed on top.

(4) Several reports of well kept formation flights varying from three to nine objects.

(5) Normally no associated sound, except in those instances a substantial rumbling roar was noted.

(6) Level flight speeds normally above 300 knots are established.

f. It is noted within the present R. S. knowledge — predicted developments concerning the development of the aircraft described in paragraph (e) above which would be capable of an appreciable range of 700 miles at schematic speeds.

g. Any developments in this country along the lines indicated would be extremely expensive, time consuming and at the considerable expense of current projects and therefore, if directed, should be set up independently of existing projects.

h. Due consideration must be given the following:

(1) The possibility that these objects are of domestic origin — the projects of some high priority project not known to AF/AS-C on this Command.

(2) The lack of physical evidence in the shape of crash recovered artifacts which would undeniably prove the existence of these objects.

Report from the Materiel Division Report Date: 10 May 1960	NDP 76 0160 5-4-71
(U) SECURE	
SAF	
HEADQUARTERS	
AIR MATERIEL COMMAND	
SUBJECT: AFM Operation Concerning "Flying Discs"	
To:	Commanding General Air Force Materiel Command Washington 25, D. C. ATTENTION: AFM, General George Schulgen AC/AS-2
SAC: SEP 2 5 INT	
SUBJ: AFM Operation Concerning "Flying Discs"	
1. As requested by AF/AS-C there is presented below the Commanding General's opinion of this Command concerning the unclassified sightings of "flying discs". This opinion is based on investigation of records, data charts, and by AF/AS-C preliminary study by personnel of the Materiel and Aircraft Laboratory, Engineering Division T-3. This opinion was arrived at by consultation with personnel from the Air Directorate of Technology, Materiel Services T-2, Office of Chief of Engineering Division, and the Aircraft, Power Plant and Propeller Laboratories of Engineering Division T-3.	
2. It is the opinion that:	
a. The phenomenon reported is something real and not visionary or fictitious.	
b. There are objects probably approximating the shape of a disc, of much appreciable size as to appear to be as large as cabin-size aircraft.	

U-2 spyplane (which first flew in 1955, but is still in service), a single-engined aircraft intended for ultra-long endurance missions (and hardly, therefore, a maneuvering type like the discs reported to Twining), had a maximum unrefueled range in excess of 3,000 miles—3,500, perhaps, on a good day.

And, needless to say, no one had ever owned up to building a flying saucer.

To construct the "aircraft" that Twining had in mind, he told General Schulgen: "Any developments in this country along the lines indicated would be extremely expensive, time-consuming and at the considerable expense of other projects and therefore, if directed, should be set up independently of existing projects."

Placing such a project outside the mainstream of aerospace science brought to mind the effort that had resulted in the development of the first atomic bomb. In 1956, George Trimble had also cited the Manhattan Project as the inspiration for his proposed antigravity program. Brown, too, had offered a highly unusual development model for Winterhaven, his Mach 3 antigravity program, one that only peripherally involved the U.S. aerospace industry.

I tried to keep my mind on the facts. The essential point of Twining's memo was that the development of an aircraft exhibiting the characteristics of the saucers that people were seeing in 1947 was "within the present U.S. knowledge," but that the craft themselves were nothing to do with the USAAF. Then whose were they?

Twining himself offered two possibilities. First, that they were of "domestic origin—the product of some high security project not known to AC/AS-2 (air force technical intelligence) or this Command." And second, "that some foreign nation has a form of propulsion, possibly nuclear, which is outside of our domestic knowledge."

The first part I actually found a little scary. Twining was entertaining the possibility that these craft had been developed and fielded by another branch of the U.S. armed forces—the Navy, perhaps—with the knowledge of the USAAF.

The second explanation was, at least, a little more conventional. The discs, Twining offered, were the result of another country's secret development effort.

I had been sucked into a story that seemed to have had its origins in 1956. Reluctantly, I now had to concede that if I was to do this thing any justice at all, I needed to cast the net back even further.

Chapter 4

History doesn't relate the mood of the crewmembers of the Northrop P-61 Black Widow patrolling the sky above the Rhineland that night, but the available clues suggest that they were none too happy. A night-fighter crew was about as close-knit a unit as you could find among the frontline squadrons ranged against Germany in late 1944. Success—their very survival when it came down to it—depended on a high level of trust, intense training, the reliability of a piece of hardware then still in its infancy—radar—and undiluted concentration.

The last thing U.S. Army Air Forces Lieutenant Ed Schlueter would have needed that night was a passenger. Worse, Ringwald wasn't even aircrew, but an intelligence officer.

Lieutenant Fred Ringwald was sitting behind and above Schlueter, the P-61's pilot, in the position normally occupied by the gunner. Schlueter's unit, the 415th Night Fighter Squadron of the U.S. 9th Army Air Force, had recently upgraded from their British-made Bristol Beaufighters to the Black Widow. They had also just transferred from the Italian theater of operations to England and from there across the Channel, deploying eastward in short hops across northwest France as the Allies pushed the Nazis toward the Rhine and back into Germany itself.

Schlueter flew his aircraft south along the Rhine, looking for "trade." The Black Widow was a heavy fighter; bigger than the Beaufighter and considerably more menacing in appearance. While its primary targets were the German night fighters sent up to intercept British bomber streams heading to and from Germany, there was always the chance, provided Schlueter was sharp-eyed enough, of hitting a Nazi train or a vehicle convoy, especially as the Germans were moving men and materiel under cover of darkness due to the Allies' overwhelming daytime air superiority. This had been pretty much indisputable since the D-Day landings five months earlier.

But night strafing operations brought their own hazards. Over the uncertain territory of the Rhineland, sandwiched between the bluffs of the wide and winding river and the rugged uplands of the Black Forest,

there was a better than even chance you could follow your own cannon-fire into the ground.

There was no official record as to why the intelligence officer Ringwald was along for the ride, but knowing something of the way in which spooks had a habit of ring-fencing intelligence from the people who needed it most, I had a feeling that Schlueter and his radar operator would have been left just as much in the dark.

In the black skies above the Rhineland, after a long period of routine activity, it was Ringwald who broke the silence.

"What the hell are those lights over there?" he asked over the RT.

"Probably stars," Schlueter said, concentrating on his instruments.

"I don't think so," Ringwald replied. "They're coming straight for us."

Now, Schlueter looked up and out of the cockpit. In the pitch-black of his surroundings, the formation of aircraft off his starboard wing stood out like a constellation of tiny brilliant suns. Instinctively, he twisted the control column, bringing the Black Widow's four cannon and four .50 cal machine guns into line with them. At the same time, he called ground radar.

The ground station was supposed to be Schlueter's eyes and ears at all times, but if each pulsating point of light represented the exhaust plumes of a German night fighter, there were anything up to ten aircraft closing on him and he hadn't heard a whisper out of them. Somebody had screwed up. Later, he would be angry. Now, he was simply frightened and confused. He urgently requested information.

"Negative," the reply came back. "There are no bogies in your sector. You're on your own."

Schlueter's radar operator, Lieutenant Don Meiers, who was crouched over the scope of the SCR540 airborne intercept (AI) radar in a well behind Ringwald, told him the same thing. The sky ahead of them was empty of any air activity.

But the lights were still there and they kept coming. Instead of running, Schlueter boosted the throttles and pointed the nose of the Black Widow at the lead aircraft of the formation.

As the twin air-cooled radials powered up either side of him, the glow from his opponents' exhausts dimmed; and then they winked out. Puzzled and alarmed at having lost the contacts, and with no help from Meiers on the Black Widow's own radar, Schlueter held the aircraft steady and the crew braced themselves for the engagement.

Schlueter eased the night fighter into the blacked-out bowl of sky where he had last seen the contacts. He craned his neck for a glimpse of

something—anything—that signaled the presence of another aircraft, jinking the plane left and right to check his blind spots.

Still nothing.

It was only when he started to execute a turn back to base that Ringwald told him the lights were back again.

Schlueter followed the line indicated by Ringwald and saw them a long way off. Impossibly far, in fact, but still within radar range. He called out to Meiers, but the radar operator was now having technical problems with his AI set.

Schlueter again prepared to engage the enemy, but the lights had already begun to glide away to the northeast, eventually retreating deep into the German lines and disappearing altogether.

Nobody said anything until shortly before the Black Widow landed. Although Schlueter and Meiers were agreed that the Germans must have been experimenting with some new kind of secret weapon, neither wanted to hazard a guess as to what this weapon might have been. There was nothing they knew of in their own inventory that approached the weird, darting performance characteristics of the aircraft they'd just seen.

Fearful they would become the target of unwelcome squadron humor—along predictable lines they were "losing it"—they decided not to report the incident. And Ringwald, the spook, went along with it.

Reports of this incident exist in a number of UFO books—books I'd not encountered before because to someone steeped in the dry reportage of nuts-and-bolts technical journalism, they'd never entered my orbit.

The incident showed that almost three years before Twining wrote his memo to General Schulgen unconventional aerial objects had appeared in German skies prior to their manifestation across the U.S.A. in 1947.

On odd days off work and at weekends, I'd begun trawling public archives for corroborating evidence of this sighting. What I found were details on the 415th Night Fighter Squadron and the aircraft Schlueter had flown at the time of the encounter; details that allowed me to fill in the gaps of the published account and to visualize the sense of bewilderment and fear that Schlueter and his crew would have experienced that night. But along the way, I discovered that Schlueter's sighting was far from unique. All that winter of 1944-45, Allied aircrew reported small, ball-shaped aircraft glowing orange, red and white over the territory of the Third Reich. While some attributed the lights to natural phenomena such as ball lightning or St. Elmo's fire, others could not dismiss the sightings so easily. The devices appeared to be able to home in on Allied aircraft as if guided to them remotely or by some built-in control system.

Bit by bit, the reports entered the realm of officialdom. In archives and on the Internet, I found dozens of them.

"At 0600, at 10,000 feet, two very bright lights climbed toward us from the ground," another pilot from the 415th told intelligence officers after an encounter on December 22, near Haguenau, close to where Schlueter, Meiers and Ringwald had been. "They leveled off and stayed on the tail of our plane. They were huge bright orange lights. They stayed there for two minutes. On my tail all the time. They were under perfect control. Then they turned away from us, and the fire seemed to go out."

Although their appearance was sporadic, aircrews increasingly reported the devices via the appropriate channels. They nicknamed them "foo-fighters," a bastardization of the French word "feu" for "fire" that had worked its way into a cartoon strip called *Smokey Stover, the Foolish Foo Fighter* which had first appeared in a Chicago newspaper several years earlier. Meiers, who was a resident of Chicago, appears to have been the first person to have coined the term.

The consensus was that foo-fighters were Nazi secret weapons of some kind, but mighty strange ones, since they did not open fire on Allied aircraft, nor did they explode on proximity to them. They simply appeared, tagged along for a while and then vanished.

Seemingly, the spooks couldn't provide any plausible explanation for what they were either, as the following account, by Major William Leet, a B-17 pilot attached to the U.S. 15th Air Force, indicated after a nighttime encounter with a foo-fighter—"a small amber disc"—that followed his bomber all the way from Klagenfurt, Austria, to the Adriatic Sea in December 1944. "The intelligence officer that debriefed us stated it was a new German fighter but could not explain why it did not fire at us or, if it was reporting our heading, altitude and airspeed, why we did not receive antiaircraft fire," he reported.

Most encounters were at night, but there were daylight sightings, too.

A B-17 pilot, Charles Odom, flying on a daylight raid into Germany, described them as being "clear, about the size of basketballs." They would approach to within 300 feet, "then would seem to become magnetized to our formation and fly alongside. After a while, they would peel off like a plane and leave."

A P-47 fighter pilot also reported seeing a "gold-colored ball with a metallic finish" west of Neustadt in broad daylight, while another saw a "three to five feet diameter phosphorescent golden sphere" in the same area.

In 1992, researchers digging into the foo-fighter mystery uncovered a wealth of buried reports within the U.S. National Archives at College

Park, Maryland. What was intriguing was that almost all of them had been filed by pilots and crewmembers of the 415th Night Fighter Squadron. Unlike the incident involving lieutenants Schlueter, Meiers and Ringwald (whose experience was relayed after the war by a former war correspondent), the 15 "mission reports" of mysterious intercepts, many of which occurred in a triangular sector of the Rhineland bordered by an imaginary line linking Frankfurt-am-Main to the north, Metz to the west and Strasbourg to the south, could be examined in their original format; logged in the dispassionate shorthand of the intelligence officers who originally noted them down.

"December 22-23, 1944—Mission 1, 1705-1850. Put on bogie by Blunder at 1750 hours, had AI contact 4 miles range at 0[^]7372. Overshot and could not pick up contact again. AI went out and weather started closing in so returned to base. Observed two lights, one of which seemed to be going on and off at 0[^]2422."

And another:

"February 13-14, 1945—Mission 2,1800-2000. About 1910, between Rastatt and Bishwilier, encountered lights at 3,000 feet, two sets of them, turned into them, one set went out and the other went straight up 2-3,000 feet, then went out. Turned back to base and looked back and saw lights in their original position again."

The reports made it clear that the sightings covered a period between September 1944 and April 1945.

September. Two, maybe three months before Schlueter's and Meiers' own encounter.

Maybe, I thought, that tied up one little loose end.

When the spook, Ringwald, had been riding with them that night he must have been looking for something he already knew to be there.

So what did we have here?

Ostensibly, these objects—they could hardly be described as "aircraft"—exhibited characteristics similar to T.T. Brown's flying discs. Their historical interest lay in the fact that their "existence," if you could call it that, had been logged by observers with impeccable credentials more than three years before the first rash of flying saucer sightings in the United States. True, it was wartime and things got misidentified. But with the foo-fighters there was none of the hysteria that accompanied the U.S.-based sightings of 1947, which tended to make the testimony, if anything, more objective and believable. These people, many of them hardened by years of combat experience, felt they were encountering something new and dangerous in German skies. And the year 1944 seemed to be key.

Their experiences echoed what USAAF General Twining had told General Schulgen in his secret memo of September 1947. That "aircraft" making no apparent sound, with metallic or light-reflecting surfaces, exhibiting extreme rates of climb and maneuverability, "were within the present U.S. knowledge."

What did that mean exactly?

I didn't know, but somehow I felt the wording was key.

Over the next few months, life at *JDW* went on pretty much as it had done since I'd joined the magazine in the mid-'80s; a routine made up of press conferences, air shows, defense exhibitions and weekly deadlines. But as I maintained a close watch on developments that were shaping the defense and security climate of the post-Cold War world, I found my mind on other things. Things I kept to myself.

Had the Germans developed a totally new form of propulsion, blended it with a radically different kind of air vehicle, and deployed it in the form of some new and secret weapon system in the latter stages of the war? I had to juggle this possibility with the fact that T.T. Brown had postulated the basis of an antigravity propulsion system as far back as the 1920s.

I decided to discover whether any of this wartime sightings data was reflected in work the Germans had been doing in their research facilities and factories.

In the late 1950s, a book written by a German who had served as the commanding officer of a German Army technical unit in the Second World War, Major Rudolf Lusar, went on to become an unlikely best-seller in Britain and the United States.

The book was called *German Secret Weapons of World War II*.

Seeking clues to the foo-fighter mystery, I discovered a copy in the reading room of the Imperial War Museum. In it, Lusar described in meticulous detail, in language that often made the depths of his bitterness clear, the technical achievements of "a small, industrious and honest nation which lost the war."

Secret Weapons made somber reading. Although German technical achievements were visible in developments such as the V-1 flying bomb, a direct forerunner of the modern-day cruise missile, and the V-2 ballistic missile, it was the vast extent of Germany's underpinning technology base, as revealed by Lusar, which showed just how far ahead of the Allies the Nazis had been in certain key areas.

Jet engines, rocket engines, infrared and thermal-imaging systems, proximity fuses, missile guidance seekers . . . technologies that are

integral to most modern aircraft and airborne weapon systems were all listed and described. In the late 1950s, when Lusar's book first appeared, these technologies were still in their infancy in Britain and America.

Yet the Germans had been working on them a decade and a half earlier.

But there was another side to the book, one which was so sensational that immediately on its appearance it had set alarm bells ringing in Washington.

This side of the book related to so-called German "wonder weapons" beyond the V-1 and V-2.

One of these was the *Fleissiges Lieschen*, or "Busy Lizzie," a 150 m-long tube assembly of non-alloyed cast steel with chambers leading off it that lent it the appearance of a giant millipede. It was designed to fire 150 mm projectiles over distances of 170 km, more than enough to hit most British cities from sites deep within France. It never went into service, but 45 years after the war the components of an almost identical weapon, the "supergun," were intercepted by British customs officials on their way to Iraq.

Other esoteric developments detailed by Lusar showed that the Germans had been working on technologies for bringing down Allied aircraft with sound waves, air vortices, intensely focused beams of light and jets of compressed air.

In 1958, the U.S. Air Force commissioned a "special studies group" within Air Force Intelligence headed by an Austrian-born technical consultant called Dr. Stefan Possny to carry out a detailed appraisal of Lusar's book. The research effort was branded "secret" and has only recently come to light. I found it during a long, hard night's trawl in the basement when it popped out of cyberspace after an unusual combination of word commands fed through a high-power search engine.

A section in Lusar's book was devoted to "flying saucers" which he asserted, in no uncertain terms, were the product of German wartime inventors. "Experts and collaborators in this work confirm that the first projects, called 'flying discs,' were undertaken in 1941," Lusar wrote. He even went on to name the key individuals involved. These were "the German experts Schriever, Habermohl and Miethe, and the Italian Bellonzo."

Lusar described the craft in some detail. There were two principal centers of disc development: one, headed by Miethe in the vicinity of the Lower Silesian city of Breslau in modern-day Poland; the other centered on Prague in Czechoslovakia, then also an integral part of the Reich. The Miethe disc was described as a discus-shaped "plate" of 42 m diameter, fitted with "adjustable" jet engines. Shortly before the plant where the

craft was being built was overrun by the Russians, the retreating Germans blew it up, destroying the disc inside. However, many of the "experts" who had worked on the project were captured and taken back to Siberia, where work on them "is being successfully continued," Lusar reported.

In the paranoid U.S. security climate of the late 1950s, it was this aspect of the account that had led to the commissioning of Possony's special intelligence report.

According to Lusar, the other disc, developed by Schriever and Habermohl, did achieve flightworthy status before the end of the war—on February 14, at a facility just outside Prague. "Within three minutes they climbed to an altitude of 12,400 m and reached a speed of 2000 km/h in horizontal flight," the author related.

At a time when the most advanced interceptors of the day (the late 1950s) struggled to achieve such velocities, this was an outrageous claim, but then it didn't have to be taken wholly at face value to contain an element of truth.

Having covered the Stealth Fighter story in the mid-1980s, when the aircraft was still deeply secret and witnesses were reporting things they couldn't square with any known aircraft development of the day, I trusted the old maxim about smoke and fire.

And in this assumption, I had an ally in Dr. Possony. Why else had he been tasked by the U.S. Air Force with writing a classified critique of Lusar's claims? Why else, according to one source I spoke to about him, a researcher by the name of Joel Carpenter who had devoted considerable time to studying Lusar and Possony, had the good doctor shredded his conclusions on the subject?

So, alongside wartime pilots' reports of flying objects that defied their understanding of aircraft performance were accounts of Germans having worked on projects that had no engineering counterpart, even within present-day aerospace knowledge. In Lusar's book were names, dates and places—crude data, admittedly, but usable nonetheless as jumping-off points for a more rigorous search. And though Lusar's claims didn't point directly to evidence of antigravity propulsion technology, in his discussion of another taboo technology area in scientific circles, the flying saucer, he had outlined a pathway to an altogether different kind of aerospace platform.

Just as T.T. Brown had done in the 1920s—and Trimble had in 1956.

The task seemed so simple. Make the link between Germany and the flying saucer and here was an opportunity to solve not only the antigravity propulsion riddle, but, in the process, perhaps, one of

the most baffling mysteries of the 20th century: the origins of the UFO. If machines like those in General Twining's memo were "within the present U.S. knowledge," no wonder Trimble and his colleagues had so quickly fallen silent on the subject of a new and exotic propulsion source.

The flying disc must have exhibited performance so in advance of its time that it had been super-classified, then hidden in plain sight—behind the UFO myth—for the best part of 60 years.

Perhaps, too, it explained why Trimble had sounded like he'd had the fear of God put in him when he had been approached by Lockheed's PR machine on my behalf. The spooks would have made it abundantly clear there would be no statute of limitations on this particular secret. That you never talked about it. Ever.

Across multiple satellite relays, I heard the anguished shrieks of his three-month-old, the background clatter of his older children as they ran wild in the tiny apartment and finally relief in Lawrence Cross' voice as he realized it was me; a moment's respite in a whirl of copy deadlines, late-night feeds and trips to theme parks.

The baby's crying faded into the static as Cross took the handset into another room.

I thanked him for the tip about LaViolette and Valone, as a result of which I had pieced together the essential elements of T.T. Brown's life and work. Without elaborating on how I'd arrived there, I asked him what he knew about the Germans' purported development of flying saucers in World War Two.

"Oh Jesus," Cross said, and I formed a mental picture of him rubbing his eyes and reaching for a cigarette, "where do I begin?"

I prompted him. One minute the Nazis were tinkering with this technology, the next people were reporting sightings of these objects, first in Germany, then in America . . . ?

"It's not as simple as that," Cross said.

"Come on, Lawrence. A book called *German Secret Weapons of World War II*, by Rudolf Lusar details names, dates and places. Has anyone ever dug into this stuff? Have you? The story must be fit to bust wide open."

"It's been around for decades," he said, "long enough to have been given a name."

"What do you mean?"

"In the trade, we call it 'the Legend.' It looks so straightforward, doesn't it? A story with a solid trail. But it's not like that at all. When you get into this stuff the trail goes everywhere and nowhere. The people who really did exist are long dead and others probably never existed at all. I

know. I've been there and I've looked for them. So have dozens of other investigators. The detail is fantastic, but it's all uncorroborated. By that I mean there isn't a shred of evidence in any archive—no official word, no plans, nothing—that any of these projects ever existed."

"Have you got anything on it?"

He laughed and it caught in his throat. Cross smoked for the entire journalistic profession. He hacked the cough into the room, away from the receiver.

"Sure. How much time have you got? I'll email it to you. There's another book you should get ahold of. It's by an Italian. Guy by the name of Vesco—Renato Vesco. The book's called *Intercept—But Don't Shoot*. There's an English translation. I think it was first published in the late 1960s. Vesco's well up there, right in the heart of the Legend. Maybe you should try and track him down—that is, if he really existed."

I said nothing. Cross continued.

"Take nothing at face value and you're off to a better start than me. *Lusty* was real enough, but that's all I know. Everything else is up for grabs."

I thought I must have heard him wrong. "*Lusty*? Did you say, *Lusty*?"

"It stands for *Luftwaffe Secret Technology*. The U.S. Army Air Forces' official file on the state of the German Air Force's secret weapons work at the end of the war. The Air Force Historical Research Agency has a copy in its archives at Maxwell Air Force Base. It's raw data. That is to say, it's never been processed by anyone, so it's uncorrupted and verifiable, but I have no idea what's in it. In all these years, doing what we do, I've never had an excuse to go there. Maybe you do."

"Where is Maxwell Air Force Base?"

"It's in Alabama. Where the skies are so blue. A distant memory for you, I guess."

I thanked him, hung up and knew that I wouldn't be tapping Cross for any more advice. He'd made it abundantly clear, more in tone than anything else, that for him this was the end of the road.

I thought about the months ahead. I had no legitimate business in the States except for a trip to Washington to cover an Air Force symposium—and that wasn't until later in the year. Stretching the trip to include a visit down to Alabama was simply out of the question.

I called the Air Force Historical Research Agency at Maxwell Air Force Base to see if I could get a copy of *Lusty* sent to me. In theory I could, but the files were held on a number of microfilm reels and, I was told, it could take months, perhaps years, for the release forms to plow through the system and for the report to reach me. Not only that, but the

Lusty files were old and corrupted, making repros difficult. The best, probably only way to view them, the administrator there told me, was in situ.

I put the phone down, kneaded my eyes and thought about smoking a cigarette again; something I hadn't done in years.

I had just hit another pluperfect dead end.

Chapter 5

Flugkapitän Rudolf Schriever, one of the four saucer engineers cited in Lusar's book, began talking to the West German media in 1950 about a truly fantastic flying machine he had worked on for the Nazis—one that would have changed the course of the war had it gone into full-scale production. It stemmed from work that he had allegedly undertaken for the Heinkel Aircraft Company at Marienehe, near Rostock on the Baltic coast.

Schriever was one of the four "scientists" mentioned in Lusar's book as having worked on the Nazi flying discs. He was also, I found out through the massively detailed file that my former journalistic colleague Cross sent me via email, central to the Legend.

Though he started out as a pilot in the Luftwaffe, Schriever appeared to have developed some highly advanced ideas about aircraft that could take off and land vertically and it was in this capacity, after he had been drafted to Heinkel's design section, that he soon came to the attention of company chairman Professor Ernst Heinkel, who in early 1940 encouraged him to construct a small flying prototype.

So far so good. Though best-known for its lumbering He 111, the Luftwaffe's mainstay bomber during the Blitz against London in 1940 and for much of the rest of the war, Heinkel was one of the most pioneering and innovative aircraft companies within Germany at the time.

Whatever the Legend said, this much was fact.

In 1936, Ernst Heinkel began funding experiments that three years later would lead to Germany's successful construction of the world's first jet-powered aircraft, despite the fact that it was the British designer, Frank Whittle, who had first invented and patented the concept. The tiny, one-off Heinkel He 178 first flew on August 27, 1939, five days before the German Army marched into Poland. Eighteen months later, Heinkel would again eclipse all other aircraft companies by flying the world's first jet-powered fighter, the He 280. If anyone was to develop something as radical as Schriever's idea, therefore, Heinkel was the company to do it.

The Legend then takes over.

In the spring of 1941, Schriever's blueprints were being used to construct a "proof-of-concept" model in a "garage" away from prying eyes. Officially known as the VI (V for "Versuchs" or "experimental version 1"), informally it was referred to as the "Flying Top." It was probably no more than two or three feet in diameter and powered by an electric motor or a small two-stroke engine. It is not known where this garage was, but within the sprawling complex of buildings at Marienehe, set in three square kilometers of the Mecklenburg State Park, there was ample room for Schriever's esoteric little engineering project to be hidden from view. Furthermore, it fitted into the Heinkel way of doing things. During the development of the world's first turbojet, Ernst Heinkel had installed Dr. Hans-Joachim Pabst von Ohain, a gifted graduate of the University of Göttingen, in the same kind of environment—a converted garage at the university—before transferring the fruits of von Ohain's labors, the revolutionary HeS 3A turbojet, to a secure facility within the Marienehe site.

By June 1942, Schriever's Flying Top had been test-flown and the results deemed sufficiently interesting to secure top secret funding from the RLM, the Reichsluftfahrtministerium or State Air Ministry.

With RLM funding, the intention was to construct a full-scale piloted version capable of controlled vertical takeoff and landing. Construction of this full-size version, the V2, began at Marienehe in early 1943. The V2, which was known as the "Flugkreisel" or "Flightwheel," had a diameter of approximately 25 feet, its power generated by one or perhaps two Heinkel-Hirth jet engines, depending on which version of the Legend you want to believe. The V2 supposedly flew with Schriever at the controls, but as a piece of technology it was deemed to be heavily overengineered and was quickly scrapped. As a proof-of-concept vehicle, though, it seems to have served its purpose, because shortly afterward, Schriever and his team relocated to Czechoslovakia where they set about constructing a larger and altogether more sophisticated prototype known as the V3.

With the Allied aerial bombing campaign now at its height, their activities were dispersed around the Prague area to minimize the exposure to the relentless air attacks, by now penetrating deep into the Reich. But the bulk of the team's work was centered on a restricted area of a satellite facility outside Prague belonging to the Munich-based Bayerische Motorenwerke engine company, better known today as BMW. Despite the existence of Heinkel's own jet engines, the real cutting edge of German gas-turbine research was centered on BMW and in particular on its Bramo division, located at Spandau in Berlin. Bramo,

the Brandenburgische Motorenwerke, had been bought by BMW from Siemens in 1939. By the middle of the war, BMW-Bramo had 5,000 staff working full-time on gas-turbine research alone—a discipline then barely a decade old—and was ultimately responsible for the BMW 003 jet engine, the best turbojet of the day, which powered the Me 262, the world's first operational jet fighter, and the Ar 234, the Luftwaffe's advanced jet-propelled reconnaissance-bomber, another aircraft far ahead of its time.

It was from Spandau, supposedly, that Klaus Habermohl, the second disc engineer mentioned by Lusar, was recruited to the Schriever team. Habermohl's job in Prague was to integrate the disc with a new and radical form of power plant called the radial-flow gas-turbine, or RFGT. Unlike Brown's electrogravitic motor, the RFGT was, at least, recognizable technology by modern standards, if extraordinary. It was essentially a jet engine. However, unlike a regular jet power plant, with its compressors, combustion chambers and turbines mounted one behind the other in what was basically a big tube, the RFGT formed part of the airframe itself, with the whirling turbomachinery rotating around the aircraft's centrally mounted cockpit. As such, "aircraft" did not adequately describe what the machine actually looked like. There was only one configuration to which an RFGT could possibly be adapted: that of a flying disc or saucer.

By the autumn of 1944, the V3 is said to have been completed. With German airfields under constant attack from Allied daylight bombing, the prospect of a fighter or bomber that could take off and land vertically from any dispersed site would have been exactly in line with Luftwaffe requirements. However, due to "an administrative change," the V3 program was abandoned in favor of a further prototype, the V7, propulsion coming from another experimental RFGT from BMW-Bramo. The V7 supposedly had a diameter of 60—70 feet and a crew of two or three.

Applying a crude rule of thumb, based on a rough estimate of the disc's weight, Habermohl's ingenious RFGT would have had to have generated around 10-15,000 pounds of thrust to have made the design of the disc in any way viable. This was the machine that supposedly test-flew on February 14 with Schriever and Habermohl at the controls, achieving 2,000 km/h in level flight.

The fastest aircraft of the day, the little rocket-powered Me 163, struggled to attain half this speed.

Lusar's third and last German saucer scientist, Dr. Richard Miethe, was supposedly working on another disc project at a subterranean facility

near Breslau under the auspices of an altogether separate contract. Toward the end of the war, the legend stated that Miethe was drafted from his activities in Breslau to assist with the Schriever/Habermohl disc in Prague, an indication, perhaps, that the Schriever/Habermohl disc was the better bet in a procurement environment that was by now desperately short of money, skilled labor and raw materials. This streamlining coincided with the "administrative change" that had led to the abandonment of Schriever's V3 design for the altogether more viable V7.

The V7, then, seems to have been the result of a three-way endeavor, although there were reports that a Miethe disc—based, perhaps, on the project he abandoned at Breslau—was captured by the Russians, along with a number of engineers and scientists, when they drove the Germans out of Poland.

As for the V7, some say it, too, was acquired by the Russians when they took Prague; others that it was blown up by the Waffen-SS on May 9, 1945, the day hostilities in Europe ended. The Legend had it both ways.

The detail in the Schriever/Habermohl/Miethe legend was rich and impressive. Here were names, dates and places—minutiae even, that seemed to corroborate everything Rudolf Lusar had written down. The trouble was, the data was based heavily on the say-so of Schriever, who was long since dead; the rest had magically appeared out of thin air, just as Cross had said. No one knew where the detail emerged from. Over the years it had been passed down from one researcher to the next, with no apparent attribution. When I approached BMW's archivists, for example, they denied that there had been any BMW factory near Prague "engaged in advanced aircraft projects including design or advanced research during World War II."

I began to see what Cross meant. Cross and I were used to sourcing every piece of information we ever came by. This stuff was unverifiable. What I needed was hard, solid proof; and within the Schriever portion of the Legend there was nothing to hang a hat on, beyond the fact that the man himself had existed.

Unfortunately, the same could not be said of the other characters in the story. All attempts by researchers to trace Miethe and Habermohl had foundered, although there had been occasional "sightings" in the lore that had grown up around them.

Miethe is supposed to have escaped Czechoslovakia in early May and to have headed west, eventually making contact with U.S. technical intelligence teams operating inside Germany. Herded into a "pen" along with Wernher von Braun and his fellow rocket scientists, Miethe was said to have been taken to the United States, ending up at Wright Field, the

USAAF's premier research and development center near Dayton, Ohio. If any of this actually happened, there is no trace of it.

Habermohl is said to have been captured by the Russians at the Letov factory, a German-administered aircraft plant outside Prague, and, after a period of detention, sent to work at a top secret Soviet aircraft design bureau east of Moscow.

Again, no one could say for sure whether he had even existed at all.

My research showed the Italian, Bellonzo, was real enough, except Lusar had misspelled his name. In 1950, Professor Giuseppe Belluzzo, a former industry minister in Mussolini's cabinet, started talking about disc-shaped "flying bombs" that he claimed to have worked on during the war and passed on to the Germans, who had subsequently developed them into working prototypes.

Belluzzo was also convinced that these weapons were the basis of the flying-saucer sightings that had gripped much of America for the best part of three years and that they were now under further development inside the Soviet Union. Beyond the fact that Belluzzo, like Schriever, was real, his claims also remain unverifiable. Interestingly, though, he started talking to the media just a few days before Schriever, leading some researchers to reason that Schriever, a man who like so many other Germans in 1950 was struggling to make ends meet, had invented his entire story.

In fact, the Legend lacked a single item of corroborating data. There was nothing in any archive or museum, no photograph, no indisputable piece of testimony, to say that any of it was true.

There were splits and schisms in the Legend, just as there are orthodox and unorthodox branches within major religions. The other strand to the myth that I had to pay attention to was Vesco's. I had managed to obtain a copy of his book, *Intercept—But Don't Shoot* (published 1971), and scrutinize it. What was beguiling about Vesco's account was the certainty with which he presented his case. Vesco, who was 22 years old when the war ended and said to have been well connected with technical experts within the Italian Air Force at the time he wrote *Intercept*, claimed that he pieced together his account of the top secret development effort behind the foo-fighter program from sources inside the Italian military and from Allied intelligence reports published after the war.

What also made Vesco's account worth more than a cursory glance was the fact that it "detailed" a completely different development effort from the Schriever/Lusar account. Vesco claimed that there were two kinds of foo-fighters. One was unpiloted and remotely controlled; in effect, a

flying bomb designed to knock down enemy aircraft. The other was manned, but flew in anger only once before hostilities ended.

The unpiloted version was called the Feuerball (Fireball) and it blended a number of highly advanced technologies.

"In the autumn of 1944," Vesco wrote, "in Oberammergau in Bavaria, the OBF—an experimental center run by the Luftwaffe—had completed a series of researches into electrical apparatus capable of interfering with the operation of an engine up to a maximum distance of about a hundred feet by producing intense electromagnetic fields."

In parallel, a separate effort was under way by the Germans to produce a "proximity radio interference" device capable of jamming or spoofing Allied radio and radar systems. Put these two technologies into a small, circular, armored airframe "powered by a special turbojet engine, also flat and circular and more or less resembling the shell of a tortoise" and "a highly original flying machine was born."

Radio-controlled at the moment of takeoff, the machine was steered toward Allied bomber streams by a ground operator, whereupon it automatically latched onto their slipstreams, "attracted by their exhaust flames, and approached close enough without collision to wreck their radar gear."

And then came the detail. The Fireball was first constructed at an aircraft plant at Wiener Neustadt, south of Vienna, with the help of the Flugfunk Forschungsanstalt of Oberpfaffenhoffen (FFO), an aircraft electronics firm near Munich. Hermann Goering, Hitler's deputy, inspected progress on the weapon "a number of times," hoping that the principle of the Fireball could also be used to produce "an offensive weapon capable of revolutionizing the whole field of aerial warfare.

"The fiery halo around the perimeter—caused by a very rich fuel mixture—and the chemical additives that interrupted the flow of electricity [in Allied aircraft] by overionizing the atmosphere in the vicinity of the plane, generally around the wing tips or tail surfaces, subjected the H2S radar on the plane to the action of powerful electrostatic fields and electromagnetic impulses (the latter generated by large klystron radio tubes protected with anti-shock and anti-heat armor). Since a metal arc carrying an oscillating current of the proper frequency—equal, that is, to the frequency used by the radar station—can cancel the blips (return signals from the target), the Feuerball was almost undetectable by the most powerful American radar of the time, despite its nighttime visibility."

Once again, I pictured the desperation felt by Schlueter as his radar operator, Meiers, failed to register the orbs of light ahead of the Black Widow on the SCR540.

Was this what they had encountered that night, the Fireball?

To believe that it might have been, I had to accept that what we had here was a weapon system that was decades ahead of its time. Years before the deployment of the surface-to-air guided missile—sufficiently perfected to shoot down the CIA U-2 pilot Gary Powers over Russia only in 1960—and decades before radar-evading stealth technology would become a household word, the Vesco portion of the legend held that the Germans had developed a single weapon system that blended these and other exotic technologies. Not only that, but the Fireball was able to home in on its intended prey, the Americans' B-17 and B-24 bombers, via an automatic guidance system—one that tracked their exhaust plumes—a sensor so advanced that it was unheard of, even with the benefit of modern-day knowledge. Once close to the bombers, it could then either disrupt their electronics or stop their engines in flight with a gas that killed their ignition systems.

At a stroke, Vesco had assembled all the evidence he needed to explain the foo-fighter conundrum. He also threw in some eyewitness testimony. "One person who saw the first short test flights of the device, without its electrical gear," he wrote, "says that 'during the day it looked like a shining disc spinning on its axis and during the night it looked like a burning globe.' "

The trouble was, it sounded like science fiction.

When the Russians advanced into Austria, Vesco maintained, the Fireball production line was moved from Wiener Neustadt into underground facilities run by the Zeppelin Werke in the Schwarzwald—the same area of Rhineland and Black Forest where Schlueter, Ringwald and Meiers, and a host of other aircrew from the 415th Night Fighter Squadron, had first encountered the inexplicable.

Vesco went on to explain that the Feuerball had an "older brother," the Kugelblitz or "Ball Lightning Fighter," that was built and flight-tested—once—in the vicinity of Kahla, site of a huge underground weapons development complex in Thuringia, a region of mountainous uplands in the heart of Germany known as the Harz. If this "fact" also was a hoax, it was a clever one, since the Harz certainly had housed a number of underground Nazi weapons factories, among them the facility at Nordhausen, where Wernher von Braun's V-2 rockets had been produced.

Vesco was notoriously reclusive. Some researchers believed he'd never existed at all; lending credence to the supposition that *Intercept* really was just an elaborate hoax. But Vesco was real, all right. I managed to trace a group of Italian researchers who had been in contact with him up until

his death in November 1999. According to them, Vesco didn't care whether people believed him or not.

"He was very correct," one of them told me over the phone, "so it's very hard to believe that he made it all up."

But where was the proof?

Vesco claimed that the evidence for Feuerball and Kugelblitz was to be found within obscure tracts of the British Intelligence Objectives Subcommittee (BIOS) reporting system and its successor, the U.S.-U.K.-administered Combined Intelligence Objectives Subcommittee (CIOS), published in the immediate aftermath of the war (a number of reports from which, however, are still withheld). BIOS and CIOS were the systems employed by the British and the Americans for assessing German high technology. But researchers had been through all the available CIOS and BIOS files with a fine-tooth comb and had found nothing that pointed to anything that described the Fireball or its "older brother" the Ball Lightning Fighter.

Vesco had stuck to his guns, never wavering from the conclusions he had espoused in his book: that the Germans had developed a truly revolutionary new form of air vehicle; that it was the British who had happened upon the technology at the end of the war and that they, together with Canadian scientists, had refined it in the frozen wastelands of British Columbia and Alberta.

Vehicles resulting from this endeavor, Vesco maintained, were responsible for the rash of U.S. flying-saucer sightings in 1947.

A familiar pattern was emerging and it wasn't helping my case. The more I looked, the more "evidence" I found that the Germans had been tinkering with technology that explained the foo-fighters sightings of that winter of 1944⁵.

From Schriever in the 1950s to Vesco's testimony in the 60s—and others before, during and since—maybe a dozen Germans had come forward to say that they had worked on flying-saucer technology under the Nazis. And some, like Viktor Schauberger, were so very nearly believable.

Schauberger, whose story was contained in notes on the Legend sent to me by Lawrence Cross, was an Austrian who had supposedly invented a totally new form of propulsion based upon a principle called "implosion." No one I spoke to seemed entirely sure what implosion meant, but according to the stories that had grown up around this man, the implosion process was at the heart of a radical turbine that Schauberger had installed in a sub-scale flying disc sometime during the war. A test of

this small flying vehicle, with its echoes of the Schriever Flying Top, had supposedly taken place and the results were said to have been highly impressive. In one account of the test, the craft had apparently risen toward the ceiling of the test facility "trailing a glow of ionization." This immediately elevated the report above the many others I had come across, for it signaled that, whatever was occurring within the implosion process, it had precious little to do with jet propulsion. If true, it could only have been an antigravity effect.

If true.

Though Schauerberger was long dead, his son Walter was still alive when I telephoned him at his home in Austria in 1991. Talking through an interpreter, I asked Walter about his father's experiments and got a disconcerting answer. I had imagined that Viktor had attended some impressive German or Austrian technical institute; that he was a professor of physics at Vienna or Salzburg or, at the very least, an eminent aerospace engineer. But no. Walter Schauerberger told me that his father had developed his radical ideas about energy and propulsion by observing what he had seen in Nature, the way rivers flowed and fish swam. His only professional training had been as a forester.

This was the trouble. While all of these stories were laced with detail, which immediately gave them a veneer of credibility, they were almost always let down either by corrupted data or a complete absence of it. In a decade of investigating the aerospace industry, I had never once come across a designer who did not have a set of initials after his name. A forester was simply absurd.

Had there been more to the Schauerberger story, I would not have hesitated to make the trip to Austria. But as decisions went, this was an easy one. I was using my training to make value judgments all the time on the data that was gathering in my basement. You didn't need a degree in physics, though, to appreciate that whatever they had or hadn't been, foo-fighters would have had to have come from the minds of engineers—not from a man who'd spent his time among Alpine forests and streams.

It was for this reason that I gently declined Walter Schauerberger's offer to visit his "biological-technical institute" in the Salzkammergut Mountains.

For any of the "evidence" about German flying saucers to be irrefutable, it had to emerge from official documentation. But Lawrence Cross and a number of other credible researchers I'd contacted on the subject had already been through the BIOS and CIOS reports and found nothing. And that should have been the end of it. Yet, I knew that if the Germans

had developed a revolutionary air weapon, based perhaps on some radical form of propulsion technology, it was inconceivable that the CIOS or BIOS intelligence teams would have documented the discovery for the world to read about.

To go the extra mile, I realized I'd have to access the original intelligence data on which the BIOS and CIOS assessment teams, most of whom were technical types sat at desks in Brussels, Paris, London and Washington, had based their analyses. Here again, I knew there would be no smoking gun. Had intelligence units in the field come across anything meaningful, these reports, too, would have been sanitized. Anything as obvious as a full-blown air vehicle or a new form of propulsion system would not have escaped the censor's attention.

But to identify evidence of a new form of aerospace technology, particularly one as far-reaching as that described in the legend or by Vesco, I wasn't searching for the obvious, because the obvious would have been picked up by the censors.

As with any new piece of technology, developments came together as a series of systems and subsystems. There would have been prime contractors and subcontractors, some of them working in new scientific areas—areas that would have been on the very edge of the technical knowledge of the deskbound BIOS and CIOS assessors.

Between the chaos of the front line and the overstretched resources of the intelligence units tasked with assessing German hardware and documentation—which the victors had shipped out of Germany by the ton-load—there might still, I figured, be some fresh evidence; something useful that had been overlooked.

I started with the British, whose initial stab at organized technology plunder was vested in the hands of a ragtag private army—composed, bizarrely, of sailors and Royal Marine commandos. This outfit was the brainchild of a certain Commander Ian Fleming, who 15 years later would go on to create the Bond persona in the double-oh-seven novels. Bond's character, it appears, drew heavily on the exploits of 30 Assault Unit RN, which rode roughshod over the conventions of the day. Following the battle for Cherbourg, in which 30 AU RN had been tasked with capturing German naval headquarters, the marines liberally enjoyed the spoils of war. Their behavior was described as that of "merry courageous, amoral, loyal, lying toughs, hugely disinclined to take no for an answer from foe *orfraulein*"

As a result of their "martial exuberance," as one assessor wrote, 30 AU RN was reined in somewhat and renamed 30 Advanced Unit RN. It was

also ordered to subordinate its activities to a new and much larger force of British tech-plunder units, known as "T-Forces." These would move forward with General Bernard Montgomery's 21st Army Group, principally in Monty's theater of operations, northwest Germany, but not exclusively so. Under the terms of the CIOS charter, with its Anglo-American task jointly to exploit German spoils of war, British T-Force teams would also be allowed to tag along with forward U.S. Army units in pursuit of their objectives.

These objectives were to locate and secure intact technical "targets" of interest; to preserve German high technology from "destruction, loot, robbery and, if necessary, counterattack," until the completion of their examination by teams of experts or until their removal. They were also to act as armed escorts in enemy territory for the "expert investigators" drawn from CIOS offices far behind the front lines. As a quid pro quo, U.S. technical teams could ride with forward British assault units. The eventual size of British T-Forces would grow to 5,000 personnel.

But from the reports generated by these units, now freely available in the U.K. Public Records Office, it was apparent that the British were desperately ill-prepared to make the most of the opportunity that lay before them.

Over the next two months, I spent every spare moment down at the Public Records Office. And when I wasn't in the Records Office, just down the road in Kew on the outskirts of London, I was gathering as much open-source reading material as I could on the Allies' systematic plunder of German high technology at the end of the war. One book, *The Paperclip Conspiracy* by Tom Bower, read like a manual on how to dismantle an entire nation's technology base. If America, Britain and their allies had applied the lessons of *Paperclip* to the Iraqi problem at the end of the Gulf War in 1991, the threat posed by Saddam Hussein's war machine would have been eradicated forever.

During the November 1944 to March 1945 planning stage of the T-Force operations, the problem facing British investigators was a fundamental lack of intelligence on what they were supposed to be looking for. A "black list" of technical targets was drawn up, the majority of them weapons-development and research centers put forward by CIOS.

But a general air of ignorance of the situation on the ground persisted, as a T-Force field commander later recorded. "It appeared that the sponsoring ministries knew little or nothing about the specific whereabouts and natures of their targets; and that investigators who would eventually come out would know even less."

If there was any sense at all that the British realized they were in a race against the other allies—the French and the Americans particularly—it came far too late. Roy Fedden, a senior and respected British aviation industrialist who flew into Germany to view the tech-plunder operation for himself, complained that his compatriots were being "lamentably slow" to take advantage of the information on offer. With many of the U.K. technical experts drawn from industry, part of the problem was a considerable reluctance on their part to accept that Germany could possibly have been as far ahead of British technology in certain pioneering areas—the jet engine, for example—as they evidently were.

There were British successes—the capture of the German navy's laboratories at Kiel, for example, where highly advanced German U-boats and torpedoes, propelled by an innovative peroxide-fueled engine, were being developed. There were also significant British finds at Krupp in Meppen, where advanced armor and artillery technology had been produced for the Wehrmacht.

But reading between the lines of the British-controlled tech-plunder operation, its disparate character was clearly a factor in its limited success. Without adequate direction from London, there was confusion as well as a palpable lack of urgency about the operation at the front. And in some cases—the activities of T-Forces of the 21st Army Group's Canadian First Army, for example, who were roaming between Germany and Holland, looking for targets—it was portrayed as being akin to a jaunty adventure.

Among the most prominent targets on First Army's T-list, according to records of its operations at the time, was the radio transmitter of Lord Haw-Haw, the British traitor and Nazi propagandist. "Two further wireless stations had been discovered on the (German) island of Borkum and another platoon was dispatched there as guard. That platoon was in the happy position of having 4,500 German marines to wait on them!"

It all seemed a very far cry from the exotic weaponry that Vesco had described in *Intercept—But Don't Shoot*.

In scanning the British documents, however, I'd found one big problem with Vesco's thesis that the British had discovered flying disc technology and removed it to Canada. By far the greater part of Germany's advanced aerospace development had taken place in Bavaria in southern Germany, a long way from the main thrust of British operations in the northwest of the country.

And while some British T-Force units had been integrated with American forward army operations in central and southern Germany, they were more often than not outwitted by the Americans, who time and

again seemed to have had the resources to take what they wanted, much to the dismay of Roy Fedden.

"The Americans," he wrote, "have fine-combed the country, removing considerable quantities of drawings, technical records and actual equipment direct to the States."

This, I could see, was no mere happenstance. American tech-plunder activities were the result of the most carefully thought out strategy in the history of U.S. military operations, its orchestration planned at the highest levels.

In November 1944, the U.S. Joint Chiefs of Staff had established a Technical Industrial Intelligence Committee to seek out anything in Germany that might be useful to the postwar American economy. Nor did the Joint Chiefs' target list just comprise military objectives. One subcommittee had a staff of 380 specially trained civilians set up specifically to represent the interests of 17 U.S. companies. Special agents from the U.S. Field Intelligence Agencies (Technical) scoured Germany for vacuum tubes a tenth of the size of the most advanced U.S. devices, and condensers made out of zinc-coated paper, which were 40 percent smaller and 20 percent cheaper than U.S. condensers—and, instead of "blowing" like the U.S. vacuum tubes, were "self-healing"—in other words, they could repair themselves. Such innovation would later prove invaluable to the postwar U.S. electronics industry.

The teams were also on the lookout for German textile and medical advances—and found them by the ton-load. At the German chemical giant LG. Farbenindustrie, notorious for its role in the development of the gas chambers of the Holocaust, investigators found formulas for the production of exotic textiles, chemicals and plastics. One American dye authority was so overwhelmed by the discovery that he declared: "It includes the production know-how and the secret formulas for over 50,000 dyes. Many of them are faster and better than ours. Many are colors we were never able to make. The American dye industry will be advanced at least ten years." German biochemists had also found ways of pasteurizing milk using ultraviolet light and their medical scientists had discovered a way of producing synthetic blood plasma on a commercial scale.

Hundreds of thousands of German patents were simply removed and brought back to America.

No wonder the Brits were struggling. It was clear from these and other accounts that CIOS, the Anglo-U.S. reporting channel and assessment office for German high technology, was little more than a front. The real American tech-plunder operation had been organized long beforehand, under separate cover, in Washington.

A year after the war ended the U.S. Office of Technical Services (a body set up to ensure that German technology, where required, was spun rapidly into American industry) was reportedly sifting through "tens of thousands of tons" of documentation. This "mother lode" of material, according to one contemporary report, "very likely contained practically all the scientific, industrial and military secrets of Nazi Germany." Had exotic propulsion science been buried somewhere within this mountain of raw material, it would have been almost impossible to locate. But after the war, the mother lode was divided between the Library of Congress, the Department of Commerce and the technical archives at Wright Field. Most of the Wright Field material had come via *Lusty*, the postwar intelligence report on Luftwaffe technology that Lawrence Cross had told me about. This was the report that was stuck, frustratingly, in Alabama—nowhere near the Air Force symposium that I was due to hit in a quickie assignment to Washington later in the month.

I called the Air Force Historical Research Agency at Maxwell Air Force Base in Alabama again to see if the administrator there could recommend any other sources of information comparable to the *Lusty* report. Since I was going to be in the vicinity of the U.S. National Archives in Maryland, I wanted to be sure I wasn't missing out on anything.

"If you're gonna be in D.C. anyway," she told me, chewing slowly on the syllables, "you might want to check out the other copy of *Lusty* that's available at our sister site down by the Old Navy Yard, just across the river from Capitol Hill."

For a moment, I thought she was joking. This was the person who'd told me, in painstaking detail, that if I wanted to obtain a hard copy of *Lusty* from the archives of Maxwell Air Force Base it would take me months or even years; that due to the poor condition of the microfilm reels, the only viable way of accessing the information was by reading the reels *in situ*.

When I pointed this out, her answer was nothing if not honest.

"You never mentioned nothin' 'bout no trip to D.C.," she said.

The Office of Air Force History maintained an out-of-the-way archive in Washington, D.C., located at Boiling Air Force Base, next to the Old Navy Yard.

I was supposed to be in and out of Washington inside a couple of days. I stretched the schedule and bought myself another 24 hours.

If there were trace elements of a suppressed technology within the U.S. documents plundered from Nazi Germany, I figured they had to be buried in *Lusty*.

Chapter 6

It was still dark when I exchanged the warmth of the Holiday Inn for the damp chill of the predawn, heaved my bags into the back of the cab and hunkered down for the 30-minute ride across the river. I had around eight hours to sieve *Lusty* for evidence of a technology Cross maintained couldn't be substantiated.

The commuter traffic was jamming the bridges, the top of the Washington Monument scraping the overcast sky, as the taxi headed across the Frederick Douglass Memorial Bridge and into a part of the nation's capital that only makes the news when the body counts from the drive-by shootings top three or four.

By the time I reported to the guard at the main gate, the wind had got up, lifting sheets of spray from a choppy confluence of gray wave caps where the Potomac and Anacostia rivers meet off Greenleaf Point. The spray snaked in rivulets this way and that in front of us as the cabdriver navigated the maze of open streets looking for the address I'd been given. It took him 15 precious minutes to find it.

The archive was tucked away in a forgotten corner of Boiling Air Force Base, home of the Defense Intelligence Agency and the USAF's Office of Special Investigations. It wasn't the sort of place you'd want to stroll or take the view.

Boiling was a large cordoned-off military district sandwiched between the black waters of the Potomac and the badland housing projects of Congress Heights and Washington Highlands. It also bordered the grounds of St. Elizabeth's—a hospital for the criminally insane, since renamed—where John Hinckley served time for his assassination attempt on President Reagan.

The reading room in the tiny Office of Air Force History was low-lit and windowless. True to her word, the senior archivist had placed a cup of coffee next to the warmed-up microfilm reader opposite the farthest row of metal book stacks. There was a clock on the wall above the door to her office. It was a little after seven-thirty. The air-conditioning had not

yet kicked in and my breath mingled as condensation with the steam rising from the cup.

The documents were contained on several reels of microfilm and, like the reels at the main repository in Alabama, they were in bad condition. My business schedule allowed for one shot at a read-through. My plane left at 6:15 that night.

To help me cut to it, the archivist had offered to open the office early and said I was welcome to turn up any time after seven. Coffee was on the house.

Time was already short when I slapped the first reel into the microfilm reader and got to work.

Lusty opened with a condensed history, the opening paragraph of which began:

"At a medieval inn near Thumersbach near Berchtesgaden [Hitler's mountaintop retreat in Bavaria], early in May 1945, the German General Air Staff patiently awaited the outcome of surrender negotiations taking place in the north. They had arrived by car and plane during the past weeks, when the fall of Berlin was imminent, and had kept in contact with Admiral Doenitz at Flensburg. Through the interception of one of these messages, their location, which had previously been unknown, was discovered. Within 24 hours, Lieutenant Colonel O'Brien and his small party, representing the Exploitation Division of the Directorate of Intelligence, USAFE (United States Air Forces in Europe), had arrived, located the party and conducted the first of a series of discussions with General Koller, who was then in command."

Colonel O'Brien's men were the advance guard of 200 officers chosen from HQ^AArmy Air Forces to oversee the USAAF tech-plunder operation and it was clear right from the start that they were in a race against time.

In the chaos of the collapsing Reich, many German scientists were dead, others had been captured by the Russians advancing from the east or the Americans, British and French in the west. Many were held in internment camps, but in a country brimming with former Reich slave-workers and displaced citizenry—almost all of whom had to be filtered by the Allies for Nazi party members and war criminals—it was hard to know who was who or where.

But the vast majority were still at their factories and laboratories when the advance units of the USAAF plunder operation screeched up in their jeeps and half-tracks. With no orders to stop working, they had carried on at their workbenches, even though the armed forces of the high

commands that had commissioned their work had either capitulated or been wiped off the map.

Many documents and blueprints were missing. German project managers would tell U.S. investigators that they had destroyed them under orders, but the USAAF officers quickly learned to apply some psychology of their own: men and women who had devoted years of commitment to technologies they considered vastly superior to those of their enemies were incapable of such vandalism.

Most of the files existed, but had been hidden.

Germany had already been divided by the Allies into zones of occupation; and many of the most technologically interesting German facilities—the underground V-1 and V-2 production sites in the Harz mountains, for example—were in the designated Russian sector.

Before the zones were locked and sealed the missing documents had to be located and brought under American control.

By a process of detective work and, where that failed, by persuasion and coercion of the German scientists and program managers, boxes of blueprints and notes were pinpointed, then recovered from the bottom of lakes, as well as caves, farms, crypts, hospitals and mines.

O'Brien's team hit pay dirt early: their sleuthing led them from Thumersbach to an air raid shelter covered with earth in the side of a mountain close to the Austrian border. There they found files belonging to the 1 Gruppe/6 Abteilung, the German Air Ministry's intelligence directorate. These detailed all the Luftwaffe's latest air weapons, from the Me 262 jet fighter and the Me 163 rocket fighter to radars, air-to-air missiles and guided bombs. It also showed that the blueprints had recently been smuggled out of Germany in U-boat shipments to Japan.

Reading through the signals transmitted by *Lusty* field team officers to their superiors, I was hit by the sheer scale of the operation and the pressures endured by those who took part in it.

On April 22, 1945, two and a half weeks before the guns in Europe fell silent, additional recruits were needed to process the data. "It is planned to expand the activities of air technical intelligence tenfold, securing the most highly qualified specialist personnel available to the Army Air Forces," Brigadier General George C. McDonald, head of USAAF Intelligence, wrote that day.

The caliber of the technological spoils was indicated by the arrival in-theater at the end of that month of a "special group of scientists" headed by Dr. Theodore von Karman, special consultant to the U.S. Army Air Forces' supreme command.

The hardware that awaited them was detailed on several pages of

microfilmed documentation from McDonald's intelligence directorate. Among them were a jet-propelled helicopter "in fly able condition accompanied by a complete set of documents and detailed drawings"; the Lippisch P-16 tailless rocket-propelled research aircraft whose advanced construction indicated "possible operation at high Mach numbers in the vicinity of 1.85"; and the Horten Ho 229 twin-jet flying wing bomber.

Nothing comparable existed in the American inventory; or anywhere else, for that matter.

But by midday, I had still found nothing that indicated the merest presence of the machines elucidated within the Schriever/Vesco legend.

As more men and resources were plowed into *Lusty*, the search for high technology widened. The files detailed the growing number of shipments back to the U.S. "Hanau: 50 tons of documents ready for shipment. Munich: 30 tons of documents ready for shipment. Teisendorf: Approximately two tons of documents pertaining to flight-path studies of guided rocket missiles are on location ready for shipment..." From all over Germany, documentation was being gathered, marshaled and hauled back to the U.S.

But just when it seemed as if the USAAF mission was breaking the back of the task, new finds and new problems added to the logjam. The biggest and most unexpected of these related to the Nazis' underground facilities, whose scale and number had not been anticipated by the investigators, because Allied reconnaissance had failed to pick up anything but the vaguest hints that the Germans were putting their factories underground.

The underground facilities were of more than passing interest to me since they figured in the Legend as places where the Nazis had developed their antigravity technology.

From the message traffic that built up around the discovery of these facilities in the late summer of 1945, three or four months after the war in Europe ended, it was clear that the discovery of these underground facilities was diverting the *Lusty* investigators from their primary goal of plundering German technology.

On August 29, General McDonald sent USAAF headquarters in Europe a list of six underground factories that had been discovered and "excavated." All of them had been churning out aircraft components or other specialized equipment for the Luftwaffe until the very last day of the war.

According to McDonald, the facilities varied in size from 5 to 26 kilometers in length, according to measurements of their tunnels and galleries. The dimensions of the tunnels varied from 4 to 20 meters in

width and 5 to 15 meters in height; the floor space from 25,000 to 130,000 square meters.

Seven weeks later, in mid-October, in a *Preliminary Report on Underground Factories and Facilities in Germany and Austria*, senior USAAF officers were told that the final tally showed "a considerably larger number of German underground factories than had hitherto been suspected."

In addition to Germany and Austria, the underground building program had been extended across France, Italy, Hungary and Czechoslovakia.

"Although the Germans did not go underground on a large scale until March 1944, they managed to get approximately 143 underground factories into production by the last few months of the war," the report stated. A further 107 facilities had been located that were either being built, excavated or planned by the end of hostilities, but another 600 sites could be added to the total if caves and mines, many of which had been turned into production lines and weapons laboratories, were taken into consideration.

The report's author was evidently taken aback by the breadth of the German underground plan. "It is a matter of conjecture what would have occurred if the Germans had gone underground before the beginning of the war," he concluded.

And then, hot on the heels of the underground investigation, a fresh directive, this time from a senior USAAF field officer to General McDonald, the air force intelligence chief at Wright Field in Ohio.

Set against the steady but predictable buildup of documentation, its tone was so unexpected and the content of the message so different that it took me a moment to come to terms with its implications. It was dated September 28, 1945:

1. It is considered that the following have been thoroughly investigated and have proven to have no basis of fact.

a. Remote Interference with Aircraft

Investigations have been completed on this subject and it is considered that there is no means presently known which was in development or use by the German Air Force which could interfere with the engines of aircraft in flight. All information available through interrogations, equipment and documents has been thoroughly investigated and this subject may be closed with negative result.

b. Balls of Fire

As far as can be determined from extensive interrogations, investigation of documents, and field trips, there is no basis of fact in the

reports made by aircrews concerning balls of fire other than that phenomena similar to balls of fire may have been produced by jet aircraft or missiles. This subject may be considered closed with negative results.

A.R. Sullivan Jr.,
Lt. Colonel, Sig. C.

This was the first hint or mention of anything connected with the foo-fighters. Sullivan was telling McDonald that investigations on the subject had been completed; and yet, there was no sign in any of the raw communications traffic that had hitherto flowed between the front and Wright Field that anyone was the slightest bit interested in the subject of "remote interference with aircraft" or "balls of fire." Furthermore, there wasn't any suggestion of interrogations having taken place or equipment or documents having been investigated pertaining to possible foo-fighter technologies—or of special "field trips" having been undertaken to track them down. But the instruction that the subject "may be considered closed with negative result" was so strident it made me sit up and wonder if I'd missed something.

I felt a knot of excitement in the pit of my stomach. The subject, as Sullivan put it, *may be closed with negative result*.

Why?

In the apparent informality of its memo format, the implication hadn't been clear on a first read-through. But it was now. This wasn't a casual instruction. It was an order.

I flicked back a few frames and found a field report of an investigation into German guided weapons technology.

It was then that my eyes were drawn to an assessment of a sensor called *Windhund* ("wind-hound"). *Windhund* was a "sniffer" device that detected the presence of aircraft by measuring differences in the polarity of the surrounding air. It then automatically directed its parent aircraft to follow the trail until it reached the bomber stream itself.

All you had to do was hard-wire *Windhund* into an autopilot and you had a mechanism to allow a pilotless aircraft—conceivably, the Fireball—to be controlled automatically.

In the same document, a second sensor from a related facility (same organization, different location) was detailed as an infrared tracking system for locating aircraft exhaust gas. Both technologies were at an early stage, but the fact that they were being investigated at all was significant. A half century later, as far as I knew, neither had been perfected or developed. But here, in 1945, were sensors that a foo-fighter

could have used to get close enough to a bomber to disrupt its engine ignition systems.

I felt for the first time like I had a foot in the door.

A few frames later and here was a large document set out in tabular format with headings running across the top of the page: *Target~, Organization, City, Activity, Assessed (date), Action Taken and Remarks*. From the first page, it was clear that this was a very different kind of file, filled with data as raw as it came, and instantly noteworthy for what it did—and didn't—say. In the remarks section of one target, listed only as a "research station" near the town of Eib See, assessed primarily by the British, curiously, on May 2, 1945, it said: "Over 400 evacuated Peenemunde personnel held. Excavations made in the mountainside close to lake for underground workshops. A very important target." Nothing else.

Mysterious as hell.

Next page.

Target: Luftfahrt forschungs [sic]. *City:* Brunswick. *Activity:* Radio-controlled aircraft. *Assessed:* April 21-24, 1945. *Action Taken:* Team has been dispatched. *Remarks:* "Evidence of radio-controlled aircraft."

I scratched some notes. Another potential foo-fighter technology.

Next page.

Target: Research. *Location:* 87, Weimarerstr. Vienna. *Activity:* Experiments with antiaircraft rays. *Remarks:* "Research activity is conducted in a house at the above address. Research personnel were not allowed to leave house (reported hermetically sealed)."

The Germans *had* been working on directed energy weapons and had sealed those working on it from the outside world.

So many radical technologies and so long ago. I had no idea.

Next page.

Target: Daimler-Benz. *City:* Unter-Turkheim/Stuttgart. *Activity:* Secret weapon. *Assessed:* April 25, 1945. *Remarks:* "Said to stop ignition system of a petrol engine. The apparatus has succeeded in stopping a motor vehicle w/magneto ignition, but not one w/battery, at a range of 2 or 3 km."

The report went on to say that the technology had been insufficiently developed to have been brought against aircraft, but it made clear that this was the goal.

The Germans, then, *had also* been working on devices that were designed to "interfere with the engines of aircraft in flight," contrary to the note sent by Colonel Sullivan to General McDonald on September 28.

Why, then, did Sullivan say that all information available through in-

THIS PAGE IS UNCLASSIFIED

Target	Location	Source	Activity	Remarks
U	Oderwerke (Northhausen)	QSS London - 90 2205, 13 April 1945.	Manufacturing parts for the HS-177 Rocket (A. Werden).	HS-177 actually taken place underground in an old river shaft.
C	Woffleben (Halle-Kassel)	ditto	Manufacturing parts for the HS-177 Rocket (A. Werden).	
L	Grosser Inselsberg (Northern end of the Thuringen Hills)	A.U.I.K. Rep. 3-8/1945. QSS London, SI-2805. 13 April 1945.	Proximity fusion (electrostatic)	
A	Pfeidrich Listerwerke / Elm	Plauen - Volkgland	Urg. remote control mech. ("Fernsteuerung") for the prob. prototype of HS-177 Rocket (AA Werden).	Research activity is continued in a house at the above address.
I	Research	87, Teufelsstr. Thomas (23), 13 April 1945.	Experiments with anti-aircraft rays.	Research personnel were not allowed to leave house (ruined) harmlessly sealed.
F	Reichen?	U.W. Report by A.-A.I. (K) No. 220/1945.	"Fernauswurfsstelle" consisting partly of a hatched pole section for observing the flight of rockets launched from Pfernsdorf.	Students received instruc- tion on the beam system used in connection with A-4 Rocket.
E	Lubian	ditto	Instruction on the beam system used in connection with A-4 Rocket.	In this report confirms the A.U.I.K. Rep. No. 6124394 that one or 3 batteries of an Abteilung was equipped with a Leitstrahl transmitter.

terrogations, equipment and documents had been "thoroughly investigated" and that the subject "may be closed with negative result"?

What negative result?

Over the years, I had conducted extensive interviews on and off the record with intelligence analysts on both sides of the Atlantic about Soviet and Russian weapons developments and knew something of their techniques. They would never have dismissed an invention such as *Windhund* as cursorily as Sullivan had in the light of clear evidence that the enemy had been developing such a radical strand of technology. At the very least, the reports would have gone into a holding file. If no other reports came in to substantiate this finding, then in time it would have been downgraded. Only then, would field agents have been told to stop looking.

Instead, Sullivan's exhortation for investigators to cease forthwith their search for one of the key component technologies of the foo-fighter sightings, a mystery that had absorbed the USAAF's intelligence community in that winter of 1944—45, was, well, odd.

Unless, that is, they had found a foo-fighter and needed to put the lid back on.

It got better. I came to the field report on the last target to have been assessed, the Institut für Elektrophysik Hermann Goering at Landsberg-am-Lech in Bavaria. Like so many of the others it was quite anodyne until the remarks section, which read:

"Experimental work in conjunction with airfield at Pensing, 9 km north (toward Munich). Activities: research on gas, aircraft, pilotless aircraft, radio communication."

Here, then, was a facility that merged in a single program all the activities that explained Vesco's Fireball drone: a pilotless, remote-controlled aircraft that disrupted the engines and electronic systems of Allied bombers. And the fact that the research activity was taking place at an airfield seemed to suggest that the tests were at an advanced stage when the *Lusty* assessment team showed up.

I sat staring at the screen for several more minutes, scouring the text for other corroborative details, but there were no further clues and I was out of time.

I photocopied what I could, thanked the archivist and made my way outside. The taxi was already waiting, engine running, wipers working against the rain. I threw my bags on the backseat and sat back for the ride, my head buzzing with thoughts about a strand of weapons technology that appeared to have been buried from view for more than half a century.

But it wasn't until the flight home that sanity prevailed.

What had I actually learned from *Lusty*'?

That the Germans had been working on technologies that far exceeded published accounts of their aeronautical achievements in the Second World War?

Certainly.

That they had developed a craft that explained the foo-fighter sightings reported by Allied airmen?

Possibly.

That they had developed antigravity technology—and that this technology had fallen into the hands of the Americans at the end of the war in Europe?

Unfortunately, not.

Besides which, I realized that *Lusty* had been declassified in the mid-1960s, several years before Vesco's book *Intercept—But Don't Shoot* was published. A skeptic could claim that Vesco had simply concocted his account from the documents.

I had hoped that by applying a few basic investigative principles, I would find some tangible evidence of a radical propulsion technology that had been overlooked by everyone else.

Of course, it was never going to be that simple.

Despite all the stories, there was no concrete proof that the Germans had even designed, let alone built, a flying saucer. As for antigravity propulsion, no one had ever found a single piece of evidence that would reliably hold up in court.

A short postscript to the German story emerged; in Canada of all places. At the time it seemed irrelevant to the main body of my investigation, but it came screaming back at me later in a way I couldn't possibly have guessed at so soon after my return from Washington. I would recall it later as the "Silverbug affair."

In April 1959, during hearings of the Space Committee of the U.S. Congress, the U.S. Department of Defense revealed that it was working with the Avro Canada company to develop a "flying saucer." According to the Pentagon's Research and Engineering Deputy Director, John Macauley, the craft would "skim the earth's surface or fly at altitudes reached by conventional aircraft." This was the first public admission that Avro was building a flying disc, the Avrocar, the brainchild of a gifted British engineer called John Frost, who'd moved to Canada shortly after the war.

Activity surrounding the Avro project was shrouded in mystery. Word of its existence had leaked as early as 1953, when the *Toronto Star*

reported that the company was working on a "flying saucer" that could take off and land vertically and would fly at 1,500 mph. British and American scientists were also reported to be involved. Both the Canadian government and Avro played down the reports.

When the Avrocar finally emerged from the shadows it was easy to see why. One look at the prototype was enough to know that it would never achieve supersonic flight. Flight tests, in fact, confirmed it was a dog.

Underpowered and unstable, the Avrocar made its first untethered flight in December 1959 and soldiered on for two more years as Frost and his team sought to overcome its failings.

By the end of 1961, at the conclusion of the U.S. development contract, the Avrocar had managed to claw no more than a few feet into the air. Funding for the world's only known flying-saucer program was not renewed and the Avrocar drifted into obscurity—although not fast enough for the Canadian aerospace industry, which has sought ever since to distance itself from the project. There and elsewhere, it has become a byword for failure; something of an aviation joke.

Yet, that may have been the intention all along.

Recently declassified papers show that from early 1952 right up until the cancellation of the Avrocar in 1961, a select group of engineers known as the Avro "Special Projects Group," led by John Frost, had been working on a highly classified set of programs that accurately reflected the *Toronto Star's* original leak of 1953.

The papers revealed that, commencing with Project Y, also known as the "Manta," via Project 1794 to Project PV 704, the SPG tested technology for a whole family of flying saucers whose projected performance was designed to eclipse that of all other jet fighters. Data on Project 1794, a design for a perfectly circular fighter-interceptor, show that it would have been capable of Mach 4 at 100,000 feet. The best fighters then on the drawing board were hard-pressed to achieve Mach 2.

On December 2, 1954, Canada's Trade Minister C.D. Howe sought to dampen speculation about the existence of an Avro flying saucer program with a statement that the Canadian government had backed studies into disc-shaped aircraft in 1952-53, but had since dropped the idea.

This, as it turned out, was true—up to a point. In fact, the Avro design, now called Project Y2, had been purchased outright by the U.S. Air Force and re-code-named "Project Silverbug."

Under pressure from the U.S. press to reveal the nature of this program—which, though classified, had again leaked—the USAF issued a bland statement in 1955 that it had a "research and development

contract with the Avro Company of Canada to explore a new aircraft design concept." But that was all.

USAF documents on Silverbug declassified as recently as the mid-1990s show just how radical the Silverbug/Y2 concept was. "The Project Y2 design proposal incorporates a number of advanced improvements brought about by the utilization of several radical ideas in fundamental areas." Among these was a "very large radial-flow gas-turbine engine, which, when covered, will form a flying wing with a circular planform, similar in appearance to a very large discus."

The entire air vehicle, in other words, was a giant flattened jet engine, the hot gases that propelled it forward exiting from slots around its rim. By directing the exhaust, Silverbug would have been capable of almost instantaneous high-speed turns in any direction; maneuvers that would have included 180-degree reversals, the craft flipping this way and that like a tumbling coin, and all with minimum discomfort to the pilot, because the g-forces would have been alleviated by the adjustable thrust.

Much like the craft that had vexed the U.S. Air Force intelligence community following the first rash of U.S. flying saucer sightings in 1947.

The only other place prior to this discovery I had ever heard mention of a radial-flow gas turbine, an RFGT, was in Germany during the war.

After the ignominy of the Avrocar program, the Avro SPG disbanded and most of its exceptional brain talent was snapped up by U.S. aerospace companies or government-run research establishments. The most notable exception was John Frost, the maverick genius and inspiration behind the saucer projects, who soon after the Avro company was dissolved in April 1962 emigrated to Auckland, New Zealand. After the exhilaration and excitement of designing highly classified, highly unconventional air vehicles for the Canadian and U.S. air forces, Frost settled for the relative obscurity of an engineering job with Air New Zealand. He rarely, if ever, spoke about his work on the Avrocar and never mentioned the existence of the supersonic saucer projects that had gone on all the while behind it. He died of a heart attack in October 1979 at the age of 63.

Soon after I got back from Washington I rang Frost's son, Tony. I had tracked him down to Auckland, New Zealand, via a small band of researchers, all with strong aerospace credentials, who had done a fantastic job in piecing together the Avro story—the real story; one in which high-performance flying saucers had been shown to be real; far from the stuff of legend, as they had been in Germany.

"Dad was very secretive about his work," Tony Frost told me.

"Everything I ever learned about what he had been doing in Canada I had to pick up from people he'd worked with. The Special Projects Group was, for a time, just about the most secret group of its kind in Canada or the U.S., but why its activities should remain shrouded in secrecy for so long is still a mystery to me."

In the brief silence that followed, my mind worked hard to fill in the gaps. Had the lumbering Avrocar been an intentional blind all along, something to draw the flak from the SPG's real, but hidden goal: the creation of a supersonic, vertical takeoff and landing interceptor?

With Avro shut down, had work on the supersonic saucers continued in the utmost secrecy in the U.S.? Was there something about the very shape and form of the saucer, indeed, that was inherently super-classified?

While the Avro program didn't entirely explain how saucer-shaped craft had been "within the present U.S. knowledge" in 1947, it came very close.

First off, Avro showed that man-made flying saucers were real; they'd existed.

Secondly, if you substituted the radial-flow gas turbine, the RFGT, for something even more exotic—a T.T. Brown—type propulsion system, for example—the specification was a pretty neat match.

In the conventional world of aerospace, testing a new kind of airframe with a known engine was standard operating procedure. Was this what the Avro program had really been about? Were the Americans using the Canadians to perfect the aerodynamics of the saucer before the arrival of an all-new propulsion source—something that Trimble and Co. had hinted at in their statements of 1956?

Or was the truth altogether more prosaic? Had the supersonic Avro disc program merely been hushed up all these years in an attempt to avoid embarrassment by those who had thrown money at it and seen it fail?

These were questions without answers, so I asked Tony Frost instead how he thought his father had come by his radical ideas.

He told me, almost in passing, that a file recently uncovered in Canada's National Archives had shown that his father had made a journey to West Germany in 1953. There, at a Canadian government installation in the company of British and Canadian intelligence officials, Frost met with a German aviation engineer who claimed to have worked on a vehicle similar to the disc-shaped aircraft on the drawing boards at Avro. The German said that the project had been under way at a site near Prague, Czechoslovakia, in 1944—45, and that the saucer had not only

been built, but flight-tested. He told Frost that both the plans and the craft itself had been destroyed in the closing weeks of the war.

How the information was used by Frost was never made clear. Like the Avro supersonic disc program, he never spoke about it—to his family or anyone.

I felt the historical data pulling me back. If the Canadians and the British had been colluding on German-derived saucer technology, it was dangerously close to the scenario that Vesco had espoused all along.

But Tony Frost had spent years trying to piece together the secret history of his father's life and he was still no closer to the truth.

I had exhausted the safe, cozy world of archives and Internet trails. It was like Cross had said; they'd succeeded only in taking me everywhere and nowhere. If a decade of training had taught me anything, I knew my only real chance of verifying Trimble's outburst back in 1956 was to search for leads in the real world.

I didn't know it yet, but I was about to get a helping hand from a scientist on the inside loop, a man I came in time to regard as a genius, who for reasons of his own had already embarked on exactly the same journey as I had.

Chapter 7

I had expected the drive to Austin, the state capital, to provide me with a rich set of impressions of the frontier that divided the Eastern Seaboard from the wilderness that started here and extended to the Pacific. But for two hours I sat on the freeway, my views of Texas obscured by bumper-to-bumper traffic, peeling trailer homes and secondhand car lots. And so I thought of the man I was about to visit instead.

Dr. Harold E. Puthoff, "Hal" to those who knew him better than I, was the director of the Institute of Advanced Studies at Austin. Puthoff had lived his whole life on a frontier; one that straddled known science, the world explained and understood, and a place that most people, scientists included, said didn't exist.

I felt drawn to Puthoff, because his academic qualifications, not to mention his connections to the military, put him squarely in a place that I could relate to; yet, he was clearly also involved in strange science.

His résumé, which I had pulled off the Internet, said that he had graduated from Stanford University in 1967 and that his professional background had spanned more than three decades of research at General Electric, Sperry, the Stanford Research Institute (SRI) and the National Security Agency, the NSA, the government's super-clandestine electronic eavesdropping organization.

It also told how he "regularly served various corporations, government agencies, the Executive Branch and Congress as a consultant on leading-edge technologies and future technology trends" and that he had patents issued in the laser, communications and energy fields. Puthoff was no lightweight.

What his résumé didn't describe was one of the strangest episodes in the history of U.S. intelligence gathering: a program, known as remote viewing or RV, that enabled the Central Intelligence Agency and the Defense Intelligence Agency to spy on America's enemies using clairvoyance. For years, depending on whom you believed, this highly organized, well-funded operation had proven remarkably successful, providing reams of usable data on Soviet military activities that were

inaccessible to more traditional intelligence gatherers, such as spy satellites and aircraft.

RV remained buried, deeply classified, until 1995, when the CIA finally admitted it had used "psychic spies" against Russia during the Cold War.

Puthoff had been the RV program's founder and first director and had run it for 13 years, finally leaving SRI in 1985 to establish his own institute, the IAS, at Austin.

And now, he was up to his ears in gravity work.

If the U.S. government had instituted a top secret antigravity program, decades before possibly, then chances were Puthoff was either directly involved, knew about it, or, at the very least, had some inkling of it. On the other hand, doing what he'd done, acting as a consultant to some of the government bodies he still advised, I knew that Puthoff breathed the same air as people for whom disinformation was a just another part of the tool kit.

Ordinarily, I would have been wasting my time and Puthoff's. My limited, workaday knowledge of physics put me at an enormous disadvantage when it came to sorting out good information from bad in the antigravity field.

But on that day, as the miles to Austin sped by, I had some help.

Deep in the heart of Britain's own aerospace and defense establishment, I had found someone to decode the intricate mysteries of gravity for me.

I wound back the clock to the time when I'd attended a trade conference many years earlier. It had been entitled "Antigravity! The End of Aerodynamics." I'd only gone along because of the caliber of the speaker: Brian Young, Professor of Physics at Salford University. Young was also Director of Strategic Projects for British Aerospace (BAe) Defence, Britain's biggest military contractor.

The lecture had been given at the London headquarters of the Institution of Mechanical Engineers, IMechE, just off Parliament Square. Handouts beforehand (which I'd clipped to my copy of the transcript) had set the theme: "Gravity is the most mysterious of all the natural forces. It appears completely indifferent to anything we try to do to control it. Nonetheless, its very presence is the driver for the whole science and business of aeronautics. Scientists from the very distinguished to the very ridiculous have tried for over 300 years to either explain it or destroy it. Technology is making tremendous strides in all directions and perhaps we are getting close. If so, is this the end of aerodynamics?"

I chewed over the words of Young's speech, according it a respect I hadn't given it on the night it was given. Now I wished I had.

I recalled the elegant marbled hallway and the low hubbub of conversation as I'd drifted among the attendees—a hodgepodge of MoD men and aerospace industry executives, some of whom were familiar to me—waiting for the doors to open into the auditorium.

Young had kicked off his talk by reiterating our general state of ignorance about gravity. "It is an incredibly weak force," he'd said, "although it probably didn't seem so to the Wright Brothers and I suspect most mountain climbers would disagree violently."

"But really gigantic quantities of matter, about 6×10^{21} (to the power of 21) tonnes in the case of the Earth, are required to produce the gravity field in which we live. If you think of a simple horseshoe magnet lifting a piece of iron, then, as far as that piece of iron is concerned, the magnet weighing a few ounces is outpulling the whole Earth."

He flashed up a set of artist's impressions showing how three different kinds of antigravity vehicles might look assuming engines could be built for them.

One, a "heavy-lifter," looking much like an airship, was depicted effortlessly transporting a giant section of roadbridge through the air. Young made the assumption that the antigravity engine of the heavy-lifter not only canceled the weight of the craft, but its underslung load as well.

Another, saucer-shaped, craft, described as being like a bus for "city hopper" applications, used a "toroidal" or doughnut-shaped engine to allow it to skip between "stops" in a futuristic-looking metropolis.

The third showed a combat aircraft with a green-glowing antigravity lift engine on its underside for vertical takeoff and landing (VTOL) and conventional jet engines for forward propulsion.

But all these things were fiction, he stressed, so the designs were simply conjecture. "The more I have read and thought about antigravity and its terrestrial applications, the more I have become convinced that as it stands today it is going nowhere. Whatever name the press may give experiments, either in someone's garage or in Geneva, most of the real work is trying to uncover basic knowledge and few people really believe it will lead to gravity control."

"To the contrary, a view I have seen expressed is that if gravity control is ever discovered it will be as an unexpected by-product of work in some completely different field."

To which he'd added a rider. Were he to have gotten it all wrong, were

an antigravity effect ever to be discovered, the growth of the industry surrounding it would be exponential—look what had happened after Faraday's discovery of electromagnetic induction in 1831. Within five years, the electric motor industry was born. Within seventy, the generation of electricity was huge business. Today, it is hard to conceive of a time when man-made electricity did not exist.

I vaguely remembered going up to Young after the lecture and chatting to him about some of the points he'd raised. But now I kicked myself. If only I'd paid more bloody attention.

Soon after I'd returned from the low-key archive in Washington that housed the *Lusty* documents, I'd picked up the phone and dialed British Aerospace. Through to the public relations department and I logged the request. Could they get me Professor Young's phone number?

Forty minutes later, the media manager called back to say that Professor Young had died, of a heart attack he thought, some years earlier.

I put the phone down, a little shocked at the news, and pinched the bridge of my nose, looking for that extra bit of concentration. What had Young and I talked about? Down into the basement and another protracted rifle through a box taped up and marked "notebooks: 1989—94." There in among my old reporter's pads, after an hour on my hands and knees, I found the one that I'd taken with me that night to IMechE. I started reading and flicking pages.

The notes I'd jotted down were piss poor, highlighting terms like "gravito magnetic permeability" and "toroidal engine" and putting big question marks against them.

I kept turning. There, right at the end, I'd written myself a memo to put in a call to a media affairs officer at BAe who would arrange for me to receive photos of the artist's impressions Young had flashed up during his speech.

This, I remembered now, was one of the things Young and I had discussed.

Below the memo were two names: Dr. Ron Evans of "BAe Defense Military Aircraft's Exploratory Studies group" and Dr. Dan Marckus, an eminent scientist attached to the physics department of one of Britain's best-known universities. The department was more than familiar to me. Its work was well known in defense circles for the advice that it gave to the government.

Beside Evans' and Marckus' names were their contact phone numbers. I called Dr. Evans first.

He turned out to be a nice, soft-spoken mathematician based at BAe's Warton plant, a facility that cranked out jet fighters close to where the bleak mudflats of Lancashire's Ribble Estuary met the Irish Sea.

When I asked if he could brief me, as per the suggestion of Professor Young, Evans said he needed to consult with the company's media managers. He'd never dealt with the press before. I got the impression it was sensitive stuff.

It was, but not for any of the reasons I had guessed or hoped. Britain's biggest aircraft manufacturer, BAe, had an antigravity department. That in itself was key.

But its activities weren't classified. Far from it. The impression I got, more than anything else, was that the management had tucked Evans' tiny department into the farthest-flung corner of the company because it didn't know what else to do with it. That and the fact no one in BAe at that level wanted to wake up to headlines that it was involved in kooky, Buck Rogers science. It would destabilize the share price.

In the end, Evans wasn't allowed formally to brief me on his work, but he was given permission to fill me in on some of the background that had led to the formation of his Exploratory Studies group. That suited me fine. BAe didn't want to break cover—nor did I. For once, we were on the same side of the secrecy divide.

In March 1990, Evans chaired a two-day University-Industry Conference of Gravitational Research, sitting around a table with a gathering of distinguished academics to identify any emerging "quantum leaps" that might impact on BAe's military aircraft work. Gravity control figured extensively on the agenda.

Imagine it. A technology, popping out of nowhere, that rendered all of its current multibillion-pound work on airliners and jet fighters redundant at a stroke. Now that, I thought, *really* would do things to the share price.

The company also undertook some practical laboratory work in a bid to investigate the properties of a so-called "inertial-thrust machine" developed by a Scottish inventor called Sandy Kidd.

In 1984, after three years' work building his device—essentially, a pair of gyroscopes at each end of a flexible crossarm—Kidd apparently turned it on and watched, startled, as it proceeded to levitate, then settle three inches above the surface of his workbench.

An ex-RAF radar technician, the Scotsman had become obsessed with the idea of inventing an antigravity machine after he'd removed a still-spinning gyroscope from a Vulcan nuclear bomber. Carrying it backward

down the aircraft's ladder, he reached the ground and was promptly flattened by a sledgehammer blow to his back. The force, he realized, had been transmitted from the gyroscope, which had reacted in some inexplicable way as his feet had touched the ground.

In May 1990, BAe began a series of trials to test whether there was anything in Kidd's claims, knowing full well that he wasn't alone in making them.

In the mid-1970s, Eric Laithwaite, Emeritus Professor of Heavy Electrical Engineering at Imperial College London, demonstrated the apparent weight loss of a pair of heavy gyroscopes by lifting the whole whirling contraption with one hand and wheeling it freely around his head. When the twin rotors were not in motion, attempting this feat was impossible. The gyroscopes were the size and weight of dumbbells. But when the rotors turned, they became as light as a feather.

Somewhere along the line, the machine had lost weight.

The accepted laws of physics said that this was not possible, out of the question—heresy, in fact. But Laithwaite's claims were supported by a top-level study into gyroscopes published by NATO's Advisory Group for Aerospace Research and Development (AGARD) in March 1990.

The authors of the AGARD report concluded that a "force-generating device" such as Laithwaite's, if integrated into a vehicle of some kind, could, in theory, counteract gravity. "Clearly if such a counteracting force was of sufficient magnitude it would propel the vehicle continuously in a straight line in opposition to said field of force and would constitute an antigravity device."

The report went on to say that there was at least one "gyroscopic propulsive device" that was known to work and that the inventor, E.J.C. Rickman, had taken out a British patent on it. The trouble was, the report concluded, the impulses generated by these machines were so slight they would be useless for all practical applications—except, perhaps, to inch a satellite into a new orbit once it had already been placed in space by a rocket.

It was hardly a quantum technological leap. But that wasn't the point, Dr. Evans told me. What was being talked about here was an apparent contravention of the laws of physics; the negation, at a stroke, of Newton's Third Law, of action-reaction. Which was why the BAe-sponsored tests on the Kidd machine had a relevance that went way beyond their immediate and apparent value.

If there were ways of generating internal, unidirectional, reactionless forces in a spacecraft, and in time they could be refined, honed and developed, the propulsion possibilities would be limitless.

In May 1990, a trial went ahead under the watchful eyes of two wind tunnel engineers from BAe's Warton aircraft plant. The first series of tests showed no antigravity effects, but to everyone's surprise, during a second series of trials, one result did appear to indicate that the machine had changed weight.

Unfortunately, the engineers failed to repeat the result; and in science and engineering, repeatability is the benchmark of success.

Had it been repeatable, the two percent weight change Kidd claimed for his device would have been scientifically verified, and BAe would have employed the world's first known antigravity device as an orbital maneuvering motor for satellites.

Shortly after the trial, Cold War defense budgets collapsed and BAe's share price went into free fall. As it fought for its survival, there were cutbacks across the company and Dr. Evans' department, with little hope of any short-term returns, was disbanded as quietly as it had come together.

That, essentially, was how things were when he and I met.

I rang Dr. Marckus a few times, but only ever got an answering machine and so I capped my involvement in the BAe antigravity story with a few low-key articles in *Jane's* about the BAe investigation. It was more an excuse than anything else to broadcast a question: Was anyone else out there engaged in similar work?

The U.S. Air Force had recently proclaimed its interest in the field with a document, published in August 1990, called the *Electric Propulsion Study*. Its objective was to "outline physical methods to test theories of inductive coupling between electromagnetic and gravitational forces to determine the feasibility of such methods as they apply to space propulsion." Stripped of the gobbledegook, it was really asking whether there was any theory out there that might permit the engineering of an antigravity device.

The existence of the *Electric Propulsion Study* gave me an excuse to slip tailored messages into a couple more stories about futuristic propulsion. It felt a little like pointing a transmitter into deep space and waiting for a response.

I waited, but nothing came back. No emails, no anonymous faxes, no spooky phone calls. Nothing.

Thereafter, I kept a weather eye on the antigravity scene, but the receiver I had rigged for the slightest trace of life out there remained silent. In the end, I switched it off.

During the mid-1990s, BAe kept its toes in the gravity field by placing a few small-scale contracts with a number of British universities,

principally to see if there might be a provable relationship between gravity and heat; a link first postulated by Michael Faraday in 1849.

Like me, the company was clearly reaching, but getting nowhere.

Then, in 1996, NASA—the U.S. National Aeronautics and Space Administration—began questioning some of the perceived limits to the advancement of air and space travel in two notable areas. One related to the unbreachability of the light-speed barrier—a throwback to Einstein, whose Theory of Relativity said that the velocity of light was a fixed constant never to be exceeded.

The other questioned the heretical notion that, somewhere in the universe, there might be an antigravity force; one that could be exploited for air and space travel.

The resultant NASA program, wrapped within something called ASTP—the Advanced Space Transportation Program—prompted BAe to resurrect its antigravity work. It has since been rebranded "Project Greenglow," a "speculative research program . . . the beginning of an adventure which other enthusiastic scientists might like to join, particularly those who believe that the gravitational field is not restricted to passivity."

Decoded, Greenglow involved little more than reseeding small amounts of company R&D money to more British universities, funding a few more experiments of the Kidd kind, and acting as a European feed-in to the NASA effort.

It was, to say the least, depressing news. I thought back to George S.Trimble, the man who had been too afraid to talk to me after Dani Abelman of Lockheed Martin had approached him on my behalf. If Trimble had cracked the gravity code back in 1956, why were BAe and NASA, with their immense know-how and resources, still struggling with the gravity conundrum today? It didn't make any sense.

And that should have been where the matter ended.

But long months after I had consigned the antigravity story to the graveyard, I got a phone call. The man at the other end of the line was bluff and unapologetic, verging on the rude. Dr. Marckus had finally gotten in touch.

Chapter 8

The sunlight reflected brightly off the thin carpet of snow that had fallen along the east coast overnight. I kept the estuary on my right and the sea defenses on my left, edging the car along the narrow road, navigating by the instructions I had been given over the phone. There was something almost primeval about the combination of the pine forest, the limbs of the trees twisted into impossible shapes by the wind, the open sky and the stillness.

The trees parted briefly and I saw the radar mast that I'd been told to look for. Around another bend and the tiny haven swung into view.

I pulled up in the parking lot beside a wooden jetty that jutted into the green, fast-flowing current toward a small settlement on the far bank. The buildings seemed run-down and dirty compared to the crisp scenery through which I'd just driven.

There were two other cars parked beside mine and both were empty. I got out, stretched my legs and shivered as the still-freezing temperature blew away the fug of the hour-and-a-half-long journey from London. Suffolk. The East Anglian coast and the cold waters of the North Sea. I was in a part of the country I didn't know.

A light breeze tugged at the rigging of a row of yachts dragged onto a thin stretch of stony beach nearby. Seabirds picked along the jetsam and detritus of the shoreline. Above the gentle pinging of the mast ropes, I heard children's voices.

There were four people on the jetty.

A woman sat on the far side, wrapped warmly against the cold, reading a book as her two small children line-fished for crabs.

The father, standing farther off, his back to me, was tracking an open motorboat chugging hard against the fast-flowing water, its helmsman preparing to tie up alongside the jetty's end.

As the boat bumped against the tire buffers, the man stepped on board and settled onto one of the bench seats.

I waited for the woman and the two children to get up and join him, but they never so much as glanced his way.

It was then that I realized the man in the boat was Marckus.

He looked up and waved me aboard. I hesitated. Taking a ride on a boat had not figured in our brief phone conversation.

Until then, I'd not considered the possibility that Marckus was anything other than the person he claimed to be.

But what did I know about him? That he was plugged firmly into Britain's aerospace and defense community, that he didn't make a habit of talking to journalists, and that he was taking a risk by talking to me.

The ferry was preparing to leave.

I ran along the jetty and dropped down into the boat. Marckus paid for both of us and the ferryman cast off.

I found myself apologizing for the mix-up; how I'd thought Marckus was with the woman and her children on the jetty. Silly, nervous jabbering.

Dr. Marckus cut me short. "Does the significance of this site mean anything to you?" he asked, his arm almost hitting my face as he waved in the direction of the road.

I shrugged and he gave a mild snort of derision. "I thought you said you were a Jane's man."

"I'm sorry," I said, "it looks like a small yacht club and a fishing village." I neglected to tell him, because I couldn't see the point, that history wasn't my bag. My work was high technology of the 21st-century kind.

Once, Dan Marckus must have been lean and ascetic-looking; the thinning hair and high forehead still gave a hint of it. With age, however, he had put on some padding, making him look a lot less severe than he sounded. A dark, scratchy beard, flecked with gray, distracted the eye on first meeting from a thick pair of lenses in tortoiseshell frames.

Behind them, Marckus' brown, beady eyes watched me intently and with a hint of irritation.

He was dressed sharply in a thick blue polo-neck, a trendier version of the kind fishermen wear, and a brown leather jacket. His voice said he was around 65, but he looked younger.

"What you saw when you drove through those trees was one of the orginal antenna masts of Bawdsey Manor. Before the Second World War broke out, the Germans sent an airship, the *Graf Zeppelin*, up and down this coastline trying to figure out what we were up to—what those tall steel lattice towers meant."

"And what did they learn?" I asked.

"They'd packed the *Graf Zeppelin* with electronics—listening gear. But the old man was so much smarter than them, thank God."

Marckus registered my bewilderment. "Robert Watson-Watt. The man who gave the British the secret of radar. It all happened here. I thought you'd know that."

Once on the other side, we jumped down onto the beach and strolled along the shoreline, past a lonely cafe, in the direction of the point.

"So how did Watson-Watt outwit the Germans?" I asked, to fill the silence more than anything else.

"They saw the *Graf Zeppelin* on their scopes when it was way out over the North Sea, realized what it was up to and switched off the radars. The Germans went away believing that they were radio masts, that Germany was the only nation that had successfully developed radar detection. A month later, they invaded Poland."

"That's interesting. Did you work for him?" The term he'd used, "old man," denoted more than a passing familiarity. "Watson-Watt, I mean."

Marckus never answered. He walked on in silence, watching the gulls soaring on the cold currents above our heads. Then, without looking at me, he said: "It's been a while since you've written about my favorite subject. I was beginning to worry that you'd lost interest in it."

"It's like Professor Young said in his speech. I have come to the conclusion—a little late in the day, perhaps—that antigravity is for the birds."

"I see," he said. "Are you hungry?"

I looked at my watch. It was close to midday and I hadn't had any breakfast.

Marckus, registering my hesitation, doubled back toward the cafe.

I pulled a copy of the U.S. Air Force's *Electric Propulsion Study* from my coat pocket and set it down on the table next to a tatty yellowing manila folder that had been clamped under his arm throughout.

Marckus and I were the only two people this side of the deep fryer. After serving us our plates of cod and chips, the cafe's proprietor left us on our own.

Marckus seemed to start to relax.

"Apart from the fact that the math and the physics are beyond me, the evidence says that this whole science, if you can call it that, is still in its infancy," I said. "One look at this report says they're still playing around with algebra. When I wrote what I wrote, I thought the story was bigger than that. I have to concede I was wrong."

"What were you hoping for, if I may ask?" Marckus asked.

I shrugged. "I thought maybe they'd have at least built some hardware by now. God knows they've had time. The U.S. aerospace industry first

started talking about this stuff in the mid-1950s. You'd think by now there might just be something to show for it. But the Air Force is just staring at its navel and doing a few sums."

I tapped the *Electric Propulsion* report. "Modifications to Maxwell's equations and all that."

"As I recall," began Marckus, "in the 1970s, the Americans tripped over some research published by a Russian mathematician, Ufimtsev, who in turn had gained his inspiration from work done by Maxwell a hundred years earlier. The end result was stealth, the biggest breakthrough in military aircraft design since the jet engine. Not bad, for a mathematician."

"The difference is, they developed stealth," I told him.

"How do you know they haven't developed antigravity?"

I studied his face. How much did Marckus really know?

He must have read my expression. "My information isn't any better than yours," he said, raising a hand in the air to signal his innocence. "The U.S. and the U.K. share a lot of sensitive data, as you know. But there's nothing come across my desk that says we're cooperating with the Americans on antigravity—or even that the Americans are doing it on their own. My point is, we don't know." He brought his hand down on the manila folder to reinforce the point.

I looked at the U.S. Air Force study again. It was hardly reassuring. If antigravity existed and people had known about it years ago, how come the Air Force—along with BAe and NASA—was only just starting to look for it in the mid-1990s?

Marckus stared into his cup. "Did they build the first atomic bomb with a few 'poxy' million bucks? Christ, no. It cost them billions of dollars, even in 1940s values. Secret dollars nobody outside the secret was supposed to even guess at."

"So, has antigravity been developed in the black or not?"

Marckus glanced up. His voice hardened. "I already told you. I don't know. None of us this side of the bloody Atlantic knows." Then he coughed, semi-apologetically. "But there are signs."

"What signs?" Despite myself, I leaned forward.

"Up until 1939, atomic energy was still a matter of conjecture, even among physicists in the field. But by the following year, some of the brighter sparks in the community realized that it was doable, that if you split a neutron into equal parts, you'd quite likely get an enormous release of energy. Things that appear impossible usually aren't, even when the physics say they are."

I knew enough to know that in the 1930s next to nothing was

understood about nuclear physics, that by the outbreak of the Second World War all experiments resulting in nuclear reactions had required more energy than had been released.

As more and more physicists escaped the Nazis, however, and debate in America was stimulated further, the conjecture spilled into the U.S. press and writers began to speculate about what they had heard. A number of sensational articles appeared about the weapons implications of a nuclear reaction that could be compressed into the blink of an eye. The word was, it would result in a massive bomb.

By 1940, when it was understood that uranium bombarded with neutrons fissioned and produced more neutrons and that a multiplying chain reaction really might occur with huge explosive force, something remarkable happened. The nuclear physics community voluntarily stopped the publication of further articles on fission and related subjects and this, in turn, dampened the media's interest.

By the time America entered the war in December 1941, you couldn't find a mention of fission. It was as if no one had ever been discussing it.

The parallel with what I had read in the U.S. media in the mid-1950s, culminating in all those statements in 1956 that antigravity was doable with a Manhattan-style effort behind it, and then the crushing silence, was extraordinary.

By 1960, you talked about antigravity and people looked at you like you were mad.

I stared out the window, trying to find Watson-Watt's once-secret radar mast, but it was lost behind the trees. I knew Marckus lived somewhere nearby. I wondered where, exactly; what his house was like, whether he was married, if he had kids.

"What's in this for you?" I asked him.

"You have freedom of movement, I don't," he said. "You can go visit these people and ask questions, I can't."

I still couldn't quite see where all this was headed.

Marckus took a moment before adding: "I think it's 1939 all over again. I think we're poised on the brink of something. The physics is mind-boggling—Christ, I don't pretend to understand half of it myself—but what we're talking is huge."

"Spell it out for me."

"Bye-bye nuclear power. Bye-bye rocket motors. Bye-bye jet engines. If we can manipulate gravity, nothing will ever be the same. But expecting British Aerospace to develop this stuff is like asking someone who'd spent the first half of the 17th century building horse-drawn carts to come up with the outline for Stephenson's bloody Rocket. People

don't really have a taste for it here, not at a senior level. They don't have the vision that the Americans do. Antigravity is too much mumbo-jumbo, not enough ROI." He rubbed his thumb and forefinger under his nose and pulled a face. "Return on investment—bean-counter talk."

"So, if you can prove that the Americans really are up to something," I said, "you raise the stakes and get something going here? You get a real program in this country with real money behind it?"

"Right—to a point," he said. "The difference is you're going to prove it."

I stared at him, thinking I hadn't heard him right.

"It's not such a bad arrangement," he said, reaching into the manila folder and pulling a bunch of papers from it. "You get a short course in physics, a few tips on who to talk to, and maybe a story out of it." He paused. "The story of the biggest breakthrough in transportation technology since the invention of the wheel."

"You said there were signs . . ." I said cautiously.

Marckus pushed the papers across the table. The cover sheet was a photocopy from some kind of index. The writing was small, but my eyes were drawn to a section of text that had been highlighted with a yellow marker pen.

I peered closer and words like "magnetic monopoles," "gravitational charge" and "linear equations" swam into my vision.

"This is taken from something called the 'Source Index,'" he said. "It's a list of doctorate papers published by universities around the world. The highlighted part shows you what is happening in one U.S. university, that's all. There are dozens, maybe hundreds, of other facilities like these across America engaged in related research. All of them probing the sorts of areas you'd want to probe if you were pulling a field-dependent propulsion program together. Some of the entries go back decades."

Field-Dependent Propulsion; yet another term for antigravity.

"This thing is a multifaceted, multilayered puzzle and you won't solve it by picking away at the edges," Marckus said. "Begin with what you can see, then search for what you can't. What NASA's doing is in the open, so get over there and start looking and listening. Then go deeper. Some people say that the B-2 Stealth Bomber uses an antigravity drive system. It's like the bomb. The physics has been around for a long time. All it requires is some imagination, a shitload of money and some technical know-how and someone who wants to put the pieces together."

On the way back home, as the shadows lengthened and the traffic bunched on the choke points into London, I thought long and hard

about the road ahead and where it was taking me. I knew absolutely nothing more about Marckus than I had gleaned from him before the meeting. The meeting had confirmed that he was a highly gifted academic with strong ties to the British government and its defense industry and that he was unquestionably up to his eyes in secret work. But that was all.

It didn't make me feel particularly comfortable. But then again, I'd already painted myself into a corner and this, at least, was a way forward. Doing nothing was not an option if I wanted to progress things.

Marckus had given me the names of two people I needed to see in the States. One was Garry Lyles at NASA; the other was Hal Puthoff.

In American defense procurement, there was a white world and a black world—one that was open to public scrutiny and one that was so firmly closed, from an outsider's perspective it was like it didn't exist.

The white world had know-how, a good sprinkling of imagination and not a little money at its disposal.

The black world, on the other hand, had all of the above in spades.

Lyles clearly worked in the white world. But with Puthoff, it was absolutely anyone's guess.

Chapter 9

A larger-than-life depiction of Wernher von Braun, whose self-assured good looks conjured up images of postwar Hollywood matinee idols, stared down at me from a mural in the gallery above the arrivals hall at Huntsville International Airport.

Von Braun is still revered in that remote corner of Alabama as the founding father of the nearby George C. Marshall Space Flight Center, the NASA "center of excellence" for space propulsion. Von Braun had died more than 20 years earlier, but American rocketry remained as a testament to the work of Operation Paperclip, the U.S.A.'s clandestine effort after World War Two to recruit Nazi V-2 specialists.

A complex of drab, rectangular buildings, "the Marshall Center Wernher von Braun administrative complex," erected at that time, stands as testimony to the decade in which the high ambient humidity routinely reverberated to test runs of engines for the Mercury-Redstone and Saturn V launch vehicles. The former took the first U.S. astronaut, Alan B. Shepard, briefly into space three weeks after Yuri Gagarin in May 1961; the latter formed the basis of the giant Apollo moon rocket.

Today, NASA Marshall scientists still work on the same basic technology that von Braun developed under the auspices of the Third Reich, even though chemical rockets, his great invention, are today regarded as too slow, too unreliable, too expensive and too inefficient.

It was mid-summer, four months after my meeting with Marckus, before I was able to wangle a trip to Marshall and do as he'd suggested.

Set in a vast area of grasslands, bordered by well-tended, lush forestation, Marshall was melting under the high heat when I eased my rental car up the long approach road that peeled off the interstate to the hub of the Center itself. As I got out of the car, a military Huey thudded low overhead, disappearing behind the trees in the direction of the Redstone Arsenal, the Army's neighboring missile test facility. As the low-frequency throb of the blades faded from earshot, it was replaced by the gentle swish of grass sprinklers and the chatter of cicadas.

I was met at the entrance of the main building of the complex by a

woman from public affairs. As we moved through the lobby, she sketched in some of the background to the program I'd come to talk about with its manager, Garry Lyles.

Under the Advanced Space Transportation Program, ASTP, officials at NASA Marshall have been working since the mid-1990s on methods to reduce the cost of space access. Using today's generation of launch vehicles—the Space Shuttle Orbiter, the Delta and Atlas rockets—it costs around \$10,000 per pound to put a satellite into low Earth orbit. One short-term payoff of ASTP will be the development of a family of reusable rocket planes that should cut these costs by a factor of ten by the end of the present decade. A mid-term goal, due to yield results in around 2015, is to develop a hybrid "combined cycle" engine—half-rocket, half-jet engine—that will reduce launch costs by a factor of a hundred. Even though the hybrid engine, if it can be mastered technologically, will give NASA space access for as little as \$100 per pound, it will never lead to anything other than a delivery system to low Earth orbit.

For missions beyond Earth, scientists will either have to rely on improvements to the chemical rocket or on something radically different.

For a mission to the planets of the inner solar system a nuclear rocket could be developed in a relatively compressed timescale to cut round-trip journey times from Earth to Mars from many months to possibly just weeks. The feasibility of nuclear rockets had been proven in two NASA-developed solid core fission reactor test programs, NERVA and ROVER, between 1959 and 1972. These never came to fruition, my PR guide told me, due to concerns over safety and also because emerging and more advanced nuclear fusion technology was seen as yielding three times the efficiencies. Like the fission rockets, however, fusion systems never happened either—not in the white world, anyhow. In the early 1990s, the Pentagon was reported to be working on a classified nuclear thermal rocket under the code name "Timberwind," but soon afterward it was canceled. Since the black world is publicly unaccountable, it is impossible to know whether this really was the case.

Lightweight nuclear fusion propulsion is one methodology being studied for a manned NASA mission to Mars, my guide continued as we drifted toward the elevators, alongside "light-sails," "magnetic sails," "antimatter fusion drives," lasers and other exotic-sounding technologies.

While the physics behind these ideas is generally well understood, she added, the challenge in turning them into hardware is enormous.

But ASTP did not end there. In addition to examining ways to provide cheaper access to low Earth orbit and coming up with propulsion systems that by the second half of the century will lead to "the commercialization of the inner solar system," its original task went a lot further.

Into an elevator, half a dozen floors up, and we emerged into a world of bright walls, carpeted floors, posters of exotic spacecraft and artists' impressions of planetary systems, constellations and nebulae.

Up ahead, through an open door, sitting behind a large desk, Garry Lyles—bespectacled, bearded and thoughtful-looking—was waiting.

"Within a hundred years," he told me over coffee, his soft southern drawl catching with emotion, "we'll be going to the next star system."

He spoke about the journey with such absolute conviction that it was difficult, in spite of the sheer fantasy at the heart of the idea, not to believe him. Lyles was a dreamer, but his dreams were rooted in work that was going on within the Wernher von Braun administrative complex as we spoke. ASTP scientists at Huntsville—people working right on the edge of what was known and understood in the harsh world of propulsion physics—were collaborating with other NASA facilities on science that would enable astronauts to make a "fast trip" to the Alpha Centauri star system, Earth's nearest stellar neighbor, 4.3 light-years away. As head of this portion of ASTP, Lyles was bound up in the immensity of the challenge.

I asked what he meant by a "fast trip."

"We have set a goal to go to Alpha Centauri in 50 years—a 50-year trip." He stared at me intently as he rocked back in his chair. "You have to really use your imagination to come up with a propulsion system that will do that."

"When will you be able to make that trip?" I asked.

He gestured to another part of the complex visible beyond the blinds of his window. "We have a group of thinkers out there that's beyond anti-matter. They're looking at space warp, ways to get to other stars in a Star Trek-like fashion. If we're not thinking about that kind of thing today, we won't be going to the stars in a hundred years' time. But I have to tell you that if you talk to these people, these engineers who are working on interstellar missions, they think they can do it. And I have no doubt that they're right."

The trip, he reiterated, would be possible before the end of the 21st century.

The phone rang and Lyles took it, allowing me a moment to consolidate my thoughts. His team, he said, had opted for a 50-year trip because it represented a time span most engineers could get their heads

around. Fifty years, however, made no sense from the point of view of a practical mission, because even if it could be accomplished technologically, old age (or boredom) would have killed most of the crew long before they ever got there, let alone back again.

But that wasn't the point. The "extra-Solar System" portion of ASTP was about getting scientists to think about realizing impossible journeys. In 1960, there were those who said that landing a man on the moon by the end of the decade would be impossible. That we weren't colonizing Mars by the end of the 20th century, they maintained, was directly attributable to the drop-off in research activity that followed the Apollo program. If we wanted to get to the stars by the end of the 21st, Lyles said, we needed to be thinking about it now.

Which, though he didn't know it, was why I was here.

In 1956, a group of engineers had talked openly about an impossible idea. Lyles' "fast trip" to Alpha Centauri was so impossible, his team was having to really reach—to think way out of the box—to come up even with half-answers to the challenge. To make the fast trip happen, NASA had no choice but to tap heavily into some taboo areas of science. That was my lead.

Supposing for a moment that NASA could come up with a vehicle capable of light-speed, a one-way trip to Alpha Centauri would take 4.3 light-years. Since the speed of light was a constant 186,000 miles per second, by extrapolation, the spaceship would be moving at almost 670 million miles per hour.

The 50-year journey time reduced the requirements to a more manageable 16,000 miles per second or 57.6 million miles per hour.

I had only once come across real hardware (aside from a laser, the ultimate "speed of light" weapon) about which comparable speeds had been mentioned.

Shiva Star, the 250-ton capacitor I'd seen at Kirtland Air Force Base in New Mexico in 1992, had had a task to fire a plasma bullet at 10,000 km per second.

And then, just a few years later, when I returned to Kirtland, it was like the program never existed.

In the carefully regulated temperature of Lyles' office, I shivered.

Shiva's plasma bullet, designed to be dumped into the soft electronics of an incoming nuclear warhead at one-thirtieth the speed of light, was but one of a handful of deep black programs I'd encountered in the course of my career.

How many more secrets existed below the waterline of accountability? What other secrets were out there? What was the ultimate weapon? The

questions churned, as they always did in a place like this, almost too fast for me to get a hold on them.

What Lyles and his counterparts in the black world shared was their absolute belief that they could engineer things other people said were impossible. It was a part of the feeling I got when I visited facilities like this and it thrilled and scared me with equal measure.

Why had Marckus sent me here? What did Lyles know?

When he got off the phone, I asked Lyles if he ever felt NASA was developing technology that had previously been developed by the Air Force in the black. He took a moment before answering, choosing his words carefully. I thought that by tweaking his professional pride I might be able to ruffle some of that soft, southern *savoir-faire*. But Lyles had seen me coming. "Yeah, sometimes we do. But you have to understand that going into space is a very hard thing to accomplish. It takes a lot more than money. I would guess—certainly when it comes to single-stage-to-orbit technology—that NASA is at the cutting edge."

This was the essence of the multilayered conundrum that Marckus had touched upon during our meeting on the estuary. While NASA was struggling to develop technology that would reduce the cost of space access, there were those that said that the Air Force had already solved the problem. Believers in Aurora, the large triangular aircraft supposedly sighted over the oil rig in the North Sea, maintained that at least one variant was capable of conducting single-stage missions to low Earth orbit. If, as George S. Trimble and his contemporaries had stated back in 1956, the U.S. was on the threshold of developing antigravity technology, why spend billions on something as conventional (by comparison) as Aurora 20 years later? Or any other advanced kind of weapon system for that matter?

But when Lyles said that he doubted the Air Force had cracked single-stage-to-orbit, I actually had no way of knowing what he really knew. There was every chance that NASA was simply repeating work the Air Force had already carried out—and this applied to all aspects of its work "on the cutting edge."

I knew of several Congressmen who were deeply pissed off about it. On their watch, they'd committed money to a number of NASA programs knowing damn well, but unable to prove it, the Air Force had duplicate technology up its sleeve.

A number of NASA scientists I had spoken to in other fields of endeavor had acknowledged a feeling—though it was often no more than this—that their work was a mirror image of some hidden effort, something they couldn't see, that was going on in the black. If NASA

scientists, therefore, were thinking "way out of the box" to come up with science that would enable them to reach another star system by the end of the century, maybe someone, somewhere had experienced that same feeling.

If anyone at Marshall had sensed such a presence, it would have been George Schmidt. Lyles recommended I go talk to Schmidt, because Schmidt, being a project engineer, was right on the cutting edge. A big, genial man with a football player's physique and a marine-style haircut, Schmidt, as head of propulsion, research and technology on the ASTP effort, waded routinely through the theory that could one day transform Lyles' interstellar vision into real hardware. Still under escort, I found him down a maze of corridors in a different part of the von Braun administrative complex.

Schmidt punctuated his remarks with references to "superluminal velocities"—faster-than-light speeds—and "space-time distortion," phenomena that weren't even accepted by mainstream physics ("although General Relativity and Quantum Mechanics don't discount the possibility"). By the same token, he said, there were some holes in current thinking—"some windows of opportunity we may be able to take advantage of—that might transition the theory sooner rather than later. One of these related to so-called "wormholes," shortcuts through the universe that could be created with large amounts of "negative-energy density matter": material that existed only on the very fringes of theory.

Since it was known from the established physics of General Relativity that gravity, electromagnetism and space-time were interrelated phenomena, he said, it followed that any distortion of space-time might well yield an antigravity effect.

Questioned on the principles of a wormhole, Schmidt quickly warmed to his theme. "It's based on the idea that you can actually curve space," he told me. "Einstein discovered that large masses curve space. Well, it's the same thing with a wormhole. You're curving and distorting space so that you don't have to follow the usual dimensions—you can go up and out, so to speak, into another dimension, if you will. And that allows you to travel much faster than the speed of light."

I asked him if he ever got the feeling that anyone was working on this stuff elsewhere; that there may even have been breakthroughs—ten, twenty, maybe forty years earlier. His reply was more candid than I had anticipated.

"That's always a possibility. I'm not one for conspiracy theories, but we've learned about things that governments and organizations have been involved in in the past."

He paused and rubbed the bristle hairs at the back of his head. I waited.

"It would be extremely frustrating to think that we're duplicating what was done maybe decades before. We like to think we're on the cutting edge, at the forefront of knowledge. It would be very discouraging to think that everything had been determined before and we were just sort of . . . here for show. I hope that's not the case. But you never know. Maybe in some aspects it is."

It was then that my PR escort interjected.

"You know," she said, "I really don't think this is anything that we can comment on in any depth."

I knew that the topic of conversation had taken an unwelcome turn. Now, I had intruded far enough.

"I'm simply trying to get a feel for the kind of capabilities that are out there—on both sides of the fence," I replied.

"Then maybe you should go talk to the Air Force," she said crisply. In any case, I was talking to the wrong people at NASA, she added. Huntsville was where the big picture behind the interstellar vision came together. But, she said, the "breakthrough physics" that would one day lead to the engineering of a "device" was being managed by the NASA Glenn Research Center at Lewis Field in Cleveland. It was from there, not here, that NASA was actively in the process of handing out contracts that aimed to prove or disprove whether there was any substance to the theory.

If I wanted to know whether the physics was doable, get up there, she told me. Talk to a guy called Millis. Marc Millis.

Before I left, I had one last question for Schmidt. I asked him if he genuinely believed that a wormhole might one day give man the ability to cross the universe.

"Yeah," he said, smiling but serious at the same time. "If you can build a wormhole, certainly. Who knows? But you've gotta be careful to make sure you've got the right end point. Who knows where you'll end up if you just generate it? You could end up anywhere in the universe. So that's something to be very careful about."

Six hundred miles to the northeast in Cleveland, Ohio, but a world away culturally, Marc Millis had indeed just received a half-million-dollar contract to test whether there was any theory out there, anything at all, that might one day enable a propulsion device to be built with the energy to take man out of the Solar System and into interstellar space.

Millis ran the NASA Breakthrough Propulsion Physics program, referred to in the trade as BPP.

"This is it," he told me, gesturing to the modest office he shared with another NASA official whose head was just visible on the other side of the screen that set the boundaries of Millis' work space. It had taken some hard negotiating after Huntsville, but I had managed to secure a visit to the Glenn Research Center, NASA's "center of excellence for aeropropulsion research and technology," on my way to Washington, where I was due to conduct some interviews for the magazine before heading on down to Puthoff's place in Austin, Texas.

With his half a million dollars, Millis had awarded five contracts, any one of which might lead to the breakthrough that gave BPP its name.

The first contract, placed with the University of Washington, Seattle, was designed to test theory which maintained that a change in inertia was possible with a sudden energy-density change, induced, say, by a massive jolt of electricity. Manipulate inertia—an object's innate resistance to acceleration—and apply it to a spacecraft and you had the ability to reduce, if not negate altogether, its need for propellant. If the sums were right, it would continue to accelerate until it reached light-speed.

The second experiment was tasked with examining the reality of the so-called zero-point energy field and whether it existed at anything like the magnitudes that had been postulated by some scientists. If these people were even half-right, Millis said, zero-point energy held enormous significance for space travel and offered entirely new sources of non-polluting energy on Earth.

What, I asked, was zero-point energy?

For some years, Millis replied, there had been a developing understanding that space was not the empty vacuum of traditional theory, but a seething mass of energy, with particles flashing in and out of existence about their "zero-point" baselines. Tests indicated that even in the depths of a vacuum chilled to absolute zero (minus 273.15°C)—the zero point of existence—this energy would not go away. The trouble was, no one knew quite where it was coming from. It was just there, a background radiation source that no one could adequately explain. With millions, perhaps billions, of fluctuations occurring in any given second, it was theoretically possible to draw some—perhaps a lot—of that energy from our everyday surroundings and get it to do useful work. If it could be "mined"—both on Earth and in space—it offered an infinite and potentially limitless energy source.

The third experiment, he continued, would set out to examine the unproven link between electromagnetism and gravity and its possible impact on space-time. Perturbations of space-time not only promised a

propulsive antigravitational effect, but opened up other more esoteric possibilities, such as time travel.

I asked Millis if this stuff was for real and for a moment he looked unsure. The mere mention of time travel and we were suddenly into science-fiction territory. It was enough to put any serious scientist on the defensive.

"The immediate utility is not obvious," he said stiffly, "but it's like a foot in the door."

The fourth experiment looked at "superluminal quantum tunneling"—faster-than-light speed. Recent laboratory tests had shown light pulses accelerating beyond the speed of light, thereby shattering Einstein's Theory of Relativity, which said that the light-speed barrier was unbreachable. If a light pulse could arrive at its destination before it left its point of departure, the theory said a spacecraft might be able to as well.

"How it translates into space travel," Millis said, anticipating my next question, "is way down the pike."

Again I was presented with a failure of fit. If NASA was still struggling with the theory now—40 years and a world of knowledge after the pronouncements of George S. Trimble and his colleagues in 1956—what on earth had led them to believe they could crack gravity then?

Something I'd written down tugged me back to the present. I flicked back over the list of experiments and did a quick inventory count.

"You said there were five contracts. I've only got four. What's the fifth?"

He looked momentarily taken aback.

"It's conducted under the auspices of the Marshall Space Flight Center itself," he said. "Since you'd just come from there I rather presumed they'd briefed you on it."

I shook my head. "Briefed me on what?"

Millis reached into a drawer and pulled out a file full of newspaper clippings and magazine articles. He rifled through them until he found what he was looking for, a clipping from Britain's *Sunday Telegraph* dated September 1, 1996. The headline read: "Breakthrough As Scientists Beat Gravity."

"I'm surprised you didn't know about this," he said, pointing to its source.

I said nothing. I had a photocopy of the same story in my files at home. I'd not taken much notice of it because it had seemed totally outlandish.

The story detailed the experiments of a Russian materials scientist called Dr. Evgeny Podkletnov, who claimed to have discovered an

Breakthrough as scientists beat gravity

by ROBERT MATTHEWS
and IAN SAMPLE


SCIENTISTS in Finland are about to reveal details of the world's first anti-gravity device. Measuring about 12in across, the device is said to reduce significantly the weight of anything suspended over it.

The claim — which has been rigorously examined by scientists, and is due to appear in a physics journal next month — could spark a technological revolution. By combatting gravity, the most ubiquitous force in the universe, everything from transport to power generation could be transformed.

The Sunday Telegraph has learned that Nasa, the American space agency, is taking the claims seriously, and is funding research into how the anti-gravity effect could be turned into a means of flight.

The researchers at the Tampere University of Technology in Finland, who discovered the effect, say it

HOW THE ANTI-GRAVITY DEVICE WORKS



Unit approx. 12in diameter

tures. The team was carrying out tests on a rapidly spinning disc of superconducting ceramic suspended in the magnetic field of three electric coils, all enclosed in a low-temperature vessel called a cryostat.

"One of my friends came in

the *Journal of Physics-D: Applied Physics*, published by Britain's Institute of Physics.

Even so, most scientists will not feel comfortable with the idea of anti-gravity until other teams repeat the experiments.

antigravity effect while working with a team of researchers at Tampere University of Technology in Finland. What made the claim different from "so-called 'antigravity' devices put forward by both amateur and professional scientists," wrote the paper's science correspondent, was that it had survived "intense scrutiny by skeptical, independent experts." It had been put forward for publication in *Journal of Physics D: Applied Physics*, published by Britain's Institute of Physics.

When the *Telegraph* hit the streets, Podkletnov's world turned upside

down. He was abandoned by his friends, ostracized by the university, which claimed that the project did not have its official sanction (and later threw him out), and the journal pulled his paper. Leaving his wife and family in Tampere, Podkletnov returned to Moscow to lick his wounds until the furor had died down.

And all because the *Sunday Telegraph* had used a single, taboo word in its opening sentence: antigravity. The term may have been vital for 1950s pulp sci-fi, but it had no place in the world of mainstream physics.

What was unfair, was that Podkletnov had been so careful to avoid the term himself. All along, he had referred to the test as a *gravity-shielding* experiment.

It was still heresy; and he had burned for it.

But here was Millis saying that NASA *had* taken Podkletnov's claims seriously. Not only that, but NASA had been apprised of the "gravity-shielding" properties of superconductors even before the *Sunday Telegraph* story broke.

I wondered why no one had told me about all this while I was in Huntsville.

In 1993, the Advanced Concepts office at the Marshall Space Flight Center was handed a copy of a paper written by two physicists, Douglas Torr and Ning Li, at the University of Alabama at Huntsville. It was called "Gravitoelectric-electric coupling via superconductivity" and predicted how superconductors—materials that lose their electrical resistance at low temperatures—had the potential to alter gravity. Even if the amounts were fractional, NASA officials quickly saw the benefits. Applied to something like a rocket, even fractional methods for cutting launch weights would, over time, lead to a considerable reduction in the space agency's fuel bills.

In their paper, Torr and Li went out of their way to avoid the term "antigravity." One whiff of it, they knew, and no one would take their work seriously.

It worked. NASA promised money for an experiment.

But when the Podkletnov story broke, Millis told me, NASA decided to take a multi-path approach instead. First it tried to contact Podkletnov to see if he would share his findings with them. When the Russian refused, it elected to do the experiments itself, setting up the "DeltaG group" at Marshall specifically to look for interactions between superconductors and gravity.

At first, Ning Li, the brilliant Chinese woman scientist who had predicted a one percent weight change in her original calculations, joined forces with the DeltaG effort. Then, apparently frustrated by the pace of

work at Marshall, she set up to pursue the experiments on her own. If she was getting anywhere with them, she wasn't saying, all her most recent tests having been conducted behind closed doors.

Since that time, according to Millis, the DeltaG group had found cause to believe that *something* unusual was happening when gravity fields were influenced by superconductors, although the precise nature of the effect was a matter of conjecture. Podkletnov had claimed that two percent weight reductions were being consistently observed during his experiments—a result, if you could call it that, that was so close to the Torr/Li prediction it was scary.

Like the whirling gyroscopes investigated by BAe, the low percentages were, in a sense, immaterial. What was being talked about here—and apparently engineered by Podkletnov—was a phenomenon that science said was impossible.

The DeltaG experiments at Marshall continued, but, without Podkletnov's input, Millis admitted that things had not gone as fast as NASA would have liked. Part of the problem had been its inability to reproduce superconductors of the size and fidelity Podkletnov said was required to achieve gravity shielding—a term with which even Millis, who was broader-minded than most, was uncomfortable.

"It's more accurate to call it an anomalous gravitational effect, but even that is jumping the gun." It could, he said, simply be an experimental error.

But what if it wasn't? If Podkletnov was right, if he had found a way of altering an object's weight, aerospace was about to change—civilization, too, maybe—and forever. Conscious, no doubt, that it might be staring at the greatest scientific discovery since the bomb, I could see why NASA was hedging its bets.

It had solicited a company in Columbus, Ohio, to try to develop superconducting ceramic discs like those built by Podkletnov. Without the Russian's secret recipe, however, it hadn't been getting on as well as it might. At the time of my visit to Ohio, the Columbus-based company had managed to build small discs, but it still hadn't acquired the knack for turning out the larger variety—the type that Podkletnov said was crucial to the success of the experiment. And this was the bittersweet irony. Being a materials scientist, Podkletnov had an insight into the way the discs were constructed that left NASA, with its \$13 billion annual budget, floundering in his wake. Tempting as it was to view Huntsville's silence over Podkletnov as a deliberately misleading omission, I suspected that NASA's impotence in the face of this irony was the real reason no one had mentioned him.

* * *

On the flight down to Texas, I picked through a pile of clippings I'd managed to track down during my brief stopover in Washington. From the little that was known of Podkletnov's Tampere experiments, it was apparent that the superconductors needed to be spun at around 5,000 rpm inside a large steel container filled with liquefied gas for cooling. Electromagnets were used to make the superconductors levitate and get them up to speed. In the absence of any firm data on Podkletnov's discs, though, attempts by NASA to duplicate his work were inconclusive as far as gravity shielding was concerned. A couple of times, the readings had blipped—a little like Dr. Evans' experience with the Kidd inertial thrust machine at BAe—but they were never consistent enough to be taken seriously.

If Podkletnov knew all this, I thought—and somehow or other, I guessed he probably did—then he was more than likely pissing himself laughing.

Either he had perpetrated a fantastic confidence trick, or he really had pulled off one of the greatest scientific discoveries of the 20th century.

So, where the hell was Podkletnov?

The best and only clues to his whereabouts had been provided by an American journalist named Charles Platt who had documented his attempts to track down Podkletnov in a lengthy article for *Wired* magazine. After the *Sunday Telegraph* story, Platt managed to get a phone number for Podkletnov and they spoke a couple of times, but only on the basis that Platt wouldn't publish anything without the Russian's consent.

"He told me how he made his discovery," Platt wrote more than a year later, quoting the Russian: " 'Someone in the laboratory was smoking a pipe and the pipe smoke rose in a column above the superconducting disc. So we placed a ball-shaped magnet above the disc, attached to a balance. The balance behaved strangely. We substituted a non-magnetic material, silicon, and still the balance was very strange. We found that any object above the disc lost some of its weight, and we found that if we rotated the disc, the effect was increased.' "

When Platt suggested that huge amounts of energy would be required for this to be remotely so, Podkletnov reportedly became irritated.

"We do not need a lot of energy," he'd snapped back. "We don't absorb the energy of the gravitational field. We may be controlling it as a transistor controls the flow of electricity. No law of physics is broken. I am not one crazy guy in a lab; we had a team of six or seven, all good scientists."

And that, for the next year or so, was that. It was only when an intermediary told Platt that Podkletnov had returned to Finland, where he was now set up as a materials scientist in a local company, that the American jumped in a plane and finally managed to confront the man who'd dared to claim he had shielded gravity.

What Podkletnov had been doing during his yearlong self-imposed exile in Moscow was never made clear, although the Russian soon hinted that he had not been idle. He was quick to explain to Platt that his original findings at Tampere had been meticulously charted using a mercury barometer. Immediately above the superconducting disc, it had registered a 4 mm drop in air pressure, because, the Russian said, emphasizing the point, the air itself had been reduced in weight.

When he'd taken the barometer upstairs, Podkletnov had found the same drop in pressure over the point where the experiment was taking place on the floor below. He went up to the top floor of the building and it was the same thing.

This showed that gravity reduction would not diminish with distance, that the effect had no limit. Podkletnov's gravity shield went on, extending upward in a 30 cm diameter column—forever.

And then, if that wasn't good enough, pay dirt. The two percent weight reduction in all the air above the disc meant that a vehicle equipped with gravity shielding would be able to levitate, buoyed up by the heavier air below.

"I'm practically sure," Podkletnov had told Platt, giving him an intense look, "that within ten years this will be done. If not by NASA, then by Russia."

During his year in Russia, he went on to reveal, he had conducted research at an unnamed "chemical scientific research center" where he had built a device that *reflected* gravity. By using superconductors, resonating fields and special coatings—and "under the right conditions"—gravitational waves had been repelled instead of blocked. Applied to an air vehicle, Podkletnov said, this "second generation of flying machines will reflect gravity waves and be small, light and fast, like UFOs. I have achieved impulse reflection; now the task is to make it work continuously."

Their meeting over, Podkletnov slipped back into the shadows and Platt returned home. While there had been sightings of Podkletnov in the years since—reports of his whereabouts had surfaced in Japan, Russia and Finland—he had, to all intents and purposes, disappeared again.

Back in the States, Platt canvassed the physics mainstream for its

reaction to the Podkletnov claims. Most scientists laughed outright. Gravity shielding, they said, was simply inconceivable.

I made a note on the flight home to ask Marckus to keep his ear to the ground for signs of the Russian's whereabouts. It was tempting to believe that Podkletnov was by now working in the bowels of a crumbling military research institute somewhere deep behind the Urals, refining his experiments to the accompaniment of a howling Siberian gale. And maybe this wasn't a million miles from the truth.

But scientists had a habit of talking to one another, whatever obstacles seemed to separate them. And since Podkletnov had shown himself to be loquacious enough when it suited him, I thought he'd probably surface again before too long.

When he did, I needed to be there.

I was intrigued by the Russian's work. It echoed something that Professor Young had said in his lecture on antigravity—something that had sounded right at the time. If gravity control is ever to be discovered, Young had said, it would most likely be as an unexpected by-product of work in some completely different field. An accident.

Podkletnov hadn't been looking for an antigravity effect. He'd been playing around with superconductors. If he'd been working in a Western laboratory, his discovery might have forever gone unnoticed. But in Russia—and Finland, too, apparently—it was OK to light up a pipe in a room full of test equipment. And, bingo, the smoke had hit the column of shielded air and snaked straight up to the ceiling.

It struck me as not impossible that Podkletnov had discovered—in a million-to-one fluke—what Trimble et al had predicted theoretically in 1956.

Had Podkletnov tripped over the answer and got in by the back door?

If anyone knew, it was Dr. Hal Puthoff, the second of the two contacts passed to me by Marckus during our meeting by the estuary.

Even in the cocooning environment of the aircraft, nursing a stiff drink, I felt anxious at the prospect of meeting him.

Up until now, I'd managed to palm off my interest in NASA's advanced propulsion activities as a quirky sideline to my regular journalistic pursuits. I knew, though, that when I pitched up at Puthoff's door, a discreet address remote from any other legitimate aerospace business, there'd be no hiding what I was really up to.

Chapter 10

I reached Puthoff's offices after a drive through the industrial suburbs of the Texas state capital, parked and double-checked that I had the right address. I had expected the Institute of Advanced Studies to be housed in another windowless science block, a modern version of the concrete edifices at NASA, replete with clean rooms and scientists swishing around in white coats. But the IAS was a small glass-fronted office in a leafy business park. Several of the other lots were vacant and at least one was boarded up.

Puthoff's reputation as a scientist of vast repute went a long way before him. His papers on gravity and inertia were viewed as seminal works by the science community. He also sat on something called the Advanced Deep Space Transport group, ADST, a low-profile organization composed of people that someone had described as being "on the frontier of the frontier." Staffed by select NASA, Air Force and industry representatives, it was the real cutting edge, but you had to know where to look for it.

Puthoff's link to NASA was bound up in theoretical work that he had been conducting since the early 1970s into the zero-point energy field.

Before getting out of the car, I was beset with a rush of all the old doubts. Did I really want to declare my interest in antigravity to a man who had clear connections to the intelligence community?

I checked my watch. It was a little after five-thirty in the morning back in England, but I needed reassurance. I called Marckus on my cell phone.

He picked up on the second ring, sounding bright and alert. I wondered for a moment whether Marckus ever slept, then dispensed with the small talk and told him of my concerns.

Marckus, showing a tendency that would become familiar over the course of our acquaintance—a tendency to impart knowledge well after the time I needed it most—inform me that NASA, being a government institution, was, in effect, the outer marker of an elaborate but invisible security system.

"How's that?" I asked.

"You show up asking the kind of questions you asked at Huntsville and

Cleveland and, believe me, someone, somewhere will have registered the fact. I have little doubt that you began pulsing away on radar scopes at government agencies too numerous to mention the moment you even logged your requests to visit NASA. You've already crossed the threshold. It's too late to go back now."

I felt the muscles in my stomach tighten, but I said nothing. This propensity to unsettle was also a part of Marckus' act. I asked him instead, as casually as I could, for a data dump on everything that was known about zero-point energy (ZPE).

I'd already grasped the principles: that space—the air that we breathed, the atoms in our bodies, the far reaches of the cosmos, everything—was filled with a churning "foam" of energy. Its presence was signified by the background static hiss of the universe—something you could hear on a transistor radio—but as far as visual detection went, was invisible. Which was what made it so controversial.

In 1948, Marckus told me, a Dutch physicist, Hendrik Casimir, came up with a theoretical model for proving that the "quantum vacuum" existed. If two aluminum plates were placed very close together, so close that their separation was less than the wavelengths of the fleeting particles in the quantum foam, it followed, Casimir reasoned, that there would be nothing between the plates.

Because the theory said that everything outside the plates would still be seething with zero-point fluctuations, the external force pushing in on the plates ought to be enough to close them together, thereby proving the existence of the ZPE field.

The difficulty, however, related to the experiment. To exclude the wavelengths of the fluctuating particles, Casimir's plates had to be exactly parallel and separated by less than one micron—one hundredth the width of a human hair. It wasn't until 1997 that science was able to come up with equipment that could satisfy the tolerances required for the Casimir experiment. When it did, it found that the Casimir Effect existed exactly as predicted.

And with it flowed the corollary that the quantum vacuum, zero-point energy—call it what you will—existed as well.

In principle, there was no reason why mechanisms couldn't be built to tap into this energy.

And since the fleeting particles were popping in and out of existence billions of times every second, on every conceivable frequency, in every possible direction, there was, theoretically speaking, no limit to the amount of energy resulting from the fluctuations.

In principle. Theoretically speaking.

"The argument now is not about whether the ZPE field exists," Marckus concluded, "but about how much energy it contains. It comes down to a difference in people's calculations and interpretations. Some say that there's enough energy in a shoe box to blow the world apart; others say that all the ZPE in the volume of the Earth couldn't boil you an egg with it." He paused. "I think you'll find with Puthoff that it's closer to the shoe-box scenario." And then he signed off.

I locked the car, walked over to the IAS office, pushed the glass and watched as it swung open. There was no one in the front office, so I called out a couple of times and, not getting any answer, moved toward the back of the building. I followed a corridor till I heard a man's voice. Poking my head around a door, I saw Puthoff sitting on the edge of his desk, a phone jammed to his ear. He was shorter than he'd appeared in photographs, but his puckish features and thick head of hair belied his 60-odd years. He waved me in, pointed with a mock grimace to the voice on the other end of the receiver, and gestured for me to grab a seat. A couple of minutes later, he hung up, muttered something about a canceled meeting in Washington, and asked if I'd like a coffee.

While we were standing by the coffee machine, I asked him which of Marckus' analogies he inclined toward.

"If we're right," he said, as I stirred creamer into my coffee, "and you have all possible directions of propagation in the universe, and all possible frequencies and wavelengths, and every one of them has this little contribution of zero-point energy, when you add it all up, the numbers are ridiculous."

"How ridiculous?" I asked.

He paused and added matter-of-factly: "There's enough energy in the volume of your coffee cup to evaporate all the world's oceans many times over."

He showed me a large workshop at the back of the building filled with workbenches, electrodes and measuring equipment. This, Puthoff said, was where he and his small team worked on the ZPE devices that people routinely sent him for analysis; tiny little reactors that supposedly drew energy out of the vacuum of space: the space that we occupied and the space, in its quantum sense, that occupied us.

The machines came in all shapes and sizes. Some were solid-state devices with no moving parts that used the pinching motion of the Casimir effect to create clusters of electrical charge with energy outputs many times greater than the kick they required to get going.

Others used the millions of imperceptible yet infinite movements of the quantum foam to stimulate a vibrating motion in magnetic fields.

These vibrating fields, their proponents claimed, served as "gates" via which unending supplies of vacuum energy could be fed through electronic circuits and put to use.

If they could be proven to work repeatedly and reliably, Puthoff said, imagine it: no more pulling power off the national grid. You'd have a reactor in a box in your backyard; a little thing the size of a microwave oven that sucked energy from the space that it occupied and belted out clean, unending power wherever it was needed. Initial machines would feed power to your house, but later, as they developed in sophistication, just as computers had developed exponentially at the end of the 20th century, these devices would shrink in size and double in output; and they'd keep on shrinking and doubling, over and over, until they'd be small and powerful enough to put in cars or aircraft or submarines. Applications would be limited only by the imagination of the user.

"If," I said. "You said, 'if.' "

"Ah, well, that's the point," he replied. "Of the 30-odd devices that have come through this door, none has passed the magic test yet: a demonstrable measurement of more energy flowing out than is flowing in."

But it would happen. Maybe tomorrow, maybe in 30 years. But for sure, one day someone was going to announce that they'd invented a machine that could extract energy from the immeasurable pulse of the universe. It'd rate a paragraph in the news-briefs columns, but clip it or keep the paper, mark it well, Puthoff said, because on that day the world would change. Nothing would ever be the same again.

As he talked, Puthoff filled in the gaps in the résumé that I'd pulled off the Net. Soon after gaining his master's degree at the University of Florida, he went on active duty as a Naval Intelligence officer seconded to the National Security Agency. When he left the Navy, he converted to civilian status at the NSA, then went on sabbatical at Stanford University in California to get his Ph.D. He gained his Ph.D. in 1967, resigned from the NSA and soon afterward joined the Stanford Research Institute, a spin-off from the university, set up, among other things, to pursue heavily classified research for the U.S. defense and intelligence communities. Puthoff's thing was lasers, but it was at SRI that he developed the notion of the remote-viewing program for the CIA and the DIA. The RV work formed the hub of the "heavily classified research" that he undertook at SRI for the next 13 years.

I asked how he'd made the jump from lasers to remote viewing.

"It sounds weird, but it was a very straightforward thing," he said. In the early 1970s, physicists were searching everywhere for evidence of

tachyons—particles that theoretically had the ability to travel faster than light. Finding them was problematic, because, if the theory was right, they couldn't travel below light-speed, making identification somewhat challenging.

Someone had given Puthoff a book called *Psychic Discoveries Behind the Iron Curtain*, which mentioned how Soviet scientists were looking for evidence of ESP in plant life. The Russians had connected up a pair of plants to polygraphs, separated them over huge distances and shielded them from all electromagnetic frequencies. They found that when one plant was burned with a cigarette, or even if someone was thinking about it, the other plant responded on the polygraph.

Since one of the postulated characteristics of tachyons was that they could penetrate all kinds of shielding, it occurred to Puthoff that one of the few places that scientists hadn't looked for tachyons was in "organic living systems." He wrote up a proposal that aimed to verify if plants were "tachyonically connected" by linking separated algae cultures with lasers. He sent the proposal to an acquaintance involved in similar work who happened to show it to Ingo Swann, a noted psychic and medium, who in turn approached Puthoff, assuming that he was interested in ESP. Even though he wasn't, Puthoff was sufficiently intrigued to invite Swann to SRI, where he proposed to immerse him in a "shielded quark magnetometer," a "tank" in a vault that was totally cut off from all known emissions and frequencies.

As Swann approached the vault, which he hadn't been forewarned about, he appeared to perturb the operation of the magnetometer. The dials and instruments inside it all flickered into life. He then added insult to injury by remote viewing the interior of the apparatus, "rendering by drawing a reasonable facsimile of its rather complex construction," according to Puthoff, even though details of it had never been published anywhere.

Puthoff wrote up his findings, circulated the report among his colleagues and got on with other work.

A couple of weeks later, the CIA showed up at his door and invited him to set up a classified research program to see if it was technically feasible to use clairvoyance to spy on the Soviet Union. It turned out that the CIA had been tracking Soviet efforts to do the same thing for several years, but didn't believe what it was seeing. It was only when Puthoff wrote up his paper that it put two and two together and thought it'd better set up something of its own. And so the U.S.-controlled remote-viewing program was born, with Puthoff as its director.

For the next decade and a half, U.S. "psychic spies" roamed the Soviet

Union, using nothing more than the power of their minds to reconnoiter some of the Russians' most secret R&D establishments. If you believed what you read in published accounts of these people's activities, they were able to roam not merely in three dimensions—up, down, left and right—but in the fourth dimension as well: time. They could go back in time to review targets and they could look at them in the future as well.

Puthoff showed me a picture that one of his team of remote viewers had drawn of an "unidentified research center at Semipalatinsk," which had been deeply involved in Soviet nuclear weapons work. He then showed me some declassified artwork of the same installation, presumably drawn from a U.S. spy satellite photograph. The two were damn near identical.

It all seemed a long way from zero-point energy, gravity and inertia, in which Puthoff had been busying himself even before he left the remote-viewing field.

Or was it?

In a recent paper written by Puthoff, which I had already seen, he'd mentioned that the Russian physicist and Nobel prizewinner Andrei Sakharov had published a paper in 1967 suggesting that gravity and inertia might be linked to what was then still a highly theoretical proposition: vacuum fluctuations of the zero-point energy field. Now that the zero-point energy field had been proven to exist, "there is experimental evidence that vacuum fluctuations can be altered by technological means," Puthoff had written in the paper. "This leads to the corollary that, in principle, gravitational and inertial masses can also be altered."

Puthoff was too smart to use the term—for all the reasons that Podkletnov, Ning Li and the others hadn't—but he was saying that antigravity was indeed possible.

And, so were the Russians. All you had to do—somehow—was perturb the zero-point energy field around an object and, hey presto, it would take off.

I asked him why NASA and the Air Force should be so interested in ZPE when it was obvious that there weren't going to be any practical applications of it for years—maybe centuries.

Puthoff looked at me in a meaningful way. "Unless we find a shortcut."

"Is that a possibility?"

"Podkletnov could have come across a shortcut."

The maverick Russian materials scientist and the idea of pipe smoke hitting a gravity shield had entered the conversation so abruptly that it

caught me off guard. Puthoff was saying that anything was possible. And this from a man who had spent most of his professional life plugged into the heart of the classified defense environment.

I probed further, but gently. Were there already forms of aerospace travel out there—in the black world, maybe—whose principles contravened, if not the laws of physics, then at least our understanding of aerodynamics?

He sucked the top of his pen, giving the question a lot of thought before responding. "I've certainly talked to people who claim that something is going on," he said, pausing to add: "I would say the evidence is pretty solid."

I felt myself rock back on my chair.

I eased back on to the more comfortable subject of the NASA BPP study. "Which of the five methods outlined in Breakthrough Propulsion Physics would you say is the one most likely to have a payoff?" I asked him.

Because of Puthoff's promotion of ZPE, I thought it inevitable that this explanation would have figured in his answer. But it didn't. Without even thinking about it, he said his money was on the third experiment—the one about perturbing space-time; antigravity from time travel.

Given Puthoff's strong connections to the military-intelligence community, it was tempting to dismiss all such talk as a deliberate blind, something to lead me away from a more probable area of breakthrough. But it didn't seem that way at all.

I liked Puthoff. He seemed on the level. Besides, one thing I had learned after years of formal briefings and interviews was how difficult scientists found it to lie when confronted with a direct question on their chosen subject. Spooks could lie at the drop of a hat. It was what they were trained to do. But scientists found it inherently unnatural, since it contravened all their instincts to match a question quid pro quo with data.

Puthoff's answer to the question had been swift and visceral. And there had been something else, too; a look on his face, a willingness to confront my gaze, that in itself conveyed information.

We'd been talking about breakthroughs—events that, at a stroke, had the power to short-circuit the evolutionary development of technology.

The important thing about breakthroughs was that they didn't happen over an extended period of time. They changed things overnight. And they could occur at any moment.

It was only later, when I was back on the freeway, crawling south in the bumper-to-bumper traffic, that I had the feeling he'd been trying to tell me that some kind of payoff from the third experiment had already produced tangible results.

Chapter 11

If antigravity had been discovered in the white world, then someone, somewhere had to be perfecting it—maybe even building real hardware—in the black. The trouble was, I had no idea where to begin looking for it. Contrary to the implicit promise of revelation contained in Dr. Dan Marckus' hints by the estuary, neither Lyles nor Puthoff had taken me any further in so far as hard-and-fast leads were concerned.

On my return, I tried to raise Marckus. I left several messages on his answering machine. In the first of them I foolishly spelled out the fact that I didn't know a whole lot more—beyond strict theory, at least—than I had before my trip to the States.

During subsequent calls, as the answering machine clicked in, I just knew that Marckus was there—listening in, but refusing to pick up. I was beginning to get the measure of this relationship. Marckus gave a little and I had to give something back. He now knew that I had returned from the States with nothing, so he was punishing me for it. Well, I didn't need to play psych games with some spook academic with a warped taste for drama. I had enough on my plate already—a major feature article on stealth technology to deliver. Antigravity, for the time being, was going to have to go on the back burner and Marckus could stew with it.

As background, I reread an account of what is probably the most definitive history of stealth to date: *Skunk Works*, the story of Lockheed Martin's super-secrective special projects facility, as told by the guy who had run it for 16 years, Ben Rich.

In the summer of 1976, America's leading radar expert, Professor Lindsay Anderson of the Massachusetts Institute of Technology, arrived at what was then the Lockheed Skunk Works in Burbank, California, with a bag of ball bearings in his briefcase. The ball bearings ranged in size from a golf ball to an eighth of an inch in diameter. Rich, a grizzled veteran of Lockheed's secret projects department for more than 25 years, led Anderson to a mocked-up design of a funny-looking diamond-shaped aircraft, poised on a plinth in the middle of a blacked-out hangar.

Over the course of the day, Anderson repeatedly attached the ball

bearings to the nose of the aircraft, starting with the largest and ending with the smallest, while Lockheed engineers fired radar signals at it. What they were looking to measure was the radar cross section, or RCS, of the prototype aircraft, code-named Have Blue. The moment the radar picked up the ball bearing instead of the aircraft behind it was the moment its RCS could be accurately plotted.

When the test was over, Anderson zipped the bearings back into his case, left the site as quietly as he'd arrived and returned to MIT to process the data.

A few days later, Rich received a telegram from the air force chief of staff. Have Blue had been classified "Top secret—Special Access Required," a security grading reserved only for "quantum leaps"—programs that jumped right over the state of the art. The first of these super-secret "black" programs had been the atomic bomb. The telegram had just pitched "stealth" into the same category of secrecy.

Have Blue's RCS had bettered every single one of the ball bearings Anderson had attached to it—even the one that was sized down to an eighth of an inch. This gave the aircraft and its operational successor, the F-117A Stealth Fighter, a radar signature equal to that of a bee-sized insect. Stealth, which up to that point had been viewed by skeptics as nothing more than puff and magic, suddenly went "deep black"—so secret that no one outside the program would ever know it was there. Lockheed had discovered a way of making aircraft invisible to radar, but the stakes were far, far higher even than that.

Soon after the late 1977 first flight of Have Blue from the U.S. Air Force's remote testing base on the dry desert lake bed at Groom Lake, Nevada, also known as Area 51, Dr. Zbigniew Brzezinski, President Jimmy Carter's national security adviser, flew into the facility in an unmarked jet and grilled Rich about the true significance of his firm's breakthrough. Rich put it on the line for him.

"It changes the way that air wars will be fought from now on. And it cancels out all the tremendous investment the Russians have made in their defensive ground-to-air system. We can overfly them any time at will."

"There is nothing in the Soviet system that can spot it in time to prevent a hit?" Brzezinski asked, jotting notes as he talked and listened.

"That is correct," Rich replied simply.

After more than three decades of standoff between the two superpowers, the realization made Carter's national security chief catch his breath. If the Russians wanted to match stealth with counterstealth, theoretically they could do it, but it would bankrupt them in the process.

Stealth had the power to end the Cold War, but it also had the power in the meantime—should the Russians realize what the Americans were up to—to trigger World War Three.

Have Blue was a strange-looking aircraft. It didn't look mean and hungry. It didn't look like the kind of plane that could start and end wars. It had the appearance of a bunch of bolted-on geometric shapes that had somehow been hammered into the vague outline of a jet. It turned out there was more than a good reason for that.

Have Blue hadn't been the creative inspiration of an aerodynamicist, but of a mathematician.

Begin with what you can see, then search for what you can't.

As I wrestled with the stealth feature, Marckus' words from our meeting battered around my head, forming as a dark thought with a black body and soft wings.

I had trawled the white world and found five possible pathways to antigravity: manipulating an object's mass and/or inertia; exploitation of the zero-point energy field; perturbations of the space-time continuum; faster-than-light travel; and gravity shielding. None of them was visible in the developmental activities of the U.S. aerospace industry, but just because you couldn't see them, it didn't mean they weren't there.

Just as it had, or hadn't been with stealth a decade and a half earlier.

Ben Rich and I had sparred on a number of occasions on the stealth question—most recently at an air show where he'd turned up, desperately ill with cancer, to promote his book. I'd respected him utterly and liked him hugely, sensing in his presence the grit and wisdom of a generation of postwar aviation pioneers that wouldn't be around us for much longer.

It was Rich who'd once told me of a place—a virtual warehouse—where ideas that were too dangerous to transpose into hardware were locked away forever, like the Ark of the Covenant in the final scene of *Raiders of the Lost Ark*. It had almost happened to stealth.

By starting with stealth, something that I knew about, something I could *see*, maybe I could pick up traces of that energy, the friction that astronomers looked for when hunting down black holes.

In the year ahead, the calendar of aerospace and defense events showed that I had multiple excuses for visiting the States—to look for the self-generated heat of a buried antigravity effort's interaction with the real world.

As I sat behind my desk, calling contacts and sources in an effort to plug holes in the incomplete, still highly secret history of stealth, I could hear the faint echo of a signal. It was almost impossible to decipher, but

the cold side of my training, the part I kept separate from the muffled wing beats of the creature let loose upon my thinking by Marckus, said antigravity had something to do with the shared origins of the two most powerful weapons of the 20th century—stealth and the atomic bomb.

Have Blue would never have happened, but for the work of a retired Lockheed math guru named Bill Schroeder and a Skunk Works software engineer named Denys Overholser. Schroeder revisited a set of mathematical formulae originally derived by Maxwell and refined by a turn-of-the-century German electromagnetics expert called Arnold Johannes Sommerfeld. Between them, these two physicists predicted the manner in which a given geometric shape would scatter or reflect electromagnetic radiation. A Russian, Pyotr Ufimtsev, honed their work into a more simplified set of equations in the early 1960s, but Ufimtsev could only apply the equations to the most basic geometric shapes. Schroeder's and Overholser's breakthrough had been to take Ufimtsev's concept and apply it to the inherently complex form of an aircraft, rejecting the regular smooth, curved lines of aerodynamic aesthetics in favor of a set of ugly flat, angled panels. If an aircraft could be broken down into hundreds, maybe thousands of these two-dimensional shapes, if each flat surface or "facet" could be angled in a way that would reflect an incoming radar beam away from its source, and if the combined shape could still create lift, then a true "stealth" aircraft was possible.

The irony, of course, was that stealth was in part a Soviet invention. The Soviets had allowed Ufimtsev's paper to be published, because no one east of the Iron Curtain had entertained the idea that such a complex set of equations could ever be applied to an aircraft. It had taken Overholser to pick it out from hundreds of other ponderous Soviet science texts and draw the necessary conclusions.

Stealth, like the bomb, then, had owed its genesis to algebraic formulae. In a sense, it was pure accident that it had been discovered within the nuts-and-bolts world of the aerospace industry.

Podkletnov's apparent discovery of gravity shielding in Tampere, Finland, had been an accidental by-product of the Russian's work with superconductors.

What they all shared was their origins in pure math and physics.

That thought gnawed at me as, working through a decade's worth of my files, I began to reimmerse myself in the world of the F-117A Stealth Fighter, Have Blue's operational successor.

Lockheed was used to black programs. It had built the U-2 spyplane that the CIA had used to overfly the Soviet Union on reconnaissance

missions in the late 1950s and it had developed the Mach 3.2 A-12 for the CIA as the U-2's successor. Both projects had come together under rules of draconian secrecy, but stealth, from a security standpoint, would need to be protected even more stringently. The problem facing Rich and his customer in the Pentagon, the Air Force Special Projects Office, was how to go about shielding an entire industry. The U-2 and the A-12 had been built in "onesies and twosies," but the F-117A was required in multiple squadron strength. Clearly, there would come a time when the program would need to be revealed, but the longer it could be shielded from public view, the less time the Russians would have to react when it finally emerged into the light.

Rich was one of five Lockheed employees cleared for top secret work and above. Everyone else connected with the program—including several thousand factory floor workers—had to be rigorously security-checked. And nothing about their lives was off-limits. "Security's dragnet poked and prodded into every nook and cranny of our operation," Rich wrote years later. It almost drove him insane.

As the F-117A began to take shape at Burbank, arrangements were made to accommodate it operationally. A secret base was constructed in the desert at Tonopah, Nevada, and pilots were asked to volunteer without being told anything about the assignment. In New Mexico, at a radar range miles from the nearest public land, the final configuration of the F-1 17A was subjected to more RCS checks to validate its insect-like radar properties. When the aircraft made its first flight, in June 1981, it did so at Area 51, shielded from the remotest scrutiny by the jagged mountains of southern Nevada and a crack air force special forces unit authorized to use "lethal force" to protect Groom Lake from intruders.

As production ramped up under the administration of the hawkish presidential incumbent, Ronald Reagan, who took office in 1980, it rapidly became clear that the Skunk Works was outgrowing its Burbank facilities and so plans were drawn up to relocate the plant eastward, to Palmdale, on the edge of California's Mojave Desert. Palmdale was also home to a giant new production facility operated by Northrop, designer of the B-2, the USAF's four-engine strategic Stealth Bomber. The B-2 was black, but not as black as the F-1 17A. This meant it was acknowledged to exist, even though the rudiments of its design were classified.

It was in this triangle of land, bordered by New Mexico to the south, California to the west and Nevada to the northeast, that people—ordinary people—started seeing things on moonlit desert nights that they could not correlate with the known facts. Speculation grew that a "stealth fighter" program of some description existed—and almost certainly at

Lockheed, where security in recent years had tightened considerably—but the detail was invariably way off the mark. And since the money for the F-117A program—to the tune of several billion dollars—had been appropriated from the Pentagon's "black budget," ring-fenced as it was from public scrutiny, that was how things remained until late 1988, when the outgoing Reagan administration revealed just what it had hushed up for so many years.

The F-117A, the bland press release stated, had gone operational in 1983. For five years, its pilots had operated at squadron strength "in the black," roaming the desert night skies of the U.S. Southwest, practicing for the nighttime attack mission they would be required to perform in a war. The disclosure had only been made at all because the air force wanted to expand the training envelope and start flying the F-117A by day. Otherwise it would have remained buried even longer.

During the F-117A's five years of secret ops, thousands of workers had been involved in the assembly process at Burbank; hundreds more in supporting the aircraft at Tonopah. And yet, not one had breathed a word about it.

The big story in November 1988, then, related to the existence and capabilities of the Stealth Fighter, a program that rivaled the scale and daring of the Manhattan Project almost 50 years earlier. Like the bomb program, the F-117A had been meticulously covered up, but unlike the bomb program, the secrecy had held. With hindsight, this was understandable. The moment the U.S. government had decided not just to develop the bomb, but to use it on the enemy, its security was compromised. Atom spies sympathetic to the Soviet Union—the liberal, heady atmosphere of Los Alamos cultivated them like seeds in a hot-house—were always going to ensure that details of the weapon would be passed to Moscow; and the traffic merely accelerated when it was dropped on the Japanese.

The logic these people had applied in betraying their country was simple. They didn't want America to have a monopoly on such a devastatingly powerful strand of weapons technology. By passing bomb secrets to the Russians, the atom spies believed they were restoring the balance of power and making the world a safer place. It was a lesson that U.S. security managers would not forget.

Stealth was a passive technology. It did not need to be "dropped" on an unsuspecting enemy, but it had the power to begin and end wars just the same. Because it didn't need to be revealed, it had been easy to cover up, until the moment it suited the U.S. government to make the revelation—to bring it out of the black. Neglected in the process of

disclosure, however, was any analysis of the system that had allowed this superhuman feat of engineering and parallel security to materialize; and the black world was a system—a huge, sprawling machine.

Between the bomb and the Stealth Fighter, a massive security structure had swung into place to protect America's ever-proliferating black programs. And in the vastness of the desert, it wasn't hard for individuals—the little cogs of the machine—to develop a sense of their frailty. The desert had been a good place to build the bomb and the perfect place to field stealth. It had swallowed both programs—the weapons and the people who'd constructed them—whole.

If antigravity was real—and by now I believed the white world contained evidence that *something* was going on—the desert was where it would have come together.

According to Marckus, the black world had known I was coming from the moment I set foot in NASA and had had plenty of time to prepare for my visits.

In London, I struggled to make the right connections.

I returned to something Marckus had said during our meeting by the estuary; something about the Northrop B-2 Stealth Bomber. There was a strand of thought, even among some quite high-profile academics, that the B-2 used some kind of antigravity drive system. I wasn't inclined to believe the story at face value. But with scant entry points to the black world, it seemed, for the moment, as good a place as any to start. If there were lessons still to be learned from the media's coverage of the F-117A in the early 1980s—before the program had been revealed, but when people were reporting things they couldn't explain in the night skies of California, Nevada and New Mexico—the most appropriate was the old one about smoke and fire.

The detour through NASA had pointed up five possible pathways to antigravity's application in the black world: manipulating an object's mass and/or inertia; exploitation of the zero-point energy field; perturbations of the space-time continuum; faster-than-light travel; and gravity shielding. Of the five, two (space-time and faster-than-light) were patently out of the question—the B-2 was not a time machine and it could not even muster the speed of sound—and two more (mass/inertia manipulation and zero-point exploitation) were still some way from a practical breakthrough. Only one, the last of the five, seemed to have resulted in any kind of payoff. And Podkletnov had tripped over it by accident because of somebody's pipe smoke, which meant it had been out there, waiting to be discovered, for as long as anyone needed it.

Applied to the B-2, using some of the many billions that had gone into

its development, the kind of weight reductions that the Russian had been reporting (anywhere from 2 percent to 5 percent) would have been highly significant. The B-2 had already shown its ability to fly halfway around the world and back—a 37-hour, 29,000-nautical-mile return flight from Whiteman Air Force Base, Missouri, to Guam in the Pacific with several air-to-air refuelings en route. With gravity shielding—if such a thing could be moved from the laboratory to a fully fledged aircraft program—either its range could be extended or the number of midair refuelings cut. Either of these would make the technology a powerful draw.

The biggest argument against the existence of antigravity technology had seemed to me to be the lack of any visible hardware from production programs. Now, I was having to confront the possibility that antigravity played a vital part in the most important technological development since the bomb.

Chapter 12

I'd been aware of rumors about the B-2 for some time, but they were so bizarre as to be quite unbelievable, so I'd ignored them completely. In following the thread that had been dropped by Marckus at the estuary, however, I discovered that it led somewhere that was already familiar. The chief proponent of the idea that the B-2 utilized an exotic propulsion source was my old friend Paul LaViolette, the Ph.D. who had tripped over the reference to antigravity in the Library of Congress.

LaViolette had written an elegant thesis about the B-2 and its anti-gravitational properties soon after the publication of a revelatory article in *Aviation Week*, the "bible" of the U.S. aerospace industry, in which a number of black-world engineers had whistle-blown to the author, the magazine's West Coast editor, Bill Scott, about a host of amazing new technologies supposedly being developed in the deeply classified environment. One of these related to "electrostatic field-generating techniques" in the B-2's wing leading edge with a view to reducing the giant Stealth Bomber's radar cross section.

LaViolette had taken this revelation and coupled it to his own knowledge of Thomas Townsend Brown's work and patents and drawn the conclusion that the B-2 was the embodiment of Project Winterhaven, T.T. Brown's proposal in 1952 to the U.S. military for a Mach 3 saucer-shaped interceptor powered exclusively by electrogravitics—antigravity.

In an essay entitled "The U.S. Antigravity Squadron," LaViolette outlined the reasoning behind his extraordinarily bold assertion.

The B-2, he held, was simply a scaled-up version of the subscale discs, the little charged saucer-shaped condensers, that had whizzed around Brown's laboratories in front of multiple witnesses from the U.S. Air Force and Navy in the late '40s and early '50s. If true, it explained why superficially the U.S. Air Force had shown such little interest in Winterhaven, while on the inside—I remembered the recovered transcripts of conversations between Air Force generals Bertrandias and Craig following the former's visit to Brown's lab in 1952—those who had seen the tests were going apoplectic with excitement.

While conventional wisdom had it that the B-2's outer skin was composed of a highly classified radar-absorbent material (RAM) that made it invisible to radar, LaViolette plausibly argued that the RAM was in fact a ceramic dielectric material able to store high amounts of electric charge. The material was said to be made from powdered depleted uranium—an incredibly hard substance, commonly used to tip armor-piercing tank shells—with three times the density of the "high-k" ceramics proposed by Brown in the 1950s. This would give the B-2 "three times the electrogravitic pull" of the Mach 3 saucer at the heart of the Winterhaven study.

Another useful bit of synchronicity was the fact that Brown had, for a short period of time, worked for General Electric, the company responsible for the B-2's engines.

But try as I might, I couldn't get LaViolette's theory to fit. For the thesis to be accurate, it hinged on something approximating a switch in the Stealth Bomber's two-man cockpit that would permit the aircraft to transition from normal jet-powered flight—something, according to LaViolette, that it used on takeoff and landing and other occasions where witnesses might be present—to antigravity cruise mode.

If there were such a thing as this switch, I knew, pound to a penny, dollar to a cent, its existence would have leaked.

The B-2 first flew in 1989 and the first operational aircraft arrived at Whiteman Air Force Base in 1993. A total of 20 aircraft had been built at \$2 billion a shot and all of them had been delivered by the end of the decade. Dozens of crewmembers had been trained in that time, and many of them had been interviewed. The B-2 had started as a black program (or more accurately, a dark-gray one, since its existence had been confirmed officially from the start, all other details being classified). But now it was in the open, albeit, still, with some highly classified features. Pilots—an inherently talkative breed—would have let it slip somewhere along the line. Rumors would have developed about such a switch. But they never had. Moreover, B-2 pilots and engineers hadn't just denied the antigravity story, they had scoffed at it openly. Intuitively, my colleagues and I were agreed, the B-2 antigravity story was wrong. In fact, the whole notion of it was absurd.

The end result was that "the B-2 as antigravity vehicle" had been consigned to the columns of conspiracy mags and tabloids; the mainstream aerospace press, afraid of an adverse reaction from its conservative readership, had refused to touch it.

With one exception.

Enter Britain's most eminent aerospace journalist, Bill Gunston, OBE,

Fellow of the Royal Aeronautical Society, and an article he'd penned entitled "Military Power."

Gunston, who served as a pilot in the RAF from 1943 to 1948, was scrupulous over his facts—as editor of Jane's yearbook on aero-engine propulsion, he had to be. "Military Power," published in *Air International* magazine, was a walk-through dissertation on the development of aero-engine technology since the end of the Second World War; good, solid Gunston stuff.

Until the last couple of pages, when, to the uninitiated, it appeared that the aerospace doyen had lost his mind. Gunston not only portrayed the B-2 antigravity drive story as fact, but went on to reveal how he had been well acquainted with the rudiments of T.T. Brown's theories for years, but had "no wish to reside in the Tower [of London], so had refrained from discussing clever aeroplanes with leading edges charged to millions of volts positive and trailing edges to millions of volts negative."

Gunston explained why he felt that there was much more to the B-2 than met the eye, drawing on a lifetime of specialized knowledge. In short, if you applied the laws of aerodynamics and basic math to the known specifications of the B-2, there was a glaring mismatch in its published performance figures.

It was clear, Gunston said, that the thrust of each GE engine was insufficient to lift the 376,500-pound listed gross weight of the aircraft at takeoff.

The only way the B-2 could get into the air, therefore, was for it somewhere along the line to shed some of its weight. And that, of course, was impossible, unless you applied the heretical principles of Thomas Townsend Brown to the aircraft's design spec. The B-2 wasn't a black world aircraft program anymore. It was operational, had been blooded in combat and was frequently exhibited to the public. If it used an antigravity drive system, the proof of the phenomenon existed in plain sight.

If it didn't, then someone, somewhere had gone to some lengths to make a number of highly respected aerospace experts believe that it did.

In October 1990, *Aviation Week* published a report based on an analysis of 45 different daylight and nighttime eyewitness observations of strange aircraft over the desert Southwest—Bill Scott's backyard. In this, he concluded that there were "at least two—but probably more—types of vehicles" beyond the F-1 17A and B-2 under test. One was a "triangular-shaped, quiet aircraft seen with a flight of F-1 17A stealth fighters several

times since the summer of 1989"; another was a "high-speed aircraft characterized by a very deep, rumbling roar reminiscent of heavy lift rockets"; a third was a "high-altitude aircraft that crosses the night sky at extremely high speed . . . observed as a single, bright light—sometimes pulsating—flying at speeds far exceeding other aircraft in the area." Because the latter was always seen above 50,000 feet, where you wouldn't routinely expect to hear engine noise, Scott deduced that the second and third aircraft might be one and the same type.

Speculation that the Reagan administration had embarked on a massive program of black world aerospace and defense research began circulating in earnest among Washington reporters and defense journalists following the publication of a "line item" called "Aurora" in the Pentagon's fiscal year 1986 budget. Aurora was supposedly a "sleeper site" used by the Pentagon for burying B-2 funds at a time when the Stealth Bomber was still highly classified. But this explanation, put about by "highly placed Pentagon sources" when the Aurora story hit the streets, simply smacked of backpedaling. As Aurora was listed under the "Other Aircraft" category and was projected to receive a huge increase in funding between fiscal years '86 and '87—a jump from \$80.1 million to \$2.272 billion—the sleuths concluded that its inclusion in the budget request was a genuine mistake, that the aircraft was real and, because it was listed in the "reconnaissance" section, must have been a replacement for the Mach 3.2 Lockheed Blackbird, which was earmarked to retire from its strategic spying duties in the early 1990s.

Then, when that dark triangular shape was seen by a trained Royal Observer Corps spotter over the oil rig in 1989, people put two and two together and deduced that Aurora wasn't just a line item, but a real flying aircraft.

One of the supreme ironies of the black budget system is that, thanks to U.S. bureaucratic procedure, you can work out, almost to the cent, what the Pentagon is spending on its most deeply classified programs. Every year its financial planners submit an unclassified version of the defense budget to Congress. It is a remarkably detailed document, listing major line items of expenditure in the military research and development (R&D) and weapons procurement arenas. Trawl down the list, however, and some of the dollar values of these line items are missing. Many of those in this category have strange-sounding code names like Forest Green, Senior Year, Chalk Eagle or Centennial. Others have curiously nondescript headings such as "special applications program" or "special analysis activities."

These are black programs. Because the unclassified budget includes

accurate total budgets for all three armed services, all you need do is subtract the total dollar value of the line items from each category, and the difference is what the Pentagon is spending on black world research and acquisition.

In 1988, the total was computed to be \$30 billion for R&D and secret weapons programs—more than the entire annual defense budget of a major European NATO nation such as Britain, France or Germany.

What we were talking here was an industry within an industry, multiple layers deep and compartmentalized to death horizontally. The black world's inaccessibility was based not just on its segregation from the outside, but on the inability of managers and engineers in one of its compartments, or on one of its layers, to have oversight of anything that wasn't within his or her "need to know."

In a system whose existence had been designed to be denied, examples were few and far between, but one shone clear. General George Sylvester, head of the Air Force's R&D program in 1977, was not "accessed" to the Lockheed Have Blue program, even though the program itself was run out of his office at Wright-Patterson Air Force Base. One of the "bright young colonels" at Air Force Special Projects in Washington who ran the Air Force's portion of the black budget had decided to cut Sylvester out of the loop.

It was small wonder, then, that reporters and investigators began to level the charge that the black world had spiraled out of control.

How deep did the layers go? How many compartments did the system contain? What kind of technologies were being pursued within it?

The truth was, nobody knew. The black world was labyrinthine and it was essentially unaccountable; its whole system of compartmentalization so convoluted, so tortuous, that if people like Sylvester had been excluded, then God only knew how many others who should have had the "need to know," didn't.

It was in this climate of hidden expenditure that Bill Scott reported what he was seeing and hearing in and around Palmdale.

The prognosis was that Aurora was a massive leap forward in aerospace terms, that it was powered by a new form of "combined-cycle" engine fueled by liquid methane or hydrogen to give it a cruise speed anywhere up to Mach 8—5,300 mph. It was also a fair bet, Scott and others reported, that the Lockheed Skunk Works, which was in the process of shifting to Palmdale, was the firm responsible for building it.

Outwardly, the Skunk Works had next to no work on the stocks, but something was keeping its 4,000 employees busy just the same and something was rattling the windows of the desert Southwest at night.

From mid-1991 until the mid-'90s, the U.S. Geological Survey, the agency that monitors earthquake activity, revealed that an unknown aircraft, with sound footprints different from the SR-71 or the Space Shuttle, was causing "airquakes" as it crossed the California coastline from the Pacific, heading inland for southern Nevada. Couple this with some fresh U.S. eyewitness sightings after the North Sea encounter of a huge, arrow-shaped aircraft that left a weird, knotted contrail in its wake and made a noise "like the sky tearing apart," and a recent program of expansion at the Air Force's test site at Area 51 in southern Nevada, and it all added up to a black world economy that was, so to speak, booming, too; a situation that continued even after the Cold War ended.

Following the dissolution of the Soviet Union, the U.S. Air Force budget, in concert with those of the other armed services, was cut and black world funding along with it. But as a proportion of the overall budget, the classified segment of the USAF's R&D and acquisition spending actually increased and, even in the mid—late 1990s, hovered comfortably at around the \$11 billion a year mark. This left more than enough money for the Air Force to pursue a range of exotic technologies for the 21st century.

In early 1992, Scott was contacted by a group of black world engineers who had decided to break cover and talk about some of the programs they were working on. Their reasoning was simple. The Cold War was over and there was no need for the U.S. government to protect some of its most highly classified technologies any longer. Some of these technologies, the engineers told the reporter, could actually be spun into the white world of commercial aerospace programs to give the U.S. an enormous economic advantage over its international aerospace trade rivals.

In revealing their presence to Scott, they acknowledged that they were breaking a "code of silence that rivals the Mafia's." The dangers of their approach were already manifest. Two engineers said that their civil rights had been "blatantly abused" by their black world controllers in a bid to prevent them from leaving the deeply classified environment.

"Once you're in, they don't let you go," one of them said.

Amidst the revelations that the engineers dangled in front of Scott were a number of technologies for reducing the visible signatures of the B-2.

To people in the business, Northrop's winning of the \$40 billion Stealth Bomber competition against Lockheed in 1981 had long been something of a mystery. It was, after all, Lockheed that had made the breakthrough in mathematically computing stealth in 1975. But then, the

F-1 17A, composed of its mass of angled flat panels, was "first-generation stealth."

The B-2, or the Advanced Strategic Penetrating Aircraft, as it was known when it was still deeply gray, was a second-generation aircraft and this, according to Northrop, was manifest in its blended aerodynamic shape. Instead of being rough and angular like the F-1 17A, it was smooth and rounded—a product, Northrop claimed, of five years' worth of improvements in computing technology between the start of the Have Blue program and the ASPA.

Improved software algorithms processing had allowed Northrop to come up with a shape—a refinement of the YB-49 flying wing jet bomber it had first flown in 1947—that allowed the aircraft to look and fly right *and* to remain invisible on enemy radar screens.

Not only that, but according to LaViolette it had the capability to fly electrogravitically as well.

In my basement, where the noise of the city couldn't intrude upon the sort of silence you needed to sift a decade's worth of documents and files on the arcane properties of stealth, I still wasn't happy about LaViolette's assertion, because of the absence of that all-important switch, the one that kicked in the B-2's antigravity drive system.

There was one other beguiling aspect to Scott's story. His sources said that charging the airframe electrostatically not only helped to make the B-2 stealthy, it also reduced friction heating of the airframe, its sonic footprint and its drag.

This last point was crucial. If it were possible to alter the "drag" or air resistance of an aircraft, you could either make it fly farther or faster or both.

The drag reduction system was supposed to work as follows: By creating an electrostatic field ahead of an aircraft, it ought in theory to repel air molecules in the aircraft's path, allowing the plane, in effect, to slip through the atmosphere like a thin sliver of soap through warm water.

Like antigravity itself, the whole notion of using electricity fields to reduce aircraft drag was heresy—it was, according to traditionalists, just smoke and mirrors. But here, in Scott's article, was a reference to Northrop having conducted high-grade research in 1968 on just such a phenomenon: the creation of "electrical forces to condition the air flowing around an aircraft at supersonic speeds." Furthermore, the California-based aircraft company had supposedly submitted a paper on the subject to the American Institute of Aeronautics and Astronautics that same year.

I hit the Net and went into the AIAA website, then did a title search. But the papers on the AIAA site went back only as far as 1992 and I was out of time.

I rubbed my eyes. In two days I was due to attend an aerospace convention in Las Vegas. And I wasn't any closer to solving the puzzle.

Or was I?

During the Second World War, I remembered Thomas Townsend Brown had been involved in experiments that sought to show how you could make a ship disappear on a radar screen by pumping it with large doses of electricity. Between 1941 and 1943, Brown had supposedly been involved in tests, I saw now, *that were identical in principle to the methodology that Northrop seemed to have applied to the B-2 to make it the ultimate word in stealth.* Researchers had never taken Brown's wartime experiments seriously because the precise nature of the work had been obscured by the myth of parallel dimensions aboard the USS *Eldridge*—the ship at the heart of the "Philadelphia Experiment." Brown had also used electrostatics to power his model discs and, by his own testimony and that of witnesses—including a general from the U.S. Air Force—they had flown by defying gravity.

Could it really be that in researching the B-2 I had picked up the threads of T. T. Brown's work again? Work that the U.S. had continued to develop in the black?


I jotted the keywords on a piece of paper, like a crossword player consigning letters to a page in the hope of unraveling an anagram. *Brown—electrostatics—stealth—antigravity—and then, against my better judgment—parallel dimension.*

I studied the words for a long time, but nothing jumped out at me.

Below: Alleged photograph of Rudolph Schriever.




Right: The Legend's rendition of the Schriever/Habermohl/Miethe disc.



Right: Richard Miethe.

Below: Artist's impression of foo-fighters engaging a formation of B-17 bombers. (All images via Bill Rose)





Aug. 16, 1960


T. T. BROWN

2,949,550

Filed July 3, 1957

ELECTROKINETIC APPARATUS

2 Sheets-Sheet 2




Above: Thomas Townsend Brown.

Right: Patent illustration showing experimental layout of the charged condenser plates Brown demonstrated to the US Air Force and Navy.

FIG. 7



FIG. 8



INVENTOR
THOMAS TOWNSEND BROWN
BY
Hanson, C. & Associates, Inc.
ATTORNEYS

THOMAS TOWNSEND BROWN: BAHNSON LAB 1958-1960

ELECTROGRAVITICS LEVITATION EXPERIMENTS
ORIGINAL SILENT FOOTAGE TRANSFERRED TO VHS VIDEO



/Above: Brown holding a disc capacitor prior to charging.


See New Styles of Electrogravitic Prototypes Designed for Vertical Flight and Hovering, Captured on Film at the Bahnson Laboratory

Left: A demonstration of electrogravitic lift at the Bahnson Laboratories in the late 1950s.
(All images via the Integrity Research Institute)




Left: John Frost.

Below: A cutaway illustration of the Avro Silverbug showing the radial flow gas-turbine.



Below: John Frost (right) with other members of the Avro SPG team in front of the Avrocar. (All images via Bill Rose)







Above: A full-scale engineering mock-up of the Project Y Manta at Avro's factory in Malton, Ontario.


Right: General arrangement of the proposed Avro Project 1794, showing arrangement for its six Viper turbojets.

Below: Avro's test-rig for the Mach 4 Project 1794 saucer.
(All images via Bill Rose)




SECTION A-A





Above: British Aerospace's concept of an advanced fighter interceptor, powered by an anti-gravity lift engine.

Below: Another British Aerospace design for a heavy-lift anti-gravity vehicle. (Both images via BAE Systems)






Above left: Tom Valone of the Integrity Research Institute. (Tom Valone)


Above right Paul LaViolette. (Paul LaViolette Ph.D.)

Below: Dr Evgeny Podkletnov (left) with Dr Ron Evans, Head of BAE Systems' Project Greenglow. (BAE Systems)






Above: The Northrop Grumman B-2 Stealth Bomber. (Northrop Grumman)




Left and below: LaViolette's schematic illustrations of the B-2's electrogravitic drive system, showing positive and negative charge and flow distribution. (Electrogravitic Systems)




Left: The Lockheed Martin F-117A Stealth Fighter over typical desert terrain.

Below: 50 years of Skunk Works Products - those that the company has owned up to.



Right: Ben Rich and the F-117A.
(All images via Lockheed Martin)






Above: Map showing key California-based sites of the Skunk Works in the late 80s/early 90s. (Lockheed Martin Skunk Works)

Right: Jack Gordon, head of the Skunk Works in 1996. (Lockheed Martin Skunk Works)

Below: An artist's rendering of Aurora, the USAF's mythical hypersonic spyplane. (Julian Cook)





Right: The concrete henge structure at the Wenceslas Mine - test-rig or cooling-tower? (M. Banas)

Below: The Wenceslas Mine, Ludwigsdorf, Poland, looking across what was once a covered railway marshalling yard. (Nick Cook)



Above: The Wenceslas Mine. The henge-like structure is situated behind the trees in the distance. (Nick Cook)

Right: The entrance to the Mittelwerk at Nordhausen. (I. Witkowski)




Below: SS General Hans Kammler - the only known photograph of him in uniform; France, 1944.




Above: A partially lined gallery at the SS-run 'Giant' underground manufacturing complex near Waldenburg, Poland. (M. Banas)




Left: A completed gallery at the 'Giant' underground complex. (M. Banas)



Above: Viktor Schauberger and his implosion-based home power generator in 1955, with interior shot.




Left and below: The Schauberger Repulsine with and without cone-shaped intake. (All images via Schauberger Archive)









Left: Bob Widmer, chief designer of General Dynamics Convair, in his office in the late 1950s. (Lockheed Martin)

Below: A grainy shot of a wind-tunnel model of Kingfish, Widmer's Mach 6 spyplane proposal. (Air International)



Right Bob Widmer in his Lockheed Martin office at Fort Worth, in 1996. (Nick Cook)





John Hutchison in his Vancouver apartment, surrounded by equipment integral to the Hutchison Effect. (John Hutchison)



Metal ingots 'disrupted' by the Hutchison Effect. (John Hutchison)

Chapter 13

Vegas. Two thumps through the 747 and I was on the ground at McCarran International, an airport, like Miami's, that crackled with illicit activity; you could just feel it. We taxied to the terminal and passed engineering shops filled with propeller planes that looked like they should have been scrapped decades ago; old classics running freight operations now, with radial engines that dripped oil on the ramp and coughed clouds of smoke, their tired but sturdy airframes lined with irregular rows of rivets and bumps.

I raised my eyes and took in the view. Vegas, sprawled there just beyond the perimeter fence under an upturned bowl of blue sky, half a dozen giant new hotels since I'd been here last, half a dozen more going up amidst a thicket of cranes.

I loved and loathed this city with equal measure.

Down on the Strip, with the sun setting on this city of broken dreams, I took in the glare of the lights from the back of the cab, half listening to the driver's lament over the mob's exodus and its replacement by corporate America.

The slipping sun played over the tops of the mountains that bordered the southern end of the Nellis Air Force Range, a closed-off piece of government land the size of Switzerland where the Air Force tested its latest military hardware and ran air warfare exercises, and the driver's voice still filling my head about the days when people in the city had lived and died by a set of rules everyone understood.

I grunted something neutral as I tracked the Convention Center where tomorrow I'd tramp the hallways, doing what I always did on these occasions: talk to old friends, meet contacts, trawl the stands, hoover up data, collate stories, file copy, hit the town, go to bed late, get up early and then go through it all over again.

The cab pulled into the Sahara, a hotel that exuded the Vegas of Sinatra, Martin and the mob; the mid-'50s, when it had opened.

It had been face-lifted so many times, it seemed like the demolition teams would be here at any moment, but through half-closed eyes you

could still see how it once had been, which gave it a strange kind of charm. The Sahara suited me just fine.

Its pile-'em-in, rack-'em and stack-'em ethos, ideal for gamblers passing through, meant it was shunned by most of the show delegates, but it had the enormous advantage of being a stone's throw from the Convention Center.

The following day, I picked up my reporter's pass and filed into the exhibition hall. It was jam-packed with aerospace and defense companies from across America, their stands and booths ranged in rows either side of the carpeted walkways.

From all over the hall, TVs plugged into videos on permanent loop flashed with images of combat aircraft going through their paces or weapons under test, the pictures backed by deadpan monologues and *Top Gun*-style soundtracks.

I slipped into the routine, walking the course, checking the stands for signs of developments or breakthroughs.

Sometimes stories came from a new aircraft, spacecraft or weapon system depicted in model form or artist's impression, but such copy as this generated was merely the low-hanging fruit—things that the public affairs people wanted you to see. As long as it constituted news, most reporters, myself included, were happy to go through the motions.

More often than not, though, the developments we looked for were more subtle: a missile fitted with a new seeker or an aircraft with a new sensor, denoted by an aerial or antenna window on the display model that hadn't been there last time around.

Such changes, though seemingly inconsequential to an outsider, pointed to performance gains that signaled real news to a defense reporter, news that the industry's primary customers generally didn't want to see in print.

But there was another side to show reporting, a side that I altogether preferred: chewing the fat with people in the business I'd known and liked for years, gathering data that gave hints of longer-term trends; what was really happening to the industry in the face of Cold War budget cuts, where the technology was headed, who was buying, who wasn't.

Three days at an aerospace and defense exhibition gave you information you couldn't get from a month's worth of phone calls. And after the intensity of weeks spent scrutinizing obscure pieces of data for hints of a black world antigravity program, the change was refreshingly good.

On the third day, long after the press conferences had dried up and I'd filed my final piece of copy to the magazine, I hit the Northrop Grumman

stand and started talking to a guy who'd worked a whole range of the company's programs, including the B-2 Stealth Bomber.

Careful not to provoke the reaction that usually greeted the antigravity question, I asked him what he made of the stones that the B-2 charged its surface electrostatically.

This time, no derogatory smirk, just a straight answer and a half smile.

It was impossible, he told me. If you charged the airframe of the B-2, you'd fry its onboard electronics. He went on to describe a secondary problem. All aircraft need to discharge the static electricity they absorb naturally as they move through the air—usually via small wire aerials on the wing tips. Deliberately adding to this static buildup with a man-made electrical charge would be enormously risky, the engineer maintained. Since it would be impossible to discharge the electricity fast enough, it would turn the aircraft into a giant lightning conductor.

I called Marckus from my hotel room the following morning. The eight-hour time difference meant that he was at home, the day's work finished. Classical music filled the background, Brahms' Symphony Number Four turned deafeningly high, giving me, in a moment of insight, a clear view of Marckus' domestic situation. I knew then the man lived alone.

"He said what?" he shouted back.

I repeated the engineer's assertion. Marckus left the phone for a few seconds to turn the music down.

"The man's an idiot," he said when he came back. "Either that or he deliberately lied to you."

"Why should he do that?"

"Why do people normally lie?"

"Come on, Dan, it's too damned early here for twenty questions."

"The guts of the B-2 are held within a Faraday Cage," Marckus explained. "It doesn't matter how much electricity you pump across the skin. The inside remains insulated from it. The guy was bullshitting you. And it's not even very good bullshit."

The rest of his argument was badly flawed also, Marckus said. If you looked at the B-2 from the top, it looked like a boomerang, with straight, sharp leading edges and a serrated trailing edge. Regular aircraft had two wing tips; two points from which to discharge their natural buildup of static. But the B-2 was a flying wing. With its sawtooth back end, it had, in effect, seven wing tips from which to discharge the continuous flow of electricity from its supposed flame-jet generators.

At low altitude and low speed, the discharges would be difficult, if not

impossible to see with the naked eye. But at cruise altitude, with enough plasma buildup around its leading edges, an aircraft—particularly a supersonic or a hypersonic platform—would glow like a lightbulb. It would pulse.

"A high-altitude aircraft that crosses the night sky at extremely high speed . . . observed as a single, bright light—sometimes pulsating—flying at speeds far exceeding other aircraft in the area," one of the nighttime eyewitness observations reported in the October 1990 edition of *Aviation Week*.

"Northrop has been playing around with electrostatics for years," Marckus said. "In 1968, it submitted a paper called 'Electroaerodynamics in Supersonic Flow' to the American Institute of Aeronautics and Astronautics."

"I know," I told him, happy for once I had something he didn't. "I tried to find it on the AIAA website, but the papers on their electronic database don't go back far enough. I'll have to request a copy from their archives instead."

Even across eight time zones, I caught Marckus' withering sigh.

"Don't bother. I applied eight years ago. It's missing. They couldn't find any record of it. Almost certainly, some special projects office in the Pentagon pulled it. Made it disappear. That happens occasionally. Someone screws up and the black world has to go clean up. Actually, it wasn't all Northrop's fault. Back in the '60s, they had some noble ideas that this technology could be applied as a drag-reduction device for supersonic airliners, cutting heat friction and fuel burn. It was only when someone clocked the fact that a plasma shield around an aircraft also reduces its radar signature that the military really woke up to it. It's been rumored for a while that Northrop had come up with something revolutionary in this field. The missing paper proves it."

Marckus carried on. I closed my eyes and tried to think.

The signal I was picking up was painfully indistinct against the background mush of interference, but I could see that it added up to something important.

The B-2 clearly used some kind of skin-charging mechanism to lower its radar (and quite possibly its heat) signature. But it was also clear, from the nature of the paper that Northrop had written in 1968, that the origins of this electric shield, this stealth cloaking device, were actually rooted in another field of research altogether—drag reduction. According to Marckus, who now admitted to having witnessed similar experiments in the U.K., the electrostatic field could also be made to behave like a big "jet flap," an extension of the wing. Deploy this invisible

jet flap on takeoff and it was the same as if the wing area had been substantially increased, with a parallel reduction in drag and a concomitant improvement in lift—by perhaps as much as 30 percent.

Bingo.

This could explain how the B-2, or an aircraft like it, staggered into the air with a full bomb-and-fuel load. Thirty percent seemed far too high, but even a few percentage points would constitute a breakthrough. A substantial lift improvement was the equivalent of a substantial reduction in weight. Maybe the B-2 lost weight, so to speak, aerodynamically, not electrogravitically. It would explain why Northrop had beaten Lockheed to the B-2 contract. While the Skunk Works had been playing around with Maxwell's equations and two-dimensional shapes, Northrop had already worked out how an aircraft could be enveloped in a shield of static electricity and how that energy could be put to work: as both a cloaking device and a method for reducing its air resistance.

No laws of physics broken, no antigravity switch in the cockpit, but an extraordinary development nonetheless.

A weight reduction device by proxy.

And in working this through, I realized something else. The B-2 antigravity story actually played into the hands of black world security. It was so fundamentally unbelievable that it discredited any serious investigation of the application of electrostatics to aircraft. What defeated me was why the spooks should have worked so hard to stop its disclosure—pulling papers from the AIAA and so forth—and yet be content for details to be released of the Lockheed stealth method.

But then, disinformation works best when you mix it in with a little truth.

It was then that the picture cleared momentarily and I glimpsed something through the snow and distortion. The keywords I'd written down and then hurled into the bin in the basement of my home a few days earlier swam into focus. If I could just hold the picture a second or two longer . . .

Markus' voice, punchy and impatient on the other end of the line, brought me back to the here and now of my room in the Sahara.

I'm still here, I told him.

He was asking me where I was headed next.

I told him there was only one possible place I could go. If Northrop had been working on electrostatics since the 1960s, *if* it was now operationally deployed on a \$2 billion frontline bomber that had been designed in 1980, there had been plenty of time in the intervening period for improvements and refinements. I knew there was no way that

Lockheed, now Lockheed Martin, would have taken Northrop's B-2 victory standing still.

Aviation Week had written about the "fast mover" that had been observed crossing the night sky as a single, pulsating point of light. And others had reported the manifest contradiction of a prominent Air Force contractor at Palmdale that was filled to capacity with workers, was hugely profitable, yet had nothing to show for it.

I told Marckus that my request to interview the head of the Skunk Works, Jack Gordon, had just been given the green light.

Better still, it was to take place within the holy of holies itself.

It was only on the road to Lancaster, my next stop, across the border in California's Antelope Valley and a three-hour drive across the Mojave Desert, that I found the quiet I needed to work through what I'd been unable to finish in my call to Marckus.

Its starting point was one of the keywords that I'd written down on the note I'd penned and then trashed five days earlier: stealth. It all started and ended with stealth. In 1943, Brown had been involved in an experiment that sought to make a ship radar-invisible. His test was the first attempt by anyone—certainly, as far as I knew—to have conducted a genuine radar-stealth experiment.

At its heart was a technique developed by Brown to wrap a ship in a field of intense electromagnetic energy, produced by powerful electrical generators.

In the mid-1970s, the U.S. Department of Defense was briefed by Lockheed on a new form of technology—essentially, to do with airframe shaping—that reduced the radar cross section of a fighter-sized aircraft to that of a large insect.

In 1976, the company was awarded a contract to develop the Have Blue stealth demonstrator. Two years later, Lockheed was given the go-ahead to develop the F-117A, the world's first "stealth fighter."

A new military science had been born—one that was about to change the defense planning of entire nations.

As a result, the technology was classified to the rooftops. Stealth had become America's biggest military secret since the bomb.

It was then that somebody must have remembered Northrop's research papers into drag-reducing plasmas and dusted them off, recalling, probably, that electrostatics had a highly beneficial spin-off effect: it could reduce—perhaps even cancel altogether—the radar signature of an aircraft.

A picture formed in my mind's eye of an insect-sized image on a radar

screen winking out altogether. The B-2, three times the size of the F-1 17, would be 100 percent radar-invisible if it worked as advertised with its plasma cloak activated.

Since the development of big-budget weapon systems is all about risk reduction, the Northrop approach, fundamentally different from Lockheed's, would have appealed to the bright young colonels of Air Force Special Projects.

If one of the methods failed to deliver, you could always rely on the other one to show you the money. It gave the colonels two bites at the problem.

Soon afterward, Northrop won the contract to build the B-2A Stealth Bomber, based on its use of electrostatics to make it ultra-stealthy. The technique, first developed by Brown, also improved its aerodynamic efficiency and reduced its drag. It represented a phenomenal breakthrough—a development that would give the U.S. a ten-year lead, maybe more, over the Soviet Union.

But herein lay a problem.

The whole science of stealth would need to be protected like no other strand of weapons science before—the lessons of the bomb being uppermost in the minds of those charged with securing the Air Force's precious new secret.

A novel protection strategy, one that overcame the flawed security model established to protect the bomb 40 years earlier, needed to be set in place.

Eradication and disinformation, coupled with the secure compartmentalisation of the program, formed the basis of that strategy.

From that moment on, past clues or pointers to stealth would have slowly and carefully been excised from officialdom. It was most likely at or around this point that the Northrop paper was pulled from the archives of the AIAA.

But then, in the archive trawl, somebody would have come across the work of T. T. Brown. Brown's stealth work for the U.S. Navy in the early 1940s, coupled with his use of electrostatics in his electrogravitic work, was an unexpected discovery that needed to be handled with great care. The man's work was a dangerous liability. Something needed to be done—something truly innovative.

In 1978, the authors of the Philadelphia Experiment, Berlitz and Moore, were apparently approached by two U.S. Air Force officers who claimed to have knowledge of a fantastic secret. They told how they had had a chance encounter with a man who'd claimed to have been on the *\JSSEldndgeinl943*.

The man had told the two officers how the warship had vanished from its moorings during a test that sought to make it invisible to radar.

In 1979, after further research by Berlitz and Moore, much of it supplied by their elusive and slippery source Carlos Allende, aka Carl Allen, *The Philadelphia Experiment* appeared in print—and Brown's name became forever linked with the myth of the *Eldridge*,

The story's primary twist was that the ship had been transported into another dimension—kooky stuff and, to any sensibly minded individual, wholly unbelievable.

In the 1980s, the decade when stealth came to be fielded operationally, the technology had acquired its own tailor-made cloak of disinformation.

Tempting as it was to assume that Winterhaven or something like it had been pursued in the black since the 1950s, there was simply no evidence for it.

There was no doubt in my mind that LaViolette's articles on the B-2 were sincere, but they had only helped to muddy the Brown story further. When the B-2 and its antigravity propulsion system appeared in print, Brown—whose work was at the heart of this technology—was already a highly discredited figure. Whichever way you approached the story of his life and work, sooner or later you hit the Philadelphia myth. It blew all hope of dispassionate reporting on the man out of the water.

It turned out that the removal of the AIAA paper wasn't the only example of censorship in this field. Before leaving Vegas, I'd called Tom Valone of the Integrity Research Institute. Valone told me that he had been trying for years to locate a copy of an appraisal of T.T. Brown's experiments conducted in 1952 by the Naval Research Lab in Washington, D.C., but had been informed that the NRL no longer had it. *The T. T. Brown Electrogravity Device: A Comprehensive Evaluation by the Office of Naval Research* had disappeared into the ether as well.

To deduce from this that Brown had cracked gravity was too big a leap for me. In his experiments, he had demonstrated to the satisfaction of many that an antigravity effect was possible. But that didn't mean he'd got to the point where the technology could power an aircraft. Project Winterhaven had promised to deliver this, but Winterhaven never got off the ground. Not in the white world, anyhow.

Somewhere between the junction at Barstow and Boron, a tiny mining community on the edge of the Mojave, the sun finally slipped behind the southern tip of the Sierra Nevadas and I knew I'd taken the disinformation angle as far as it would go. There were just two things that wouldn't fall into place. One of them related to the words on that self-

penned note. Brown, electrostatics, stealth and antigravity: all of these elements were bound together in the disinformation story—those who had assembled it operating on the age-old principle that a little truth really does go a long, long way.

Why, then, did they introduce the fifth element? The part about the *Eldridge* slipping into a parallel dimension? How had such an improbable notion popped into the heads of the disinformation gurus?

Somehow, I had the feeling Marckus knew about all of this, yet hadn't passed the knowledge on. As I hit the freeway and turned west toward Lancaster, night falling as fast as rain, it was this thought more than any other that troubled me most.

The Desert Inn motel, its lobby replete with signed photographs of test pilots, display-case models of their aircraft and other Right Stuff memorabilia, was located on an invisible confluence of routes connecting the various nodes of the Air Force's classified desert kingdom. Twenty-five miles to the north lay Edwards Air Force base, home of the USAF's premier test pilots from the days when Chuck Yeager had broken the sound barrier in the Bell X-1. Proof of a link between Edwards and Area 51, the Nevada test base that didn't officially exist, 250 miles away as the crow flew, had been ascertained in a neat piece of detective work by Glenn Campbell, the leader of an outfit called the Desert Rats, a group of dedicated "stealth watchers" who hung out in the scrubland around Area 51, hoping for a glimpse of some exotic piece of classified hardware. Campbell had come by a procedures manual for security personnel assigned to Area 51 in which they were advised to tell anyone who professed any interest in them that they were attached to a facility called "Pittman Station." Going through local county records, Campbell found papers attesting to the fact that these personnel had been sworn in as local deputies. One of them had listed his mailing address as "Pittman Station, Henderson 89044," but all of them said that they were affiliated to something called "DET AFFTC." Campbell knew that AFFTC stood for the Air Force Flight Test Center at Edwards.

A powerful Web search engine did the rest. It turned up a 1990 NASA résumé for an Air Force astronaut candidate called Captain Carl Walz who was employed at "Pittman Station, Nevada," which in reality, Campbell knew, was nothing more than an abandoned electricity substation on the Boulder Highway. Another biography, found by a Net sleuth colleague of his, ascertained that Walz had served as a flight-test program manager at "Detachment 3 of the Air Force Flight Test Center at Edwards." Because there was no Det. 3 at Edwards itself, Campbell

deduced that it had to be Area 51 and that Pittman Station was the mail drop.

Edwards, then, was the white world mirror image of Area 51, the super-classified facility at Groom Lake.

Both the "pulser" and the aircraft tracked by the U.S. Government Survey's seismic equipment (more than likely one and the same aircraft type) were viewed on a heading that either led into or out of Groom Lake.

Twenty miles to the west, somewhere behind the palms surrounding the Desert Inn's swimming pool, lay the "Tejon Ranch," Northrop's RCS range, a remote facility buried in a gulley on the southern end of the Tehachapi Mountains. People who found themselves in the vicinity of this place in the dead of night had reported seeing orbs of light within the confines of the range and, inevitably, stories that Northrop was testing recovered alien spacecraft began to circulate among UFO buffs. Based on Northrop's highly classified work applying electrostatics to the B-2, these pieces of eyewitness testimony now perhaps had a more plausible explanation.

UFO stories had grown up around the two other radar test ranges in the area—Lockheed Martin's and McDonnell Douglas' (now Boeing's)—a score of miles to the east. Closer still, was Plant 42, the site on the airfield at Palmdale where the Skunk Works' huge manufacturing facility was located. It was there that I was due to meet with and interview Jack Gordon, bright and early after breakfast.

It was sometime in the dead hours that I heard the sound of scratching coming from the lock. I turned toward the handle, no more than four feet from the head of the bed, and saw it move, fractionally, but perceptibly, in the thin sliver of light spilling from a crack in the bathroom door. In my head, I knew exactly what was happening even before the action started. Someone was about to hard-key the front door lock and there was nothing I could do about it. As I braced for the bang from the hammer that would drive the platinum tool into the barrel of the lock and core it out in a split instant, it happened, but so fast, all I could do was lie there, paralyzed, as the door flew in and the bolt cutters took down the chain. Then, I saw them: three shapes silhouetted momentarily in the doorway—black jumpsuits, body armor, kevlar helmets and semiautomatic weapons. A glint of reflected moonlight on the barrel of the gun as it moved and then it was hard against my forehead, grinding against the thin skin between my eyebrows. More pressure, killing pressure, as the snub-barreled M16 was pushed down

harder and harder, the force driving my head deep into the mattress. And above it all, the double click of the bolt being cocked and locked, a shock of white teeth in a face otherwise hidden by the shadows as he braced to take the shot.

It was then that I came up for air, sucking it down in great gulps, my hands outstretched, pushing against bodies I could still see but weren't there and never had been.

Slowly, the ropey, '70s charm of my motel room at the Desert Inn pulled into focus. I was bathed in sweat and my hands were shaking. I lay there motionless until the sweat dried on my skin and then I tried to get back to sleep, but against the rattle of the air-conditioning unit, I knew I was righting a losing battle. Around four-thirty, I gave up, got out of bed and stepped outside, a bottle of Coke in hand, into the enveloping warmth of the tinder-dry Mojave night air.

The liquid shimmer of the night sky had been diluted by the first traces of dawn light, dimming the intensity of the stars a little, but not the overall effect. I lay back on the hood of my car and tried to lose myself in thoughts of the cosmos and Garry Lyles' dreams of a "50-year fast trip" to Alpha Centauri in a craft whose modus operandi was beyond the comprehension of the best brain talent in NASA.

After six months of sniffing around the black world, even longer reporting on its periphery, I had found nothing within the Air Force or its attendant industry to suggest the remotest link with any of the five methodologies espoused by NASA for cracking the gravity code. All I had found was a possible link to T. T. Brown and, significant though this was, it left too many gaps in the overall picture.

While Brown's early work might conceivably have explained how an aircraft with the performance characteristics of a UFO had come to be "within the present U.S. knowledge" in 1947—the footnote to the memo I'd found between USAAF generals Twining and Schulgen (now stained with the coffee I'd spilled across it)—it begged other questions: why NASA had studiously chosen to ignore his work, for example, as a pathway to antigravity.

The official reason, according to Valone, LaViolette and others, was that the Air Force had carried out its own investigation into the Brown effect as recently as 1990 and had found no cause to believe it was real. In a report on "Twenty-first Century Propulsion Concepts," prepared for the Air Force Systems Command Propulsion Directorate, the author, Robert Talley, had concluded from his experiments that "no detectable propulsive force was electrostatically induced." This failure to replicate

Brown's work was cited by both NASA officials and Dr. Hal Puthoff as a valid reason for excluding the Brown effect in their search for exotic propulsion technologies.

But what the Air Force report did not make clear was that Brown had consistently advocated using very high voltage to induce an electro-gravitic effect—50 kilovolts in the case of the 2-foot diameter discs and 150 kV in the follow-up experiment that had supposedly exhibited results "so impressive as to be highly classified." In Winterhaven, Brown had urged further tests using a 10-foot diameter disc charged at 500 kV—the Air Force, by contrast, had not raised the voltage in its tests above 19 kV.

During two out of dozens of readings, it conceded that it had recorded some "anomalous" results, but it then made no attempt to explain these anomalies in any subsequent analysis.

I tried to figure why this should be.

Was the Air Force afraid of the truth—or merely trying to suppress it?

In the clarity of thought that can come in the aftermath of dreams, the other notion I latched onto was this: Ever since I'd stepped into the black world I'd not been moving forward, as I'd hoped, but backward; to a period of research I'd already covered, rooted in work that had gone on a half century earlier.

I was trying to find answers in high-tech hot spots like the Skunk Works, when something told me I was looking in the wrong place; the wrong time, even.

And Marckus, I felt sure of it, had known about this from the beginning. He'd been in possession of the facts surrounding the B-2 and for all I knew he was bang up to date on NASA's advanced propulsion activities as well. For the past God knows how long all I'd done was duplicate his vast knowledge in a couple of key areas.

I'd been set some kind of practical exam and gone along with it.

After breakfast, I set off for my appointment with Gordon, keeping the railway on my left and the sun behind me. In the distance, through the early morning haze, a giant hangar pulled into view. It had been built in the '70s as a final assembly hall for Lockheed L1011 airliners, but subsumed since the early '90s by the Skunk Works for the only official work on its stocks: a bunch of "onesie-twosie" upgrade programs associated with the U-2 and F-117A.

The hangar dominated the desert scrubland for miles.

I pulled into the parking lot across the road from the main facility and was met by Ron Lindeke, a seasoned veteran of Lockheed's PR machine.

Since the interview was to take place on the main site itself—a rare sanction, Lindeke reminded me—I was reminded to leave my camera and tape recorder in the PR building. This functioned as an air lock between the outside world and the organization's highly secure operations on the other side of the road.

We drove in his car to the main gate, where our documentation was double-checked and triple-checked. It took an age before we were finally admitted to aerospace's inner sanctum.

Gordon's office was located in an administration block in the lee of the giant hangar. There was a low-slung building to the left of the gate and a cluster of busy-looking facilities to the right. These were the factories where the bulk of the company's classified work went on.

The giant hangar was what you were meant to see. The real work went on in the buildings in its immense shadow.

I followed Lindeke to Gordon's office. The cool, low-lit atmosphere chilled the sweat on my back and I wondered what it must be like to work in this windowless environment all the time, day in, day out; and whether it was true, as the engineer had told Bill Scott, that they really didn't let you go once you were inducted.

We stepped into an office, as dimly lit as the corridor I had just left. A tall, powerfully built man got up from a table on the edge of the room. Jack Gordon, the fourth president of the Skunk Works, smiled hesitantly and shook my hand.

The Skunk Works and its dynastic leadership system was founded in 1943 by Kelly Johnson, a hugely talented, opinionated and irascible aerospace engineer, in response to an urgent U.S. Army Air Forces requirement for a jet fighter to counter the emergent threat of the Messerschmitt 262. The Me 262, a twin-jet swept-wing fighter-bomber of revolutionary performance, had recently been identified by Allied intelligence as being under development for the Luftwaffe.

Johnson had long been badgering the Lockheed leadership for a special engineering department—small, agile and free of the overweening overheads and administrative baggage that encumbered technical progress in every large aerospace corporation (including Lockheed); and with the XP-80 project he finally got it. The XP-80 was built in just 143 days and, despite arriving too late for a showdown against the 262, went on to become an aviation classic.

The unorthodox methods Johnson pioneered to bring the XP-80 in on time and under budget—all the while under conditions of intense secrecy—set the standard for everything that followed. Ten years later, when the CIA drafted a requirement for a spyplane that could overfly the

Soviet Union, it was Lockheed that came up with the solution. When the Agency needed an aircraft to surpass the performance of that aircraft, the glider-like U-2, again it was Lockheed that furnished the answer, this time with the extraordinarily graceful Mach 3 A-12 Blackbird.

Like the U-2, the A-12 started as a black program, but gradually moved into the light, emerging finally as the USAF SR-71, the fastest operational aircraft in the world until its retirement in 1990. To this day, there is nothing faster, but that, of course, presupposes the "Aurora" stories aren't true.

Ben Rich, who succeeded Johnson in 1975, denied in his book *Skunk Works*, which serves as its semiofficial history, that Aurora ever existed. But Rich did not offer any plausible explanation for what his company had been up to since it produced the last F-117A Stealth Fighter in 1990. There had been rollouts of two YF-22 prototypes—the forerunner of the USAF's new F-22A Raptor advanced tactical fighter—and a couple of unmanned DarkStar reconnaissance vehicles, but neither of these could account for the comings and goings of 4,000 workers or the busy appearance of the low buildings I had passed on my way in.

I was questioning Gordon about the Skunk Works' operating charter when he let slip that in the course of his 23 years with the company he had worked on 15 "real flying aircraft"—adding hastily that he could only talk about 12 of them.

It was in the ensuing seconds that the thought formed and rapidly solidified in my sleep-starved brain.

Whatever was going on here, however many secret programs the Skunk Works had on its books, none of them had anything to do with antigravity.

Perhaps Aurora was real. Maybe it was the "fast mover" that people had described crossing the high frontier above the Southwestern U.S. as a pulsating point of light—as reported in *Aviation Week* in the late 1980s. Maybe it employed a drag-reduction device of even greater sophistication than the one apparently on the B-2. But whatever it was or wasn't, Aurora was an aircraft, designed and developed by aerospace engineers.

Forty years earlier, T. T. Brown had proposed specifically that Project Winterhaven should be built outside the aerospace industry. Maybe Brown knew intuitively what it had taken me months to grasp. The aerospace industry had more to lose from antigravity than it could ever gain from it. Gordon's entire business base relied on the science of aerodynamics. Take that away and he, it—everything about the aerospace industry, in fact—ceased to have any role or purpose.

Antigravity threatened to wipe the aerospace industry off the map.

Was this the reason for the crushing silence that had followed the industry's initial burst of enthusiasm for antigravity in 1956? If so, it was the simplest explanation of all. And I had spent months on the wrong heading.

I pressed Gordon on the missing three aircraft, but in view of the direction my thoughts had taken, I must have unintentionally signaled my indifference.

When he raised his eyes to mine, they told me that the meeting was over.

Minutes later, Lindeke escorted me back toward the high heat that was baking the floor of the Antelope Valley.

Just before I left the building, I stopped in front of a large chart on the wall of the lobby area. I hadn't noticed it on the way in.

I proudly illustrated the lineage of every Skunk Works aircraft since the XP-80. Past the picture of the U-2, past the SR-71 Blackbird and the F-117A Stealth Fighter, past the YF-22 and DarkStar, and there was something called "Astra."

Sitting at the top of the tree, Astra looked like an ultra-high-speed reconnaissance aircraft; every pundit's dream of how Aurora ought to look.

It was difficult to gauge Lindeke's reaction because he did not remove his sunglasses, but he knew as well as I that the Skunk Works had built nothing since DarkStar; and DarkStar, being unmanned, looked nothing like "Astra."

In the car, I pressed him for an answer and he promised to come back to me. It was several weeks later, long after I returned to England, that he finally did.

"Astra was a high-speed airliner we studied in the 1970s that got canned," he announced without any preamble. "I checked."

It took me a moment to absorb this. "The thing I saw was at the very top of the chart, Ron, and it was most certainly not an airliner."

"Hey," he interjected testily, "the only Astra anyone here ever heard of was something we considered more than 20 years ago and never built. Take it or leave it." For a moment, the receiver filled with the sound of his breathing. "Call me if you got any further questions, won't you?" And with that he was gone.

Chapter 14

There were two final way points on the invisible network of nodes and links that held the black world together and here was the first of them, the Pentagon, with its long, wide corridors leading to the offices of Lieutenant General George Muellner, principal deputy, Office of the Assistant Secretary of the Air Force for Acquisition. After a month's interlude in the U.K., I was back in Washington on official business.

It was Muellner's last day as the service's chief procurement officer. Tomorrow he retired, the culmination of a service career that had begun in the late '60s as a pilot flying F-4s out of Cam Ranh Bay, South Vietnam.

According to his bio, he had more than 5,300 hours on F-4s, A-7s, F-15s and F-16s. Muellner completed 690 combat missions over Vietnam and logged another 50 combat sorties during the Gulf War in 1991, at which time he commanded a deployment of Joint STARS battlefield surveillance aircraft. In 1991, Joint STARS was still officially under experimental test, but its belly-mounted radar proved invaluable in picking out Iraqi tanks and Scud launchers against the flat backdrop of the desert. It was, to use the jargon, an "enabling technology" that had helped the Coalition to win the war.

Another "enabler" had been stealth, a technology with which Muellner was intimately familiar, although you would never guess it from his official past. The merest clue that he had once been bound up in the stealth revolution came in the fourth paragraph of the "assignments" section of his bio, the part that outlined the key milestones of his career. While the maximum period of time contained in each of the 17 other paragraphs covered a year, two at the most, the fourth paragraph spanned an unusually long length of time: May 1973 to July 1982.

During these nine years, Muellner's activities were described as "F-15 operational test pilot, 4486th Test Squadron, Edwards Air Force Base, Calif.; chief of systems testing, USAF Test Pilot School; test pilot with the F-16 Combined Test Force; and operations officer and commander, 6513th Test Squadron, testing classified aircraft." It was kind of like Pittman Station all over again.

So, Muellner had the Right Stuff in spades and he had lived a great deal of his professional life in the black. His nine years in the test community, it turned out, covered one of the most intense periods of development in the U.S. Air Force's history—a time when it had had to take a long hard look at its equipment and tactics and rapidly come up with a raft of new ideas and hardware to ensure that it was as ready for the next conflict as it had been ill-prepared for the one it had just fought and lost in Vietnam. It was this period that presaged the biggest hike in black world spending since the days of the first and most grandiose black world project of all: the bomb.

By even stepping into the building—the great edifice constructed by General Leslie Groves prior to his appointment as director of the atomic bomb project—I knew I had entered a world of Catch-22s. If the Air Force was developing antigravity, there was no way, of course, that Muellner was simply going to come clean and admit it. Then again, if it was going on, it was also possible Muellner had been excluded from it. If this was the case, Muellner's answers to my questions, whether they were true or not, would have "plausible deniability"—the earnest and sincere response of an official whose résumé said he should be in the know, even though in reality he wasn't.

It seemed an apposite way to start, so I asked the general whether he was satisfied that he'd been briefed on everything that was being procured for the Air Force in his name.

He replied without missing a beat. "I believe so. You know, it's one of those questions you never know the answer for sure, but I think the answer is 'yes.' "

So, just what was the Air Force developing in the black?

"Things that push the envelope beyond where we're at," he replied, again without hesitation. Next-generation stealth, certain breakthrough weapons technologies, information warfare, spy satellites. That sort of thing.

Civilians who made it their business to keep tabs on such affairs, sleuths like the Desert Rats who ventured as deep as it was possible go into black world territory without getting their heads blown off, claimed that there might be as many as four secret "platforms" flying out there, I probed, including the fabled Aurora—or was it Astra now?

Suddenly, Muellner looked uncomfortable in his stiff, blue uniform; a shirtsleeves airman who'd have gladly swapped the formality of his Pentagon suite for something a little closer to the front line of advanced flight test.

"If they are," he said, "then my earlier answer about knowing about

all our programs may be faulty, because I'm certainly not aware of them."

The Air Force, he added, had no need for a reconnaissance aircraft that could fly around the globe at Mach 7 or Mach 8 when the satellites it already had in place could, for the most part, do the same job at a fraction of the cost.

Stealth, however, was a different matter and here the Air Force was developing a range of new technologies in the black.

Muellner wouldn't be specific, but he hinted that at least part of this effort entailed the development of "reactive skins" that would enable a next-generation stealth fighter to change color like a chameleon, blending against its background from any perspective. A cloaking device that could, in effect, make an aircraft disappear into the surrounding ether. Visual stealth to add to the radar and infrared variety already in place.

None of these efforts had resulted in any new operational aircraft yet, but there were test programs under way, he said, designed to "mature technology."

From this it was acceptable to deduce that *something* had been developed at the time of the Aurora sightings over the North Sea oil rig and elsewhere, that something real had been tripping the seismology sensors of the USGS, but that for whatever reason the vehicle had not been procured for front-line service. Why? Because Muellner didn't need to be saying any of this. The Air Force never talked about black programs; but on this, his last full day in office, had the general decided to invoke his other responsibility as the service's "chief information officer" and set the record straight?

The black world had taken a pasting of late. The rumors about what the Air Force was doing away from the public light of accountability were starting to spiral out of control. There were certain things that the Air Force would always need to develop in the black to maintain America's readiness in an uncertain world, Muellner said.

"And the rest?" I asked him, easing between the agreed subject matter of the interview and the antigravity question. What about the people—apparently sane and credible witnesses—who said that they saw things in the desert night skies that seemed to defy the laws of physics? Craft that hovered right over them sometimes, huge, motionless, silent?

Muellner smiled. Get him the name of the contractor, he said, because he'd love to put them under contract. Whatever was out there, it had nothing to do with the Air Force.

But here was the disconnect. People *were* seeing things that defied explanation in the vicinity of Area 51, Nevada, the U.S. Air Force base

that didn't exist a hundred miles north of Las Vegas, north of a bend on Highway 375—the "Extraterrestrial Highway," as it had been dubbed by the Nevada state legislature.

Here, Tikaboo Valley, 25 miles wide and six times as long, as lonely a place as you could find on the planet, stretched into the distance till it merged with the horizon.

It was Tikaboo Valley that connected the outside world to Area 51, hidden as it was by the jagged peaks and ridges of the Groom Range and the Jumbled Hills. Right above it, high on top of Bald Mountain, elevation 9,380 feet, sat the radar site and observation post that gave security officers an unrestricted view of all traffic entering and transiting the area. It was the first of several security rings and sensor networks employed by the Air Force to ensure that nothing penetrated the discreet *cordon sanitaire* around the base. The others, if you were unlucky, could be a lot less passive, more in-the-face: the sworn-in deputies from "Pittman Station" who patrolled the valley looking for miscreants and who occasionally locked them up and were said to throw away the key; and the Black Hawk helicopters that had been known to sandblast the odd Desert Rat from his hidey-hole on the high ground overlooking Area 51's 27,000-foot runway.

Already one of the longest paved runways in the world, it had been specially extended in the late 1980s in a move that supported the widely held view in the "stealth-watcher" community that a very high-speed aircraft was under test at Area 51. Couple this with the unmarked charter planes and buses with their windows blacked out that were ferrying up to 4,000 workers in and out of the base from Los Angeles and Las Vegas every week and word soon got around that Area 51 was as busy as it had ever been in its 40-year history—ever since, in fact, Kelly Johnson, the Skunk Works' founder, brought the U-2 here for testing in the mid-1950s.

But alongside the stealth-watchers, another set of pilgrims regularly made the trip to Tikaboo Valley. Many of these people had been drawn to Area 51 by the revelations of a man named Robert Lazar, who in 1989 claimed to have worked on recovered alien spacecraft at a site adjacent to Area 51 known as S-4. While Lazar's claims seemed wild, his followers consistently reported sightings of strange orbs of light over the base; objects that seemed to defy the way aircraft—even the top secret variety at Groom Lake—ought to fly.

But tread the fine line between the world of black programs and the world of UFOs and you entered another minefield of disinformation.

In 1997, the CIA admitted in a set of released documents that it had

encouraged reports of flying saucers in the 1950s and 1960s to obscure the flight trials of its then top secret U-2 and A-12 spyplanes. The implication was that if you saw one of these things overflying the highway and reported it, there was something wrong with your reasoning. It was a tactic that worked well.

Add this to the highly compartmentalized, watertight environment in which black program engineers and scientists worked, and the fear they were reportedly subjected to, and here, as with the Philadelphia Experiment, was a cheap, but highly effective means of maintaining program secrecy. By the cusp of the millennium, there was so much bad information mixed in with the good about black programs that it was often impossible to tell them apart.

Was Astra another case in point?

On my return from California I'd left a message on Marckus' answering machine. Maybe, I'd conceded, talking to the machine like it was a confessional, I had been expressly meant to see that chart. Maybe Astra was a subtle piece of disinformation; a misleading fragment planted ahead of my visit. Then again, maybe it wasn't. Maybe it was genuine.

Faced as I was with the official denial, Astra was every bit as effective as a hit team equipped with Colt Commandos and a platinum hard-key. Less messy, too.

This time, Marckus had called me back. I told him then what I thought of his wild-goose chase. He never attempted to deny it. I got the impression he truly believed it was good for me, like it was an extension of my education or something.

In the small hours of a particularly hot night, three days after I'd sat in Muellner's office, I watched from my mountaintop lookout as a ball of shining golden light rose above the hills that separated Groom Lake from Tikaboo Valley. The light appeared to hang in the air for a few moments before drifting lazily downrange and disappearing behind a mountain peak 50 miles to the south.

It left me strangely unmoved.

In the kind of silence that you can shred with a razor when you're a hundred miles from anywhere, I found myself thinking again about the memo from Lt. Gen. Nathan Twining, head of the U.S. Army Air Forces' Air Materiel Command, to Brig. Gen. George Schulgen, chief of USAAF intelligence in September 1947. The memo I'd spilled coffee over in my basement back in London.

Twining's observation—that objects "approximating the shape of a disc, of such appreciable size as to appear to be as large as a man-made aircraft" were neither "visionary nor fictitious"—was written just three

months after the supposed crash of a UFO at Roswell, New Mexico, in July 1947: the period that most people tout as the real start of the modern UFO sightings era.

What was it that Twining had written? That it was possible, "within the present U.S. knowledge—provided extensive detailed development is undertaken—to construct a piloted aircraft which has the general description of the object above which would be capable of an approximate range of 7,000 miles at subsonic speeds."

In mid-August 1945, I recalled, too, that 300 railroad freight cars of V-2 components captured in the European theater of operations had trundled westward across the wilderness from the Gulf of Mexico to the Organ Mountains, deep inside the dry, desert hinterland. Every railroad siding from El Paso, Texas, to Belen, New Mexico, a distance of 210 miles, had been seen to be full of cars all headed for the White Sands Proving Ground, 125 miles to the west of Roswell. Following on behind them were approximately 40 captured German scientists recruited under the covert selection program known as "Operation Paperclip." They were led by Wernher von Braun.

Maybe Roswell had happened, maybe it hadn't. But if something had come down in that summer of 1947, something extraterrestrial, there was no definitive proof, more than half a century later, to say that what had crashed there was alien; and dozens, if not hundreds of researchers had sought to discover the truth about it.

But if Roswell's extraterrestrial dimension was simply a piece of fiction, then why not turn the thing around, look a little to the left or the right, and see what swam into focus.

When I looked at Roswell this way, I saw the White Sands Proving Ground. And then I saw something else.

The V-2 rocket program had started under the aegis of the German Army, the Wehrmacht, but had been transferred in the final stages of the war on Hitler's orders to a higher authority: Reichsführer-SS, Heinrich Himmler. The same had been true of the Me 262 jet fighter and just about every other German high-tech weapons program. Most of them had started under Luftwaffe control and then migrated in the same direction. In short, anything that had shown any real promise as a weapon system—in particular, anything that appeared to represent a quantum leap over the then state of the art—had ended up under the oversight of the SS.

Eight thousand feet above sea level, I felt a constriction in my throat. I was so keyed up, my breath was coming in short, sharp gasps. But the thoughts kept on tumbling.

Just as it seemed I'd been wrong in trying to connect antigravity with the USAF, so I realized, in a moment of clarity, I'd made the same mistake in Germany.

What had *Lusty* actually stood for? *Luftwaffe* Secret Technology.

In looking for evidence of an antigravity program in the raw intelligence reports of Project *Lusty*, had I been heading up a blind alley?

If antigravity had been developed by the Nazis, then perhaps it hadn't been under Luftwaffe control at all, but that of the SS.

I called Marckus from a public phone in a windswept diner just after I turned off the Extraterrestrial Highway for Vegas. I told him the conclusions I'd reached and that the available clues said I should take another look at Germany.

"I think you're making a mistake," Marckus said. "If anyone has cracked it, it's the Americans. Stay in the States, use your contacts, go deeper."

I neglected to tell him that, thanks to his field test which showed that my research in the U.S. was merely duplicating his knowledge, I'd wasted enough time already.

Chapter 15

In the reading room of London's Imperial War Museum, under a skylight battered by autumn winds and icy rain, I immersed myself in books and papers that plotted the SS's gradual takeover of the German aerospace and weapons industry in the last two years of the war. But as the evidence mounted, I found myself increasingly distracted by thoughts of Marckus. Why had he warned me off the German research angle? Why did Marckus, with his incisive mind and razor-sharp reasoning, insist on belittling information that seemed more beguiling by the hour?

It was the intractable aspect of this contradiction that led me to pledge quietly to myself that from now on I'd be much more careful about what I told him.

It was in August 1943, just three months after the first successful test shot of the A-4 rocket, better known as the V-2 vengeance weapon, that Himmler, head of the SS, managed to wrest the V-2 from the control of the Wehrmacht, the German Army. Himmler was at the Wolf's Lair, Hitler's remote headquarters in the forests of East Prussia, when the Führer, in the company of his deputy Martin Bormann, brought up the subject of A-4 manufacture and his concerns about program secrecy. Hitler, as was his wont, was demanding the impossible.

The recent test of the A-4, which had landed just five kilometers off-target over a distance of 265 kilometers, had finally convinced Hitler, after months of vacillation, that the rocket would thwart the Allies' developing invasion plans and quite possibly bring Britain to its knees. He demanded that 5,000 rockets should go into immediate production and be delivered in the shortest possible time.

Himmler saw his chance. Quick to exploit an opportunity, the Reichsführer knew that neither the Wehrmacht nor Albert Speer, Hitler's young and energetic minister of armaments and war production, could deliver the project under existing arrangements. Hitler had decreed that "only Germans" should be employed in the production of

the A-4, but the sheer size of the task meant, practically speaking, that foreign civilian workers—Poles, Hungarians, Czechs and others from the conquered east—would have to be involved sooner rather than later in the process. But with foreign workers came the risk of the security leaks that Hitler had expressly sought to eliminate. It was to this apparently insoluble conundrum that Himmler claimed he had the answer.

What he offered Hitler, he promised, was watertight: his proposed workforce would be drawn not from the diminishing manpower resources of the Reich's civilian labor pool, but from the SS-controlled concentration camps, from which there was no escape. The same workforce, Himmler added, could be used to build the underground manufacturing facilities that would keep the A-4 secure from Allied bombing. Faced with the logic of this proposal, Hitler agreed, but informed Himmler that he must cooperate with Speer.

Though the SS controlled the camps, it was the armaments ministry that had masterminded the revitalization of the German manufacturing economy, achieving month-on-month growth, in spite of the ferocity of the Allies' strategic bombing campaign, since Speer's appointment the previous year. Yet, when he was informed of Himmler's proposal, Speer couldn't help but object to it, knowing that Himmler, whose dreams invariably exceeded his abilities, would never be able to deliver what he had promised. This was undoubtedly the case, but it was not what Hitler wanted to hear. The following day, Himmler informed Speer that he had taken charge of the manufacture of the "A-4 instrument" (as it was known to conceal its identity and purpose to outsiders) and demanded Speer's full cooperation as a subordinate. This the outmaneuvered armaments minister was forced to provide.

Thus, within 24 hours, did Himmler not only acquire control of the V-2 program, but in the process lay the foundations for the SS takeover of all secret weapons programs within the Third Reich.

For years, the Reichsführer-SS had harbored a grand scheme to turn the SS, originally established as Hitler's bodyguard, into a self-sustaining economic entity, capable of funding and running its affairs without reliance on the everyday financial and administrative mechanisms that maintained and sustained the Reich. It was via the camps that Himmler and his leading economic authority, Oswald Pohl, realized that they could achieve their aims. By 1943, population of the camps had surged—from around 25,000 on the eve of the war, to well over half a million, a figure that did not begin to reflect the millions that had already died in death camps like Auschwitz, Dachau and Mauthausen and the millions

more that would die of disease, starvation and the gas chambers before hostilities were over.

In a perverse reflection of the growth of the camps, the size of the SS itself surged from an eve-of-war strength of 240,000 to an end-of-war tally numbering 800,000. Aside from its task to maintain security within Hitler's state and its brutal administration of the death and labor camps, the SS also provided 38 fighting divisions alongside the Wehrmacht.

After the abortive bomb plot against Hitler in July 1944, a scheme hatched by anti-Hitler elements within the German Army General Staff, the control of the SS and its grip upon Germany would become absolute. But it was in August 1943, at the Wolf's Lair, that the seeds of the SS's takeover of the armaments industry were sown. Skilled workers and scientific specialists, Himmler told Speer, would immediately be transferred from the camps to begin work on the A-4. Furthermore, the Reichsführer-SS added, he had asked a "young energetic construction expert, who had already proved his outstanding ability, to take charge of the enterprise."

Speer must have had an overwhelming sense of inevitability as to the identity of this individual, but he inquired anyway. The man who was to assume direct oversight of the A-4's manufacture, Himmler told him, was Dr. Ing. (Engineer) Hans Kammler, head of the SS's Building and Works Division—the entity that had masterminded and built the camps.

Kammler's rise inside the SS had been meteoric. Up until 1941, he had been an unremarkable civil servant within the RLM, the Reich Air Ministry, with special responsibility for construction projects—hangars, barracks, administrative offices and the like. In the autumn of that year, realizing that he was never going to rise to the heights to which he aspired, he transferred to the SS, where he received the rank of Brigadeführer and was immediately put in charge of building projects. By the end of that year, Kammler had already drafted a five-year RM13-billion program of construction for SS barracks and concentration camps stretching from the newly captured territories of the Soviet Union to Norway.

The sense that the SS gave Kammler free rein to indulge his ideas and ambitions—suppressed during his years of obscure service at the RLM—is unavoidable. No sooner had his appointment taken effect, than he was drawing up detailed plans for the rapid expansion of the camps, egged on by Pohl, who had impressed upon him their value as the engines of growth for his and Himmler's grotesque vision.

To turn it into reality, Kammler immediately saw what needed to be

done and requested to expand the capacity of the camps to four million; three months later, he increased this figure again to 14 million.

While the building program got under way, he set about industrializing the existing camps, but progressively found himself thwarted by Speer, who, irrespective of any feelings he may or may not have had about the moral consequences of such a scheme, believed that the camps were not the answer to Germany's manufacturing needs. Yet it was a confrontation that Speer was bound to lose, as he tacitly admitted in his memoirs; for, in Kammler, he was essentially battling against a twisted image of himself.

"He too came from a solid middle-class family, had gone through university, had been 'discovered' because of his work in construction, and had gone far and fast in fields for which he had not been trained." Speer, it is clear, was jealous of Kammler, but while Speer patently wrestled with his conscience, especially during these years, there is no sign of any moral doubt on the part of the 40-year-old SS officer.

Kammler was cut from the same cloth as SS-Obergruppenführer Reinhard Heydrich, the author of the "Final Solution of the Jewish Problem," who was assassinated by Czech agents in Prague in 1942.

"Both were blond, blue-eyed, long-headed, always neatly dressed, and well bred. Both were capable of unexpected decisions at any moment, and once they had arrived at them would carry them through with a rare obstinacy," Speer noted.

With Heydrich dead, Himmler had found in Kammler a cold, ruthless and energetic executor of his wishes; the man who would carry though his simple, yet grandiose vision.

Kammler set about his task as overseer of the A-4 manufacture program with clinical zeal. One week after Himmler's fateful meeting with Hitler at the Wolf's Lair, he dispatched the first group of concentration camp inmates from Buchenwald to Nordhausen in the Harz Mountains of central Germany, where work on a giant underground construction facility for the A-4/V-2 commenced.

Within a month, Kammler and Speer, the latter in his new capacity on the project as junior partner to the SS, established the low-profile Mittelwerk GmbH (Central Works Ltd.) to run the rocket venture.

By the end of the year, 10,000 Buchenwald prisoners—mostly Russians, Poles and Frenchmen—had been dispatched into the limestone cliffs of the Kohnstein, the mountainous ridge close to the village of Nordhausen into which the facility was tunneled, to bring about the impossible: the construction of the largest underground factory in the world, a facility a kilometer and a half long containing 20 kilometers of

tunnels and galleries, dedicated to the construction of rockets, missiles and other top secret weapons. It was finished in a year—20,000 prisoners dying in the process.

Plans show that it was the first phase in a construction program that would have made the Mittelwerk part of an enterprise three times the size.

Thereafter, Kammler's ascent to the top levels of the bureaucracy surrounding Hitler's inner cabinet was unstoppable. As 1943 drifted into 1944, the SS's steadfast refusal to believe in anything other than final victory ensured its deepening involvement at all levels of state. In March 1944, at the formation of a combined "Fighter Aircraft Staff composed of senior Luftwaffe and armaments ministry officials tasked with increasing fighter production in the face of the Allies' determination to wipe it out with carpet bombing, Kammler represented the SS as Himmler's special delegate, offering two primary areas of expertise: his rapidly developing knowledge of underground factory construction and his ability to mobilize vast numbers of concentration camp laborers. In recognition of this, Reichsmarschall Hermann Goering, head of the Luftwaffe and Hitler's nominal successor, tasked Kammler with transferring strategic aircraft factories underground. Just over three months later, Himmler was able to report to Hitler that ten aircraft factories with an underground floor space of tens of thousands of square meters had been constructed in just eight weeks.

As the Reich's decay accelerated under the relentless news of defeat on all fronts—the D-Day invasion had placed Allied troops in France, the Red Army had just crossed into Poland and British and American bombers were pummeling German cities daily from the air—Kammler's influence spread. In the air of paranoia that followed the bomb plot of July 20, his ascendancy entered its final and most decisive phase. On August 6, following Himmler's appointment as head of armaments, adding to his portfolio as chief of internal security and commander in chief of the reserve army, Kammler was given responsibility for all aspects of V-2 operations—from design through manufacture to deployment and offensive operations against England and the Low Countries. In effect, Kammler now had sole charge of the project from planning to firing; a position that, thanks to his obsessive eye for detail, gave him an insight into the running of a strategic weapons program that no other individual has enjoyed before or since.

In January 1945, with the Red Army in Budapest and Warsaw and the Western Allies turning the tide of the Germans' Ardennes offensive, Kammler was appointed head of all missile programs—defensive as well

as offensive. On February 6th, Hitler transferred cradle-to-grave responsibility for all air weapons—fighters, missiles and bombers—to the man that many Party insiders now viewed as the most powerful and influential state official outside Hitler's inner cabinet.

Kammler had first come to Hitler's attention with a carefully crafted hand-colored design for the Auschwitz concentration camp. It was his perverse capacity to retain control of the minutiae of a project without ever relinquishing grip on the strategic objectives, a trait shared by Hitler himself, that drew him increasingly into the Führer's orbit; a characteristic that came to the fore when Kammler "reengineered" the Warsaw Ghetto after the crushing of the uprising in August 1944. He not only leveled the "contaminated area," but made sure that each of the 34 million bricks from the Ghetto was used for other essential building projects. He was also called in to advise on methodologies for increasing the daily output of the camps' gas chambers and ovens from 10,000 to 60,000. For this and his other work he was rewarded with an SS general's rank of Obergruppenführer.

By the beginning of April 1945, with the Red Army driving toward Berlin and Hitler now ensconced in the Führerbunker below the Party Chancellery, Kammler had acquired control of every single major aircraft—or rather, *aerospace*—program of any importance within the Third Reich. His empire now included any project or facility that had either an SS influence or was aircraft or missile related. Geographically, this gave him the authority to enter and operate any weapons plant within the shrinking borders of the Reich, by now restricted to Norway, Denmark, northern Germany, southern Bavaria, Czechoslovakia, Austria and northern Italy.

What is astounding is the belief of the Nazi leadership, even at this terminal stage of the war, that Kammler would bring about miracles and deliver the German people from the relentless advances of the Soviets in the east and the British and Americans in the west. "The Führer has had prolonged discussions with Obergruppenführer Kammler who now carries responsibility for the reform of the Luftwaffe," propaganda minister Joseph Goebbels wrote in his diary on April 3. "Kammler is doing excellently and great hopes are placed on him."

Yet, for all the power vested in Kammler at this late stage of the war he barely rates a mention in standard works of reference on the Luftwaffe or its major programs. There is so little data on him, in fact, that archivists at London's Imperial War Museum confessed that their files contained next to nothing on him.

Just when I thought I was going to have to look elsewhere, perhaps in

Germany itself, I received a phone call from a diligent researcher at the Imperial War Museum who'd become as engrossed as I was by the lack of information on Kammler.

The researcher told me, a note of triumph in his voice, that he *had* turned up a book on Kammler after a rigorous search: a slim volume, with a curious title—*Blunder! How the U.S. Gave Away Nazi Supersecrets to Russia*—penned by a British Cambridge graduate-turned-foreign correspondent named Tom Agoston.

What *Blunder!* contained, however, was dynamite, for Agoston had established what no other researcher has managed to before or since.

While Kammler carried out his job to the letter, churning out the rockets and jet aircraft that Hitler hoped would turn the tide against the Allies in the closing weeks of the war, he also set up, unbeknownst to anyone connected with those projects, a top secret research center tasked with the development of follow-on technology, a place where work on "second-generation" secret weapons was already well advanced.

"In modern high-tech jargon, the operation would probably be referred to as an 'SS research think tank,'" Agoston wrote.

In the jargon of the U.S. aerospace and defense industry, however, Kammler's "think tank" would have been known by a different term—one that would have been familiar, though, to anyone who had ever been consigned to the black world.

What Kammler had established was a "special projects office," a forerunner of the entity that had been run by the bright young colonels of the USAF's stealth program in the 1970s and 1980s; a place of vision, where imagination could run free, unfettered by the restraints of accountability. Exactly the kind of place, in fact, you'd expect to find antigravity technology, if such an impossible thing existed.

Chapter 16

According to Agoston, Kammler set up his special *Stab* (staff) for these visionary projects in a highly compartmentalized section of the Skoda Works in the Reich Protectorate of Bohemia and Moravia in the vassal Nazi state of Czechoslovakia. Through the brutal administration of Heydrich, the Protector of Bohemia-Moravia until his assassination in June 1942, the SS had come to treat the industry of the entire Czech region as its own private domain. In March 1942, Himmler formally took over the operations of the Skoda Works, a giant industrial complex centered on the Czech towns of Pilsen and Brno, as an SS-run operation. In the process, he managed to bypass Speer, who knew nothing of the arrangement until Hitler told him it was a *fait accompli*. With the SS in charge of Skoda, Hitler tasked Himmler with evaluation and, where appropriate, duplication of captured enemy weapon systems whose performance outstripped the Germans' own. Under the leadership of Skoda's director general, honorary SS-Standartenführer (colonel) Wilhelm Voss, this project was completed in record time. On May 11, 1942, Voss was able to explain to Himmler what it was that had enabled him to progress the development of these back-engineered weapon systems so successfully—"by concentrating all developments for the Waffen-SS in the liaison staff of Skoda and Bruenn [Brno] weapons plants, and assuring systematic, intense cooperation with the SS Ordnance Office."

In effect, Voss was telling Himmler that he had been able to move as quickly and effectively as he had thanks to the removal of the overweening bureaucracy that went hand in hand with projects administered by Speer's armaments office.

As far as the regular mechanisms of the Reich procurement machine were concerned, then, the Skoda operation was to all intents and purposes beyond oversight. This would set the scene for the Kammler special projects group, which was so deeply buried, so beyond regular control, it was as if it did not exist.

Voss, "tall, quietly reserved in manner, graying, upright and elegant,"

would become Kammler's alter ego in the administration of the special projects group. Seven years older than the SS general and with a reputation as an industrialist that was as solid as a rock, Voss, with his nuts-and-bolts background, was the ideal complement to Kammler's driven bureaucrat. Even better, the two men had a task from Hitler and Himmler to run the special projects group without the least interference from Goering or Speer. Apparently, neither Goering, head of the Luftwaffe, nor Speer, head of all armaments programs, would have the least idea it existed.

But for Tom Agoston, in fact, the reality of the *Kammlerstab* might never have emerged at all.

Agoston was a Cambridge graduate who had served during the war as an air photo interpreter. After the war, he traveled to Germany to report on the Nuremberg war crimes trials. It was there that he ran into Voss.

Though not indicted by the war crimes tribunal, Voss was so close to the Nazi elite that his name was constantly on the lips of the prosecutors.

"Everyone wanted to interview Voss, but they couldn't find him," Agoston told me over the phone from his home in Germany a half century later, the satisfaction still evident in his voice. "As he was sheltering in my house at the time, this was hardly surprising."

In the course of several extended interviews in 1949, Voss unburdened himself of the whole Skoda story, speaking of his activities there with "unique frankness."

Tired and disillusioned, Voss was patently grateful for Agoston's company and when the subject of the Kammler Group surfaced, he told the reporter everything.

The Kammler special projects office was regarded by the select few who knew about it as the most advanced high-technology research and development center within the Third Reich. It was totally independent of the Skoda Works' own R&D division, but used Skoda for cover.

All funding for the programs at the *Kammlerstab* was channeled through Voss, who reported alongside Kammler directly to Himmler. The scientists were culled from research institutes all over the Reich, chosen for their acumen as engineers and scientists, not for their allegiance to the party.

"Many scientists, anxious to see their work in print, even if it was kept top secret, prepared papers for a central office of scientific reports, which circulated them to specific recipients. Some of these reports were used as the basis for selecting candidates for employment at Skoda," Agoston wrote.

Once recruited to the special projects group, whether they liked it or

not, the scientists set about their work, their activities protected by a triple ring of security provided by SS counter-intelligence specialists assigned specially for the task.

These security rings were established by Himmler around Skoda's sites at Pilsen and Brno and their administrative center in Prague.

Voss described the activities of the scientists at the *Kammlerstab* as beyond any technology that had appeared by the end of the war—working on weapon systems that made the V-1 and the V-2 look pedestrian. Among these were nuclear power plants for rockets and aircraft, highly advanced guided weapons and antiaircraft lasers. The latter were so far ahead of their time that by the end of the 20th century there was still no official confirmation of laser weapons having entered service, despite the best efforts of the Russians and the Americans.

In March 1949, several weeks, after he briefed Agoston, Voss was pulled in for questioning by the U.S. Counter-intelligence Corps.

Shortly afterward, he wrote in secret to Agoston and requested that *die heiße Angelegenheit*—"the hot matter"—they had discussed earlier that year was not for publication.

Voss had told the CIC agents about the range of research activities that had been pursued by the special projects group and they informed him, in no uncertain terms, that he was never to speak about the *Kammlerstab* or its programs to anyone.

Voss gave them his word, neglecting to tell them, of course, that he had already briefed Agoston.

Agoston duly honored Voss' request to put a lid on the matter, but with Voss' death in 1974 he was absolved of his obligation and began picking up the threads of the story again. Voss' interrogation reports should have been freely available under the U.S. Freedom of Information Act, but despite persistent requests, Agoston was told they were unavailable; that there was no record, in fact, of any such interrogations ever having taken place.

I recalled from the *Lusty* documents I had examined in Washington a report that had been filed by OSS agents, forerunners of the CIA, from Austria in April 1945. Bundled into a collection of unprocessed notes on Nazi high-technology weapons "targets" examined by U.S. intelligence specialists, the report had mentioned a location in Vienna where experiments had supposedly been conducted on "antiaircraft rays"—laser weapons.

"Research activity is conducted in a house at the above address (87, Weimarerstr. Vienna). Research personnel were not allowed to leave house (reported hermetically sealed)."

Funny, I thought, how this much later on in the investigation I saw things through a markedly different prism. At the time, I'd been focused on the technology. Now, the fact that the Germans had been working on directed energy weapons was subordinate to a subtly different observation: that the Nazis had operated a compartmentalized security structure like the one now operated by the black world.

It was then that I saw the Kammler special projects group for what it was; not as a facility where experiments were conducted, but as an R&D coordinating center. Testing wouldn't have been performed at Skoda itself, but in the field.

It was exactly as Agoston said it was: a think tank, not an experimental site.

I put out feelers into the Czech Republic and Poland. I now saw that if there had been any truth at all in the reports of flying saucers developed by the Germans at the end of the war, this was the place to start looking. If the legend that had grown up around the supposed activities of Messrs. Schriever, Habermohl, Miethe and Bellonzo was in any shape, way or form based on truth—and while my earlier research had shown there wasn't a shred of solid evidence to say they were, I was convinced there was no smoke without fire—two places always seemed to recur in so-called testimony.

The first was Prague, not a million miles from Skoda's twin hubs at Pilsen and Brno (Bruenn). The second was Breslau in the German province of Lower Silesia. Then part of Germany, Breslau—renamed Wroclaw—is today in southwest Poland.

I turned my attention back to Kammler.

In March 1945, Kammler moved his headquarters from Berlin to Munich, working out of the regional Waffen-SS and Reich Police Construction Office. On April 16, three weeks after he had received his mission from Hitler to change the course of the war, Kammler delegated Gerhard Degenkolb, industrial plenipotentiary for the manufacture of jet aircraft at Speer's ministry for armaments and war production, to assume special responsibility for manufacture of the Messerschmitt 262, Germany's last-ditch hope in the war in the air. On April 17, with the Third Reich crashing around his ears, Kammler sent a message to Himmler at SS Command Headquarters in Berlin denying Himmler the use of a heavy truck that the Reichsführer-SS had requested from the Junkers aircraft factory motor pool—an order so obsessive in its attention to detail, given the inferno that was burning around Kammler, that Speer later described it as "both terrifying and laughable."

Be that as it may. It was the last official message anyone ever received

from Kammler, who by April 18 had dropped off the map—effectively without trace.

"In the course of my enforced collaboration with this man," wrote Speer of Kammler after the war, "I discovered him to be a cold, ruthless schemer, a fanatic in pursuit of a goal, and as carefully calculating as he was unscrupulous."

If there was one thing I had learned about Kammler in the short time I had spent analyzing his activities, it was that nothing in the clockwork routine of his life happened by chance. By the beginning of April 1945, while Hitler was heaping new responsibilities on his shoulders almost by the day, the general would have been acutely aware that it was just a matter of time before Germany collapsed.

In contrast to Himmler, who vainly believed that he could enter into armistice negotiations with the Allies and emerge from them with a leadership role in the reconstituted Germany, Kammler was a realist. His part in the construction of the concentration camps, the clinical way in which he had sought to boost the throughput of the gas chambers and ovens, not to mention the methods he had employed in leveling the Warsaw Ghetto, would have placed him high on the list of SS officers sought by the Allies for war crimes.

Unlike Himmler, however, Kammler had something of value to deal—something tangible. By early April, Hitler and Himmler had placed under his direct control every secret weapon system of any consequence within the Third Reich—weapons that had no counterpart in the inventories of the three powers that were now bearing down on central Germany from the east and the west.

A man who had no trouble transforming the grand visions of others into reality while retaining control of the minutiae, would have found it just as easy to talk victory and plot an exit strategy at the same time.

The clues, which double as the countdown to Kammler's disappearance, were there from the beginning of that month.

On April 3, Kammler had his meeting with Hitler in Berlin. As Goebbels reveals, following their discussions about the "miracle" weapons that could still win Germany the war, the Führer retained great faith in him. But Kammler had already moved his headquarters out of Berlin to Munich for reasons that Speer guesses at—accurately, it seems—in his memoirs: given the expected failures of these projects, it would have been dangerous for him to remain within Hitler's reach.

Before he left Berlin for the last time, soon after his final meeting with Hitler, Kammler paid Speer a visit. He had come to say good-bye.

"For the first time in our four-year association, Kammler did not display his usual dash," Speer recalled. "On the contrary, he seemed insecure and slippery with his vague, obscure hints about why I should transfer to Munich with him."

Speer might have put this down to the fact that the central part of Kammler's kingdom, at least as far as he, Speer, was aware—the giant Mittelwerk underground A-4 construction complex at Nordhausen in central Germany's Harz Mountains—was about to be overrun by the U.S. First Army. With most of the A-4/V-2 program team still in place at Nordhausen, Kammler's nervousness was understandable. He needed to get them out.

But then, Kammler told the armaments minister that "efforts were being made in the SS to get rid of the Führer."

As Speer was also in the throes of a desperate scheme to assassinate Hitler, using poison gas to kill him and his entourage in the bunker beneath the Reich Chancellery, this perhaps came as no great surprise, except that it came from Kammler's lips. Speer did not know it, but SS General Walther Schellenberg, Germany's combined intelligence chief, had already opened secret surrender negotiations on behalf of Himmler with Count Folke Bernadotte, vice chairman of the Swedish Red Cross. These were transmitted to the Western Allies, but promptly and roundly rejected.

But then Kammler told Speer he was planning to contact the Americans and that in exchange for a guarantee of his freedom he would offer them everything—"jet planes, as well as the A-4 rocket and other important developments." He informed Speer that he was assembling all the relevant experts in Upper Bavaria in order to hand them over to U.S. forces.

"He offered me the chance to participate in this operation," Speer said, "which would be sure to work out in my favor." Speer turned him down. In a late display of remorse for his years of unquestioning devotion to Hitler, the armaments minister now saw it as his solemn duty to prevent Hitler carrying out his scorched-earth policy in the face of the advancing Allies. As soon as Hitler was issuing orders for factories to be burned to the ground or blown up, Speer was countermanding them, arguing with frightened and bewildered plant managers that the war was lost and these facilities would be needed for the economic revival of the "new Germany."

On April 10, Army General Walter Dornberger, onetime head of the A-4/V-2 program, now subordinated to Kammler, ordered 450 key A-4 scientists to evacuate the Nordhausen complex and head for the

mountains south of Munich. Those that could not board a special south-bound train, nicknamed by those on it the "Reprisal Express," would be left behind at Bleicherode close by, where Dornberger and von Braun had established their headquarters.

Von Braun argued against this move on the grounds that it would mean deserting his men in the Harz, but Dornberger reminded him that Kammler had issued the order and if they did not obey it, "the SS would shoot them all."

Kammler did not come with von Braun and Dornberger on the journey south, but he did make arrangements to meet them at the Hotel Lang, near Oberammergau, which was already serving as an evacuation site for Messerschmitt's advanced projects team, evacuated late the previous year from the company's bombed-out factory at Augsburg.

Von Braun and Dornberger made their way by car to Oberammergau, arriving there on April 11, the day that U.S. forces entered and overran the underground factory they had just left. The two men, who had co-operated on the rockets from the beginning, were settled into a compound that had been prepared for them by the SS. On his arrival, von Braun remarked on the beauty of his new surroundings, the snow on the peaks, the lush pasturelands, but noted, too, that the barbed wire had been designed to keep them in, not the Americans out.

Soon after he arrived, von Braun made his way to the Hotel Lang.

As he sat in the lobby, he could hear two people talking in an adjoining room. Straining to catch what they were saying, the rocket engineer recognized the voices as those of Kammler and his chief of staff, SS-Obersturmbannführer Starck.

The two men were discussing a plan to burn their uniforms and lie low for a while in the 14th-century abbey at Ettal, a few kilometers down the road. From the tone of the conversation, von Braun was unsure if they were serious or joking. It was at that moment that an SS guard appeared and ushered him into the room.

Kammler, sitting next to Starck, appeared relaxed and affable—very different from the man that Speer had described a week earlier in Berlin.

He asked von Braun how he found the accommodation, whether his team was comfortable, if the broken arm that von Braun had suffered in a car crash the previous month was causing him any great distress, whether the team would be able to begin its design work again soon.

Von Braun, noting the machine pistol propped beside Starck's chair, replied as enthusiastically as he could to Kammler's questions, careful to maintain the charade. To have done anything less might have been fatal

given Kammler's ruthless nature; besides which, von Braun had made provisions of his own.

Before leaving Nordhausen, he had had his men stash ten tons of blueprints relating to the A-4 and its successor, the intercontinental A-9/A-10 rocket, in a deserted iron-ore mine several kilometers from the complex. Had Kammler any inkling of this, von Braun had no doubt as to the fate that awaited them all.

But with the A-4 team settled in Bavaria, the general didn't appear overly concerned about the scientists anymore. He had other matters to attend to, he told von Braun, matters relating to his duties as Hitler's special plenipotentiary for jet aircraft production. He would shortly be leaving Oberammergau for an indeterminate period. SS Major Kummer would assume command in his absence. Kammler got to his feet. The meeting was over.

They said their good-byes and von Braun left.

It was the last time anyone of any standing could corroborate seeing Kammler.

Yet there are traces of his movements in the days, even weeks, that followed; and some of his subsequent actions are decipherable—to a point.

Kammler told von Braun that he had to leave Oberammergau to oversee production of the Me 262 jet fighter according to Hitler's wishes. Yet, five days later, on April 16, the general appointed Degenkolb to the job, thereby delegating his responsibility; to someone eminently capable, as it turned out. Kammler communicated this decision from his office in Munich, dispatching it to Speer, Goering, Himmler, Hitler's Luftwaffe liaison officer Colonel Nikolaus von Below and SS General Hermann Fegelein, Himmler's liaison officer in the Führerbunker. Describing this communiqué years later as a bafflingly "unimportant message," even Speer signals his bewilderment at Kammler's determination to spread the word to all and sundry about Degenkolb's appointment.

It seemed obvious with the benefit of hindsight that Kammler wanted as many people as possible to see that he was on top of his assignment—that he was in Munich doing what he had been told to do.

With Degenkolb in charge of Me 262 production, a job that required constant supervision, it left him free to pursue other matters.

When Kammler told Speer that he would offer the Americans "jet planes and the A-4 rocket," he must have known that their currency was devalued by the sheer numbers of people associated with them. The Me 262, Arado Ar 234 and the Heinkel He 162 jets were in widespread

service with the Luftwaffe at the end of the war and plans for hundreds of other jet aircraft were on drawing boards at aircraft factories across the Reich. The Americans and the Russians could take the blueprints for these projects and the design staff associated with them irrespective of Kammler.

The same went for the A-4. As von Braun's actions testify, the rocket had been secretly earmarked for handover to the Americans for some time by the people who had designed and built it. Whether Kammler was aware of these moves or not is immaterial; the fact that there were hundreds of V-2s in various stages of construction at Nordhausen meant that the Americans would simply take them as war booty, with only the fate of the design engineers left to negotiate.

If Kammler did have plans to hand the Dornberger/von Braun design team over to the Americans, these came to naught. It appears, though, that he never even tried.

On April 13, von Braun succeeded in persuading Major Kummer to disperse the scientists into the surrounding villages, ostensibly to cut down the risk of their being wiped out by air attacks on Oberammergau. Von Braun's real fear, of course, was that it would be Kammler issuing the orders for their annihilation.

The move may well have saved their lives, but it also left von Braun free to make his own deal with the U.S. agents that had been dispatched into Bavaria to seek him out.

On April 17, Kammler sent his message to Himmler about the "truck"—the last signal he was known to have dispatched from his Munich headquarters. Under German communications procedure, his teletyped "signature" was preceded with the letters "GEZ"—an abbreviation for "*Gezeichnet*" or "signed by"—denoting to the recipient that he was physically where he claimed to be.

The fact is, he could have been anywhere—anywhere within a narrow corridor of territory then still in the Reich's possession.

As I immersed myself in as many sources as I could regarding Kammler's mercurial existence, a single question pounded repeatedly above the others.

Where had he gone?

Degenkolb was taking care of jet aircraft production. The inner circle in the Führerbunker, cut off from reality and increasingly from the world, had its own problems. No one seemed to care anymore about the whereabouts of the golden boy—the keeper of the miracle weapons who would save Germany from annihilation.

They could have had little idea that he was planning to bargain them for his freedom—for his life.

What did Kammler hope to trade? The rockets were gone. The jets were gone. That left the "other developments" he had told Speer about.

On April 17, the GIs of General Devers' 6th Army Group were just days—hours, perhaps—from Munich. All Kammler would have to do was hole up and wait for them.

But what evidence there is suggests he did something quite else.

There is testimony that Kammler moved on from there to Prague, a journey that would have taken him right past the Skoda Works at Pilzen.

Voss had spoken about nuclear propulsion, highly advanced guided weapons and lasers. What other secrets resided within the *Kammlerstab*?

If the witness testimony was accurate, if Kammler did move into Czechoslovakia, what was so precious about the inner workings of the "special projects group"—this Pandora's box of advanced weaponry—that it compelled him to go east, not west? Toward the Russians, of all people?

Nothing in Kammler's carefully coordinated life happened by chance. I knew that much. The documentation led nowhere. Even Agoston had reached the end of the trail.

All that was left, was Kammler's lasting legacy: the underground factories that riddled his kingdom, built to his carefully exacting standards, forged by the Holocaust that was also largely of his making.

I requested some time off work and booked myself a fly-drive package to Hanover. From there, with a good rental car, I would be in the Harz Mountains less than two hours after the wheels of the aircraft greased the airport tarmac.

Chapter 17

I entered the complex from the southern end of the Kohnstein—a long ridge dotted with pine and deciduous trees that rose like the hackles of a crouched beast above a sleepy plateau in the Harz Mountains close to the small town of Nordhausen. Inside, I hesitated before drawing breath. Twenty thousand people had died in the construction of Nordhausen, many of them right here, inside the mountain itself.

Anywhere else and you could half pretend that you were in the bowels of an archaeological site; where lazy reveries about the past of the place were part of the experience. But the imprint of what had happened here within the bare limestone walls was indelible and overwhelming.

I was standing at the entranceway of Gallery 46, a long tunnel with a semicircular roof. It was still littered with wartime debris. The records said that it had been used in the construction of the V-1 flying bomb.

Every now and again it was possible to recognize a piece of hardware—components of an Argus pulse jet destined for a V-1, a section of fuel tank or a part of a wing—but, for the most part, in the dim yellow electric light, it could have been any old scrap metal. Next to nothing had been touched, my guide informed me, since this part of the facility had been reopened to visitors in 1994.

Soon after the war, when eventually the Americans left the site—but not before they'd shipped 100 fully assembled A-4/V-2s back to the States—the Soviets sealed Nordhausen by blowing the entrances to the long galleries. In time, the trees had returned to the Kohnstein and East Germans could almost forget that the place had ever been here.

There was a kind of symmetry to this; the Nazis had gone to very elaborate lengths to conceal it during the war, even from their own people.

The Mittelwerk was vast. At the height of its activities, it had been turning out thousands of V-1s and V-2s a month and had consisted of almost 100,000 square meters of floor space. Built in the 1930s as an underground storage facility for fuel oil, the complex—already huge—

gave Speer and Kammler a head start in the construction program they began and completed between 1943 and 1944.

The site consisted of two main tunnels running in parallel between the northern and southern faces of the ridgeline. One of these, Tunnel B, the main V-2 production line, was a kilometer and a half long.

Between the tunnels, like the rungs of an enormous ladder, were 50 "galleries," each between 100 and 200 meters long. Many had been component manufacturing facilities for the giant rocket assembly line. Some, such as those given over to the manufacture of jet engines, were assembly lines in their own right.

Until the construction of concentration camp Dora nearby, the slave workers who had prepared Nordhausen for the V-1 and V-2 had lived in situ.

Six thousand prisoners had crammed into three tunnels, Number 46 being one of them. Initially, they had slept on straw, their surroundings lit only by a few carbide lamps. Then, they had been allowed to build bunks, four-story affairs that touched the ceiling, but which still failed to accommodate everyone. Since work went on around the clock, this did not matter; the laborers went to work and slept in shifts.

Beyond Gallery 18, deep inside the mountain, there was no ventilation, water or heating. Every explosion set off by the tunnelers filled the galleries with blinding, choking dust and gases.

This, mixed with the smell of the open latrines—oil drums sawed in half with two planks on top—made the air almost unbreathable. They also made the tunnels a place of terrible disease and infection. It was a privilege to be on the evening rota that loaded the latrines onto train wagons and emptied them in the open air.

A little more than 50 years later, as I walked the galleries beneath the Kohnstein, I found the silence deeply troubling. I'd wanted to concentrate on the science and engineering, to look at the Mittelwerk as a place that had been responsible for the most awesome piece of technology to have come out of Europe during the war; to breathe the same air as those who'd created the A-4 rocket's assembly line, to soak up that atmosphere in the way that a profiler would use a murder scene to build a picture of the killer. This place, after all, was one of the final staging posts to Kammler's disappearance. I needed to think as he had thought. But uppermost in my mind was the sheer scale of the crime that had been committed down here and its unexpected proximity to my own ordered existence. The people who'd toiled and died down here to create this Dantean design from Kammler's notebook, the same one perhaps that

had spawned his delicate sketches of Auschwitz, were just a generation away from the safe confines of my own world. And all I had succeeded in doing was immersing myself in the fate of its slave workers, not in the thoughts of the icy technocrat who'd driven them on. In the subzero chill of the tunnels, all I could hear were the cries of the dying, not Kammler's whispered escape plans.

Only when I reemerged from the tunnels could I reapply my mind back to the problem.

When the building work on the tunnels stopped, those that survived went on to work on the rockets. At first, quality control was poor and many of the rockets exploded or went off course.

Suspecting sabotage, the SS ordered mass executions.

On one day in March 1945, the guards hanged 52 people in Gallery 41, tying a dozen at a time to a beam which was then pulled up by an electrical crane. Those next in line to die were forced to watch.

Then there was Gardelegen, 80 miles to the north. On April 13, two days after advance U.S. units entered Nordhausen, the SS forced a thousand evacuated prisoners from Dora into a barn that had been pre-soaked with gasoline, locked the doors and set light to it. When American troops came across the scene, they found some of the bodies still in the barn, some in a partially dug mass grave. Twenty men had survived by sheltering in holes they had dug beneath the charred corpses of those that had perished in the first moments of the conflagration.

These were Kammler's hallmarks. Others might have pulled the trigger, operated the cranes or poured the gasoline, but these acts had been carried out according to his wishes. More than 50 years later, you could sense it.

This was something about the man that I had pulled from the dank, subterranean air.

It was curious, then, that so few people after the war—the many tens of thousands whose lives had been touched by Kammler—could recollect him in any detail. I found this incomprehensible at first, remembering occasions in which concentration camp survivors had identified their tormentors decades after the war.

Even those who had not been directly traumatized by Kammler commented upon this mercurial ability to blend in. A U.S. diplomat who had served in Berlin until the U.S. entered the war in 1941, whom Agoston had interviewed, recalled his ability to "soothe and tame an unruly horse, using a magically gentle touch, and then minutes later order a negligent groom to be brutally horsewhipped."

"I always thought he was a man to watch," the American told Agoston.

"But I never saw him again. When I got back to Berlin after the war, no one seemed to remember him."

How could this man, this monster, the most powerful individual in Nazi Germany outside Hitler's inner circle in the last days of the war, come to be quite so easily forgotten?

It was only outside Nordhausen's giant bombproof doors, as I gazed again at the single wartime picture that exists of Kammler—in his general's uniform, striding for the camera, his cap with death's head badge enough to one side to betray more than a hint of vanity—that I began to understand. Kammler was fair, as so many Germans are. There was a gleam of raw intelligence behind those expressionless eyes and a hint of cruelty about the mouth. But take away the uniform, take away the job, the jut to the jaw and the energy in the stride, and he could have been any average 40-year-old European male.

In the chaos of the collapsing Reich, Kammler could have gone anywhere he wanted, assumed any persona he liked, and no one would have been the wiser.

On April 11, 1945, when the first elements of the Third U.S. Armored Division of the First Army swept into Nordhausen, Kammler's legacy would have been plain for the Americans to see.

Clearly, however, his war crimes were not uppermost in their minds.

When Voss was picked up by agents of the U.S. Counter-intelligence Corps soon after hostilities ended, he told them about the existence of the *Kammlerstab* at Skoda. Their reactions surprised him. They appeared so unmoved by the notion that the special projects group contained a Pandora's box of exotic military secrets, he concluded that U.S. intelligence already knew.

When Voss went on to propose that they should spare no effort in finding Kammler before the Russians got to him, the CIC agents appeared equally uninterested.

This, from a nation that had mounted the biggest plunder operation of all time—Involving the Army, Navy and Air Force, as well as civilian agents—to bring advanced German technology back to the United States.

At first, I wondered if their behavior had to do with the fact that Pilsen had fallen within the designated Russian zone of occupation. Maybe the CIC recognized that Kammler was beyond their reach and had simply abandoned any hope of finding him?

Then I came upon the final piece of the jigsaw.

The rapid eastward push that had brought the Third Armored Division to Nordhausen did not stop there. Disregarding agreements

signed by the exiled Czech government and the Soviet Union, troops of Patton's Third Army to the east of Nordhausen crossed the Czech frontier on May 6.

Deep into the Soviet-designated zone of occupation, a forward unit of Patton's forces entered Pilsen that morning. Records I'd had sent to me from the U.S. National Archives before I left England showed that U.S. forces had run the Skoda Works for six days, until the Red Army showed up on May 12. Following protests from Moscow, the U.S. Third Army was eventually forced to withdraw.

Six days is a long time if you're retrieving something you already know to be there.

Had Kammler already done his deal with the Americans? Speer, during his initial interrogation at the hands of the U.S. Strategic Bombing Survey Mission on May 21, less than two weeks after the German surrender, intuitively believed so.

Asked by the mission for technical details of the V-2, he replied: "Ask Kammler. He has all the facts."

As I pushed eastward toward the Czech border, using wartime documents I'd gathered in London and a worn copy of *Blunder!* as my guide, I found myself following a scent of collusion that was still choking more than 50 years later.

Chapter 18

On the morning of Sunday May 6, 1945, five days after the announcement of Hitler's death in Berlin, troops of the U.S. 16th Armored Division pushed cautiously through an eerie, soulless landscape that its soldiers described as "neither German nor Czech." The local people—German-speaking, terrified by their Slav neighbors to the east and undecided whether to remain in their homelands or flee westward—were powerless to resist the U.S. troops.

The war was all but over and everyone knew it; everyone except a hard core of Waffen-SS in and around Prague, southern Czechoslovakia and Austria that had signaled their intention to defend this, one of the final remaining pockets of the Reich, to the very last.

As the half-tracks rolled deeper into Czech territory, the U.S. soldiers passed ragged columns of Wehrmacht troops rolling westward to the comparative safety of the U.S. zone of occupation. The U.S. objective was Pilsen, a place known to the soldiers of the 16th Armored for its beer, not the giant armaments factory where Skoda had cranked out guns and munitions for the Nazis for the best part of seven years.

Inside the U.S. half-tracks, you could cut the atmosphere with a knife. No one wanted to end up a statistic in the last few days of the war. But still, the pace was relentless. Rumor had it that Patton wanted to reach Prague ahead of the Russians. And that meant a confrontation with the SS.

Approaching Pilsen, the atmosphere changed. The U.S. troops noticed a scattering of red, blue and white Czech flags in place of white flags of surrender. As the half-tracks left the Sudetenland behind and pushed on into the heartland of Czechoslovakia proper, the people came out into the streets, waving and cheering as the 16th Armored swept past. "It was Paris all over again, with the same jubilant faces, the same delirium of liberation," one soldier recalled in the 16th Armored Division's official account of the liberation of Pilsen.

By 0800 hours, the U.S. troops were in the center of Pilsen, their advance all but halted by the crowds that had spilled off the pavements

and into the streets. The situation was chaotic. It was only when German snipers opened up in many sections of the city an hour after the Americans arrived that sanity of a kind was restored. The civilians fled back into their homes, leaving the U.S. soldiers to mop up what resistance remained and carry out house-to-house searches.

One unit from the 16th Armored soon came upon Lt. Gen. Georg von Majewski, the German army commandant in Pilsen, with 35 members of his staff in the Wehrmacht's town headquarters. As American troops entered von Majewski's office, the general surrendered his forces to the officer in charge, then pulled a pistol from the drawer of his desk and shot himself through the head in front of his wife.

Von Majewski was one of 14 enemy dead that day. Around 4,500 more surrendered. The Americans had suffered one fatal casualty in the whole Pilsen operation.

A second 16th Armored unit, Task Force "Casper," captured the airport and a third, Task Force "Able," occupied the Skoda Works.

An air raid by U.S. B-17s several weeks previously had rained destruction on a large part of the facility. The forward units of Task Force Able reported that the central administrative building had been 80 percent destroyed.

The news that Pilsen had fallen to the Americans was picked up on the radio that night by the onetime head of the Skoda Works, Dr. Wilhelm Voss, who was hiding out with his wife and family at a hunting lodge on a Skoda estate south of Prague.

Voss had been out of a job since January, when Goering had dismissed him for refusing to accept two of his nominees on the Skoda main board.

For weeks, Voss had been monitoring the Allies' radio reports for news of their advance, knowing that it was touch and go whose forces would get to Pilsen first. Armed with the news he wanted to hear, Voss seized the moment soon after the German surrender on May 7 and set out for Skoda.

Given Voss' former position and the chaos that engulfed the former Protectorate, this was a hazardous undertaking. Czech partisans were summarily executing all the Germans they could find. While the bulk of the Waffen-SS units had officially surrendered, some were reported still to be holding out against the Russians in and around Prague. But as Voss told Tom Agoston during their conversations in 1949, he was determined to hand over the blueprints of Kammler's SS special projects group to the Americans, not the Russians.

According to *Blunder!*, Voss finally pulled up in front of Skoda's main gates on May 10. He bluffed his way past the American guard and found

a very different place from the one he remembered; a Skoda in transition, with Germans, Czechs and Americans milling about the plant. In the confusion, he located a member of the *Kammlerstab*, who told him that survivors from the group had already recovered some documents, marked them up as personnel and payroll records, and loaded them onto a truck. They were hoping to get them to a place of safekeeping before the Russians arrived. The facility was due for handover to the Red Army in two days' time.

Sensing that this was the time to act, Voss sought out the U.S. officer in charge, informed him who he was, then told him what was in the truck. The American listened, unimpressed by what he was hearing, and finally informed Voss he was under orders to hand everything over to the Russians. Voss insisted, but the American was adamant; nothing was to be moved. An American ordnance team had already inspected the plant earlier that week, he told the exasperated industrialist, and had picked up everything it needed.

Two days later, Voss watched as the truck was driven away by a Red Army transport officer.

In *Blunder!*, Tom Agoston had traced records showing that a British-led Combined Intelligence Objectives Sub-Committee team entered the Skoda Works on May 12 and promptly got the runaround from Skoda's newly ensconced Czech managers, who wanted to cooperate with the Russians and no one else.

I had found another set of CIOS reports, from a British team that entered Skoda one week later on May 19 with the plant now firmly under Soviet jurisdiction. This report, CIOS Trip No. 243, summarized a combined U.S.-U.K. operation—the U.S. team, it said, having entered the plant on May 16—in which nothing of any real interest was discovered, except for certain new projects in the ammunition field and some advanced German design and production methods for guns and munitions.

The triple ring of security put in place by Himmler and Kammler around the *Kammlerstab* had done its job, even after hostilities had ended.

Voss' story implies that Kammler never made it to the Skoda Works to retrieve the blueprints of the *Kammlerstab*—that instead the papers were ignominiously removed by the Russians and taken back to the Soviet Union.

It didn't look that way to me.

Based on what was happening in Germany, where U.S. technical intelligence agents were systematically targeting any weapons facility of interest and plundering the contents, it is inconceivable that they would

have ignored Skoda, even if they had been unaware of the SS-run special projects group in its midst.

The U.S. officer who stonewalled Voss over the papers in the truck told him that a U.S. "ordnance team" had inspected the plant earlier that week. "Earlier that week" could only mean that the "ordnance team" had walked into the plant alongside or directly behind the combat units of 16th Armored's Task Force Able on May 6.

There is no official record of any such unit's visit to Skoda. There are, however, reports that a U.S. team with nuclear expertise entered the plant and gained access to documents that outlined the SS cell's work in conjunction with the Junkers company at Dessau on nuclear propulsion for aircraft. I had been informed by someone that an account of this activity was contained in *Atomic Shield*, the official history of the U.S. Atomic Energy Commission. But after extensive inquiries with archivists at the U.S. Department of Energy, the AEC's successor, I was assured that there were no references in that book or any other to Skoda, its purported SS think tank or German nuclear research at Pilsen or Dessau.

Given Voss' claims that nuclear propulsion work had been one of three primary areas of research within the *Kammlerstab*, alongside lasers and guided weapons, it is inherently credible that some such work had gone on there.

It is at odds, however, with the official view that Germany was nowhere near as advanced in nuclear weapons technology as the Americans, who by May 10 were less than two months away from detonating their first atomic bomb in the New Mexico desert.

According to Speer, official Nazi sanction for attempts by Werner Heisenberg, the father of the would-be German atomic bomb, were rescinded in 1942 on the grounds that the earliest timetable for an available weapon was 1946—too late, Speer concluded, to be of any use in the war.

In late 1944 and early 1945, U.S. technical intelligence agents from the Alsos Mission ("alsos," in a quirky piece of coding, being ancient Greek for "grove," i.e. General Leslie Groves, who was in charge of the U.S. bomb project) tore into France, Belgium and Germany searching for signs of German atomic bomb work and quantities of German-held uranium oxide ore—a material that is inert until it has been subjected to neutron bombardment in a nuclear reactor.

On April 23, an Alsos team led by the unit's commander, Lt. Col. Boris T. Pash, a former high school teacher turned Army G-2 security

officer trained by the FBI, arrived at the locked steel door of a box-like concrete entranceway in the side of a cliff above the picturesque German town of Haigerloch, near Stuttgart.

Pash, who had been in a race against the French to get there, shot off the lock and found himself in a chamber with a concrete pit about ten feet in diameter in the middle. Within the pit hung a heavy metal shield covering the top of a thick metal cylinder. The latter contained a pot-shaped vessel, also of heavy metal, about four feet below the floor level.

What Pash had discovered is what history describes as one of only two German reactors assembled, or partially assembled, before the end of the war—the other, at Kummingsdorf, being captured by the Russians.

Using heavy water as a moderator and 664 cubes of metallic uranium as its fuel, the Haigerloch reactor had achieved a sevenfold neutron multiplication just a few weeks earlier.

Heisenberg had calculated that a 50 percent increase in the size of the reactor would produce a sustained nuclear reaction.

Had he done so, Hitler would have had the means to enrich enough fissionable material to construct a bomb.

This is the official view of the extent of the German atomic bomb program and it is terrifying enough. The Nazis had come close to developing a weapon.

But look a little to the left and right of the official view, as I had before leaving London, and an even more frightening scenario pulled into view.

On May 19, a German U-boat, the U-234, docked at the U.S. Navy port of Portsmouth, New Hampshire, having surrendered to a U.S. Navy destroyer two days earlier off the eastern seaboard. Her commander had received high-level orders to sail from Kristiansand in Nazi-held Norway to Japan on April 16.

On board, the Americans found an Aladdin's Cave of technical equipment and blueprints, most of them related to advanced German jet aircraft. The U-234 was also ferrying technical experts, a nuclear technician among them.

This fact alone tipped off U.S. investigators that U-234 was no ordinary U-boat. A secondary examination showed she had been modified to carry a very dangerous cargo. Set in her six adapted mine-laying tubes were 560 kilos of uranium held in ten gold-lined containers. The loading manifest maintained that it was uranium oxide, the state in which uranium is found when it is extracted from the earth and safe enough to carry around in a paper bag. That the U-234's converted mine-laying tubes were gold-lined indicates that its cargo was emitting gamma

radiation. This, in turn, meant the uranium oxide ore had been subjected to enrichment from a working nuclear reactor. The gold lining was to prevent the radiation from penetrating the U-boat's hull.

And yet, officially, there had been no nuclear reactor in Germany capable of fulfilling this task.

Not in Speer's orbit of operations. But what about Kammler's?

The journey to this point had revealed the extent to which the SS had ring-fenced its top secret weapons activity from the rest of the German armaments industry. Could the *Kammlerstab* have contained the plans of a working nuclear reactor? Its secret location, even? It was a possibility I had to confront.

The lack of data in U.S. archives concerning the fate of the U-234, its cargo and its personnel is astonishing. But then, everywhere I'd looked prior to leaving London, a pattern had been emerging.

Protracted searches by archivists at the U.S. National Archives for any data on Kammler had failed to locate a single entry for him.

It was the same for Skoda.

Given Kammler's range of responsibilities in the final months of the war, this absence of evidence was remarkable; so much so, that one archivist at Modern Military Records, College Park, Maryland, said that it "redlined" him in her book.

I asked what she meant.

Somebody, she replied matter-of-factly, had been in and cleaned up.

Tom Agoston had found evidence of much the same thing. Despite Voss' extensive debriefings on Skoda by U.S. CIC agents at the U.S. European Command Counter-intelligence Center at Camp Oberursel, near Frankfurt, no one had been able to trace any records or transcripts of the interrogation.

When Agoston ran checks on Kammler, it transpired that his name hadn't even registered at the Nuremberg war crimes trials—the very least that might have been expected of someone who had played such a prominent role in the Holocaust.

Unlike Martin Bormann, whose body was never found after he was reported killed during his escape from the Führerbunker on May 1, Kammler was never tried *in absentia* at Nuremberg. There is no evidence that anyone ever bothered to look for him. This, despite the fact that the four versions of his death conflict with each other so markedly that none stacks up under scrutiny.

The first version says he committed suicide in a forest between Prague and Pilsen on May 9, two days after the German unconditional surrender.

The second says that he died in a hail of bullets as he emerged from the cellar of a bombed-out building to confront Czech partisans that same day.

The third version says that he shot himself in a wood near Karlsbad (Karlovy Vary in today's Czech Republic), also on May 9.

What these variations on a theme do manage to assert is that Kammler was in or around Prague just prior to the German capitulation.

One witness quoted by Agoston, an official with the Prague regional office of the Buildings and Works Division of the SS Economic Main Office, recalled the date: "Kammler arrived in Prague early May. We did not expect him. He gave no advance notice of his arrival. Nobody knew why he had come to Prague, when the Red Army was closing in."

I could see only two reasons why Kammler should have made such a journey. First to retrieve the mother lode of documentation on the special projects group at Skoda and the Skoda administrative offices in Prague; and secondly to bury it somewhere prior to setting up the deal that would save him from the gallows and buy him his freedom.

Where in the crumbling, collapsing edifice of the Reich would Kammler have considered hiding this priceless cargo? Certainly not in Czechoslovakia, not with the Russians closing in. And it was unlikely that he would have headed west through the partisan-infested forests between Prague and the German border.

The fourth version of his death, I felt, offered the best possible clue.

The German and Austrian Red Cross, according to Agoston, held two relevant documents after the end of the war, the first posting Kammler as "missing," the second updating the first and listing him "dead." The first report had been filed by a relative. It placed Kammler's last known location as Ebensee in Steiermark, Austria. The updated report was based on the testimony of unnamed "comrades." No burial place was specified.

I did some thinking of my own, using the reference books I had brought with me as pointers.

In early May, the province of Steiermark was putting up a vigorous defense against the Russians, while bracing itself for the American advance, under the leadership of Gauleiter Siegfried Ueberreither. The gauleiter—the Nazi party's chief administrator for the Steiermark "gau"—was backed in his decision to hold the province to the last by large numbers of Wehrmacht and SS troops.

Kammler knew Ebensee well. Under his orders and supervision, work had commenced there in 1943 on a giant underground complex for the intercontinental A-9/A-10 rocket—the weapon that Hitler had intended

to use against New York. The facility was code-named Zement and a large section of it had been completed by the end of the war. Zement was tucked into the mountains abutting the shoreline of Lake Traunsee.

On May 8, 1945, Lake Traunsee could be considered the epicenter of the Nazi empire on the eve of its surrender; a seething confluence of terrified citizenry, refugees from the east, heavily armed German troops and Nazi party members looking for places to lie low until the frenzy of capitulation had subsided. Gauleiter Ueberreither himself, having urged his people to fight to the last bullet, slipped out of his office in Graz and disappeared. Years later, he emerged in Argentina.

I began to make arrangements that would get me to Ebensee. But shortly before I could complete them, I received a call from Poland. A researcher there had responded to my request for information about German wartime research activities around Breslau, one of the two purported homes of German flying saucer research.

The guy's name was Igor Witkowksi and he had been recommended to me by Polish sources through my work at Jane's as someone who was both highly knowledgeable and reliable. Witkowsi, they told me, had self-published a series of books in Polish detailing his findings.

For years, Witkowsi had been sifting through archives related to the many secret Nazi scientific establishments in and around Breslau—modern-day Wroclaw—looking for traces of wartime secret weapons work. What he'd uncovered was evidence of a systematic SS-controlled R&D operation in a rat run of interconnected facilities, many of them underground, in the remote Sudeten Mountains in the southwest of Poland—Lower Silesia as it had been, before its handover to Poland under postwar reparations by Germany.

Witkowsi's research led him to conclude that a number of these facilities had been used by the SS for nuclear research work. But one site did not conform to the pattern. In this case, he said, a series of experiments had taken place in a mine in a valley close to the Czech frontier. They had begun in 1944 and carried on into the April of the following year, under the nose of the advancing Russians.

The experiments required large doses of electricity fed via thick cabling into a chamber hundreds of meters belowground. In this chamber, a bell-shaped device comprising two contra-rotating cylinders filled with mercury, or something like it, had emitted a strange pale blue light. A number of scientists who had been exposed to the device during these experiments suffered terrible side effects; five were said to have died as a result.

Word had it that the tests sought to investigate some kind of anti-

gravitational effect, Witkowsky said. He wasn't in a hurry to agree with this assessment, he added, but he was sufficiently intrigued by the data to alert me to the possibilities.

Like Witkowsky, I was skeptical, a feeling that grew when he told me who had presided over the work: Walther Gerlach, Professor of Physics at the University of Munich and head of nuclear research at the Reich Research Council in 1944-45.

Wilhelm Voss, joint head of the SS special projects groups inside the Skoda Works, had made it clear to Tom Agoston that German nuclear work had absorbed a large part of the group's research activities. By early May, Kammler should have had adequate intelligence that Pash, the FBI-trained former teacher and head of the U.S. Alsos Mission, had located most, if not all Germany's nuclear research centers in the western zone of influence along with the scientists that went with them. Kammler might also have known that the U-234 had sailed with its cargo of uranium ore for Japan; maybe he'd been responsible for the order himself.

While he could not have known that the Americans were close to perfecting their own nuclear weapon, or that the U-234 would fall into U.S. hands, he didn't seem like a man who would gamble his life on providing America with a technology it had already acquired via diligent detective work. It didn't tally with his profile.

My feeling was that Kammler would offer them something so spectacular they'd have no choice but to enter into negotiations with him; and, intriguing as it was, this experiment down the mine didn't seem to fit the bill. The sickness suffered by the scientists appeared to be the result of radiation poisoning. As for the device itself, I said, by now thinking out loud, since it fell within the Soviet zone of influence, it hardly conformed to the plan that Kammler had outlined to Speer in Berlin.

It was then that Witkowsky, who had listened for several minutes patiently enough, interrupted me. His research had uncovered the existence of an SS-run "Special Evacuation Kommando (command, team or unit)" that had evacuated the "Bell" and its supporting documentation prior to the Russians' overrunning the facility. The evidence said it had been shipped out, destination God only knew where; except it wasn't there when the Russians arrived.

Despite the warmth of my hotel room, I felt a sudden chill. For all the months I had been following Kammler, he had been a remote figure, even at his closest in the wet, freezing-cold underground galleries at Nordhausen. Now, fleetingly, I felt him in the same room as me.

"And the scientists?" I asked. "What happened to them?"

I thought the line had gone dead, but it was just Witkowski taking his time, choosing his words.

"They were taken out and shot by the SS between April 28 and May 4, 1945. Records show that there were 62 of them, many of them Germans. There were no survivors, but then that's hardly surprising."

He paused. "It's quite clear that someone had gone to very great lengths to clean up."

It was the second time in as many weeks that this expression had been used in the context of Kammler. Within the hour, I had booked myself on a flight to Warsaw.

It was thus—simply and irrevocably—that I was pulled into the extraordinary story of the Bell.

Chapter 19

We left the Polish capital in the haze of an overcast dawn, moving quickly past drab living blocks half-hidden by a frenzy of construction work.

Witkowski drove, apologizing in advance for the schedule ahead. The Wenceslas Mine was 50 kilometers or more the other side of Wrocław, formerly Breslau of the German *gau* of Lower Silesia, was around 350 km from Warsaw. The drive would be hard, he said, much of the route consisting of poorly maintained single-lane roads, but all being well we would reach our destination before sunset. In the meantime, he would bring me up to speed on his research, which had started with a phone call from a Polish government official.

Witkowski, a former defense journalist, did not want to tag his informant, but he assured me that his integrity was impeccable. The official had permitted him to view some documents, to make notes, but not to take anything away with him.

Witkowski had taken him at his word and transcribed most of what he had been allowed to view verbatim.

At the airport, his stocky appearance was just as I had imagined him. He was younger than I, with a thickset expression, high Slavic cheekbones and dark, intensely serious eyes. From the word go, we talked about little else but the research that united us. Had Witkowski been in any way a lightweight, I would have turned around and got on the first plane home. But when I saw him, I knew he was OK. There were aspects to him that I saw in myself. Like me, Witkowski was consumed by a thirst for knowledge. Knowledge of a highly specialized kind.

After months on the road it felt good to be sharing data again. And the very fact he'd let me in on his discovery told me that Witkowski felt the same way. We were walking a tightrope. If either of us were wrong about the data we'd collated, it'd be a long fall. On the phone, Witkowski had convinced me that he could take me to places that only he and a select group of local researchers knew about.

The papers that Witkowski had been allowed to transcribe detailed the

activities of a special unit of the Soviet secret intelligence service, the NKVD, forerunner of the KGB. Attached to this unit were two Polish officers, General Jakub Prawin, head of the Polish military mission in Berlin, and Colonel Wladyslaw Szymanski, a senior member of his staff.

It was during their "debriefing" of a former high-ranking official from the *Reichssicherheitshauptamt*, the Reich Central Security Office or RSHA, that Prawin and Szymanski learned of "General Plan 1945."

Their interrogation subject, Rudolf Schuster, had worked at the RSHA—one of four security organizations run by the SS-controlled *Sicherheitsdienst* or Security Service—until his deployment on June 4, 1944, to the Special Evacuation *Kommando*, an organization unknown to the NKVD, which prided itself on knowing everything. Schuster, it transpired, had been responsible for the ELF's transportation arrangements.

Schuster's information must have set alarm bells ringing at the highest Soviet levels, for Prawin and Szymanski had learned that the man behind General Plan 1945 was none other than Martin Bormann, Hitler's deputy, last seen fleeing the Führerbunker on the night of May 1. Though supposedly killed in the attempted breakout, Bormann's body had never been formally identified. Tying Bormann into a scheme for some kind of evacuation procedure, therefore, had disturbing implications, but pumping Schuster was evidently pointless as it was clear he was only privy to the transportation details. The SS, operating its traditional compartmentalization system of security, had kept him from the broader aspects of the plan. It was apparent, however, that much of the *Kommando*[^] activities centered on the territory of Lower Silesia, now southwest Poland.

The next piece of the jigsaw was provided by SS-Obergruppenführer Jakob Sporrenberg, who from June 28, 1944 had commanded a section of the *Kommando* attached to the Gauleiter of Lower Silesia, Karl Hanke. Hanke, like all gauleiters, who were the party's representatives in the gaus or regions, was answerable directly to Bormann, the party chief.

Sporrenberg had been captured by the British in May 1945, but was handed over to the Poles because he was a general in the SS and had been active in Poland, the scene of many of the SS-run death camps. Had they realized Sporrenberg's true brief, it is unlikely they would have been quite so quick to release him.

Sporrenberg was a big shot, one of a scattering of senior SS police generals of exalted rank and status positioned to give Himmler absolute power throughout Germany in the closing years of the war. To put his seniority in perspective, he carried the same rank as Ernst Kaltenbrunner, the brutal Austrian who succeeded Heydrich as head of the

Reich Security Office, the Nazis' central agency for internal counter-espionage and repression. For Sporrenberg to have been assigned the command of the Special Evacuation *Kommando* unit attached specifically to Hanke's gau of Lower Silesia shows how important Bormann's evacuation plan was considered to be.

And, I thought, just how secret. There has never been any official acknowledgment of the existence of the Special Evacuation *Kommando*.

Exactly what, I asked Witkowski, was the *Kommando* tasked with evacuating?

The taciturn Pole, concentrating on the road ahead, flashed me a look. Patience. Everything in its own time.

Sporrenberg, he continued, was sent for trial and sentenced to death in 1952, but not before he had testified in secret before the Polish courts about his role as one of Lower Silesia's main plenipotentiaries for the evacuation of high-grade technology, documents and personnel or his part in the murder of 62 scientists and lab workers associated with a top secret SS-run project in a mine near Ludwigsdorf, a village in the hills southeast of Waldenburg, close to the Czech border.

Sporrenberg, operating under the political oversight of Gauleiter Hanke, had been in charge of a *Kommando* cell tasked with "northern route" evacuations via Norway, which remained in German hands until the very end of the war.

The NKVD/Polish intelligence team learned that SS-Obersturmbannführer Otto Neumann, who had commanded a *Kommando* detachment in Breslau, had been responsible for all southbound shipments to Spain and South America.

Neumann was never caught, but was reputed to have escaped to Rhodesia, where he was supposedly sighted after the war.

According to some estimates, Witkowski said, the air bridge established by the *Kommando*'s southern command between the Reich's remaining occupied territories and neutral, but Axis-sympathetic Spain managed to evacuate 12,000 tons of high-tech equipment and documentation in the final months of the war using any commandeered Luftwaffe transport it could lay its hands on.

The *Kommando* southern command had one other exit route available to it up to the end of the war—albeit an even more hazardous one. Using some of the northern Adriatic ports that remained in German hands up to the surrender, a brave or foolish U-boat commander could conceivably have run the gauntlet of Allied air and maritime superiority to evacuate cargo and personnel by sea.

But it was Sporrenberg's northern route operations that underpinned

the reason for the trip, because it was Sporrenberg who provided the only details that have ever come to light on the Bell.

The experiment started out at a top secret SS-run facility near Leubus (Lubiaz in modern-day Poland), northwest of Breslau, in early-mid 1944. With the Soviets' rapid push into Poland during late 1944/early 1945, the unit was transferred to a castle on a hill above the ancient village of Fuerstenstein (Ksiaz), 45 kilometers to the south, close to the Lower Silesian coal-mining center of Waldenburg. From there, it was moved again to the mine near Ludwigsdorf (Ludwikowice), 20 kilometers the other side of Waldenburg, nestling in the northern reaches of the Sudeten Mountains. The Wenceslas Mine, where the Bell ended up, had been requisitioned by the SS as part of a neighboring underground weapons complex, code-named *Riese*—"Giant."

Riese, only part-completed by the end of the war, was an attempt to transform an entire mountain into an underground weapons production center. The many tens of kilometers of galleries that had been tunneled by the end of the war had been clawed from the rock by inmates drafted in from the nearby concentration camp of Gross-Rosen. Modern excavations of *Riese* show that the SS had been attempting to link it to the Wenceslas mine via a tunnel almost ten kilometers long.

By mid-afternoon, Witkowski and I had reached Wroclaw, a smoky industrial city on the banks of the River Oder that in 1945, as the "fortress" city of Breslau, had held out against the Red Army for more than 70 days, despite total encirclement.

In September 1944, long before the Russians were at Breslau's gates, the German military commander, General Krause, proposed the evacuation of 200,000 civilians, but Gauleiter Hanke refused, seeing it as a sign of weakness.

Four months later, as the Soviet ring of steel closed around the city, Hanke finally relented and women and children were permitted to leave.

But in the depths of a winter that was cold enough to freeze the sweeping waters of the Oder, those that weren't cut down, killed or raped by the Russians succumbed in their thousands to the subzero temperatures.

In February, the troops of SS Fortress Battalion "Besselein" launched a ferocious counterattack in an attempt to break the siege. While the German troops met the Russians head-on in some of the heaviest street-fighting since Stalingrad, Hanke exhorted his citizens to victory under the slogan "Every house a fortress." He then ordered a ruthless conscription of manpower that produced five regiments of Volkssturm—old men beyond normal fighting age—and Hitler Youth.

Following the briefest of training regimes, principally in the use of antitank "panzerfausts," they went out and died in the thousands.

We edged forward in the heavy traffic, soon finding ourselves on a long wide thoroughfare, the former Kaiserstrasse, which by April 1945 had been transformed into a landing strip for medium transport aircraft after the Russians captured the airport. It signaled the beginning of the end, but still the defenders fought on.

German Red Cross flights of Junkers Ju 52s removed 6,000 of the more seriously wounded and beneath the dense overcast that merged with the factory smoke a half century later, I could almost hear the rumble of trimotors as they swept into land, touching down at night under the guiding light of flame torches, and the crump-wump of artillery explosions in the suburbs.

By this stage, the defenders—soldiers and civilians alike—were forced to battle the Russians on all conceivable fronts, including the sewers beneath the streets.

On May 4, two days after the Russians took Berlin, a delegation of churchmen begged General Niehoff, Krause's replacement, to surrender. Two days later, he agreed and on May 7 the Red Army marched in.

Furious at having been held at bay by such a vastly inferior force, the Soviets launched heavy reprisals against the soldiers and citizenry of Breslau; a fate that Gauleiter Hanke chose not to share. He had flown out of the city on May 4 in a Fieseier Storch liaison aircraft to take up his post as successor to the disgraced Himmler, whose surrender negotiations had been uncovered by Hitler shortly before his suicide.

Hanke, local administrator of Bormann's enigmatic General Plan 1945, the man to whom the Breslau-based detachment of the Special Evacuation *Kommando* was directly accountable, was never seen again.

When I told Witkowski about Kammler, how he too had vanished off the face of the earth, he was unmoved. It was simply part of a pattern, he told me. Hanke, Kammler . . . there were dozens of high-ranking former SS or Party members that had never been called to account. They had simply disappeared. Many of them shared the distinction of having had access to highly advanced technology.

In the immediate aftermath of his death sentence, Sporrenberg, head of the Special Evacuation unit's Breslau operations, was smuggled to Russia, milked by the NKVD in one last brutal round of interrogation, then disposed of. Schuster, in charge of the unit's transportation arrangements, died "mysteriously" in 1947 and Szymanski and Prawin, the two Polish intelligence officers who'd interrogated them, also met

untimely ends after their release from the special NKVD investigation cell—Szymanski in an air crash and Prawin in a drowning accident.

"What was it that killed them?" I asked. "Their knowledge of the Special Evacuation *Kommando* or the fact they knew about the mine?"

Witkowski kept his eyes on the road. "The Germans maintained a special detachment of aircraft, probably Ju 290s, and a single Ju 390, both rare heavy transport types, based at Opeln—Opole today—a hundred kilometers from here. Witnesses say the planes were well camouflaged and that some were painted with yellow and blue identification markings, which suggests they were masquerading as Swedish planes. If so, it was almost certainly a unit of KG 200, the Luftwaffe special operations wing, which frequently flew aircraft under the flag of enemy or neutral countries. The point is, the Evacuation *Kommando* had the ability to move thousands of tons of documentation, equipment and personnel, and they could move it north or south. Sporrenberg's brief was to go north. The whole operation was highly secret. But nothing, it seems, was more secret than the Bell."

We crawled out of Wroclaw in a heavy convoy of trucks bound for the Czech border. I focused on a faint ribbon of sky between the clouds gathering to the south and the outline of the Sudeten Mountains, the inner core of Kammler's high-tech kingdom, indistinct yet somehow threatening, on the distant horizon.

Chapter 20

In his interrogation sessions with the NKVD and his deposition to the Polish courts, SS General Jakob Sporrenberg could only tell the interrogators and the prosecutors so much, Witkowski told me. Sporrenberg was a policeman and an administrator, not a technician. He could relay what he had seen and heard, little snippets he had picked up, but the technology itself was beyond him. He had been tasked with getting men, materiel and documents out of Lower Silesia before the Russians reached them; he had had no need to know much else. This was standard SS operating procedure.

I asked what else he knew about Sporrenberg.

Witkowski stared straight ahead as he talked. First, I had to understand that Sporrenberg was very high up the chain of command. Records showed, he said, that Sporrenberg was appointed deputy commander of the Waffen-SS VI Korps under SS-Obergruppenführer Walter Krueger in 1944. Krueger, as far as Witkowski had been able to ascertain, had been involved in a series of top secret SS-conceived operations in the closing months of the war, including a plan to occupy neutral Sweden, the evacuation of Nazi wealth to South America and other neutral or nonaligned states, a harebrained scheme to strike at New York with V-1s launched by submarine and, of course, the special evacuation of Nazi secret weapons and high technology. Sporrenberg was appointed plenipotentiary in charge of northern-route "special evacuations" to Norway on June 28, 1944. Like Kammler, it seemed that General Sporrenberg was recognized and rewarded for his ability to organize, not for his ability to fight.

Following his capture, as much as Sporrenberg was able to divulge to Soviet intelligence and the Polish courts about the Bell was this, Witkowski said. The project had gone under two code names: "*Laternestrager*" and "*Chroms*" and always involved "*Die Glocke*"—the bell-shaped object that had glowed when under test. The Bell itself was made out of a hard, heavy metal and was filled with a mercury-like substance,

violet in color. This metallic liquid was stored in a tall thin thermos flask a meter high encased in lead three centimeters thick.

The experiments always took place under a thick ceramic cover and involved the rapid spinning of two cylinders in opposite directions. The mercury-like substance was code-named "Xerum 525." Other substances used included thorium and beryllium peroxides, code-named *Leichtmetall*.

The chamber in which the experiments took place was situated in a gallery deep belowground. It had a floor area of approximately 30 square meters and its walls were covered with ceramic tiles with an overlay of thick rubber matting. After approximately ten tests, the room was dismantled and its component parts destroyed. Only the Bell itself was preserved. The rubber mats were replaced every two to three experiments and were disposed of in a special furnace.

Each test lasted for approximately one minute. During this period, while the Bell emitted its pale blue glow, personnel were kept 150 to 200 meters from it. Electrical equipment anywhere within this radius would usually short-circuit or break down. Afterward, the room was doused for up to 45 minutes with a liquid that appeared to be brine. The men who performed this task were concentration camp prisoners from Gross-Rosen.

During the tests, the scientists placed various types of plants, animals and animal tissues in the Bell's sphere of influence. In the initial test period from November to December 1944, almost all the samples were destroyed. A crystalline substance formed within the tissues, destroying them from the inside; liquids, including blood, gelled and separated into clearly distilled fractions.

Plants exposed to the Bell included mosses, ferns, fungi and molds; animal tissues included egg white, blood, meat and milk; the animals themselves ranged from insects and snails to lizards, frogs, mice and rats.

With the plants, chlorophyll was observed to decompose or disappear, turning the plants white four to five hours after the experiment. Within eight to fourteen hours, rapid decay set in, but it differed from normal decomposition in that there was no accompanying smell. By the end of this period, the plants had usually decomposed into a substance that had the consistency of axle grease.

In a second series of experiments that started in January 1945, the damage to the test subjects was reduced to around 12—15 percent following certain modifications to the equipment. This was reduced to two to three percent after a second set of refinements. People exposed to the program complained of ailments, in spite of their protective clothing. These ranged from sleep problems, loss of memory and balance, muscle

spasms and a permanent and unpleasant metallic taste in the mouth. The first team was said to have been disbanded as a result of the deaths of five of the seven scientists involved.

This, Witkowski said, was all contained in the documents he had been shown. What impressed him—and what I now confessed had me intrigued as well—was their level of apparent detail. It told him—and me—that in all likelihood *something* had happened down the mine; something mysterious. But what? Witkowski insisted that it had all the hallmarks of an antigravity experiment. But I wasn't so sure. Before I could begin to believe this was the Holy Grail, I needed to check off a set of more rational explanations first. And the most glaring of these was the probability that this had been some kind of test involving nuclear material. The documents that Witkowski had seen had mentioned the involvement of Professor Walther Gerlach, the man charged with oversight of Germany's atomic weapons programs. Disturbingly, they also cited Dr. Ernst Grawitz, head of the euphemistically labeled SS Medical Service. Grawitz had been the boss of Josef Mengele, the infamous doctor of Auschwitz. Inevitably, there were reports that the Bell had been tested on humans as well. Again, if the Bell had emitted radiation, the Nazis—given their record on war crimes in this area—would have probably monitored its effects.

I said nothing, however, out of deference to Witkowski, who told me that much of the rest of his evidence was visible at the Wenceslas Mine itself.

As we approached Waldenburg, Walbrzych as it is today, the landscape altered. After nothing but lowlands and flatlands all day, the road now started to twist and climb as we headed into the foothills of the Sudeten range.

Waldenburg itself was bleak, its imposing mix of late nineteenth- and early twentieth-century architecture, Germanic to the rooftops, blackened with coal dust and beset by decay. Though it had escaped the ravages of the war, the Soviet advance passing it by to the north and the south, it had been less fortunate in the wake of the communist collapse. By the early 1990s, almost all mining operations in the region had been shut down, transforming it overnight from one of Poland's most prosperous areas to its poorest.

Men with nothing to do gathered on street corners and intersections and passed bottles of cheap vodka among one another, eyeing us suspiciously as we swept by.

On the narrow road leading out of Waldenburg to Ludwigsdorf, I asked Witkowski what had led him to believe that the Bell experiments had been an attempt by the Nazis to manipulate gravity.

There were scant clues, he admitted, but those that Sporrenberg had provided in his testimony seemed to add up to something. Many of the descriptions used by the Bell scientists did not gel with any of the accepted terms associated with nuclear physics, nor were there any obvious radioactive materials used in the experiments themselves. One of the terms Sporrenberg had picked up had been "vortex compression"; another was "magnetic fields separation." These were physical principles that had come to be associated with the new wave of gravity and anti-gravity pioneers—people like Dr. Evgeny Podkletnov, Witkowski said.

Podkletnov. Now, I began to pay more attention. If I'd interpreted Witkowski correctly, there was some kind of a relationship between the Russian's experiments with spinning superconductors—the effect he had tripped over in Finland when his assistant's pipe smoke had hit that column of gravity-shielded air—and the effect produced by the Bell.

I needed to call Marckus and run some of this stuff past him; if, that is, he was still talking to me after my refusal to dig deeper in America.

Witkowski also claimed there were anomalies in the curriculum vitae of Professor Walther Gerlach that placed him firmly in the orbit of the gravity scientists, despite the fact that, ostensibly, his discipline was nuclear physics. In the '20s and '30s, Witkowski discovered, Gerlach had immersed himself in phenomena such as "spin polarization," "spin resonance" and the properties of magnetic fields—areas that had little to do with the physics of the bomb, but much to do with the enigmatic properties of gravity.

A student of Gerlach's at Munich, O.G. Hilgenberg, published a paper in 1931 entitled "About Gravitation, Vortices and Waves in Rotating Media"—putting him in the same ballpark as Podkletnov and the Bell. And yet, after the war, Gerlach, who died in 1979, apparently never returned to these subject matters, nor did he make any references to them; almost as if he had been forbidden to do so.

"The Germans ignored Einstein and developed an approach to gravity based on quantum theory," Witkowski said. "Don't forget that Einsteinian physics, relativity physics, with its big-picture view of the universe, represented Jewish science to the Nazis. Germany was where quantum mechanics was born. The Germans were looking at gravity from a different perspective to everyone else. Maybe it gave them answers to things the pro-relativity scientists hadn't even thought of."

Kammller, Witkowski told me, had the ability to vacuum up all scientific activity, whether it was theoretical or practical, through an SS-run organization called the FEP, for *Forschungen, Entwicklungen und Patente*—researches, developments and patents. The FEP introduced

the last major player in the story of the Bell and the Special Evacuation *Kommando*, an SS-Obergruppenführer called Mazuw. A high-ranking general, Emil Mazuw had been able to acquire any significant technology, science theory or patent application that had come to the attention of the SS—and via its prolific security arms, there wasn't much that passed it by. The FEP, according to Witkowski's researches, operated independently of the *Reichsforschungsrat*, the Reich Research Council, but would unquestionably have had oversight of it.

My guess was that the FEP had been administered by Kammler's secret research cell within the Skoda Works. After the war, the Allied powers seized 340,000 German-held patents from the captured records of the Reich Research Council.

But here was something else: the cream of Germany's wartime scientific research that had been skimmed off by the SS and compartmentalized for its own use. Its value would have been immeasurable. Perhaps this really was what Kammler had returned to Czechoslovakia for in the death throes of the Reich?

There were other clues, too, but these, Witkowski repeated, were the physical traces at the mine itself and were best explained when we got there.

At Ludwigsdorf we followed the line of an old railway that had been built before the war to connect the region's network of industries to the outside world. As the road climbed, we found ourselves in a valley bordered on both sides by tall trees. A patchy mist clung to the upper branches as we ascended into the hills.

Then we rounded a bend and the scenery opened up. We were in a valley, one that had appeared out of nowhere. Witkowski parked on a patch of gravel overlooking some level ground. The railway track, overgrown from years of disuse, followed its median line, eventually disappearing from view behind a large derelict building whose tall arched windows rose cathedral-like into the mist. It was next to this building, Witkowski said, that the shaft of the mine disappeared belowground.

I stood on the edge of a steep bank a short walk from the car. The valley was around 300 yards across, the trees either side of it so dense you'd never know the place existed unless you happened upon it.

The bank, which was 20 feet high, was mirrored by an identical feature on the opposite side. During the war, this expanse of land had been an underground marshaling yard, Witkowski told me, thick wooden planks called sleepers topped with turf hiding a six-lane section of track where the railhead met the workings of the mine.

Even the buildings disguised the facility's true purpose.

High on the opposite bank was a large redbrick house, an original 19th-century feature of the Wenceslas Mine. But look a little closer, Witkowski said, and you could see where the Germans had tunneled into the ground underneath and constructed a large concrete bunker, one of the many blockhouses scattered around the complex. All of the bunkers had been covered with earth and trees planted on top.

In short, the Germans had gone to a great deal of trouble to ensure that the place looked pretty much as it had always looked since mining operations began here at the turn of the last century, a clear indication that whatever had happened here during the war had been deeply secret. Now, half a century after the Germans had left, second-guessing the use to which it had been put, simply from the available physical evidence, was far from easy.

When, immediately after the war, the Poles repatriated the local German population, substituting German place-names for Polish ones in the process, they removed any witnesses who could have shed light on the SS' activities. Almost everything that was known about the Wenceslas Mine had been handed down from Sporrenberg.

It had been run by the SS, had employed slave labor and had been sealed from the outside world by a triple ring of checkpoints and heavily armed guards.

Beyond that, Witkowski said, we were into the realm of analysis, interpretation and detective work; the skills required for such a task being more akin to those of an archaeologist.

Although the locals had used the site for small-scale industrial activity in the years since the war—I could see freshly milled lumber stacked on a flat piece of land over by the valley entrance—it had been entirely neglected by the history books; its wartime use known only to the handful of people Witkowski had brought here.

That, I could understand. The place was as far off the beaten track as I'd ever been on the continent of Europe. I felt like I was tramping on ground that had hardly been touched since the Germans had pulled out more than half a century earlier.

The first thing to point out, Witkowski said, was the structure at the end of the valley. This, built before the war, was a power station capable of burning a thousand tons of coal a day—enough to provide the valley with vast quantities of electricity. It had more than likely been this single feature that had drawn the SS to the site, Witkowski said, since it had helped to make the place entirely self-sufficient. After the war, when the Russians eventually made it into these hills, they would have found an

abandoned complex given over to some quasi-military purpose, a mine shaft that had been flooded, possibly deliberately, and little more.

"With the help of Sporrenberg's testimony, we have an edge that was denied to the first Red Army units to enter this place," Witkowski said. "Come, I want to show you something; something very strange."

We clambered back into the car and made our way down the track that led past the abandoned power station. It was now dark enough for Witkowski to use his lights, the beams catching in the weeds that tumbled through the cracks in the prefab concrete roadway laid by the Germans during the war.

Whatever the SS had been moving around here, it had been pretty heavy-duty, Witkowski remarked.

The track cut through a wood, emerging in a clearing on the far side of the power station. Caught in the headlights, rising out of the ground straight ahead, was a circular concrete construction 30 meters wide and 10 high. With its 12-meter-thick columns and horizontal beams, it was part-reminiscent of some ritual pagan edifice.

Witkowski parked the car, but left the headlights on, the beams bathing the columns of the object in bright white light.

"What is that thing?" I asked.

But Witkowski was already out of the door, crouching over something a few feet from the car. He was studying what looked like a partly exposed underground drain. Its concrete cover had cracked to reveal a duct about a foot across.

"This carried the electricity cable from the power station," he said. "It disappears into the ground just beyond the car, but diverts via this thing."

"What is it?" I asked again.

"I am not sure. But whatever it is—whatever it *was*—I believe the Germans managed to complete it. In this light it is difficult to see, but some of the original green paint remains. You do not camouflage something that is half-finished. It makes no sense." He paused. "There is something else. The ground within the structure has been excavated to a depth of a meter and lined with the same ceramic tiles that Sporrenberg describes in the chamber that contained the Bell. There are also high-strength steel hooks set into the tops of the columns. I think they were put there to support something; to *attach* to something. Something that must have exerted a lot of power."

I looked at him. "What are you saying?"

He took a moment before replying. "I'm saying I think it's a test rig. A test rig for a vehicle or an engine of some kind. A very powerful one."

That night I tried to call Marckus on my cell phone, but the isolated farmhouse where we were staying was buried in a deep valley, every bit as remote as the one that housed the nearby Wenceslas Mine complex, and I couldn't get a signal.

The Polish couple who owned the small farm let out a room to Witkowski whenever he came south from Warsaw to visit. They shared his curiosity about the mine and the neighboring rat run of uncompleted tunnels that were part of the SS-run "Giant" underground weapons complex. For all they or anyone else knew, the tunnels extended under the ground I was standing on. But being only second-generation inhabitants of this region, their parents having been shipped in to replace the Germans who'd lived here for three hundred years previously, they knew as little about the history of the place as Witkowski and I did. This, undoubtedly, helped to give the place a detached, soulless air.

There was no anecdotal evidence to fall back on; the only evidence was history's imprint on the topography itself.

I needed Marckus. I needed to run every last detail of the mine past him to see if it rang any bells in that weird analytical mind of his. I needed his guiding expertise again. Without some input from a physicist, the data on the Bell was nothing more than a bunch of science mumbo jumbo.

As I collected my thoughts under the swath of clear, starlit sky between the ridgelines, I elected to concentrate on a matter I did know something about; something that had been nagging at me ever since Witkowski had mentioned it: the matter of the KG 200 unit at Opeln, the base that had operated the Ju 290 and 390 transport aircraft on detachment to the *Kommando*. It was a highly significant development.

The Ju 290 was quite a rare bird; the 390 even more so. The 290 had been a big four-engined aircraft designed for the maritime reconnaissance, transport and bombing roles. The first transports had been used in the Stalingrad air bridge in 1942, but toward the end of the war, the final variants were modified for extreme-long-range operations. Three such aircraft are known to have made flights to Japanese bases in Manchuria. The Ju 390 was a six-engined modification of the Ju 290 with even more impressive operating characteristics. Though only two prototypes were built, they clearly demonstrated the Ju 390's ability to mount ultra-long-range operations of up to 32 hours endurance. On one occasion, a Ju 390 flew to within 12 miles of New York and back again. It could also carry a very heavy payload.

Above all, though, the Luftwaffe had referred to these aircraft as "trucks."

A book in Witkowski's possession, *The Nuclear Axis* by Philip Henshall, had provided this one essential detail. Henshall had picked up on Tom Agoston's story of Himmler's request for a "truck." But Henshall had arrived at a markedly different interpretation of the data. When Kammler had telexed Himmler on April 17 from his Munich office—the last signal anyone ever received from him—refusing the Reichsführer-SS the use of a "heavy truck" from the Junkers "motor pool," he hadn't been referring to a truck in the conventional sense. He had been referring to an aircraft. A long-range one with a heavy payload.

Kammler had been telling Himmler, his superior, that he couldn't have a Junkers 290 or 390 for his own use, because it was committed elsewhere. It could only have been for use by the Special Evacuation Kommando. Why else stamp such an ostensibly bland, innocent message *geheim*—secret? Why else would Himmler have been bothering himself with trucks?

With Bormann, Hitler's deputy, in charge of the evacuation plan, Kammler would have been in a unique position to call the shots. Although Hitler viewed Himmler as a valued and trusted ally—certainly up until the moment his secret surrender negotiations were discovered—it was Bormann, Hitler's gray eminence, who had his ear.

Kammler could safely refuse Himmler permission to use a Ju 290 or 390 without fear of recrimination. With such aircraft at its disposal, the Kommando could have flown its cargo of documents, personnel and technology pretty much anywhere it wanted. Spain, South America—Argentina, even—would have represented no problem to such a long-range platform.

The realization pulled me up short. What was the point of chasing Kammler, if he'd already shipped everything out?

After dinner, a good but simple meal of boiled eggs and local ham, I wandered back outside to take the air, leaving Witkowski inside to talk to our hosts.

I gazed at the ridgelines again. A few miles away, lay the Wenceslas Mine.

Part of me was transported back to the desert around Groom Lake, to the time when I'd stood vigil over the hills that shielded Area 51 from the outside world.

The part-completed Giant underground manufacturing complex and the Wenceslas Mine were a lot closer than I'd ever been able to get to the 27,000-feet-long paved runway on the desert lakebed. But the feeling they evoked in me was curiously similar.

The SS had installed a security system around the mine every bit as

impenetrable as the cordon around Groom Lake. That, undoubtedly, was a part of it.

As was the fact that the technologies at the heart of both places were essentially a mystery to everyone but the people who'd worked on them.

It was not knowing what lay beyond those ridgelines that made me do this. It was the not knowing that drove me on.

All these sensations contributed to the almost tangible atmosphere that imbued both places.

I was consumed with the need to go on until the picture steadied.

I knew then that I was tired—more tired than I'd ever been. I had got to the point where my every living minute was filled with a need to know the truth of something this insoluble, and that it had got ahold of me so badly that my subconscious mind was trying to solve it in my sleep, failing in the process and turning in on itself, twisting raw thoughts and images into the worst kind of dreams. Here, in the depths of Kammler's kingdom, a thousand miles from the warm archives of the Imperial War Museum and the Public Records Office, the scale of the Holocaust had begun to cling to me like a second skin.

Amidst it all, though, a particular sequence of images had stuck in my mind since my visit to the mine. Try as I might to stop it, I couldn't. I saw it with my eyes open and I saw it with them shut. In the chill air between the ridgelines, I saw it now.

It started with the arrival of a convoy of Opel trucks a week before the end of the war.

The scientists would have had mixed feelings about leaving: a sense of regret at having to abandon the project; relief to be escaping the oppressive feeling of the place; joy at the knowledge they were getting out ahead of the Russians.

A late spring day, the guns a long way off, the scientists talking and smoking in threadbare suits, their bundled possessions to hand, as the convoy was readied for departure. Sixty-two scientists meant three trucks' worth; perhaps another five to transport the crates. Eight trucks in all, moving west, gears grinding on the steep roads through the mountains. A disciplined unit operating to a fixed schedule, the details drawn up by Sporrenberg's bureaucrats in Breslau.

Two kinds of troops attached to the unit: drivers and logistics personnel to oversee the evacuation procedure and an armed *sonder-kommando* group acting as escort.

To these special-action troops, the scientists were simply "subjects." The trick was to maintain their docility and compliance until the last possible moment.

After multiple "operations" in the ghettos of Lodz, Warsaw, Minsk and the other occupied towns and cities of Central and Eastern Europe, the SS had found that by sticking to a preprepared book, the subjects could be kept as passive as cattle.

For some of the scientists, there may have been a tremor of foreboding as the three trucks peeled off from the main body of the convoy and headed into the forest.

But for most it would only have hit them when the tailgates dropped away exposing a line of troops with SS runes on their collars, their guns raised.

The Waffen-SS would have outnumbered the subjects two to one, their crushing shock and bewilderment combining with the SS's weight of numbers to make them respond without question to the barked orders as they moved toward the edge of the ditch. At no point, special-action orders decreed, should subjects feel that the executioners offered the remotest kind of hope.

And so, it had come to this. Two shifts, 31 bullets per shift, each shot three, maybe four seconds apart. The first victims facedown in the dirt for each member of the second shift to witness in a terrible moment of finality as he stepped up to the ditch, their grave, the proportions worked out in advance in some warm administrative office a long way from the front.

Sixty-two men meant a ditch ten meters long, two meters wide and one meter deep, the bodies stacked two high.

If the executioner tasked with placing his pistol at the base of each head observed the guidelines laid down for his benefit, the act of pulling the trigger was calculated to be no more arduous than switching off an electric light.

When the trucks left, the only thing to denote the scene of the crime would have been a mound of freshly dug earth half a meter high.

By the following spring, the mound would have collapsed to the exact same level as the ground around it. This, too, would have been carefully calculated according to the known decay rates of human tissue and the ambient conditions of the ditch.

To preserve the secret of the mine, the SS had consulted not even a page, but a paragraph or two from the execution manual it had drawn up for the Holocaust. It had tapped on the crime of the century to do this to a handful of its own people.

Though none of this was proven fact, remember this, I told myself, should you ever be tempted to view this as a technology hunt in a remote place that has little or nothing to do with you.

Chapter 21

The next day, I was back into Germany, on my way to Austria, making good progress on the autobahns in the rental car I'd picked up across the border, when the phone rang. It was Marckus. Hearing his voice, I felt a wave of relief, the cellular link between us suddenly feeling like a lifeline.

I spent the next 20 minutes briefing him on the mine. To begin with, he put up some resistance, trotting out his old belief that it was pointless pursuing the Germans for anything remotely as advanced as antigravity technology. But I had a feeling that Marckus was simply going through the motions.

The longer I talked, the more he listened. By the end, he was firing the questions at me.

I told him I had a hunch; that the Wenceslas Mine complex had been some kind of nuclear installation, the strange, henge-like construction being redolent of a reactor housing, with a tile-lined coolant pool in its midst. I didn't buy Witkowski's test-rig thesis, but then again I wasn't dismissing it either.

As for the Bell, I had absolutely no idea what it could have been, but without my notes at hand, I couldn't give Marckus the kind of details he required to conduct an in-depth analysis.

I promised to send him an email with the facts that night, as soon as I got to my hotel in Bad Ischl, across the Austrian border.

"What happens in Bad Ischl?" Marckus asked.

I told him about the Junkers "truck" business and how it had altered my plans. With long-range transport aircraft at his disposal, Kammler had had the means to ship the core secrets of the Skoda Works anywhere he wanted. And for all I knew, he had gone with them. It was pointless trying to pick up the clues of his disappearance in Ebensee or anywhere else for the time being. It was enough to know that the Nazis had, via Kammler's special projects group, a repository of technical secrets that had gone way beyond the V-weapons they had developed and used by the end of the war.

It had taught me other things, too. For years, I had read of the SS'

involvement in exotic weapons technology, but dismissed it as inadmissible claptrap peddled by people with an unhealthy interest in its warped ritualistic ideology; this, despite Speer's solid documentation of Himmler's penchant for unconventional scientific solutions to Germany's dire military position in the closing years of the war. In *Infiltration*, Speer's account of how the SS eventually succeeded in establishing its own industrial empire within the visible economic framework of the Reich, he had listed some of Himmler's more absurd ideas: producing fuel from fir-tree roots, tapping the exhaust fumes of bakery chimneys for the manufacture of alcohol and producing oil in abundance from geraniums.

Speer also attests to Himmler's willingness to entertain any radical ideas for new weapons—especially those that leaped the current state of the art. One such was a proposal for turning the upper atmosphere into a giant high-voltage conductor, presumably for the purpose of frying the Allies' B-17, B-24 and Lancaster bombers as they flew into Reich airspace. These ideas, many of which were based on unsound science, came to naught, but it would have taken only one far-reaching proposal underpinned by some solid scientific reasoning for this vast array of funded research activity to deliver the payoff so desperately sought by the Nazi leadership.

Like lasers, for example. Or an atomic bomb.

It had also taught me just how effective—and this word, of course, glosses over the grotesque cruelty of its methods—the SS security machine had been.

The *Kammlerstab* had been protected by a triple ring of counter-intelligence agents to prevent word of its activities from leaking. If it hadn't been for Voss and Agoston, the existence of the special projects group at Skoda might never have surfaced at all.

It had been the same story at the Wenceslas Mine. A combination of active security measures and a neighboring populace that had been rounded up and removed *en masse*, and its wartime activities had all but disappeared from existence.

But there was an even more powerful reason why none of this had ever properly surfaced before.

By involving the concentration camps, the SS had unwittingly set the seal on any serious postwar investigation of the science and technology it had pursued during the conflict. Because it had been the SS, not so much Speer's Armaments Ministry, that had backed so-called high-payoff, "visionary" projects with funding, German industry found itself in league up to its collective neck with the perpetrators of the Holocaust.

Many of the companies that had benefited from SS money, and employed concentration camp labor in the process, are still in existence today.

Bosch, Siemens, Zeiss and AEG all maintained highly secretive research and development operations in Lower Silesia.

It remains in their interests for their activities at this time to stay largely undocumented. I doubted, in any case, even if they were to open up their archives, whether anything of real value remained in them. Like the records pertaining to Kammler's past, I suspected they'd long since been cleaned out.

This, I said to Marckus, was how secrets came to be locked away, buried for all time, leaving nothing but myths and legends in their aftermath.

Maybe the Nazis had initiated a flying saucer program; maybe some of the technology had borne fruit. There was certainly enough anecdotal evidence to support the view that a variety of disc-shaped craft had flown before the end of the war. It was simply that there was *no proof*.

My journey into Kammler's kingdom had opened my mind to possibilities I wouldn't have begun to entertain a few months earlier. And in the process of this willingness at least to confront new ideas, I had found myself drawn to Bad Ischl.

It was there, before leaving London, that I had arranged to meet with the family of Viktor Schuberger.

I had picked up on the Schuberger story via the Legend forwarded to me by my friend and former colleague, Lawrence Cross, in Australia. I had initially dismissed it out of hand because not being an engineer but a simple forester, what this Austrian inventor was said to have achieved technically hadn't made any sense. And the phone conversation I had had with his son Walter, who back in 1991 had urged me to visit the family-run "bio-technical institute" in the Salzkammergut mountains so I could make up my own mind, had only served to alienate me further.

But Bad Ischl was less than 15 minutes' drive time from Ebensee. And a lot had changed in the interim. I knew, for example, thanks to Podkletnov and his lab assistant's pipe smoke, that whatever antigravity was or wasn't, it had to be induced by something highly unconventional. And Schuberger's approach to the energy-generation process he referred to as "implosion"—if the Legend as relayed to me by Cross contained even a grain of truth—was certainly unconventional.

The Schubergers maintained a large archive dedicated to Viktor's work in their house, the location of the institute, on the outskirts of Bad Ischl. If they would allow me to take a look at it, perhaps I would be able

to find something, some proof, that the Germans really had developed a new and exotic propulsion medium.

They were reluctant to do so and I could understand why: It was well documented in the legends that had grown up around him that Viktor Schauberger had worked for the SS.

In my initial assessment of Schauberger's work, this association seemed to have been drawn straight from the mythology surrounding SS activities during the war and had merely compounded the case against him.

Now, of course, it was quite the reverse.

I'd reached the end of the road; told Marckus everything I'd learned. It was only when I finished that I realized he hadn't interjected once.

I thanked him for listening. He stopped me just before I cut the link.

"Call me, won't you, if you need to talk. I mean it. Day or night. Sometimes it can be good to download, clear the mind for what lies ahead."

As I journeyed on toward the mountains, I felt better about Marckus than I had in months. It was difficult to put my finger on, but it had something to do with the fact that, for once, I was telling him about material of which he had no prior knowledge. I wasn't entirely sure why, but for the first time I felt I could trust him.

Something about Marckus' final tone of voice told me something else.

I felt sure that he knew exactly what I was going through.

As soon as I crossed the Austrian border, I took a break at a rest stop, got out the laptop and pulled up Cross' notes on the Legend. The relevant portion, the part devoted to Viktor Schauberger, boiled down to the following precis:

Late in the war, despite being close to pensionable age, Schauberger was called up for active duty in the German Army. Soon afterward, he received orders to report to an SS institution in Vienna. From there, he was taken to the nearby concentration camp of Mauthausen, informed by the camp commandant that his inventions had received the blessing of Reichsführer Himmler himself, and ordered to handpick a group of engineers from among the prisoners. This would be the "team" that would help him complete his work on an energy device of radical design that Schauberger had begun working on before the war. If he did not comply with this order, the commandant informed him, he would be hanged and reprisals instituted against his family. Schauberger did as he was told.

By 1944, serious work had begun on a Schauberger machine that

appeared to have a dual purpose: first as an energy generator and second as a power plant for an aerospace vehicle of saucer-like appearance. Descriptions of the workings of this generator, sometimes referred to as a "trout turbine," were always woefully inadequate. One such relayed the apparent fact that "if water or air is rotated into a twisting form of oscillation known as 'colloidal,' a buildup of energy results, which, with immense power, can cause levitation."

Details of the chronology were hazy, Cross explained, but Schauberger's team had apparently claimed success a few days before Germany surrendered. One of the scientists who'd worked with Schauberger reportedly said that at the first attempt to run the machine "the flying saucer rose unexpectedly to the ceiling and then was wrecked. The apparatus functioned at the first attempt. . . and rose upward trailing a blue-green and then a silver-colored glow."

The craft in question, Cross said, had a diameter of 1.5 meters, weighed 135 kilograms and was started by a small electric motor with takeoff energy supplied by the so-called trout turbine.

The next part, Cross' text alerted me, was really weird. "A few days later an American group reportedly appeared, who *seemed to understand what was happening and seized everything* (his emphasis, not mine)." Schauberger was kept under "protective U.S. custody" for six months to a year, Cross maintained, and some of his work was reportedly branded by the Americans as "atomic energy research." Cross ended up saying he had no idea what happened to Schauberger after this period.

I kept all of this in mind as I drove through Ebensee, scanning the mountains for a glimpse of the entrance to Kammler's tunnel system: the Zement complex that would have developed and manufactured the A-9/A-10 intercontinental rocket. I saw nothing but for some road signs directing visitors to a Holocaust Memorial. Zement had become a tourist attraction.

Following directions from Schauberger's grandson, Joerg, I pushed through Bad Ischl, following the River Traun through a patchy expanse of woodland until, on the left-hand side of the road, I spotted the institute, a large imposing house with a tower rising from one of its wings, the initials "PKS" painted in letters six feet high on one of its walls. It was here that the Schauberger family organized periodic lectures and seminars devoted to the heretical scientific principles promulgated by Viktor and his son Walter. Walter had since died, but his son Joerg continued to push the ideas that his father and grandfather had promoted for the creation of a better world.

All I knew about the Schaubergers was what I had read; that Viktor,

through his early work as a forester, had developed a passionate set of theories about the essential characteristics of air and water as living, energizing media and that he viewed Nature as a complex interaction of forces that constantly created or reinvigorated matter—against the orthodox view that matter is in a natural state of decay.

Walter, a trained mathematician, had taken his father's theories a step further, merging Viktor's views with Pythagorean concepts of harmonics and the laws of planetary motion developed by Johannes Kepler. It was no coincidence, Walter believed, that a carefully cut cross section through a Pythagorean hyperbola or the elliptical planetary orbits plotted by Kepler happened to be egg-shaped, a form that was one of Nature's highest expressions of energy and strength. PKS, I remembered now, stood for Pythagoras-Kepler System.

It sounded more than a little like New Age craziness, the last place that someone with my background and training should have gone out of his way to visit; but then again, I reminded myself, maybe it was actually no crazier than NASA's Garry Lyles telling me that man would be journeying to the stars by the end of the 21st century in a spacecraft that used some kind of warp drive or wormhole.

I eased the car onto the Schaubergers' property. The institute lay at the end of a long drive. The snow-capped tops of the mountains beyond were dazzlingly bright against the blue sky.

Joerg Schauberger, Viktor's grandson, greeted me on the steps of the house and led me to the basement area where a series of small-scale experiments had been set up to demonstrate the theories developed by his father and grandfather. Viktor's maxim had been "comprehend and copy Nature" and central to this thesis had been his understanding of the vortices that occurred naturally in the environment.

These "energy spirals," as he saw them, creative whirlpools of Nature, were visible everywhere: from the spiral galaxies of the outer cosmos to the power of a tornado or simply in the whorled horn of a kudu's antler or the double helix of a DNA strand.

Nature, Viktor Schauberger had believed, employed the vortex as its most efficient conduit for the transmission of energy.

When this three-dimensional spiraling energy pattern was channeled inward, not outward, in a process Viktor called "implosion," it became endowed with "higher order" properties—characteristics which Viktor himself described as "atomic" (although his understanding of the word was quite different from that of a nuclear physicist's). These properties were capable of generating phenomenal levels of force. Transposed into machinery, Viktor had even coined a term for it: "bio-technology."

One such device, an implosion-based generator, stood disused and rusting in the corner of the basement area where Joerg Schauberger and I now viewed the fruits of his forebears' lifework.

Joerg Schauberger had signaled during our phone conversations his reluctance to prize open the family archives to a complete stranger. But he was content for me to come to Bad Ischl to plead my case. I was glad that I had. He was roughly my age and had an open, honest face. I liked him immediately. I could see that he was weighing me up every bit as carefully as I was assessing him. Both of us were entering uncertain territory. For me, his grandfather's experiments bordered on the incredible. But Joerg wasn't interested in publicity. This in itself gave me confidence. Whatever lay in the archives, I knew it would be fresh, untainted evidence, free from any kind of interpretation or bias.

Joerg was proud of his family's achievements. In the lofty, baronial entranceway of the family institute, he had shown me a cutaway section of a log flume that his grandfather had built in the forests for the efficient removal of lumbered trees.

It was then that I realized the impression painted of Viktor Schauberger in the Legend was a misleading one. Schauberger wasn't a forester in the strict sense of the word. He was an engineer; and from the look of the machinery around the house an extremely accomplished one. His milieu happened to be the forest.

Now, as we stood beside the implosion machine, his grandson spelled out his fears. Others had documented Viktor Schauberger's experiments, he told me; there was little, he believed, I or anybody else could add to this body of knowledge—the more so in my case, as, by my own admission, the science of the processes involved wasn't principally what had led me here.

I was after proof of a technology that conventional science said was impossible.

If, as the Legend had it, his grandfather had cooperated with the SS, what did his family and its reputation possibly have to gain from my seeing the archives?

I had thought long and hard about this during my drive through Germany, but whichever way I cut it, there was nothing reassuring to say.

So I told him instead about the journey that had brought me to his house; about the series of statements in 1956 by U.S. aerospace companies that the conquest of gravity was imminent, that all that was required to make it happen—to usher in an era of clean, fuelless propulsion technology, of free energy—was money and a little application from the U.S. government.

I told him about Thomas Townsend Brown's efforts at much the same time to interest the U.S. Air Force and Navy in an electrogravitically powered Mach 3 flying saucer and how aspects of Brown's ideas had emerged in the B-2 Stealth Bomber more than three decades later.

I told him how Evgeny Podkletnov had defied the theoreticians of NASA's Breakthrough Propulsion Physics program by engineering a device that shielded gravity by as much as five percent and how, officially, Brown's work had been discredited by the U.S. military—despite its very real application to the B-2.

We spoke, as well, about the glaring discrepancy between General Twining's view in 1947 that the development of a manned aircraft with the operating characteristics of a UFO was "within the present U.S. knowledge" and the fact that, even in the black world of modern U.S. aerospace technology, there were so few traces of this knowledge it was as if it had never existed.

Which had brought me to the Germans.

There was, via the Kammler trail, a mounting body of evidence that the Nazis, in their desperation to win the war, had been experimenting with a form of science the rest of the world had never remotely considered. And that somewhere in this cauldron of ideas, a new technology had been born; one that was so far ahead of its time it had been suppressed for more than half a century.

Joerg Schuberger's grandfather, I felt sure, was integral to the truth of what had happened at that time. But unlike the Wenceslas Mine, where few traces of Sporrenberg's court testimony remained on view, Viktor Schuberger had left records. That, I told his grandson, was why I was here.

At this, he gave an almost imperceptible nod and indicated we should head upstairs. It was getting late and there was a lot of material we needed to get through in the short space of time I was in Bad Ischl.

Like me, he said, pausing at the base of the steps that led to the upper levels of the house, he was only interested in the truth. In the complex world his grandfather had lived in, it was a commodity that was not always so easy to get a handle on.

It was then that I understood. Joerg Schuberger had been trying to assemble his own jigsaw puzzle—one that explained the intricate mystery of his grandfather's life.

I held pieces of the puzzle he didn't possess.

In regulating the movement of water via dams and irrigation schemes, man had obstructed the complex, vortex flow patterns that water

naturally adopted for itself, Viktor Schauberger said, causing widespread degeneration and decay in the ecosystem. Harnessed in this unnatural state, water, he came to believe, acted as a pollutant upon the earth, not a life force.

In seeking to come up with methods to restore the earth's ecological balance, he turned to the development of "bio-technical" machinery and promoted his ideas through books, some of which came to the attention of Berlin.

In 1934, a year after Hitler came to power, Schauberger was summoned to the German capital to explain to his Führer and fellow countryman how processes of natural motion and temperature and the vital relationship between soil, water and vegetation combined to create a sustainable and viable society; a society, in effect, that was at ease with itself.

At the meeting was Max Planck, the great German physicist and pioneer of quantum theory, who when asked by Hitler at the end of Schauberger's talk what he thought of his theories replied testily, "Science has nothing to do with Nature" and withdrew from the discussion. Hitler nonetheless asked his technical and economic advisers what could be done to incorporate Schauberger's ideas into the country's four-year economic plan, but nothing ever came of it.

For his part, Schauberger told Hitler that the short-termism of Germany's economic strategy would undermine and ultimately destroy the country's biological foundations.

As a result, he said, the thousand-year Reich would be lucky if it lasted ten.

By using an "impeller"—a propeller that induced an inward-, instead of outward-flowing motion—to draw water through a tube, a flow pattern Schauberger referred to as "centripetal [the opposite of centrifugal] force," he found that the output he was getting was nine times greater than could be achieved with a conventional pressure turbine.

His early egg-shaped implosion machines were also generating extremely high vacuum effects.

By substituting air for water, Schauberger began to envisage a device that with some refinement could be put to use as a radical form of aero-engine; one that sucked rather than pushed its way through the atmosphere.

In 1939, he conceived of a device that could be put to use either as an energy generator or as a power plant for aircraft or submarines. In the application he lodged to the Reich Patent Office the following year,

Schauberger described the essential characteristics of this machine as a "multistage centrifuge with concentrically juxtaposed pressure chambers."

Shortly afterward, he wrote to his cousin that he had invented an aircraft that didn't make any noise.

With these devices, Schauburger realized he had created an entirely new methodology for propelling vehicles through air and water.

As Joerg Schauburger and I settled into the archive, which was crammed full with the old man's files and papers, it was clear that Viktor Schauburger had documented every turn of his career in meticulous detail.

Through his letters, duplicates of which he always placed on file, it was possible to paint an intricate profile of the man and his inventions. The challenge, even for his family, lay in decoding the shorthand Schauburger had used to describe his work.

During the war, he had concentrated on the development of several types of machine—the Repulsator and Repulsine for water purification and distillation, the Implosion Motor for electricity generation, the Trout Turbine for submarine propulsion and a "flying saucer" that used air instead of water as its driving medium. Because they all worked on the same principles, Schauburger tended to interchange the names of these devices and their applications when it suited him.


So, when Viktor wrote in 1940 that he had commissioned a company in Berlin called Kaempfer to build a "Repulsator," it wasn't immediately apparent what this machine was for. It was only when he ran into contractual difficulties with Kaempfer, which was having enormous problems manufacturing the machine to Viktor's demanding specifications, that its function became clear.

By February 1941, Viktor had switched contractors to the Kertl company in Vienna, and here, in correspondence, he described the prototype (which he was building at his own expense) as having a two-fold purpose: to investigate "free energy production" and to validate his theories of "levitational flight."

The machine relied on a turbine plate of waviform construction that fitted onto a similarly molded base plate. The gap between the plates was whorl-shaped, mimicking the corkscrew action of a kudu's antler.

Having drawn air in via the intake, the rapidly rotating turbine propelled it to the rim of the rotating mass under centrifugal force.

The vortex movement of air created by the waviform gap between the plates led to its rapid cooling and "densation," producing a massive reduction in volume and generating a vacuum of enormous pressure, which sucked more air into the turbine.



The machine required a small starter motor to commence the process (as depicted in the Legend), but having whipped the turbine up to around 15,000-20,000 rpm the motor was turned off and the operation became self-sustaining.

By connecting the machine to a gear shaft, electricity could be generated from it; or left to its own devices, it could be made to take off.

This capacity to fly Schauberger partly attributed to the creation of the vacuum in the rarefied region immediately above the plates. But the primary levitating force, he claimed, was due to some other process altogether—a reaction between the air molecules in their newly excited state and the body of the machine itself.

Here, we had touched on the heart of the matter: Had Viktor Schauberger created an "antigravity" device?

If the files were anything to go by, the answer was unequivocal.

Soon after work on the device restarted at Kertl, an associate of Schauberger's made an unauthorized test run while Schauberger was absent from the plant. During this experiment, the machine generated such a powerful levitational force that it shot upward, smashing against the roof of the hangar.

Schauberger's correspondence makes it clear that March 1941 was when things really started to happen. It was then, while work began on repairing the Kertl device, that the Gestapo, the secret police arm of the SS, became aware of his work.

That same month, he reported that Professor Ernst Heinkel, inventor of the world's first jet aircraft, was also showing an interest in his technology.

In fact, Professor Heinkel had illegally obtained sight of Schauberger's patent application at the Reich Patent Office and, without the Austrian inventor's knowledge, had begun to incorporate his ideas into a Heinkel project—one that is presumed by some who have studied the Schauberger archive to have been the He 280 fighter, which made its maiden jet-powered flight on March 30, 1941.

While it is known that development of the He 280 ran into technical problems, especially in the area of its troubled HeS 8 turbojets, there is no confirmation that the He 280 was the aircraft in question. And so I found myself wrestling with a heretical notion. It could just as easily have been the "Flying Top" proof-of-concept vehicle supposedly being built by Rudolf Schriever in a "garage" under Heinkel's guidance at Marienehe, near Rostock on the Baltic coast. Try as I might to resist the

notion, it would be bolstered by additional data on Heinkel in the space of a few more minutes.

The Schriever story was part of the legend that had grown up around Rudolf Lusar's *German Secret Weapons of World War II*, the book that had drawn me into the contentious area of German flying disc research. In it, Lusar, who had been well informed on all other aspects of German secret weapons activity, made the bold assertion that the first Nazi flying saucer projects commenced in 1941 and that the research was centered on two places: Prague and Breslau.

Heinkel coerced the patent office into restricting the use of Schauerberger's technology to water purification and distillation applications, leaving Heinkel free to make fraudulent use of Schauerberger's innovations in his aircraft projects. When the Reich Patent Office offered Schauerberger a patent on this basis, the Austrian refused and consequently had his application turned down.

In May 1941, Schauerberger was formally approached by the "Gestapo," who served him notice (in the company of his patent lawyers) that they would not restrict his work—far from it—but in future he had to pursue it in secret.

This meeting appears to be the trigger for a period of highly covert activity in Schauerberger's life, about which he leaves few clues in his letters. Weeks after his meeting with the Gestapo (which, in Schauerberger-speak, could just as easily have been any SS personnel), he wrote to his son Walter saying he was in Gablonz (present-day Jablonec) in Czechoslovakia and that "what I am doing is secret."

Today, Jablonec is almost exactly where the Czech, German and Polish borders meet. During the war, it was bang in the heart of Kammler's high-tech kingdom.

In a letter the following month, Schauerberger told his son that he had an agreement with a "factory" in the Sudeten region which was undertaking unspecified "research." This factory appears to be one and the same place Schauerberger had mentioned earlier: Gablonz.

Looking at the map and trying not to be seduced by the Legend, I noted that Gablonz was conveniently located between Prague and Breslau.

In the same correspondence, Schauerberger reveals two other facts of significance: first that work at Kertl (apparently on repairs to the device that smashed against the roof of the hangar) is "going so slowly it is as if someone is deliberately trying to slow it down"; and that he has become aware that Heinkel has stolen his ideas.

Schauerberger also reveals that his source in the Heinkel matter is

a member of the "secret police"—a term Schauberger used interchangeably with "SS."

In July 1941, Schauberger describes how "an intermediary" has approached him in a bid to rectify certain problems Heinkel was experiencing with the technology he had lifted from Schauberger's patent application. With ill-concealed *schadenfreude*, Schauberger reveals that Heinkel's engineers have made a mistake copying his design, and that he knows what the error is. Through his dialogue with the intermediary, Schauberger believes that Heinkel is trying to substitute conventional engine technology with technology of his own.

On the face of it, this ought to mean one of two things: that the compressor technology of the troubled He 280's HeS 8 turbojets had been marked for replacement by Schauberger turbine components, the "centripetal compressors" of his successful 1936 patent application; or that Heinkel was attempting to swap complete Schauberger power-plant units for the underperforming HeS 8 turbojets.

The first explanation makes little sense. Adding a "centripetal compressor"—one that causes air to flow radially inward—would have destroyed the carefully tailored flow dynamics of the HeS 8 engine, which was a centrifugal gas turbine based on a principle diametrically opposed to the Schauberger implosion technique.

The second explanation is equally vexing. Strapping Schauberger's engines, which worked on a suction principle, onto the He 280 would have made a total nonsense of the aircraft's graceful aerodynamics.

History, in any case, tells us that the Air Ministry (RLM) would give the He 280 a production order only if Heinkel substituted the HeS 8 engine for BMW 003 or Jumo 004 axial-flow turbojets. Neither of these would fit, so the entire project was scrapped.

There has never been any mention in Heinkel files, so far as I am aware, of any association between the He 280 project and Viktor Schauberger.

This process of elimination allowed me to entertain the notion with greater confidence than I had earlier that Heinkel had been working on some altogether different form of air vehicle—one that would have worked best with a Schauberger engine, not a gas turbine; and one that has never come to light.

It is interesting to note that the legend of the Schriever Flying Top makes specific mention of the fact that construction of full-sized disc-shaped vehicles was transferred from Heinkel's Marienehe facility at Rostock to Czechoslovakia in 1943.

From late 1941, Schauberger's correspondence makes it plain that he

is busily engaged in secret work in Czechoslovakia. In late 1941, he found himself in a "weapons factory" at Neudek (now Nejdek) near Karlsbad (Karlov Vary), around 120 km southwest of Gablonz. In December 1941, he was back in Gablonz eagerly awaiting the arrival of a starter motor—presumably, the little electric engine that was needed to kick-start his implosion device. At one point he tells his son that he is having to make frequent visits to the train station at Reichenberg (today, Libérée in the Czech Republic) to see if the "apparatus" has arrived.

In early 1942, Schauberger transferred to the Messerschmitt prototype works at Augsburg in Bavaria. There, he worked on the further development of a Repulsator implosion machine. In his correspondence, it is not clear if this is the same machine as the ones on which he was working in secret at the two locations in the Czech Republic, although it seems unlikely. What appears to be the case is that Schauberger's work was central to a number of parallel activities during 1941 and 1942: Heinkel's mysterious work at Marienehe, the two secret weapons factories in Czechoslovakia, the Kertl company in Vienna and Messerschmitt at Augsburg.

From Schauberger's writings, we know what happened to the Kertl and Messerschmitt machines. Work on the former ceased due to the company's inability to obtain essential supplies for the repair of the damaged device. This is not unreasonable, given Germany's growing inability to get hold of strategic raw materials from this period on, but Schauberger still voices his doubts, being convinced that somebody has ordered the work to be slowed deliberately.

The Messerschmitt device suffered a catastrophic failure due to the use of substandard casting techniques (and probably low-grade metal alloys) during the manufacture of the turbine. According to Schauberger it suffered a "meltdown" as soon as it reached its maximum rotational velocity. A company in Vienna, Ernst Kubiznak, was ordered to repair the device under a "Führer directive," but this, like the Kertl work, appears to have come to naught.

This leaves the Heinkel work and the secret activities in Czechoslovakia.

After the war, when Schauberger picked up on rumors of German flying saucers that supposedly flew near Prague during the closing stages of the conflict, he voiced his conviction that these were developments of his own ideas—and, given the way Heinkel had behaved, this may have been the case. It was already quite clear that the Sudeten region was the epicenter of some of the Nazis' most secret and unconventional research of the war period. But flying saucers? Antigravity propulsion?

In April 1944, after a yearlong period of inactivity, Schauberger was summoned before a "draft panel" of the Waffen-SS.

In June, he traveled to Breslau, ostensibly to join an SS-Panzergruppe division. But more than six months before the Red Army had battled its way to the gates of the city—before Gauleiter Hanke had sifted the citizenry of Breslau for men far fitter than Schauberger—it is ludicrous to suppose that the SS would have recruited an Austrian of his age to fight in the defense of a patch of the Fatherland hundreds of kilometers from his home. In April 1944, Schauberger was 59 years old.

Given the SS penchant for extreme secrecy, coupled to the fact that Schauberger had already been up to his eyes in secret program work in Czechoslovakia, it is infinitely more likely that the Panzergruppe story was a cover and that Schauberger was taken to Breslau for a specific purpose; one that played to his primary talent as an inventor.

This would have been an ill-disciplined leap of faith, a lapse in cold reasoning on my part, but for one thing.

By June 1944, the SS was almost a year into its takeover of the German secret weapons industry. Bit by bit, the Luftwaffe and the Wehrmacht were being eased out of arms production. In spite of the failures at Kertl and Messerschmitt, there is ample justification—based on what happened next—to assume that the Czech-based projects had validated at least some of Schauberger's ideas.

A month previously, a month *before* Schauberger went to Breslau, he was ordered to Mauthausen concentration camp to handpick a team from among the inmates to start building as many as five different types of Schauberger machine. This is confirmed by the Schauberger archive. The SS wanted him to stop "tinkering around with prototypes and begin serious construction work."

Somewhere along the line, someone must have been convinced that the technology worked.

The essential point, the point that made the hairs on the back of my neck stand on end as I settled into the part of Schauberger's diary that dealt with the last six months of the war, was one of which Viktor himself would have been unaware as he arrived outside the stone walls of Mauthausen camp in that summer of 1944. The big picture, which was in the possession of a mere handful of individuals in Berlin, was that the concentration camps were the designated *production* engines of the "state within the state," as Speer had called the SS; the grim new powerhouses of the Reich economy.

Viewed this way, it shed light on Schauberger's frantic activities

during 1941, when his feet had hardly touched the ground as he dashed between project locations in Germany, Austria and Czechoslovakia, a time that could now be reinterpreted as the start date for prototype work of a diverse and highly secret nature; work that continued until early 1944, when the SS became satisfied that the machines were ready for manufacture.

The immediate task of Schauberger's handpicked team at Mauthausen was to produce engineering drawings from which production work could begin.

Schauberger's diaries tell us what these machines were. One was a water purifier. Another was an energy device capable of generating high-voltage electricity. A third was for "biosynthesizing" hydrogen fuel from water. A fourth was a machine that "naturally" produced intense heat or cold. The fifth, dubbed the "*fleigende scheibe*," or "flying saucer," was the unconventional aero-engine that had come to the attention of Heinkel and others during 1941. It was this latter device that I now focused on.

Schauberger set up design offices in the SS technical engineer school, in the Rosenhugel district of Vienna.

Working under oppressive conditions, effectively at gunpoint, he managed to gain a number of key concessions from the SS. Telling them that he needed personally to select his team from Mauthausen's inmates and that these men should be removed from the camp environment, fed and clothed properly if they were to work productively, he was able to extract around 25 former technicians from the camp and set them up in workshops nearby.

The first set of preproduction drawings were signed off on in August 1944 and delivered from the engineer school to these facilities. Schauberger moved between the two sites, his visits to Mauthausen increasing in frequency as metal was cut on the propulsion device, which he referred to sporadically in the text as the Repulsine (even though this name had come to be associated in former years with other projects).

By early October 1944, Schauberger's diary shows that he was already starting to mill components for the Repulsine. On October 6, he made the first preproduction drawings of a part of the device he refers to as the *Schaufelentwurf*, a design for a cone-shaped air intake that bolted onto the top of the disc.

Five days later, work is interrupted by the first of many air raid alerts as Vienna and its environs are hit by Flying Fortresses on daylight missions against local industrial targets. Mauthausen was close to Linz, home of the Hermann Goering Works, a giant heavy engineering com-

plex. It is clear from these and other entries that Schauberger was racing against the clock to complete his task.

On October 14, he wrote: "I wish I could work longer, but I have to stop at four in the afternoon. There is a shortage of materials and tools. I had to construct parts today from an old tank." Because much of the raw material was of low quality, he instructed his team to be particularly mindful of heat stresses and welding quality.

It was also clear that he was being forced to shift between projects as demands required. On October 23, he turned his attention to developing a more efficient carburetor for Opel Blitz army trucks, presumably on the instructions of the SS, an activity that distracted him for the best part of ten days. On November 4, one of the water-based devices needed his design input. On November 26, it was the turn of the Repulsine again—the electric starter motor needed to be checked and fitted.

The team worked on the Repulsine on Christmas Day and on into the New Year, all the while having to contend with power cuts and air raids. Finally, on February 28, Schauberger moved his operations center to the little village of Leonstein in Upper Austria. The decision was a fateful one on two counts. It not only took him away from the bombing, but placed him squarely in what would later become the U.S. zone of occupation. Had he stayed in Vienna, Schauberger would have found himself in the Soviet zone of influence.

In light of what happened to him in the days, weeks and months that followed the end of the war, it might have been the softer option.

On April 5, 1945, the diary reveals that final assembly of the Repulsine had commenced. A month later, it was ready to go. But the end, when it came, was almost surreal. Instead of firing up the turbine for its first test run as planned on May 6, Schauberger awoke to find that the SS officers charged with oversight of the operation had fled into the night. The team stopped work on May 8, hours before the surrender of German forces took effect at a minute past midnight on May 9.

The SS machine had not flown in the last days of the war, as the Legend had maintained, but in almost all other respects it had held up remarkably well.

The diary had made it clear that Viktor Schauberger had built a machine that had flown earlier in the war at Kertl (and almost certainly during Schauberger's secret period of research in Czechoslovakia). It was also quite clear that the device's modus operandi was wholly unconventional—that is to say, the method by which it generated lift was insufficiently explained by current scientific knowledge.

The diary had given me something I could believe in at long last.

Via Schauberger, the Nazis had been deeply involved—no question—in what can only be described as flying saucer technology. And from this flowed the corollary that other parts of the Legend were perhaps also based on fact.

Couple the Schauberger evidence with what I had seen in the *Lusty* files and suddenly it wasn't so hard anymore to believe that the Germans had developed prototype remote-controlled air vehicles—craft that had the ability to latch onto the exhaust wakes of Allied bombers or the infrared signatures of the aircraft themselves—that explained many of the foo-fighter sightings of 1944 and 1945.

It was also highly likely that some of these "conventionally powered" German disc-shaped craft were test beds for Schauberger-type propulsion systems.

And therein lay another subset to the mystery.

Something about this whole strand of development had conspired to make it the most classified form of technology in existence. Even more so than the bomb.

Unlike the bomb, however, this was a secret that had held for more than 50 years.

Days after the end of the war, U.S. intelligence agents found Schauberger in Leonstein and apprehended him. Exactly as the Legend had it, the agents, who were almost certainly Counter-intelligence Corps—the same outfit that had detained and interrogated Skoda's director, Wilhelm Voss—were remarkably well informed about his entire operation. It was as if, Schauberger noted later, someone had guided them directly to him.

Across the country, at almost exactly the same moment, Russian intelligence agents located and entered his apartment in Vienna. They removed anything of value they could lay their hands on, then blew the place up, presumably to prevent anyone else from discovering information they had overlooked.

For the next nine months, until March 1946, Schauberger was held under close house arrest and extensively debriefed by U.S. technical intelligence about his activities during the war. Years later, he disclosed that it was his knowledge of "atomic" energy that had got the Americans so vexed. This is understandable. The merest suggestion that Schauberger had engaged in nuclear research would have set alarm bells ringing at the local headquarters of Boris Pash's Alsos agents, the U.S. team charged with scouring the former Reich for atomic weapons activity.

And yet, it would have taken the CIC and Alsos days, if that, to

ascertain that what Schauerberger had really been up to had nothing to do with a Nazi bomb or even atomic energy production—at least, not in the sense Los Alamos understood it.

That night, back at my hotel, I applied myself to a study of the Schauerberger effect that had been written up by a researcher named Callum Coats. I had picked up a copy of one of his books at the Schauerberger institute. In it, there was a description of what happened when a Repulsine was rotated at 20,000 rpm. The high rotation speeds appeared to cause the air molecules passing through the turbine to pack so tightly together that their molecular and nuclear binding energies were affected in a way that triggered the antigravity effect. "A point is reached where a large number of electrons and protons with opposite charges and directions of spin are forced into collision and annihilate with one another," Coats wrote. "As lower rather than higher orders of energy and the basic building blocks of atoms, they are upwardly extruded as it were out of the physical and into virtual states."

Virtual states? What the hell did that mean?


A few paragraphs on, Coats elaborated on this theme: "Through the interaction between centrifugal and centripetal forces functioning on a common axis, he was able impulsively to return or re-transmute the physical form (water or air) into its primary energetic matrix—a non-spatial, 4th, or 5th dimensional state, which has nothing to do with the three dimensions of physical existence."

"I stand face-to-face with the apparent 'void,' the compression of dematerialization that we are wont to call a 'vacuum,'" Schauerberger had written in his diary on August 14, 1936. "I can now see that we are able to create anything we wish for ourselves out of this 'nothing.'"

More than 60 years after Schauerberger had consigned these thoughts to paper, I had stood in the Austin, Texas, offices of Dr. Hal Puthoff and listened to the American's descriptions of a near-identical scientific process. *All you had to do—somehow—was perturb the zero-point energy field around an object and, hey presto, it would take off.*

Long before the term had ever been coined, Schauerberger had been describing the interaction of his machines with vacuum energy—the zero-point energy field.

Part of the contribution to the craft's ascent, Coats explained, was due to a quasi-aerodynamic phenomenon—something I knew as the "Coanda effect." But another part came from the expulsion of the densely compressed "emulsion" of molecules and atoms that had not been "virtualized" as they passed through the gill-like slits of the craft's



Teil Nr.	Gegenstand	Stückzahl	Material	Norm Nr.	Gewicht	Bemerkung
Buchstabe	Buchstabe kommt vor	ÄNDERUNG			Datum	Name
Gezeichnet	m. 2. 1941 V. Schauberger	VIKTOR SCHAUBERGER			Einsatz für	
Geprüft					Eingesetzt durch	
Genehmigt	Mauthausen					
Maßstab:	1:1;	Apparat - Repulsine Type B. E. Z. Nr. 70 Ro or Oberteil		Nr d-Zeichng		71

compressor blades. This would have produced a glowing bluish-white luminescent discharge akin to ionization, exactly as described in the Legend.

I closed the book and reached for the large manila envelope that Joerg Schauberger had given me before I left his home.

"Drawing No. 71," signed off on by Schauberger at Mauthausen, showed the simple innards of the *Schaufelentwurf*, the design for the shovel intake, that had bolted onto the top of the craft. I pressed the drawing flat, amazed that a thing of such organic grace should have come from a place that had once been the embodiment of evil. I thought, too, of the effect this must have had on the fragile psyche of the onetime forester, the man who had wanted to bequeath his grandchildren's generation—my generation—a limitless source of safe energy, to do good for the world, but who'd ended up working for the machinery of the Holocaust.

Chapter 22

The sun was just hitting the peaks of the Salzkammergut and the roads were all but empty. Two and a half hours to Munich airport via Bavaria, the cradle of Nazism, and I could box up this whole thing and head home. The beauty of the topography and the shadows of the past were inextricable. It was time to move forward again.

I now had a sequence of events that allowed many of the pieces to fall into place. I had seen the plans of a craft that had drawn on a source of energy that could not be explained by conventional science. The Germans, a people who knew the value of good engineering when they saw it, had had sufficient confidence in the technology to throw money and resources at it. When all else fails, follow the money. It rarely lies.

It must have freaked the Americans out.

At the end of the war, the U.S. Counter Intelligence Corps, the CIC, had spent nine months debriefing Schuberger. They would rapidly have come to realize that implosion technology had nothing to do with "nuclear" energy in the bomb sense, even though Schuberger himself described the processes at work as "atomic."

What they would have learned, however, would have been no less unsettling.

Schuberger believed that his machines were generating an antigravity effect whose power potential was unlimited.

The CIC agents would have had little comprehension of the mechanics of the device—why should they?—but somebody must have been spooked by it. Why else did they incarcerate the old man for nine months? Why else did they threaten him?

By the early part of 1946, following the debriefings and their admonition to Schuberger never again to involve himself in "atomic" science, the CIC would have had in its possession the rudiments of an entirely new propulsion medium—one that 18 months later would have allowed General Nathan Twining, head of the USAAF's Air Materiel Command, to admit to a subordinate in a classified memorandum that it was possible "within the present U.S. knowledge—provided extensive

detailed development is undertaken—to construct a piloted aircraft" with the operating characteristics of a UFO; a craft that seemingly defied the laws of physics.

What had happened to this knowledge?

Part of the answer, I felt sure, lay in some curious synchronicity between the final months of Viktor Schauberger's life and events that had started to unfold at around the same time in quiet corners of the North American aerospace industry.

In the spring of 1958, Schauberger, by now 72 years old and in poor health, was visited by Karl Gerchsheimer, a German-American acting as an intermediary for a U.S. financier and multimillionaire named Robert Donner, who had heard of Schauberger's inventions and wanted to develop and implement them in America.

Gerchsheimer, a 55-year-old German living in the U.S.A., was a man with a colorful past. He left Germany in 1922, eventually settling in Texas in 1937. During the war, he appears to have been deeply involved in U.S. counterespionage activities, ending up as an intelligence agent, almost certainly for the CIC. From the war's end in 1945 to 1950, he was the U.S. civilian property administrator-in-chief in charge of all civil administration, logistics, transport and accommodation for the U.S. army of occupation, and in this role was described as "the most powerful nonmilitary individual in the U.S. zone." In 1950, he returned to the U.S., where he set up a metal fabrication business that ended up doing significant business, ironically, with NASA.

It was during the '50s, according to Callum Coats' book, that Gerchsheimer met Donner, the retired former owner of the Donner Steelworks in Philadelphia. Donner, Coats wrote, was a "patriot who waged constant war against subversive activity in the United States." Between them, Gerchsheimer and Donner became sold on the idea that there had to be an alternative to the use of explosive forces to generate power and motion—a view that Gerchsheimer, through his NASA work, even took up with the master of rocketry himself, Wernher von Braun.

In 1957, Gerchsheimer came to Schauberger's home in Austria, telling him that there were millions of research dollars waiting for him in the United States and an engineering facility in Sherman, Texas, that was all ready to implement his ideas for safe, clean energy production. All he had to do was cross the Atlantic and make it happen. Schauberger, who was suffering from emphysema and a bad heart, took some persuasion, but eventually agreed, the stipulation being that he would stay just long enough to get things up and running—around three months—before

returning home to Austria. His son Walter, a trained physicist and mathematician, would remain on in the U.S.A. for a year to turn his ideas into blueprints from which the Americans could put his implosion technology into production.

Throughout this dialogue, it is clear that Schauberger felt that Gerchsheimer had come to him as some kind of representative of the U.S. government. And while Gerchsheimer subsequently denied this, his evident connections with U.S. law enforcement and intelligence agencies, connections that he never hid from the Schaubergers, merely reinforced the view in the inventor's mind that the offer carried official U.S. sanction. This impression was compounded when Viktor and Walter arrived in the States and Gerchsheimer and Donner tapped physicists at the National Atomic Research Laboratories at Brookhaven, Long Island, for expert advice on implosion technology. Seven weeks after the Schaubergers' arrival in Texas, Gerchsheimer instructed them to write up separate reports about implosion, then forwarded them to Brookhaven for analysis.

At first, Viktor Schauberger was happy to go along with the Gerchsheimer/Donner plan because at long last he believed here was something that would allow him to implement his free-energy ideas for the benefit of mankind. But as the weeks dragged into months, and negotiations in the high heat of Texas turned sticky, Schauberger's health became ever more fragile and he yearned increasingly to go home.

Gerchsheimer and Donner, meanwhile, had become frustrated at the painfully slow pace with which Schauberger was transcribing his ideas onto paper, and the whole relationship broke down.

Prepared to do anything that would enable him to get home quickly, Schauberger made the cardinal error of signing a document that he had not bothered to translate properly. When, later, he fully absorbed the text of it, he realized that everything he owned—models, sketches, prototypes, reports, even intellectual property—had become the sole possession of the Donner-Gerchsheimer consortium.

The release form also stipulated—just as U.S. intelligence agents had, 12 years earlier—that Schauberger was to commit himself to total silence on the subject of implosion and that any further ideas he might develop were never to be discussed with anyone other than designated U.S. personnel.

It was the last straw. On September 25, five days after arriving back in Linz, Viktor Schauberger died a broken man.

There was no point looking for any explanation of this tragedy on the

U.S. side. Other researchers had petitioned U.S. intelligence agencies under the Freedom of Information Act for papers relating to Schauberger's period in Texas, but had been blocked by officialdom, which would neither confirm nor deny the existence of such records, always citing the National Security Act of 1947.

In seeking to find something that explained this final sequence of events in Schauberger's life from the Austrian's side, however, I had come across a small, almost throwaway reference in the Schauberger archive that had seemed to act as the trigger for it.

Shortly before the Gerchsheimer approach, Schauberger records that he was contacted by two aircraft companies, one American, the other Canadian, and that they both offered him substantial sums of money—the U.S. company, according to Schauberger, pitched its bid at \$3.5 million—for the rights to acquire his propulsion ideas.

Schauberger never referred to either firm by name, but as this period coincided exactly with the rise of John Frost's shadowy Special Projects Group and its design of a top secret Mach 3-4 flying saucer for the U.S. Air Force—Project Y2/Silverbug—the Canadian company could only have been Avro. No other Canadian aerospace firm at that time was remotely engaged in such advanced technology work.

I recalled my long phone conversation with Frost's son and his revelation that his father had once made a secret visit to West Germany in 1953. There, at a "Canadian/U.K. government installation," according to declassified documents in Canada, John Frost had met with an engineer who claimed to have worked on a flying saucer project at a site near Prague in 1944-45. Since Frost was probing in all the right areas, he would inevitably have come across Schauberger's work; the engineer whom he debriefed might even have known him personally.

Schauberger rejected both offers because neither company would meet his twin overarching demands: that his turbine should be used "for the common good"—i.e. for commercial aviation purposes—and that the deal should be made public.

This period, of course, also coincided with the period in which George S. Trimble of the Martin Aircraft company, backed by his counterparts in other aerospace firms across the U.S.A., was talking up the conquest of gravity and the impact this breakthrough would have on air and space flight. The world, they announced, was poised on the brink of an era in which free, clean energy would be the norm; and one day, they predicted, it would take us to the stars. In 1955, Trimble had set up RIAS—Martin's Research Institute for Advanced Studies—whose charter had been "to observe phenomena of Nature and to encourage,

promote and support investigations in search of underlying knowledge of these phenomena . . . to discover fundamental laws . . . and to evolve new technical concepts for the improvement and welfare of mankind."

RIAS' charter could have been Schauberger's own.

And look at what had happened next.

No sooner had people started showing an interest in Schauberger again than a man with clear ties to the U.S. intelligence community turns up on his doorstep, tempts him over to the United States and shuts his operation down—permanently.

In 1956, Trimble and others announce to the world that gravity can be harnessed in the same space of time it took to develop the atomic bomb.

Within two more years, all of the companies that had been waxing to the rooftops about the coming antigravity revolution had fallen silent on the subject. By 1960, it was as if none of them had ever even thought of it.

In the meantime, Avro Canada, the only company in the world to have openly admitted to working on a flying saucer—a dog of a thing that has trouble even getting off the ground—has had the rights to the real but highly secret technology, the supersonic models that show phenomenal promise, acquired by the United States. Then that operation is shut down, too.

John Frost, the man who headed up Avro's Special Projects Group, takes its secrets to his grave and George Trimble, many years into his retirement, gets spooked by a seemingly mundane journalistic inquiry 40 years on from these events.

I returned to that central question. What was it about these complementary technologies—the saucer airframe and antigravity propulsion—that had caused such extreme and violent reactions in people directly linked to them?

For the next three decades, no one in scientific circles would have dared discuss these twin strands of development for fear of the ridicule it would have brought upon them. The cult of the UFO—a phenomenon that has clearly been "spun" by the military-intelligence community from time to time—simply made this worse. It was only in the 1990s, as NASA and other scientists began to strive for propulsion methods that far outstrip today's state of the art, that the whole antigravity business started to gain currency again outside the black world. But even then, you had to proceed with caution.

But the real mystery was this. If the principles of antigravity technology had been available since the war—earlier, perhaps, if T.T. Brown's contribution should be given credit—then why were some of

the best math and science brains on the planet still struggling to develop the theory underpinning it?

It was this question, I knew, that would take me back to the States one last time.

Shortly before I boarded my flight at Munich, I checked my cell phone for messages. There were four and Dan Marckus had left three of them. Whatever was on his mind, I knew I was on to something, because, for once, Marckus was chasing me.

With one eye on the departure gate and another on the clock, I called him back.

Even over the bustle of movement in the departure hall and the static of a bad line, I could tell something was definitely up.

"You need to get yourself to Sheffield the middle of next week," Marckus said, his voice scarcely audible. "Podkletnov's come out of hiding. He's going to be talking about his gravity-shielding experiments to a small group of invitees at the university there— aerospace industry and defense ministry types mainly. It isn't a closed session, so you should be able to get yourself on the list. Go talk to him. Something tells me he knows a lot more than he's letting on."

An airline official at the desk was looking at her watch and beckoning for my ticket. I was about to sign off, when Marckus stopped me.

"There's something else," he said.

I showed the girl my ticket and passport and carried on talking while I proceeded to the aircraft.

Marckus started to talk about the experiment down the mine. He'd got my email and now he wanted to talk physics.

"Dan, I'm about to get on the bloody airplane. This can wait, can't it?"

"I know what they were trying to do," he said simply.

I stopped a few paces short of the door of the aircraft.

"I know what the Bell was really about," he repeated.

My tone softened. "OK, go ahead. I'm listening." I was staring out of a small window at the end of the jetway. The rain was beating down on the wings of the aircraft. I could see passengers settling into their newspapers through the fuselage windows.

"They were trying to generate a torsion field."

"What is a torsion field?"

"Laterneentrager means 'lantern holder.' But it's the second code name that's the giveaway. *Chronos*. You know what it means, don't you?"

"Yes, Dan. I know what it means. What is a torsion field? What does it do?"

"If you generate a torsion field of sufficient magnitude the theory says you can bend the four dimensions of space around the generator. The more torsion you generate, the more space you perturb." He paused momentarily, long enough for me to make the connection. "When you bend space, you also bend time."

Chronos.

The third experiment in NASA's BPP study and the one that Puthoff had predicted would be the first to yield a result.

""Now, do you understand what they were trying to do?"

I said nothing. It was Marckus who closed the loop.

"They were trying to build a fucking time machine," he said.

Chapter 23

Viewed from the back of the lecture theater, Dr. Evgeny Podkletnov, the man who claimed to have nullified gravity, cut a slight, shy figure as he gazed up at his audience, a mixed gathering of academics and defense industry types.

Wearing my Jane's hat, I had managed to get myself invited to the Russian's presentation by Dr. Ron Evans of BAE Systems, formerly British Aerospace or plain "BAe." It was Evans who single-handedly ran the giant aerospace and defense company's antigravity effort, Project Greenglow. BAE, I was given to understand, had sponsored Podkletnov's trip to the U.K.

Evans' name was one of the two I'd scribbled down in my notes after Professor Brian Young's lecture, "Antigravity: The End of Aerodynamics?", in London in 1991. The other name, of course, had been that of Dr. Dan Marckus.

Though Podkletnov's English was immaculate, he talked so softly that I found myself leaning right forward to catch his words, which were marked by a strong accent. He didn't look like anyone's idea of a rebel.

He was, I supposed, in his mid-to-late 40s and thin of stature. His dark eyes, which had an intensely haunted look about them, radiated seriousness and conviction. I knew the day his lab assistant's pipe smoke had hit the wall of the gravity shield he claimed to have created, he had found himself locked on a course against the science mainstream. Its reaction had been so vehement and overwhelming that he had gone into exile for the best part of four years to recover from the shock of it—and, word had it, to refine his experiments.

For a heretic, Podkletnov had acquired for himself a nice, quiet audience in this, his first appearance since the personal trauma of that time. Aside from a handful of graduates and undergraduates from Sheffield University's Department of Electronic and Electrical Engineering, the small group of invitees included observers from the U.K. Ministry of Defense, Rolls-Royce aero-engines and BAE Systems.

I let my mind wander back to the day I'd ambled, innocent of all knowledge on the subject, into the marbled hallway of the Institution of Mechanical Engineers in the center of London eight years earlier to hear Professor Young's lecture. Some of the faces in the audience today looked familiar from that evening, though I couldn't exactly say who or why. If I'd known then what I knew now, I wondered if I'd have ended up where I was sitting today. It had been a long haul.

Podkletnov started by recapping the events that had shaped his life and work over the past decade.

By using rotating magnets to spin up his superconducting doughnut-shaped discs to speeds of around 5,000 rpm, he had found that any object placed in the area of influence above them was losing a part of its weight.

Weight reductions of between 0.3 and 2 percent were found to be easily repeatable, he claimed. Five percent reductions had been recorded, although not with the same repeatability. "For a physicist, this is an enormous weight loss, for a practical engineer it is not," he noted. I thought of the NASA engineers at Huntsville struggling to reduce the cost of space access. For every hundred kilos of rocket, Podkletnov was offering them a launch weight reduction of two kilos.

He was right. It wasn't a whole lot. But then again, gravity shields weren't supposed to exist.

I jotted everything down as best I could, knowing that the better my notes, the better Marckus' efforts at interpreting them for me in plain-talk later. He was reluctant for us both to show in the same place, especially with so many members of the establishment present. Once again, I was acting as his eyes and ears.

The Russian peppered his talk with references to General Relativity and Quantum Theory, the unique properties of superconducting materials and his admiration for other pioneers in his field, among them Hal Puthoff, whose idea that gravity might be a "zero-point fluctuation force" Podkletnov professed, with characteristic Russian understatement, to find "rather interesting."

He started to talk about torsion fields, a term still fresh in my mind from Marckus' analysis of the experiment down the mine. Igor Witkowski, my Polish guide, had also talked of "vortex compression," a term used by SS police general Jakob Sporrenberg in his deposition to the Polish courts. Witkowski had remarked how this had linked in with Podkletnov's gravity-shielding experiments.

"The observed effect," Podkletnov said, talking about his own tests, "might be caused by a torsion field excitation of the physical vacuum inside and outside the superconductor."

The idea came from somewhere far and remote, then steadied long enough for me to begin processing it.

Marckus and I had spoken at length since my return from Austria about the Bell experiment and Schuberger's ideas. There was no doubt at all in his mind that what the Nazis had been trying to do at the Wenceslas Mine constituted an investigation into the arcane properties of space-time. And the fact that they had tested the effect of the torsion field on vegetation, lizards, mice and, God only knew, humans as well, suggested that there had been a practical aspect to the experiment as well. There was evidence, Marckus had told me, that particles appeared to slow down when they entered a torsion field.

"Since the zero-point energy field is composed of billions of tiny fluctuations of energy that pop in and out of existence every split second, relentlessly and infinitely," he'd explained, "anything that can mesh with those fluctuations, so the theory goes, can tap into them and extract energy from the field. That's what some people, myself included, believe that a torsion field does."

A torsion field, he'd continued, was best imagined as a rotating whirlpool. If you created one of these whirlpools, dipped it into the zero-point energy field, the seething mass of latent energy that existed on an almost undetectable level all around us, there was evidence to show that it reacted in an almost magical way by directing the flow of energy. Anything that rotated, even a child's spinning top, Marckus said, was capable of generating a torsion field, albeit a very small one.

"There's evidence of this?" I'd interjected.

Marckus had looked at me and nodded. "If you imagine the zero-point field as a giant vat of treacle and man-made torsion field generators like the Bell and Schuberger's Repulsine as mix-masters—food-blenders—then you can begin to visualize their effect."

Some things were better at stirring the zero-point field than others, Marckus had said. Anything that generated an electromagnetic field was a case in point.

"The Bell would have been radiating like hell," according to Marckus, "generating electromagnetic energy on all frequencies, from radio waves to light—no wonder they'd buried the damned thing so deep." The fact that the Germans had also filled the rotating cylinders with a mix of different metals was also significant, he believed. If you could get the proportions exactly right you stood an even better chance of interacting with the zero-point field. But it would have been a very hit-and-miss process—an observation that was borne out by Sporrenberg's

testimony. Each test had been very short, lasting on average of around a minute.

It seemed very much as if the scientists had been trying to "tune" the Bell, much as you would a radio set.

"Get it right and you've got a very interesting piece of hardware," Marckus had told me, "get it wrong and all you've got is some expensive junk."

Sixty-two scientists shot dead by the SS told both of us that the Bell tests had to have been at least partially successful.

It had been the same with Schauerberger's machine. The air molecules whirling around within it had been spun into such a state of super-excitement that something very strange seems to have happened.

Compounded by collisions of electrons and protons—"the building blocks of atoms," as Callum Coats had called them—the result had been the creation of a vortex—a torsion field—similar to the Bell's, only without the electromagnetic component.

This was no ordinary "twister" in a three-dimensional sense, but a coupling device to the zero-point field—a pump, if you will—that not only acted as a conduit for its infinite source of energy, the seething mass of fluctuations, but had combined with an aerodynamic lift component and the Coanda Effect to produce lévitation.

If, as Puthoff and others were suggesting, gravity and inertia were component forces of the zero-point field, along with electromagnetism, then the vortex/torsion field was also responsible for interacting with the zero-point field's gravity and inertia properties. Tune the machine right and you could manipulate them.

Manipulate the inertia of an object and you removed its resistance to acceleration. Put in space and it would continue to accelerate all the way up to light-speed—and maybe even beyond.

Manipulate the local gravity field around an object and you could get it to levitate.

Both of these pathways to "advanced propulsion" were being explored within NASA's Breakthrough Propulsion Physics initiative.

But here was the truly wild part. The vortex, Marckus said, wasn't a three-dimensional phenomenon or even a four-dimensional one. It couldn't be. For a torsion field to be able to interact with gravity and electromagnetism it had to be endowed with attributes that went beyond the three dimensions of left, right, up and down and the fourth-dimensional time field they inhabited; something that the theorists for convenience sake labeled a fifth dimension—hyperspace.

It was here, they said, beyond the vacuum of absolute space, beyond what we now know to be a "plenum" (the opposite of a vacuum) flooded with zero-point energy, that the binding mechanisms of the universe actually lay.

To date, we know of four of these fundamental forces: gravity, electromagnetism, the strong nuclear force and the weak nuclear force.

If a whirling torsion field with or without an electromagnetic component was binding with gravity to produce a levitational effect—an *antigravity* effect—it wasn't doing so in the four dimensions of this world, but somewhere else.

It explained why the Germans had attempted to use a torsion field to act upon the fourth dimension of time. Time, like gravity, the theorists said, was simply another variable stemming from the hyperspace.

"Say," Marckus had told me, "to use an extreme example, the Germans had been able to slow time within the area of the Bell's torsion field, this ceramic-lined chamber, to one thousandth the rate at which it was progressing outside and you sat inside the chamber for a year. What you've done is slow time down on the inside, while on the outside it progresses at its normal rate. Step outside the chamber after a year has ticked by on your calendar and you find yourself a thousand years into the future."

The chances were, Marckus said, the Germans had only been able to generate a tiny time perturbation—perhaps of the order of a hundredth or a thousandth of one percent. Its area of influence, mercifully, would also have been very localized. God only knew, Marckus said, what effect a larger-scale manipulation of the space-time matrix would have had otherwise. Science itself could offer few clues.

I believed from what I had seen in the Schauberger archive that antigravity was real and that the Germans had thrown considerable resources into cracking the problem.

It struck me then that they wouldn't have pursued a single pathway to antigravity, but several. In the same way that the Americans had pursued several different theoretical and applied approaches to the creation of an atomic bomb.

And in the same way that NASA was investigating multiple different routes to breakthrough propulsion physics for taking us to the stars.

In my mind's eye I imaged the strange henge-like edifice next to the power station. Maybe, just maybe, Witkowski was right. Maybe it had been a test rig of some sort. A test rig for a highly unconventional engine or a large circular aircraft.

If the Germans had been looking for an antigravity effect at the

Wenceslas Mine, they would have needed something robust to test the forces generated.

And what of the weirdest variable of all? The gateway to the hyperspace itself?

There was a burst of applause in the lecture theater. If I'd got the Russian right, he was saying that gravity shielding—antigravity by another name—was a torsion field "excitation" of the physical vacuum, soaked as it was with zero-point energy.

The Bell, Schuberger's work and Podkletnov's were all linked by torsion fields and torsion fields appeared to be the key to a host of effects science was at a loss to explain; effects that came from somewhere, some other place, that lay beyond the vacuum of nothingness.

In the Q&A session after the formal part of his talk, Podkletnov had made a reference to something that I needed to follow up on. He'd mentioned that during a recent round of experiments he had generated an even more pronounced shielding effect, one that demonstrated a *consistent* weight loss in objects exposed to the beam of five percent—double his previous best—for short periods of time.

No doubt mindful that NASA had been unable to produce superconductors of the required size, strength and fidelity to duplicate the Russian's experiments, a member of the audience asked him what size discs he'd been using. With a look of evident pride, Podkletnov replied that he'd been using 30-centimeter discs—so strong that the president of Toshiba had been able to stand on one without breaking it.

Half an hour later, during a buffet organized by the university for its unconventional guest and the assortment of government and industry types who'd come to hear him speak, I managed to talk to Podkletnov in a quiet corner of the room.

He must have known that I wanted to pick up on the Japanese angle, because no sooner had I hit him with my business card, than he announced that most regrettably the topic of Japanese sponsorship wasn't for discussion. He had said more than he ought on the subject and that was that.

This in itself, of course, was interesting. But the block on this avenue of inquiry was so abrupt that for a moment it wrong-footed me.

I grappled for some new purchase on the conversation. "How about aerospace applications?" I asked.

I half expected Podkletnov to tell me what he had previously told Charles Platt, the American journalist who'd tracked him down in the wilds of Finland. Platt had revealed that Podkletnov was working on a

spin-off of the shielding effect that had allowed him to achieve "impulse reflection"—using gravity waves as a repelling force. These, Podkletnov had told the writer from *Wired*, would one day permit the development of a "second generation of flying machines," beyond the brute-thrust types we rode on today.

But Podkletnov went off on a different tack altogether.

"I have discovered," he said slowly, watching and weighing me as he spoke, "a new and truly bizarre effect."

"What kind of effect?" I probed. The low-key nature of the Russian's lecture had already signaled that he wasn't in the habit of using hyperbole.

"I'm not sure that I am ready to talk about it publicly yet," Podkletnov said, smiling awkwardly.

I was about to return to the subject of the Japanese for leverage, but in the end I didn't have to.

"I am, however, happy to share this news with you," he said. He paused to give the room a quick sweep. There was no one close enough to overhear us.

"If the superconductors are rotated considerably faster than the 5,000-rpm speeds I've been mainly using until now, perhaps five to ten times as fast, the disc experiences so much weight loss that it actually takes off."

"Have you experimented?" I asked.

"Yes," he said meaningfully, "with interesting results."

It was then that I asked him if he had heard of Viktor Schauberger. It kind of came from nowhere but in actual fact it wasn't such a wild shot in the dark. There was something incredibly familiar about those rotation speeds.

Schauberger had been generating a lévitation effect using rotational velocities of 15-20,000 rpm. If these were rapid enough to produce a usable torsion field, no wonder Podkletnov had been getting "interesting results" from 25-50,000 rpm.

Podkletnov weighed his response before replying.

"You should understand that I come from a family of academics," he said after a long moment of reflection. "Both my father and grandfather were scientists. Shortly after the war, my father came to acquire a set of Schauberger's papers. Some time later, when I was old enough to understand them, he showed them to me."

"Original papers?" I asked.

Podkletnov said nothing, but in itself this told me as much as I needed to know.

When the Red Army entered Vienna, Russian intelligence agents

had made a beeline for Viktor Schuberger's apartment. There, they had found documents and certain component parts that Schuberger had spirited out of Mauthausen and the SS technical school at Vienna-Rosenhugel shortly before the SS transferred him and his design team away from the bombing to the little village of Leonstein.

Afterward, the Russians had blown up the apartment. But they had evidently taken what they could from it. And for some reason, Podkletnov's father had acquired some of the plunder.

"Perhaps you know that the Germans experimented quite extensively with gyroscopic forces for aircraft during World War Two," Podkletnov continued.

I wanted to tell him about the evidence I had uncovered in Austria and Poland, to share some data quid pro quo so I could learn more about his father's work, but out of the corner of my eye I could see one of the civil servants from the Ministry of Defense making his way toward us, a plate of sandwiches in one hand, a notepad in the other. A discussion about Podkletnov Senior and German "gyroscopic aircraft" would have to wait till another time.

On the way to the train station, I called Marckus and told him about the three significant facts I had learned that day. First, that the Japanese were in on the antigravity/free-energy act; and Toshiba wasn't routinely in the business of throwing good money at crackpot ideas. Second that Podkletnov was generating a free-floating levitational effect by rotating his superconductors at 25-50,000 rpm. And third, that Evgeny Podkletnov's father had come to acquire original Schuberger documents soon after the end of the Second World War.

"What would Evgeny's dad have been doing with them, Dan? It's a hell of a coincidence, isn't it?"

"Leave it with me," he said, "I'll call you back."

He did so within the hour. He'd been on the Internet, he said, scanning, processing stuff, eventually finding what he was looking for on the IBM patent-server, a voluminous website. I could tell from his voice that this was Marckus' kingdom; the place where he was happiest.

"I grind away at this material in low gear like a Stalin tank, but I always get there in the end."

"What did you find?" I asked.

"Something . . . curious. More than a coincidence, I believe. It seems that Podkletnov's father filed a master patent in 1978. It's funny, because I damn near quipped to you earlier that he'd probably turn out to be the father of the Moscow sewerage system. Well, guess what? The patent relates to industrial techniques for the continuous pure enameling of

metal pipes. Evgeny's dad, in other words, had come up with a novel way of lining steel water pipes. It tells us two things: that he was a materials scientist, just like his son; and that he knew about water. My guess is that the Podkletnovs would have been quite comfortably off under the old Soviet system."

"How did he come to acquire the Schauberger papers?"

"Ah, well, let's postulate a little. Russian intelligence agents enter Schauberger's apartment in Vienna at the end of the war and find a bunch of papers, the contents of which they really don't begin to understand. But they know they're important, because they've already tied Schauberger to the SS secret weapons program. So they bring the documents back to Moscow and look around for someone who can decode them. Most of Schauberger's work, remember, related to machines that used air and water as their driving medium. What was Evgeny's dad? He was a leading academician in the hydro-engineering field—a pipes man. Not exactly the first person you'd think of approaching to do the work had he lived over here, but then again, in Schauberger's field, remember, there weren't any dedicated experts. So, Podkletnov Senior, a water engineer, got the job."

"You think the Soviets took the technology further?"

"Who knows? The Russians have never been afraid to exploit weird science. Remember the remote-viewing program kicked off by the CIA under Hal Puthoff? The reason the CIA was into it at all was because it picked up on the fact that the Soviets had started to dabble in the psychic spying business. There is ample evidence that the Russians have been tinkering with torsion fields for a long time, too. But my guess is that they never found enough in Schauberger's apartment to make practical use of his theories."

"But enough for Evgeny's dad to be intrigued by them," I suggested.

Markus paused.

"What is it?" I asked.

"Podkletnov's discovery probably wasn't the accident he's cracked it up to be."

"You mean, when his assistant's pipe smoke hit the column of gravity-shielded air he was already looking for an effect?"

"That's certainly how it appears now we know what we know about his father."

"Jesus," I said, "if he can do it on a shoestring, anyone can."

"Right. Scary, isn't it?" He sounded like he meant it, too.

But I was still left staring at the end-wall of the same blind alley in which I'd found myself on the way back from Bad Ischl.

"Here's the problem I'm facing, Dan. If Schuberger and Podkletnov can crack antigravity, more or less in their garden sheds, then why can't NASA, for Christ's sake, with billions of dollars at its disposal?"

He thought about it for a long moment, then lightly cleared his throat.

"My best shot, and it doesn't help you much, is that what you and I have learned tells me that there are two kinds of science. The stuff they teach you in college and all the weird shit they don't. The aerospace and defense industry is inherently conservative. It doesn't like change. This . . . knowledge, if you can call it that, is dangerous stuff. It's change with a capital C and it's not easy to get your head around. The aerospace and defense industry says it likes people who think out of the box, because they're the guys who give us the breakthroughs . . . radar, the bomb, stealth and that. But think this far out and they look at you like you're crazy. They might even put you away."

I thought back to the article that had landed on my desk all those years ago and my failed attempt to interview George S. Trimble.

Only a maverick, an enlightened one at that, would have pushed for the foundation of something as bold as RIAS—the research institute spun out of the Martin Aircraft Company that Trimble had helped to set up in 1955, at exactly the time he was making his public pronouncements about the imminent conquest of gravity.

I liked to think that all those people who'd been talking up antigravity in the mid-1950s had been forced into silence by some murky intelligence outfit tasked with guarding the truth about the Germans' achievements during World War Two.

Yet, they might just as easily have been silenced by their own chief executives—people who realized they were talking up the premature death of the aerospace industry itself. A science that made their whole world redundant.

Like white goods, fast-food chains and cars, the aerospace and defense industry of the 21st century has become a global enterprise. Back in the 1950s, though, as a modern entity, one that we would recognize today, it was only just getting going.

The 1950s was the decade that spawned the intercontinental ballistic missile, the Mach 2 jet fighter, guided weapons and transatlantic jet passenger planes. It was a time when the industry needed brilliant individuals—mavericks like Kelly Johnson and Ben Rich of the Skunk Works—to bring about those revolutionary advances.

Maybe that was why one name had kept popping into my head since my return from Austria.

When I had first inquired about Trimble and RIAS at Lockheed

Martin, my contacts there had recommended I interview a man who routinely talked about the kinds of things that had once been integral to the RIAS charter.

A man who tended to talk about Nature, not science; a physicist who looked at things quite differently from other people.

Boyd Bushman was a senior scientist for Lockheed Martin's Fort Worth Division, the part of the corporation that turned out F-16 and F-22 fighters for the U.S. Air Force. If you're interested in antigravity, then talk to Bushman, they'd said.

Bushman, the guy who'd levitated paper clips on his desk.

I hadn't, because back then Bushman's theories had sounded just a little too weird. And back then, I hadn't begun to accept the fact that antigravity and the multibillion-dollar interests of the aerospace industry went together like chalk and cheese.

But it was all a question of perspective. I saw things very differently now from the way I'd seen them then.

What, I wondered at this moment, was an antigravity man doing at the world's most prolific manufacturer of good old-fashioned, jet-powered combat aircraft?

If Bushman was talking up antigravity in the midst of an industry that didn't want to know, then maybe, just maybe, he could provide the answer to the \$64,000 big one; the discrepancy between clear evidence in the real world of the existence of antigravity technology and the people at NASA who were, at this very moment, spending ten times that amount scratching around for theory to make it work.

In two weeks' time, I was due on another defense industry junket to the U.S. West Coast. From there, it would be easy to route back to England via Texas.

Journeying into Kammler's kingdom had taught me the value of analyzing data *in situ*. Now, as I prepared for one last trip to the States, I needed something to help me acquire a different kind of mind-set.

I considered how, in July 1986, an F-117A Stealth Fighter crashed at dead of night into a desert canyon just outside Bakersfield, California. For me, the crash had always held a special kind of significance, since it was one of those rare occasions when the black world had been flushed into the light.

In 1986, the very existence of the Stealth Fighter was still a classified secret even though the aircraft had been in full squadron service for three years.

What did Bakersfield say about the system that had been set up to protect America's most deeply classified secrets? Had it been a murder

site, what would a profiler have deduced from the traces this incident had left in the environment?

Back at my desk, I started sifting through a pile of newspaper clippings that I'd kept since the time of the crash. Through the clippings, I had traced the whereabouts of a number of witnesses to the events of that night. One of them, Amelia Lopez, now a local law enforcement officer, had reluctantly promised to act as my guide should I ever find myself in that neck of the woods.

I checked to see that I still had Lopez's numbers. Then I picked up the phone and called the public affairs department of Lockheed Martin, Fort Worth.

Chapter 24

The flight was several hours out of Heathrow, when I reached into my hand luggage and extracted the book that Marckus had made me promise I'd read before I got to the States. He'd sent it to me ten days earlier, but in the hurly-burly of stories that needed to be written before I disappeared off to America, and in the course of preparations for the weeklong trip itself, I simply hadn't had the time to pick it up.

Not only that, but the book, which was called *Blowback—The First Full Account of America's Recruitment of Nazis, and Its Disastrous Effect on Our Domestic and Foreign Policy*, looked rather academic and heavy-going. Even now, as I gazed at the cover, I found myself reluctant to make a start on it, the more so as Marckus never told me what it was about the book that made it essential reading. But I'd read the newspaper, established that there were no movies on the entertainment system worth watching and had even plundered the in-flight magazine for items of interest. With nowhere left to go, I turned to *Blowback* and started reading.

It began promisingly enough, although the territory itself was more than familiar. By Chapter 3, the author, Christopher Simpson, an American, had got his teeth into his country's recruitment program of German scientists after the war—an operation that had started under the code name "Overcast." As early as July 1945, barely two months after the end of the war in Europe, the U.S. Joint Chiefs of Staff gave its authorization for Overcast to commence. Its job was to "exploit. . . chosen rare minds whose continuing intellectual productivity" America wanted to use in the continuing war against the Japanese and in the gathering Cold War against the Soviet Union. Under the top secret project, the JCS directed that 350 specialists, mainly from Germany and Austria, should be brought immediately to the United States. Among the Germans who came to the U.S. at this time were rocket specialists like von Braun and his boss, Army General Walter Dornberger.

I did as I always did with books that held details of Germany's wartime technology effort: I flicked to the index and looked for references to

Kammler. More often than not, Kammler never rated a mention, such was the ease with which he had blended into the bland backdrop of the Nazi administrative machine. Here, in *Blowback*, however, Kammler did score a single entry, but only in the context of his relationship with Dornberger at the Nordhausen underground V-2 plant. An arrangement was struck, the book stated, in which the SS, under Kammler, assumed the day-to-day administration of Nordhausen, leaving Dornberger in charge of the rocket technology and the V-2's production schedules.

By 1946, the Pentagon's Joint Intelligence Objectives Agency wanted to recruit even more former Nazi scientists in the gathering standoff against the Russians, but because U.S. immigration laws forbade entry into the U.S. of any former Nazi Party officials, President Truman was forced to authorize a more extensive effort than Overcast, code-named Paperclip, on condition that it was kept secret and that only "nominal" Nazis—opportunistic party members who had not committed any war crimes—were to be allowed into the country.

I read on, but the rest of the chapter told me nothing I didn't already know. Simpson's research was excellent, but it was marred by the fact I'd previously read Tom Bower's compelling work, *The Paperclip Conspiracy*. From this, I'd gleaned as much as I needed to absorb about Paperclip: that by September 1947 it had officially been canceled, but in actuality had been replaced by a new "denial program"; one so secret that Truman himself was unaware it existed. Paperclip continued as a "deep black" program (as it would be described today), recruiting its last scientist in the mid-1950s. Through Paperclip, hundreds, if not thousands of former Nazi engineers, many of them with highly undesirable backgrounds, were permitted entry to the United States and employed on cutting-edge defense and aerospace projects, not the least of which was NASA's moon program.

I pressed on, but from Chapter 4 onward, Simpson's book took a new turn, describing the CIA's recruitment of former Nazi intelligence agents, the spy networks they set in place against the Soviet Union, and the State Department's elaborate efforts to cover up its wholesale recruitment of these people.

I laid the book down and took a pull of my drink. When Simpson had written *Blowback* in the late 1980s, I could see it had been hard-hitting, revelatory stuff. But I couldn't see why Marckus had gone to such lengths to persuade me to read it.

I continued to wonder about this long after I landed in Los Angeles and arrived at my hotel. But given the time difference between California and the U.K., it was too late to call Marckus and by the following

morning, as I journeyed up the coast to Santa Barbara, the scene of several appointments with a number of defense electronics companies that day, my mind was preoccupied with other things.

After Santa Barbara, I headed inland to Bakersfield, met up with Amelia Lopez and proceeded to the site where the F-117A Stealth Fighter had crashed in 1986, two years before it was unveiled. That the security surrounding the plane had remained in place after the crash was mainly down to the fast footwork of special Air Force crash-recovery teams, whose policy included sieving the earth up to a thousand yards from the crash site for pieces, then scattering bits of a '60s-vintage Voodoo fighter prior to their departure to throw the curious off the scent.

To me, Bakersfield said two things. It said that the black world would stop at nothing to prevent America's "quantum leap technologies" from spilling into the public domain until its bright young colonels were satisfied nothing more could be gained by holding the lid of secrecy in place. And two, in scattering bits of Voodoo, an aircraft that hadn't been flown actively for years, it wasn't averse to sending a signal.

When restrictions on the site were lifted, aircraft enthusiasts went into the hills and picked over the site to see if they could find any evidence that for the first time would point to the reality of the Stealth Fighter's existence. When they brought back little pieces of aircraft and had them analyzed, they knew exactly who'd been yanking the chain. The pieces of Voodoo were the telltale flourish that anonymous artists leave to denote the authorship of their work to cognoscenti of their talent.

It was in the canyon where the Stealth Fighter had gone down that I saw the book again in a flickering moment of clarity. When Amelia Lopez and I were clambering back across the rocks to our cars, the picture steadied and I knew why Marckus had wanted me to read it. The book was the key to everything, not least Marckus' role in all this. His fatalism, humor, cynicism and cunning clicked neatly into place. It was so simple yet overwhelming it almost took my breath away.

Fort Worth owes its very existence to the military, which in 1849 established a small outpost on the Clear Fork of the Trinity River some 30 miles to the west of Dallas.

A quarter of a century later, the coming of the railroad turned what until then had been little more than a cluster of tents and wooden huts into the biggest cattle railhead west of Chicago, a status it retained until shortly after World War Two.

By the 1940s, Fort Worth had begun to profit from other industries aside from cattle and oil, and as you drive out of the city, heading west, it

is impossible not to notice Air Force Plant No. 4, a low building a mile long that rises from the Texas plains like an enormous rock escarpment.

Plant 4 had been set up during World War Two to manufacture the B-24 Liberator heavy bomber. In 1944, when output reached its peak, the plant was turning out almost ten aircraft a day; a feat of mass manufacturing that would never be equaled again.

Today, Plant 4 produces the Lockheed Martin F-16 at just a handful of aircraft per month. Lockheed Martin inherited the F-16 program from General Dynamics, whose aircraft and missile businesses were purchased by what was then simply the Lockheed company in the early 1990s.

GD had, in turn, acquired the Fort Worth operation from Convair, with which it merged in 1954. During World War Two, Convair was known as Consolidated Vultee Aircraft. Originally centered on San Diego, Convair, together with the USAAF, decided to set up the B-24 production line about as far from the Pacific as you could get, as a precaution against bombing by the Japanese.

In 1942, a young engineer called Robert Widmer moved from the San Diego operation to Fort Worth to begin work on the B-36, a massive intercontinental bomber that would make even the B-24 look tiny. It had been specified in 1939 to bomb Germany from the United States in the event that Hitler succeeded in invading Britain. It rolled out of Plant 4 in 1946 and went on to become the backbone of Strategic Air Command during the 1950s.

Bob Widmer spent much of the late '40s and early '50s lobbying to save the B-36 from cancellation before going on to design the B-58, SAC's first supersonic bomber. But by the late 1950s, Widmer had also begun throwing his energies into another project—one that was so advanced that the details would still be beyond the public domain four decades later.

The project was called Kingfish and it emerged out of a requirement by the CIA for a very fast high-altitude reconnaissance aircraft to replace the U-2, which was rapidly running out of places to hide from the Soviets' SA-2 Guideline surface-to-air missiles in the rarefied air over the Soviet Union.

As chief engineer of Fort Worth at the time of the Kingfish program and, almost 40 years later, still a consultant for Lockheed Martin Fort Worth, Bob Widmer was someone I very much wanted to meet.

What was so special about Kingfish that it had remained out of the limelight all these years? Questions about this aircraft take on an even more bizarre twist, when you consider the official history of the program that spawned it.

I was interested in Widmer and Kingfish, because to me it appeared to offer a link—one that had been overlooked by everyone else—to Aurora, the large black triangular aircraft that had supposedly been sighted over the North Sea in 1989.

In August 1959, the CIA's contract for a high-flying, super-fast reconnaissance aircraft went not to GD Convair, but to Kelly Johnson's Skunk Works, which went on to deliver an aircraft called the A-12 under a program code-named "Oxcart."

The A-12 flew for the first time in 1962 from the USAF's remote test center at Groom Lake, Nevada, better known today as Area 51. The 15 A-12s that were built conducted three years' worth of operational reconnaissance missions up to the program's cancellation in 1968. By then, an Air Force version of the plane, the Lockheed SR-71 Blackbird, had entered service.

With a top speed of Mach 3.4, the Blackbird remained the world's fastest operational aircraft until its retirement in the early 1990s. If you exclude the evidence for Aurora, there is, even today, nothing remotely as fast.

In official versions of the Kingfish/A-12 contest (declassified by the CIA only in the early 1990s) the Convair and Lockheed proposals were remarkably evenly matched. Both aircraft were optimized for cruise speeds of Mach 3.2, but the Lockheed proposal beat its rival's by tiny margins on key parameters such as range and operating altitude. On this evidence, then, it was only right that Lockheed got the deal.

But in the same way, as Marckus observed, that there are two kinds of science, so there are two kinds of history.

In 1995, a small photo of a wind-tunnel model of Kingfish appeared for the first time in the pages of an aerospace magazine. Scrutiny showed the model to be much more advanced than the paltry details in the caption suggested.

When I made inquiries at Lockheed Martin, under whose historical aegis both of these programs (Kingfish and the A-12/SR-71) now fell, I discovered that Bob Widmer, Kingfish's designer, was still employed at Fort Worth as a consultant. At 80 years of age, Widmer was busying himself in unmanned combat air vehicle technology. UCAVs—cheap, pilotless fighter planes—are seen very much as the future. It says a great deal about Widmer's skills as a designer that he was still employed at the cutting edge of aerospace and defense engineering almost 60 years after he joined Consolidated; but then Widmer, I was about to learn, was no ordinary engineer. Like Lockheed's Kelly Johnson and other "engineer-

leaders" of the day, he'd helped to shape the modern U.S. aerospace industry.

His official biography describes him as an "innovator and a maverick" and the "first engineering leader to totally integrate airframe, propulsion and avionics into a single weapon system, the B-58."

What it does not say, but what I had gleaned from others, was that he blended these skills with those of a pastor, acting as guide and mentor to his team. In the 1950s, the pressure on GD Convair to outperform its rivals was intense, but the atmosphere inside the plant was informal and shirtsleeves—almost familial. To many of the younger engineers, Bob Widmer wasn't just a brilliant designer, but someone they could turn to for advice—something of a father figure, in fact.

We started by talking about Convair's own black ops capability, a special projects facility that stayed in the shadows, unlike Johnson's entity over at Lockheed, which reveled in its status as black world aviation's one-stop shop for special programs.

"Kelly was the catalyst, the guy who did things differently," Widmer said, "but, hey, we had a 'skunk works,' too; the difference was, we never spoke about it."

Widmer's slight, sinewy body, so much frailer than it had been in the black-and-white shots from the '60s, trembled with excitement. His small eyes shone.

As the talk drifted into the early days of the Cold War, a time when GD Convair came into its own as a reconnaissance specialist, I could see that the minder assigned to me from the PR department was beginning to lose interest. An individual I hadn't come across before, he was there as a matter of routine to ensure that I did not encroach upon matters that would put Lockheed Martin in a bad light or compromise national security. But since this was history, history from way back at that, it hardly constituted a threat to either. It was late and the minder looked tired. His eyes were closed, to focus on the conversation more intently perhaps? But if I hadn't known better, I'd have said he was asleep.

In this atmosphere of near-conviviality, I asked Widmer about Kingfish.

At first, he was reluctant to say a great deal, but seeing that I knew some of the details already, he began to open up. As we talked, I could hear the anger in his voice; anger, I thought, that stemmed from the fact that the CIA had chosen Lockheed's plane, not his, for the contract.

This portrait of Kingfish—the one that Widmer now began to talk about—was markedly different from the official CIA version. And here

was one element of the mystery. In its final design form, Kingfish had been at least twice as capable as the variant that is recorded in official agency files.

The final iteration of Kingfish was optimized to cruise at 125,000 feet at a speed of Mach 6.25. It had contained a great many "firsts," including two powerful ramjets—a radical form of power plant that, to this day, has never been deployed on an operational aircraft—a blended, stealthy airframe shape composed of high-temperature steel and a special heat-resistant substance called "pyro-ceram" that also acted as a highly effective radar-absorbent material.

Kingfish would have streaked far above the Soviets' air defenses, twice as fast as any aircraft the Russians ever consigned to the drawing board; and it would have been all but invisible to radar.

Kind of chilling, too, was the fact that its specification was a near-perfect match for most people's idea of what Aurora should look like—and Widmer had designed the aircraft almost 30 years before rumors of a high-speed Mach 6-8 reconnaissance plane had begun circulating in *JDWand Aviation Week*.

That the CIA bought the A-12 instead of Kingfish is, of course, an enormous tribute to the Lockheed project team, which went on to overcome unprecedented design challenges to make the A-12/SR-71 Blackbird a reality. But while Blackbird is now a museum piece, the finer details of Kingfish, the loser in the CIA contest of 1958-59, will forever remain locked inside Widmer's head.

At the end of the competition, the CIA moved in and ordered all the blueprints, plans and files to be shredded. They also told him never to talk about it again.

I asked what effect this had had upon him.

"At the time, the CIA were in their glory," he replied. "I don't want to go into detail about the kind of life I had to lead . . . how I had to conduct myself, my family and so forth."

"Was it very restrictive?"

Widmer looked over his shoulder at the minder, who must have detected the change in mood. The minder looked at me and I looked at Widmer.

The former chief designer continued. I could see that he was determined to say his piece. "It was unbelievable. It wasn't just restrictive. It was counter to my conscience and morals and so forth."

"The whole business of black program secrecy?"

He nodded and I got the sense he was speaking not so much for himself as the younger members of his team, the people who'd looked to him for

guidance. For an aerospace engineer the black world was a regime that ought to be endured only for a small window in a person's career—and even that was a crime, Widmer said. It was not, he added, what he would call a good formula over a long haul.

When I noticed his hands, I saw that they were shaking.

After a pause that was long enough for me to image the mixture of suspicion and draconian security that had once stalked him, he added: "I mean, the degree they went to in order to hold the secrecy was unbelievable."

The look in his eyes was so far-off and intense that I was reluctant to intrude further. But I needed to ask him one last question.

"Did it frighten you, ever?" I said.

"Yes," he replied, his voice scarcely a whisper.

As I left the room, heading under escort for my appointment with Bushman in the cavernous interior of Plant 4, the minder and I said nothing. Perhaps this way, it was easier to pretend that nothing had happened. But as we walked across the shop floor that had consistently churned out combat aircraft for the past 50-odd years and then stepped back into the strip-lit corridors and past endless security-sealed doors, the air-conditioning cold on our faces, I couldn't help but wonder at the senselessness of it all. One look at Widmer, another at his résumé, told you everything you needed to know. His ambition had been to design and build aircraft that made GD Convair the best plane-maker in the business and which protected America from its enemies. In most people's eyes, that made him a patriot, not a security threat.

But the CIA, or whoever the hell had terrified the bejeezus out of him and his family, had insisted on making the point. And what for?

Had Kingfish, the official loser in the CIA's high-speed spyplane contest, gone on as a "deep black" development program, emerging years later as Aurora?

Or was the truth more prosaic?

Had Kingfish simply been too expensive to be developed, forcing someone, somewhere to consign its design principles into a black hole of ignorance, the mythic warehouse that Ben Rich had told me about, where no one would ever be able to get at them?

If so, how many other technologies were in that place? How many more Widmers had been terrified into keeping that perpetual oath of silence?

Hundreds. Thousands in all probability.

Whether George S. Trimble had developed antigravity in the black or

not, as head of advanced programs at Martin Aircraft, he would have been forced to abide by the same austere security oath. No wonder, then, he'd had second thoughts about our meeting all those years ago. No wonder my PR contact at Lockheed Martin had relayed the observation that he'd sounded scared.

Trimble must have been scared for the same reasons Widmer had been.

God only knew what they thought might happen to them.

Like an unsinkable ship, the black world had been built up around multiple, layered compartments, each securely sealed. Some of those compartments, it is now clear, had been designed never to be opened again. Ever.

The Kingfish story provided a tantalizing glimpse of the contents of one such compartment. By approaching the black world from another direction altogether, I had come to learn of the contents of at least one other.

At the end of the Second World War, via Schauberger's work and, no doubt, a wealth of other captured technology from Germany that has never seen the light of day either, America acquired knowledge of the most dangerous kind.

The Germans' secret weapons program had yielded a power plant—a free-energy source, to boot—that had given America the ability to design a radically different form of air vehicle, one that reflected General Twining's assessment to General Schulgen that a "circular or elliptical" craft that normally made no sound, which could turn on a dime and was capable of extreme acceleration, could be constructed, "provided extensive, detailed development" was undertaken.

Something of the order of the Manhattan Project that had yielded the atomic bomb.

But there is no evidence that the Pentagon ever initiated such a program.

The closest thing to it that has reached the light of public accountability was the recently declassified Project Silverbug, the USAF's development of the supersonic saucers designed by Avro Canada in the mid-late 1950s. But Silverbug, being jet-powered, was conventional compared to what the SS had tried to develop in Czechoslovakia, Lower Silesia and Austria.

If the U.S. agents who debriefed Viktor Schauberger in 1945-46 so much as half appreciated the ability of his or other devices to cause disturbances, however small, in the fabric of space and time, to unleash untold "variables" hidden in dimensions that conventional science hadn't

begun to see or understand, then someone, somewhere would have taken the view that it should be kept secret for a very long time.

There were other even more disturbing possibilities.

During their time in America, it had dawned on Viktor and Walter Schauberger that the implosion process could be harnessed to create a massive weapon—one that was infinitely more destructive than the hydrogen bomb. More than anything else, they were terrified that their "hosts" would latch onto the same idea.

I filed this away in the back of my mind as something to put to Marckus.

Bomb or not, it appears that antigravity was boxed up and put away, ready for the day when American science could harness it consistently enough to control it. Besides, in the 1940s and 1950s, it wasn't as if the world really needed it. The global economy—and none more than America's—has been reliant on oil and the technology that supports it for more than a century. It is only now that we're having to confront the end of the petro-chemical era and seek alternative power sources.

I was left with an image of Widmer's hand, shaking with anger, and a cold feeling I hadn't had since Witkowski told me about the murders of the 62 Bell scientists at the hands of the SS. It didn't take a whole lot of introspection to get to the root of it.

For years, I'd labored under the assumption that it had been the bomb program that had kick-started America's black world into being. It hadn't, of course. From a security standpoint, the Manhattan Project was a flawed model. It had leaked like a sieve and the Soviets had acquired the most powerful technology on earth, the atom bomb, a scant handful of years after the Americans.

In 1944, when the Joint Chiefs of Staff established their high-level committee to hunt down on America's behalf every military and industrial secret that Nazi Germany possessed, they couldn't have begun to foresee the full consequences of that directive. America, more than any other ally, had acquired a vast array of German-derived technology. Much of it, like von Braun's V-2 rocketry, was highly advanced, but essentially recognizable. Some of it, however—notably the nonconventional science pursued by the SS—came from a different culture altogether.

To master the weapons it had acquired from the Germans, America found that it had to recruit the engineers responsible for them. Some were former Nazis, but many were just scientists, no more no less, who'd simply been doing their job.

The trouble was, it didn't end there. With the help of German-derived

science, America's technological lead over the rest of the world accelerated exponentially after the Second World War. But the black world was a low-grade reflection of the system that had been employed to protect the secrets of the *Kammlerstab* within the confines of the Skoda Works.

The state within a state had been transported four thousand miles to the west and somehow, I just knew, Kammler had come with it.

The intuitive feeling I'd experienced all these years in obscure corners of the U.S. aerospace and defense industry had suddenly acquired a face.

Throughout the next interview, I felt the presence of the info-minders like a low-intensity headache. After the near-informality of my time with Bob Widmer, the proximity of three company caretakers in the room gnawed away at my concentration. They were here to watch over the man who could levitate paper clips on his desk.

But now that the interview was over, the minders were looking the other way. They'd done their job and I'd done mine. Boyd Bushman had spilled no secrets either on or off the record and that was what mattered. Nor had he in any way tarnished the reputation of the Lockheed Martin company.

That I had learned nothing of value must have shown on my face, for without warning, Bushman leaned forward and put his hand on my shoulder.

He asked me what was wrong and I told him.

"It's a lonely walk, but a rewarding one," he said, so quietly that I almost missed it. I looked into his eyes, which were quite blue but for that superficial milkiness that sometimes denotes the onset of old age.

He smiled at me. "Keep traveling the road and you might just find what you're looking for."

"What do you mean?" I asked cautiously.

"In all my years with this company, no one has asked me the questions you came here with today." He paused a moment, then said: "Here, I want to show you something."

He produced a videotape from the folds of his loose-hanging suit and gave me the same smile that had crossed his face a number of times during the interview; usually, I noticed, when he needed a moment or two to think about a particularly awkward question I'd thrown at him. He looked not unlike Walter Matthau, the Hollywood actor, with a kindly, hangdog expression and thick straight hair to match. Instinctively, I'd liked him. But more to the point, I knew that I could trust him. Unusually for someone who toiled in the heart of the U.S. defense-

industrial complex, he looked like someone who was incapable of telling a lie.

Bushman appeared beyond the retirement age of most good man-and-boy company men, which in itself spoke volumes about his ability.

Before I met him, I'd been told that he was a one-off, one of those people who liked to think out of the box—defense jargon for a guy in the trade who looked at things differently. Just how differently, I was about to find out.

The picture on the TV screen steadied and I found myself staring at a scene of such ordinariness I thought for a moment that Bushman had plugged in the wrong tape.

From the graininess of the film and the way the camera trembled, it rapidly became clear that this was some kind of a home movie. It depicted the interior of a garage or workshop. In the middle of the frame was a saw on a pockmarked wooden work surface. Beside it was a pair of pliers and a handful of nails.

For a while, we both just stood there watching a picture in which nothing happened. The minders had taken a quick look and apparently decided it was an eccentric end to a quirky afternoon's work, because they went back to their own conversation.

My gaze drifted back to the TV just as one of the nails in the center of the screen began to twitch and the pliers moved—jumped, rather—a centimeter to the left. Within seconds, everything was shaking, as if rocked by a series of earth tremors. And then, quite suddenly, the nails stood up, like hairs on a cold forearm.

I looked at Bushman, who was studying the screen intently. "Watch," he urged, "this is the best bit."

One by one, the nails took off and shot off the top of the screen. A moment later, the pliers started to do a crazy nose-down dance across the work surface and the saw began flapping around like a fish out of water. Then they both went the way of the nails, shooting out of frame toward the ceiling.

Then, the picture switched: same background, different still life.

Ice cream was creeping up the sides of a see-through tub. A lump of it broke away and rose toward the ceiling trailing little pearls of cream. Seconds later, the whole tub jumped off the bench.

My skepticism was working overtime. But Bushman read my thoughts. "It's no fake," he said. "This is as real as you or I." He cast a look over his shoulder at our companions. "The man responsible is a civilian, nothing to do with the defense business. That he's a genius goes without saying, of course. You needed to see this, because it's

confirmation of the *data*." He emphasized the word as if somehow it would help me understand what I had just seen. "You must get to see this man. I think, then, you will be closer than you think to the end of your journey."

"Data? What data?" I looked at him more closely, hoping for more, but he said nothing.

As I was ushered out of the room, the scientist looked down at the thick pack of material he'd handed me before the interview. I hadn't had time to look at it. Now, I wished I had.

That evening, I sat down with Bushman's file in my hotel room. He'd loaded me down with so many papers, I didn't know where to begin or what I was really looking for. There was a stack of photostats relating to the patents he'd filed during a long and illustrious career that had included stints at some of the giants of the postwar U.S. aerospace and defense industry: Hughes Aircraft, Texas Instruments, General Dynamics and Lockheed.

There was his résumé, which noted his top secret security clearance, as well as copies of scientific papers, stapled neatly at the corner, with exotic-sounding titles that meant little to me.

My eyes swam over pages of complex mathematical equations and algebra sets. I was looking for clues, but seeing nothing that made much sense.

I pressed the "play" button on my tape recorder.

From the tenth floor of my Fort Worth hotel room, the lights of the city shimmered in the sizzling 90-degree heat, even though it was a good half an hour after the sun had slipped below the skyline. Inside, the air-conditioning was cold enough to prickle my skin.

". . . I went back to some of the first work on gravity done by Galileo. . . ."

Bushman's voice crackled in the darkness, clashing for a moment with that of a CNN reporter on the TV. I adjusted the volume on the tape recorder, hit the fast-forward button and pressed play again.

"So, what I think is: Follow the data and log your data very well and don't throw it away because you have a theory. The theory of gravity is just a theory. Einstein improved on the original theories of Newton and he was verified by data—and data gave precedence. No one believed Einstein until the data arrived. Well, our data's arrived . . ."

I scrolled the tape again. As I did so, I spotted something among the collection of papers Bushman had given me. Tucked beneath the patents and company brochure material on weapons technology was a grainy

photocopy of a UFO flying low over a straight stretch of desert road. A handwritten caption underneath identified the location as Santa Ana, California, and the date as 1966.

Flicking past it, I found another cluster of photostats taken from the "Ramayana of Valmiki—Translated from the Original Sanskrit." A stamp on the title page identified the book as belonging to "Lockheed/Fort Worth."

Puzzled, I toggled with the buttons on the tape recorder and finally found the part of the interview I'd been looking for.

"Nature does not speak English," Bushman had been telling me. "Not only that, but if we verbalize it, we're probably approximating, but not telling the truth. Math comes close, but it isn't there either. What Nature tells us is what must be honored. It has been talking to us on many domains."

I let the words echo in my head for a moment.

Bushman was a senior scientist for one of the world's biggest defense contractors; a member of the fraternity that I'd looked to for expert comment all my professional life, and he was telling me to trust in what I saw. Follow data, he'd said, not theory. He worked on programs that lay beyond the cutting edge of known science.

"I can't talk to all the theoreticians," he'd told me, "because there don't exist theories where I am."

For the past ten years, I *had* been following data and the data had shown me the existence of something real. And yet some of the brightest minds on the planet were struggling to produce theory to underpin it. How could there be no theory for effects that were genuinely manifesting themselves in the environment?

What Nature tells us is what must be honored. It has been talking to us on many domains.

And we'd only understood a fraction of what it had been saying.

If we verbalize it, we're probably approximating, but not telling the truth.

And then it hit me. That's all theory was or ever had been—an approximation of the truth. Sometimes that approximation worked out and held fast, becoming scientific currency; other times it didn't and was superseded by another theory.

In this case, there were so many holes in our knowledge, it had allowed for people like Podkletnov and Schuberger to fall through the gaps, to show us things that science couldn't account for. Because science theory hadn't mapped them yet.

Not outside Nazi Germany, at any rate.

Einstein had followed Newton and data had verified them both. But in this field, as with other fields of scientific endeavor, the data was still coming in.

It rendered NASA's theory, in this particular area, at least, quite meaningless.

I flipped another piece of paper in Bushman's file and saw a man with a mop of unkempt hair staring at me from a grainy photostat. He was standing in the same garage I'd witnessed in the film Bushman had shown me at Plant 4.

A note told me his name and that he lived in Vancouver—a place, if Bushman was right, that one way or another looked like being where the road ran out.

Chapter 25

George Hathaway, a science and engineering Ph.D. who ran a consulting firm on "nonconventional propulsion technology" in Ontario, Canada, offered a suitable staging post between the reality of sorts that exists in London and the world inhabited by the man in the photostat that Bushman had shown me: self-trained scientist and inventor, John Hutchison.

I met Hathaway in the coffee shop of my Toronto hotel and was gratified to note that there was nothing remotely flaky about him. Hathaway, whom I guessed to be in his early 50s, looked like most people's idea of a slightly unconventional nine-to-five scientist: a tall man, dressed in jeans and a tartan shirt, with wire-framed glasses, Bill Hickok mustache and hair over the collar. The briefcase gave him a suitably businesslike air as well; he was someone you could talk turkey with.

Hearing him talk, I began to feel a lot better.

I was booked on a flight that afternoon to Vancouver, but had been filled with doubts about the whole trip ever since my arrival in Canada.

It was three weeks after my visit to Fort Worth and I had flown halfway around the world on my own ticket to meet a man who claimed to be able to levitate objects with equipment he'd salvaged from electrical shops and thrift stores.

I had managed to grab myself two days off work. Two days to get to Vancouver and back via Toronto. The ticket had cost me a small fortune.

Faced with a sizable hole in my finances, the whole thing suddenly seemed to stretch credulity, even if it had come with an endorsement from Bushman.

Worse, I'd been unable to make any kind of contact with Marckus.

As was his wont, he had vanished off the face of the planet. I'd called, left messages, sent emails, but he wasn't answering. In the end, I had been unable to put off the opportunity to travel any longer.

Hathaway's ability, therefore, to converse in hard-and-fast science and engineering terms about Hutchison came as an enormous relief. The key point was that he and a colleague, a South African called Alexis Pezarro,

had believed in Hutchison sufficiently to back him financially for a considerable chunk of the 1980s. And even though they weren't linked in any business sense anymore, Hathaway still professed to be a fan of his work.

"There are three kinds of people involved in the advanced propulsion field in my experience," Hathaway had told me over the phone a few weeks earlier. "There are the professionals, people like Marc Millis at NASA."

Millis. The guy who ran the space agency's Breakthrough Propulsion Physics program out of a lonely office at the Glenn Research Center in Cleveland, Ohio. I cast my mind back. It seemed like a lifetime ago.

"Then there are the Tesla types who have an intuitive grasp of electromagnetics who come in from the ground up," Hathaway continued. "And then there's John Hutchison who doesn't fit into either category."

Hutchison, it was more accurate to say, fitted into no real category at all.

He had been born in 1945 in North Vancouver and had grown up an only child, surrounding himself with the only things that he seemed to show any real empathy for—machines. Reading between the lines, it was clear that he had not fitted in as a kid. He'd dropped out of school at Grade 10 and had gone on to complete an education of sorts with the aid of a private tutor (mirroring, I recalled, the schooling regime that Thomas Townsend Brown had endured 40 years earlier).

Amassing a large collection of machine tools, steam engines, old guns, chemical equipment and electromagnetic gear, Hutchison became infatuated from a young age with the theories of Einstein and Faraday, even though he was not remotely academic. His interest in them stemmed from his intuitive grasp of science in general and of electromagnetism in particular. His great hero was the turn-of-the-century electrical engineering pioneer, Nikola Tesla.

I had done my homework on Tesla, because Bushman's papers had been peppered with references to him.

Tesla was a Serb engineering graduate from Prague University who in 1884 at the age of 28 emigrated to America in the hope of finding work in the then fledgling electricity industry. With nothing more to his name when he arrived in New York than a letter of introduction to the man widely regarded as the father of the electricity industry, Thomas Edison, Tesla rapidly established himself in Edison's eyes as an engineer of great skill.

But while Edison used him primarily as a glorified Mr. Fixit to repair and refine his power generators, which were by then beginning to proliferate across the eastern seaboard, Tesla had set his mind on introducing

an electricity supply system that was infinitely better than Edison's. If Edison would only use his, Tesla's, method of alternating or AC current instead of direct or DC current, which was notorious for its inability to travel over long distances, Tesla knew he could deliver a system that was far more efficient.

Yet, by the mid-1880s, Edison's entire infrastructure—indeed, his entire business base—revolved around the supply of DC power, whatever its manifest deficiencies. As a result, Tesla left the Edison Electric Company to set up on his own.

From then on, his career was dogged by a series of business disasters, but as an inventor—and the man who gave the world AC current into the bargain—Tesla was without parallel. In 1891, he developed the Tesla Coil, a remarkable invention that remains the basis for radios, televisions and other means of wireless communication.

But it was the wireless transmission of energy that became his great goal and for most of his professional life Tesla dedicated himself to the pursuit of a system that promised to give the world a free supply of energy. It hardly seemed to enter his head that big business, which by the early 20th century was starting to make handsome profits out of electricity, would be resolutely opposed to the idea.

Tesla's wireless energy transmission system was initially based upon technology for sending power through the air, but he soon developed a far more efficient transmission medium: the ground. In a series of experiments in Colorado, Tesla showed that the earth could be adapted from its customary role as an energy sinkhole—a place to dump excess electricity—into a powerful conductor; a giant planetwide energy transmission system that obviated the need for wires.

Tesla died penniless in 1943, mumbling to the end about fantastic beam weapons that could blast aircraft from the skies or be bounced off the ionosphere to strike at targets anywhere on the globe. One of the wilder theories that still cling to his memory revolves around a supposed experiment to beam some kind of message to the Arctic explorer Robert Peary, who in mid-1908 was making an attempt to reach the North Pole. According to legend, Tesla, who had built a powerful transmitter at Wardenclyffe on Long Island, beamed the "message" on June 30 and awaited news from Peary. The explorer saw nothing, but thousands of miles away, in a remote corner of Siberia, a massive explosion equivalent to 15 megatons of TNT devastated 500,000 acres of land centered on a place called Tunguska.

The Tunguska "incident" is generally ascribed to the impact of a comet fragment or a meteorite, but the absence of evidence for the impact

theory has enabled Tesla proponents to maintain the line that it was Tesla's "death ray" that caused the blast. Certainly, Tesla himself seemed to believe he was responsible, for immediately afterward he dismantled the "weapon" and reverted to other pursuits.

What Tesla and Hutchison shared was a complete sense of ease at being around electricity. Tesla, for example, knew it was perfectly safe to pass high voltages through his own body, provided the current was kept low. Using himself as a conductor, he would stage dramatic demonstrations aimed at validating the safety of AC power. In one of them, he would hold a wire from a Tesla Coil in one hand while drawing a spark from the fingers of the other to light a lamp. The newspapers of the day would frequently run photographs of him in morning dress, his whole body alive with sparks. It must have seemed like magic, but was nothing more than straightforward evidence of Tesla's deep-seated feel for his subject.

It was the same with Hutchison. By the age of 15, he had already assembled his own laboratory, cramming it with Tesla machines he had constructed himself; signal generation equipment, static generators and a uranium source housed in a copper tube that used a magnetron pulsed by an old rotary spark plug system to transmit microwaves. During the 1970s, Hutchison set up shop in a bigger, more sophisticated laboratory in the basement of a house in Lynn Valley, North Vancouver, which contained racks full of signal generators, low-power radar systems and phasing equipment coupled to refinements of his earlier devices.

One evening in 1979, Hutchison was in his laboratory, sitting amid the sparks and high voltage effects of his equipment, when he was struck on the shoulder by a piece of metal. He threw it back and it hit him again. This was the ignominious beginning of the Hutchison Effect. In the ensuing months, Hutchison found that by tweaking the settings on the equipment, he could get things—ordinary household objects—to levitate, move horizontally, bend, break and even explode. In the latter case, this portion of the effect came to be known as "disruption."

In 1980, Hutchison went into partnership with a company called Pharos Technologies, which had been formed by Hathaway and the South African Pezarro. Pharos Technologies was designed exclusively to promote the Hutchison Effect, which in these early days was billed as a "lift and disruption" system. This, however, barely begins to describe some of the extraordinary phenomena Hutchison was manifesting in his Lynn Valley basement.

Using only tiny amounts of power—between 400 and 4,000 watts via a 110 AC wall socket—Hutchison found he could raise any kind

of material—sheet metal, wood, styrofoam, lead, copper, zinc . . . whatever—from a few ounces to a hundred pounds in weight at distances of up to 80 feet. From a physics standpoint, it was meant to be impossible, of course, but Hutchison was doing it—regularly, if not exactly on demand. Nine times out often, nothing would happen. The rest of the time it was like a bad day on the set of the film *Poltergeist'*, objects rising slowly from the area of influence in the center of the equipment and looping back somewhere else; or shooting skyward in a powerful ballistic arc and striking the ceiling, kicked there by a massive energy impulse; or simply levitating and hovering, continuously, for ages.

What was more, Hathaway said, it had been captured on hours of film and videotape and had been demonstrated in front of hundreds of witnesses.

As for what might be producing the Effect, there were many theories. Some said that it was triggered by opposing electromagnetic fields that canceled each other out, creating a powerful flow of zero-point energy to any object in the zone of influence; others that Hutchison was causing electromagnetic fields to spin or swirl in some unknown way and that this was the trigger for lévitation—shades of the torsion fields generated by Schuberger, Podkletnov and the Bell. Another camp maintained that Hutchison was generating the Effect himself by psychokinesis.

What was interesting, Hathaway said, was that Hutchison himself did not know what caused it and he had no control over it once it started. Sometimes the Effect produced lévitation, sometimes it caused objects to shred, tear apart or evaporate; other times, it would cause transmutation—the alteration of an object's molecular composition; Hutchison, like an alchemist, could change one metal into another.

"You're shitting me," I said.

But Hathaway shook his head. "John's a wild and crazy guy. He lives in his own world and he can't readily express what he's doing in terms an engineer or a physicist might always understand. You should go see him, no question." He paused, then added: "A visit to John is definitely a once-in-a-lifetime thing."

As the Canadian Airlines A320 tracked westward above the Great Lakes, I sat alone at the back of the aircraft, my mind still beset by doubts, in spite of Hathaway's no-nonsense assessment of the man I was about to meet.

The one thing that kept me going was the germ of a mystery rooted in some hard, incontrovertible fact I'd managed to glean separately from Hutchison and Hathaway: in 1983, a Pentagon team spent money on

Hutchison—tens of thousands of dollars over a period of several months—analyzing the Effect.

In promoting Hutchison's technology to potential investors—people to exploit its potential as a propulsion or "disruption" system—Hathaway and Pezarro had come into contact with a unit inside the Pentagon that was prepared to take their claims at face value.

Led by Col. John B. Alexander of the U.S. Army's Intelligence and Security Command, INSCOM, the team put up money to move Hutchison's cramped lab at Lynn Valley to a large empty warehouse on the edge of North Vancouver.

The facility was set up under tight security and anything pertaining to it was classified.

The "INSCOM group" tasked with analyzing the Hutchison Effect comprised five people besides Hutchison, Hathaway and Pezarro—two scientists from Los Alamos National Laboratory, a representative of the Office of Naval Research, an Army R&D specialist and Alexander himself.

The presence of the Los Alamos and ONR representatives appeared to be significant, since both of these organizations were driving forces behind the Strategic Defense Initiative. SDI—"Star Wars," as it had been dubbed by the media—had been the brainchild of Ronald Reagan's science advisers and Reagan had hawked it to the rooftops as the be-all and end-all arms program of his presidency: a space-based ring of steel comprising sensors, laser weaponry and missiles that would protect the United States from an all-out nuclear strike by the Soviets.

SDI was unveiled in March 1983 and from then until its death later in the decade was heavily staffed by experts from Los Alamos and ONR.

For the life of me, I couldn't think what Hutchison's work had to do with it.

Soon after Hutchison's lab was moved to the large warehouse on the edge of North Vancouver, the experiments began in readiness for the INSCOM visit. The Pentagon's funding was good for four months. Alexander had provided Hutchison, Hathaway and Pezarro with objects on which to test the Hutchison Effect in its new location. One of these was a six-by-quarter-inch molybdenum rod. Molybdenum is an incredibly strong but light metal used in nuclear reactors. It has a "sag" temperature of 3,700 degrees C. Alexander had deliberately marked the rod so it couldn't be switched. Give it a whirl, he told Hathaway. See what happens.

A few days later, the colonel was contacted by Hathaway and told that the test runs had been spectacular. The test specimens were dispatched to the INSCOM team for analysis forthwith.

The Effect had snaked the molybdenum rod into an S shape as if it were a soft metal—lead, tin or copper. There were other successes to report, too. A length of high-carbon steel had shredded at one end and transmuted into lead the other. A piece of PVC plastic had literally disappeared into thin air before the eyes of the evaluators. Bits of wood had become embedded in the middle of pieces of aluminum.

And, of course, things had taken off—all over the laboratory, items had been levitated; some deliberately, others in parts of the lab that were nothing to do with the experiment. The lab at one point had also become self-running, drawing its electricity from God only knew where—since the power was down at the time.

Keep everything exactly as it is, Alexander told him. We're on our way up.

When the INSCOM team got off the plane in Vancouver, everyone was gibbering with excitement about the looming set of tests—everyone, that is, except for the more senior of the two Los Alamos scientists, a dour man who'd been given the unaffectionate nickname "Big Bad Bob" by the other team members.

Bob remained convinced that somehow it was all just a bunch of jiggery-pokery.

Someone, he'd told Alexander, was being had here.

A shimmy through the airframe as the A320 thumped down on the tarmac at Vancouver and here I was, same place as the INSCOM group, different time.

Like them, I was thinking of the encounter to come; and for a reason I couldn't entirely put my finger on, I felt anxious.

I set out from the hotel for Hutchison's apartment 12 hours later.

It was a crisp morning, warm air from the Pacific mixing with the cold air of a blue sky to form a thick layer of cloud in the waterways of the harbor below. I followed Hutchison's directions, turning left on a wide run-down street, eventually spotting what I was looking for: a brown apartment building with an antenna farm on one of the balconies.

I could see Hutchison in silhouette at one of the windows, scanning the street for my arrival.

He met me at the entrance and we shook hands. He was dressed in jeans and a denim jacket. A strand of thick black hair fell over one lens of his glasses. The other had a crack running through it. He was tall and powerfully built, but his voice was softer than I remembered it on the phone. Through the handshake I could sense he was as restless as I was. We exchanged a few pleasantries, then I followed him inside. A minute

later, there was a rattle of keys and tumbling of locks as Hutchison pushed back the door of his apartment.

The door traveled about 18 inches, then stopped abruptly.

"Sorry, it's kind of a squeeze," he said, forcing his way through the gap and disappearing inside. I followed suit and found myself in a dark, narrow corridor with what looked like crates lining the walls. The gap down the middle was barely wide enough for a child. Hutchison pushed the door shut and somehow managed to squeeze past me, heading for the back.

He pulled a drape off the window and light flooded the room. Outside, the aerials and dishes I'd spotted from the street pointed skyward at crazy angles. It was only when I turned back to study the apartment interior that I realized the crates I'd seen were radios—huge, metallic, gray, lifeless things, some the size of tea chests. I peered into the kitchen, the bathroom and bedroom and saw more radios.

Every available inch of floor and wall space, in fact, was filled with radios and radar screens, gear that had gone out with the Stone Age, complete with dials, valves and analogue readouts.

It was like stepping into a cross between a military surplus store and Radio Shack circa 1952.

Hutchison lowered his huge frame to the floor and assumed the lotus position on the chocolate-colored carpet. He had an almost childlike innocence about him. He'd never once asked why I'd come all this way to speak to him and he gave me the impression that only a small part of his mind was here in the room with me.

The equipment, he told me, had been requisitioned from some '50s-vintage Canadian warships he was helping to dismantle in the harbor. In return for his salvage services, the Canadian military said he could have all the stuff he was able to carry away with him. Clearly, they'd underestimated his strength.

Radios, sonar systems and electronic countermeasures devices lined shelves suspended from the ceiling. Hutchison told me he'd consulted the plans of the apartment block and had drilled holes in all the right places.

"There's a ton and half of the stuff above your head, but it's safe enough," he smiled. I glanced up nervously and noticed a heavy-looking cluster of boxes suspended over me, then asked if this was the equipment that had been used in the INSCOM tests. Hutchison shook his head. Much of it had been confiscated by the Canadian police and after their intervention, he'd found it hard to carry on, though recently he'd rebuilt the lab and was readying to start operations again.

He showed me an aluminum ingot that looked as if it had blown apart

from the inside and a file filled with letterheads I recognized: Boeing and McDonnell Douglas had been but two of Hutchison's backers.

But he was getting ahead of himself and so was I. First things first. "What happened after the arrival of the INSCOM team?" I asked.

He scrolled back 15 years and told me about the plow.

The building had been unremarkable, another storage facility in the outer suburbs. There was no security, just an eight-foot wire fence and the warehouse's relative obscurity amid the smoke and grime of Vancouver's industrial quarter. It was not optimum, but guards would only have attracted attention.

The equipment was housed behind a wall of breeze blocks in the middle of the warehouse. The breeze blocks extended halfway to the roof—around 15 feet.

This was the target area.

Behind the wall was the large Tesla Coil containing the uranium source, a device around four and half feet tall with a doughnut-shaped metal coil on top.

Diagonally across the room, on the other side of the target area, sat a powerful Van de Graaf generator capable of producing a 250,000-volt DC static charge.

The other prominent piece of technology was a three-and-a-half-foot double-ended Tesla Coil, known as the "dumbbell," suspended from supports running across the top of the wall.

Between these three main equipment items were an assortment of smaller devices, all connected to each other by multiple coils of wire and cabling.

There were tuning capacitors, high-voltage transmission caps, RF coils and a spark gap that would snap every 40 seconds or so sending an earsplitting shock wave throughout the building while it was up and running.

Now, though, as the equipment warmed up prior to the test, all it emitted was a low-intensity hum.

Outside the target area lay a pile of unclaimed scrap metal. Leaning against the wall were three old streetlights and a spool of wire hawser. Set in front of them was a rusting horse-drawn plow, three times the weight of a man.

In the flickering light of the monitoring screens members of the team would see their shadows moving on the edge of their vision. Several said it made it feel like there were people there, watching them.

Between the scrap metal and the target area a bank of receivers and monitoring equipment rested on a workbench.

Hutchison, then in his late 30s, sat at the bench, watching monitors and tuning dials.

The Pentagon and DoE evaluation team had flown in from across the border the previous night. The atmosphere was relaxed as it could be—shirtsleeves, first-name stuff—but everyone was tense. No one knew what to expect; no one, that is, except Big Bad Bob, the crusty old skeptic from the Los Alamos National Laboratory.

What was happening here, Bob told anyone who'd listen, was just a bunch of smoke and mirrors, a waste of fuckin' time, effort and money.

His sidekick, John, was blessed with a more open mind and, given the nature of the assignment, he expressed the view, though not when Bob was around, that this was no bad thing.

Only a few days earlier, tongues of fire had licked their way up through the concrete floor, a manifestation of the Effect that had not been encountered before.

Everyone except Bob had accepted the Canadians' version of events at face value.

As Hutchison adjusted the monitoring equipment, he could hear the ladder behind him groaning under Bob's considerable bulk. Bob was gazing down on the equipment from the top of the wall. (Hutchison never made clear to me the reasons for the wall. Had it been put there to contain the Effect or possible fallout from it—an explosion perhaps?) Colonel Alexander, who was pacing in the shadows to the left of the workbench, asked him what he could see.

The equipment was humming and the cameras were turning. The target, a length of steel rod, wasn't moving—wasn't even quivering—so no friggin' surprise there, Bob replied.

Just then, the radio crackled. Bob's deputy was running some checks with special monitoring equipment outside the building. He was 300 feet from the warehouse and picking up some kind of distortion. Something, he said, was happening.

Alexander stepped out of the shadows and studied one of the monitors. He peered at the readings, then glanced at the video image of the steel bar, but it was just as Bob had said: nothing was moving. Nothing had changed.

Suddenly, without anyone touching anything, the ceiling lights switched on and began to glow intensely bright. For a moment, the entire floor area was bathed in incandescent white light. Then the bank of bulbs blew, sending a shower of red-hot filaments and glass onto the target area.

Except for the glow from the monitors, the warehouse was plunged

back into darkness. Bob, who was standing a little to the right of Hutchison, started to laugh.

Out of the corner of his eye, over by the scrap area, the Canadian caught the movement, real movement this time, and instinctively ducked his head.

The crash was followed by a cry from Big Bad Bob, then silence.

The plow had sailed across the room at shoulder height and buried itself into the wall close to where he had been standing.

After the INSCOM team departed, Hutchison's work continued. The lab moved again to a different part of Vancouver and people came and went.

This much I know to be true, because Hutchison showed me copies of the reports.

Among Hutchison's visitors at this time was Jack Houck of the McDonnell Douglas aerospace corporation, who in 1985 spent two days analyzing the Effect. Houck came away satisfied that no fraudulence had occurred during the experiments he attended, during which, in a subsequent report, he pronounced that "some very interesting events" had been captured on videotape. However, he went on, "some of the biggest events occurred outside the target area. The first evening a gun barrel and a very heavy (60-pound) brass cylinder were hurled from a shelf in the back corner of the room onto the floor. Simultaneously, on the opposite side of the room toward the back, three other objects were hurled to the ground. One was a heavy aluminum bar (% inch by 2% inch by 12 inch). It was bent by 30 degrees."

Houck attested to the randomness of the Hutchison Effect and postulated that it might in part be attributable to psychokinesis—that Hutchison, in other words, was either boosting the energy generated by the machinery with his mind or that his mind alone was responsible for the manifestations produced.

These phenomena, Hutchison was now able to report, included time-dilation—pockets of altered space-time within the target area—and, most extraordinary of all, the capacity of the Effect to turn metal ingots transparent. It was as if, momentarily, the ingots were there, but not there; visible in outline yet totally see-through.

It had been Hathaway's dream in the early stages of their work to develop Hutchison's machinery into a "stationary launch-assist" device: an antigravity aerospace platform.

This was based on his initial belief that the inertial properties of the machinery itself appeared to have altered during some of the experiments; this, in addition to the clearly altered weight condition of the

objects placed in the target area—objects that I'd seen in the video Bushman had shown me in Texas.

But in 1986, Hutchison and Pharos Technologies went their separate ways, Pezarro and Hathaway having failed to secure the kind of backing for the Effect they'd always hoped for. It all fell apart, Hathaway said, when it was suggested that Hutchison "was an integral part of the apparatus"; that the Effect, in other words, was unachievable without the presence of Hutchison himself.

For his part, Hutchison vehemently denies this, but the fact that the most spectacular manifestations of the Effect always seemed to occur when he was around is hard to refute. The psychokinesis proponents hold this up as evidence that their theory is the right one; the technologists counter that it is Hutchison's intuitive feel for the machinery, his ability to tune it to the hairsbreadth tolerances required for things to start happening, that makes his presence necessary.

The fact that two aerospace companies—Boeing and McDonnell Douglas—felt it necessary to investigate the Hutchison Effect is at least telling.

Boeing partly funded a series of experiments during the late 1980s. By then, however, Hutchison had become disenchanted with the Canadian scene and in 1989 went on a scouting tour of Europe intent on moving his lab to either Austria or Germany. When he came back to Vancouver at the turn of the decade, he found the Canadian government had confiscated half of his equipment. The Royal Canadian Mounted Police maintained that it constituted an environmental hazard; Hutchison thought the Canadians had been leaned on by the Americans.

Hutchison's suspicions were compounded when attempts to have the Los Alamos report released under the Freedom of Information Act resulted in failure.

No one in officialdom had been able to locate the files. It looked as if the report had been buried.

Later, Colonel Alexander, the head of the INSCOM group, told me that the report had been classified, but was subsequently downgraded to "confidential." In the end, it had been "routinely destroyed."

Alexander's take on the "plow incident" was also markedly different from Hutchison's. On the day in question, he said, nothing mysterious had happened. That, from INSCOM's point of view, was the problem.

I asked Alexander what he felt about it all.

"Oh, it's real all right," he told me, "the trouble is, sometimes it works and sometimes, it doesn't. But four out of five of us came away believing."

Chapter 26

I met Marckus in a country pub close to the haven on the estuary where we'd hooked up at the very beginning. It had been several years since our initial rendezvous, but when I walked in and saw him in the corner, time might as well have stood still. He looked up from his paper, gave me a wave and clicked his fingers for some attention from the bar. Other than a bored-looking girl behind the pumps, there was no one else in the room.

After she'd taken my order, she came over and set my beer down on Marckus' paper. Marckus clucked like an old hen and handed the glass to me. The circle of condensation, I noticed, had partly ringed the word "doom" on a headline about the forthcoming earth summit. It felt about right. It had been raining solidly for months.

"So you know," he said with an air of finality. "Good. I can't stand awkward explanations."

Before departing for Canada, when I'd been desperate for him to brief me on Hutchison, I'd left messages on his answering machine telling him I wanted to talk about *Blowback*—the book he had lent me about U.S. recruitment of high-value Nazis. Marckus must have figured out that I'd known then.

"Where have you been?" I asked him.

"I decided to make use of some vacation time that's been owing to me." He held up his hands. "I'm sorry. I should have told you."

"So where did you go?"

He looked down at his beer. "I've been following in your footsteps, paying a few respects. Bavaria is beautiful this time of year. From the little hotel where I was staying you can see the snow on the mountains. Maybe on a clear day, if you could only see through the wire, you might have glimpsed the same view from Dachau."

"How many members of your family did you lose?" I asked.

"A couple that I know of. An uncle and aunt on my father's side. Most of me is as English as you are." He paused. "I'm sorry I put you through the hoops. It must all seem a little trite, I suppose. But it was important to me that you found things out for yourself."

"When you started out on the path that you did, sooner or later the trail was going to lead you to Germany. I was happy to help you with the science, but it was important that I held back certain kinds of information. It has taken me decades to build the knowledge that you have acquired in just a few years. But if I'd told you the full story from the outset, you'd have written me off as a crank."

"Do you think the Americans have mastered the technology?" I asked.

He shrugged theatrically. "We're going in circles again, aren't we? I told you when we first met. I don't know. I know—you know now—that they have access to it, but whether they've *mastered* it or not . . . it depends on what you mean by 'mastered.' "

He took a sip from his drink and continued. "When the Americans tripped over this mutant strain of nonlinear physics and took it back home with them, they were astute enough to realize that their home-grown scientific talent couldn't handle it. That it was beyond their cultural terms of reference. That's why they recruited so many Germans. The Nazis developed a unique approach to science and engineering quite separate from the rest of the world, because their ideology, unrestrained as it was, supported a wholly different way of doing things. Von Braun's V-2s are a case in point, but so was their understanding of physics. The trouble was, when the Americans took it all home with them they found out, too late, that it came infected with a virus. You take the science on, you take on aspects of the ideology, as well."

"And the black world?" I prompted.

"Well, there are different shades of black, of course. But what interests me is that the bit that is truly unaccountable is the part of the system that's been exposed longest to the virus. So in time, it will simply destroy itself. That, at least, is my earnest hope."

Marckus stared out of the window a while, then shook himself like a dog. "Tell me about Canada."

I took a pull of my drink and set the glass down. I felt tired. "There isn't a whole lot to go by. After the Royal Canadian Mounted Police paid him a visit, Hutchison doesn't even have the equipment to do lévitations or transmutations anymore."

He glanced up. "So, you believe him."

"I believe that he caused objects to lose their weight. I also believe he made iron bars turn momentarily transparent and I believe that he transmuted metals." I tried to rub some of the tiredness away, but the alcohol wasn't helping. "Beyond that, I don't know."

Marckus fell silent for a few seconds, then he said: "Do you have any idea how much energy it takes to transmute a metal?"

At the back of my head, I seemed to recall Hathaway talking in terms of a gigawatt or two. I found it difficult to visualize a gigawatt and said as much to Marckus.

I didn't want another physics lesson, but I realized too late I was going to get one.

"At the low end of the scale, Hutchison was using 400 watts to achieve the effects you're talking about," Marckus said. "Four hundred watts are the heat and light you get from four light bulbs." A kilowatt—a thousand watts—represents the heat you get from a one-bar electric fire. A thousand kilowatts are a megawatt and a thousand megawatts are a terawatt. A thousand terawatts, Marckus told me, were a gigawatt.

"You know now what a gigawatt is?" he said. "A gigawatt is the equivalent energy release of the Nagasaki bomb. And that's what Hutchison was pulling out of his 110-volt AC wall socket to turn steel into lead."

"Transmutation is real. The reason most people don't use it to turn lead into gold is that the power requirements cost more than the gold."

"Why would INSCOM be interested in alchemy?" I asked.

Marckus smiled again, but the look in his eyes was intense. "Transmutation has other purposes. You can use it to change the rate at which radioactive elements decay or you can turn the whole process around."

Marckus was putting me through the hoops again.

"Go on," I told him.

"Think of it," he said, leaning forward, "transmutation, if you can do it the Hutchison way, is a cheap method of enriching uranium, for example. There's another thing, too. Remember when you were heading for Puthoff's place and you asked me for a no-brainer on the power potential of zero-point energy? I gave you the shoebox analogy and how much untapped energy there is in it?"

He swilled the dregs of his beer around, drained them, then set the glass down and looked at me. "Maybe Hutchison doesn't know it. Release that energy slowly and you've got a safe, clean reactor that can go on pumping out power forever. Speed up the process and make all the right connections and you've got a bomb; one that'll make a thermonuke look like a child's firecracker. No wonder they shredded the report at Los Alamos."

I was about to remind him that it had merely been "routinely destroyed," but stopped myself. Standing next to the coffee machine at the Institute of Advanced Studies in Austin, Puthoff had coolly relayed his belief that there was enough energy in my cup to boil the world's oceans many times over. And then I remembered the Schaubingers' fears over the Americans' hijacking of the implosion process—its potential use in the development of a giant bomb.

I could feel my heart in my throat.

"When he does what he does," Marckus said, "Hutchison is reaching into a place that you can't see or touch and he's pulling things out we call 'phenomena.' " He turned his index fingers into quote marks to make the point. "But things don't happen by magic. Phenomena happen because there are laws of physics we don't understand yet. The fact we haven't found the answer yet is our problem, not Nature's. But we're working on it." He stared at me and frowned. "Why on Earth are you smiling?"

"Because you just nailed something down for me. Something that's bugged the hell out of me almost from the very beginning."

"What's that?"

"The Philadelphia Experiment. A neat piece of disinformation, I think you'll agree, but stuffed full of real elements that make it impossible for people like you or me to conduct a meaningful investigation of anything remotely related to it. I'm thinking of T. T. Brown, electrostatics and stealth, of course, but there was always that one part of the story that didn't seem to fit—the part about the ship slipping into another dimension. Even that, it turns out, has a grain of truth in it."

Marckus nodded. "The best disinformation always does."

Bushman *was* right. Everything had been right there in Hutchison's apartment.

No wonder INSCOM's weird-shit division had shown up at his door.

Sooner or later, someone would succeed in developing a zero-point reactor—a machine that would be able to exploit the fluctuations in the quantum sea as they blinked in and out of existence millions of times every second. It was just a matter of time. "We are at the stage that the fathers of the atomic bomb were at when they put together their first test nuclear reactor in the early Forties—and look what happened a few years later," Puthoff had been quoted as saying recently.

I offered Marckus another drink, but he looked at his watch and said that he had to get going. The wind had started to lash against the window, bringing more rain with it. Marckus pulled up the collar of his coat and readied himself to leave.

But I hadn't quite finished with Dan Marckus just yet.

"So what's it to be?" I asked him. "Are you going to build a bomb or a reactor?"

Marckus made an exaggerated display of busying himself with his coat. He said nothing, so I asked him again.

"I'd like to know what you're going to do with this knowledge, Dan. You know as well as I do that radical technology that destroys existing technology isn't welcome."

"Let me put it another way," he said. "Years ago, a story appeared in one of those pulp American science-fiction rags, *Astounding Science Fiction* I think it was, in which the U.S. military authorities gathered America's best scientific brains around a table and told them to develop and build an antigravity machine, because it was known that the Russians had already developed something similar. The twist in the tale, of course, was that the business about the Russians was a total piece of fiction, but not knowing this the scientists went away, put their heads together, and came up with a crude, but workable device.

"The point of the story is that nothing is impossible, once it is known to be practicable. I have something infinitely better than that. Physicists have proven the existence of the zero-point energy field and your data shows that people have actually developed antigravity devices that flew. I want you to know that you weren't simply duplicating knowledge that I already had. Let's say I had the big picture and you provided the details. It's the details in this business that are important. What we're faced with here is a simple set of alternatives. Free, clean energy for everyone on the one hand or the biggest bloody bomb you can imagine. I want the good guys to be in the van here, because a day doesn't go by that I don't think about my family and what happened to them under the Nazis."

Marckus strolled across the parking lot, heading for a pathway through the woods on the other side. There was something nagging at the back of my head that I'd meant to ask him, but I couldn't think of it. I closed my eyes, but the inspiration wouldn't come. When I opened them again, he had vanished among the trees.

It was then, of course, that it came to me. I'd meant to ask if he had sent the article that had landed on my desk at Jane's all those years ago.

I smiled to myself. That would have been too Machiavellian; even by Marckus' impressive standards. But I would ask him all the same.

Twenty minutes later, I eased my car along the narrow road that ran beside the abandoned radar establishment and parked up close to the jetty where Marckus and I had first met. The rain had passed, leaving the unmistakable smell of spring in its wake. I got out of the car and gazed skyward. High above a checkered layer of cirrocumulus, an aircraft contrail tracked westward toward America, the silver arrowhead at the point of its creation looking tiny and vulnerable against the deep blue. For years, I had banked on the presumption that technology would pursue a set course—one that had been preordained ever since someone had had the wit to fashion a wheel out of a piece of flat stone. Now I knew that it didn't have to be that way, that there were shortcuts in the process

that would allow us to leap the state of the art in moments of searing inspiration. I thought back to the day that Garry Lyles at NASA had outlined his vision of interstellar travel, the fast trip to Alpha Centauri that he earnestly hoped man would make within the next 100 years.

If we looked in the right places, we might just be making it a lot sooner.

Epilogue

A few final thoughts, the first of them regarding George S. Trimble Jr., the man whose comments on the imminent "reality" of gravity control provided the trigger for the investigation behind this book. In 1967, Trimble left the Martin Company for NASA, where he became director of the Advanced Manned Missions Program. His reasons for not wanting to be interviewed about his activities at Martin Aircraft deep into his retirement remain unclear, although, of course, they may be entirely innocent. As an ironic twist on everything that followed, I recently came across a comment of his: "The biggest deterrent to scientific progress is a refusal of some people, including scientists, to believe that things that seem amazing can really happen," the *New York Herald Tribune* quoted him as saying on November 22, 1955. "I know that if Washington decides that it is vital to our national survival to go where we want and do what we want without having to worry about gravity, we'd find the answer rapidly."

Enough said.

In October 2000, the U.S. magazine *Popular Mechanics* revealed that Ning Li, the Chinese-American scientist working in an area of endeavor similar to Dr. Evgeny Podkletnov, was close to developing an operational machine capable of exerting an attractive or repulsive force on all matter. By using 12-inch disc-shaped superconductors and around a kilowatt of electricity, Li's device would produce a force field, the magazine stated, that would "effectively neutralize gravity above a 1-foot-diameter region extending from the surface of the planet to outer space." Li left the University of Alabama at Huntsville in mid-2000 to pursue the commercial development of her invention and has kept a low profile since. While her eye is said to be on civilian spin-offs of this remarkable breakthrough—NASA scientists at the Marshall Space Flight Center believe her "AC Gravity" machine could be used as a shield for protecting the International Space Station against impacts by small meteorites—it goes almost without saying that it could also be hijacked as a weapon. The same force that can deflect a meteorite or a piece of

orbiting space junk can also stop a reconnaissance satellite or a ballistic missile dead in its tracks. Hit a foot-wide gravity-reflecting beam at an orbiting velocity of 17,000 miles per hour and it would be no different to hitting a brick wall, except there would be no evidence to say that you had hit anything. The results, to borrow an expression from science-fiction writer Arthur C. Clarke, would be "indistinguishable from magic," the ultimate in weapons technology.

Over recent years, a wealth of articles have appeared outlining the view of an emerging breed of physicists that we may indeed reside in a multi-dimensional cosmos. The August 2000 edition of *Scientific American*, for example, generated a mass of copy in newspapers here and in the States on this growing belief—that parallel universes may exist alongside our own and how additional, unseen dimensions would help to unify the fundamental forces of Nature that inhabit our four-dimensional space-time. According to this thinking, much of it wrapped within so-called "string theory," our universe may simply exist as a membrane floating within a higher-dimensional space, with gravity—this impossibly weak, little-understood yet massively influential force—the only one of the four fundamental forces capable of propagating across the dimensions. Over the course of the next decade, experimentation may provide real answers to these tantalizing and complex suppositions. For me, it has simply served to show how taboo notions—taboo *science*—can, in time, enter the mainstream of ideas.

As I sit here, picking over clippings related to these and other areas of research connected with this book, it has dawned on me only now that attitudes *have* changed in the relatively short space of time it has taken me to write it. Broadsheet newspapers from the *New York Times* to the *Sunday Times* and the *Observer* have all carried thought-provoking features over the past few years on zero-point energy, faster-than-light travel and other contentious areas of science. Suddenly, the idea of gravity having an *antigravity* component—this heresy that terrified the professional daylights out of me a decade ago—doesn't seem so strange after all.

On that note, I should add that this book is in not intended to be a catchall explanation for UFOs. While it may go some way toward explaining some of the thousands of sightings that have occurred since the Second World War—many of them documented in official files—the subject is too complex, too multifarious, in my opinion, to conform to a single explanation. Although the data that Boyd Bushman encouraged me to follow to glean the truth about antigravity is sufficient for me to reach some definitive conclusions on that subject, it is inconclusive, to

my mind at least, on the question of UFOs. For the time being, it must remain a loose end.

Regarding the definitive fate of SS General Dr. Hans Kammler, there are rumors that he died in Virginia—some have it as Texas—but it amounts to insubstantial testimony and to date there is no hard evidence to say that he came to the States, let alone died there. The U.S. National Archives remain devoid of any meaningful files on him.

In April 2001, the CIA provided confirmation of just about everything Christopher Simpson wrote in *Blowback* when under Congressional directive it released 20 files on Nazi war criminals recruited by U.S. intelligence at the end of the Second World War. One of these was Emil Augsburg, an SS officer instrumental in drawing up the "Final Solution to the Jewish Question" at the Wannsee conference in 1941. Another was Gestapo captain Klaus Barbie, the infamous "Butcher of Lyons."

Yet to be confirmed at official levels, though likely to be just as true, is the even less salutary story of Gunter Reinemer, an SS lieutenant in charge of death squads at the Treblinka concentration camp. After the war, the CIA gave Reinemer a whole new identity—as a Jewish Holocaust survivor of all people—before dispatching him into East Germany as a spy. In 1988, a few days before he was found dead, Reinemer was unmasked by a German financial investigator named Dieter Matschke. "You have to ask yourself," Matschke was quoted in the London *Times* on December 15, 2000, "how many other Reinemers did America spirit to safety?"

If the U.S. recruitment program did this for Augsburg, Barbie and Reinemer, it would have bent over backward to accommodate Kammler, keeper of the Third Reich's most exotic military secrets.

Marckus did not place the article on my desk. I checked.

Since the publication of the U.K. edition of *The Huntfor Zero Point*, one other significant postscript to the story is worth mentioning—in time, the word "momentous" may even be appropriate. On March 26, 2002, a U.S. patent was granted for a device called the "Motionless Magnetic Generator," or MEG, that its supporters say will be the world's first commercially available free-energy home-generator. The MEG has been developed by a team of inventors led by long-time zero point energy pioneer and proponent Dr. Thomas E. Bearden, a retired U.S. Army lieutenant colonel and former nuclear engineer. The MEG is designed to provide an indefinite output of 2.5 kilowatts—enough to run a room or two in your house—by tapping into the infinite energy of the quantum sea. Link three or four MEGs together, proponents claim, and you get

enough output to run an entire house. The event is significant, at least, because the U.S. Patent Office traditionally shuns so-called "free-energy" devices; de facto, it has accepted there must be something to this one. As I write this, scant weeks after the patent's acceptance, it remains to be seen whether claims for the MEG will indeed transcend into commercially available hardware. But maybe, just maybe—as I sit here recalling Hal Puthoff's exhortation—3.26.02 will be the date I put in my diary as the day the world changed—forever.

Bibliography

PRINCIPAL PUBLISHED SOURCES:

Books

- Agoston, Tom. *Blunder! How the U.S. Gave Away Nazi Supersecrets to Russia.* New York: Dodd Mead, 1985.
- Barrow, John D. *Impossibility: The Limits of Science and The Science of Limits.* London: Vintage, 1999
- Berlitz, Charles and Moore, William. *The Philadelphia Experiment.* London: Souvenir Press, 1979
- Bower, Tom. *The Paperclip Conspiracy.* London: Michael Joseph, 1987
- Coats, Callum. *Living Energies.* Bath, UK: Gateway Books, 1996
- Darlington, David. *The Dreamland Chronicles.* London: Little Brown, 1997
- Good, Timothy. *Above Top Secret.* London: Grafton Books, 1989
- Greene, Brian. *The Elegant Universe.* London: Vintage, 1999
- Harwood, William. B. *Raise Heaven & Earth. The Story of Martin Marietta People & Their Pioneering Achievements.* New York: Simon & Schuster, 1993
- Henshall, Philip. *The Nuclear Axis.* Stroud, UK: Sulton, 2000
- . *Vengeance. Hitler's Nuclear Weapons, Fact or Fiction.* Stroud, UK: Alan Sutton Publishing, 1995
- Hirschfeld, Wolfgang and Brooks, Geoffrey. *Hirschfeld: The Secret Diary of a U-Boat.* London: Orion, 1997
- Jane's Fighting Aircraft of World War Two.* London: Jane's Publishing Company, 1946-47
- Jones, R.V. *Most Secret War.* London: Coronet, 1979
- Lomas, Robert. *The Man Who Invented the 20th Century—Nikola Tesla, Forgotten Genius of Electricity.* London: Headline Book Publishing, 1999

- Lucas, James. *Last Days of the Reich*. London: Grafton Books, 1987
- Lusar, Rudolf. *German Secret Weapons of World War II*. London: Neville Spearman Ltd., 1959
- Manning, Jeane. *The Coming Energy Revolution*. New York: Avery Publishing Group, 1996
- Masters, David. *German Jet Genesis*, London: Jane's Publishing Company, 1982
- Miller, Jay. *Lockheed's Skunk Works—The Official History*. Leicester, UK: Midland Publishing, 1995
- Overy, Richard. *The Penguin Historical Atlas of the Third Reich*. London: Penguin Group, 1996
- Pisziewicz, Dennis. *The Nazi Rocketeers—Dreams of Space and Crimes of War*. New York: Praeger Publishers, 1995
- Pitt, Barrie (editor). *The Military History of World War Two*. Twickenham, UK: Temple Press, 1988
- Rhodes, Richard. *The Making of the Atomic Bomb*. New York: Touchstone, 1988
- Rich, Ben R. and Janos, Leo. *Skunk Works*. London: Little Brown, 1994
- Simpson, Christopher. *Blowback*. New York: Weidenfeld & Nicolson, 1988
- Speer Albert. *Infiltration: How Heinrich Himmler Schemed to Build an SS Industrial Empire*. New York: Macmillan Publishing Co., 1981
- Speer Albert. *Inside the Third Reich*. London: Weidenfeld & Nicolson, 1970
- Trevor-Roper, Hugh (editor). *The Goebbels Diaries: The Last Days*. —. *The Last Days of Hitler*. 7th Edition. London: Macmillan, 1995
- Vajda, Ferenc A. and Dancey, Peter. *German A/C Industry & Production 1933-45*. Shrewsbury, UK: Airlife Publishing, 1998
- Valone, Thomas MA, PE, (editor). *Electrogravitics Systems: Reports on a New Propulsion Methodology*. Washington: 2nd Edition. Integrity Research Institute, 1995
- Vesco, Renato. *Intercept—But Don't Shoot*. New York: Grove Press, 1971
- Wiener, Tim. *Blank Check*. London: Warner Books, 1990
- Witkowski, Igor. *Supertajne Bronie Hitlera*. Warsaw: WIS, 2000
- Zuk, Bill. *Canada's Flying Saucer Projects. The Story of Avro Canada's Secret Projects*. Boston Mills Press, 2001

Articles

- "Black World Engineers, Scientists Encourage Using Highly Classified Technology for Civil Applications," William B. Scott, *Aviation Week*, 9/3/92
- "Breaking the Law of Gravity," Charles Platt, *Wired*, 3/98
- "Military Power," Bill Gunston OBE, FRAeS, *Air International*, 1/00
- "Nordhausen," Karel Mevgrly, *After the Battle*, No. 101
- "Secret Advanced Vehicles Demonstrate Technologies for Military Use," William B. Scott, *Aviation Week*, 1/10/90
- "Secrets by the Thousands," C. Lester Walker, *Harper's Magazine*, 10/46
- "Taming Gravity," Jim Wilson, *Popular Mechanics*, 10/00
- "The Universe's Unseen Dimensions," Nima Arkani, Savas Dimopoulos and Georgi Dvali, *Scientific American*, 8/00
- "The U.S. Antigravity Squadron," Paul LaViolette, Ph.D., *Electrogravitics Systems* (Op Cit)
- "Without Stress or Strain . . . or Weight," Intel, *Interavia*, 5/56

Reports

- Electro aero dynamic s in Supersonic Flow*, M.S. Cahn and G.M. Andrew, The Northrop Corporation, AIAA, 1/68
- Electrogravitics Systems: An Examination of Electrostatic Motion, Dynamic Counterbary and Barycentric Control*, Aviation Studies (International Limited) 1956
- Project LUSTY*, U.S. Air Force History Office 1945
- Project Winterhaven*, The Townsend Brown Foundation 1952
- USAF Electric Propulsion Study*, Dennis J. Cravens
- USAF Report: 21st Century Propulsion Concepts*, R.L. Talley, 5/91

Index

Page numbers of illustrations appear in italics.

- A-12 (Mach 3.2 A-12 Blackbird), 8, 18, 14, 119, 126, 144, 246, 248. *See also* SR-71 Blackbird
Abelman, Daniella "Dani," 7-9, 21, 85
"About Gravitation, Vortices and Waves in Rotating Media" (Hilgenberg), 194
Advanced Deep Space Transport (ADST), 108
AEG, 204
Aerospace industry, 224-28, 239-40, 247, 252, 254
Agoston, Tom, 159, 160-69, 172, 176, 177, 180, 181, 183, 199, 203
Air Force Office of Special Investigations (AFOSI), 32
Air International magazine, 125
Alexander, Col. John B., 262, 263, 266, 268
Alsos Mission, 178-79, 220-21
American Institute of Aeronautics and Astronautics (AIAA), 129-30, 134, 137, 138
Anderson, Lindsay, 115-16
Antigravity, 14
 aerospace applications, 235-36
 article, 1996, 101-3, 702
 articles on, 1950s, 1-4, 5-7, 12, 17-19, 208
B-2 Stealth Bomber and, 91, 121-22, 123-26, 129-30, 132-37
"Bell, the," Germany, WWII, 182-84, 188, 191-201, 202, 228-29, 232, 234
black programs, 14, 16, 33, 79, 89, 90, 115
 economic effect of, 81, 144-45, 239, 251
experiments, 5, 7, 23, 24, 58, 83-84
 five pathways to, 117, 121, 141
 gravity shielding as, 103, 104, 107, 117, 121, 228, 231, 234, 275
Hutchison Effect and, 265-67
lévitation and, 10-11, 211, 213, 233, 236, 237
Lockheed Martin and, 10-11
NRL, 24
Philadelphia Experiment, 24-27, 130, 137-38, 272
Podkletnov discovery, 101-7, 113-14
propulsion system, 1-4, 44, 208-22, 222, 223
Puthoff and, 79, 221
Schauberger and, 208-22, 222, 223-27, 250-51
Sheffield, England, talk by Podkletnov, 228, 230-31, 235-37
silence concerning research or pursuit of, 12
U.S. shutting down of white programs, 223-27, 251
weight reduction of object associated with, 102, 103, 104, 121-22, 231, 235, 236, 269
See also Trimble, George S.
"Antigravity: The End of Aerodynamics?" (Young), 230
Area 51 (Groom Lake, NV), 116, 119, 128, 199
 A-12 test flight, 246
alien spacecraft stories and, 149
link with Edwards, 139-40
sightings of aircraft around, 148-49
Tikaboo Valley and Extraterrestrial Highway, 149
Astra, 145, 150
Atomic Shield, 178
Augsburg, Emil, 277
Aurora, 14-15, 16, 126, 127, 144, 148, 246, 249
 link to Kingfish, 245[^]9
 sighting, 1989 and 1990, Belgium, 16
 sighting, 1992, North Sea, 15, 246
Austria
 Bad Ischl, Schauberger home, 204, 238
 Ebensee (Zement), underground Nazi complex, 181-82, 206
Leonstein, 219, 220
Russian army in, 219, 236-37
Vienna, Schauberger's apartment, 236-37

- Vienna, secret Nazi research facility, 162
Vienna, SS technical engineer school, 218, 237
Aviation Studies (International), Ltd., 19
Aviation Week, 5, 123, 125-26, 134, 144, 248
Avro company, Canada, 226, 250
Avrocar, 73-74, 76
Project Silverbug/Y2 (U.S. Air Force flying saucer project), 73-77, 226, 250
- B-2 Stealth Bomber, 119, 121, 122, 123-26, 128-29, 132-37, 142
B-24 Liberator bomber, 245
B-36, 245
B-58 (first supersonic bomber), 245
BAE Systems (formerly British Aerospace [BAe] Defence), 79, 82, 83-85, 230
Bahnson Company, 34
Bakersfield, CA, F-117A Stealth Fighter crash, ix-xi, 240^1, 244
Voodoo fighter remains scattered at, 244
Barbie, Klaus, 277
Bearden, Thomas E., 277
Belgium, sightings of aircraft, 1989, 1990, 16
Bell, antigravity device (*Die Glocke*, code names *Laternenträger* and *Chronos*), Germany, WWII, 182-84, 188, 191-201, 202
massacre of Nordhausen scientists and, 183-84, 187, 200-201, 233, 251
meaning of code names, 228
as time machine, 229, 232
zero-point energy field and, 232-33
Bell, Lawrence D., 3, 4, 7, 12, 20
Bell Aircraft, 3, 20, 35
Belluzzo, Giuseppe, 45, 54, 163
Berlitz, Charles, 25, 26, 137
Bernadotte, Count Folke, 165
Bertrandias, Maj. Gen. Victor E., 29-30, 32, 33, 123
Biefeld, Paul Alfred, 23
Black operations, 15, 96-97, 121, 250, 269 A-12, 144, 150, 248
Air Force projects, 147-48
antigravity development, 14, 16, 33, 89, 115, 121
Aurora, 126
budget, 120, 126-27
"deep black," 116, 243, 249
disinformation and, 17, 26, 27, 76, 137, 138, 139, 149-50, 227, 272
F-1 17A Fighter crash at Bakersfield and, ix-xi, 240-41, 244
- German, 159, 160-69, 177, 178, 179-80, 187, 220
Kingfish program, 245-49, 250
Operation Paperclip becomes, 243
origins, 251, 269
Pentagon and, 146
"quantum leap technologies" and, 116, 244
remote viewing (RV), 78-79, 112-13, 238
Shiva Star, 14, 15, 96
Skunk Works (and Plant 42), 8-9, 10, 115-16, 119, 127, 135, 136, 139, 140, 142-45, 149, 239, 246, 247
stealth development, x-xi, 46, 115-22, 124, 240
Timberwind or nuclear thermal rocket, 94
See also Area 51
- Blowback—The First Full Account of America's Recruitment of Nazis, and Its Disastrous Effect on Our Domestic and Foreign Policy* (Simpson), 242^43, 269, 277
- Blunder! How the U.S. Gave Away Nazi Supersecrets to Russia* (Agoston), 159, 160-69, 174, 176, 177
- BMW (Bayerische Motorenwerke), Bramo, 51-52, 53
Boeing Corp., 20, 140, 265, 268 Model 15A, 23
Boiling Air Force Base, Washington, D.C., 63, 64
Bormann, Martin, 153, 180, 186, 187, 199
Bosch, 204
Bower, Tom, 60, 243
"Breakthrough As Scientists Beat Gravity" (Matthews & Sample), 101-3, 702
Breakthrough Propulsion Physics program (BPP), 99-100, 114, 229, 233, 258
Breslau, Germany (Wroclaw, Poland), 45, 163, 182-84, 185-90, 217
subterranean facility, 52-53
- Britain
Bawdsey Manor, antenna masts, 87-88
Nazi technology, Canadian testing of, 57, 61
Nazi technology, obtaining, 59-62
Sheffield talk by Podkletnov, 228, 230-31, 235-37
U.K. Public Records Office, 60
- British Intelligence Objectives
Subcommittee (BIOS), 57, 58-59
- Brown, Thomas Townsend, 7, 20-30, 33-34, 35, 44, 46, 76, 123, 124, 125, 130, 136, 137, 141, 142, 144, 227, 258, 272
- Gravitor, 23, 29

- Brown, Thomas Townsend (*cont'd*)
 Philadelphia Experiment and, 24-27,
 130,138,272
 Project Winterhaven, 30, 31-33, 123,
 138,144
 Brzezinski, Zbigniew, 116
 Bushman, Boyd, 10-11, 240, 249, 252-56,
 257,276
 videotape from, 253
- Cady, Willoughby M, 32-33
 Campbell, Glenn, 139-40
 Canada
 Nazi technology, testing of, 57, 61
 "Silverbug affair," 73-77
 Carpenter, Joel, 46
 Carroll, Maj. Gen. Joseph F., 32
 Casimir, Hendrik, 109
 Casimir Effect, 109
 CIA (Central Intelligence Agency)
 Kingfish program and need for high-flying, super-fast spy plane, 245-49, 250
 Mach 3.2 A-12 Blackbird and, 246
 Nazis recruited by, 243, 251-52, 277
 strong-arm tactics by, Kingfish program, 248-49
See also Aurora
 Clarke, Arthur C., 276
 Clarke Electronics, 20
 Coanda effect, 221-22, 233
 Coats, Callum, 221,224, 233
 Combined Intelligence Objectives
 Subcommittee (CIOS), 57, 58-59, 60, 62, 177
 Convair (formerly Consolidated Vultee Aircraft), 20, 35, 245
 B-24 Liberator bomber, 245
 B-36, 245
 B-58 (first supersonic bomber), 245
 black operations at, 247
 Widmer at, 245
 Craig, Gen., 29-30, 32, 33, 123
 Cross, Lawrence, 15-17, 21, 26, 47-48, 53, 57, 58, 63, 77, 204, 205, 206
 Curtiss-Wright, 20
 Czechoslovakia
 Brno (Bruenn), 160, 163
 Gablonz (Jablonec), 214
 Neudek (Nejdek) weapons factory, 216
 Pilsen, 160, 163, 172, 175, 178
 Prague, German WWII research, 45, 51, 52,76-77,163
 Skoda Works, 160, 169,174, 175-84, 195,202,252
- U.S. Third Army (Patton's troops) in, 173-74,175-76
- Degaussing, 24, 26-27
 Degenkolb, Gerhard, 163, 167, 168
 Desert Rats, 139,147
 Disinformation, 17, 26, 27, 76,137, 138, 139, 149-50,227
 Philadelphia experiment as, 272
 UFOs as, 149-50, 227
- Doenitz, Adm. Karl, 65
 Donner, Robert, 224, 225
 Dornberger, Walter, 165-66, 242, 243
 Douglas Aircraft Company, 19
 "Drag" or air resistance of aircraft, 129, 132-33,134-35
- Edison, Thomas, 258-59
 Edwards Air Force Base, 139
 Einstein, Albert, 85, 98, 194, 254, 256
 Electricity
 AC current, 258-59, 260
 gigawatt, 271
 Hutchinson and, 260-61
 measurement of, 271
 Tesla and, 258-60
 Van de Graaf generator, 265
- Electric Propulsion Study* (U.S. Air Force), 84, 88, 89
 "Electroaerodynamics in Supersonic Flow" (Northrop), 134, 137, 138
- Electrogravitics Systems—An Examination of Electrostatic Motion, Dynamic Counterbary and Barycentric Control*, 17-20, 30
- Electromagnetism, 100-101, 232, 234, 258, 261
- Electrostatic fields, 129-30, 132-33, 141-42
- Eradication (clean-up, red-lining) of information, 137, 138, 180
 Los Alamos report on Hutchinson Effect, 268,271
- Evans, Ron, 81-82, 83, 105, 230
- F-16 fighter, 240, 245
 F-22 fighter, 240
 F-111 bomber, 15
 F-117A Stealth Fighter, x-xi, 46, 116, 118, 119, 120, 125, 129, 136, 144, 145, 240
- Faraday, Michael, 85
 Fedden, Roy, 61, 62
 Fegelein, Hermann, 167
 Fifth dimension (hyperspace), 233-34
 Fleming, Ian, 59
Flight International, 5

- Flying saucer
 Brown's, 23, 24, 31-33, 34, 43, 124
 Canada efforts to create, 73-77, 226, 227
 German, 45-46, 47, 51-54, 58, 77, 163,
 204-22, 222
 Schauberger and, 211, 272, 213-22, 222
 U.S. Air Force's, 226
See also Foo-fighters
 "Flying Top," 51, 52, 58, 213, 215
 Marienehe, Germany, 213, 215
 Foo-fighters, 40-44, 54-57, 68-70, 72, 220
 Kugelblitz (Ball Lightning Fighter), 56
 Feuerball (Fireball), 55
 Fort Worth, 244-45, 254
 Air Force Plant No. 4, 245
 Lockheed Martin Division at, 240, 241,
 245
 Frost, John, 73-77, 226
 Frost, Tony, 75-76

 Galileo, 254
 General Dynamics (GD Convair), 245, 247,
 249, 254
 loss of Kingfish contract, 246
 General Electric, 6, 20, 34, 124
 "G-Engines Are Coming!, The" (Gladych),
 14, 3, 6
 Gerchsheimer, Karl, 224-25
 Sherman, TX, facility, 224-25
 Gerlach, Walther, 183, 194
German Secret Weapons of World War II
 (Lusar), 44-46, 47, 214
 Germany (Nazi Germany)
 A-4 rocket, 153-54, 155, 156, 165, 168
 A-9/A-10, 167
 antigravity research, 182-84, 188,
 191-201, 202, 234-35
 anti-Hitler elements, 155, 165
 atomic bomb and nuclear development,
 178-80, 182, 183, 202, 203, 220-21
 "Bell, the," 182-84, 188, 191-201, 202,
 228-29
 concentration camps, for labor pool for
 weapons projects, 154-57, 170, 171,
 172, 188, 204, 217, 218
 evacuation of secret weapons, 187-90,
 199, 202, 204
Fleissiges Lieschen ("Busy Lizzie"), 45
 "Flying Top," 51, 52, 58, 213, 215
 foo-fighters, 40-44, 54-57, 68-70, 72,
 220
 General Plan 1945, 186, 189
Graf Zeppelin, 87-88
 guided weapons technology and
Windhund, 69-70, 72

 Kammlerstab, 160-61, 172, 177, 180, 203,
 252
 massacre of Nordhausen scientists,
 183-84, 187, 200-201, 233, 251
 Mittelwerk GmbH, 156, 165, 170
 naval advances, 61
 remote interference with aircraft, 68,
 70-71, 203, 220
 RLM (State Air Ministry), 51
 Skoda Works, Czechoslovakia, 160, 169,
 174, 175-84, 195, 202, 252
 surrender, 176, 219
 SS, 155, 172, 175, 200-201
 SS FEP (*Forschungen, Entwicklungen und*
 Patente), 194-95
 SS oversight of secret weapons research,
 151-52, 153-59, 160-69, 178,
 199-200, 202-3, 217, 238, 243
 SS "Special Evacuation Kommando"
 183, 186, 187, 189, 190, 195, 198, 199
 tech-plunder of, 59-63, 64-73, 71, 151,
 162, 172, 178, 195, 206, 250, 251
 underground facilities, 66, 67-68, 157,
 181-82, 188, 193, 196-201
 underground weapons factory,
 Nordhausen, 56, 156-57, 165-66, 167,
 170-74, 183-84, 187, 200-201, 243
 V-1 flying bomb, 44, 170
 V-2 ballistic missile, 44, 56, 151, 157,
 168, 170, 171, 251, 269
 Vienna, SS technical engineer school,
 218, 237
 Vienna secret research facility, 162
 World War II secret weapons, 44-47, 51,
 187, 250, 251, 269
 Gladych, Michael, 1, 3
 Glenn Research Center, Lewis Field,
 Cleveland, 99-100, 258
 Goebbelsjoseph, 158, 164
 Goering, Hermann, 55, 157, 161, 167, 176
 Gordon, Jack, 136, 142-45
 "Gravitoelectric-electric coupling via
 superconductivity" (Torr & Li), 103
 Gravity, 11, 13-14, 79-80, 234
 electromagnetism and, 100-101
 heat and, 85
 reduction and distance, 106
 shielding (antigravity), 103, 104, 107,
 117, 121, 228, 231, 234, 235, 275-76
 superconductivity and, 103, 106-7, 194,
 231
 ZPE and, 113, 117, 121, 231
 Grawitz, Ernst, 193
 Groom Lake, NV. *See* Area 51
 Groves, Gen. Leslie, 147, 178

- Gunston, Bill, 124-25
 Gyroscopes, 82, 83, 104, 237
- Habermohl, Klaus, 45, 52, 53, 54, 163
 Hanke, Karl, 186, 187, 188
 Hathaway, George, 257, 258, 260, 261, 262, 267, 268
 Heinert, Ernst, 50, 213, 215, 214, 215
 Heinkel Aircraft Company, Marienehe, Germany, 50, 215
 He 111, 50
 He 162, 167
 He 178, 50
 He 280, 50, 213, 215
 HeS 3 A turbojet, 51
 Heisenberg, Werner, 178
 Henshall, Philip, 199
 Hermann Goering Works, Linz, 218—19
 Heydrich, Reinhard, 156, 160, 186
 Hilgenberg, O. C., 194
 Hiller Company, 19-20
 Himmler, Heinrich, 151, 153, 154, 155, 157, 160, 161, 163, 164, 167, 168, 177, 186, 199, 203, 205
 Hitler, Adolf, 153, 158, 159, 161, 164, 175, 179, 209
 Holocaust, 200-201, 206, 222, 277
 Houck, Jack, 267
 Howe, C. D., 74
 "How I Control Gravitation" (Brown), 23
 Hughes Aircraft, 254
 Hutchison, John, 256, 257-58, 260-67
 Intelligence and Security Command (INSCOM) visit to, 262-63, 264, 265-67, 272
 laboratory, last, 267
 laboratory, Lynn Valley, 260
 laboratory, North Vancouver, 262, 265-67
 phenomena and, 272
 psychokinesis and, 261, 267, 268
 videotape of experiments, 253
 ZPE and, 261
 Hutchison Effect, 260, 261
 Boeing and McDonnell Douglas interest in, 265, 267, 268
 description of various results, 264-67
 lévitation by, 253, 260, 261, 263, 267
 Los Alamos report on, 268, 271
 molybdenum rod results, 262-63
 Pentagon funding, 261-62
 time-dilation and, 267
 transmutation of metal and, 261, 267, 269-70, 271
 Hyperspace, 233-34
- I. G. Farbenindustrie, 62
 IBM patent-server website, 237
 Implosion energy generation, 57-58, 204, 205, 207-8, 210, 211, 223, 225, 251
 bomb potential of, 271
 Inertia, 100, 113, 117, 121, 233, 267
 Inertial-thrust machine, 82-83
Infiltration (Speer), 203
 Institute for Advanced Studies (IAS), Austin, TX, 78, 79, 108, 271
 Integrity Research Institute, 17, 18, 21, 138
 Intelligence and Security Command (INSCOM), 262-63, 264, 265-67, 271, 272
Interavia Aerospace Review, 5—7, 32
Intercept—But Don't Shoot (Vesco), 48, 54-57, 61, 72, 73, 77
- Jane, Fred T., 5**
Jane's All the World's Aircraft, 5
Jane's Defence Weekly (JDW), 2, 45, 15, 44, 84, 248
Jane's Fighting Ships, 5
 Japan, antigravity research, 235, 236, 237
 Johnson, Kelly, 143, 149, 239, 246
 Joint STARS, 146
 Junkers company, Dessau, 178
- Kaltenbrunner, Ernst, 186-87
 Kammler, Hans, 155-59, 160-69, 171, 172-73, 174, 177, 180-81, 183, 189, 191, 194-95, 199, 202, 242-43
 denies Himmler's request for a "heavy truck," 163, 168, 190, 198-99, 202
 fate of, 252, 277
Kammlerstab, 160-61, 172, 177, 180, 203, 252
 special projects, 159, 160-69, 172, 176, 178
 Kepler, Johannes, 207
 Kidd, Sandy, 82-83, 84, 105
 Kingfish, 245-49, 250
 Mach 6.25 capability, 248
 ramjets and pyro-ceram, 248
 Kirtland Air Force Base, NM, 14
 Koller, Gen., 65
 Krupp, Meppen, Germany, 61
 Kummer, SS Major, 167
- Laithwaite, Eric, 83
 Las Vegas, NV, 131-32
 LaViolette, Paul, 17-19, 30, 33, 34, 47, 123, 129, 138
 Lazar, Robert, 149
 Lear, William P., 3-4, 7, 12

- Lear Inc., 3, 20
Legend, the, 47, 50-53, 54, 56, 57-58, 204, 208, 213, 214, 219-20
Leet, Maj. William, 42
Leoning, Grover, 20
Lévitation, 10-11, 211, 213, 233, 236, 237, 260, 261, 263
Li, Ning, 103, 275
Light-speed barrier, 85, 96
 faster-than-light travel, 101, 117, 121, 233, 276
Lindeke, Ron, 142-43
Lockheed Martin, 7, 20, 21, 85, 254
 acquisition of General Dynamics, 245
 anti-gravity and, 10-11, 240
 Astra, 145
 black programs, 118-19, 124, 144
 DarkStar, 145
 F-16 fighter, 240, 245
 F-22 fighter, 240
 F-1 17A Stealth Fighter, x-xi, 46, 116, 118, 119, 120, 125, 129, 136, 144, 145, 240
 Fort Worth Division, 240, 241, 245
 Kingfish project, 246-49
 Mach 3.2 A-12 Blackbird (Oxcart program), 8, 18, 14, 119, 126, 144, 246, 248
 radar test ranges, 140
 Skunk Works (and Plant 42), 8-9, 10, 115-16, 119, 127, 135, 136, 139, 140, 142-45, 149, 239, 246, 247
 SR-71 Blackbird, 145, 246
 U-2 spyplane, 36, 38, 56, 118-19, 145, 149, 150, 245
 XP-80, 143, 145
 YF-22, 145
Longhurst, John, 19
Lopez, Amelia, ix-xi, 241
Los Alamos National Laboratory, 262
 "Big Bad Bob" scientist observing
 Hutchison Effect, 263, 266-67
 report on Hutchison Effect, 268, 271
Luftwaffe
 air weapons, 66
 Ar 234, 52, 167
 He 162, 167
 Ju 290 and 390, 190, 198-99, 202
 Me 163, 52, 66
 Me 262, 52, 66, 143, 151, 163, 167
 surrender of, 65
Lusar, Maj. Rudolf, 44-46, 47, 214
Lusty (Luftwaffe Secret Technology), 48-49, 63, 64-73, 77, 152, 162
Lyles, Garry, 92, 95-97, 207, 274
Macauley, John, 73
Magnetic fields separation, 194
Manhattan Project, 10, 20, 31, 38, 89-90, 120, 178, 250
Markus, Dan, 81, 84, 85, 86-92, 97, 108-10, 117, 121, 123, 133-36, 142, 152, 153, 198, 202, 204, 205, 228-29, 230, 231, 232, 233, 234, 237-39, 242, 243, 244, 251, 257, 269-73
Marshall Space Flight Center, Huntsville, AL, 93-95, 98-99, 101, 103, 275
 DeltaG group, 103-4
Martin-Marietta (formerly Martin Company, Glenn L. Martin Aircraft Company), 2-3, 8, 10, 20, 35, 226
 Research Institute for Advanced Studies (RIAS), 9-10, 226-27, 239, 240
 See also Lockheed Martin
Matschke, Dieter, 277
Mauthausen concentration camp, Austria, 217, 218, 237
Maxwell Air Force Base, AL, 48-19, 63
Mazuw, Emil, 195
McDonald, Gen. George C, 66, 67, 68
McDonnell Douglas, 265, 267, 268
Meiers, Don, 40, 41, 55, 56
Mengele, Josef, 193
Men-in-black, x, 17, 140-41
Messerschmitt works, Augsburg, Bavaria, 216
Miethe, Richard, 45, 52-54, 163
"Military Power" (Gunston), 125
Millis, Marc, 99-104, 257
Molybdenum, 262-63
Montgomery, Gen. Bernard, 60
Moore, William, 25, 26, 137
Motionless Magnetic Generator, 277
Muellner, George, 146-48
NASA (U.S. National Aeronautics and Space Administration), 85, 92
AMMP (Advanced Manned Missions Program), 275
antigravity research, 101, 103-7, 141-42, 227, 239, 240, 275
ASTP (Advanced Space Transportation Program), 85, 94-96
black programs and, 97-98
Breakthrough Propulsion Physics program (BPP), 99-100, 114, 229, 233, 258
Gerchsheimer at, 224
Glenn Research Center, Lewis Field, Cleveland, 99-100
interstellar travel and, 95-96

- NASA (*cont'd*)
 Marshall Space Flight Center, 93-95, 98-99, 101,103
 Nazis recruited for, 243
 nuclear rocket development (NERVA and ROVER), 94
 Podkletnov's experiment and, 235
 "Project Greenglow," 85, 230
 National Atomic Research Laboratories, Brookhaven, Long Island, NY, 225
 National Investigations Committee on Aerial Phenomena (NICAP), 34
 NATO's Advisory Group for Aerospace Research and Development (AGARD), 83
 Naval Research Laboratory (NRL), 24
 Negative-energy density matter, 98
 Nellis Air Force Range, 131
 Neumann, Otto, 187
New York Herald Tribune, 275
New York Times, 211
 Nondifferential linear equations, 11
 North American Inc., 20
 Northrop Grumman
 B-2 Stealth Bomber, 119, 121, 122, 123-26,128-29,132-37, 142
 electrostatic field research, 129-30, 135, 137
 P-61 Black Widow, 39
 RCS range at Tejon Ranch, 140
Nuclear Axis, The (Henshall), 199
 Nuremberg war crimes trial, 180
 O'Brien, Lt. Col., 65
Observer(London), 276
 Odom, Charles, 42
 Office of Naval Research (ONR), 32, 35, 262
 Operation Paperclip, 60, 93, 151, 243
 as deep black program, 243
 Oppenheimer, Robert, 10
 "Overcast," 242
 Overholser, Denys, 118
 Oxcart, A-12 development program, 246, 248
 P-61 Black Widow, 39
 Pabst von Ohain, Hans-Joachim, 51
Paperclip Conspiracy, The (Bower), 60, 243
 Parallel universes, 276
 Pash, Boris T., 178-79, 183, 220
 Patton, Gen. George, 173-74, 175-76
 Peary, Robert, 259
 Pentagon
 black operations and, 146
 Hutchison Effect and, 261-62
 Joint Intelligence Objectives Agency, 243
 Pezarro, Alexis, 257-58, 260, 262, 268
 Pharos Technologies, 260, 268
 Phenomena, 272
Philadelphia Experiment, The (Berlitz & Moore), 24-27,130,137-38, 150, 272
 Planck, Max, 209
 Platt, Charles, 105-7, 235
 Plenum, 234
 Podkletnov, Evgeny, 101-7, 113-14, 121, 194, 204, 209, 228, 255,275
 Bell and, 194,231
 father's patent and, 237-38
 gravity-shielding and, 103, 104, 107, 117, 121,228,235
 Schauberger's papers and, 236-38
 Sheffield, England, talk, 228, 230-31, 235-37
 Pohl, Oswald, 154
 Poland
 Breslau (Wroclaw), 45, 52-53, 163, 182-84,185,188-90, 217
 Leubus (Lubiaz) secret facility, 188
 Opeln (Opole), Ju 290 and 390 base, 190, 198-99
 Waldenburg (Walbrzych), 193
 Wenceslas Mine, Riese underground complex, 188, 193, 196-201, 209, 232, 234-35
 See also Bell, antigravity device
Popular Mechanics, 215
 Possony, Stefan, 45, 46
 Prawinjakub, 186,189-90
 Project Greenglow, 85, 230
 Project 1794, 74
 Project Silverbug/Y2 (U.S. Air Force flying saucer project), 73-77, 226, 250
 Project Winterhaven, 20, 30-33, 123, 138, 144
Psychic Discoveries Behind the Iron Curtain, 112
 Puthoff, Harold E. "Hal," 78-79, 92, 107, 108,110-14,142,229,231,233,238, 271,272
 Quantum Theory, 231
 Quantum vacuum (foam), 109, 110-11, 272
 Radford, Adm. Arthur W., 29
 Radial-flow-gas-turbine (RFGT), 52, 75, 76
 Reagan, Ronald, 119, 120, 126, 262
 Reinemer, Gunter, 277
 Remote viewing (RV), 78-79, 111-12, 238

- Research Institute for Advanced Studies (RIAS), 9-10, 226-27, 239, 240
- Reynolds, Josh, 29
- Rich, Ben, 115, 116, 117, 119, 144, 239, 249
"virtual" warehouse of too-dangerous projects, 117, 249
- Rickman, E. J. C., 83
- Ringwald, Fred, 39-42, 56
- Rolls Royce aero-engines, 230
- Russia, 56
advance into Nazi territory, 56, 157, 181, 188-89
- Cold War and, 115, 242
- intelligence agents search and destroy Schaefer's Vienna apartment, 236-37
- remote viewing (RV), 78, 238
- SA-2 Guideline, 245
- Soviet secret intelligence service, 186
- stealth and, 116-17
- torsion fields, 238
- Sakharov, Andrei, 113
- Schauberger, Joerg, 206, 207, 208, 209, 222
- Schauberger, Viktor, 57-58, 204-22, 212, 222, 255
American custody of, 206, 220, 223, 250
- bio-technology, 207
- centripetal force, 210, 215, 221
- contacted by American and Canadian aircraft companies, 226
- death of, 225
- flying saucer (antigravity device), 211, 272, 213-22, 222, 233, 236
- Heinkel, 213, 215, 214, 215
- implosion energy generation, 204, 205, 207-8, 210, 211, 223, 225, 251
- Red Army search and destruction of Vienna apartment, 236-37
- Repulsine and Repulsator, 211, 272, 213, 216, 218, 219, 221, 222, 223, 232, 250-51
- Sherman, TX, work with Gerchsheimer, 224-25, 226
- SS associations, 205, 213, 214-15, 217
- trout turbine, 206, 211, 226
- zero-point energy and, 221-22, 222, 233
- Schauberger, Walter, 58, 204, 206, 214, 225, 251
- Schellenberg, Walther, 165
- Schlüter, Lt. Ed, 39-42, 55, 56
- Schmidt, George, 98-99
- Schriever, Rudolf, 45, 50-52, 53, 57, 58, 163, 213, 215
- Schroeder, Bill, 118
- Schulgen, Gen. George, 35, 38, 44, 141, 150, 250
- Schuster, Rudolf, 186, 189
- Scientific American*, 276
- Scott, Bill, 123, 125, 127, 128, 129, 143
- Sequoia National Forest, ix-xi. *See also* Bakersfield, CA
- Shiva Star, 14, 15, 96
- Siemens, 204
- Sikorsky, 20
- Simpson, Christopher, 242-43
- Skunk Works (Rich)*, 115, 144
- Sommerfeld, Arnold Johannes, 118
- Source Index, 91
- Space-time continuum, 98, 117, 121, 229
Hutchison Effect and, 267
torsion field and, 232, 234
zero-point energy and, 232-33
- "Special Projects Group," 9, 226
facilities, 8
See also Lockheed Martin, Skunk Works
- Speer, Albert, 153, 154, 155, 156, 161, 164-65, 167, 171, 178, 203, 217
- Sperry-Rand, 6, 20
- Sporrenberg, Jakob, 186, 187-89, 191, 194
- SR-71 Blackbird, 145, 246
Mach 3.2 A-12 Blackbird, 8, 18, 14, 119, 126, 144, 246, 248
- Stealth, 8, 89, 115-22, 136
B-2 Bomber, 91, 121-22, 123-26, 128-29, 132-37, 142
as deep black program, 116, 136
- F-117A Fighter, x-xi, 46, 116, 118, 119, 120, 125, 129, 136, 144, 240
- Have Blue, 116, 117, 118, 127, 136
- RCS (radar cross section), 116
- reactive skins, 148
- Skunk Works and, 8, 115
- Strategic Air Command (SAC), 245
- Strategic Defense Initiative (SDI or Star Wars), 262
- String theory, 276
- Strong nuclear force, 234
- Sullivan, Lt. Col. A. R., Jr., 68-69, 70, 72
- Superluminal quantum tunneling, 101
- Swann, Ingo, 112
- Sylvester, Gen. George, 127
- Szymanski, Wladyslaw, 186, 189-90
- Tachyons, 112
- Talley, Robert, 141
- Tampere University of Technology, Finland, 102, 105

- Tesla, Nikola, 258-60
 Tunguska and, 259-60
 Tesla Coil, 259, 265
 Texas Instruments, 254
 T-Forces, 60-62
 Theory of Relativity and General
 Relativity, 85, 98, 101, 231, 254
 30 AURN, 59-60
 Timberwind or nuclear thermal rocket, 94
 Time, 234
 Time machine, 228-29
 torsion field and, 232, 234
 See also Bell, antigravity device; Zero-point energy
 Time travel, 101
Times (London), 276, 277
 Topology, 11
 Torr, Douglas, 103
 Torsion field, 228-29, 231, 232, 233, 234,
 238, 261
 Toshiba, 235, 237
 Townsend Brown Foundation, 29—30, 32,
 33
 Transmutation of metal, 261, 267, 269—70,
 271
 Trimble, George S., 3, 4, 7, 8, 9, 11-12,
 17, 21, 31, 33, 38, 46, 47, 76, 77,
 85, 97, 101, 226, 227, 239, 249-50,
 275
 Truman, Harry, 243
*T. T. Brown Electrogravity Device, The: A
 Comprehensive Evaluation by the Office
 of Naval Research*, 138
 Tunguska, Siberia, 259-60
 "Twenty-first Century Propulsion
 Concepts" (Talley), 141-42
 Twining Lt. Gen. Nathan, 35-36, 37, 38,
 44, 141, 150, 209, 223, 250
 UCAVs (unmanned combat air vehicle
 technology), 246
 Ueberreither, Siegfried, 181, 182
 Ufimtsev, Pyotr, 89, 118
 UFO
 antigravity aircraft as, 11, 34
 disinformation and, 149-50, 227
 explanation for, 276
 Northrop and, 140
 origins, 47
 photostat, Bushman's, 255
 Roswell crash and, 151
 sightings, 35, 43, 57, 75, 125-26, 128,
 134, 140, 148, 150, 151
 Twining memo, 35-36, 37, 38, 44, 47,
 141, 150-51, 209, 223-24, 250
- United States
 aerospace industry, 224-28, 239^0, 247,
 252, 254
 Alsos Mission, 178-79, 220-21
 antigravity technology, state of in, 269
 Kammler and, 165-69, 174
 Nazis recruited by, 93, 151, 242-43,
 251-52, 269, 277
 tech-plunder activities, Nazi Germany,
 62-63, 64-73, 71, 151, 162, 172, 178,
 195, 206, 250, 251
- U.S. Air Force
 Air Material Command, 36
 anti-gravity and, 84, 88, 89, 141-42
 black programs, 97, 147-48
 crash recovery teams, 244
 F-22A Raptor, 144
 plasma-bullet, 14, 96
 Project Silverbug/Y2 (flying saucer
 development), 74-77, 226, 250
 "special studies group," 45
 SR-71, 144
 top-secret classification, 116
 Twining memo, 35—36, 37, 38, 44,
 47, 141, 150-51, 209, 223-24,
 250
 "vision," 11
 "U.S. Antigravity Squadron, The"
 (LaViolette), 123-24
 U.S. Army, Intelligence and Security
 Command (INSCOM), 262-63, 264,
 265-67, 271, 272
 U.S. Army Air Force (USAAF)
 9th, 415th Night Fighter Squadron, 39,
 43
 15th, 42
 Convair and, 245
 U.S. Counter-intelligence Force (CIC),
 162, 220-21, 223, 224
 U.S. Joint Chiefs of Staff (JCS)
 Operation Paperclip and, 251
 recruitment of Nazis and, 242
 U.S. Office of Technical Services, 63
 U.S. Patent Office, 277
USS Andrew Furuseth, 27
USS Eldridge, 25-27, 130, 137-38, 139
 Valone, Tom, 18, 19, 21, 30-31, 33, 34, 47,
 138
 Vancouver, Canada, 257, 260
 Vesco, Renato, 48, 54-57, 61, 72, 73, 77
 Von Below, Nikolaus, 167
 Von Braun, Werner, 53, 56, 93, 95, 151,
 166-68, 242, 269
 Von Karman, Theodore, 66

- Von Majewski, Georg, 176
Voodoo fighter, 244
Vortex compression, 194, 231
Voss, Wilhelm, 160-61, 162, 169, 173, 176-78, 180, 183, 203, 220
- Walz, Capt. Carl, 139
Wardencliff, Long Island, NY, 259
Watson-Watt, Robert, 88
Weak nuclear force, 234
Weight reduction of object associated with antigravity, 102, 103, 104, 121-22, 231, 235, 236, 269
- Whiteman Air Force Base, 124
Whittle, Frank, 50
Widmer, Robert, 245-49, 251, 252
Kingfish, 245-49
Winterhaven, 31-32. *See also Project Winterhaven*
Wired, 236
Witkowski, Igor, 182-84, 185-90, 193-94, 195-97, 231, 234, 251
Worcester, Richard "Dicky," 19
World War II
 German surveillance of British coast, 87-88
 Operation Paperclip, 93, 151, 243
 Russian advance into Nazi territory, 56, 157, 181, 188-89, 236-37
U.S. First Army, 165
U.S. Third Army (Patton), 173-74, 175-76
U.S. invasion after D-Day, 157
- U.S. pilots spotting foo-fighters, 39-43, 55, 220
See also Germany (Nazi Germany)
Wormholes, 98
Wright brothers, 19
Wright-Patterson Air Force Base, Dayton, OH, 18, 30, 53-54, 63, 68, 127
- Yeager, Chuck, 139
Young, Brian, 79-81
 "Antigravity: The End of Aerodynamics?" (lecture, 1991), 230, 231
- Zeiss, 204
Zement, underground Nazi complex, 181-82, 206
Zeppelin Werke, Schwarzwald, Germany, 56
- Zero point, 100
Zero-point energy (ZPE), 100, 108, 109, 110-11, 113-14, 117, 121, 232, 273, 276
 Bell, the, and, 232-33
 electromagnetism and, 232, 261
 gravity and, 113, 117, 121, 231
 Hutchinson and, 261
 hyperspace and, 234
 potential of, pro and con, 273
 reactor, 272
 Schauberger and, 221-22, 222, 233
 space-time and, 232-33

ABOUT THE AUTHOR

Nick Cook is the Aviation Editor of *Jane's Defence Weekly*, the world's leading military affairs journal, a position he has held for twelve years. He has also written for national U.K. newspapers, including the *London Times*, *Sunday Times* and *Sunday Telegraph*, and has had many of his stories syndicated worldwide.